

A Local Energy Identity for Parabolic Equations with Divergence-Free Drift

Francis Hounkpe*

Mathematical Institute, University of Oxford, Oxford, UK

March 7, 2024

Abstract

We prove a local energy identity for a class of distributional solutions, in $L_{2,\infty} \cap W_2^{1,0}$, of parabolic equations with divergence-free drift.

1 Introduction

We are considering the parabolic equations of the type

$$\partial_t u - \operatorname{div}(a \nabla u) + b \cdot \nabla u = 0,$$

where a is a bounded, symmetric and uniformly elliptic matrix and b a divergence-free vector field belonging to $L_\infty(BMO^{-1})$. We say that a divergence-free vector field b belongs to the space BMO^{-1} if there exists a skew symmetric matrix d belonging to BMO such that $b = \operatorname{div}(d)$. Therefore, the above equation can be rewritten as follows:

$$\partial_t u - \operatorname{div}(A \nabla u) = 0, \tag{1}$$

where $A = a + d$, with a as before and $d \in L_\infty(BMO)$ a skew symmetric matrix.

G. Seregin and co-authors introduced, in their paper [1], the notion of *suitable weak solutions* to equation (1), which are distributional solutions that belong to the energy class $L_{2,\infty} \cap W_2^{1,0}$ and that satisfy a particular local energy inequality. In this paper, we establish a local energy identity for distributional solutions of (1) which belong to the energy class $L_{2,\infty} \cap W_2^{1,0}$, and therefore, we prove at the same time that the local energy inequality required in the definition of *suitable weak solutions*, introduced in [1], is a direct consequence of being a distributional solution in the above energy class.

*Email adress: hounkpe@maths.ox.ac.uk;

2 Preliminaries

In what follows, we will use the following abbreviated notations: $B := B(0, 1)$ (the unit ball of \mathbb{R}^n), $Q := B \times (-1, 0)$, as well as $z := (x, t)$.

We recall that a function d is in the space $BMO(\Omega; \mathbb{R}^{n \times n})$ if the following quantity

$$\sup \left\{ \frac{1}{|B(0, r)|} \int_{B(x_0, r)} |d - [d]_{x_0, r}| dx : B(x_0, r) \subset \subset \Omega \right\},$$

with $[d]_{x_0, r}$ the average of d over $B(x_0, r)$, is bounded; and a function u belongs to the Hardy space $\mathcal{H}^1(\mathbb{R}^n)$ if there exists a function $\phi \in C_0^\infty(B)$ such that

$$u_\phi \in L_1(\mathbb{R}^n),$$

where $u_\phi(x) := \sup_{t>0} |(\phi_t \star u)(x)|$, and $\phi_t(x) := t^{-n} \phi(x/t)$.

For simplicity we adopt the following notation convention $\partial_i f = f_{,i}$. We have the following classical div-curl type lemma for Hardy spaces, which is a direct consequence of Theorem II.1 in [5].

Lemma 1. *Let $u \in W_p^1(\mathbb{R}^n)$ and $v \in W_q^1(\mathbb{R}^n)$, with $1 < p < \infty$ and $1/p + 1/q = 1$. Then $u_{,j} v_{,i} - v_{,j} u_{,i} \in \mathcal{H}^1(\mathbb{R}^n)$ for all $i, j = 1, \dots, n$ and we have*

$$\|u_{,j} v_{,i} - v_{,j} u_{,i}\|_{\mathcal{H}^1} \leq C \|\nabla u\|_{L_p} \|\nabla v\|_{L_q}, \quad \forall i, j = 1, \dots, n.$$

We recall also some basic facts related to the spectral decomposition of the Laplace operator on a bounded domain Ω of \mathbb{R}^n , with smooth boundary. The Laplacian viewed as an unbounded operator from $L_2(\Omega)$ into itself has a discrete spectrum; we denote by $0 < \lambda_1 < \lambda_2 < \dots < \lambda_n < \dots$ (with $\lambda_n \rightarrow \infty$), its eigenvalues and $\{\phi_k\}_{k=1}^\infty$ the corresponding eigenvectors which form a Hilbert basis of $L_2(\Omega)$. Setting $\dot{L}_2^1(\Omega)$ to be the completion of $C_0^\infty(\Omega)$ with respect to the Dirichlet semi-norm $\|u\|_{L_2^1}^2 := \int |\nabla u|^2$, and $H^{-1}(\Omega)$ to be the dual of $\dot{L}_2^1(\Omega)$, we have the following classical lemma, which gives us a Hilbert basis of $\dot{L}_2^1(\Omega)$ and a representation of the norm of H^{-1} by means of the eigenvectors and eigenvalues of the Laplace operator.

Lemma 2. *$(\phi_k / \sqrt{\lambda_k})_{k=1}^\infty$ is a Hilbert basis of $\dot{L}_2^1(\Omega)$ and as a direct consequence, we have that, if $f \in H^{-1}(\Omega)$, then*

$$\|f\|_{H^{-1}(\Omega)}^2 = \sum_{k=1}^\infty f_k^2 / \lambda_k,$$

where $f_k = \langle f, \phi_k \rangle$.

Proof. The proof of this result is quiet classical, therefore we skip it. □

3 Main Theorem

We now state the main result of this paper.

Theorem 3.1. *Let u belonging to the energy class*

$$L_{2,\infty}(Q) \cap W_2^{1,0}(Q),$$

such that

$$\int_Q u \partial_t \phi dz = \int_Q (A \nabla u) \cdot \nabla \phi dz \quad \forall \phi \in C_0^\infty(Q), \quad (2)$$

where $A = a + d$, with $a \in L_\infty(Q; \mathbb{R}^{n \times n})$ a symmetric matrix satisfying

$$\nu \mathbb{I}_n \leq a \leq \nu^{-1} \mathbb{I}_n$$

and $d \in L_\infty(-1, 0; BMO(B; \mathbb{R}^{n \times n}))$ a skew symmetric matrix. Then the following energy identity holds for all $t_0 \in (-1, 0)$ and for all test functions $\phi \in C_0^\infty(B \times (-1, 1))$:

$$\begin{aligned} \frac{1}{2} \int_B \phi(x, t_0) |u(x, t_0)|^2 dx + \int_{-1}^{t_0} \int_B \phi \nabla u \cdot a \nabla u dz &= \frac{1}{2} \int_{-1}^{t_0} \int_B |u|^2 \partial_t \phi dz \\ &\quad - \int_{-1}^{t_0} \int_B (A \nabla u) \cdot \nabla \phi u dz. \end{aligned}$$

4 Proof of Theorem 3.1

The method we use for this proof are due to Seregin, in his lecture notes: "Parabolic Equations". We start by proving a simple regularity result for the time derivative of u defined as in Theorem 3.1.

Proposition 4.1. *Let u defined as in Theorem 3.1. Then*

$$\partial_t u \in L_2(-1, 0; H^{-1}(B))$$

Proof. Step 1. Let us set

$$g(x, t) = A(x, t) \nabla u(x, t),$$

and consider the problem

$$\begin{cases} -\Delta U(\cdot, t) = \operatorname{div} g(\cdot, t) & \text{for a.e } t \in (-1, 0) \\ U(\cdot, t)|_{\partial B} = 0. \end{cases} \quad (3)$$

Let $v \in C_0^\infty(B)$, we have:

$$\begin{aligned} \int_B \operatorname{div} g(\cdot, t) v dx &= - \int_B g(\cdot, t) \cdot \nabla v dx \\ &= - \int_B (a \nabla u) \cdot \nabla v dx - \int_B (d \nabla u) \cdot \nabla v dx \\ &=: A_1 + A_2. \end{aligned}$$

We have by a straightforward computation that

$$|A_1| \leq \|a\|_{L_\infty(Q)} \|\nabla u(\cdot, t)\|_{L_2(B)} \|\nabla v\|_{L_2(B)} \quad \text{for a.e } t \in (-1, 0).$$

On the other hand, we have thanks to the skew symmetry of d , that A_2 can be rewritten as follows

$$-A_2 = \frac{1}{2} \sum_{i,j=1}^n \int_B d_{ij}(u_{,j} v_{,i} - v_{,j} u_{,i}) dx.$$

Denote by \bar{u} the extension of u from B to \mathbb{R}^n such that

$$\|\bar{u}(\cdot, t)\|_{W_2^1(\mathbb{R}^n)} \leq c \|u(\cdot, t)\|_{W_2^1(B)} \quad \text{for a.e } t \in (-1, 0),$$

where c depends only on n . Similarly, let us denote by \bar{d} the extension of d from B to \mathbb{R}^n such that

$$\|\bar{d}(\cdot, t)\|_{BMO(\mathbb{R}^n; \mathbb{R}^{n \times n})} \leq c \|d(\cdot, t)\|_{BMO(B; \mathbb{R}^{n \times n})} \quad \text{for a.e } t \in (-1, 0),$$

where, again, c depends only on n . In the later case, to construct such an extension, one can use a reflection on the boundary (See, e.g., Theorem 2 in [2], where this is done for very general domains $\Omega \subset \mathbb{R}^n$). Therefore, because v is compactly supported in B , we have that

$$-A_2 = \frac{1}{2} \sum_{i,j=1}^n \int_{\mathbb{R}^n} \bar{d}_{ij}(\bar{u}_{,j} v_{,i} - v_{,j} \bar{u}_{,i}) dx.$$

We have from Lemma 1 that $\bar{u}_{,j} v_{,i} - v_{,j} \bar{u}_{,i} \in \mathcal{H}^1(\mathbb{R}^n)$ with

$$\|\bar{u}_{,j} v_{,i} - v_{,j} \bar{u}_{,i}\|_{\mathcal{H}^1(\mathbb{R}^n)} \leq C \|\nabla \bar{u}\|_{L_2(\mathbb{R}^n)} \|\nabla v\|_{L_2(\mathbb{R}^n)},$$

and since $BMO(\mathbb{R}^n)$ is the dual of the Hardy space $\mathcal{H}^1(\mathbb{R}^n)$, we derive that

$$|A_2| \leq C \|\bar{d}\|_{L_\infty(-1,0; BMO(\mathbb{R}^n; \mathbb{R}^{n \times n}))} \|\nabla \bar{u}\|_{L_2(\mathbb{R}^n)} \|\nabla v\|_{L_2(\mathbb{R}^n)},$$

and a fortiori

$$|A_2| \leq C \|d\|_{L_\infty(-1,0; BMO(B; \mathbb{R}^{n \times n}))} \|\nabla u\|_{L_2(B)} \|\nabla v\|_{L_2(B)},$$

(with C depending only on n). Hence, we have that $\operatorname{div} g(\cdot, t) \in H^{-1}(B)$, with

$$\|\operatorname{div} g(\cdot, t)\|_{H^{-1}(B)} \leq C(n, a, d) \|\nabla u(\cdot, t)\|_{L_2(B)} \quad \text{for a.e } t \in (-1, 0).$$

Therefore, there exists a unique $U(\cdot, t) \in \dot{L}_2^1(B)$ which solves (3) and such that

$$\|\nabla U(\cdot, t)\|_{L_2(B)} \leq C(n, a, d) \|\nabla u(\cdot, t)\|_{L_2(B)} \quad \text{for a.e } t \in (-1, 0).$$

We also deduce that $\nabla U \in L_2(Q)$ and

$$\|\nabla U\|_{L_2(Q)} \leq C(n, a, d) \|\nabla u\|_{L_2(Q)}.$$

Step 2. Now, we can rewrite (2) as follows

$$\int_Q u \partial_t \phi dz = \int_Q \nabla U \cdot \nabla \phi dz \quad \forall \phi \in C_0^\infty(Q). \quad (4)$$

By a density arguments, we can test (4) with functions $\phi(x, t) = \chi(t)\phi_k(x)$, where $\chi \in C_0^\infty(-1, 0)$ and ϕ_k is an eigenfunction (introduced in the second part of the preliminaries section; here we choose $\Omega = B$). Since $(\phi_k)_{k=1}^\infty$ is a Hilbert basis of $L_2(B)$, we can write u as follows

$$u(x, t) = \sum_{k=1}^{\infty} d_k(t)\phi_k(x),$$

where $d_k(t) = \int_B u(\cdot, t)\phi_k dx$; we also have

$$U(x, t) = \sum_{k=1}^{\infty} b_k(t)\phi_k(x),$$

where $b_k(t) = \int_B U(\cdot, t)\phi_k dx$. So we have, thanks to Lemma 2, that

$$\|\nabla U\|_{L_2(Q)}^2 = \int_{-1}^0 \sum_{k=1}^{\infty} b_k^2(t)\lambda_k dt \leq C(n, a, d)\|\nabla u\|_{L_2(Q)}^2 < \infty.$$

We have now

$$\begin{aligned} \int_{-1}^0 d_k(t)\chi'(t)dt &= \int_{-1}^0 \chi(t) \int_B \nabla U \cdot \nabla \phi_k dx dt \\ &= \int_{-1}^0 \chi(t) b_k(t) \lambda_k dx dt. \end{aligned}$$

So, $d'_k(t) = -b_k(t)\lambda_k$ and we derive that

$$\partial_t u(x, t) = \sum_{k=1}^{\infty} d'_k(t)\phi_k(x),$$

where the convergence of this sum occurs in the space of distributions; thus we have, for every $w \in C_0^\infty(B)$ and $\chi \in C_0^\infty(-1, 0)$, that

$$\begin{aligned} \int_{-1}^0 \langle \partial_t u(\cdot, t), w \rangle \chi(t) dt &= - \lim_{N \rightarrow \infty} \int_{-1}^0 \sum_{k=1}^N b_k(t) \lambda_k \int_B \phi_k(x) w(x) dx \chi(t) dt \\ &= \lim_{N \rightarrow \infty} \int_{-1}^0 \sum_{k=1}^N b_k(t) \int_B \nabla \phi_k(x) \cdot \nabla w(x) dx \chi(t) dt \\ &\leq \|\nabla w\|_{L_2(B)} \int_{-1}^0 \left(\sum_{k=1}^{\infty} b_k^2(t) \lambda_k \right)^{1/2} |\chi(t)| dt \quad \text{by Lemma 2} \\ &\leq c(n, a, d) \|\nabla u\|_{L_2(Q)} \|\chi\|_{L_2(-1, 0)} \|\nabla w\|_{L_2(B)}, \end{aligned}$$

and the statement follows. \square

Remark 1. Let $\varphi \in C_0^\infty(B)$. Then, we readily deduce from the above proposition that

$$\partial_t(u\varphi) \in L_2(-1, 0; H^{-1}(B)).$$

Since obviously $u\varphi \in L_2(-1, 0; \mathring{L}_2^1(B))$, we conclude that

$$u\varphi \in C([-1, 0]; L_2(B)).$$

Proof of Theorem 3.1. Step 1. Consider the following auxiliary equation

$$\begin{cases} \partial_t w - \operatorname{div}(A \nabla w) = F - \operatorname{div} G & \text{in } B \times (-1, 0), \\ w|_{\partial' Q} = 0, \end{cases} \quad (5)$$

where $F = u \partial_t \varphi - (A \nabla u) \cdot \nabla \varphi$ and $G = u(A \nabla \varphi)$. We have that the distribution $F - \operatorname{div} G$ belongs to $L_2(-1, 0; H^{-1}(B))$. This is a direct consequence of the fact that $u\varphi$ is distributional solution of (5) together with Remark 1 and Step 1 of the proof of Proposition 4.1. But, for our purpose we are interested, more precisely, in the bounds of the terms which appear in the definition of $F - \operatorname{div} G$ and which belong to $L_2(-1, 0; H^{-1}(B))$. Therefore let us consider a function $w \in C_0^\infty(B)$; then for a.e $t \in (-1, 0)$

$$\begin{aligned} \int_B F w dx + \int_B G \cdot \nabla w dx &= \int_B u \partial_t \varphi w dx - \int_B (a \nabla u) \cdot \nabla \varphi w dx - \int_B (d \nabla u) \cdot \nabla \varphi w dx \\ &\quad + \int_B (a \nabla \varphi) \cdot \nabla w dx + \int_B (d \nabla \varphi) \cdot \nabla w dx \\ &= J_1 + J_2 + J_3, \end{aligned}$$

where

$$\begin{aligned} J_1 &:= \int_B u \partial_t \varphi w, \\ J_2 &:= - \int_B (a \nabla u) \cdot \nabla \varphi w dx + \int_B (a \nabla \varphi) \cdot \nabla w dx, \\ J_3 &:= - \int_B (d \nabla u) \cdot \nabla \varphi w dx + \int_B (d \nabla \varphi) \cdot \nabla w dx. \end{aligned}$$

We rewrite the term J_3 as follows

$$\begin{aligned} J_3 &= - \int_B d \nabla(uw) \cdot \nabla \varphi dx + \int_B (d \nabla w) \cdot \nabla \varphi dx + \int_B (d \nabla \varphi) \cdot \nabla w dx \\ &= - \int_B d \nabla(uw) \cdot \nabla \varphi dx \quad (\text{by the skew symmetry of } d). \end{aligned}$$

But again, thanks to the skew symmetry of d , we have that

$$J_3 = - \frac{1}{2} \sum_{i,j=1}^n \int_B b_{i,j} [(uw)_{,j} \phi_{,i} - \phi_{,j} (uw)_{,i}] dx$$

and making the same computations as for A_2 in Step 1 of the proof of Proposition 4.1, with the only difference being that we keep p arbitrary (instead of choosing $p = 2$ as in the proof of Proposition 4.1), we obtain

$$|J_3| \leq C(n, a, d) \|\nabla(uw)\|_{L_p(B)} \|\varphi\|_{L_{p/(p-1)}(B)},$$

(for $1 < p < \infty$ to be suitably chosen in function of n) which implies that

$$|J_3| \leq C(n, a, d, \varphi) (\|w \nabla u\|_{L_p(B)} + \|u \nabla w\|_{L_p(B)}).$$

If $\mathbf{n} \geq \mathbf{3}$, we steadily have for

$$1 < p < \min(2, \frac{n}{n-1}),$$

that (here $2^* = \frac{2n}{n-2}$)

$$\begin{aligned} \|w \nabla u\|_{L_p(B)} + \|u \nabla w\|_{L_p(B)} &\leq c(n) [\|w\|_{L_{2^*}(B)} \|\nabla u\|_{L_2(B)} + \|u\|_{L_{2^*}(B)} \|\nabla w\|_{L_2(B)}] \\ &\leq c(n) (\|\nabla u\|_{L_2(B)} + \|u\|_{L_2(B)}) \|\nabla w\|_{L_2(B)}, \end{aligned}$$

where Sobolev embedding and Poincaré's inequality are used in the last estimate.

The case $\mathbf{n} = \mathbf{2}$ is a straightforward adaptation of the previous (since $H^1(B)$ embeds continuously in every $L_s(B)$, $1 \leq s < \infty$), whereas for the case $\mathbf{n} = \mathbf{1}$, we take $p = 2$, use the fact that $H^1(B)$ is continuously embedded in $L_\infty(B)$ and Poincaré's inequality for the term in w .

Next, we have the following easy bound for the terms J_1 and J_2 :

$$|J_1 + J_2| \leq C(n, \varphi) (\|\nabla u\|_{L_2(B)} + \|u\|_{L_2(B)}) \|\nabla w\|_{L_2(B)}.$$

So in conclusion, we have, for a.e $t \in (-1, 0)$ that

$$\|F(\cdot, t) - \operatorname{div} G(\cdot, t)\|_{H^{-1}(B)} \leq C(n, a, d, \varphi) (\|\nabla u(\cdot, t)\|_{L_2(B)} + \|u(\cdot, t)\|_{L_2(B)}),$$

and we get a fortiori

$$\|F - \operatorname{div} G\|_{L_2(-1, 0; H^{-1}(B))} \leq C(n, a, d, \varphi) (\|\nabla u\|_{L_2(Q)} + \|u\|_{L_{2, \infty}(Q)}) \quad (6)$$

Step 2. Let us now tackle the question of well-posedness of (5). Consider the time-indexed family of bilinear forms

$$\delta_t(w, v) := \int_B (A \nabla w) \cdot \nabla v dx.$$

Let us first notice that the map $t \in (-1, 0) \mapsto \delta_t(w, v)$ is measurable for every $w, v \in \dot{L}_2^1(B)$. Furthermore, we have by similar computations as those made in Step 1 in the proof of Proposition 4.1, that there exists a constant $C = C(n, a, d) > 0$ independent of t such that

$$|\delta_t(w, v)| \leq C \|\nabla w\|_{L_2(B)} \|\nabla v\|_{L_2(B)},$$

for all $w, v \in \dot{L}_2^1(B)$ i.e δ_t is a bounded bilinear operator on $\dot{L}_2^1(B)$. We have, additionally, the following coercivity estimate

$$\begin{aligned} \delta_t(w, w) &= \int_B (A \nabla w) \cdot \nabla w dx \\ &= \int_B (a \nabla w) \cdot \nabla w dx \quad (\text{by the skew symmetry of } d), \\ &\geq \nu \int_B |\nabla w|^2 dx. \end{aligned}$$

In view of these previous estimates and the regularity proved for the right-hand side of (5) and considering the evolution triple $\dot{L}_2^1(B) \subset L_2(B) \subset H^{-1}$, we have by applying J-L. Lions abstract theorem for well-posedness of evolution equations (see e.g., [4], Theorem 4.1, Chapter 3, section 4) that there exists a unique solution

$$w \in C([-1, 0]; L_2(B)) \cap L_2(-1, 0; \dot{L}_2^1(B)),$$

with

$$\partial_t w \in L_2(-1, 0; H^{-1})$$

such that

$$\int_Q \partial_t w v dz + \int_Q (A \nabla w) \cdot \nabla v dz = \int_Q (F - \operatorname{div} G) v dz \quad (7)$$

for any $v \in C_0^\infty(Q)$. Let us notice that, from Remark 1, the fact that $u\varphi$ is a distributional solution of (5) and by the above uniqueness result:

$$w = u\varphi.$$

On another hand, by the regularity obtained for w , we can extend identity (7) to functions v in $L_2(-1, 0; \dot{L}_2^1(B))$ and therefore, test (7) with w itself. Thus, we get

$$\int_Q (A \nabla w) \cdot \nabla w dz = \int_Q (F - \operatorname{div} G) w dz. \quad (8)$$

Denote by L , the left-hand side of the above identity. By the skew symmetry of d , we obtain that

$$L = \int_Q (a \nabla w) \cdot \nabla w dz;$$

therefore coming back to u , we get

$$\begin{aligned} L &= \int_Q a(\varphi \nabla u + u \nabla \varphi) \cdot (\varphi \nabla u + u \nabla \varphi) dz \\ &= \int_Q \varphi^2 (a \nabla u) \cdot \nabla u dz + 2 \int_Q u \varphi (a \nabla u) \cdot \nabla \varphi dz + \int_Q u^2 (a \nabla \varphi) \cdot \nabla \varphi. \end{aligned}$$

Now, denote by R the right hand side of (8); we easily obtain that

$$\begin{aligned} R &= \int_Q u \partial_t \varphi w dz - \int_Q (a \nabla u) \cdot \nabla \varphi w dz + \int_Q (a \nabla \varphi) \cdot \nabla w u - \int_Q d \nabla (u w) \cdot \nabla \varphi dz \\ &= \int_Q u^2 \varphi \partial_t \varphi dz - \int_Q u \varphi (a \nabla u) \cdot \nabla \varphi dz + \int_Q \varphi u (a \nabla u) \cdot \nabla \varphi dz + \int_Q u^2 (a \nabla \varphi) \cdot \nabla \varphi dz \\ &\quad - 2 \int_Q u \varphi (d \nabla u) \cdot \nabla \varphi dz. \end{aligned}$$

Therefore, (8) implies that

$$\int_Q \varphi^2 (a \nabla u) \cdot \nabla u dz + 2 \int_Q u \varphi (a \nabla u) \cdot \nabla \varphi dz = \int_Q u^2 \varphi \partial_t \varphi dz - 2 \int_Q u \varphi (d \nabla u) \cdot \nabla \varphi dz, \quad (9)$$

for all $\varphi \in C_0^\infty(Q)$. Let us notice that all the integrals in the above identity are finite, especially the last one of the right hand side of (9). To see this we rewrite

$$\int_Q u\varphi(d\nabla u).\nabla\varphi dz = \int_Q d\nabla(u^2/2).\nabla(\varphi^2/2)dz$$

and use the same method as in the estimation of J_3 in the previous step.

Step 3. Now, we choose $\varphi(x, t) = \chi_\epsilon(t)\phi(x, t)$, where $\phi \in C_0^\infty(B \times (-1, 1))$ and $\chi_\epsilon(t) = 1$ if $t \leq t_0 - \epsilon$, $\chi_\epsilon(t) = (t_0 + \epsilon - t)/(2\epsilon)$, if $t_0 - \epsilon < t < t_0 + \epsilon$, and $\chi_\epsilon(t) = 0$ when $t \geq t_0 + \epsilon$, with $t_0 \in (-1, 0)$. Therefore passing to the limit $\epsilon \rightarrow 0$ in (9), we have that Theorem 3.1 is proved.

□

Acknowledgement This work was supported by the Engineering and Physical Sciences Research Council [EP/L015811/1]. The author would like to thank Siran Li and Ghazlane Yahiaoui for their careful reading and insightful discussion.

References

- [1] G. Seregin, L. Silvestre, V. Šverák and A. Zlatoš (2012). On divergence-free drifts. *Journal of Differential Equations*. **252**, 505-540.
- [2] P.W Jones (1980). Extension theorems for BMO. *Indiana Univ. Math. J.*, **29**, no 1, 41-66.
- [3] E. Stein (1970). *Singular Integrals and Differentiability Properties of Functions*. Princeton, NJ: Princeton University Press.
- [4] J-L. Lions (1965). *Problèmes aux limites dans les équations aux dérivées partielles*. Presses de l'Université de Montréal.
- [5] R. Coifman, P-L. Lions, Y. Meyer and S. Semmes. Compensated Compactness and Hardy Spaces. *J. Maths. Pures Appl.*, **72**, 1993, 247-286.