

Spin-orbit precession along eccentric orbits for extreme mass ratio black hole binaries and its effective-one-body transcription

Chris Kavanagh,¹ Donato Bini,² Thibault Damour,¹ Seth Hopper,³ Adrian C. Ottewill,⁴ and Barry Wardell^{5,6}

¹*Institut des Hautes Études Scientifiques, 91440 Bures-sur-Yvette, France.*

²*Istituto per le Applicazioni del Calcolo “M. Picone”, CNR, 00185 Rome, Italy.*

³*CENTRA, Departamento de Física, Instituto Superior Técnico - IST, Universidade de Lisboa, 1049 Lisboa, Portugal*

⁴*School of Mathematics and Statistics and Institute for Discovery, University College Dublin, Belfield, Dublin 4, Ireland.*

⁵*School of Mathematics and Statistics, University College Dublin, Belfield, Dublin 4, Ireland.*

⁶*Department of Astronomy, Cornell University, Ithaca, NY 14853, USA*

In this work we present an analytical gravitational self-force calculation of the spin-orbit precession along an eccentric orbit around a Schwarzschild black hole, following closely the recent prescription of Akcay, Dempsey, and Dolan. We then transcribe this quantity within the Effective-One-Body (EOB) formalism, thereby determining several new, linear-in-mass-ratio, contributions in the post-Newtonian expansion of the spin-orbit couplings entering the EOB Hamiltonian. Namely, we determine the second gyro-gravitomagnetic ratio $g_{S_*}(r, p_r, p_\phi)$ up to order p_r^2/r^4 included.

I. INTRODUCTION

The development of accurate waveform templates for compact binaries is essential for the science of gravitational waves. For example, the extraction of physical information from the first LIGO detections [1, 2] has made a key use of a bank of $\sim 200,000$ semianalytical templates [3, 4], describing the inspiral, merger and ringdown of two comparable-mass black holes, that were developed within the Effective-One-Body (EOB) formalism [5–8]. For future detectors such as LISA to reach their full potential one needs to describe systems with mass ratios varying from 1:1 to $\sim 1 : 10^6$, evolving over long inspirals into plunge, merger and ringdown phases. When attempting to model such orbital evolutions one typically relies on several approximate ways of solving Einstein’s equations, valid in different asymptotic regimes: post-Newtonian (PN) theory in the slow-motion, weak-field regime; post-Minkowskian (PM) theory in the weak-field regime; gravitational self-force (SF) theory for small mass ratios; numerical relativity (NR) for strong-field comparable mass binaries; and EOB theory for analytically interpolating between various regimes.

Recent years have witnessed a fruitful crossbreeding between these various methods. Notably, the EOB formalism has provided, through its natural theoretical flexibility, a common ground for incorporating the results of other approaches. Examples of recent works contributing to the crossbreeding between EOB theory and other approximation methods are: $\text{EOB} \cup \text{PN}$ [9–12]; $\text{EOB} \cup \text{PM}$ [13]; $\text{EOB} \cup \text{SF}$ [14–17]; and $\text{EOB} \cup \text{NR}$ [18–22].

The primary focus of this paper is on the third of these strategies: the extraction of physical information from SF results, and their EOB transcription. The SF approach — in which Einstein’s equations are solved perturbatively with the mass-ratio as small parameter — is ideally suited to describing the motion of compact binary systems with a large discrepancy in the masses. An im-

portant theme (initiated in Refs. [23, 24]) within the SF community over the past ten years has been the extraction of physically meaningful quantities through the computation of gauge-invariant SF quantities. These quantities are typically defined within the conservative sector, with dissipative effects of the self-force ignored or turned off. See Refs. [25, 26] for the first corresponding EOB transcriptions of gauge-invariant SF quantities. In the literature there now exists a wide array of gauge invariant quantities, each with varying dependencies on the perturbed metric and its derivatives. The utility of these include: insights into the physical effects of the self-force (see, e.g., [27]); comparisons within SF theory between calculations in differing gauges (e.g. [28]); comparisons with independent PN calculations (e.g. [29–31]) and with NR codes (e.g. [32]); and the extraction of high-PN-order contributions to the potentials of EOB theory (e.g. [33–35]).

In a recent paper, Akcay, Dempsey and Dolan [36] presented a methodology for calculating the gauge invariant self-force correction to the spin-orbit precession of a spinning compact body along an eccentric orbit in Schwarzschild spacetime, as well as a numerical calculation of the precession using a Lorenz gauge code. Their presentation is the first example of a gauge invariant quantity for an eccentric orbit binary which depends on derivatives of the metric perturbation, and gives access for the first time to spin-orbit effects along an *eccentric* orbit.

The first aim of the present work is to complement the (mostly numerical) results of Ref. [36] by presenting an *analytical* calculation of their invariant as a PN expansion within SF theory. To do so we rely on a low eccentricity assumption in a manner following closely that of Ref. [15]. By contrast, both, with Ref. [15] (which used a Regge-Wheeler gauge), and with Ref. [36] (which used a Lorenz gauge), we will work in a so-called radiation gauge, which unlike the Regge-Wheeler gauge is readily

extendible to a Kerr spacetime. This will provide an independent check of the gauge invariance of the spin-precession quantity defined in [36].

The second aim of the present work is to explicitly derive the relationship between the spin precession invariant along eccentric orbits, and the various potentials parametrizing spin-orbit effects within the EOB formalism. [For the corresponding relationship in the simpler case of circular orbits see Ref. [34].] We shall then use this relationship to show how the knowledge of the $O(e^2)$ [respectively, $O(e^4)$] eccentric corrections to the spin precession translates into new information about the terms quadratic (resp., quartic) in the radial momentum in the spin-orbit potentials of the EOB Hamiltonian.

The organisation of this paper is as follows. In Sec. II we review the formalism for calculating the eccentric spin-precession, discuss eccentric geodesics in Schwarzschild spacetime and their perturbation and review the radiation gauge approach to reconstructing the perturbed metric. In Sec. III we describe the post-Newtonian approach we take to calculating the retarded metric perturbation, the self-force, give our regularization and metric completion and finally the eccentric spin precession. Then in Sec. IV, after briefly recalling the EOB parametrization of spin-orbit effects, we show how to transcribe the spin-precession invariant $\Delta\psi(p, e)$ into a knowledge of the $O(\nu)$ contribution to the second gyrogravitomagnetic ratio $g_{S*}(u, p_r, p_\phi; \nu)$.

II. GEODETIC SPIN-PRECESSION

A. Overview

We start by giving a brief summary of the prescription of Akcay, Dempsey and Dolan to calculate the gravitational self-force (SF) correction to the spin precession. For a more detailed description we refer the reader to [36].

We wish to calculate the amount of precession angle a test spin vector accumulates over one radial period compared to the accumulated azimuthal angle. This precession is conveniently measured by the quantity (in units where $G = c = 1$)

$$\psi(m_2\Omega_r, m_2\Omega_\varphi; q) = \frac{\Phi - \Psi}{\Phi}. \quad (2.1)$$

Here, we consider a binary system with masses m_1 and m_2 (with $q \equiv \frac{m_1}{m_2} \ll 1$); Φ is the accumulated azimuthal phase from periapsis to periapsis (i.e. during a radial period), and Ψ the corresponding accumulated phase of the spin vector (both being computed along an eccentric orbit perturbed by SF effects), and, finally; Ω_r , and Ω_φ are, respectively the radial and (mean) azimuthal angular frequencies. The question is then how to define Ψ .

The spin vector s^a is parallelly transported along an equatorial geodesic of the (regularized [37]) $O(q)$ -perturbed spacetime with four velocity u^a and proper

time τ : $\frac{Ds_a}{d\tau} = 0$. Projecting this equation onto a particular (polar-type) reference frame e_α^a , its spatial components satisfy

$$\frac{ds}{d\tau} = \boldsymbol{\omega} \times \mathbf{s} \quad (2.2)$$

where

$$(\mathbf{s})_i = e_i^a s_a, \quad (\boldsymbol{\omega})_i = -\frac{1}{2}\epsilon_{ijk}\omega^{jk}, \quad \omega_{ij} = -g_{ab}e_i^a \frac{De_j^b}{d\tau}. \quad (2.3)$$

The spin precession is then entirely defined by the relative motion of the tetrad throughout the orbit. Choosing the basis so that only the (θ -like) $(\boldsymbol{\omega})_2$ component is non-zero, one finds that the spin vector accumulates an angle Ψ over one radial period (periapsis to periapsis)

$$\Psi(m_2\Omega_r, m_2\Omega_\varphi; q) = \oint \omega_{13}(\tau) d\tau. \quad (2.4)$$

The aim is then to explicitly calculate the $O(q)$, SF contributions to ψ , Eq. (2.1), and to Ψ , Eq.(2.4), i.e. the quantities

$$\Delta\Psi = \Psi(\Omega_r, \Omega_\varphi, q) - \Psi(\Omega_r, \Omega_\varphi, 0), \quad (2.5)$$

and

$$\Delta\psi = \psi(\Omega_r, \Omega_\varphi, q) - \psi(\Omega_r, \Omega_\varphi, 0) = -\frac{\Delta\Psi}{\Phi}, \quad (2.6)$$

where we recall that $q = m_1/m_2$ denotes the small mass ratio, and where we used the fact that

$$\Phi \equiv \Omega_\varphi \mathcal{T} \equiv 2\pi \frac{\Omega_\varphi}{\Omega_r}, \quad (2.7)$$

is the same on the perturbed ($q \neq 0$) and background ($q = 0$) orbits. Here, differently from Ref. [36], we use the letter $\mathcal{T} = 2\pi/\Omega_r$ to denote the coordinate-time radial period.

In practice, we shall work below with an intermediate $O(q)$ variation (denoted δ) which does not keep fixed the values of the two frequencies $(\Omega_r, \Omega_\varphi)$. We can then recover the correct value of $\Delta\psi$, Eq. (2.6), by first ‘subtracting’ the induced frequency shifts

$$\Delta\Psi = \delta\Psi - \frac{\partial\Psi}{\partial\Omega_r}\delta\Omega_r - \frac{\partial\Psi}{\partial\Omega_\varphi}\delta\Omega_\varphi, \quad (2.8)$$

and then computing

$$\Delta\psi = -\frac{\Delta\Psi}{\Phi}. \quad (2.9)$$

With this broad outline in mind, the next few sections will focus on the explicit details of this calculation when using post-Newtonian (PN) expansions.

B. Motion on the background ($q = 0$): Schwarzschild spacetime

In the usual Schwarzschild coordinates, the unperturbed ($q = 0$) metric takes the form

$$ds^2 = -f dt^2 + f^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2) \quad (2.10)$$

where $f \equiv (1 - 2m_2/r)$.

1. Background equatorial geodesics

We start by recalling some of the defining equations for a particle undergoing bound equatorial motion in a Schwarzschild spacetime. Here and henceforth we use a subscript p to denote evaluation at the position of the particle. Such motion is parameterised by two constants of motion, the specific energy and angular momentum (\mathcal{E}, \mathcal{L}) respectively. The tangent four velocity is then given by

$$u_p^\mu = \left(\frac{\mathcal{E}}{f}, u_p^r, 0, \frac{\mathcal{L}}{r_p^2} \right). \quad (2.11)$$

Here we have made the standard restriction to equatorial motion setting $\theta_p = \frac{\pi}{2}$, $u_p^\theta = 0$. The radial motion can be parametrised using Darwin's relativistic anomaly χ [38]

$$r_p(\chi) = \frac{pM}{1 + e \cos \chi} \quad (2.12)$$

where $\chi = 0$ corresponds to periapsis, p is the semilatus rectum, and e the eccentricity. Using these parameters,

$$\mathcal{E}^2 = \frac{(p-2)^2 - 4e^2}{p(p-3-e^2)}, \quad \mathcal{L}^2 = \frac{p^2 M^2}{p-3-e^2} \quad (2.13)$$

and

$$\frac{dt_p}{d\chi} = \left[\frac{(p-2)^2 - 4e^2}{p-6-2e \cos \chi} \right]^{1/2} \times \frac{p^2 M}{(p-2-2e \cos \chi)(1+e \cos \chi)^2}, \quad (2.14)$$

$$\frac{d\varphi_p}{d\chi} = \left[\frac{p}{p-6-2e \cos \chi} \right]^{1/2}, \quad (2.15)$$

$$\frac{d\tau_p}{d\chi} = \frac{Mp^{3/2}}{(1+e \cos \chi)^2} \left[\frac{p-3-e^2}{p-6-2e \cos \chi} \right]^{1/2}. \quad (2.16)$$

With these one can compute the unperturbed radial period as the coordinate time taken between successive periapses

$$\mathcal{T} \equiv \int_0^{2\pi} \left(\frac{dt_p}{d\chi} \right) d\chi. \quad (2.17)$$

The characteristic orbital frequencies are then

$$\Omega_r = \frac{2\pi}{\mathcal{T}}, \quad \Omega_\varphi = \frac{\Phi}{\mathcal{T}}, \quad (2.18)$$

with $\Phi = \int_0^{2\pi} \left(\frac{d\varphi_p}{d\chi} \right) d\chi$.

2. Background reference tetrad

As recalled above, in order to define the precession of the spin vector, one needs to choose a reference frame. Following [36], a suitable polar-type tetrad is that given explicitly (when $q = 0$) by Marck [39]:

$$e_0^a = u^a = \left(\frac{\mathcal{E}}{f}, u^r, 0, \frac{\mathcal{L}}{r^2} \right), \quad (2.19)$$

$$e_1^a = \frac{1}{f\sqrt{1+\mathcal{L}^2/r^2}} (u^r, f\mathcal{E}, 0, 0), \quad (2.20)$$

$$e_2^a = (0, 0, 1/r, 0), \quad (2.21)$$

$$e_3^a = \frac{1}{r\sqrt{1+\mathcal{L}^2/r^2}} \left(\frac{\mathcal{E}\mathcal{L}}{f}, \mathcal{L}u^r, 0, 1 + \frac{\mathcal{L}^2}{r^2} \right). \quad (2.22)$$

Using (2.3) the key frequency determining the precession function is

$$\omega_{13} = \frac{\mathcal{E}\mathcal{L}}{r^2 + \mathcal{L}^2}. \quad (2.23)$$

It is straightforward then to calculate the background spin precession using (2.12), (2.13) with (2.1), (2.4).

C. Motion and spin-precession in the perturbed spacetime ($q \neq 0$)

Ref. [36], generalizing previous results by Barack and Sago [40], has derived an explicit integral expression for the SF contribution $\delta\Psi$ to Ψ (using a specific variation δ which does not fix the frequencies). Their final result reads

$$\delta\Psi = \int \left(\frac{\delta\dot{\Psi}}{\dot{\Psi}} - \frac{\delta u^r}{u^r} \right) \dot{\Psi} \frac{d\tau}{d\chi} d\chi. \quad (2.24)$$

The term proportional to δu^r appears since the proper time in (2.4) also needs to be varied. The evaluation of this expression requires further definitions. We have introduced $\dot{\Psi} \equiv \omega_{13} = \omega_{[13]}$ to be the integrand of (2.4) in favour of the ω of [36] to avoid overlap with the frequency notation for the frequency domain solutions of the Teukolsky equation. Its variation is given by Akcay *et al* as

$$\delta\dot{\Psi} = \frac{1}{2} \dot{\Psi} h_{00} + \delta\Gamma_{[31]0} + (c_{01}e_1^b + c_{03}e_3^b) e_{a[3} \nabla_b e_{1]}^a \quad (2.25)$$

where we have neglected a total derivative term which vanishes upon integration over the orbit, $\delta\Gamma_{[31]0}$ is a tetrad component of $\delta\Gamma_{\mu\nu\rho} = \frac{1}{2}(h_{\mu\nu,\rho} + h_{\mu\rho,\nu} - h_{\nu\rho,\mu})$ and

$$c_{01} = \frac{1}{f\sqrt{1+\mathcal{L}^2/r^2}} (\mathcal{E}\delta u_{BS}^r - u^r \delta\mathcal{E}_{BS}), \quad (2.26)$$

$$c_{03} = \frac{\delta\mathcal{L}_{BS}}{r\sqrt{1+\mathcal{L}^2/r^2}} \quad (2.27)$$

are terms arising from perturbing the tetrad. In this expression the subscript BS refers to the perturbations in energy, angular momentum and radial velocity defined by Barack and Sago in [40]. They differ from those used by Akcay *et al* by the normalisation of the 4-velocity, because Barack and Sago normalise with respect to the background metric while Akcay *et al* normalise with respect to the perturbed spacetime. This leads to the following relation valid to linear order in the mass-ratio

$$\delta\mathcal{E}_{BS} = \delta\mathcal{E} - \frac{1}{2}\mathcal{E}h_{00}, \quad (2.28)$$

$$\delta u_{BS}^r = \delta u^r - \frac{1}{2}u^r h_{00}, \quad (2.29)$$

$$\delta\mathcal{L}_{BS} = \delta\mathcal{L} - \frac{1}{2}\mathcal{L}h_{00}. \quad (2.30)$$

where $h_{00} \equiv h_{\mu\nu}e_0^\mu e_0^\nu$. The $O(q)$ perturbations $\delta\mathcal{E}_{BS}$, $\delta\mathcal{L}_{BS}$ (which are denoted ΔE and ΔL by Barack and Sago) of the energy and angular momentum $u_t \equiv -\mathcal{E}$, $u_\varphi \equiv \mathcal{L}$ entering the above equations have been shown in Sec II. C of [40] to be determined by the following quadratures

$$\delta\mathcal{E}_{BS}(\chi) = \delta\mathcal{E}_{BS}(0) - \int_0^\chi F_t^{\text{cons}} \frac{d\tau}{d\chi} d\chi, \quad (2.31)$$

$$\delta\mathcal{L}_{BS}(\chi) = \delta\mathcal{L}_{BS}(0) + \int_0^\chi F_\varphi^{\text{cons}} \frac{d\tau}{d\chi} d\chi. \quad (2.32)$$

Here $\delta\mathcal{E}(0)$ and $\delta\mathcal{L}(0)$ are the energy and angular momentum shift at periapsis, given in Eq. (37) and (38) of [40], and F_μ^{cons} is the conservative part of the self-force which we discuss below (see Eq. (3.12)). [By contrast to the Detweiler-Whiting formulation of SF we used in our presentation above, Barack and Sago use the formulation where the perturbed motion satisfies the forced-motion equation $\frac{D_0 u^\mu}{d\tau_0} = F^\mu(\tau_0)$.] Using the normalisation condition of Akcay *et al* the perturbed radial velocity is then calculated from the relation

$$\frac{1}{2}h_{00} - \frac{\mathcal{E}\delta\mathcal{E}}{f_p} + \frac{u^r\delta u^r}{f_p} + \frac{\mathcal{L}\delta\mathcal{L}}{r_p^2} = 0. \quad (2.33)$$

Finally the gauge invariant precession function is calculated using Eqs. (2.8) and (2.9). The frequency shifts calculated by perturbing (2.18) are given in Eq. (75)-(76) of

[40]. We however do not require their α term, which they added to ensure the asymptotic flatness of their metric perturbation, since we will work in an asymptotically flat gauge.

D. Radiation gauge metric perturbation

Our strategy for computing a post-Newtonian expansion of the retarded metric perturbation in many ways follows closely that laid out in [15], in that we will use the method of extended homogeneous solutions to construct a particular solution of a particular partial differential equation whose solutions are related to the metric perturbation. The key difference however is that we shall use the tetrad formalism and radiation gauge to construct the metric, using the CCK procedure, so named after its development by Chrzanowski [41] and Cohen and Kegeles [42, 43]. Specifically this involves building inhomogeneous solutions to the Bardeen-Press-Teukolsky (BPT) equation ($a = 0$ Teukolsky equation), from this a Hertz potential and finally the metric perturbation and all its first derivatives. Since much of the details are covered by a variety of authors e.g [44–48] we will give an abridged overview of the strategy and refer the reader to the given references for details.

1. Bardeen-Press-Teukolsky (BPT) equation

The description of perturbations to a black hole spacetime can be reduced to a single partial differential equation for the tetrad components of the perturbed Weyl tensor $C_{\mu\nu\rho\sigma}$. In particular (essentially) all information is simultaneously contained in the two quantities ψ_0 and ψ_4 defined by:

$$\psi_0 = -C_{\alpha\beta\gamma\delta}l^\alpha m^\beta \bar{l}^\gamma \bar{m}^\delta, \quad (2.34)$$

$$\psi_4 = -C_{\alpha\beta\gamma\delta}n^\alpha \bar{m}^\beta n^\gamma \bar{m}^\delta, \quad (2.35)$$

where l^μ , n^μ , m^μ and \bar{m}^μ are the Kinnersley tetrad legs given in Appendix A. The dynamics of the perturbed Weyl scalars on a Schwarzschild background are described by the BPT equation:

$$\begin{aligned} \frac{r^4}{\Delta} \frac{\partial^2 \psi}{\partial t^2} - \frac{1}{\sin^2 \theta} \frac{\partial^2 \psi}{\partial \varphi^2} - \Delta^{-s} \frac{\partial}{\partial r} \left(\Delta^{s+1} \frac{\partial \psi}{\partial r} \right) - \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \psi}{\partial \theta} \right) - 2is \frac{\cos \theta}{\sin^2 \theta} \frac{\partial \psi}{\partial \varphi} - 2s \left[\frac{m_2 r^2}{\Delta} - r \right] \frac{\partial \psi}{\partial t} \\ + (s^2 \cot^2 \theta - s) \psi = 4\pi r^2 T. \end{aligned} \quad (2.36)$$

with

$$\Delta \equiv r^2 - 2m_2 r. \quad (2.37)$$

Here, for $s = +2$

$$\psi = \psi_0, \quad T = 2T_0 \quad (2.38)$$

and for $s = -2$

$$\psi = \varrho^{-4}\psi_4, \quad T = 2\varrho^{-4}T_4, \quad (2.39)$$

while the source terms are

$$\begin{aligned} T_0 = & (\delta + \bar{\omega} - \bar{\alpha} - 3\beta - 4\tau) \times \\ & [(\mathbf{D} - 2\epsilon - 2\bar{\varrho})T_{13} - (\delta + \bar{\omega} - 2\bar{\alpha} - 2\beta)T_{11}] \\ & + (\mathbf{D} - 3\epsilon + \bar{\epsilon} - 4\varrho - \bar{\varrho}) \times \\ & [(\delta + 2\bar{\omega} - 2\beta)T_{13} - (\mathbf{D} - 2\epsilon + 2\bar{\epsilon} - 2\bar{\varrho})T_{33}], \end{aligned} \quad (2.40)$$

$$\begin{aligned} T_4 = & (\Delta + 3\gamma - \bar{\gamma} + 4\mu + \bar{\mu}) \times \\ & [(\bar{\delta} - 2\bar{\tau} + 2\alpha)T_{24} - (\Delta + 2\gamma - 2\bar{\gamma} + \bar{\mu})T_{44}] \\ & + (\bar{\delta} - \bar{\tau} + \bar{\beta} + 3\alpha + 4\varpi) \times \\ & [(\Delta + 2\gamma + 2\bar{\mu})T_{24} - (\bar{\delta} - \bar{\tau} + 2\bar{\beta} + 2\alpha)T_{22}], \end{aligned} \quad (2.41)$$

where $\mathbf{D} = l^\mu \partial_\mu$, $\Delta = n^\mu \partial_\mu$ and $\delta = m^\mu \partial_\mu$. Here T_{ij} are the Kinnersley-tetrad projections of the point particle source. Now and henceforth we will focus on the $s = 2$ solutions for ψ_0 (a similar procedure could be followed with ψ_4 since it contains the same information as ψ_0).

This equation is fully separable by means of a Fourier transform and projection over spin-weighted spherical harmonics. Due to the double-periodicity of the eccentric orbits the Fourier transform reduces to a Fourier series labelled by the discrete frequencies $\omega = m\Omega_\varphi + n\Omega_r$.

$$\psi_0(t, r, \theta, \varphi) = \sum_{\ell m} \psi_0^{\ell m}(t, r) {}_2Y_{\ell m}(\theta, \varphi), \quad (2.42)$$

$$\psi_0^{\ell m}(t, r) = \sum_{n=-\infty}^{\infty} e^{-i\omega t} {}_2R_{\ell m \omega}(r). \quad (2.43)$$

The radial functions ${}_sR_{\ell m \omega}(r)$ satisfy

$$\begin{aligned} \left[\Delta^{-s} \frac{d}{dr} \left(\Delta^{s+1} \frac{d}{dr} \right) + \frac{r^4 \omega^2 - 2is(r - m_2)r^2 \omega}{\Delta} \right. \\ \left. + 4is\omega r - {}_s\lambda_{\ell m} \right] {}_sR_{\ell m \omega}(r) = {}_sT_{\ell m \omega}, \end{aligned} \quad (2.44)$$

where ${}_s\lambda_{\ell m} = \ell(\ell + 1) - s(s + 1)$. Assuming a pair of homogenous solutions ${}_s\hat{R}_{\ell m \omega}^+$, ${}_s\hat{R}_{\ell m \omega}^-$ to the above, the corresponding inhomogeneous solution is written

$${}_2R_{\ell m \omega}(r) = c^+ {}_2\hat{R}_{\ell m \omega}^-(r) + c^- {}_2\hat{R}_{\ell m \omega}^+(r) \quad (2.45)$$

where

$$c^+ = \frac{1}{W_{\ell mn}} \int_r^{r_p} \Delta^2 {}_2\hat{R}_{\ell m \omega}^+(r) T_{\ell m \omega} dr, \quad (2.46)$$

$$c^- = \frac{1}{W_{\ell mn}} \int_{r_p}^r \Delta^2 {}_2\hat{R}_{\ell m \omega}^-(r) T_{\ell m \omega} dr \quad (2.47)$$

and $W_{\ell mn} = \Delta^{s+1} \left({}_2\hat{R}_{\ell m \omega}^+ {}_2\hat{R}_{\ell m \omega}'^- - {}_2\hat{R}_{\ell m \omega}^- {}_2\hat{R}_{\ell m \omega}'^+ \right)$. The source term here is

$$T_{\ell m \omega} = \frac{1}{\mathcal{T}} \int_0^{2\pi} \int_0^{2\pi} \int_0^\pi T_0 e^{i\omega t} {}_2Y_{\ell m}^*(\theta, \varphi) \sin \theta \frac{dt}{d\chi} d\theta d\varphi d\chi. \quad (2.48)$$

2. Hertz potential and the retarded metric perturbation

As is standard we reconstruct the metric perturbation by means of an auxiliary function known as the Hertz potential Ψ_H , as in for example [45]. In the outgoing radiation gauge we work in, Ψ_H satisfies the spin-2 BPT equation, as well as the angular equation

$$\psi_0 = \frac{1}{8} (\mathcal{L}^4 \bar{\Psi}_H + 12m_2 \partial_t \Psi_H). \quad (2.49)$$

Here $\mathcal{L}^4 = \mathcal{L}_1 \mathcal{L}_0 \mathcal{L}_{-1} \mathcal{L}_{-2}$ and $\mathcal{L}_s = -(\partial_\theta - s \cot \theta + i \csc \theta \partial_\varphi)$. Spectrally decomposing the Hertz potential as

$$\Psi_H = \sum_{\ell mn} e^{-i\omega t} \Psi_{\ell m \omega} {}_2Y_{\ell m}(\theta, \varphi) \quad (2.50)$$

Eq. (2.49) can be algebraically inverted to give

$$\Psi_{\ell m \omega} = 8 \frac{(-1)^m D {}_2\bar{R}_{\ell, -m, -\omega} + 12im_2 \omega {}_2R_{\ell m \omega}}{D^2 + 144M^2 \omega^2} \quad (2.51)$$

where $D = \ell(\ell + 1)(\ell - 1)(\ell + 2)$. The metric is then obtained from Ψ_H by applying a set of differential operators:

$$\begin{aligned} h_{\alpha\beta} = & -\varrho^{-4} \{ n_\alpha n_\beta (\bar{\delta} - 3\alpha - \bar{\beta} + 5\varpi)(\bar{\delta} - 4\alpha + \varpi) \\ & + \bar{m}_\alpha \bar{m}_\beta (\Delta + 5\mu - 3\gamma - \bar{\gamma})(\Delta + \mu - 4\gamma) \\ & - n_{(\alpha} n_{\beta)} [(\bar{\delta} - 3\alpha + \bar{\beta} + 5\varpi + \bar{\tau})(\Delta + \mu - 4\gamma) \\ & + (\Delta + 5\mu - \bar{\mu} - 3\gamma - \bar{\gamma})(\bar{\delta} - 4\alpha + \varpi)] \} \Psi_H + \text{c.c.}, \end{aligned} \quad (2.52)$$

where c.c. denotes complex conjugation. The first derivatives of the metric which appear in the formula for the spin precession can then be written in terms of three derivatives of the inhomogeneous solution (2.45).

Ultimately the decomposition in spin-weighted spherical harmonics of ψ_0 , Eq. (2.42), generates a corresponding decomposition of $h_{\alpha\beta}$ in *tensorial* spherical harmonics, with ℓ (together with the parity) labelling each irreducible representation of the rotation group. In turn, this generates a corresponding decomposition of the spin-precession ψ . We shall often refer to the irreducible pieces of these decompositions as “ ℓ -modes”.

III. POST-NEWTONIAN APPROACH

We now wish to proceed with the calculation laid out in the previous sections analytically, using a PN assumption that the orbital separation between the two bodies is large, i.e. $p \gg 1$. For simplicity, we will additionally assume small eccentricities ($e \ll 1$), and so our results will appear as double expansions in e and $1/p$.

In practice, to achieve the required accuracy of the spin precession invariant we will work to 7 orders in $1/p$ and 6 orders in e . This will yield the result to 5 PN orders and

accurate to order e^2 (since the invariant is defined as the ratio of two angles one loses two powers of e throughout the calculation, i.e. e^4 accuracy is needed for e^2 results). Our extra orders are kept to reduce potential systematic errors.

A. Background orbit

The various background orbital elements of IIB can be easily calculated in the PN regime, see e.g. [15]. For example the orbital period, \mathcal{T} , can be calculated by expanding (2.14)

$$\begin{aligned} \frac{dt_p}{d\chi} = p^{3/2} & \left[(1 - 2e \cos \chi + 3e^2 \cos^2 \chi + \mathcal{O}(e^3)) \right. \\ & \left. + 3(1 - e \cos \chi + e^2 \cos^2 \chi + \mathcal{O}(e^3)) \frac{1}{p} + \mathcal{O}(p^{-2}) \right] \end{aligned} \quad (3.1)$$

and integrating order by order. This can be extended with ease to the desired orders in $1/p$ and e . Repeating this for $\frac{d\varphi_p}{d\chi}$, we can obtain the two orbital frequencies

$$m_2 \Omega_r = \left(\frac{1 - e^2}{p} \right)^{3/2} \left[1 - 3 \frac{1 - e^2}{p} + \mathcal{O}(p^{-2}) \right] \quad (3.2)$$

$$m_2 \Omega_\varphi = \left(\frac{1 - e^2}{p} \right)^{3/2} \left[1 + 3 \frac{e^2}{p} + \mathcal{O}(p^{-2}) \right] \quad (3.3)$$

These are equivalent in the Newtonian limit.

B. PN-expanded BPT equation

Before we compute the perturbed orbital elements we need the self-force and thus the metric perturbation and its derivatives. For our set of homogeneous solutions we use exactly those described in Ref. [47], with the rotation parameter a limiting to zero. That is, dropping the $s = 2$ subscript and translating notation,

$$\hat{R}_{\ell m \omega}^{+/-} = R_{\ell m \omega}^{\text{up/in}}(a \rightarrow 0). \quad (3.4)$$

Note that while the computation of [47] is aimed at circular orbits, the homogeneous solutions obtained therein are derived with only the assumptions that the orbital radius is large and that the frequency scales as $\omega \sim r^{-3/2}$. Both of the assumptions are satisfied in our current study, as can be seen explicitly by Eq. (3.3).

We would also like to emphasise the nature of these solutions as a function of ℓ . Since the regularised self force is convergent in ℓ , in numerical studies of the self force a finite number of ℓ values are computed, which amounts to some corresponding accuracy when computing the full sum over spherical harmonics, as in say Eq. (2.42). In the case of post-Newtonian expansions, the situation is somewhat different. It turns out one can compute homogeneous solutions leaving ℓ as a parameter, the drawback being that they typically breakdown for low ℓ . Thus the strategy is to compute explicit PN expansions for $\ell = 2 \dots 6$, the rest being captured by the general expansions. Typically as one increases the order of our PN expansion more low ℓ 's are needed.

As a particular example we will run through the procedure with $\ell = 2$. In Sec.II B of [47], the homogeneous solutions to the radial Teukolsky equation are computed (similarly to Ref. [49]) as expansions in $\eta = \frac{1}{\bar{c}}$ with coefficients in terms of the two variables $X_1 = \bar{G}M/r$, and $\sqrt{X_2} = \omega r$. These expansions are simplified by writing them as a product of an exponential factor with radial dependence entirely contained in logarithmic terms, and a remaining series in η . After limiting $a \rightarrow 0$ these are

$$\begin{aligned} \hat{R}_{\ell=2m\omega}^+ = & -\frac{72iX_1^4\eta^7}{\sqrt{X_2}} \left(1 + i\sqrt{X_2}\eta + \left(5X_1 - \frac{X_2}{2} \right) \eta^2 \right. \\ & - \frac{1}{6}iX_2^{3/2}\eta^3 + \left(\frac{120X_1^2}{7} + \frac{7X_1X_2}{2} + \frac{X_2^2}{24} \right) \eta^4 \Big) \\ & + \mathcal{O}(\eta^{12}) \end{aligned} \quad (3.5)$$

$$\begin{aligned} \hat{R}_{\ell=2,m\omega}^- = & \frac{3}{2} - i\sqrt{X_2}\eta - \frac{11X_2}{28}\eta^2 + \frac{3}{28}iX_2^{3/2}\eta^3 \\ & + \left(-\frac{37X_1X_2}{42} + \frac{23X_2^2}{1008} \right) \eta^4 + \mathcal{O}(\eta^5) \end{aligned} \quad (3.6)$$

These solutions, while coming from usual Mano-Suzuki-Takasugi (MST) expansions, have been normalised to remove certain radius independent factors that are unimportant for constructing the inhomogeneous solution. The solutions can now be converted to series expansions in $1/p$ and e as a function of χ by using Eqs. (2.12) and (3.3) with the frequency $\omega = m\Omega_\varphi + n\Omega_r$ i.e. by evaluating (3.5),(3.6) at the position of the particle for the relevant frequency values. In the above, the η factors are simply an order counting tool and in practice can be dropped when converting to the expansion in $1/p$. For now we will hold off on fully switching to p, e and χ and instead swap p for the dimensionless frequency variable

$$y \equiv (m_2 \Omega_\varphi)^{2/3}. \quad (3.7)$$

The resulting double expansion in y and e for our example is

$$\begin{aligned}\hat{R}_{\ell=2,m\omega}^+ &= (1 + 5e \cos \chi + \mathcal{O}(e^2)) y^5 + (i(m+n) + 4i(m+n)e \cos \chi + \mathcal{O}(e^2)) y^{11/2} + \left(5 - \frac{1}{2}(m+n)^2\right. \\ &\quad \left.+ \left(30 - \frac{3}{2}(m+n)^2\right) e \cos \chi + \mathcal{O}(e^2)\right) y^6 - \left(3in + \frac{1}{6}i(m+n)^3 + \left(12in + \frac{1}{3}i(m+n)^3\right) e \cos \chi + \mathcal{O}(e^2)\right) y^{13/2} \\ &\quad + \mathcal{O}(y^7)\end{aligned}\quad (3.8)$$

$$\begin{aligned}\hat{R}_{\ell=2,m\omega}^- &= \frac{3}{2} + (-i(m+n) + i(m+n)e \cos \chi + \mathcal{O}(e^2)) y^{1/2} + \left(-\frac{11}{28}(m+n)^2 + \frac{11}{14}(m+n)^2 e \cos \chi + \mathcal{O}(e^2)\right) y \\ &\quad + \left(\frac{3}{2}\left(2in + \frac{1}{14}i(m+n)^3\right) + \frac{3}{2}\left(-2in - \frac{3}{14}i(m+n)^3\right) e \cos \chi + \mathcal{O}(e^2)\right) y^{3/2} + \mathcal{O}(y^2)\end{aligned}\quad (3.9)$$

where we have removed a constant factor from $R_{\ell m \omega}^+$.

C. Hertz potential and retarded metric perturbation

The modes of the Hertz potential and thus the metric perturbation are then straightforwardly given using Eqs. (2.51) and (2.52). The main new feature in this construction as compared to the circular case is the sum over the radial frequencies, i.e. the infinite sum over n . The sum is expected to be exponentially convergent for the bound geodesics we are considering. This manifests in the convergence of the small eccentricity expansion. What one finds is that for an expansion valid to e^k , one needs only to sum $n = -k, \dots, k$ with $h_{\mu\nu}^{\ell mn} = \mathcal{O}(e^{k+1})$ for $|n| > k$. In other words, to capture higher eccentricity orbits one needs more and more n -modes.

Computationally speaking the n -sum can be time consuming and a potential bottleneck. We find it therefore more convenient to sum the n -modes of the Hertz potential to give $\Psi_{\ell m}(t, r) = \sum_{n=-\infty}^{\infty} e^{-i\omega t} \Psi_{\ell m \omega}(r)$. We must also compute the n -sum for each of the t derivatives that then appear in (2.52), for example

$$\partial_t \Psi_{\ell m}(t, r) = -i \sum_{n=-\infty}^{\infty} \omega e^{-i\omega t} \Psi_{\ell m \omega}(r). \quad (3.10)$$

as well as the relevant r derivatives (which are obtained from Eq. (2.51)). With these in hand the metric perturbation as a function of ℓ and m is more or less trivially computed. For instance, using Eq. (2.52)

$$\begin{aligned}h_{tt}^{\ell m} &= \frac{1}{8}(r - 2m_2)^2 \left(\partial_{\theta}^2 {}_2Y_{\ell m}(\theta, \varphi) + 2m \partial_{\theta} {}_2Y_{\ell m}(\theta, \varphi) \right. \\ &\quad \left. + (m^2 - 2) {}_2Y_{\ell m}(\theta, \varphi) \right) \Psi_{\ell m}(t, r)\end{aligned}\quad (3.11)$$

(which can be recognized as being a pure scalar harmonic). Each metric component and its derivatives are in practice evaluated here at the position of the particle. At this stage we sum over m to get $h_{\mu\nu}^{\ell} = \sum_{m=-\ell}^{\ell} h_{\mu\nu}^{\ell m}$.

As will be discussed in more detail below, the singular nature of $h_{\mu\nu}(t, r, \theta, \varphi)$ in the vicinity of the source world-line requires us to separately evaluate the two different radial limits $r \rightarrow r_p^{\pm}$, from above or below, of the modes. These limits are indicated below by a \pm subscript: e.g. $h_{\mu\nu, \pm}^{\ell}$, or $F_{\mu, \pm}^{\ell}$.

For ease of reading we omit explicit expressions for each of the components of the metric and their first order partial derivatives.

D. Perturbed geodesic and self-force

Before calculating the perturbed orbit quantities $\delta \mathcal{E}$, $\delta \mathcal{L}$ and δu^r we must explicitly compute the t and φ components of the conservative self-force. The self-force is given by [50]

$$\begin{aligned}F^{\mu} &= \mathcal{P}^{\mu\nu\lambda\rho} (2h_{\nu\lambda; \rho} - h_{\lambda\rho; \nu}), \\ \mathcal{P}^{\mu\nu\lambda\rho} &= -\frac{1}{2}(g^{\mu\nu} + u^{\mu}u^{\nu})u^{\lambda}u^{\rho}.\end{aligned}\quad (3.12)$$

where the covariant derivatives here are taken with respect to the background metric. Formally this equation requires the regularised metric perturbation; we will however use it with the ℓ -modes of the retarded metric and leave regularization to later.

At linear order in the mass ratio we can uniquely define the dissipative and conservative parts of the self-force via

$$F_{\text{diss}}^{\mu} = \frac{1}{2}(F^{\mu}[h_{\mu\nu}^{\text{ret}}] - F^{\mu}[h_{\mu\nu}^{\text{adv}}]), \quad (3.13)$$

$$F_{\text{cons}}^{\mu} = \frac{1}{2}(F^{\mu}[h_{\mu\nu}^{\text{ret}}] + F^{\mu}[h_{\mu\nu}^{\text{adv}}]). \quad (3.14)$$

Noting the symmetry relation of Eq. (2.80) of [51], the authors of [52] rewrote these for the case of equatorial geodesics purely in terms of the retarded solution:

$$F_{\text{diss}}^{\mu} = \frac{1}{2}(F_{\text{ret}}^{\mu}(\tau) - \epsilon_{(\mu)} F_{\text{ret}}^{\mu}(-\tau)) \quad (3.15)$$

$$F_{\text{cons}}^{\mu} = \frac{1}{2}(F_{\text{ret}}^{\mu}(\tau) + \epsilon_{(\mu)} F_{\text{ret}}^{\mu}(-\tau)), \quad (3.16)$$

where $\epsilon_{(\mu)} = (-1, 1, 1, -1)$. With our parameterization

$F_{\text{ret}}^\mu(\tau \rightarrow -\tau) \equiv F_{\text{ret}}^\mu(\chi \rightarrow -\chi)$. We compute this analytically, e.g. for $\ell = 2$ we have

$$q^{-1}F_{t,+}^{\text{cons}} = (3e \sin(\chi) + 3e^2 \sin(2\chi) + \mathcal{O}(e^3)) p^{-5/2} + \left(-\frac{85}{14}e \sin(\chi) - \frac{127}{14}e^2 \sin(2\chi) + \mathcal{O}(e^3) \right) p^{-7/2} + \mathcal{O}(p^{-9/2}), \quad (3.17)$$

$$q^{-1}F_{\varphi,+}^{\text{cons}} = \left(-\frac{9}{7}e \sin(\chi) - \frac{9}{7}e^2 \sin(2\chi) + \mathcal{O}(e^3) \right) p^{-2} + \left(\frac{1237}{126}e \sin(\chi) + \frac{114}{7}e^2 \sin(2\chi) + \mathcal{O}(e^3) \right) p^{-3} + \mathcal{O}(p^{-4}). \quad (3.18)$$

In this double expansion the corresponding integrals needed for Eq. (2.32) are trivially computed order by order. Explicitly,

$$\delta\mathcal{E}_{\text{BS}}^+ = \left(\frac{3}{2} + 3e \cos(\chi) + \mathcal{O}(e^2) \right) p^{-1} + \left(\frac{1}{14} - \frac{11}{7}e \cos(\chi) + \mathcal{O}(e^2) \right) p^{-2} + \mathcal{O}(p^{-3}), \quad (3.19)$$

$$\delta\mathcal{L}_{\text{BS}}^+ = \frac{3}{2}\sqrt{p} + \left(\frac{1}{14} + \frac{9}{7}e \cos(\chi) + \mathcal{O}(e^2) \right) p^{-1/2} + \mathcal{O}(p^{-3/2}), \quad (3.20)$$

and the corresponding $\delta u_{\text{BS}}^{r,+}$ is found using the normalisation condition.

Eq. (2.25) is now straightforwardly computed here to give the $\ell = 2$ values

$$\begin{aligned} \delta\dot{\Psi}^+ &= \left(\frac{3}{2} + 3 \cos \chi e + \mathcal{O}(e^2) \right) p^{-3/2} \\ &\quad - \left(\frac{19}{28} + \frac{11}{7} \cos \chi e + \mathcal{O}(e^2) \right) p^{-5/2} + \mathcal{O}(p^{-3}), \\ \delta\dot{\Psi}^- &= - (1 + 2 \cos \chi e + \mathcal{O}(e^2)) p^{-3/2} \\ &\quad + \frac{4}{7} (1 + 6 \cos \chi e + \mathcal{O}(e^2)) p^{-5/2} + \mathcal{O}(p^{-3}). \end{aligned} \quad (3.21)$$

The integration of Eq. (2.24) is as usual applied order by order.

The only remaining perturbed quantities are the frequency shifts $\delta\Omega_r$, $\delta\Omega_\varphi$ which appear in (2.9). Using Eq. (75)-(76) of [40] (without the α factor) we find

$$\begin{aligned} m_2\delta\Omega_r^+ &= \left(\frac{3}{2} - \frac{9}{4}e^2 \right) p^{-3/2} \\ &\quad + \left(-\frac{115}{28} + \frac{575}{56}e^2 + \mathcal{O}(e^4) \right) p^{-5/2} + \mathcal{O}(p^{-7/2}) \end{aligned}$$

$$\begin{aligned} m_2\delta\Omega_\varphi^+ &= \left(\frac{3}{2} - \frac{9}{4}e^2 \right) p^{-3/2} \\ &\quad + \left(-\frac{103}{28} + \frac{77}{8}e^2 + \mathcal{O}(e^4) \right) p^{-5/2} + \mathcal{O}(p^{-7/2}) \end{aligned} \quad (3.22)$$

Using Eq. (2.9) we give illustrative sample expansions for the retarded $\Delta\psi$ for $\ell = 2$:

$$\begin{aligned} \Delta\psi_{\ell=2}^+ &= \frac{43}{48}p^{-1} - \left(\frac{2047}{1008} + \frac{1427}{1008}e^2 \right) p^{-2} + \mathcal{O}(p^{-3}), \\ \Delta\psi_{\ell=2}^- &= \frac{43}{48}p^{-1} - \left(\frac{5827}{1008} + \frac{1427}{1008}e^2 \right) p^{-2} + \mathcal{O}(p^{-3}). \end{aligned} \quad (3.23)$$

Let us also give sample expansions for the generic ℓ expressions. These are valid for all ℓ greater than a value determined by the given PN order (see Sec II B. of Ref. [35]). These are useful for clearly seeing the divergent behaviour discussed in the next sections

$$\begin{aligned} \Delta\psi_\ell^+ &= \frac{3(1+7\ell+7\ell^2)}{4(-1+2\ell)(3+2\ell)} p^{-1} + \left(\frac{3(1920-1197\ell-852\ell^2+120\ell^3-653\ell^4-402\ell^5+90\ell^6+64\ell^7)}{16\ell(1+\ell)(-3+2\ell)(-1+2\ell)(3+2\ell)(5+2\ell)} \right. \\ &\quad \left. - \frac{3(-480-102\ell-466\ell^2-583\ell^3+71\ell^4+435\ell^5+145\ell^6)e^2}{32\ell(1+\ell)(-3+2\ell)(-1+2\ell)(3+2\ell)(5+2\ell)} \right) p^{-2} + \mathcal{O}(p^{-3}), \\ \Delta\psi_\ell^- &= \frac{3(1+7\ell+7\ell^2)}{4(-1+2\ell)(3+2\ell)} p^{-1} + \left(-\frac{3(-1920+1377\ell+1104\ell^2-848\ell^3-467\ell^4+402\ell^5+358\ell^6+64\ell^7)}{16\ell(1+\ell)(-3+2\ell)(-1+2\ell)(3+2\ell)(5+2\ell)} \right. \end{aligned}$$

$$- \frac{3(-480 - 102\ell - 466\ell^2 - 583\ell^3 + 71\ell^4 + 435\ell^5 + 145\ell^6)e^2}{32\ell(1+\ell)(-3+2\ell)(-1+2\ell)(3+2\ell)(5+2\ell)}p^{-2} + \mathcal{O}(p^{-3}). \quad (3.24)$$

E. Regularization and completion of the metric perturbation

1. Metric completion (non-radiative multipoles $\ell \leq 1$)

The reconstructed retarded metric perturbation obtained by the CCK procedure laid out in the previous section is well known to give the full metric perturbation modulo perturbations to the spacetime due to the mass and angular momentum of the small body. In Schwarzschild spacetime this amounts to the absence of the spherical harmonic $\ell = 0, 1$ modes in the reconstructed metric. We include these modes using the corrected Regge-Wheeler-Zerrilli low multipoles given in Appendix A of [15]. Ultimately, the contribution of these modes to the spin precession is, at leading orders,

$$\begin{aligned} \Delta\psi_{\ell=0}^+ &= \frac{-\frac{3}{4} + \frac{3e^2}{4}}{p^2} + \frac{-\frac{73}{16} + \frac{45e^2}{8} + \mathcal{O}(e^4)}{p^3} + \mathcal{O}(p^{-4}), \\ \Delta\psi_{\ell=0}^- &= \frac{-\frac{3}{4} + \frac{3e^2}{4}}{p^3} + \frac{-\frac{97}{16} + \frac{199e^2}{16} + \mathcal{O}(e^4)}{p^4} + \mathcal{O}(p^{-5}), \end{aligned} \quad (3.25)$$

$$\begin{aligned} \Delta\psi_{\ell=1}^+ &= \frac{1}{p} + \frac{\frac{9}{2} - e^2}{p^2} + \frac{\frac{163}{8} - 6e^2 + \mathcal{O}(e^4)}{p^3} + \mathcal{O}(p^{-4}), \\ \Delta\psi_{\ell=1}^- &= \frac{1}{p} + \frac{\frac{9}{4} + \frac{5e^2}{4}}{p^2} + \frac{\frac{143}{16} + \frac{51e^2}{8} + \mathcal{O}(e^4)}{p^3} + \mathcal{O}(p^{-4}). \end{aligned} \quad (3.26)$$

2. Regularization

So far in the calculation we have been constructing the ℓ -modes of the spin precession invariant from the retarded metric perturbation. The full retarded metric perturbation is, however, a singular quantity due to the point particle delta function source term. This singular nature manifests itself as a *direction-dependent* divergent sum over ℓ -modes (each individual ℓ -mode being, however, finite), which for our spin-precession invariant takes the following form for large values of ℓ

$$\Delta\psi_{\pm}^{\ell \rightarrow \infty} = \pm A_{\infty}(2\ell + 1) + B_{\infty} + \mathcal{O}(\ell^{-2}). \quad (3.27)$$

Here, the sign of the A_{∞} term is dependent on whether one takes the limit $r \rightarrow r_p^{\pm}$ from above or below. By definition, the two coefficients A_{∞} and B_{∞} parametrizing the large- ℓ singular behavior Eq. (3.27) of $\Delta\psi$ are

independent of ℓ . [We have shown above (see, e.g., Eq. (3.24)) how our PN-expanded method allows us to analytically extract the values of A_{∞} and B_{∞} .]

Calculating the regularized finite value for $\Delta\psi$ involves calculating, and subtracting out, the ℓ -modes of the corresponding singular piece $\Delta\psi^S$ of $\Delta\psi$, say

$$\Delta\psi_{\ell,\pm}^S = \pm A_S(2\ell + 1) + B_S. \quad (3.28)$$

Detweiler and Whiting [37] have shown how to define the singular pieces of $h_{\mu\nu}^S$, $h_{\mu\nu,\lambda}^S$, and thereby $\Delta\psi^S$, in the Lorenz gauge. Recently, Ref. [53] emphasized that, when using (as we do here, following most of the analytical work on SF corrections to gauge-invariant quantities starting with Ref. [23]) a decomposition of $h_{\mu\nu}$ in *tensor-harmonics* ℓ -modes, there are subtleties concerning the value of the coefficient A_S of $(2\ell + 1)$ in Eq. (3.28). Indeed, while the value of B_S was found to be independent of ℓ (and therefore equal to the large- ℓ value B_{∞} entering Eq. (3.27)), Ref. [53] found that the value of A_S stabilized to its asymptotic value A_{∞} only when $\ell \geq 2$. The (nonradiative) modes $\ell = 0, 1$ involve A_S coefficients that differ from A_{∞} . This difference comes from the peculiar low- ℓ dependence of certain radial derivatives of $h_{\mu\nu}$, such as $h_{t\varphi,r}$ or $h_{\varphi\varphi,r}$, see notably Eqs. (6.11j) and (6.11l) in Ref. [53]. [Although the mentioned equations deal with the case of circular orbits, nothing fundamentally changes in the eccentric orbit case, and the same conclusions hold.]

In the Lorenz gauge, the correct regularized spin-precession invariant would then be given by

$$\Delta\psi = \sum_{\ell=0}^{\infty} (\Delta\psi_{\pm}^{\ell} - \Delta\psi_S^{\ell,\pm}) = \sum_{\ell=0}^{\infty} (\Delta\psi_{\pm}^{\ell} \mp A_S(2\ell + 1) - B_S). \quad (3.29)$$

This result can be simplified by working with the average over the limits $r \rightarrow r_p$ from both sides, thereby avoiding the need to keep track of the low ℓ dependence of A_S , and being able to use the value of B_{∞} extracted from the large ℓ behavior of $\Delta\psi$, Eq. (3.27):

$$\Delta\psi = \sum_{\ell=0}^{\infty} \left(\frac{1}{2} (\Delta\psi^{\ell,+} + \Delta\psi^{\ell,-}) - B_{\infty} \right). \quad (3.30)$$

There is one remaining subtlety, which is that the regularization procedure explained above was derived in the Lorenz gauge while our calculation of the tensorial ℓ modes was done in a radiation gauge. There are two ways to deal with this additional subtlety. On the one hand, as was shown by Barack and Ori in the case of the gravitational self-force [54], the procedure should also

work in a gauge which can be reached from Lorenz gauge by a gauge transformation of a smooth enough nature. The radiation gauge we work in does not, however, fall within this class (for a discussion of the singular structure of the various radiation gauges we refer the reader to [55]). Much work has been focussed recently on formally defining the correct regularization procedure in radiation gauge. The strategy employed is to construct a gauge transformation ξ^μ which is defined locally in the vicinity of the worldline, and which can be used to transform from the radiation gauge to a “locally Lorenz” gauge. It is then argued that this generates a correction to the mode-sum formula, Eq. (3.29), which has an overall sign that differs on either side of the $r \rightarrow r_p$ limit. A straightforward solution, then, is to take the average of both limits (as we are doing), thus eliminating the troublesome singular gauge contribution.

On the other hand, it is likely that a much simpler

solution resolves the issue in our case. The quantity we are computing is a gauge invariant, and so we are free to use the regularization procedure derived in the Lorenz gauge (with averaging to compensate for the fact that we are not decomposing into scalar spherical harmonics) and apply it to any other gauge in the same invariance class. We stress the point, however, for two reasons: (i) while we have partially checked the correctness of this fact (notably by checking the continuity of $\Delta\psi_\pm^\ell - \Delta\psi_S^{\ell,\pm}$ across $r = r_p$), we have not analytically shown that $\Delta\psi$ is invariant under the transformation from Lorenz to radiation gauge; and (ii) things would not be so straightforward if one were interested in computing non-gauge-invariant quantities such as the self-force (in which case a more careful analysis along the lines of Ref. [55] may be required).

Finally, taking the large- ℓ limit of our expression (3.24) (to all orders we computed) we can read off the A_∞ and B_∞ coefficients:

$$A_\infty = \frac{3}{8}p^{-2} + \left(\frac{49}{32} - \frac{27e^2}{32} + \mathcal{O}(e^4)\right)p^{-3} + \left(\frac{1007}{128} - \frac{221e^2}{32} + \mathcal{O}(e^4)\right)p^{-4} + \left(\frac{23441}{512} - \frac{25437e^2}{512} + \mathcal{O}(e^4)\right)p^{-5} \\ + \left(\frac{575983}{2048} - \frac{354297e^2}{1024} + \mathcal{O}(e^4)\right)p^{-6} + \mathcal{O}(p^{-7}) \quad (3.31)$$

$$B_\infty = \frac{21}{16}p^{-1} + \left(-\frac{201}{128} - \frac{435e^2}{512} + \mathcal{O}(e^4)\right)p^{-2} + \left(\frac{529}{1024} - \frac{1155e^2}{1024} + \mathcal{O}(e^4)\right)p^{-3} + \left(\frac{152197}{16384} - \frac{352849e^2}{65536} + \mathcal{O}(e^4)\right)p^{-4} \\ + \left(\frac{17145445}{262144} - \frac{5100243e^2}{131072} + \mathcal{O}(e^4)\right)p^{-5} + \left(\frac{886692225}{2097152} - \frac{2456459237e^2}{8388608} + \mathcal{O}(e^4)\right)p^{-6} + \mathcal{O}(p^{-7}). \quad (3.32)$$

Although we do not use the A_∞ parameter due to our averaging, we have checked that both of these expressions agree with independently calculated A_S and B_S . These were computed from expansions in p and e of the singular metric perturbation calculated using the methods described in [53, 56, 57] with a projection onto scalar spherical harmonics. This provides a valuable check that our radiative modes ($\ell \geq 2$) are demonstrating the cor-

rect singular behaviour.

F. Spin-precession invariant

Upon regularization, our generic- ℓ terms can be seen analytically to converge as ℓ^{-2} . This allows us to explicitly compute the infinite series over ℓ , giving one of our main results

$$\Delta\psi = -p^{-1} + \left(\frac{9}{4} + e^2\right)p^{-2} + \left(\left(\frac{739}{16} - \frac{123\pi^2}{64}\right) + \left(\frac{341}{16} - \frac{123\pi^2}{256}\right)e^2 + \mathcal{O}(e^4)\right)p^{-3} + \left(\left(-\frac{587831}{2880} + \frac{1256\gamma}{15}\right) \right. \\ + \frac{31697\pi^2}{6144} + \frac{296\log(2)}{15} + \frac{729\log(3)}{5} - \frac{628\log(p)}{15}\Bigg) + \left(-\frac{164123}{480} + \frac{536\gamma}{5} - \frac{23729\pi^2}{4096} + \frac{11720\log(2)}{3}\right. \\ \left.- \frac{10206\log(3)}{5} - \frac{268\log(p)}{5}\right)e^2 + \mathcal{O}(e^4)\Bigg)p^{-4} + \left(\left(-\frac{48221551}{19200} - \frac{22306\gamma}{35} + \frac{2479221\pi^2}{8192} + \frac{22058\log(2)}{105}\right) \right. \\ \left.- \frac{31347\log(3)}{28} + \frac{11153\log(p)}{35}\right) + \left(-\frac{164123}{480} + \frac{536\gamma}{5} - \frac{23729\pi^2}{4096} + \frac{11720\log(2)}{3} - \frac{10206\log(3)}{5}\right)$$

$$\begin{aligned}
& -\frac{268 \log(p)}{5} e^2 + \mathcal{O}(e^4) \Big) p^{-5} + \left(\frac{49969\pi}{315} + \frac{319609\pi}{630} e^2 + \mathcal{O}(e^4) \right) p^{-11/2} + \left(\left(-\frac{1900873914203}{101606400} - \frac{344021\gamma}{1890} \right. \right. \\
& + \frac{7230119267\pi^2}{2359296} - \frac{7335303\pi^4}{131072} - \frac{2514427 \log(2)}{270} + \frac{234009 \log(3)}{70} + \frac{9765625 \log(5)}{9072} + \frac{344021 \log(p)}{3780} \Big) \\
& + \left(-\frac{464068669129}{5080320} - \frac{2508913\gamma}{945} + \frac{32088966503\pi^2}{2359296} - \frac{146026515\pi^4}{1048576} + \frac{273329813 \log(2)}{945} - \frac{159335343 \log(3)}{8960} \right. \\
& \left. \left. - \frac{17193359375 \log(5)}{145152} + \frac{2508913 \log(p)}{1890} \right) e^2 + \mathcal{O}(e^4) \Big) p^{-6} + \mathcal{O}(p^{-13/2}). \tag{3.33}
\end{aligned}$$

We could perform several checks on our result Eq. (3.33). First, the first three terms of our expansion can be seen immediately to agree with the coefficients of p^{-1}, p^{-2}, p^{-3} derived in Ref. [36].

Second, we have compared the numerical values predicted by our analytical result Eq. (3.33) to the numerical estimates of $\Delta\psi$ obtained in Ref. [36]. More precisely, Table II in Ref. [36] lists a sample of numerical estimates $\Delta\psi^{\text{num}}(p_i, e_k)$ of $\Delta\psi(p, e)$, together with estimates of the corresponding numerical error $\sigma^{\text{num}}(p_i, e_k)$. This sample includes nineteen values, $10 \leq p_i \leq 100$, of the semilatus rectum p , and, for each value of p , nine values of e , namely: $e_k = 0.050 + 0.025k, k = 0, 1, \dots, 8$. As our main aim was to compare the eccentricity dependence of these numerical data to the one predicted by our analytical result Eq. (3.33), we extracted a numerical estimate of the $\mathcal{O}(e^2)$ contribution, say $e^2 \Delta\psi^{(2)}(p)$, in $\Delta\psi(p, e)$,

$$\Delta\psi(p, e) = \Delta\psi^{(0)}(p) + e^2 \Delta\psi^{(2)}(p) + e^4 \Delta\psi^{(4)}(p) + \mathcal{O}(e^6), \tag{3.34}$$

in the following way. First, we used the analytical knowledge of the leading-order $\mathcal{O}(e^4)$ contribution, $e^4 \Delta\psi_{\text{LO}}^{e^4}(p) = -\frac{1}{2} e^4 p^{-3}$ (derived in Ref. [36]) to work with the $\mathcal{O}(e^4)$ -corrected¹ inclusion of numerical data $\Delta\psi^{\text{num}'}(p_i, e_k) \equiv \Delta\psi^{\text{num}}(p_i, e_k) - e_k^4 \Delta\psi_{\text{LO}}^{e^4}(p_i)$. Then, we fitted, for each value of p_i the nine numerical data $\Delta\psi^{\text{num}'}(p_i, e_k); k = 0, 1, \dots, 8$ to a linear function of e^2 , say

$$\Delta\psi^{\text{fit}}(p_i, e_k) = m_i e_k^2 + q_i. \tag{3.35}$$

We used a least-squares fit, weighted by the (inverse squares of the) corresponding numerical errors $\sigma^{\text{num}}(p_i, e_k)$ listed in Table II of [36]. This fitting procedure gave us (for each p_i) estimates of $m_i (\approx \Delta\psi^{(2)}(p_i))$ and $q_i (\approx \Delta\psi^{(0)}(p_i))$, together with corresponding fitting errors (obtained from the covariance matrix). In addition, the goodness of each fit is measured by the corresponding reduced $\chi_{\text{red}}^2(p_i) = \chi_{\text{min}}^2 / (N_{\text{data}} - N_{\text{param}})$, with $N_{\text{data}} = 9$ and $N_{\text{param}} = 2$.

¹ We have checked that taking into account a next-to-leading-order $\mathcal{O}(e^4)$ contribution, $e^4 \Delta\psi_{\text{NLO}}^{e^4}(p) = c_{\text{NLO}} e^4 p^{-4}$, with $|c_{\text{NLO}}| \leq 3$ did not significantly change our results.

TABLE I. Comparison between the values $m^{\text{thy}}(p)$, predicted by our analytical result Eq. (3.33), of the coefficient of e^2 in $\Delta\psi(p, e) = q(p) + m(p)e^2 + \mathcal{O}(e^4)$, to the numerical estimates $m^{\text{num}}(p)$ of $m(p)$ obtained by least-squares fitting the numerical data for $\Delta\psi(p, e)$ obtained in Ref. [36]. The last entry is the ratio (3.36) between the difference $m^{\text{num}} - m^{\text{thy}}$ and our estimate of the total error $\sigma_m^{\text{tot}} \equiv \sqrt{(\sigma_m^{\text{num}})^2 + (\sigma_m^{\text{thy}})^2}$ on m . See text for more details (notably about the estimates of the errors σ_m^{thy} and σ_m^{num}).

p	χ_{red}^2	$m^{\text{num}}(\sigma_m^{\text{num}})$	$m^{\text{thy}}(\sigma_m^{\text{thy}})$	r_m
10	0.221	$2.83892(11) \times 10^{-2}$	$3.9(2.1) \times 10^{-2}$	-0.50
15	4.119	$9.12787(61) \times 10^{-3}$	$10.0(1.9) \times 10^{-3}$	-0.45
20	2.591	$4.40237(32) \times 10^{-3}$	$4.54(34) \times 10^{-3}$	-0.41
25	0.0357	$2.561664(35) \times 10^{-3}$	$2.956(92) \times 10^{-3}$	-0.38
30	0.867	$1.66508(23) \times 10^{-3}$	$1.677(31) \times 10^{-3}$	-0.37
35	0.646	$1.16553(20) \times 10^{-3}$	$1.170(12) \times 10^{-3}$	-0.33
40	0.372	$8.5943(13) \times 10^{-4}$	$8.611(56) \times 10^{-4}$	-0.30
45	0.102	$6.58803(62) \times 10^{-4}$	$6.597(28) \times 10^{-4}$	-0.32
50	0.227	$5.20356(82) \times 10^{-4}$	$5.211(15) \times 10^{-4}$	-0.48
55	0.481	$4.2124(14) \times 10^{-4}$	$4.2171(85) \times 10^{-4}$	-0.55
60	1.249	$3.4798(15) \times 10^{-4}$	$3.4813(50) \times 10^{-4}$	-0.29
65	3.630	$2.9178(37) \times 10^{-4}$	$2.9215(31) \times 10^{-4}$	-0.75
70	1.800	$2.4793(43) \times 10^{-4}$	$2.4859(20) \times 10^{-4}$	-1.40
75	3.113	$2.1347(49) \times 10^{-4}$	$2.1405(13) \times 10^{-4}$	-1.14
80	6.313	$1.8583(90) \times 10^{-4}$	$1.86200(91) \times 10^{-4}$	-0.41
85	4.832	$1.6289(97) \times 10^{-4}$	$1.63430(64) \times 10^{-4}$	-0.56
90	1.707	$1.4552(39) \times 10^{-4}$	$1.44578(45) \times 10^{-4}$	2.42
95	3.687	$1.292(10) \times 10^{-4}$	$1.28797(33) \times 10^{-4}$	0.37
100	1.882	$1.130(14) \times 10^{-4}$	$1.15456(24) \times 10^{-4}$	-1.70

The results for our estimates of the e^2 slope $m_i \approx \Delta\psi^{(2)}(p_i)$ are given in Table I below.

Namely, the first four entries of Table I are: p_i , $\chi_{\text{red}}^2(p_i)$, the numerical estimate m_i^{num} of the slope $m_i \approx \Delta\psi^{(2)}(p_i)$ obtained from our fit, and the corresponding theoretical prediction $m_i^{\text{thy}} \equiv \Delta\psi^{(2)\text{PN}}(p_i)$, as obtained from our PN-expanded analytical result Eq. (3.33). In addition, we have indicated in parentheses, both for the numerical estimates of m_i , and their theoretical ones, estimates of the corresponding uncertainties in their values. The estimates of the numerical uncertainty on m_i^{num} was obtained by renormalizing the fitting error

by a factor $\sqrt{\chi_{\text{red}}^2(p_i)}$. Indeed, though the goodness-of-fit parameters $\chi_{\text{red}}^2(p_i)$ were always (as tabulated) of order unity, they were not always numerically close to 1. In view of the difficulty, in numerical SF computations, to accurately estimate the numerical error, we considered that a value of $\chi_{\text{red}}^2(p_i)$ different from 1 indicated an inaccurate estimate of the numerical errors $\sigma_m^{\text{num}}(p_i, e_k)$, which we (coarsely) corrected by multiplying $\sigma_m^{\text{num}}(p_i, e_k)$ (and correlatively $\sigma_{m_i}^{\text{fit}}$) by a factor $\sqrt{\chi_{\text{red}}^2(p_i)}$. Concerning the theoretical error $\sigma_{m_i}^{\text{thy}}$, it was estimated by the value of the last analytically computed contribution to $\Delta\psi^{(2)}(p_i)$ in Eq. (3.33), i.e. $\sigma_m^{\text{thy}}(p) = |(18101.8418 + 1327.467196 \ln(p))1/p^6|$.

Finally, the last entry in Table I displays the values of the ratios

$$r_{m_i} \equiv \frac{m_i^{\text{num}} - m_i^{\text{thy}}}{\sigma_{m_i}^{\text{tot}}}, \quad (3.36)$$

where $\sigma_i^{\text{tot}} \equiv \sqrt{(\sigma_{m_i}^{\text{num}})^2 + (\sigma_{m_i}^{\text{thy}})^2}$ is an estimate of the combined numerical-analytical error on $m_i \approx \Delta\psi^{(2)}(p_i)$. The most significant fact (for our purpose) in Table I is that the values of the latter ratios are all of order unity. This is a valuable, independent check on our analytical computations.

Our fitting procedure has also given use numerical estimates of $q_i \approx \Delta\psi^{(0)}(p_i)$ that we have satisfactorily compared (with corresponding ratios $r_{q_i} \equiv (q_i^{\text{num}} - q_i^{\text{thy}})/\sigma_{q_i}^{\text{tot}}$ found to be of order unity) to the 9.5PN current analytical knowledge of $\Delta\psi^{(0)}(p)$ (given in Appendix A of [36]). [We recall (see the end of section IV for more details) that $\Delta\psi^{(0)}(p) = \lim_{e \rightarrow 0} \Delta\psi(p, e)$ differs from $\Delta\psi^{\text{circ}}(p)$, and that the difference $\Delta\psi(p, e \rightarrow 0) - \Delta\psi^{\text{circ}}(p)$ was related in Section IIIB of [36] to the EOB function $\rho(x)$ measuring periastron precession. While $\Delta\psi^{\text{circ}}(p)$ is known to very high PN orders [35, 58], $\rho(x)$ is currently known to the 9.5-PN level [59].]

IV. IMPROVING THE SPIN-ORBIT SECTOR OF THE EOB HAMILTONIAN

In this section, we shall show how to transcribe the new SF results contained in Eq. (3.33) above into an improved knowledge of the spin-orbit sector of the EOB Hamiltonian. The inclusion of spin couplings in the EOB Hamiltonian was initiated in Ref. [8] and developed in Refs. [3, 19, 20, 34, 60–70]. Here, we will focus on the case of non-precessing spins (parallel or antiparallel to the orbital angular momentum), and only consider effects *linear* in spins. Following the formulation of Refs. [60, 66], the spin-orbit couplings are described by two phase-space-dependent gyrogravitomagnetic ratios g_S and g_{S*} .

We would like to emphasise that throughout this section all coordinate variables will be referring to EOB variables which, despite overlapping labelling, are not to be confused with those of the previous section. For example

we will encounter an EOB eccentricity e , which is distinct to that used in Eq. (3.33) to parameterize the noncircular dependence of the SF spin precession invariant. To avoid issues when relating the EOB spin precession function to its previous SF version, we shall henceforth relabel all independent variables from the previous section with an additional overbar, i.e. the variables entering Eq. (3.33) will be now written as \bar{p} and \bar{e} . In addition, in order to better explicate the introduction of various dimensionless quantities in the EOB formalism, we shall often return, in this section, to the use of physical units where G and c are not set to unity.

A. EOB notation and reminders

Let us first recall the standard EOB results and notation, which we shall follow here. The total Hamiltonian of the system is expressed as

$$H(\mathbf{R}, \mathbf{P}, \mathbf{S}_1, \mathbf{S}_2) = Mc^2 \sqrt{1 + 2\nu \left(\frac{H_{\text{eff}}}{\mu c^2} - 1 \right)} \equiv Mc^2 h, \quad (4.1)$$

where (with the convention $m_1 < m_2$, and $m_1 \ll m_2$ in the extreme-mass-ratio limit)

$$M = m_1 + m_2, \mu = \frac{m_1 m_2}{(m_1 + m_2)}, \nu = \frac{\mu}{M} = \frac{m_1 m_2}{(m_1 + m_2)^2}, \quad (4.2)$$

and where the *effective EOB Hamiltonian* H_{eff} is decomposed as

$$H_{\text{eff}} = H_{\text{eff}}^{\text{O}} + H_{\text{eff}}^{\text{SO}}. \quad (4.3)$$

Here

$$H_{\text{eff}}^{\text{O}} = c^2 \sqrt{A \left(\mu^2 c^2 + \mathbf{P}^2 + \left(\frac{1}{B} - 1 \right) P_R^2 + Q \right)}, \quad (4.4)$$

denotes the *orbital* part of the effective Hamiltonian, expressed in terms of the squared linear momentum

$$\mathbf{P}^2 = \frac{P_R^2}{B} + \frac{\mathbf{L}^2}{R^2} = \frac{P_R^2}{B} + \frac{P_\phi^2}{R^2}, \quad (4.5)$$

with $\mathbf{L} = \mathbf{R} \times \mathbf{P}$ denoting the orbital angular momentum (with magnitude $L \equiv P_\phi$), and in terms of the EOB radial potentials parametrizing the effective metric (specialized here to equatorial motions)

$$ds_{(\text{eff})}^2 = g_{\mu\nu}^{(\text{eff})} dX^\mu dX^\nu = -Ac^2 dT_{\text{eff}}^2 + BdR^2 + R^2 d\phi^2. \quad (4.6)$$

The last (quartic in momenta) contribution Q on the right-hand-side (rhs) of Eq. (4.4) will be defined below. In addition

$$H_{\text{eff}}^{\text{SO}} = G_S^{\text{phys}} \mathbf{L} \cdot \mathbf{S} + G_{S*}^{\text{phys}} \mathbf{L} \cdot \mathbf{S}_* \quad (4.7)$$

denotes the *spin-orbit* part of the effective Hamiltonian, expressed in terms of the following two symmetric combination of the spin vectors \mathbf{S}_1 and \mathbf{S}_2 of the system

$$\mathbf{S} = \mathbf{S}_1 + \mathbf{S}_2, \quad \mathbf{S}_* = \frac{m_2}{m_1} \mathbf{S}_1 + \frac{m_1}{m_2} \mathbf{S}_2. \quad (4.8)$$

In the parallel-spin case that we consider here $\mathbf{L} \cdot \mathbf{S} = LS = P_\phi S$ and $\mathbf{L} \cdot \mathbf{S}_* = LS_* = P_\phi S_*$. It is convenient to work with the following dimensionless, rescaled variables

$$r = \frac{c^2 R}{GM}, u = \frac{GM}{c^2 R} \equiv \frac{1}{r}, j \equiv p_\phi = \frac{c P_\phi}{GM \mu}, p_r = \frac{P_R}{\mu c}, \quad (4.9)$$

and quantities

$$\hat{H}_{(\text{eff})} = \frac{H_{(\text{eff})}}{\mu c^2} \equiv \hat{H}_{(\text{eff})}^{\text{O}} + \hat{H}_{(\text{eff})}^{\text{SO}}$$

$$\begin{aligned} g_S^{\text{PN}}(u, p_r, p_\phi) &= 2 + \eta^2 \left[-\frac{5}{8} \nu u - \frac{27}{8} \nu p_r^2 \right] + \eta^4 \left[\nu \left(-\frac{51}{4} u^2 - \frac{21}{2} u p_r^2 + \frac{5}{8} p_r^4 \right) + \nu^2 \left(-\frac{1}{8} u^2 + \frac{23}{8} u p_r^2 + \frac{35}{8} p_r^4 \right) \right] + \mathcal{O}(\eta^6), \\ g_{S*}^{\text{PN}}(u, p_r, p_\phi) &= \frac{3}{2} + \eta^2 \left[-\frac{9}{8} u - \frac{15}{8} p_r^2 + \nu \left(-\frac{3}{4} u - \frac{9}{4} p_r^2 \right) \right] + \eta^4 \left[-\frac{27}{16} u^2 + \frac{69}{16} u p_r^2 + \frac{35}{16} p_r^4 \right. \\ &\quad \left. + \nu \left(-\frac{39}{4} u^2 - \frac{9}{4} u p_r^2 + \frac{5}{2} p_r^4 \right) + \nu^2 \left(-\frac{3}{16} u^2 + \frac{57}{16} u p_r^2 + \frac{45}{16} p_r^4 \right) \right] + \mathcal{O}(\eta^6). \end{aligned} \quad (4.11)$$

Here $\eta \sim 1/c$ is a place-holder for keeping track of the PN order, which we shall generally ignore in the following. The values of g_S and g_{S*} cited above have been expressed in the Damour-Jaranowski-Schaefer (DJS) spin gauge [60, 71], which is defined so that these quantities do not actually depend on p_ϕ .

Note that, at the PN order indicated above, g_S and g_{S*} depend on the symmetric mass ratio ν in the following way

$$\begin{aligned} g_S(u, p_r, p_\phi) &= 2 + \nu g_S^{(\nu^1)}(u, p_r, p_\phi) + \nu^2 g_S^{(\nu^2)}(u, p_r, p_\phi) \\ &\quad + \mathcal{O}(\nu^3), \\ g_{S*}(u, p_r, p_\phi) &= g_{S*}^{(\nu^0)}(u, p_r, p_\phi) + \nu g_{S*}^{(\nu^1)}(u, p_r, p_\phi) \\ &\quad + \nu^2 g_{S*}^{(\nu^2)}(u, p_r, p_\phi) + \mathcal{O}(\nu^3). \end{aligned} \quad (4.12)$$

Analytical gravitational self-force theory allowed one to improve the knowledge on the first gyrogravitomagnetic ratio g_S , to linear order in ν and for circular orbits. Namely, Ref. [70] derived (along circular orbits) the PN expansion of $g_S^{(\text{circ})}(u) = 2 - \frac{5}{8} \nu u \delta G_S^{\text{resc}} + \mathcal{O}(\nu^2)$ to the 7.5PN level, see Eq. (4.3) there. For concreteness, let us quote here only the first few terms of this expansion

$$\begin{aligned} g_S^{(\text{circ})}(u) &= 2 - \frac{5}{8} \nu u \left[1 + \frac{102}{5} u + \left(\frac{80399}{720} - \frac{241}{120} \pi^2 \right) u^2 \right. \\ &\quad \left. + \mathcal{O}(u^3) \right] + \mathcal{O}(\nu^2). \end{aligned} \quad (4.13)$$

$$g_S = R^3 G_S^{\text{phys}}, \quad g_{S*} = R R_c^2 G_{S*}^{\text{phys}} \quad (4.10)$$

where $R_c^2 = R^2 + \mathcal{O}(\text{spin}^2)$ [66]. Here, as we work linearly in spins, we can neglect the spin quadratic contribution to R_c^2 . In the following, we shall sometimes set, for simplicity, the velocity of light to 1.

B. Present knowledge of the EOB gyrogravitomagnetic ratios g_S and g_{S*}

Let us describe the present knowledge of the two (phase-space-dependent) dimensionless gyrogravitomagnetic ratios g_S and g_{S*} . First, from the PN-expanded point of view, g_S and g_{S*} are known at the next-to-next-to-leading-order (NNLO) level [63, 64]

Concerning the second gyrogravitomagnetic ratio g_{S*} , it was emphasized in Ref.[62] that the ν -independent piece of g_{S*} , $g_{S*}^{(\nu^0)}$, could be exactly determined from considering a spinning particle in an external background [61]. Taking as external background a Schwarzschild metric (consistently with our working linearly in spins) this leads to the explicit expression

$$\begin{aligned} g_{S*}^{(\nu^0)}(u, p_r, p_\phi) &= \frac{1}{1 + \sqrt{1 + p_\phi^2 u^2 + (1 - 2u)p_r^2}} \\ &\quad + \frac{1}{\sqrt{1 + p_\phi^2 u^2 + (1 - 2u)p_r^2}} \frac{2}{1 + \frac{1}{\sqrt{1 - 2u}}}. \end{aligned} \quad (4.14)$$

This exact expression for $g_{S*}^{(\nu^0)}$ has introduced an explicit dependence on p_ϕ , corresponding to being in a different spin gauge than the DJS one used in the PN-expanded expressions (4.11).

Gravitational self-force theory allowed, starting in 2014, to acquire new knowledge on spin precession in extreme-mass-ratio binaries [34, 35, 68, 72]. The knowledge acquired in the latter references was limited to the case of circular orbits and was transcribed within the EOB formalism in [34]. When considering circular orbits, it is natural to decompose g_{S*} (within self-force theory) in the following way

$$g_{S*}^{\text{circ}}(u, \nu) = g_{S*}^{\text{circ}(\nu^0)}(u) + \nu g_{S*}^{\text{circ}(\nu^1)}(u) + \mathcal{O}(\nu^2), \quad (4.15)$$

where

$$g_{S*}^{\text{circ}(\nu^0)}(u) \equiv \frac{3}{1 + \frac{1}{\sqrt{1-3u}}}, \quad (4.16)$$

is defined by replacing p_r by 0 and p_ϕ by $p_\phi^{(\text{circ})}(u) = [u(1-3u)]^{-1/2}$ in (4.14). The additional $\mathcal{O}(\nu)$ correction $\nu g_{S*}^{\text{circ}(\nu^1)}(u)$ in Eq. (4.15) has been analytically determined as a PN expansion, up to the 9.5PN level in [34, 68]. [From the analytical results of [35] one could further determine $g_{S*}^{\text{circ}(\nu^1)}(u)$ to the 23PN level]. Moreover, Ref. [34], combining analytical knowledge with a fit to numerical SF data from Ref. [72] [together with numerical SF data from [73]] derived a simple representation of $g_{S*}^{\text{circ}(\nu^1)}(u)$ as a rational function of u in the interval $0 \leq u < \frac{1}{3}$, see Eqs. (6.39)–(6.40) in Ref. [34]. Let us quote here only the first terms of the PN expansion of $g_{S*}^{\text{circ}(\nu^1)}(u)$

$$\begin{aligned} g_{S*}^{\text{circ}(\nu^1)}(u) = & -\frac{3}{4}u - \frac{39}{4}u^2 + \left(\frac{41}{32}\pi^2 - \frac{7627}{192} \right) u^3 \\ & + \left(-24\ln(u) - \frac{1017}{20} - \frac{1456}{15}\ln(2) - 48\gamma \right. \\ & \left. + \frac{23663}{2048}\pi^2 \right) u^4 + \mathcal{O}(u^5). \end{aligned} \quad (4.17)$$

See Eq. (A1) in the Appendix of Ref. [68] for the 9.5PN accurate extension of this expression.

C. Improving the analytical knowledge of g_S and

g_{S*}

In the present paper we shall improve the SF knowledge of g_{S*} by computing the $\mathcal{O}(p_r^2)$ corrections to Eq. (4.15). To this end, let us decompose g_{S*} in the following way

$$\begin{aligned} g_{S*}(u, p_r, p_\phi; \nu) = & g_{S*}^{(\nu^0)}(u, p_r, p_\phi) + \nu [g_{S*}^{\text{1SF0}}(u) \\ & + p_r^2 g_{S*}^{\text{1SF2}}(u) + \mathcal{O}(p_r^4)] + \nu^2 g_{S*}^{(\nu^2)}(u) \\ & + \mathcal{O}(\nu^2 p_r^2) + \mathcal{O}(\nu^3). \end{aligned} \quad (4.18)$$

Here $g_{S*}^{(\nu^0)}(u, p_r, p_\phi)$ is the phase-space function defined in Eq. (4.14) above, while we have written the additional $\mathcal{O}(\nu)$ and $\mathcal{O}(\nu^2)$ contributions as p_ϕ -independent functions of u and p_r , expanded in powers of p_r^2 . [We use here the freedom allowed by DJS-type spin gauge to eliminate any p_ϕ dependence in the $\mathcal{O}(\nu)$ terms.] In the following we shall use the SF result Eq. (3.33) to determine the PN expansions of $g_{S*}^{\text{1SF0}}(u)$ and $g_{S*}^{\text{1SF2}}(u)$,

namely

$$\begin{aligned} g_{S*}^{\text{1SF0}}(u) = & g_{*01}u + g_{*02}u^2 + g_{*03}u^3 + (g_{*04}^c + g_{*04}^{\text{ln}} \ln u) u^4 \\ & + \dots \\ g_{S*}^{\text{1SF2}}(u) = & g_{*20} + g_{*21}u + g_{*22}u^2 + (g_{*23}^c + g_{*23}^{\text{ln}} \ln u) u^3 \\ & + (g_{*24}^c + g_{*24}^{\text{ln}} \ln u) u^4 + \dots \\ g_{S*}^{\text{1SF4}}(u) = & g_{*40} + g_{*41}u + g_{*42}u^2 + (g_{*43}^c + g_{*43}^{\text{ln}} \ln u) u^3 \\ & + (g_{*44}^c + g_{*44}^{\text{ln}} \ln u) u^4 + \dots \end{aligned} \quad (4.19)$$

The first step towards the determination of the coefficients in Eq. (4.19) is to relate the circular limit of Eq. (4.18) to the previous circular result, Eq. (4.15). Indeed, the circular limit of (4.18) reads

$$\begin{aligned} g_{S*}(u, p_r, p_\phi) \Big|_{\text{circ}} = & g_{S*}^{(\nu^0)}(u, 0, p_\phi^{(\text{circ})}(u, \nu)) + \nu [g_{S*}^{\text{1SF0}}(u)] \\ & + \mathcal{O}(\nu^2). \end{aligned} \quad (4.20)$$

where the expression of the ν -dependent value of the square of $p_\phi^{(\text{circ})}(u, \nu)$, which is well known from EOB theory [5, 25], reads

$$\begin{aligned} [p_\phi^{(\text{circ})}(u, \nu)]^2 = & -\frac{\partial_u A(u; \nu)}{\partial_u (u^2 A(u; \nu))} \\ = & \frac{1}{u(1-3u)} \left[1 - \nu \frac{2a(u) + (1-2u)a'(u)}{2(1-3u)} \right] \\ & + \mathcal{O}(\nu^2). \end{aligned} \quad (4.21)$$

Here $A(u; \nu) = 1 - 2u + \nu a(u) + \mathcal{O}(\nu^2)$ is the ν -expansion of the main radial EOB potential $A(u; \nu) = -g_{00}^{(\text{eff})}$, see Eq. (4.6). This yields

$$p_\phi^{(\text{circ})}(u, \nu) = j_{(\text{circ})}(u) + \nu \delta j(u) + \mathcal{O}(\nu^2), \quad (4.22)$$

with

$$j_{(\text{circ})}(u) = \frac{1}{\sqrt{u(1-3u)}}, \quad \delta j(u) = -\frac{2a(u) + (1-2u)a'(u)}{4\sqrt{u(1-3u)}^{3/2}}. \quad (4.23)$$

Inserting this result in Eq. (4.20), we see that the first term on the rhs contributes an additional $\mathcal{O}(\nu)$ contribution, namely

$$\begin{aligned} g_{S*}^{(\nu^0)}(u, 0, p_\phi^{(\text{circ})}(u, \nu)) = & \frac{3}{1 + \frac{1}{\sqrt{1-3u}}} + \nu \frac{\partial g_{S*}^{(\nu^0)}(u, 0, p_\phi)}{\partial p_\phi} \Big|_{p_\phi=j_{(\text{circ})}} \delta j(u) + \mathcal{O}(\nu^2). \end{aligned} \quad (4.24)$$

This implies the following link

$$g_{S*}^{\text{circ}(\nu^1)}(u) = \frac{\partial g_{S*}^{(\nu^0)}(u, 0, p_\phi)}{\partial p_\phi} \Big|_{p_\phi=j_{(\text{circ})}} \delta j(u) + g_{S*}^{\text{1SF0}}(u), \quad (4.25)$$

which determines the value of $g_{S*}^{\text{1SF0}}(u)$ from the previously known results on $g_{S*}^{\text{circ}(\nu^1)}(u)$, given to 4PN fractional accuracy in Eq. (4.17) and to 9.5PN accuracy in

Eq. (A1) of Ref. [68]. Let us cite explicitly here only the first coefficients in the PN expansion (4.19) of $g_{S*}^{\text{1SF0}}(u)$

$$\begin{aligned} g_{*01} &= -\frac{3}{4} \\ g_{*02} &= -\frac{39}{4} \\ g_{*03} &= \frac{41}{32}\pi^2 - \frac{7987}{192} \\ g_{*04}^c &= -\frac{11447}{120} - \frac{1456}{15}\ln(2) - 48\gamma + \frac{26943}{2048}\pi^2 \\ g_{*04}^{\ln} &= -24, \end{aligned} \quad (4.26)$$

leading to

$$\begin{aligned} g_{S*}^{\text{1SF0}}(u) &= -\frac{3}{4}u - \frac{39}{4}u^2 + \left(\frac{41}{32}\pi^2 - \frac{7987}{192}\right)u^3 \\ &+ \left(-24\ln(u) - \frac{11447}{120} - \frac{1456}{15}\ln(2) - 48\gamma \right. \\ &\left. + \frac{26943}{2048}\pi^2\right)u^4 + \mathcal{O}(u^5). \end{aligned} \quad (4.27)$$

In a second step, we can determine the coefficients $g_{*20}, \dots, g_{*24}^{\ln}$ in the PN expansion, Eq.(4.19), of the $\mathcal{O}(p_r^2)$ contribution to $g_{S*}(u, p_r, p_\phi)$, Eq. (4.18). The technicalities of this determination will be explained in the next subsection. For clarity, let us quote in advance the result we shall obtain

$$\begin{aligned} g_{*20} &= -\frac{9}{4} \\ g_{*21} &= -\frac{9}{4} \\ g_{*22}^c &= -\frac{717}{32} \\ g_{*22}^{\ln} &= 0 \\ g_{*23}^c &= \frac{1447441}{960} - \frac{4829}{256}\pi^2 - \frac{16038}{5}\ln(3) + \frac{46976}{15}\ln(2) \\ &\quad - \frac{512}{5}\gamma \\ g_{*23}^{\ln} &= -\frac{256}{5} \\ g_{*24}^c &= -\frac{185195453}{38400} + \frac{19162}{35}\gamma + \frac{2097479}{8192}\pi^2 \\ &\quad + \frac{454167}{20}\ln(3) - \frac{1081966}{35}\ln(2) \\ g_{*24}^{\ln} &= +\frac{9581}{35}. \end{aligned} \quad (4.28)$$

Note that only the values of the first two coefficients were known before, see Eq. (4.11). The results for $g_{*22}, g_{*23}^c, g_{*23}^{\ln}, g_{*24}^c$ and g_{*24}^{\ln} are new with this work. At the level $\mathcal{O}(\nu, p_r^4)$, instead, the only known coefficient [63, 64] is

$$g_{*40} = \frac{5}{2}, \quad (4.29)$$

as shown in Eqs. (4.11).

D. EOB computation of Ω_r and Ω_ϕ as functions of energy and angular momentum

We have seen above how SF theory led to a determination of the functional link between the spin precession quantity ψ and the two gauge-invariant frequencies of the orbital motion, Ω_r and Ω_ϕ :

$$\begin{aligned} \psi(Gm_2\Omega_r, Gm_2\Omega_\phi; q) &= \\ \psi_0(Gm_2\Omega_r, Gm_2\Omega_\phi) &+ q\Delta\psi(Gm_2\Omega_r, Gm_2\Omega_\phi) + \mathcal{O}(q^2), \end{aligned} \quad (4.30)$$

where $q = m_1/m_2 = \nu + \mathcal{O}(\nu^2)$. In order to relate the SF result, Eq. (3.33), on $\Delta\psi(Gm_2\Omega_r, Gm_2\Omega_\phi)$ to the PN expansion of the EOB gyrogravitomagnetic ratio $g_{S*}(u, p_r, p_\phi; \nu)$, Eq. (4.18), we need, as a first task, to compute the functional link predicted by EOB theory between Ω_r and Ω_ϕ and the (gauge-invariant) total energy E_{tot} and orbital angular momentum $L = P_\phi$ of the corresponding motion of the binary system.

Having in mind the link, Eq. (4.1), between $E_{\text{tot}} = H$ and the effective energy $\mathcal{E}_{\text{eff}} = H_{\text{eff}}$ (or, equivalently, $\hat{\mathcal{E}}_{\text{eff}} = \mathcal{E}_{\text{eff}}/(\mu c^2)$ and $\hat{H}_{\text{eff}} = H_{\text{eff}}/(\mu c^2)$), together with the definition (4.9) of the rescaled angular momentum j , our first task will be to compute Ω_r and Ω_ϕ as functions of $\hat{\mathcal{E}}_{\text{eff}}$ and j .

The two frequencies we are interested in can be written as

$$\Omega_r = \frac{2\pi}{\mathcal{T}}, \quad \text{with} \quad \mathcal{T} = \oint dT, \quad (4.31)$$

and

$$\Omega_\phi = \frac{\Phi}{\mathcal{T}} = \frac{\oint d\phi}{\oint dT} = \frac{\oint \dot{\phi} dT}{\mathcal{T}}. \quad (4.32)$$

Here T denotes the physical² time (to be distinguished from the effective time T_{eff} entering Eq. (4.6)), while \oint denotes a periapsis-to-periapsis integral.

Using the rescaled quantities (4.9) and (4.10), together with the dimensionless, rescaled physical time

$$t = \frac{c^3 T}{GM}, \quad (4.33)$$

the above frequencies become

$$GM\Omega_r = \frac{2\pi}{\oint dt}, \quad GM\Omega_\phi = \frac{\oint d\phi}{\oint dt}. \quad (4.34)$$

The time integration in these integrals can be replaced by radial integration using Hamilton's equations for the rescaled radial variable

$$\frac{dr}{dt} = \frac{1}{\nu} \frac{\partial h}{\partial p_r}, \quad (4.35)$$

² T is the standard Schwarzschild-like coordinate time, as observed at infinity.

together with Hamilton's equation for the azimuthal variable

$$\frac{d\phi}{dt} = \frac{1}{\nu} \frac{\partial h}{\partial j}. \quad (4.36)$$

In order to turn Eq. (4.35) into a relation of the type

$$\hat{\mathcal{E}}_{\text{eff}} = \hat{H}_{\text{eff}}(u, p_r, j; S, S_*) = \sqrt{A(u) \left(1 + j^2 u^2 + A(u) \bar{D}(u) p_r^2 + \hat{Q}\right)} + \frac{j u^3}{M^2} (g_S S + g_{S*} S_*) \quad (4.37)$$

where we replaced the metric potential $B = g_{RR}^{\text{eff}}$, Eq. (4.6), by $\bar{D} \equiv (AB)^{-1}$.

As our aim here is to compute the coupling coefficients g_S, g_{S*} parametrizing effects linear in spins, it is easily seen that it is enough to compute Ω_r and Ω_ϕ to *zeroth* order in spins. In other words, we can neglect the spin-dependent terms in the energy conservation law (4.37) and work with the simplified mass-shell condition

$$\hat{\mathcal{E}}_{\text{eff}}^2 = A(u) \left(1 + j^2 u^2 + A(u) \bar{D}(u) p_r^2 + \hat{Q}\right) + \mathcal{O}(\text{spin}). \quad (4.38)$$

The SF expansions of the EOB potential A, \bar{D} and $\hat{Q} = Q/\mu^2$ read

$$A(u) = 1 - 2u + \nu a(u) + \mathcal{O}(\nu^2),$$

$dt = f(r)dr$ we need the explicit expression of the (rescaled) radial momentum p_r as a function of r . The latter relation is obtained by writing the law of conservation of energy

$$\begin{aligned} \bar{D}(u) &= 1 + \nu \bar{d}(u) + \mathcal{O}(\nu^2), \\ \hat{Q}(u) &= \nu q_4(u) p_r^4 + \nu q_6(u) p_r^6 + \nu q_8(u) p_r^8 + \mathcal{O}(\nu^2, p_r^{10}). \end{aligned} \quad (4.39)$$

The PN expansions of the first SF-order radial functions $a(u), \bar{d}(u), q_4(u), \dots$ entering the latter equations have been determined by SF theory to very high PN-orders: see Refs. [33, 35, 49, 74–77] for $a(u)$, Refs. [14] for $\bar{d}(u)$ and Refs. [14, 15, 59] for $q_4(u), q_6(u)$ etc...

For concreteness, let us quote here the beginning of these expansions

$$\begin{aligned} A(u) &= 1 - 2u + 2\nu u^3 + \left(\frac{94}{3} - \frac{41}{32}\pi^2\right) \nu u^4 \\ &\quad + \left[\left(\frac{2275}{512}\pi^2 - \frac{4237}{60} + \frac{128}{5}\gamma + \frac{256}{5}\ln(2)\right) \nu + \left(\frac{41}{32}\pi^2 - \frac{221}{6}\right) \nu^2 + \frac{64}{5}\nu \ln(u)\right] u^5 + \mathcal{O}(u^6) \\ \bar{D}(u) &= 1 + 6\nu u^2 + (52\nu - 6\nu^2)u^3 + \left[\left(-\frac{533}{45} - \frac{23761}{1536}\pi^2 + \frac{1184}{15}\gamma - \frac{6496}{15}\ln 2 + \frac{2916}{5}\ln 3\right) \nu \right. \\ &\quad \left. + \left(\frac{123}{16}\pi^2 - 260\right) \nu^2 + \frac{592}{15}\nu \ln(u)\right] u^4 + \mathcal{O}(u^5) \\ \hat{Q}(u, p_r) &= \left[2(4 - 3\nu) \nu u^2 + \left(\left(-\frac{5308}{15} + \frac{496256}{45}\ln 2 - \frac{33048}{5}\ln 3\right) \nu - 83\nu^2 + 10\nu^3\right) u^3 + \mathcal{O}(u^4)\right] p_r^4 \\ &\quad + \left[\left(-\frac{827}{3} - \frac{2358912}{25}\ln 2 + \frac{1399437}{50}\ln 3 + \frac{390625}{18}\ln 5\right) \nu - \frac{27}{5}\nu^2 + 6\nu^3\right] u^2 p_r^6 + \mathcal{O}(u^3, p_r^8). \end{aligned} \quad (4.40)$$

Inserting the SF expansion, Eqs. (4.39), into the mass-shell condition allows us to compute the functional dependence of p_r on $u, \hat{\mathcal{E}}_{\text{eff}}$ and j . To the first SF-order, i.e.,

$$p_r(u, \hat{\mathcal{E}}_{\text{eff}}, j; S, S_*) = p_r^{(0)}(u, \hat{\mathcal{E}}_{\text{eff}}, j) + \nu p_r^{(1)}(u, \hat{\mathcal{E}}_{\text{eff}}, j) + \mathcal{O}(\nu^2, \text{spin}) \quad (4.41)$$

the explicit expressions of $p_r^{(0)}$ and $p_r^{(1)}$ read (when neglecting spins)

$$p_r^{(0)}(u, \hat{\mathcal{E}}_{\text{eff}}, j) = \frac{\sqrt{\hat{\mathcal{E}}_{\text{eff}}^2 - (1-2u)(1+u^2j^2)}}{1-2u}$$

$$p_r^{(1)}(u, \hat{\mathcal{E}}_{\text{eff}}, j) = -\frac{\hat{\mathcal{E}}_{\text{eff}}^2}{p_r^{(0)}} \frac{a(u)}{(1-2u)^3} - p_r^{(0)} \left[\bar{d}(u) + \frac{a(u)}{(1-2u)} + (p_r^{(0)})^2 \frac{q_4(u) + q_6(u)(p_r^{(0)})^2 + q_8(u)(p_r^{(0)})^4 + \dots}{1-2u} \right]. \quad (4.42)$$

We can then obtain an explicit expression for replacing time integration by radial integration by inserting Eqs. (4.41) and (4.42) in the relation

$$dt = \frac{\sqrt{1 + 2\nu(\hat{\mathcal{E}}_{\text{eff}} - 1)\hat{\mathcal{E}}_{\text{eff}}}}{Ap_r(A\bar{D} + 2p_r^2\nu q_4 + 3p_r^4\nu q_6 + 4p_r^6\nu q_8 + \dots)} dr, \quad (4.43)$$

which follows from the radial equation of motion (4.35). Inserting Eq. (4.43) in the t -integral expressions for Ω_r and Ω_ϕ , Eqs. (4.35) and (4.36), and formally expanding p_r in powers of ν according to Eqs. (4.41) and (4.42), leads to r -integral expressions for the frequencies of the type

$$GM\Omega_{r,\phi} \sim \sum_n \int_{u_{(\min)}}^{u_{(\max)}} du \frac{f_n(u)}{\left(\hat{\mathcal{E}}_{\text{eff}}^2 - (1-2u)(1+u^2j^2)\right)^{n+\frac{1}{2}}} \quad (4.44)$$

where $f_n(u)$ involves a combination of $(1-2u)^k$, $a(u)$, $\bar{d}(u)$, etc. Note that these radial integrals are divergent for $n \geq 1$. The appearance of singular integral is due to our formal replacement of the expansion (4.41) in originally convergent integrals of the type $\oint dr f(r)/p_r$. As shown in Ref. [78], the correct result for these expansions is obtained simply by taking Hadamard's partie finie (Pf) of the singular integrals, Eq. (4.44).

We are interested here in computing the PN-expansions of the frequencies. These PN-expansions can be conveniently obtained: i) by replacing the various first SF-order EOB potentials $a(u)$, $\bar{d}(u)$, etc., by their PN expansion illustrated in Eq. (4.40); and, using again the general result of Ref. [78], ii) by formally expanding the radicals entering the denominators above, namely

$$\hat{\mathcal{E}}_{\text{eff}}^2 - (1-2u)(1+u^2j^2) = \hat{\mathcal{E}}_{\text{eff}}^2 - 1 + 2u - j^2u^2 + 2j^2u^3, \quad (4.45)$$

around the Newtonian-like, quadratic radical

$$\mathcal{R}_0(u) = \hat{\mathcal{E}}_{\text{eff}}^2 - 1 + 2u - j^2u^2. \quad (4.46)$$

In other words, the combination of the ν -expansion and the PN-expansion leads to integral expressions for the

frequencies of the type

$$GM\Omega_{r,\phi} \sim \sum_n \text{Pf} \int_{u_{(\min)}}^{u_{(\max)}} du \frac{\tilde{f}_n(u)}{\mathcal{R}_0^{n+\frac{1}{2}}}(u). \quad (4.47)$$

Here the symbol Pf denotes Hadamard's partie finie. Note that the values of the end points of the u -integrations are different from the ones in Eq. (4.44) above and now denote the two roots of the Newtonian-like quadratic radical $\mathcal{R}_0(u)$, Eq. (4.46). Finally, we are left with evaluating Newtonian-like radial integrals of the type (4.47). When evaluating the partie finie of these singular integrals it is convenient to replace $\hat{\mathcal{E}}_{\text{eff}}$ and j by the quantities³ u_p and e defined so that the two roots of $\mathcal{R}_0(u)$ are $u_p(1 \pm e)$, namely

$$\hat{\mathcal{E}}_{\text{eff}}^2 - 1 = -u_p(1 - e^2), \quad j^2 = \frac{1}{u_p}. \quad (4.48)$$

As we are interested in slightly eccentric⁴ motions, we can further expand the latter (singular) Newtonian-like integrals in powers of e .

When doing so, we use the following Newtonian-like parametrization of the inverse radius $u = 1/r$

$$u = u_p(1 + e \cos \chi_0), \quad (4.49)$$

so that

$$du = -u_p e \sin \chi_0 d\chi_0, \quad \mathcal{R}_0(u) = u_p e^2 \sin^2 \chi_0. \quad (4.50)$$

We have then shown that the e -expansion of the partie finie of the integrals (4.47) is correctly obtained by taking the ϵ^0 term in the Laurent expansion in ϵ of the integrals of the type

$$\int_{\epsilon}^{\pi-\epsilon} \frac{g_n(\chi_0)}{\sin^{2n} \chi_0} d\chi_0, \quad (4.51)$$

that are generated by the expansions (4.47).

Finally, the combined PN-, SF- and eccentricity-expansions of the frequencies yield

$$GM\Omega_r = \Omega_r^{(0)}(u_p, \nu) + e^2 \Omega_r^{(2)}(u_p, \nu) + e^4 \Omega_r^{(4)}(u_p, \nu) + \mathcal{O}(e^6), \quad (4.52)$$

with

³ As said above, the so-defined EOB variables $p \equiv 1/u_p$ and e

$$\begin{aligned}
\Omega_r^{(0)}(u, \nu) &= u_p^{3/2} + \left(\frac{1}{2}\nu - \frac{3}{2}\right) u_p^{5/2} + \left(\frac{19}{8}\nu - \frac{33}{8}\right) u_p^{7/2} + \left[-\frac{523}{16} + \left(-\frac{41}{64}\pi^2 + \frac{2591}{48}\right)\nu\right] u_p^{9/2} \\
&+ \left[-\frac{40389}{128} + \left(\frac{692771}{1152} - \frac{26735}{1536}\pi^2 + \frac{1168}{15}\gamma + \frac{584}{15}\ln(u_p) + \frac{1458}{5}\ln(3) - \frac{2096}{15}\ln(2)\right)\nu\right] u_p^{11/2} \\
&+ \left[-\frac{849573}{256} + \left(\frac{50524}{105}\gamma - \frac{611631}{70}\ln(3) + \frac{13140979}{2304} + \frac{25262}{105}\ln(u_p) + \frac{255236}{15}\ln(2) + \frac{481}{96}\pi^2\right)\nu\right] u_p^{13/2} \\
&+ \frac{11770}{63} u_p^7 \pi \nu + \mathcal{O}(u^{15/2}, \nu^2) \\
\Omega_r^{(2)}(u_p, \nu) &= -\frac{3}{2} u_p^{3/2} + \left(-\frac{5}{4}\nu + \frac{15}{4}\right) u_p^{5/2} + \left(-\frac{109}{16}\nu + \frac{171}{16}\right) u_p^{7/2} + \left[\frac{2607}{32} + \left(\frac{123}{64}\pi^2 - \frac{4609}{32}\right)\nu\right] u_p^{9/2} \\
&+ \left[\frac{186639}{256} + \left(-\frac{3208}{15}\gamma - \frac{1604}{15}\ln(u_p) - \frac{28431}{5}\ln(3) + \frac{128728}{15}\ln(2) + \frac{273443}{6144}\pi^2 - \frac{18601901}{11520}\right)\nu\right] u_p^{11/2} \\
&+ \left[\frac{3564729}{512} + \left(\frac{26578611}{224}\ln(3) - \frac{1657307}{6144}\pi^2 - \frac{92786}{105}\gamma - \frac{2905678}{7}\ln(2) - \frac{46393}{105}\ln(u_p) \right. \right. \\
&\quad \left. \left. + \frac{21484375}{224}\ln(5) - \frac{8195349553}{806400}\right)\nu\right] u_p^{13/2} - \frac{92341}{315} u_p^7 \pi \nu + \mathcal{O}(u_p^{15/2}, \nu^2) \\
\Omega_r^{(4)}(u_p, \nu) &= \frac{3}{8} u_p^{3/2} + \left(-\frac{45}{16} + \frac{15}{16}\nu\right) u_p^{5/2} + \left(\frac{401}{64}\nu - \frac{495}{64}\right) u_p^{7/2} + \left[-\frac{6489}{128} + \left(-\frac{123}{64}\pi^2 + \frac{13927}{128}\right)\nu\right] u_p^{9/2} \\
&+ \left[-\frac{333831}{1024} + \left(\frac{5218813}{5120} + \frac{436}{5}\ln(u_p) - \frac{566832}{5}\ln(2) + \frac{872}{5}\gamma - \frac{59563}{2048}\pi^2 + \frac{1953125}{96}\ln(5) \right. \right. \\
&\quad \left. \left. + \frac{6647751}{160}\ln(3)\right)\nu\right] u_p^{11/2} \\
&+ \left[-\frac{3194019}{2048} + \left(\frac{451817433}{4480}\ln(3) - \frac{30997}{420}\ln(u_p) - \frac{4017578125}{2688}\ln(5) + \frac{99386957}{30}\ln(2) \right. \right. \\
&\quad \left. \left. - \frac{30997}{210}\gamma + \frac{18324577}{24576}\pi^2 - \frac{24403514197}{3225600}\right)\nu\right] u_p^{13/2} - \frac{11660111}{53760} u_p^7 \pi \nu + \mathcal{O}(u^{15/2}, \nu^2). \tag{4.53}
\end{aligned}$$

Note that the dependence on $\ln u_p$ starts at $\mathcal{O}(u_p^{11/2})$. Similarly, the azimuthal frequency reads

$$GM\Omega_\phi = \Omega_\phi^{(0)}(u_p, \nu) + e^2 \Omega_\phi^{(2)}(u_p, \nu) + e^4 \Omega_\phi^{(4)}(u_p, \nu) + \mathcal{O}(e^6), \tag{4.54}$$

with, for example,

$$\begin{aligned}
\Omega_\phi^{(0)}(u_p, \nu) &= u_p^{3/2} + \left(\frac{3}{2} + \frac{1}{2}\nu\right) u_p^{5/2} + \left(-\frac{17}{8}\nu + \frac{111}{8}\right) u_p^{7/2} + \left[\frac{2099}{16} + \left(-\frac{5935}{48} + \frac{205}{64}\pi^2\right)\nu\right] u_p^{9/2} \\
&+ \left[\frac{172059}{128} + \left(-\frac{3557387}{1920} + \frac{38195}{1024}\pi^2 - \frac{448}{5}\gamma - \frac{224}{5}\ln(u_p) - \frac{896}{5}\ln(2)\right)\nu\right] u_p^{11/2} \\
&+ \left[\frac{3720501}{256} + \left(-\frac{33940}{21}\gamma - \frac{174231}{70}\ln(3) - \frac{1332163499}{57600} - \frac{16970}{21}\ln(u_p) - \frac{74468}{105}\ln(2) \right. \right. \\
&\quad \left. \left. + \frac{291883}{1536}\pi^2\right)\nu\right] u_p^{13/2} - \frac{13696}{105} u_p^7 \pi \nu + \mathcal{O}(u_p^{15/2}, \nu^2) \\
\Omega_\phi^{(2)}(u_p, \nu) &= -\frac{3}{2} u_p^{3/2} + \left(-\frac{3}{4} - \frac{5}{4}\nu\right) u_p^{5/2} + \left(-\frac{129}{16} - \frac{49}{16}\nu\right) u_p^{7/2} + \left[-\frac{1407}{32} + \left(-\frac{369}{128}\pi^2 - \frac{303}{32}\right)\nu\right] u_p^{9/2} \\
&+ \left[-\frac{20793}{256} + \left(-\frac{2656}{15}\gamma - \frac{1328}{15}\ln(u_p) - \frac{5832}{5}\ln(3) + \frac{12416}{15}\ln(2) + \frac{148949}{6144}\pi^2 - \frac{6765029}{11520}\right)\nu\right] u_p^{11/2}
\end{aligned}$$

should be distinguished from the SF-defined variables \bar{p} and \bar{e} used as independent variables in Eq. (3.33) above.

⁴ The parameter e defined in Eq. (4.48) does not measure the exact eccentricity of the corresponding EOB orbits that would vanish

when p_r vanishes. Nevertheless, when used in PN-expansion, it will allow us to correctly evaluate the gauge-invariant quantities we are interested in.

$$\begin{aligned}
& + \left[\frac{1631139}{512} + \left(\frac{788049}{20} \ln(3) - \frac{15875}{1536} \pi^2 - \frac{100046}{105} \gamma - \frac{1566262}{21} \ln(2) - \frac{50023}{105} \ln(u_p) \right. \right. \\
& \quad \left. \left. - \frac{7980766891}{806400} \right) \nu \right] u_p^{13/2} - \frac{262043}{315} \pi \nu u_p^7 + \mathcal{O}(u_p^{15/2}, \nu^2) \\
\Omega_\phi^{(4)}(u_p, \nu) = & \frac{3}{8} u_p^{3/2} + \left(-\frac{27}{16} + \frac{15}{16} \nu \right) u_p^{5/2} + \left(\frac{581}{64} \nu - \frac{855}{64} \right) u_p^{7/2} + \left[-\frac{20799}{128} + \left(-\frac{123}{64} \pi^2 + \frac{30193}{128} \right) \nu \right] u_p^{9/2} \\
& + \left[-\frac{2037927}{1024} + \left(\frac{4235449}{1024} + \frac{1046}{5} \ln(u_p) - 19324 \ln(2) + \frac{2092}{5} \gamma - \frac{395017}{4096} \pi^2 + \frac{63423}{5} \ln(3) \right) \nu \right] u_p^{11/2} \\
& + \left[-\frac{49225221}{2048} + \left(-\frac{1153657809}{4480} \ln(3) + \frac{655871}{420} \ln(u_p) - \frac{654296875}{2688} \ln(5) + \frac{29552521}{30} \ln(2) \right. \right. \\
& \quad \left. \left. + \frac{655871}{210} \gamma + \frac{4564433}{12288} \pi^2 + \frac{130198554749}{3225600} \right) \nu \right] u_p^{13/2} + \frac{21293}{28} u_p^7 \pi \nu + \mathcal{O}(u_p^{15/2}, \nu^2). \tag{4.55}
\end{aligned}$$

Note again that the dependence on $\ln u_p$ starts at $\mathcal{O}(u_p^{11/2})$.

In the expansions (4.53) and (4.55) above we have only displayed the beginning of the PN expansions of $\Omega_r^{(0,2,4)}(u_p, \nu)$ and $\Omega_\phi^{(0,2,4)}(u_p, \nu)$. Actually we have computed them to the highest PN order currently known from SF calculations. Similarly, we have also computed the eccentricity expansions (4.52) up to $\mathcal{O}(e^{10})$.

It will be convenient in the following to trade the two dimensionless frequencies $GM\Omega_r$ and $GM\Omega_\phi$ by two other related dimensionless (equally gauge-invariant) parameters, namely the (fractional) periastron advance per orbit

$$k \equiv \frac{\Omega_\phi}{\Omega_r} - 1 = \frac{\Phi}{2\pi} - 1, \tag{4.56}$$

and the dimensionless azimuthal frequency variable

$$y \equiv (Gm_2\Omega_\phi)^{2/3} = \left(1 - \frac{2}{3} \nu + \mathcal{O}(\nu^2) \right) (GM\Omega_\phi)^{2/3}. \tag{4.57}$$

The combined eccentricity-, PN- and SF-expansions of these quantities read

$$k(u_p, e; \nu) = k^{(0)}(u_p, \nu) + e^2 k^{(2)}(u_p, \nu) + e^4 k^{(4)}(u_p, \nu) + \mathcal{O}(e^6), \tag{4.58}$$

with

$$\begin{aligned}
k^{(0)}(u_p, \nu) = & 3u_p + \left(\frac{45}{2} - 6\nu \right) u_p^2 + \left[210 + \left(-205 + \frac{123}{32} \pi^2 \right) \nu \right] u_p^3 \\
& + \left[\frac{17325}{8} + \left(-\frac{2512}{15} \gamma + \frac{191671}{3072} \pi^2 - \frac{592}{15} \ln(2) - \frac{1458}{5} \ln(3) - \frac{1256}{15} \ln(u_p) - \frac{559223}{180} \right) \nu \right] u_p^4 \\
& + \left[\frac{189189}{8} + \left(-\frac{364664}{21} \ln(2) + \frac{739519}{2048} \pi^2 - \frac{45188}{35} \ln(u_p) + \frac{172773}{35} \ln(3) - \frac{90376}{35} \gamma - \frac{7826703}{200} \right) \nu \right] u_p^5 \\
& - \frac{99938}{315} \pi \nu u_p^{11/2} + \mathcal{O}(u_p^6, \nu^2) \\
k^{(2)}(u_p, \nu) = & \left(-\frac{3}{2} \nu + \frac{15}{4} \right) u_p^2 + \left[\frac{315}{4} + \left(-106 + \frac{123}{128} \pi^2 \right) \nu \right] u_p^3 \\
& + \left[\frac{10395}{8} + \left(-\frac{519697}{240} - \frac{23440}{3} \ln(2) - \frac{1072}{5} \gamma + \frac{111797}{2048} \pi^2 + \frac{20412}{5} \ln(3) - \frac{536}{5} \ln(u_p) \right) \nu \right] u_p^4 \\
& + \left[\frac{315315}{16} + \left(\frac{1791937}{4096} \pi^2 - \frac{7486}{5} \ln(u_p) - \frac{14661947}{400} + \frac{29100532}{105} \ln(2) - \frac{14972}{5} \gamma - \frac{7611489}{160} \ln(3) \right. \right. \\
& \quad \left. \left. - \frac{21484375}{224} \ln(5) \right) \nu \right] u_p^5 \\
& - \frac{319609}{315} \pi \nu u_p^{11/2} + \mathcal{O}(u_p^6, \nu^2) \\
k^{(4)}(u_p, \nu) = & -\frac{3}{2} u_p^3 \nu
\end{aligned}$$

$$\begin{aligned}
& + \left[\frac{3465}{64} + \left(\frac{411686}{5} \ln(2) - \frac{3620943}{160} \ln(3) - \frac{37}{5} \ln(u_p) - \frac{74}{5} \gamma + \frac{13529}{48} - \frac{1953125}{96} \ln(5) + \frac{33601}{8192} \pi^2 \right) \nu \right] u_p^4 \\
& + \left[\frac{135135}{64} + \left(\frac{455078125}{448} \ln(5) - \frac{16271}{70} \ln(u_p) - \frac{16271}{35} \gamma - \frac{1358100027}{2240} \ln(3) + \frac{4363277}{16384} \pi^2 \right. \right. \\
& \quad \left. \left. - \frac{148342613}{105} \ln(2) + \frac{10223249}{16800} \right) \nu \right] u_p^5 \\
& - \frac{1089581}{2560} \pi \nu u_p^{11/2} + \mathcal{O}(u_p^6, \nu^2). \tag{4.59}
\end{aligned}$$

In the following it will be convenient to work with the rescaled periastron advance per orbit \hat{k} , defined by

$$\hat{k} \equiv \frac{k}{3}. \tag{4.60}$$

Similarly

$$y(u_p, e; \nu) = y^{(0)}(u_p, \nu) + e^2 y^{(2)}(u_p, \nu) + e^4 y^{(4)}(u_p, \nu) + \mathcal{O}(e^6), \tag{4.61}$$

with

$$\begin{aligned}
y^{(0)}(u_p, \nu) &= \left(1 - \frac{2}{3} \nu \right) u_p + \left(-\frac{1}{3} \nu + 1 \right) u_p^2 + \left(9 - \frac{91}{12} \nu \right) u_p^3 + \left[83 + \left(-\frac{9967}{72} + \frac{205}{96} \pi^2 \right) \nu \right] u_p^4 \\
&+ \left[\frac{1671}{2} + \left(-\frac{1792}{15} \ln(2) + \frac{12185}{512} \pi^2 - \frac{896}{15} \gamma - \frac{1686559}{960} - \frac{448}{15} \ln(u_p) \right) \nu \right] u_p^5 \\
&+ \left[\frac{17835}{2} + \left(-\frac{164996}{315} \ln(u_p) + \frac{971767}{9216} \pi^2 - \frac{26024}{63} \ln(2) - \frac{329992}{315} \gamma - \frac{58077}{35} \ln(3) \right. \right. \\
&\quad \left. \left. - \frac{1768379519}{86400} \right) \nu \right] u_p^6 - \frac{27392}{315} \pi \nu u_p^{13/2} + \mathcal{O}(u_p^6, \nu^2) \\
y^{(2)}(u_p, \nu) &= \left(\frac{2}{3} \nu - 1 \right) u_p - \frac{2}{3} u_p^2 \nu + \left(-1 - \frac{23}{12} \nu \right) u_p^3 + \left[\frac{21}{2} + \left(-\frac{41}{48} \pi^2 - \frac{457}{9} \right) \nu \right] u_p^4 \\
&+ \left[\frac{667}{2} + \left(-\frac{1967801}{1728} + \frac{128815}{4608} \pi^2 - \frac{3328}{45} \ln(u_p) + \frac{22144}{45} \ln(2) - \frac{6656}{45} \gamma - \frac{3888}{5} \ln(3) \right) \nu \right] u_p^5 \\
&+ \left[6043 + \left(-\frac{4996270991}{302400} + \frac{180792}{7} \ln(3) - \frac{168544}{315} \ln(u_p) - \frac{15795776}{315} \ln(2) - \frac{337088}{315} \gamma \right. \right. \\
&\quad \left. \left. + \frac{121213}{4608} \pi^2 \right) \nu \right] u_p^6 - \frac{565174}{945} \pi \nu u_p^{13/2} + \mathcal{O}(u_p^6, \nu^2) \\
y^{(4)}(u_p, \nu) &= (-1 + \nu) u_p^2 + \left(-\frac{63}{8} + \frac{119}{12} \nu \right) u_p^3 + \left[-\frac{343}{4} + \left(\frac{1443}{8} - \frac{369}{256} \pi^2 \right) \nu \right] u_p^4 \\
&+ \left[-\frac{7687}{8} + \left(\frac{11677919}{4320} + \frac{8888}{45} \gamma + \frac{4444}{45} \ln(u_p) + \frac{40338}{5} \ln(3) - \frac{1686137}{36864} \pi^2 - \frac{113864}{9} \ln(2) \right) \nu \right] u_p^5 \\
&+ \left[-\frac{42707}{4} + \left(\frac{1620873859}{67200} - \frac{72938151}{448} \ln(3) + \frac{7624295}{24576} \pi^2 - \frac{654296875}{4032} \ln(5) + \frac{200895602}{315} \ln(2) \right. \right. \\
&\quad \left. \left. + \frac{70123}{105} \ln(u_p) + \frac{140246}{105} \gamma \right) \nu \right] u_p^6 + \frac{372467}{1890} \pi \nu u_p^{13/2} + \mathcal{O}(u_p^6, \nu^2). \tag{4.62}
\end{aligned}$$

Finally, we will need in the following to invert the functional link between (u_p, e) and (\hat{k}, y) , i.e., to compute the functions

$$u_p = f_{u_p}(\hat{k}, y), \quad e^2 = f_{e^2}(\hat{k}, y). \tag{4.63}$$

This inversion requires some care, because the Jacobian $\partial(\hat{k}, y)/\partial(u_p, e^2)$ is of order u_p near the origin of the u_p ,

e^2 plane. If we provisionally introduce the quantity $\epsilon \equiv e^2 u_p$, the Jacobian $\partial(\hat{k}, y)/\partial(u_p, \epsilon)$ will be of order unity near the origin of the u_p, ϵ plane. This shows that we can invert the link $X^i = (u_p, \epsilon) \rightarrow Y^i = (\hat{k}, y)$ by standard Taylor expansions of the symbolic type

$$X^i = A^i_j Y^j + A^i_{jk} Y^j Y^k + \dots \tag{4.64}$$

When going back from the pair $(u_p, \epsilon \equiv e^2 u_p)$ to the orig-

inal pair (u_p, e^2) we obtain the following transformation

$$\begin{aligned}
u_p &= \hat{k} + \left[\left(\frac{5}{2} + \frac{1}{3} \frac{y}{\hat{k}} \right) \nu - \frac{35}{4} + \frac{5}{4} \frac{y}{\hat{k}} \right] \hat{k}^2 \\
&\quad + \left[\left(\frac{41}{128} \pi^2 \frac{y}{\hat{k}} - \frac{151}{8} \frac{y}{\hat{k}} + \frac{25}{12} \left(\frac{y}{\hat{k}} \right)^2 + \frac{50}{3} - \frac{205}{128} \pi^2 \right) \nu - \frac{145}{16} \frac{y}{\hat{k}} + \frac{45}{16} \left(\frac{y}{\hat{k}} \right)^2 + \frac{455}{8} \right] \hat{k}^3 + \mathcal{O}(\hat{k}^4, y^3) \\
e^2 &= 1 - \left(1 + \frac{2}{3} \nu \right) \frac{y}{\hat{k}} + \left[-\frac{27}{4} \frac{y}{\hat{k}} + \frac{1}{4} \left(\frac{y}{\hat{k}} \right)^2 + \left(-2 \frac{y}{\hat{k}} + \frac{1}{6} \left(\frac{y}{\hat{k}} \right)^2 \right) \nu \right] \hat{k} \\
&\quad + \left[\left(-\frac{45}{8} \frac{y}{\hat{k}} + \frac{7}{8} \left(\frac{y}{\hat{k}} \right)^2 + \frac{15}{16} \left(\frac{y}{\hat{k}} \right)^3 - \frac{5}{16} \right) \right. \\
&\quad \left. + \left(-\frac{205}{128} \pi^2 \frac{y}{\hat{k}} + \frac{41}{128} \pi^2 \left(\frac{y}{\hat{k}} \right)^2 + \frac{1225}{24} \frac{y}{\hat{k}} - \frac{473}{24} \left(\frac{y}{\hat{k}} \right)^2 + \frac{9}{4} \left(\frac{y}{\hat{k}} \right)^3 + \frac{1}{8} \right) \nu \right] \hat{k}^2 + \mathcal{O}(k^3, y^4). \tag{4.65}
\end{aligned}$$

Again we have only indicated, for concreteness, the beginning of these expansions.

E. EOB computation of the spin precession frequency

When considering, as we do here, spin couplings to linear order, i.e., an Hamiltonian of the form $H = H_{\text{orbital}} + \mathbf{\Omega}_{S_1} \cdot \mathbf{S}_1 + \mathbf{\Omega}_{S_2} \cdot \mathbf{S}_2$, Hamilton's equations of motion for the spins ($a = 1, 2$)

$$\frac{d\mathbf{S}_a}{dt} = \{\mathbf{S}_a, H\} \tag{4.66}$$

yields

$$\frac{d\mathbf{S}_a}{dt} = \mathbf{\Omega}_{S_a} \times \mathbf{S}_a, \tag{4.67}$$

showing that $\mathbf{\Omega}_{S_a} = \frac{\partial H}{\partial \mathbf{S}_a}$ is the vectorial precession frequency of \mathbf{S}_a . Restricting to the case of interest of parallel spins we conclude that the (algebraic) magnitude of the spin frequency of body 1 is given by

$$\Omega_{S_1} = \frac{\partial H}{\partial S_1} = \frac{M\nu}{h} \frac{\partial \hat{H}_{\text{eff}}}{\partial S_1} = \left(\frac{1}{GM} \right) \frac{\nu}{h} j u^3 \left(g_S + g_{S*} \frac{m_2}{m_1} \right). \tag{4.68}$$

At first order in ν , this reads

$$\begin{aligned}
GM\Omega_{S_1}(u, p_r, j; \nu) &= j(1 - \nu)[1 - \nu(\hat{\mathcal{E}}_{\text{eff}} - 1)]u^3[\nu g_S + (1 - \nu)g_{S*}] + \mathcal{O}(\nu^2) \\
&= j(1 - \nu\hat{\mathcal{E}}_{\text{eff}})u^3[\nu g_S + (1 - \nu)g_{S*}] + \mathcal{O}(\nu^2). \tag{4.69}
\end{aligned}$$

The averaged spin frequency is then given by

$$\langle \Omega_{S_1} \rangle(\hat{\mathcal{E}}_{\text{eff}}, j; \nu) = \frac{1}{T} \oint \Omega_{S_1} dT = \frac{\Omega_r}{2\pi} \oint \Omega_{S_1} dT. \tag{4.70}$$

As indicated, the averaged spin frequency $\langle \Omega_{S_1} \rangle$ is a function of the conserved dynamical quantities, $\hat{\mathcal{E}}_{\text{eff}}$ and j . Replacing as above the T -integration by a radial integration, say $dT = f(r)dr = \tilde{f}(u)du$, see Eq. (4.43), we see that the computation of $\langle \Omega_{S_1} \rangle$ amounts to computing radial integrals of the type

$$\oint du \tilde{f}(u) u^3 [\nu g_S + (1 - \nu)g_{S*}(u, p_r, j; \nu)]. \tag{4.71}$$

In the radial integral involving g_S we can simply replace $g_S = 2$ because of the ν prefactor. By contrast, in the radial integral involving g_{S*} we need to insert the expression (4.18) which involves undetermined coefficients at the p_r^2 and p_r^4 levels, such as g_{*22}, g_{*23}^c , etc. This radial integral can be computed by the same technique we used above for computing Ω_r and Ω_ϕ . Replacing as above $\hat{\mathcal{E}}_{\text{eff}}$ and j by the Newtonian-like quantities u_p and e^2 , defined by Eqs. (4.49), we found

$$\begin{aligned}
GM\langle \Omega_{S_1} \rangle(u_p, e^2; \nu) &= \\
&\Omega_{S_1}^{(0)}(u_p, \nu) + e^2 \Omega_{S_1}^{(2)}(u_p, \nu) + e^4 \Omega_{S_1}^{(4)}(u_p, \nu) + \mathcal{O}(e^6), \tag{4.72}
\end{aligned}$$

with, for example,

$$\begin{aligned}
\Omega_{S_1}^{(0)}(u_p, \nu) &= \left(\frac{3}{2} - \nu\right) u_p^{5/2} + \left(\frac{63}{8} - \frac{15}{4}\nu\right) u_p^{7/2} + \left(-\frac{207}{4}\nu + \frac{1131}{16}\right) u_p^{9/2} \\
&\quad + \left(-\frac{174979}{192}\nu + \frac{1271}{128}\nu\pi^2 + \frac{90939}{128}\right) u_p^{11/2} \\
&\quad + \left(\frac{387023}{2048}\nu\pi^2 - \frac{8368}{15}\nu\ln(2) - \frac{1392}{5}\nu\gamma - \frac{544259}{40}\nu + \nu g_{*22} + \frac{1946997}{256} - \frac{696}{5}\nu\ln(u_p)\right) u_p^{13/2} \\
&\quad + \mathcal{O}(u_p^7, \nu^2) \\
\Omega_{S_1}^{(2)}(u_p, \nu) &= \left(-\frac{9}{4} + \frac{3}{2}\nu\right) u_p^{5/2} + \left(-\frac{105}{16} + \frac{15}{8}\nu\right) u_p^{7/2} + \left(\frac{135}{16}\nu - \frac{1167}{32}\right) u_p^{9/2} \\
&\quad + \left(\frac{1}{2}\nu g_{*22} - \frac{38361}{256} - \frac{21319}{128}\nu - \frac{615}{256}\nu\pi^2\right) u_p^{11/2} \\
&\quad + \left(8\nu g_{*22} + \frac{636407}{4096}\nu\pi^2 - \frac{1708303}{256}\nu - 696\nu\gamma + \frac{3}{2}\nu g_{*41} + \frac{1}{2}\nu g_{*23}^c - 348\nu\ln(u_p) + \frac{1}{2}\nu g_{*23}^{\ln}\ln(u_p)\right. \\
&\quad \left.+ \frac{331875}{512} - \frac{13122}{5}\nu\ln(3) + \frac{18952}{15}\nu\ln(2)\right) u_p^{13/2} + \mathcal{O}(u_p^7, \nu^2) \\
\Omega_{S_1}^{(4)}(u_p, \nu) &= \left(\frac{9}{16} - \frac{3}{8}\nu\right) u_p^{5/2} + \left(-\frac{315}{64} + \frac{135}{32}\nu\right) u_p^{7/2} + \left(\frac{1281}{16}\nu - \frac{9255}{128}\right) u_p^{9/2} \\
&\quad + \left[-\frac{971463}{1024} + \left(\frac{437543}{256} - \frac{3813}{256}\pi^2 - \frac{3}{8}g_{*22}\right)\nu\right] u_p^{11/2} \\
&\quad + \left[-\frac{24272901}{2048} + \left(\frac{224683}{8} - \frac{7}{2}g_{*22} + \frac{9}{2}g_{*41} + \frac{3}{8}g_{*42}^{\ln}\ln(u_p) + \frac{6642}{5}\gamma - \frac{187742}{5}\ln(2) - \frac{3855321}{8192}\pi^2\right.\right. \\
&\quad \left.+ \frac{3321}{5}\ln(u_p) + \frac{505197}{20}\ln(3) + \frac{3}{8}g_{*42}^c + \frac{7}{16}g_{*23}^{\ln}\right)\nu\left. \right] u_p^{13/2} + \mathcal{O}(u_p^7, \nu^2). \tag{4.73}
\end{aligned}$$

As above, we have only indicated here the first terms in the PN-expansions we computed.

Let us now consider the ratio

$$\psi \equiv \frac{\langle \Omega_{S_1} \rangle}{\Omega_\phi}. \tag{4.74}$$

It is easily checked (see, e.g., Refs. [34, 72]) that the

so-defined ratio is identical to the spin-precession measure ψ introduced in Eq. (2.1) above. The combined eccentricity-, PN- and SF-expansions of the function $\psi(u_p, e; \nu)$ is then found to be of the type

$$\psi(u_p, e) = \psi^{(0)}(u_p, \nu) + e^2 \psi^{(2)}(u_p, \nu) + e^4 \psi^{(4)}(u_p, \nu) + \mathcal{O}(e^6), \tag{4.75}$$

where the beginnings of the expansions of $\psi^{(0)}(u_p, \nu)$ and $\psi^{(2)}(u_p, \nu)$ are given by

$$\begin{aligned}
\psi^{(0)}(u_p, \nu) &= \left(\frac{3}{2} - \nu\right) u_p + \left(\frac{45}{8} - 3\nu\right) u_p^2 + \left(-33\nu + \frac{663}{16}\right) u_p^3 + \left(-\frac{1537}{3}\nu + \frac{41}{8}\nu\pi^2 + \frac{47805}{128}\right) u_p^4 \\
&\quad + \left(\frac{951309}{256} - \frac{4336}{15}\nu\ln(2) + \nu g_{*22} - 144\nu\gamma + \frac{109897}{1024}\nu\pi^2 - \frac{1162607}{160}\nu - 72\nu\ln(u_p)\right) u_p^5 + \mathcal{O}(u_p^6, \nu^2) \\
\psi^{(2)}(u_p, \nu) &= \left(3 - \frac{3}{2}\nu\right) u_p^2 + \left(-39\nu + \frac{75}{2}\right) u_p^3 + \left[\left(-\frac{56179}{64} + \frac{615}{64}\pi^2 + \frac{1}{2}g_{*22}\right)\nu + \frac{3639}{8}\right] u_p^4 \\
&\quad + \left[\frac{89751}{16} + \left(\frac{3}{2}g_{*41} + \frac{1}{2}g_{*23}^c - \frac{6176}{15}\ln(2) + \frac{35}{4}g_{*22} - \frac{3232}{5}\gamma + \frac{282651}{1024}\pi^2\right.\right. \\
&\quad \left.- \frac{5618027}{384} + \frac{1}{2}g_{*23}^{\ln}\ln(u_p) - \frac{1616}{5}\ln(u_p) - \frac{4374}{5}\ln(3)\right)\nu\left. \right] u_p^5 + \mathcal{O}(u_p^6, \nu^2) \\
\psi^{(4)}(u_p, \nu) &= \left(-\frac{3}{4}\nu + \frac{3}{16}\right) u_p^3 + \left[\frac{1101}{32} + \left(-\frac{52051}{512} + \frac{123}{256}\pi^2 + \frac{3}{8}g_{*22} + \frac{3}{8}g_{*41}\right)\nu\right] u_p^4 \\
&\quad + \left[\frac{124533}{128} + \left(-\frac{1074}{5}\gamma - \frac{45358}{5}\ln(2) + \frac{675243}{8192}\pi^2 - \frac{537}{5}\ln(u_p) + \frac{19683}{4}\ln(3) + \frac{145}{16}g_{*22}\right.\right.
\end{aligned}$$

$$\begin{aligned}
& + \frac{99}{16}g_{*41} + \frac{3}{4}g_{*23}^{\ln} \ln(u_p) + \frac{3}{8}g_{*42}^{\ln} \ln(u_p) + \frac{3}{4}g_{*23}^c + \frac{3}{8}g_{*42}^c + \frac{7}{16}g_{*23}^{\ln} - \frac{17185101}{5120} \Big) \nu \Big] u_p^5 \\
& + \mathcal{O}(u_p^6, \nu^2). \tag{4.76}
\end{aligned}$$

Though the so obtained function $\psi(u_p, e; \nu)$ is gauge-invariant (u_p and e being functions of the gauge-invariant dynamical quantities $\hat{\mathcal{E}}_{\text{eff}}$ and j), it cannot be directly compared with the (equally gauge-invariant) function $\psi(m_2\Omega_r, m_2\Omega_\phi; q)$ obtained in the SF computation above.

In order to compare our EOB-derived result with the SF result we first need to transform the dependence on u_p and e into a dependence on $Gm_2\Omega_r$ and $Gm_2\Omega_\phi$, or equivalently, on $y = (Gm_2\Omega_\phi)^{2/3}$ and $\hat{k} = \frac{1}{3}(\frac{\Omega_\phi}{\Omega_r} - 1)$.

Using Eqs. (4.65) above we then find that the EOB-derived functional dependence of ψ on (\hat{k}, y) is given, at first order in ν , by

$$\psi(\hat{k}, y) = \psi_0(\hat{k}, y) + \nu \Delta\psi(\hat{k}, y) + \mathcal{O}(\nu^2), \tag{4.77}$$

where the beginnings of the expansions of $\psi_0(\hat{k}, y)$ and $\Delta\psi(\hat{k}, y)$ read

$$\begin{aligned}
\psi_0(\hat{k}, y) &= \frac{3}{2}\hat{k} - \left(\frac{9}{8}\hat{k}y + \frac{9}{2}\hat{k}^2\right) + \left(\frac{75}{32}\hat{k}^2y - \frac{75}{32}y^2\hat{k} + \frac{27}{2}\hat{k}^3\right) \\
&+ \left(-\frac{135}{128}\hat{k}^3y - \frac{45}{128}\hat{k}^2y^2 - \frac{45}{8}\hat{k}y^3 - \frac{5229}{128}\hat{k}^4\right) + \dots \\
\Delta\psi(\hat{k}, y) &= -\hat{k} + \left(8\hat{k}^2 - \frac{5}{4}\hat{k}y\right) + \left(\frac{123}{256}\hat{k}^2\pi^2y - \frac{615}{256}\hat{k}^3\pi^2\right. \\
&- \frac{93}{8}\hat{k}^2y - \frac{53}{16}y^2\hat{k} + \frac{69}{4}\hat{k}^3\Big) + \left(c_6^c + c_6^{\ln} \ln(\hat{k})\right)\hat{k}^4 \\
&+ \dots \tag{4.78}
\end{aligned}$$

with

$$\begin{aligned}
c_6^c &= -\frac{36735959}{23040} + \left(-\frac{1}{16}g_{*41} + \frac{177067}{1440} - \frac{3995649}{320} \ln(3) - \frac{11518508}{45} \ln(2) + \frac{68359375}{576} \ln(5)\right) \left(\frac{y}{\hat{k}}\right)^3 \\
&+ \left(-\frac{5137157}{7680} + \frac{9}{16}g_{*41} + \frac{10900979}{15} \ln(2) + \frac{37}{5}\gamma - \frac{33203125}{96} \ln(5) + \frac{3}{8}g_{*22} - \frac{4081}{16384}\pi^2 + \frac{1560789}{32} \ln(3)\right) \left(\frac{y}{\hat{k}}\right)^2 \\
&+ \left(-\frac{15}{16}g_{*41} - \frac{5}{4}g_{*22} + \frac{1111433}{768} + \frac{11195}{8192}\pi^2 - 689470 \ln(2) + \frac{21484375}{64} \ln(5) - 122\gamma - \frac{18575649}{320} \ln(3)\right) \frac{y}{\hat{k}} \\
&- \frac{1953125}{18} \ln(5) + \frac{7}{8}g_{*22} + \frac{7}{16}g_{*41} + \frac{9842609}{45} \ln(2) + \frac{595}{3}\gamma + \frac{438129}{20} \ln(3) + \frac{1580185}{49152}\pi^2 \\
c_6^{\ln} &= \frac{595}{6} - 61\frac{y}{\hat{k}} + \frac{37}{10} \left(\frac{y}{\hat{k}}\right)^2. \tag{4.79}
\end{aligned}$$

Of most interest for our present work is the function $\Delta\psi(\hat{k}, y)$, Eq. (4.78). We can directly compare the latter function with the one computed with SF theory above in Eq. (3.33), modulo the fact that the SF computed one above was expressed not in terms of \hat{k} and y , but instead in terms of eccentricity and semi-latus rectum parameters, \bar{e} and $\bar{p} = 1/\bar{u}_p$, different from the ones, e and u_p , used here (recalling the notational change discussed at the beginning of this section).

The transformation between (\bar{e}, \bar{u}_p) and (e, u_p) is only needed at order ν^0 and can be obtained by identifying the μ -rescaled Schwarzschild energy and angular momentum used to define \bar{e} and \bar{u}_p to the EOB quantities $\hat{\mathcal{E}}_{\text{eff}}$ and j . In other words, while (\bar{e}, \bar{u}_p) were defined in Eqs. (2.12), (2.13), i.e., equivalently, by writing

$$\hat{\mathcal{E}}_{\text{eff}} = \sqrt{\frac{(1 - 2\bar{u}_p)^2 - 4\bar{u}_p^2\bar{e}^2}{1 - 3\bar{u}_p - \bar{u}_p\bar{e}^2}}, \tag{4.80}$$

$$j = \frac{1}{\sqrt{\bar{u}_p(1 - 3\bar{u}_p - \bar{u}_p\bar{e}^2)}}, \tag{4.81}$$

the other pair (e, u_p) , used in our EOB computation, was defined by writing

$$\hat{\mathcal{E}}_{\text{eff}} = \sqrt{1 - u_p(1 - e^2)}, \quad j = \frac{1}{\sqrt{u_p}}. \tag{4.82}$$

The comparison between these two expressions implies the following transformation law

$$\begin{aligned}
u_p &= \bar{u}_p(1 - 3\bar{u}_p - \bar{u}_p\bar{e}^2), \\
1 - e^2 &= (1 - \bar{e}^2) \frac{1 - 4\bar{u}_p}{(1 - 3\bar{u}_p - \bar{u}_p\bar{e}^2)^2}. \tag{4.83}
\end{aligned}$$

Using either this transformation (together with intermediate equations given above) or directly the well known elliptic-integrals expressions giving Ω_ϕ and k in a Schwarzschild background, consistently with Eqs.

(2.13)-(2.16) in Ref. [36], we obtain

$$\begin{aligned}
\hat{k} &= \bar{u}_p + \left(\frac{9}{2} + \frac{1}{4}\bar{e}^2\right)\bar{u}_p^2 + \left(\frac{45}{2} + \frac{15}{4}\bar{e}^2\right)\bar{u}_p^3 + \left(\frac{35}{64}\bar{e}^4 + \frac{945}{8} + \frac{315}{8}\bar{e}^2\right)\bar{u}_p^4 + \left(\frac{2835}{8}\bar{e}^2 + \frac{945}{64}\bar{e}^4 + \frac{5103}{8}\right)\bar{u}_p^5 \\
&+ \left(\frac{31185}{128}\bar{e}^4 + \frac{93555}{32}\bar{e}^2 + \frac{385}{256}\bar{e}^6 + \frac{56133}{16}\right)\bar{u}_p^6 + \left(\frac{405405}{128}\bar{e}^4 + \frac{729729}{32}\bar{e}^2 + \frac{312741}{16} + \frac{15015}{256}\bar{e}^6\right)\bar{u}_p^7 \\
&+ \left(\frac{18243225}{512}\bar{e}^4 + \frac{10945935}{64}\bar{e}^2 + \frac{14073345}{128} + \frac{675675}{512}\bar{e}^6\right)\bar{u}_p^8 + \mathcal{O}(\bar{u}_p^9) \\
y &= (1 - \bar{e}^2)\bar{u}_p + (2\bar{e}^2 - 2\bar{e}^4)\bar{u}_p^2 + \left(-\frac{23}{8}\bar{e}^4 - \frac{55}{16}\bar{e}^6 + 6\bar{e}^2\right)\bar{u}_p^3 + \left(-\frac{13}{4}\bar{e}^4 + 24\bar{e}^2 - \frac{209}{12}\bar{e}^6\right)\bar{u}_p^4 \\
&+ \left(-\frac{1237}{16}\bar{e}^6 - \frac{1}{4}\bar{e}^4 + 120\bar{e}^2\right)\bar{u}_p^5 + \left(-\frac{3069}{8}\bar{e}^6 + 672\bar{e}^2 + 51\bar{e}^4\right)\bar{u}_p^6 + \left(730\bar{e}^4 - \frac{52265}{24}\bar{e}^6 + 3936\bar{e}^2\right)\bar{u}_p^7 \\
&+ \left(-\frac{26279}{2}\bar{e}^6 + 8032\bar{e}^4 + 23424\bar{e}^2\right)\bar{u}_p^8 + \mathcal{O}(\bar{u}_p^9). \tag{4.84}
\end{aligned}$$

Inserting these expressions in the EOB-derived func-

tion $\Delta\psi(\hat{k}, y)$ obtained above gives

$$\Delta\psi(\bar{u}_p, \bar{e}) = \Delta\psi^{(0)}(\bar{u}_p) + \bar{e}^2\Delta\psi^{(2)}(\bar{u}_p) + \bar{e}^4\Delta\psi^{(4)}(\bar{u}_p) + \dots \tag{4.85}$$

where $\Delta\psi^{(0)}(\bar{u}_p)$ will be discussed below, and where the $\mathcal{O}(\bar{e}^2)$ contribution (which is new with this work) reads

$$\begin{aligned}
\Delta\psi^{(2)}(\bar{u}_p) &= \bar{u}_p^2 + \left(\frac{341}{16} - \frac{123}{256}\pi^2\right)\bar{u}_p^3 \\
&+ \left(-\frac{317491}{960} + \frac{268}{5}\ln(\bar{u}_p) - \frac{23729}{4096}\pi^2 + \frac{1}{2}g_{*22} + \frac{536}{5}\gamma + \frac{11720}{3}\ln(2) - \frac{10206}{5}\ln(3)\right)\bar{u}_p^4 \\
&+ \left(\frac{21797069}{49152}\pi^2 - \frac{131386901}{57600} - \frac{10957}{15}\ln(\bar{u}_p) + \frac{5}{4}g_{*22} + \frac{9765625}{1344}\ln(5) + \frac{1}{2}g_{*23}^c + \frac{1}{2}g_{*23}^{\ln}\ln(\bar{u}_p)\right. \\
&\quad \left.- \frac{21914}{15}\gamma - \frac{333378}{7}\ln(2) + \frac{4943349}{320}\ln(3)\right)\bar{u}_p^5 \\
&+ \frac{319609}{630}\pi\bar{u}_p^{11/2} \\
&+ \left(\frac{5}{4}g_{*23}^c - \frac{9232551768011}{101606400} - \frac{33}{16}g_{*22} - \frac{225143631}{8960}\ln(3) + \frac{1}{2}g_{*24}^c + \frac{1}{2}\ln(\bar{u}_p)g_{*24}^{\ln} - \frac{33}{16}\right. \\
&\quad \left.+ \frac{5}{4}g_{*23}^{\ln}\ln(\bar{u}_p) - \frac{17193359375}{145152}\ln(5) + \frac{284236994}{945}\ln(2) - \frac{529328}{189}\gamma - \frac{146026515}{1048576}\pi^4 + \frac{31842559607}{2359296}\pi^2\right. \\
&\quad \left.- \frac{264664}{189}\ln(\bar{u}_p)\right)\bar{u}_p^6 + \mathcal{O}(\bar{u}_p^{13/2}), \tag{4.86}
\end{aligned}$$

Here the terms up to $\mathcal{O}(u_p^3)$ coincide with those given in Eq. (5.4) of [36], and the undetermined coefficients g_{*22}, g_{*23}^c , etc. which parametrized the unknown $\mathcal{O}(p_r^2)$ dependence of the EOB gyrogravitomagnetic ratio

$g_{S*}(u, p_r, p_\phi)$ first enter $\Delta\psi^{(2)}(\bar{u}_p)$ at order $\mathcal{O}(u^4)$.

Comparing with Eq. (3.33) above we then find unique values for the so far undetermined EOB coefficients, namely

$$\begin{aligned}
g_{*22} &= -\frac{717}{32} \\
g_{*23}^c &= \frac{1447441}{960} - \frac{4829}{256}\pi^2 - \frac{16038}{5}\ln(3) + \frac{46976}{15}\ln(2) - \frac{512}{5}\gamma \\
g_{*23}^{\ln} &= -\frac{256}{5} \\
g_{*24}^c &= -\frac{185195453}{38400} + \frac{19162}{35}\gamma + \frac{2097479}{8192}\pi^2 + \frac{454167}{20}\ln(3) - \frac{1081966}{35}\ln(2) \\
g_{*24}^{\ln} &= +\frac{9581}{35}.
\end{aligned} \tag{4.87}$$

We also computed the $\mathcal{O}(\bar{e}^4)$ contribution to $\Delta\psi(\bar{u}_p, \bar{e})$ which we give below in its parametrized form

$$\begin{aligned}
\Delta\psi^{(4)}(\bar{u}_p) &= -\frac{1}{2}\bar{u}_p^3 \\
&+ \left(-\frac{366857}{1536} + \frac{3}{8}g_{*41} + \frac{1953125}{192}\ln(5) + \frac{37}{10}\ln(\bar{u}_p) - \frac{23761}{16384}\pi^2 + \frac{3}{8}g_{*22} + \frac{37}{5}\gamma - \frac{205843}{5}\ln(2) \right. \\
&\quad \left. + \frac{3620943}{320}\ln(3) \right) \bar{u}_p^4 \\
&+ \left(-\frac{524787}{65536}\pi^2 + \frac{720900871}{537600} - \frac{7143}{56}\ln(\bar{u}_p) + \frac{3}{8}g_{*42}^c + \frac{21}{16}g_{*41} + \frac{57}{16}g_{*22} + \frac{72265625}{5376}\ln(5) \right. \\
&\quad \left. + \frac{7}{16}g_{*23}^{\ln} + \frac{3}{4}g_{*23}^c + \frac{3}{4}g_{*23}^{\ln}\ln(\bar{u}_p) + \frac{3}{8}g_{*42}^{\ln}\ln(\bar{u}_p) - \frac{7143}{28}\gamma - \frac{30009569}{420}\ln(2) + \frac{210887307}{8960}\ln(3) \right) \bar{u}_p^5 \\
&+ \frac{1089581}{5120}\pi\bar{u}_p^{11/2} \\
&+ \left(\frac{55}{32}g_{*23}^{\ln} - \frac{255}{64}g_{*41} + \frac{41}{8}g_{*23}^c - \frac{797597168303}{16934400} + \frac{661}{64}g_{*22} - \frac{678223072849}{153600}\ln(7) + \frac{4018699514169}{358400}\ln(3) \right. \\
&\quad \left. + \frac{9}{16}g_{*24}^{\ln} + \frac{5}{4}g_{*24}^c + \frac{21}{16}g_{*42}^c + \frac{3}{8}g_{*43}^c + \frac{5}{4}\ln(\bar{u}_p)g_{*24}^{\ln} + \frac{3}{8}g_{*43}^{\ln}\ln(\bar{u}_p) + \frac{21}{16}g_{*42}^{\ln}\ln(\bar{u}_p) \right. \\
&\quad \left. + \frac{41}{8}g_{*23}^{\ln}\ln(\bar{u}_p) - \frac{17944462890625}{1161216}\ln(5) + \frac{2304030315373}{75600}\ln(2) - \frac{48044911}{15120}\gamma - \frac{17998485}{524288}\pi^4 \right. \\
&\quad \left. + \frac{151169660653}{18874368}\pi^2 - \frac{48044911}{30240}\ln(\bar{u}_p) \right) \bar{u}_p^6 + \mathcal{O}(\bar{u}_p^{13/2}).
\end{aligned} \tag{4.88}$$

As soon as some SF data on ψ at order $\mathcal{O}(e^4)$ become available, this result will allow one to determine the coefficients parametrizing the $\mathcal{O}(p_r^4)$ EOB gyrogravitomagnetic ratio g_{S*} .

Concerning the quantity $\Delta\psi(\bar{u}_p, \bar{e} \rightarrow 0)$, it can be obtained (as pointed out in Ref. [36]) by combining the previous SF results on $\Delta\psi_{(\text{circ})}(\bar{u}_p)$ [34, 35, 68]

$$\begin{aligned}
\Delta\psi_{(\text{circ})}(\bar{u}_p) &= \bar{u}_p^2 - 3\bar{u}_p^3 - \frac{15}{2}\bar{u}_p^4 + \left(-\frac{6277}{30} - \frac{496}{15}\ln(2) \right. \\
&\quad \left. - 16\gamma - 8\ln(\bar{u}_p) + \frac{20471}{1024}\pi^2 \right) \bar{u}_p^5 + \dots,
\end{aligned} \tag{4.89}$$

with the knowledge of the EOB function $\rho(x)$ measuring periastron precession at the 1SF-level [25, 59], namely

$$\Delta\psi(\bar{u}_p, \bar{e} \rightarrow 0) =$$

$$\Delta\psi_{(\text{circ})}(\bar{u}_p) - \frac{1}{2} \frac{(1 - 3\bar{u}_p)^{1/2}(1 - 6\bar{u}_p)^{5/2}}{1 - \frac{39}{4}\bar{u}_p + \frac{43}{2}\bar{u}_p^2} k_{(\text{circ})}^{1\text{SF}}(\bar{u}_p), \tag{4.90}$$

where

$$k_{(\text{circ})}^{1\text{SF}}(y) = -\frac{\rho(y) - 4y}{2(1 - 6y)^{3/2}}, \tag{4.91}$$

as obtained from the definition $(1 + k_{(\text{circ})}(x, \nu))^{-2} = 1 - 6x + \nu\rho(x) + \mathcal{O}(\nu^2)$, i.e., using the link $x = (GM\Omega_\phi)^{2/3} = (1 + \frac{2}{3}\nu)y + \mathcal{O}(\nu^2)$ and

$$\begin{aligned}
k_{(\text{circ})}(y, \nu) &= -1 + \frac{1}{\sqrt{1 - 6(1 + \frac{2}{3}\nu)y + \nu\rho(y)}} \\
&= k_{(\text{circ})}^{(0)}(y) + \nu k_{(\text{circ})}^{1\text{SF}}(y) + \mathcal{O}(\nu^2),
\end{aligned} \tag{4.92}$$

with

$$k_{(\text{circ})}^{(0)}(y) = -1 + \frac{1}{\sqrt{1 - 6y}}. \tag{4.93}$$

V. RESUMMATION OF g_{S*}

SF technology allows one to reach high PN orders when working linearly in the symmetric mass ratio ν [14, 15, 33, 35, 49, 59, 74–77]. We have given here one more example of this feature by deriving the $\mathcal{O}(e^2)$ (respectively, $\mathcal{O}(p_r^2)$) piece in the spin precession function ψ (respectively, gyrogravitomagnetic ratio g_{S*}) to many more PN-orders than was previously known. Past work has shown that an efficient way of using the high-PN information obtained by SF computations is to incorporate it, together with known PN contributions which are higher order in ν , within some suitably resummed coupling functions of the EOB formalism. Let us show how such an approach can be applied to the g_{S*} EOB coupling function.

To blend and resum SF and PN information concerning g_{S*} let us start from the structure of the test-mass value of g_{S*} , as written (when neglecting effects quadratic in spins) in Eq. (4.14). First, we note that this structure can be rewritten as

$$g_{S*}^{(\nu^0)}(u, p_r, p_\phi) = \frac{\sqrt{A_{\text{schw}}(u)}}{\hat{\mathcal{E}}_{\text{schw}} + \sqrt{A_{\text{schw}}(u)}} + \frac{2A_{\text{schw}}(u)}{\hat{\mathcal{E}}_{\text{schw}}(1 + \sqrt{A_{\text{schw}}(u)})}, \quad (5.1)$$

where $A_{\text{schw}}(u) = 1 - 2u$ and where $\hat{\mathcal{E}}_{\text{schw}} = \sqrt{(1 - 2u)(1 + p_\phi^2 u^2 + (1 - 2u)p_r^2)}$ is the conserved energy of a test particle in a Schwarzschild background. Taking this reformulation as a model, let us define the following EOB-compatible, ν -deformed g_{S*} coupling

$$g_{S*}^{(\text{EOB-like})}(u, p_r, p_\phi; \nu) = \frac{\sqrt{A_{3PN}(u; \nu)}}{\hat{H}_{(\text{orb, eff})}^{3PN} + \sqrt{A_{3PN}(u; \nu)}} + \frac{2A_{3PN}(u; \nu)}{\hat{H}_{(\text{orb, eff})}^{3PN}(1 + \sqrt{A_{3PN}(u; \nu)})}, \quad (5.2)$$

where $A_{3PN}(u; \nu)$ is defined as

$$A_{3PN}(u; \nu) \equiv 1 - 2u + 2\nu u^3 + \left(\frac{94}{3} - \frac{41}{32}\pi^2\right)\nu u^4, \quad (5.3)$$

and where

$$\hat{H}_{(\text{orb, eff})}^{3PN} \equiv \sqrt{A_{3PN}(u; \nu)} \times \sqrt{\left(1 + p_\phi^2 u^2 + A_{3PN}(u; \nu)\bar{D}_{3PN}(u; \nu)p_r^2 + \hat{Q}_{3PN}\right)}, \quad (5.4)$$

with

$$\begin{aligned} \bar{D}_{3PN}(u; \nu) &= 1 + 6\nu u^2 + (52\nu - 6\nu^2)u^3, \\ \hat{Q}_{3PN}(u; \nu) &= 2(4 - 3\nu)\nu u^2 p_r^4. \end{aligned} \quad (5.5)$$

We propose to use the ν -deformed ratio $g_{S*}^{(\text{EOB-like})}$, Eq. (5.2), as a reference value for the (unknown) exact g_{S*} , and to incorporate the current PN and SF information of g_{S*} in the form of a *correcting factor* $r_{S*}(u, p_r, p_\phi; \nu)$ of the type $r_{S*} = 1 + \mathcal{O}(\nu)$. [We have also explored the possibility of translating g_{S*} in terms of the gyrogravitomagnetic ratio of a test-mass in some spin-effective metric. However, this led to an effective metric whose ν -deformation was drastically different (starting at the 1PN-order) from the one of the usual orbital EOB effective metric, and which did not seem to offer a good starting point for resumming g_{S*} .] In other words, we propose to define a resummed g_{S*} of the form

$$g_{S*}^{\text{resum}}(u, p_r, p_\phi; \nu) = g_{S*}^{(\text{EOB-like})}(u, p_r, p_\phi; \nu) r_{S*}(u, p_r, p_\phi; \nu). \quad (5.6)$$

When considering fast-spinning black holes, one can still use the factorized form (5.6), but with suitably spin-quadratic-extended values of A_{3PN} and $\hat{H}_{(\text{orb, eff})}^{3PN}$ in the definition (5.2) of $g_{S*}^{(\text{EOB-like})}$ (e.g., as defined in Ref. [66]).

We will suggest several possible estimates of the correcting factor r_{S*} in Eq. (5.6). All these estimates will be constructed from the PN expansion of the correcting factor $r_{S*} = g_{S*}^{SF+PN}/g_{S*}^{(\text{EOB-like})}$, which is of the form

$$r_{S*}^{\text{PN}}(u, p_r, p_\phi; \nu) = 1 + \nu \hat{g}_1(u, p_r, p_\phi) + \nu^2 \hat{g}_2(u, p_r, p_\phi), \quad (5.7)$$

where

$$\begin{aligned} \hat{g}_1(u, p_r, p_\phi) &= \eta^2 \left(-\frac{1}{2}u - \frac{3}{2}p_r^2 \right) + \eta^4 \left[\frac{25}{24}p_r^4 + \left(-\frac{5}{8}p_\phi^2 u^2 - \frac{53}{24}u \right) p_r^2 - \frac{5}{24}p_\phi^2 u^3 - \frac{20}{3}u^2 \right] \\ &+ \eta^6 \left[\left(-\frac{2297}{144}u^2 + \frac{17}{96}p_\phi^4 u^4 - \frac{97}{144}p_\phi^2 u^3 \right) p_r^2 + \frac{17}{288}p_\phi^4 u^5 - \frac{8771}{288}u^3 - \frac{199}{72}p_\phi^2 u^4 + \frac{41}{48}u^3 \pi^2 \right] \\ &+ \eta^8 \left[\left(-\frac{1483}{192}p_\phi^2 u^4 - \frac{10692}{5}u^3 \ln(3) - \frac{1024}{15}u^3 \gamma - \frac{14077}{1152}u^3 \pi^2 + \frac{93952}{45}u^3 \ln(2) - \frac{55}{576}p_\phi^6 u^6 \right) \right. \\ &\left. + \frac{77}{576}p_\phi^4 u^5 + \frac{17436377}{17280}u^3 - \frac{512}{15}u^3 \ln(u) \right] p_r^2 - \frac{354523}{4320}u^4 - \frac{42575}{3456}p_\phi^2 u^5 + \frac{113}{144}p_\phi^4 u^6 \end{aligned}$$

$$\begin{aligned}
& - \frac{55}{1728} p_\phi^6 u^7 - \frac{2912}{45} u^4 \ln(2) + \frac{85421}{9216} u^4 \pi^2 + \frac{205}{576} p_\phi^2 u^5 \pi^2 - 32 u^4 \gamma - 16 u^4 \ln(u) \Big] \\
& + \eta^{10} \left[\left(\frac{144261}{10} u^4 \ln(3) - \frac{256}{9} u^5 \gamma p_\phi^2 - \frac{433}{10368} p_\phi^6 u^7 - 891 u^5 \ln(3) p_\phi^2 + \frac{23488}{27} u^5 \ln(2) p_\phi^2 \right. \right. \\
& - \frac{128}{9} u^5 \ln(u) p_\phi^2 - \frac{6281068}{315} u^4 \ln(2) + \frac{103604}{315} u^4 \gamma + \frac{869}{13824} p_\phi^8 u^8 \\
& - \frac{505706219}{172800} u^4 + \frac{22577}{6912} p_\phi^4 u^6 + \frac{8367161}{20736} p_\phi^2 u^5 + \frac{146005}{864} u^4 \pi^2 - \frac{75223}{13824} p_\phi^2 u^5 \pi^2 + \frac{51802}{315} u^4 \ln(u) \Big) p_r^2 \\
& + \frac{5248}{35} u^5 \gamma - \frac{2887}{96} p_\phi^2 u^6 + \frac{145379}{41472} p_\phi^4 u^7 - \frac{2197}{5184} p_\phi^6 u^8 + \frac{869}{41472} p_\phi^8 u^9 - \frac{4577190091}{3628800} u^5 \\
& + \frac{13059635}{110592} u^5 \pi^2 - \frac{40}{3} u^6 \gamma p_\phi^2 + \frac{2624}{35} u^5 \ln(u) - \frac{20}{3} u^6 \ln(u) p_\phi^2 \\
& + \frac{353392}{945} u^5 \ln(2) - \frac{697}{6912} p_\phi^4 u^7 \pi^2 - \frac{728}{27} u^6 \ln(2) p_\phi^2 - \frac{486}{7} u^5 \ln(3) + \frac{412673}{110592} p_\phi^2 u^6 \pi^2 \Big] + \mathcal{O}(u^6), \quad (5.8)
\end{aligned}$$

and

$$\hat{g}_2(u, p_r, p_\phi) = \left(\frac{19}{8} p_r^2 u - \frac{1}{8} u^2 + \frac{15}{8} p_r^4 \right) \eta^4. \quad (5.9)$$

In Eq. (5.8) we have displayed only the beginning of the PN expansion of \hat{g}_1 , including in particular all the $\mathcal{O}(p_r^2)$ terms that have been derived in the present paper. We kept in Eq. (5.8) only the terms that are fully known, i.e., the terms that do not involve any yet undetermined coefficient such as $g_{*41} u p_r^4 \eta^6$, etc. [More precisely, in \hat{g}_1 we see that the p_r^4 contribution is fully known only at $\mathcal{O}(\eta^4)$; by contrast, the p_r^2 one is now known from $\mathcal{O}(\eta^2)$ up to $\mathcal{O}(\eta^{10})$; in \hat{g}_2 only terms at $\mathcal{O}(\eta^4)$ are known.] Beyond

the contributions indicated, only the circular limit of \hat{g}_1 is known and it can be straightforwardly computed (in the DJS gauge) by PN-expanding the prescription given above, using for instance either the explicit 9.5PN accurate g_{S*}^{circ} result of Ref. [68] or the implicit 22PN-accurate result of Ref. [35] on ψ_{circ} .

In the rest of this subsection we will motivate various ways (namely, Taylor-like or inverse-Taylor-like, at various PN-approximations) of defining the correcting factor r_{S*} in Eq. (5.7) by studying the circular limit of g_{S*}^{resum} , Eq. (5.6). For concreteness, let us exhibit the explicit 4PN-accurate value of r_{S*} when setting $p_r \rightarrow 0$

$$\begin{aligned}
r_{S*}^{4PN}(u, p_\phi; \nu) = & 1 - \frac{1}{2} u \nu \eta^2 + \left(-\frac{20}{3} \nu u^2 - \frac{1}{8} \nu^2 u^2 - \frac{5}{24} \nu p_\phi^2 u^3 \right) \eta^4 \\
& + \left(-\frac{8771}{288} u^3 + \frac{41}{48} u^3 \pi^2 - \frac{199}{72} p_\phi^2 u^4 + \frac{17}{288} p_\phi^4 u^5 \right) \nu \eta^6 \\
& + \left(-\frac{354523}{4320} u^4 - \frac{42575}{3456} u^5 p_\phi^2 - \frac{55}{1728} p_\phi^6 u^7 + \frac{113}{144} p_\phi^4 u^6 - 16 u^4 \ln(u \eta^2) \right. \\
& \left. - \frac{2912}{45} u^4 \ln(2) - 32 u^4 \gamma + \frac{85421}{9216} u^4 \pi^2 + \frac{205}{576} u^5 \pi^2 p_\phi^2 \right) \nu \eta^8. \quad (5.10)
\end{aligned}$$

Note the explicit appearance of p_ϕ in r_{S*}^{4PN} . We wish to study an estimate of g_{S*} that would be approximately valid when considering (circularized) inspiralling and coalescing binary black holes. Following the results of EOB theory [5, 6] we can approximately replace p_ϕ by a function of u having the following properties.

Above the last stable orbit we estimate $p_\phi \approx p_\phi^{(\text{circ})}(u) = \sqrt{-\partial_u A(u; \nu) / \partial_u (u^2 A(u; \nu))}$, which we simply approximate in our present qualitative study by $p_\phi \approx [u(1 - 3u)]^{-1/2}$. Beyond the last stable orbit we instead approximate p_ϕ by a constant equals to its value at

the last stable orbit (say $p_\phi \approx \sqrt{12}$). This approximation defines the evolution of r_{S*} , and correlatively of g_{S*} , during coalescence, as a function of u . The results of this exploratory study of the evolution of g_{S*} during coalescence are illustrated in Fig. 1, which considers the equal-mass case, $\nu = 1/4$. This figure compares the u -evolution of various approximations to $g_{S*} = g_{S*}^{(\text{EOB-like})} r_{S*}$: 1) the one defined by using the Taylor-like 4PN approximant for r_{S*} , Eq. (5.10) together with the just explained p_ϕ replacements; 2) the one defined by using inverse resummation of r_{S*}^{4PN} , i.e., by taking the $P^{[1,4]}$ Padé approxi-




FIG. 1. Comparison of various estimates of the EOB gravitomagnetic ratio g_{S*} in the equal-mass case ($\nu = 1/4$), and in the circularized inspiralling and coalescing approximation. The resummed g_{S*} defined by Eq. (5.6) is plotted as a function of $u = GM/(c^2 R)$ for the four cases explained in text.

mant of the PN-expansion of Eq. (5.10); 3) *ditto* with the 9.5 PN-accurate generalization of (5.10); and finally 4) *ditto* with the inverse resummation of the latter $r_{S*}^{9.5PN}$ expansion.

The main message of Fig. 1 is that, in confirmation of what had been already found at the next-to-leading order [60], and whatever be the approximation used, g_{S*} significantly decreases as the separation $R = GM/u$ between the two bodies decreases. This decrease becomes more pronounced when one uses higher PN-approximants. When using Taylor-approximants the decrease of g_{S*} is so extreme that it formally changes a sign below a certain separation R_{crit} . For instance, at 4PN we have $R_{\text{crit}} \approx GM/0.37 \approx 2.7GM$, while at 9.5PN we have $R_{\text{crit}} \approx GM/0.31 \approx 3.2GM$. As this sign change is physically unwarranted, we advise, when using Taylor-approximants, to replace g_{S*} by zero beyond the critical separation R_{crit} (i.e., to replace $g_{S*} \rightarrow \frac{1}{2}(g_{S*} + |g_{S*}|)$).

If one considers that such a vanishing of g_{S*} is too extreme a behaviour, one might consider using one of the inverse-resummed estimates of r_{S*} . At this stage the spread between the curves in Fig. 1 is a measure of our uncertainty on the true value of g_{S*} in the strong-field

domain. One will need comparisons between numerical relativity simulations and EOB computations using various g_{S*} functions (possibly including some free parameter parametrizing strong-field effects) to learn more about the exact extent to which g_{S*} decreases in the strong-field domain.

ACKNOWLEDGMENTS

CK thanks the authors of [36] for generously providing an early draft of their work, and in particular Sarp Akcay and Sam Dolan for many informative discussions. DB thanks ICRANet and the italian INFN for partial support and IHES for warm hospitality at various stages during the development of the present project. SH acknowledges financial support provided under the European Union's H2020 ERC Consolidator Grant “Matter and strong-field gravity: New frontiers in Einstein's theory” grant agreement no. MaGRaTh-646597.

Appendix A: Kinnersley tetrad

Using Boyer-Lindquist coordinates, the Kinnersley tetrad upon which the Weyl tensor is projected is given by

$$\begin{aligned} e_1^\alpha &= l^\alpha = \frac{1}{\Delta}(r^2 + a^2, \Delta, 0, a), \\ e_2^\alpha &= n^\alpha = \frac{1}{2\Sigma}(r^2 + a^2, -\Delta, 0, a), \\ e_3^\alpha &= m^\alpha = -\frac{\bar{\varrho}}{\sqrt{2}}(ia \sin \theta, 0, 1, \frac{i}{\sin \theta}), \\ e_4^\alpha &= \bar{m}^\alpha = -\frac{\varrho}{\sqrt{2}}(-ia \sin \theta, 0, 1, \frac{-i}{\sin \theta}). \end{aligned}$$

where $\Sigma \equiv r^2 + a^2 \cos^2 \theta$, $\Delta \equiv r^2 - 2m_2 r + a^2$. The non-zero spin-coefficients which appear in Sec. II D are

$$\begin{aligned} \varrho &= \frac{-1}{r - ia \cos \theta}, & \tau &= \frac{-ia \sin \theta}{\sqrt{2}\Sigma}, & \beta &= -\frac{\bar{\varrho} \cot \theta}{2\sqrt{2}}, \\ \gamma &= \mu + \frac{r - m_2}{2\Sigma}, & \mu &= \frac{\Delta \varrho}{2\Sigma}, & \varpi &= \frac{ia \varrho^2 \sin \theta}{\sqrt{2}}, \\ \alpha &= \varpi - \bar{\beta}. \end{aligned}$$

We give these in Kerr spacetime, our situation is recovered by setting a to zero in the above.

-
- [1] B. P. Abbott *et al.* (Virgo, LIGO Scientific), “Observation of Gravitational Waves from a Binary Black Hole Merger,” *Phys. Rev. Lett.* **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
 - [2] B. P. Abbott *et al.* (Virgo, LIGO Scientific), “GW151226: Observation of Gravitational Waves from a 22-Solar-

Mass Binary Black Hole Coalescence,” *Phys. Rev. Lett.* **116**, 241103 (2016), arXiv:1606.04855 [gr-qc].

- [3] Andrea Taracchini *et al.*, “Effective-one-body model for black-hole binaries with generic mass ratios and spins,” *Phys. Rev.* **D89**, 061502 (2014), arXiv:1311.2544 [gr-qc].

- [4] Michael Pürrer, “Frequency domain reduced order model of aligned-spin effective-one-body waveforms with generic mass-ratios and spins,” *Phys. Rev.* **D93**, 064041 (2016), arXiv:1512.02248 [gr-qc].
- [5] A. Buonanno and T. Damour, “Effective one-body approach to general relativistic two-body dynamics,” *Phys. Rev.* **D59**, 084006 (1999), arXiv:gr-qc/9811091 [gr-qc].
- [6] Alessandra Buonanno and Thibault Damour, “Transition from inspiral to plunge in binary black hole coalescences,” *Phys. Rev.* **D62**, 064015 (2000), arXiv:gr-qc/0001013 [gr-qc].
- [7] Thibault Damour, Piotr Jaranowski, and Gerhard Schafer, “On the determination of the last stable orbit for circular general relativistic binaries at the third postNewtonian approximation,” *Phys. Rev.* **D62**, 084011 (2000), arXiv:gr-qc/0005034 [gr-qc].
- [8] Thibault Damour, “Coalescence of two spinning black holes: an effective one-body approach,” *Phys. Rev.* **D64**, 124013 (2001), arXiv:gr-qc/0103018 [gr-qc].
- [9] Thibault Damour, Piotr Jaranowski, and Gerhard Schäfer, “Fourth post-Newtonian effective one-body dynamics,” *Phys. Rev.* **D91**, 084024 (2015), arXiv:1502.07245 [gr-qc].
- [10] Thibault Damour, Piotr Jaranowski, and Gerhard Schäfer, “Conservative dynamics of two-body systems at the fourth post-Newtonian approximation of general relativity,” *Phys. Rev.* **D93**, 084014 (2016), arXiv:1601.01283 [gr-qc].
- [11] Félix-Louis Julié and Nathalie Deruelle, “Two-body problem in Scalar-Tensor theories as a deformation of General Relativity : an Effective-One-Body approach,” (2017), arXiv:1703.05360 [gr-qc].
- [12] Francesco Messina and Alessandro Nagar, “Back to post-Newtonian: parametrized-4.5PN TaylorF2 approximant(s) and tail effects to quartic nonlinear order from the effective one body formalism,” (2017), arXiv:1703.08107 [gr-qc].
- [13] Thibault Damour, “Gravitational scattering, post-Minkowskian approximation and Effective One-Body theory,” *Phys. Rev.* **D94**, 104015 (2016), arXiv:1609.00354 [gr-qc].
- [14] Donato Bini, Thibault Damour, and Andrea Geralico, “Confirming and improving post-Newtonian and effective-one-body results from self-force computations along eccentric orbits around a Schwarzschild black hole,” *Phys. Rev.* **D93**, 064023 (2016), arXiv:1511.04533 [gr-qc].
- [15] Seth Hopper, Chris Kavanagh, and Adrian C. Ottewill, “Analytic self-force calculations in the post-Newtonian regime: eccentric orbits on a Schwarzschild background,” *Phys. Rev.* **D93**, 044010 (2016), arXiv:1512.01556 [gr-qc].
- [16] Sarp Akcay and Maarten van de Meent, “Numerical computation of the effective-one-body potential q using self-force results,” *Phys. Rev.* **D93**, 064063 (2016), arXiv:1512.03392 [gr-qc].
- [17] Donato Bini and Thibault Damour, “Conservative second-order gravitational self-force on circular orbits and the effective one-body formalism,” *Phys. Rev.* **D93**, 104040 (2016), arXiv:1603.09175 [gr-qc].
- [18] Alessandro Nagar, Thibault Damour, Christian Reisswig, and Denis Pollney, “Energetics and phasing of non-precessing spinning coalescing black hole binaries,” *Phys. Rev.* **D93**, 044046 (2016), arXiv:1506.08457 [gr-qc].
- [19] Alejandro Bohé *et al.*, “Improved effective-one-body model of spinning, nonprecessing binary black holes for the era of gravitational-wave astrophysics with advanced detectors,” *Phys. Rev.* **D95**, 044028 (2017), arXiv:1611.03703 [gr-qc].
- [20] Stanislav Babak, Andrea Taracchini, and Alessandra Buonanno, “Validating the effective-one-body model of spinning, precessing binary black holes against numerical relativity,” *Phys. Rev.* **D95**, 024010 (2017), arXiv:1607.05661 [gr-qc].
- [21] Tim Dietrich and Tanja Hinderer, “Comprehensive numerical relativity – effective-one-body comparison to inform improvements in waveform models for binary neutron star systems,” (2017), arXiv:1702.02053 [gr-qc].
- [22] Alessandro Nagar, Gunnar Riemenschneider, and Geraint Pratten, “Impact of Numerical Relativity information on effective-one-body waveform models,” (2017), arXiv:1703.06814 [gr-qc].
- [23] Steven L. Detweiler, “A Consequence of the gravitational self-force for circular orbits of the Schwarzschild geometry,” *Phys. Rev.* **D77**, 124026 (2008), arXiv:0804.3529 [gr-qc].
- [24] Leor Barack and Norichika Sago, “Gravitational self-force correction to the innermost stable circular orbit of a Schwarzschild black hole,” *Phys. Rev. Lett.* **102**, 191101 (2009), arXiv:0902.0573 [gr-qc].
- [25] Thibault Damour, “Gravitational Self Force in a Schwarzschild Background and the Effective One Body Formalism,” *Phys. Rev.* **D81**, 024017 (2010), arXiv:0910.5533 [gr-qc].
- [26] Enrico Barausse, Alessandra Buonanno, and Alexandre Le Tiec, “The complete non-spinning effective-one-body metric at linear order in the mass ratio,” *Phys. Rev.* **D85**, 064010 (2012), arXiv:1111.5610 [gr-qc].
- [27] Marta Colleoni, Leor Barack, Abhay G. Shah, and Maarten van de Meent, “Self-force as a cosmic censor in the Kerr overspinning problem,” *Phys. Rev.* **D92**, 084044 (2015), arXiv:1508.04031 [gr-qc].
- [28] Norichika Sago, Leor Barack, and Steven L. Detweiler, “Two approaches for the gravitational self force in black hole spacetime: Comparison of numerical results,” *Phys. Rev.* **D78**, 124024 (2008), arXiv:0810.2530 [gr-qc].
- [29] Luc Blanchet, Steven L. Detweiler, Alexandre Le Tiec, and Bernard F. Whiting, “Post-Newtonian and Numerical Calculations of the Gravitational Self-Force for Circular Orbits in the Schwarzschild Geometry,” *Phys. Rev.* **D81**, 064004 (2010), arXiv:0910.0207 [gr-qc].
- [30] Sarp Akcay, Alexandre Le Tiec, Leor Barack, Norichika Sago, and Niels Warburton, “Comparison Between Self-Force and Post-Newtonian Dynamics: Beyond Circular Orbits,” *Phys. Rev.* **D91**, 124014 (2015), arXiv:1503.01374 [gr-qc].
- [31] Maarten van de Meent, “Self-force corrections to the periastron advance around a spinning black hole,” *Phys. Rev. Lett.* **118**, 011101 (2017), arXiv:1610.03497 [gr-qc].
- [32] Alexandre Le Tiec, Abdul H. Mroue, Leor Barack, Alessandra Buonanno, Harald P. Pfeiffer, Norichika Sago, and Andrea Taracchini, “Periastron Advance in Black Hole Binaries,” *Phys. Rev. Lett.* **107**, 141101 (2011), arXiv:1106.3278 [gr-qc].
- [33] Donato Bini and Thibault Damour, “Analytical determination of the two-body gravitational interaction potential at the fourth post-Newtonian approximation,” *Phys. Rev.* **D87**, 121501 (2013), arXiv:1305.4884 [gr-qc].

- [34] Donato Bini and Thibault Damour, “Two-body gravitational spin-orbit interaction at linear order in the mass ratio,” *Phys. Rev.* **D90**, 024039 (2014), arXiv:1404.2747 [gr-qc].
- [35] Chris Kavanagh, Adrian C. Ottewill, and Barry Wardell, “Analytical high-order post-Newtonian expansions for extreme mass ratio binaries,” *Phys. Rev.* **D92**, 084025 (2015), arXiv:1503.02334 [gr-qc].
- [36] Sarp Akcay, David Dempsey, and Sam Dolan, “Spin-orbit precession for eccentric black hole binaries at first order in the mass ratio,” *Class. Quant. Grav.* **34**, 084001 (2017), arXiv:1608.04811 [gr-qc].
- [37] Steven L. Detweiler and Bernard F. Whiting, “Selfforce via a Green’s function decomposition,” *Phys. Rev.* **D67**, 024025 (2003), arXiv:gr-qc/0202086 [gr-qc].
- [38] C. Darwin, “The Gravity Field of a Particle,” *Proc. R. Soc. Lond. A* **249**, 180–194 (1959).
- [39] J.-A. Marck, “Solution to the equations of parallel transport in kerr geometry; tidal tensor,” *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* **385**, 431–438 (1983), <http://rspa.royalsocietypublishing.org/content/385/1789/431.full.pdf>.
- [40] Leor Barack and Norichika Sago, “Beyond the geodesic approximation: conservative effects of the gravitational self-force in eccentric orbits around a Schwarzschild black hole,” *Phys. Rev.* **D83**, 084023 (2011), arXiv:1101.3331 [gr-qc].
- [41] P. L. Chrzanowski, “Vector Potential and Metric Perturbations of a Rotating Black Hole,” *Phys. Rev.* **D11**, 2042–2062 (1975).
- [42] J.M. Cohen and L.S. Kegeles, “Space-time perturbations,” *Physics Letters A* **54**, 5 – 7 (1975).
- [43] L. S. Kegeles and J. M. Cohen, “Constructive procedure for perturbations of space-times,” *Phys. Rev.* **D19**, 1641–1664 (1979).
- [44] Carlos O. Lousto and Bernard F. Whiting, “Reconstruction of black hole metric perturbations from Weyl curvature,” *Phys. Rev.* **D66**, 024026 (2002), arXiv:gr-qc/0203061 [gr-qc].
- [45] Tobias S. Keidl, Abhay G. Shah, John L. Friedman, Dong-Hoon Kim, and Larry R. Price, “Gravitational Self-force in a Radiation Gauge,” *Phys. Rev.* **D82**, 124012 (2010), [Erratum: *Phys. Rev.* **D90**, no.10, 109902 (2014)], arXiv:1004.2276 [gr-qc].
- [46] Abhay G. Shah, Tobias S. Keidl, John L. Friedman, Dong-Hoon Kim, and Larry R. Price, “Conservative, gravitational self-force for a particle in circular orbit around a Schwarzschild black hole in a Radiation Gauge,” *Phys. Rev.* **D83**, 064018 (2011), arXiv:1009.4876 [gr-qc].
- [47] Chris Kavanagh, Adrian C. Ottewill, and Barry Wardell, “Analytical high-order post-Newtonian expansions for spinning extreme mass ratio binaries,” *Phys. Rev.* **D93**, 124038 (2016), arXiv:1601.03394 [gr-qc].
- [48] Maarten van de Meent and Abhay G. Shah, “Metric perturbations produced by eccentric equatorial orbits around a Kerr black hole,” *Phys. Rev.* **D92**, 064025 (2015), arXiv:1506.04755 [gr-qc].
- [49] Donato Bini and Thibault Damour, “High-order post-Newtonian contributions to the two-body gravitational interaction potential from analytical gravitational self-force calculations,” *Phys. Rev.* **D89**, 064063 (2014), arXiv:1312.2503 [gr-qc].
- [50] Barry Wardell and Achamveedu Gopakumar, “Self-force: Computational Strategies,” *Proceedings, 524th WE-Heraeus-Seminar: Equations of Motion in Relativistic Gravity (EOM 2013): Bad Honnef, Germany, February 17-23, 2013*, *Fund. Theor. Phys.* **179**, 487–522 (2015), arXiv:1501.07322 [gr-qc].
- [51] Tanja Hinderer and Eanna E. Flanagan, “Two timescale analysis of extreme mass ratio inspirals in Kerr. I. Orbital Motion,” *Phys. Rev.* **D78**, 064028 (2008), arXiv:0805.3337 [gr-qc].
- [52] Leor Barack and Norichika Sago, “Gravitational self-force on a particle in eccentric orbit around a Schwarzschild black hole,” *Phys. Rev.* **D81**, 084021 (2010), arXiv:1002.2386 [gr-qc].
- [53] Barry Wardell and Niels Warburton, “Applying the effective-source approach to frequency-domain self-force calculations: Lorenz-gauge gravitational perturbations,” *Phys. Rev.* **D92**, 084019 (2015), arXiv:1505.07841 [gr-qc].
- [54] Leor Barack and Amos Ori, “Gravitational selfforce and gauge transformations,” *Phys. Rev.* **D64**, 124003 (2001), arXiv:gr-qc/0107056 [gr-qc].
- [55] Adam Pound, Cesar Merlin, and Leor Barack, “Gravitational self-force from radiation-gauge metric perturbations,” *Phys. Rev.* **D89**, 024009 (2014), arXiv:1310.1513 [gr-qc].
- [56] Leor Barack and Amos Ori, “Mode sum regularization approach for the selfforce in black hole space-time,” *Phys. Rev.* **D61**, 061502 (2000), arXiv:gr-qc/9912010 [gr-qc].
- [57] Anna Heffernan, Adrian Ottewill, and Barry Wardell, “High-order expansions of the Detweiler-Whiting singular field in Schwarzschild spacetime,” *Phys. Rev.* **D86**, 104023 (2012), arXiv:1204.0794 [gr-qc].
- [58] “Electronic archive of post-Newtonian coefficients,” <http://www.barrywardell.net/research/code>.
- [59] Donato Bini, Thibault Damour, and Andrea Geralico, “New gravitational self-force analytical results for eccentric orbits around a Schwarzschild black hole,” *Phys. Rev.* **D93**, 104017 (2016), arXiv:1601.02988 [gr-qc].
- [60] Thibault Damour, Piotr Jaranowski, and Gerhard Schafer, “Effective one body approach to the dynamics of two spinning black holes with next-to-leading order spin-orbit coupling,” *Phys. Rev.* **D78**, 024009 (2008), arXiv:0803.0915 [gr-qc].
- [61] Enrico Barausse, Etienne Racine, and Alessandra Buonanno, “Hamiltonian of a spinning test-particle in curved spacetime,” *Phys. Rev.* **D80**, 104025 (2009), [Erratum: *Phys. Rev.* **D85**, 069904 (2012)], arXiv:0907.4745 [gr-qc].
- [62] Enrico Barausse and Alessandra Buonanno, “An Improved effective-one-body Hamiltonian for spinning black-hole binaries,” *Phys. Rev.* **D81**, 084024 (2010), arXiv:0912.3517 [gr-qc].
- [63] Alessandro Nagar, “Effective one body Hamiltonian of two spinning black-holes with next-to-next-to-leading order spin-orbit coupling,” *Phys. Rev.* **D84**, 084028 (2011), [Erratum: *Phys. Rev.* **D88**, no.8, 089901 (2013)], arXiv:1106.4349 [gr-qc].
- [64] Enrico Barausse and Alessandra Buonanno, “Extending the effective-one-body Hamiltonian of black-hole binaries to include next-to-next-to-leading spin-orbit couplings,” *Phys. Rev.* **D84**, 104027 (2011), arXiv:1107.2904 [gr-qc].
- [65] Andrea Taracchini, Yi Pan, Alessandra Buonanno, Enrico Barausse, Michael Boyle, Tony Chu, Geoffrey Lovelace, Harald P. Pfeiffer, and Mark A. Scheel, “Proto-

- type effective-one-body model for nonprecessing spinning inspiral-merger-ringdown waveforms,” *Phys. Rev.* **D86**, 024011 (2012), arXiv:1202.0790 [gr-qc].
- [66] Thibault Damour and Alessandro Nagar, “New effective-one-body description of coalescing nonprecessing spinning black-hole binaries,” *Phys. Rev.* **D90**, 044018 (2014), arXiv:1406.6913 [gr-qc].
- [67] Simone Balmelli and Philippe Jetzer, “Effective-one-body Hamiltonian with next-to-leading order spin-spin coupling,” *Phys. Rev.* **D91**, 064011 (2015), arXiv:1502.01343 [gr-qc].
- [68] Donato Bini and Thibault Damour, “Analytic determination of high-order post-Newtonian self-force contributions to gravitational spin precession,” *Phys. Rev.* **D91**, 064064 (2015), arXiv:1503.01272 [gr-qc].
- [69] Simone Balmelli and Thibault Damour, “New effective-one-body Hamiltonian with next-to-leading order spin-spin coupling,” *Phys. Rev.* **D92**, 124022 (2015), arXiv:1509.08135 [gr-qc].
- [70] Donato Bini, Thibault Damour, and Andrea Geralico, “Spin-dependent two-body interactions from gravitational self-force computations,” *Phys. Rev.* **D92**, 124058 (2015), [Erratum: *Phys. Rev.* D93,no.10,109902(2016)], arXiv:1510.06230 [gr-qc].
- [71] Thibault Damour, Piotr Jaranowski, and Gerhard Schafer, “Hamiltonian of two spinning compact bodies with next-to-leading order gravitational spin-orbit coupling,” *Phys. Rev.* **D77**, 064032 (2008), arXiv:0711.1048 [gr-qc].
- [72] Sam R. Dolan, Niels Warburton, Abraham I. Harte, Alexandre Le Tiec, Barry Wardell, and Leor Barack, “Gravitational self-torque and spin precession in compact binaries,” *Phys. Rev.* **D89**, 064011 (2014), arXiv:1312.0775 [gr-qc].
- [73] Sarp Akcay, Leor Barack, Thibault Damour, and Norichika Sago, “Gravitational self-force and the effective-one-body formalism between the innermost stable circular orbit and the light ring,” *Phys. Rev.* **D86**, 104041 (2012), arXiv:1209.0964 [gr-qc].
- [74] Luc Blanchet, Steven L. Detweiler, Alexandre Le Tiec, and Bernard F. Whiting, “High-Order Post-Newtonian Fit of the Gravitational Self-Force for Circular Orbits in the Schwarzschild Geometry,” *Phys. Rev.* **D81**, 084033 (2010), arXiv:1002.0726 [gr-qc].
- [75] Leor Barack, Thibault Damour, and Norichika Sago, “Precession effect of the gravitational self-force in a Schwarzschild spacetime and the effective one-body formalism,” *Phys. Rev.* **D82**, 084036 (2010), arXiv:1008.0935 [gr-qc].
- [76] Donato Bini and Thibault Damour, “Analytic determination of the eight-and-a-half post-Newtonian self-force contributions to the two-body gravitational interaction potential,” *Phys. Rev.* **D89**, 104047 (2014), arXiv:1403.2366 [gr-qc].
- [77] Donato Bini and Thibault Damour, “Detweiler’s gauge-invariant redshift variable: Analytic determination of the nine and nine-and-a-half post-Newtonian self-force contributions,” *Phys. Rev.* **D91**, 064050 (2015), arXiv:1502.02450 [gr-qc].
- [78] T. Damour and Gerhard Schafer, “Higher Order Relativistic Periastron Advances and Binary Pulsars,” *Nuovo Cim.* **B101**, 127 (1988).