

A simple test for stability of black hole by S -deformation

Masashi Kimura

*CENTRA, Departamento de Física, Instituto Superior Técnico,
Universidade de Lisboa, Avenida Rovisco Pais 1, 1049 Lisboa, Portugal*

(Dated: March 12, 2022)

We study a sufficient condition to prove the stability of a black hole when the master equation for linear perturbation takes the form of the Schrödinger equation. If the potential contains a small negative region, usually, the S -deformation method was used to show the non-existence of unstable mode. However, in some cases, it is hard to find an appropriate deformation function analytically because the only way known so far to find it is a try-and-error approach. In this paper, we show that it is easy to find a regular deformation function by numerically solving the differential equation such that the deformed potential vanishes everywhere, when the spacetime is stable. Even if the spacetime is almost marginally stable, our method still works. We also discuss a simple toy model which can be solved analytically, and show the condition for the non-existence of a bound state is the same as that for the existence of a regular solution for the differential equation in our method. From these results, we conjecture that our criteria is also a necessary condition.

PACS numbers: 04.50.-h, 04.70.Bw

I. INTRODUCTION

It is known that the equations for gravitational perturbation around a highly symmetric black hole spacetime usually reduce to decoupled master equations [1–15] in the form

$$\left[-\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} - V(x) \right] \tilde{\Phi} = 0. \quad (1)$$

Using the Fourier transformation with respect to time coordinate, $\tilde{\Phi}(t, x) = e^{-i\omega t} \Phi(x)$, the master equation takes the form of the Schrödinger equation

$$\left[-\frac{d^2}{dx^2} + V \right] \Phi = \omega^2 \Phi. \quad (2)$$

When we wish to prove the stability of the black hole spacetime, we need to show the non-existence of $\omega^2 < 0$ solution, *i.e.*, non-existence of exponentially growing solution, under the boundary conditions,¹ $\Phi \rightarrow 0, d\Phi/dx \rightarrow 0$ at $x \rightarrow \pm\infty$, and the conditions: Φ and $d\Phi/dx$ are continuous and bounded everywhere. From Eq.(2), we obtain

$$-\left[\bar{\Phi} \frac{d\Phi}{dx} \right]_{-\infty}^{\infty} + \int dx \left[\left| \frac{d\Phi}{dx} \right|^2 + V |\Phi|^2 \right] = \omega^2 \int dx |\Phi|^2, \quad (3)$$

where $\bar{\Phi}$ is the complex conjugate of Φ . The first term in LHS vanishes from the above boundary conditions. If the effective potential V is non-negative everywhere, this implies $\omega^2 \geq 0$.

In some cases, even if the effective potential contains a small negative region, the space-time still can be shown to be stable against linear perturbation in a following way. From Eq.(2), we can also show

$$-\frac{d}{dx} \left[\bar{\Phi} \frac{d\Phi}{dx} + S |\Phi|^2 \right] + \left| \frac{d\Phi}{dx} + S \Phi \right|^2 + \left(V + \frac{dS}{dx} - S^2 \right) |\Phi|^2 = \omega^2 |\Phi|^2, \quad (4)$$

where S is an arbitrary function of x . We impose continuity on S everywhere, then the integral

$$\int dx \frac{d}{dx} (S |\Phi|^2), \quad (5)$$

¹ In this paper, since we mainly focus on asymptotically flat (or de Sitter) black holes, we assume that the range of x , which is the tortoise coordinate, is $-\infty < x < \infty$. Here, $x \rightarrow -\infty$ and $x \rightarrow \infty$ correspond to the black hole horizon and the spatial infinity (or cosmological horizon), respectively. If we consider asymptotically anti-de Sitter black holes, the range of x becomes $-\infty < x < x_{\max}$ with the finite value x_{\max} .

becomes a surface term. Integrating Eq.(4), we obtain

$$-\left[\bar{\Phi}\frac{d\Phi}{dx} + S|\Phi|^2\right]_{-\infty}^{\infty} + \int dx \left[\left|\frac{d\Phi}{dx} + S\Phi\right|^2 + \left(V + \frac{dS}{dx} - S^2\right)|\Phi|^2\right] = \omega^2 \int dx |\Phi|^2, \quad (6)$$

Also, we assume that S is not divergent at $x \rightarrow \pm\infty$ so that the above surface term vanishes only from the boundary condition for Φ . The potential term is deformed as

$$\tilde{V} := V + \frac{dS}{dx} - S^2. \quad (7)$$

This is called the S -deformation of the potential V . If the deformed effective potential \tilde{V} is non-negative everywhere by choosing an appropriate function S , we can also say $\omega^2 \geq 0$. In [5–10, 13, 16, 17], the stability of spacetime was shown analytically by using this S -deformation method. However, the only way known so far to find an appropriate function S analytically is a try-and-error approach. When it is hard to find an appropriate S deformation, a numerical study is needed. In [18–23], stability and instability were investigated by numerically solving the two dimensional partial differential equation (1) in time domain.

In this paper, we propose a simple way to show the stability by showing the existence of the S -deformation such that the deformed potential \tilde{V} vanishes. For this purpose, we need to solve the equation

$$V + \frac{dS}{dx} - S^2 = 0. \quad (8)$$

The approximate solution near $x \rightarrow \pm\infty$ is $S \simeq -1/(c_{\pm} - x)$ with constants c_{\pm} if the potential rapidly decays to zero. Since Eq.(8) is a first order ordinary differential equation, all solutions near $x \rightarrow \pm\infty$ should behave this approximate solution. To obtain a solution with finite S at $x \rightarrow \pm\infty$, we need to find a boundary condition so that S is positive near $x \rightarrow -\infty$ and S is negative near $x \rightarrow \infty$, then S behaves $-1/x$ and finite at $x \rightarrow \pm\infty$. While S diverges at some point for an inappropriate boundary condition, we can find the continuous range of appropriate boundary conditions, which corresponds to bounded S , for typical examples.

One may think that to solve the equation $V + dS/dx - S^2 = W$ for a positive function $W(> 0)$ is easier than to solve Eq.(9). However, since this equation can be written in the form $(V - W) + dS/dx - S^2 = 0$, the problem is to find a solution of Eq.(9) for a deeper potential $V - W$, *i.e.*, more dangerous case. This suggests that to solve Eq.(9) is the most efficient way to find an appropriate function S .

This paper is organized as follows. In the next section, we discuss the existence of the solution of Eq.(8) in the case of positive potential, and that in a simple toy model which can be solved analytically. We also discuss the relation between the solution of Eq.(8) in marginally stable case and the solution of Eq.(2) with $\omega^2 = 0$. In Sec.III, we numerically solve Eq.(8) for the higher-dimensional spherically-symmetric black holes, and construct appropriate deformation functions from various boundary conditions. Sec.IV is devoted to summary and discussion.

II. ON EXISTENCE OF SOLUTION

A. Local existence

Let us consider the differential equation

$$V(x) + \frac{dS(x)}{dx} - S(x)^2 = 0. \quad (9)$$

We assume that V is continuous and bounded in $-\infty < x < \infty$. From the uniqueness theorem of the ordinary differential equation, there exists a solution of the above equation, at least, locally.

We can see this by considering the Taylor expansion if S and V are analytic functions. The series around some point $x = x_0$ are

$$S = \sum_{n=0}^{\infty} s_n (x - x_0)^n, \quad V = \sum_{n=0}^{\infty} v_n (x - x_0)^n, \quad (10)$$

From the differential equation, we obtain the coefficients of S as

$$s_{n+1} = \frac{1}{n+1} \left[-v_n + \sum_{m=0}^n s_m s_{n-m} \right], \quad (11)$$

where s_0 is an arbitrary constant which corresponds to the integration constant (or boundary condition). This shows that we can find a solution S locally.

In general, while $dS/dx - S^2$ is finite, S might be divergent at a finite coordinate value x . The problem is to find a function S which is continuous and bounded everywhere. In some cases, we can show the existence of such the regular function S .




FIG. 1. The relation among $\pm\sqrt{V}$, the value of S and the sign of dS/dx in $V > 0$ region. $dS/dx < 0$ if S is in $-\sqrt{V} < S < \sqrt{V}$ (shaded region), and $dS/dx > 0$ if S is in $S < \sqrt{V}$ or $\sqrt{V} < S$ (white region).

B. Positive potential

First, we consider that the potential is positive and bounded above in $-\infty < x < \infty$. While this corresponds to the manifestly stable case, to show the existence of a continuous and bounded solution of Eq.(9) is not trivial. We would like to show the following proposition.

Proposition 1. *If the potential V is positive and bounded above in $-\infty < x < \infty$, there exists a continuous and bounded solution of Eq.(9) in $-\infty < x < \infty$.*

The proof is given in Appendix.A. We can understand this proposition as follows: Since we have $dS/dx = (S - \sqrt{V})(S + \sqrt{V})$, the relation among $\pm\sqrt{V}$, the value of S and the sign of dS/dx becomes like Fig.1. If we choose the value of S in $-\sqrt{V}|_{x=x_0} < S|_{x=x_0} < \sqrt{V}|_{x=x_0}$ at some point $x = x_0$ as a boundary condition, we can say that the solution of Eq.(9) satisfies $-\sqrt{V_{\max}} < S < \sqrt{V_{\max}}$. Thus, S is bounded above and below in the region $-\infty < x < \infty$.

C. Toy model

To obtain a qualitative understanding, we consider a toy model of the potential, which can be solved analytically,

$$V = \begin{cases} 0 & (-\infty < x \leq x_1) \\ -h_1^2 (< 0) & (x_1 < x \leq x_2) \\ h_2^2 (> 0) & (x_2 < x \leq x_3) \\ 0 & (x_3 < x < \infty) \end{cases} \quad (12)$$

with constants $h_1 > 0, h_2 > 0$. The local solutions of Eq.(9) are²

$$S = \begin{cases} \frac{1}{c_1 - x}, & (-\infty < x \leq x_1) \\ h_1 \tan(h_1 x + c_2), & (x_1 < x \leq x_2) \\ -h_2 \tanh(h_2 x + c_3), & (x_2 < x \leq x_3) \\ \frac{1}{c_4 - x}, & (x_3 < x < \infty) \end{cases} \quad (13)$$

where c_1, c_2, c_3, c_4 are integration constants. From the continuity of S at $x = x_1, x_2, x_3$, we obtain the conditions

$$\frac{1}{c_1 - x_1} - h_1 \tan(c_2 + h_1 x_1) = 0, \quad (14)$$

$$h_1 \tan(c_2 + h_1 x_2) + h_2 \tanh(c_3 + h_2 x_2) = 0, \quad (15)$$

$$\frac{1}{-c_4 + x_3} - h_2 \tanh(c_3 + h_2 x_3) = 0. \quad (16)$$

From the conditions $S|_{x=x_1} \geq 0, S|_{x=x_2} > 0, S|_{x=x_3} \leq 0$ and the finiteness of S , we obtain the inequalities

$$0 < c_1 - x_1, \quad 0 \leq h_1 x_1 + c_2 < \frac{\pi}{2}, \quad (17)$$

$$0 < h_1 x_2 + c_2 < \frac{\pi}{2}, \quad h_2 x_2 + c_3 < 0, \quad (18)$$

$$0 \leq h_2 x_3 + c_3, \quad c_4 - x_3 < 0. \quad (19)$$

² Defining $\alpha := e^{2c_3}$, then S in $x_2 < x \leq x_3$ becomes $S = -h_2(-\alpha + e^{2h_2-x})/(\alpha + e^{2h_2-x})$. We can see that $\alpha < 0$ case corresponds a complex value of c_3 , and S might be divergent in that case. The solutions with $\alpha < 0$ satisfy $S^2 > h_2^2$. To construct continuous S in this subsection, we need to consider $S^2 < h_2^2$ (*i.e.*, real c_3) so that S can take both negative and positive values in $x_2 < x \leq x_3$.




FIG. 2. The effective potential V (thick black) and the typical solution S (solid blue) for the parameter of the potential $x_1 = 0, x_2 = 1, x_3 = 2, h_1 = 1, h_2 = 2$ and the boundary condition for S as $S|_{x=1.9} = 0$.

If the above equations and inequalities are satisfied, then S is continuous and bounded everywhere. We plot the typical profile of V and S in Fig.2. Note that the derivative of S is not necessary to be continuous, because we only impose the condition such that Eq.(5) becomes a surface term.

We would like to derive an inequality between the areas of the potential in negative and positive regions. From the above matching conditions and inequalities, we can show³

$$h_1 \tan(h_1(x_2 - x_1)) \leq h_2 \tanh(h_2(x_3 - x_2)) \quad (20)$$

Since we have $X < \tan X$ for $0 < X < \pi/2$, and $\tanh Y < Y$ for $0 < Y$, we obtain an inequality

$$h_1^2(x_2 - x_1) < h_2^2(x_3 - x_2). \quad (21)$$

The area of the negative region is smaller than that of the positive region. This is consistent with the result in [24] where the existence of an unstable mode is shown if $\int_{-\infty}^{\infty} dx V < 0$.

³ The derivation is as follows:

$$\begin{aligned} h_1 \tan(h_1(x_2 - x_1)) &\leq h_1 \tan(c_2 + h_1 x_1 + h_1(x_2 - x_1)) \quad (\because 0 \leq c_2 + h_1 x_1) \\ &= h_1 \tan(c_2 + h_1 x_2) \\ &= -h_2 \tanh(c_3 + h_2 x_2) \quad (\because \text{Eq.(15)}) \\ &= h_2 \tanh(-c_3 - h_2 x_2) \\ &\leq h_2 \tanh(h_2 x_3 - h_2 x_2). \quad (\because 0 \leq h_2 x_3 + c_3) \end{aligned}$$

D. Relation between existence of S and non-existence of bound state in toy model

To check the relation between the existence of regular S and the non-existence of bound state, we further study the toy model in the previous subsection and the dependence of their existence on the ratio of the areas

$$\Gamma = \frac{h_2^2(x_3 - x_2)}{h_1^2(x_2 - x_1)}. \quad (22)$$

First, for simplicity, we consider the case

$$x_1 = 0, \quad x_2 = 1, \quad x_3 = 2, \quad h_1 = 1. \quad (23)$$

If $\Gamma = 0$, this is just a single well problem. In that case, there is only a bound state in the solution of the Schrödinger equation whose energy is $\omega^2|_{\Gamma=0} \simeq -0.43$. After some calculation, we obtain the condition for the existence of the bound state, which can be calculated from $\omega^2 < 0$, as

$$\Gamma < \Gamma_{\text{cr}} \quad (24)$$

where $\Gamma_{\text{cr}} \simeq 2.79$ is defined by

$$\tan(1) = \sqrt{\Gamma_{\text{cr}}} \tanh(\sqrt{\Gamma_{\text{cr}}}). \quad (25)$$

Also, we can calculate the condition for the existence of continuous and bounded solution S of Eq.(9). That condition, which is derived from $S|_{x=x_1} \geq 0$ with $S|_{x=x_3} = 0$, becomes $\Gamma \geq \Gamma_{\text{cr}}$ with the same critical value in Eq.(25).

For general parameters, we can derive the same relations. The critical value Γ_{cr} in general case is defined by ⁴

$$\frac{\sqrt{x_3 - x_2}}{\sqrt{x_2 - x_1}} \tan(h_1(x_2 - x_1)) = \sqrt{\Gamma_{\text{cr}}} \tanh(h_1 \sqrt{\Gamma_{\text{cr}}} \sqrt{x_2 - x_1} \sqrt{x_3 - x_2}). \quad (26)$$

Note that if $h_1(x_2 - x_1) \geq \pi/2$, there exists at least one bound state regardless of the value of h_2 , in that case, Γ_{cr} is not defined. Thus, in this toy model, the condition for the non-existence of bound state coincides with that for the existence of continuous and bounded solution S .

⁴ From $X < \tan X$ for $0 < X < \pi/2$, and $\tanh Y < Y$ for $0 < Y$, we can see $\Gamma_{\text{cr}} > 1$. If the arguments of both tangent and hyperbolic tangent are small, $\Gamma_{\text{cr}} \simeq 1$.

E. Relation between regular S in marginally stable case and onset of unstable mode

Since Eq.(9) is the Riccati equation, defining

$$\frac{1}{\phi} \frac{d\phi}{dx} := -S, \quad (27)$$

we can write Eq.(9) in a second order linear ordinary differential equation

$$-\frac{d^2\phi}{dx^2} + V\phi = 0. \quad (28)$$

This is the master equation (2) with $\omega^2 = 0$. If this equation has a non-trivial regular solution, it probably corresponds to the onset of unstable mode.⁵ We can expect that there is some relation between the solutions of Eq.(9) in marginally stable case and the onset of unstable mode. In this subsection, we briefly discuss this.

Suppose that the potential contains a parameter α , and the master equation has a property such that there is no unstable mode if $\alpha \leq \alpha_{\text{cr}}$ and there are unstable modes if $\alpha > \alpha_{\text{cr}}$. In the case of $\alpha > \alpha_{\text{cr}}$, we define $\Phi_0(x; \omega(\alpha))$ as the ground state of the Schrödinger equation (2), then the energy takes negative value $\omega^2 < 0$. If we assume that the potential rapidly decays in $x \rightarrow \pm\infty$, then $\Phi_0 \sim e^{\mp x\sqrt{-\omega^2}}$ at $x \rightarrow \pm\infty$. In that case, $\Phi_0^{-1} d\Phi_0/dx$ takes finite values at the boundary $x \rightarrow \pm\infty$. Since it is known that the ground state for the one-dimensional Schrödinger equation has no nodes (for example, see [25, 26]), Φ_0 cannot become zero except at the boundary. So, we can define a function

$$\mathcal{S} := -\frac{1}{\Phi_0(x; \omega(\alpha))} \frac{d\Phi_0(x; \omega(\alpha))}{dx}, \quad (29)$$

and this is regular in $-\infty < x < \infty$. We can show that \mathcal{S} satisfies

$$V + \frac{d\mathcal{S}}{dx} - \mathcal{S}^2 = \omega^2. \quad (30)$$

If we assume the function \mathcal{S} has a smooth limit in $\alpha \rightarrow \alpha_{\text{cr}} + 0$, this becomes a regular solution of Eq.(9) since $\omega^2 \rightarrow 0$ in this limit.

In fact, the toy model in the previous subsections satisfies the above assumptions. This is the reason why the critical values are the same in the toy model. In general, the validity of the above assumption is not clear, but it seems to be reasonable.

⁵ We should note that we cannot construct a regular solution of Eq.(28) from a regular solution of Eq.(9) except for the marginally stable case. This is because the asymptotic behavior of the regular solution of Eq.(9) is $S \sim -1/x$, but the corresponding ϕ behaves $\phi \sim cx$ with a constant c , which is not a regular solution of Eq.(28).




FIG. 3. The effective potential V of the $\ell = 2$ vector perturbation in the 10-Dimensional Schwarzschild black hole (thick black), and the numerical solution S for various boundary conditions (solid blue and dashed red). Solid blue lines correspond to bounded S , solid red lines correspond to unbounded S . The boundary conditions for blue lines are $S = 0$ at $r/r_H = 2, 4, 6, 8$, respectively, and that for red lines are $r_H S = 4$ at $r/r_H = 2, 4, 6, 8$, respectively.

III. NUMERICAL CALCULATION

If the effective potential V rapidly decays as $x \rightarrow \pm\infty$, the approximate solution of Eq.(9) becomes $-1/(c_{\pm} - x)$ with constants c_{\pm} . To obtain a solution with finite S at $x \rightarrow \pm\infty$, we need to find a boundary condition so that S is positive near $x \rightarrow -\infty$ and S is negative near $x \rightarrow \infty$. Then, S behaves $-1/x$ and finite at $x \rightarrow \pm\infty$. A bounded S usually changes the value from positive to negative at some point in $V > 0$ region so that S takes negative value near $x \rightarrow \infty$ like S in Fig.2. We can expect that the appropriate boundary conditions can be found by setting $S = 0$ at a point where V is positive. In this section, we study the stability of 10-dimensional spherically symmetric black holes and the 5-dimensional Schwarzschild black string, as typical examples. The explicit form of the metrics and the master equations, which was derived in [5–7, 21, 28, 29], is given in Appendices.B and C.

In the following examples, we used the function `NDSolve` in *Mathematica* to solve the equation numerically. In a stable case, for an appropriate boundary condition, we could obtain a bounded function S without a special technic, as far as we confirmed. We estimated the numerical error by $\epsilon := (V + dS/dx - S^2)/(|V| + |dS/dx| + |S^2|)$ and confirmed $\epsilon < 10^{-6}$ in the following calculations.

A. 10-Dimensional Schwarzschild black hole

Let us consider the $\ell = 2$ vector perturbation in the 10-Dimensional Schwarzschild black hole. In fact, in this case, the existence of the S -deformation such that the deformed potential is positive is already known [6], but this is still a good example to check that our method works well. In Fig.3, we plot the effective potential, which contains a small negative region near the horizon, and the numerical solution of Eq.(9) for various boundary conditions. Note that we use the radial coordinate r in the black hole spacetime, the relation between r and x is in Appendix.B. If we plot $-1/x$ as a function of r , it seems to be rapidly decrease to zero from a finite value near the horizon since $-1/x \sim -1/\ln(r - r_H)$. There are the two attractors of solutions in $V > 0$ region, and they are almost $\pm\sqrt{V}$. This is because $dS/dx = (S + \sqrt{V})(S - \sqrt{V})$ in $V > 0$ region (see also Fig.1). We can see that the bounded solutions can be found by choosing the boundary condition as $S = 0$ at the points where $V > 0$ (solid blue lines in Fig.3). If we choose the boundary condition of S larger than \sqrt{V} in $V > 0$ region, then S is divergent at some point (dashed red lines in Fig.3).

Unlike the toy model in Sec.II C, the potential is negative near the horizon. Since the potential behaves $V \simeq -a^2(r - r_H) \sim -a^2e^{b^2x}$ with constants $a > 0, b > 0$ near the horizon, it rapidly decays in x coordinate. In that case, if the S takes positive value near the horizon, S will be bounded above and below as x decreases (see Appendix.D).

B. 10-Dimensional Schwarzschild-de Sitter black hole

As an another example, we consider the $\ell = 2$ scalar perturbation in the 10-Dimensional Schwarzschild-de Sitter black hole. In this case, the existence of the S -deformation such that the deformed potential is positive is not known, but there is a numerical proof of stability based on the quasi normal mode [18]. In Fig.4, we plot the effective potential and the numerical solution of Eq.(9) for various boundary conditions when the cosmological constant is $r_H^2\lambda = 0.05$. The effective potential also contains a small negative region near the horizon. We can see that the solutions are bounded above and below if we choose boundary condition as $S = 0$ at the points where $V > 0$ (blue lines in Fig.3). Since our method is different from the previous work [18], this is a complementary result which also supports the stability of the spacetime.




FIG. 4. The effective potential V of the $\ell = 2$ scalar perturbation in the 10-Dimensional Schwarzschild-de Sitter black hole (thick black) with $r_H^2 \lambda = 0.05$, and the numerical solution S for various boundary conditions (solid blue). Solid blue lines correspond to bounded S and the boundary conditions are $S = 0$ at $r/r_H = 2, 2.5, 3$, respectively. Here, we only plot bounded solutions.




FIG. 5. The effective potential V of the $\ell = 2$ scalar gravitational perturbation in the 10-Dimensional Reissner-Nordström-de Sitter black hole (thick black) with the parameter $r_{dS}/r_H = 5/4$, $Q/Q_{\text{extremal}} = 0.74$, and the bounded solution S with the boundary condition $S|_{r=r_{dS}-10^{-3}r_H} = 0$ (solid blue).

C. 10-Dimensional Reissner-Nordström-de Sitter black hole

We consider $\ell = 2$ scalar gravitational perturbation in the 10-Dimensional Reissner-Nordström-de Sitter black hole. In this case, it is known that there exists unstable mode for large values of electric charge and cosmological constant [22, 23, 27]. According to Fig.4 in [22], for the parameter $r_H = 1, r_{dS} = 5/4$, if the electric charge is larger than the critical value around $Q/Q_{\text{extremal}} \simeq 0.75$, the spacetime is unstable. In Fig.5, we plot the effective potential and the numerical solution of Eq.(9) for $r_H = 1, r_{dS} = 5/4, Q/Q_{\text{extremal}} = 0.74$. We




FIG. 6. The effective potential V of the scalar perturbation in the 5-Dimensional Schwarzschild black string (thick black) with the parameter $r_H k = 0.877$, and the bounded solution S with the boundary condition $S|_{r=5r_H} = 0$ (solid blue).

can find a continuous and bounded S which is positive at $r \simeq r_H$ and negative at $r \simeq r_{dS}$ by setting the boundary condition $S|_{r=r_{dS}-10^{-3}r_H} = 0$. We can see that our method still works even in the almost marginally stable case.

D. 5-Dimensional Schwarzschild black string

Finally, we discuss the case of the 5-Dimensional Schwarzschild black string spacetime. This spacetime has an unstable mode (Gregory-Laflamme instability) for a long wave perturbation $k < k_{cr} \simeq 0.876$ [28]. The master equation for the scalar perturbation also takes the Schrödinger form Eq.(2) as shown in [21, 29]. We should note that the effective potential Eq.(C2) is positive in $r > r_H$ if $k > (1 + \sqrt{3})/\sqrt{2} \simeq 1.93$. In Fig.6, we plot the effective potential and the numerical solution of Eq.(9) for $r_H k = 0.877$ with the boundary condition $S|_{r=5r_H} = 0$. In this case, the value of the potential asymptotes to a positive constant at large distance. From the same discussion in the proof of the proposition 1.(see Appendix.A), it is guaranteed that the solution is continuous and bounded in $r > r_{ini} = 5r_H$. Also, our method works for the almost marginally stable mode.

IV. SUMMARY AND DISCUSSION

We have studied a sufficient condition to prove the stability of a black hole when the linear perturbative equation takes the Schrödinger form by showing the S -deformation such that the deformed potential vanishes everywhere. While our method is just a sufficient

condition for the non-existence of bound state $\omega^2 < 0$, it also becomes a necessary condition in a simple toy model. We have also found the numerical solution S for the vector and scalar perturbation on 10-dimensional spherically symmetric black holes, and the scalar perturbation on the 5-dimensional Schwarzschild black string. While S diverges at some point for an inappropriate boundary condition, we found the continuous range of appropriate boundary conditions, which corresponds to bounded S . Furthermore, as shown in Sec.III C and Sec.III D, our method can work even in almost marginally stable cases. From these results, we conjecture the following:

Conjecture 1. *A continuous and bounded solution of Eq.(9) exists in $-\infty < x < \infty$ if and only if there is no bound state ($\omega^2 < 0$ mode) in the Schrödinger equation.*

At least, the present results suggest that our method is a good test for the stability of black hole. As for the existence of the solution, we gave a proof for the positive and bounded potential which corresponds to a manifestly stable case. If the potential V contains negative regions, we can still show the existence from a continuous and bounded solution for a different potential $V_0(\leq V)$, under some assumptions (see Appendix.E). The case of marginally (un)stable parameter was discussed in Sec.II E.

If there exists a regular solution of Eq.(9), the Schrödinger equation (2) becomes

$$\left(-\frac{d}{dx} + S\right) \left(\frac{d}{dx} + S\right) \Phi = \omega^2 \Phi. \quad (31)$$

This is known as the supersymmetric quantum mechanics system where the energy ω^2 is manifestly non-negative. If the above conjecture is correct, the quantum mechanics system which does not have a bound state with negative energy becomes supersymmetric.

ACKNOWLEDGMENTS

The author would like to thank Vitor Cardoso, Tsuyoshi Houri, George Pappas, Jiro Soda and Kentaro Tatsumi for their useful comments. M.K. acknowledges financial support provided under the European Union’s H2020 ERC Consolidator Grant “Matter and strong-field gravity: New frontiers in Einstein’s theory” grant agreement no. MaGRaTh-646597, and under the H2020-MSCA-RISE-2015 Grant No. StronGrHEP-690904.

Appendix A: Proof of the proposition 1

We give a proof of the proposition 1. in Sec.II B.

Proof. We consider to solve Eq.(9) from the boundary condition $S|_{x=x_0} = 0$ at a point $x = x_0$. We already know the local existence of the Eq.(9), so we only need to exclude possibility that S is divergent at some point.⁶ At $x = x_0$, we have $S|_{x=x_0} = 0$ and $dS/dx|_{x=x_0} = -V|_{x=x_0} < 0$, so S takes positive value at $x_0 - \delta_0 < x < x_0$, and negative value at $x_0 < x < x_0 + \delta_0$ for a small constant $\delta_0 > 0$.

First, we consider the region $x_0 < x$. We can say that once S becomes negative, S cannot be zero as x increases in $x_0 < x$. If $S = 0$ at some point $x_1 (> x_0)$ and $S < 0$ in $x_0 < x < x_1$, then $dS/dx|_{x=x_1} = -V|_{x=x_1} < 0$. However, this is a contradiction, because $dS/dx|_{x=x_1} < 0$ implies that S is already positive in $x_1 - \delta_1 < x < x_1$ for a small constant $\delta_1 > 0$. Thus, S is bounded above in the region $x_0 < x$.

We also have $dS/dx = -V + S^2 > -(V_{\max} + \delta_2) + S^2$ for a small constant $\delta_2 > 0$. If the value of S changes from the value larger than $-\sqrt{V_{\max} + \delta_2}$ into the value $-\sqrt{V_{\max} + \delta_2}$ at $x = x_2 (> x_0)$, dS/dx becomes positive at $x = x_2$. However this is a contradiction because, $S|_{x=x_2} = -\sqrt{V_{\max} + \delta_2}$ and $dS/dx|_{x=x_2} > 0$ implies $S > -\sqrt{V_{\max} + \delta_2}$ in $x_2 > x > x_2 - \delta_3$ for a small constant $\delta_3 > 0$. Since δ_2 is an arbitrary constant, S cannot be smaller than $-\sqrt{V_{\max}}$. This shows that S is bounded below in the region $x_0 < x$.

Next, we consider the region $x < x_0$. Defining $\bar{S} := -S$, $\bar{x} := -x$, then Eq.(9) becomes

$$\frac{d\bar{S}}{d\bar{x}} = -V + \bar{S}^2, \quad (\text{A1})$$

then $x = x_0$ corresponds to $\bar{x} = -x_0 =: \bar{x}_0$. We need to show \bar{S} is bounded above and below in $\bar{x}_0 < \bar{x}$, but this is formally the same problem as in the case of $x_0 < x$. Thus, S is bounded above and below in the region $-\infty < x < \infty$. \square

⁶ In fact, we also need to exclude the possibility that S is bounded above and below but it oscillates infinitely many times near some point $x = x_s$ and does not have a limiting point like $\sin(1/(x - x_s))$. However, this does not happen, because if S oscillates infinitely many times in a finite interval, its derivative should be unbounded above and below. In that case $dS/dx - S^2$ also becomes unbounded above and below, but this contradicts with that V is bounded above.

Appendix B: Effective potential for D -dimensional Reissner-Nordström-de Sitter black hole

The metric for the D -dimensional Reissner-Nordström-de Sitter black hole is

$$ds^2 = -f dt^2 + \frac{dr^2}{f} + r^2 d\Omega_{S^n}, \quad (\text{B1})$$

$$f = 1 - \frac{2M}{r^{n-1}} + \frac{Q^2}{r^{2n-2}} - \lambda r^2, \quad (\text{B2})$$

where $n = D - 2$, $\lambda = 2\Lambda/(n(n+1))$ with a cosmological constant Λ , and $d\Omega_{S^n}$ denotes the metric of the unit n -dimensional sphere. As shown in [5–7], the linear gravitational and electromagnetic perturbation reduces to the decoupled single master equations with the same form as Eq.(1). The effective potentials for the vector and scalar gravitational modes ⁷ are

$$V_{\text{vector}} = \frac{f}{r^2} \left[\ell(\ell + n - 1) + \frac{n^2 - 2n}{4} - \frac{n(n-2)}{4} \lambda r^2 + \frac{n(5n-2)Q^2}{4r^{2n-2}} + \frac{1}{r^{n-1}} \left(-\frac{n^2+2}{2} M - \sqrt{(n^2-1)^2 M^2 + 2n(n-1)(\ell+n)(\ell-1)Q^2} \right) \right], \quad (\text{B3})$$

$$V_{\text{scalar}} = \frac{f}{64r^2} \frac{U}{H^2}, \quad (\text{B4})$$

with

$$\begin{aligned} U = & \left[-4n^3(n+2)(n+1)^2(1+m\delta)^2 X^2 + 48n^2(n+1)(n-2)m(1+m\delta)X \right. \\ & - 16(n-2)(n-4)m^2 \left. \right] \lambda r^2 - n^3(3n-2)(n+1)^4 \delta(1+m\delta)^3 X^4 \\ & - 4n^2(n+1)^2(1+m\delta)^2 \left[(n+1)(3n-2)m\delta - n^2 \right] X^3 \\ & + 4(n+1)(1+m\delta) \left[m(n-2)(n-4)(n+1)(m+n^2)\delta + 4n(2n^2-3n+4)m \right. \\ & + n^2(n-2)(n-4)(n+1) \left. \right] X^2 - 16m((n+1)m(-4m+3n^2(n-2))\delta \\ & + 3n(n-4)m + 3n^2(n+1)(n-2))X + 64m^3 + 16n(n+2)m^2, \end{aligned} \quad (\text{B5})$$

$$H = m + \frac{n(n+1)}{2}(1+m\delta)X, \quad (\text{B6})$$

$$X = \frac{2M}{r^{n-1}}, \quad (\text{B7})$$

$$\delta = \frac{\mu - M}{2mM}, \quad \mu = \sqrt{M^2 + \frac{4MQ^2}{(n+1)^2}}, \quad m = \ell(\ell + n - 1) - n, \quad (\text{B8})$$

⁷ The effective potentials for other modes, tensor perturbation and electromagnetic vector and scalar perturbation, can be seen in [5–7].

where ℓ is a positive integer, $\ell \geq 1$ for vector mode, $\ell \geq 0$ for scalar mode. The relation between x and r is $dx = dr/f(r)$. x behaves $x \sim \ln(r - r_H)$ near the horizon. In the coordinate r , Eq.(9) becomes

$$V + f \frac{dS}{dr} - S^2 = 0. \quad (\text{B9})$$

To normalize the all quantities by the radius of the black hole horizon r_H , we set the mass parameter as

$$M = \frac{1}{2} (1 + Q^2 - \lambda), \quad (\text{B10})$$

then the black hole horizon locates at $r = r_H = 1$. Also, setting

$$\lambda = \frac{1 - \rho^{1-n} + Q^2(1 - \rho^{n-1})}{1 - \rho^{-n-1}}, \quad (\text{B11})$$

the location of the de-Sitter horizon becomes $r = r_{dS} = 1/\rho$. The electric charge for the extremal black hole is given by

$$Q_{\text{extremal}}^2 = \frac{(n-1)(1 - \rho^{n+1}) - \rho^2(n+1)(1 - \rho^{n-1})}{(n-1)(1 - \rho^{n+1}) - \rho^{n+1}(n+1)(1 - \rho^{n-1})}, \quad (\text{B12})$$

These useful normalization was used in [18].

Appendix C: Effective potential for 5-dimensional black string

The metric of the 5-dimensional black string is

$$ds^2 = -f dt^2 + \frac{dr^2}{f} + r^2(d\theta^2 + \sin^2 \theta d\phi^2) + dz^2, \quad (\text{C1})$$

with $f = 1 - r_H/r$. The effective potential for the scalar perturbation is given by [21]

$$V = \frac{f}{r^3} \frac{k^6 r^9 + 3k^4 r^6 (2r - 3r_H) + r_H^3 + 3k^2 r^3 r_H (-4r + 3r_H)}{(k^2 r^3 + r_H)^2}, \quad (\text{C2})$$

where k is the wave number along z direction. In this case, Eq.(9) also becomes

$$V + f \frac{dS}{dr} - S^2 = 0. \quad (\text{C3})$$

Appendix D: $V = -a^2 \exp(b^2 x)$ case

In some cases, the potential behaves $V \simeq -a^2(r - r_H)$ with a constant $a > 0$ near the horizon. Since $r - r_H \simeq e^{b^2 x}$ with a constant $b > 0$ near the horizon, the potential in x coordinate becomes $V \simeq -a^2 \exp(b^2 x)$. For this potential, we can solve Eq.(9) by using the Bessel functions

$$S = b^2 X \frac{C_1 J_1(X) + 2Y_1(X)}{2C_1 J_0(X) + 4Y_0(X)}, \quad (\text{D1})$$

where $X = 2ae^{b^2 x/2}/b^2$ and C_1 is a constant. For $X \ll 1$, the Bessel functions behave

$$J_0(X) = 1 + \mathcal{O}(X^2), \quad (\text{D2})$$

$$J_1(X) = \frac{X}{2} + \mathcal{O}(X^3), \quad (\text{D3})$$

$$Y_0(X) = \frac{2}{\pi} \ln X + 2 \frac{\gamma - \ln 2}{\pi} + \mathcal{O}(X^2), \quad (\text{D4})$$

$$Y_1(X) = -\frac{2}{\pi} \frac{1}{X} + X \frac{1}{\pi} \ln X - \frac{1 - 2\gamma + 2 \ln 2}{2\pi} X + \mathcal{O}(X^3), \quad (\text{D5})$$

where $\gamma \simeq 0.577$ is the Euler's constant. So, the approximate solution of S with the integral constant becomes

$$S \simeq -\frac{b^2}{2} \frac{1}{C_2 + \ln X}, \quad (\text{D6})$$

where $C_2 = \gamma - \ln 2 + C_1 \pi/4$. Once S takes positive value in $X \ll 1$ regime, S will be bounded above and below as X decreases towards zero. In that case, near $X \simeq 0$, S behaves $S \sim -1/\ln X \sim -1/x$ like $V = 0$ case.

Appendix E: A potential with compact support

Let us consider a continuous and bounded potential with compact support

$$V = \begin{cases} 0 & (-\infty < x \leq x_1) \\ V_- (< 0) & (x_1 < x < x_2) \\ 0 & (x = x_2) \\ V_+ (> 0) & (x_2 < x < x_3) \\ 0 & (x_3 \leq x < \infty). \end{cases} \quad (\text{E1})$$

We assume that V is smooth in $x_1 < x < x_3$ and $\lim_{x \rightarrow x_3-0} dV/dx > 0$. Also, we define another potential which is not necessary continuous

$$V_0 = \begin{cases} 0 & (-\infty < x \leq x_1) \\ v_- \ (v_- < V_- < 0) & (x_1 < x < x_2) \\ v_+ \ (0 \leq v_+ < V_+) & (x_2 \leq x < x_3) \\ 0 & (x_3 \leq x < \infty), \end{cases} \quad (\text{E2})$$

where v_{\pm} are functions and we assume $V > V_0$ in $x_1 < x < x_3$. In this set up, we would like to prove the following proposition.

Proposition 2. *If there exists a continuous and bounded solution of Eq.(9) for the potential Eq.(E2) in $-\infty < x < \infty$, there also exists a continuous and bounded solution of Eq.(9) for the potential Eq.(E1) in $-\infty < x < \infty$.*

Proof. Let S_0 be a continuous and bounded solution of $V_0 + dS_0/dx - S_0^2 = 0$. Since the behavior in $x \leq x_1$ and $x_3 \leq x$ are the same as the toy model in Sec.IIC, we can say $S_0|_{x=x_1} > 0$ and $S_0|_{x=x_3} < 0$. In the region $x_1 < x < x_2$, since $-V_- < -v_-$, we obtain an inequality

$$(S_0 - S)(S_0 - S + 2S) < \frac{d(S_0 - S)}{dx}. \quad (\text{E3})$$

We assume $S|_{x=x_1} = \eta S_0|_{x=x_1}$ with a positive constant $\eta (< 1)$.⁸ Since $dS/dx = -V_- + S^2 > 0$, S is a monotonically increasing function in $x_1 < x < x_2$. From this and the condition $S|_{x=x_1} = \eta S_0 > 0$, we can say $S > 0$ in $x_1 < x < x_2$. In the limit $x \rightarrow x_1 + 0$, the above inequality becomes

$$(1 - \eta^2)S_0^2 < \frac{d(S_0 - S)}{dx}, \quad (\text{E4})$$

and LHS is positive. Thus $S_0 - S$ is an increasing function near $x = x_1$. Once $S_0 - S$ becomes positive, it cannot be zero in $x_1 < x < x_2$ because $d(S_0 - S)/dx > 0$ for positive $S_0 - S$ in $x_1 < x < x_2$, i.e., $S_0 - S$ cannot decrease as x increases. So, we can say $S < S_0$ in $x_1 < x < x_2$. Since S_0 is bounded above, S cannot be divergent in $x_1 < x < x_2$.

⁸ Note that this is an appropriate boundary condition for S .

In the region $x_2 < x < x_3$, since $-V_+ < -v_+$ we also have the same inequality

$$(S_0 - S)(S_0 - S + 2S) < \frac{d(S_0 - S)}{dx}. \quad (\text{E5})$$

If we assume $S \geq 0$ in $x_2 < r < x_3$, we can also say $S < S_0$ from the same discussion above. However, now we also assumed $S_0 < 0$ at some point because S_0 should be connected with $1/(c_4 - r)$, which is negative, at $x = x_3$, then this contradicts with the first assumption $S > 0$. Thus, we can say $S < 0$ at some point $x = y_0$ which satisfies $x_2 < y_0 < x_3$. In $y_0 < x < x_3$, from the same discussion in the proof of the proposition 1, we can show that S is a continuous and bounded negative function.

There is still a possibility that $S = 0$ at $x = x_3$ where $V_+ = 0$. In that case, $S|_{x=x_3} = 0$, $dS/dx|_{x=x_3} = 0$, and $d^2S/dx^2|_{x=x_3} = \lim_{x \rightarrow x_3-0} dV/dx (= \beta^2) > 0$. So, the approximate behavior becomes $S \simeq \beta^2(x - x_3)^2$, but this means $S > 0$ in $x_3 - \delta < x < x_3$ with a small constant $\delta > 0$. This contradicts with $S < 0$ in $y_0 < x < x_3$. Thus, $S|_{x=x_3} < 0$ and S can be matched with $1/(c_4 - r)$ at $x = x_3$ with a negative value. This shows the existence of continuous and bounded S in $-\infty < x < \infty$. \square

-
- [1] T. Regge and J. A. Wheeler, Phys. Rev. **108**, 1063 (1957).
 - [2] C. V. Vishveshwara, Phys. Rev. D **1**, 2870 (1970).
 - [3] F. J. Zerilli, Phys. Rev. Lett. **24**, 737 (1970).
 - [4] F. J. Zerilli, Phys. Rev. D **2**, 2141 (1970).
 - [5] H. Kodama and A. Ishibashi, Prog. Theor. Phys. **110**, 701 (2003) [hep-th/0305147].
 - [6] A. Ishibashi and H. Kodama, Prog. Theor. Phys. **110**, 901 (2003) [hep-th/0305185].
 - [7] H. Kodama and A. Ishibashi, Prog. Theor. Phys. **111**, 29 (2004) [hep-th/0308128].
 - [8] G. Dotti and R. J. Gleiser, Class. Quant. Grav. **22**, L1 (2005) [gr-qc/0409005].
 - [9] G. Dotti and R. J. Gleiser, Phys. Rev. D **72**, 044018 (2005) [gr-qc/0503117].
 - [10] R. J. Gleiser and G. Dotti, Phys. Rev. D **72**, 124002 (2005) [gr-qc/0510069].
 - [11] T. Takahashi and J. Soda, Prog. Theor. Phys. **124**, 911 (2010) [arXiv:1008.1385 [gr-qc]].
 - [12] T. Takahashi and J. Soda, Phys. Rev. D **79**, 104025 (2009) [arXiv:0902.2921 [gr-qc]].
 - [13] T. Takahashi and J. Soda, Phys. Rev. D **80**, 104021 (2009) [arXiv:0907.0556 [gr-qc]].
 - [14] M. Kimura, K. Murata, H. Ishihara and J. Soda, Phys. Rev. D **77**, 064015 (2008) [arXiv:0712.4202 [hep-th]].

- [15] R. Nishikawa and M. Kimura, *Class. Quant. Grav.* **27**, 215020 (2010) doi:10.1088/0264-9381/27/21/215020 [arXiv:1005.1367 [hep-th]].
- [16] M. Beroiz, G. Dotti and R. J. Gleiser, *Phys. Rev. D* **76**, 024012 (2007) [hep-th/0703074].
- [17] T. Takahashi and J. Soda, *Prog. Theor. Phys.* **124**, 711 (2010) [arXiv:1008.1618 [gr-qc]].
- [18] R. A. Konoplya and A. Zhidenko, *Nucl. Phys. B* **777**, 182 (2007) [hep-th/0703231].
- [19] R. A. Konoplya and A. Zhidenko, *Phys. Rev. D* **77**, 104004 (2008) [arXiv:0802.0267 [hep-th]].
- [20] H. Ishihara, M. Kimura, R. A. Konoplya, K. Murata, J. Soda and A. Zhidenko, *Phys. Rev. D* **77**, 084019 (2008) [arXiv:0802.0655 [hep-th]].
- [21] R. A. Konoplya, K. Murata, J. Soda and A. Zhidenko, *Phys. Rev. D* **78**, 084012 (2008) [arXiv:0807.1897 [hep-th]].
- [22] R. A. Konoplya and A. Zhidenko, *Phys. Rev. Lett.* **103**, 161101 (2009) [arXiv:0809.2822 [hep-th]].
- [23] R. A. Konoplya and A. Zhidenko, *Phys. Rev. D* **89**, no. 2, 024011 (2014) [arXiv:1309.7667 [hep-th]].
- [24] W. F. Buell and B. A. Shadwick, *Am. J. Phys.* **63**, 256. (1995)
- [25] A. Messiah, *Quantum Mechanics Vol.I* (North Holland, 1967)
- [26] M. Moriconi, *Am. J. Phys.* **75**, 284. (2007) [quant-ph/0702260]
- [27] V. Cardoso, M. Lemos and M. Marques, *Phys. Rev. D* **80**, 127502 (2009) [arXiv:1001.0019 [gr-qc]].
- [28] R. Gregory and R. Laflamme, *Phys. Rev. Lett.* **70**, 2837 (1993) [hep-th/9301052].
- [29] J. L. Hovdebo and R. C. Myers, *Phys. Rev. D* **73**, 084013 (2006) [hep-th/0601079].