

$R_{K^{(*)}}$ and related $b \rightarrow s\ell\bar{\ell}$ anomalies in minimal flavor violation framework with Z' boson

Cheng-Wei Chiang,^{1,2,3,4,*} Xiao-Gang He,^{1,5,3,†} Jusak Tandean,^{1,3,‡} and Xing-Bo Yuan^{3,§}

¹*Department of Physics, National Taiwan University, Taipei 10617, Taiwan*

²*Institute of Physics, Academia Sinica, Taipei 11529, Taiwan*

³*Physics Division, National Center for Theoretical Sciences, Hsinchu 30013, Taiwan*

⁴*Kavli IPMU, University of Tokyo, Kashiwa, 277-8583, Japan*

⁵*INPAC, Department of Physics and Astronomy,
Shanghai Jiao Tong University, Shanghai 200240, China*

(Dated: July 22, 2022)

Abstract

A recent LHCb measurement of the ratio R_{K^*} of $B \rightarrow K^*\mu\bar{\mu}$ to $B \rightarrow K^*e\bar{e}$ branching fractions has shown evidence of deviation from the standard model (SM) expectation. This adds to the known anomalies also induced by the $b \rightarrow s\ell\bar{\ell}$ transitions, resulting in a confidence level now as high as 4σ . We analyze whether the parameter space preferred by all the $b \rightarrow s\ell\bar{\ell}$ anomalies is compatible with a heavy Z' boson assumed to have nonuniversal couplings to SM fermions dictated by the principle of minimal flavor violation (MFV). We deal with the MFV couplings of the Z' to leptons in the context of the type-I seesaw scenario for generating neutrino masses. The flavor-violating Z' interactions are subject to stringent constraints from other processes, especially B - \bar{B} mixing, charged lepton decays $\ell_i \rightarrow \ell_j\ell_k\ell_l$ occurring at tree level, and the loop induced $\mu \rightarrow e\gamma$. We perform scans for parameter regions allowed by various data and predict the ranges for a number of observables. Some of the predictions, such as the branching fractions of lepton-flavor violating $\tau \rightarrow 3\mu$, $B \rightarrow Ke\mu$, $K_L \rightarrow e\mu$, and $Z \rightarrow \ell\ell'$, are not far below their experimental bounds and therefore could be probed by searches in the near future. The viable parameter space depends strongly on the neutrino mass hierarchy, with a preference for the inverted one.

*e-mail: chengwei@phys.ntu.edu.tw

†e-mail: hexg@phys.ntu.edu.tw

‡e-mail: jtandean@yahoo.com

§e-mail: xbyuan@cts.nthu.edu.tw

I. INTRODUCTION

In addition to direct searches for new physics (NP) at the energy frontier, the CERN LHC has been testing the standard model (SM) of particle physics through studies of flavor physics. While up to date there is still no strong evidence of nonstandard particles or interactions predicted by various NP models, LHC experiments have, however, turned up quite a few anomalous results in the lower energy regime. In particular, a pattern of discrepancies from SM expectations has recently been emerging from observables in a number of $b \rightarrow s\ell^+\ell^-$ transitions, mostly at around or above the 3σ level. Such coherent deviations call for special attention, as many of them are sensitive to contributions from new particles and/or new interactions.

The aforementioned indications of anomalous $b \rightarrow s\ell^+\ell^-$ interaction showed up in the binned angular distribution of the $B \rightarrow K^*\mu^+\mu^-$ decay, first found by the LHCb Collaboration [1, 2] and later on confirmed by the Belle Collaboration [3, 4], and in the branching fraction deficit of the $B_s \rightarrow \phi\mu^+\mu^-$ decay [5, 6]. Another set of observables that have manifested unexpected values are

$$\begin{aligned} R_K &\equiv \frac{\mathcal{B}(B \rightarrow K\mu^+\mu^-)}{\mathcal{B}(B \rightarrow Ke^+e^-)} , \\ R_{K^*} &\equiv \frac{\mathcal{B}(B \rightarrow K^*\mu^+\mu^-)}{\mathcal{B}(B \rightarrow K^*e^+e^-)} , \end{aligned} \tag{1}$$

which are of great interest because a lot of hadronic uncertainties are canceled out in the ratios and they provide a test of lepton-flavor universality (LFU). These ratios are both predicted to be very close to unity in the SM. However, the former was determined by LHCb to have a value of $R_K = 0.745_{-0.074}^{+0.090}(\text{stat}) \pm 0.036(\text{syst})$ [7]. Very recently, LHCb also reported a measurement on R_{K^*} in two dilepton invariant squared mass (q^2) bins [8]:

$$R_{K^*} = \begin{cases} 0.66_{-0.07}^{+0.11}(\text{stat}) \pm 0.03(\text{syst}) & \text{for } 0.045 < q^2 < 1.1 \text{ GeV}^2 , \\ 0.69_{-0.07}^{+0.11}(\text{stat}) \pm 0.05(\text{syst}) & \text{for } 1.1 < q^2 < 6 \text{ GeV}^2 , \end{cases} \tag{2}$$

in disagreement with SM expectations at the 2.4σ and 2.5σ levels, respectively. The data on R_K and R_{K^*} together reveal consistent breaking of LFU at an even higher confidence level (CL) of about 4σ [9]. This has added to the tantalizing tentative hints of the presence of NP in these processes which has the feature of violating LFU. Thus, unsurprisingly the new R_{K^*} anomaly has stimulated a new wave of theoretical studies about lepton-flavor-nonuniversal $b \rightarrow s\ell^+\ell^-$ interactions [9–35].

In this paper, we entertain the possibility that these anomalies arise from LFU-violating NP and explore some of its implications. When addressing flavor physics beyond the SM, the usual problem one faces is that there are too many model-dependent parameters. On one hand, this provides an opportunity of having rich phenomenology in the flavor sector. On the other hand, the sizable number of parameters tends to complicate the analysis, in some cases making the situation arbitrary. If there is a way to treat the flavor structure systematically, it may simplify the analysis and provide some guide for theoretically understanding the potential NP. One of the efficient means to this end is the framework of so-called minimal flavor violation (MFV), which

we will adopt. The application of the MFV idea to an effective field theory approach at low energies offers a natural model-independent solution for TeV-scale NP to evade flavor-changing neutral current (FCNC) restrictions by demanding that all flavor and CP violations have their origin in the known structure of Yukawa couplings [36, 37]. Although initially motivated by the successful SM description of quark FCNCs, the notion of MFV can be extended to the lepton sector [38]. However, as the SM strictly does not accommodate lepton-flavor violation and it remains unknown whether neutrinos are Dirac or Majorana particles, there is currently no unique way to implement MFV in the lepton sector, and to do so will usually involve picking a particular scenario for endowing neutrinos with mass.

Our interest here is in studying within the MFV framework whether the parameter space preferred by all the $b \rightarrow s\ell^+\ell^-$ anomalies have any conflict with other related observables. After revisiting the case of the relevant dimension-six operators satisfying the MFV criterion in both their quark and lepton parts, we will focus on a scenario in which the flavor violations are induced by an electrically neutral vector particle, such as a Z' boson, which has effective fermionic interactions consistent with the MFV principle. We will look at a variety of constraints on its couplings to quarks and leptons and subsequently evaluate a number of predictions from the allowed parameter space associated with this particle.

Recent global analyses [9–12] have demonstrated that the dimension-6 operators that can produce some of the best fits to the anomalous $b \rightarrow s\ell^+\ell^-$ findings are given by

$$\begin{aligned} \mathcal{L}_{\text{eff}} &\supset \sqrt{8} G_F V_{ts}^* V_{tb} (C_9^\ell O_9^\ell + C_{10}^\ell O_{10}^\ell) + \text{H.c.}, \\ O_9^\ell &= \frac{\alpha_e}{4\pi} \bar{s} \gamma^\eta P_L b \bar{\ell} \gamma_\eta \ell, \quad O_{10}^\ell = \frac{\alpha_e}{4\pi} \bar{s} \gamma^\eta P_L b \bar{\ell} \gamma_\eta \gamma_5 \ell, \end{aligned} \quad (3)$$

with $C_i^\ell = C_i^{\text{SM}} + C_i^{\ell, \text{NP}}$ ($i = 9, 10$) being Wilson coefficients and the NP entering mainly the $\ell = \mu$ terms. In these formulas, α_e and G_F represent the usual fine structure and Fermi constants, $V_{ts, tb}$ are elements of V_{CKM} , the Cabibbo-Kobayashi-Maskawa mixing matrix, $P_L = (1 - \gamma_5)/2$, and at the m_b scale $C_9^{\text{SM}} \simeq -C_{10}^{\text{SM}} \simeq 4.2$ universally for all charged leptons.

As will be seen below, the dimension-6 operators with MFV considered in this work generate interactions that are chiral and feature the relation $C_9^{\ell, \text{NP}} = -C_{10}^{\ell, \text{NP}}$. With the NP effect on the electron channel taken to be vanishing, the 1σ allowed range of $C_9^{\mu, \text{NP}}$ has been found to be $[-0.81, -0.48]$ in this scenario [11]. Assuming that the new interactions in the MFV framework are mediated by a putative Z' gauge boson, we will examine whether the implied parameter space is consistent with existing data on processes such as the B - \bar{B} mixing, neutrino oscillations, and lepton-flavor violating (LFV) processes.

The paper is arranged as follows. Section II briefly reviews the idea of MFV and explains what type of dimension-6 operators with MFV are compatible with the $b \rightarrow s\ell^+\ell^-$ anomalies. In Section III, we introduce a Z' gauge boson that can effectively induce the desired flavor-changing interactions. Subsequently, we discuss how they can account for the $b \rightarrow s\ell^+\ell^-$ anomalies and must respect various constraints, especially from measurements of B - \bar{B} and neutrino oscillations and bounds on LFV processes. In Section IV, we scan the parameter space subject to these requirements and illustrate the viable regions. Among the restraints, we find that the $\mu \rightarrow e\gamma$ and $\tau \rightarrow 3\mu$ may play the most constraining role, depending on the

ordering of light neutrinos' masses. Section V is dedicated to our predictions for a number of processes based upon our parameter scan results. Section VI summarizes our findings.

II. OPERATORS WITH MINIMAL FLAVOR VIOLATION

Since the quark masses and mixing angles are now well-determined, the application of MFV in the quark sector is straightforward. In contrast, there is no unique way to formulate leptonic MFV because our knowledge about the nature and absolute scale of neutrino masses is far from complete. Given that flavor mixing among neutrinos has been empirically established [39], it is attractive to implement leptonic MFV by integrating new ingredients that can account for this fact [38]. One could consider a minimal field content where only the SM fermionic doublets and singlets transform nontrivially under the flavor group, with lepton number violation and neutrino masses being ascribed to the dimension-five Weinberg operator [38]. Less minimally, one could explicitly introduce right-handed neutrinos [38], or alternatively right-handed weak-SU(2)-triplet fermions [40], which transform nontrivially under an expanded flavor group and are responsible for the seesaw mechanism giving Majorana masses to light neutrinos [41, 42]. One could also introduce instead a weak-SU(2)-triplet of unflavored scalars [40, 43] which take part in the seesaw mechanism [44].¹ Here we apply MFV to leptons by invoking the type-I seesaw scenario involving three heavy right-handed neutrinos.

The renormalizable Lagrangian for the masses of SM fermions plus the right-handed neutrinos, denoted by $N_{1,2,3}$, can be expressed as

$$\begin{aligned} \mathcal{L}_m = & -(Y_u)_{jk} \overline{Q}_j P_R U_k \tilde{H} - (Y_d)_{jk} \overline{Q}_j P_R D_k H - (Y_e)_{jk} \overline{L}_j P_R E_k H \\ & - (Y_\nu)_{jk} \overline{L}_j P_R N_k \tilde{H} - \frac{1}{2} (M_N)_{jk} \overline{(N_j)^c} P_R N_k + \text{H.c.}, \end{aligned} \quad (4)$$

where summation over the generation indices $j, k = 1, 2, 3$ is implicit, $Y_{u,d,e,\nu}$ are Yukawa coupling matrices, the quark, lepton, and Higgs doublets are given by

$$Q_k = \begin{pmatrix} U_k \\ D_k \end{pmatrix}, \quad L_k = \begin{pmatrix} \nu_k \\ E_k \end{pmatrix}, \quad H = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}(h + v) \end{pmatrix}, \quad \tilde{H} = i\tau_2 H^* \quad (5)$$

after electroweak symmetry breaking, with $v \simeq 246$ GeV being the vacuum expectation value of H and τ_2 the second Pauli matrix, M_N is the Majorana mass matrix for $N_{1,2,3}$ which without loss of generality can be chosen to be diagonal, the superscript in $(N_j)^c$ refers to charge conjugation, and $P_R = \frac{1}{2}(1 + \gamma_5)$. Hereafter, we entertain the possibility that $N_{1,2,3}$ are degenerate, and so $M_N = \mathcal{M} \text{diag}(1, 1, 1)$. It is then realized that \mathcal{L}_m is formally invariant under the global flavor rotations $Q \rightarrow V_Q Q$, $P_R U \rightarrow P_R V_U U$, $P_R D \rightarrow P_R V_D D$, $L \rightarrow V_L L$, $P_R E \rightarrow P_R V_E E$, and $N = (N_1 \ N_2 \ N_3)^T \rightarrow \mathcal{O}_N N$, with $V_{Q,U,D,L,E} \in \text{SU}(3)_{Q,U,D,L,E}$ and \mathcal{O}_N being a real orthogonal matrix, provided that the Yukawa couplings behave like spurions transforming as $Y_u \rightarrow V_Q Y_u V_U^\dagger$, $Y_d \rightarrow V_Q Y_d V_D^\dagger$, $Y_e \rightarrow V_L Y_e V_E^\dagger$, and $Y_\nu \rightarrow V_L Y_\nu \mathcal{O}_N^T$.

¹ Some other possibilities of leptonic MFV have been discussed in the literature [45–47].

The right-handed neutrinos' mass, \mathcal{M} , is assumed to be very large compared to the elements of $vY_\nu/\sqrt{2}$, triggering the type-I seesaw mechanism [41] which brings about the light-neutrinos' mass matrix $m_\nu = -(v^2/2)Y_\nu M_N^{-1}Y_\nu^\text{T} = U_{\text{PMNS}} \hat{m}_\nu U_{\text{PMNS}}^\text{T}$, where U_{PMNS} is the Pontecorvo-Maki-Nakagawa-Sakata [48] mixing matrix and $\hat{m}_\nu = \text{diag}(m_1, m_2, m_3)$ contains the light neutrinos' eigenmasses, $m_{1,2,3}$. This suggests adopting the interesting form [49]

$$Y_\nu = \frac{i\sqrt{2}}{v} U_{\text{PMNS}} \hat{m}_\nu^{1/2} O M_N^{1/2}, \quad (6)$$

where O is a generally complex orthogonal matrix satisfying $OO^\text{T} = \mathbb{1} \equiv \text{diag}(1, 1, 1)$.

The MFV framework presupposes that the Yukawa couplings are the only sources of flavor violation [36, 37]. Accordingly, to construct effective Lagrangians beyond the SM with MFV built-in, one inserts products of the Yukawa matrices among the pertinent fields to devise operators that are singlet under the SM gauge group and invariant under the flavor rotations described above [37]. Of potential interest here are the combinations

$$\mathbf{A}_q = Y_u Y_u^\dagger, \quad \mathbf{B}_q = Y_d Y_d^\dagger, \quad \mathbf{A}_\ell = Y_\nu Y_\nu^\dagger, \quad \mathbf{B}_\ell = Y_e Y_e^\dagger. \quad (7)$$

Given that the maximum eigenvalues of \mathbf{A}_q and \mathbf{B}_q are, respectively, $y_t^2 = 2m_t^2/v^2 \simeq 0.99$ and $y_b^2 = 2m_b^2/v^2 \simeq 3.0 \times 10^{-4}$ at the mass scale $\mu = m_Z$, for our purposes we can form objects containing up to two powers of \mathbf{A}_q and drop contributions with \mathbf{B}_q , as higher powers of \mathbf{A}_q can be linked to lower ones by means of the Cayley-Hamilton identity [50]. As for \mathbf{A}_ℓ , we select \mathcal{M} to be sufficiently large to make the maximum eigenvalue of \mathbf{A}_ℓ equal unity. Thus, as in the quark sector, we will keep terms up to order \mathbf{A}_ℓ^2 and ignore those with \mathbf{B}_ℓ , whose elements are at most $y_\tau^2 = 2m_\tau^2/v^2 \simeq 1.0 \times 10^{-4}$. It follows that the relevant spurion building blocks are

$$\Delta_q = \zeta_0 \mathbb{1} + \zeta_1 \mathbf{A}_q + \zeta_2 \mathbf{A}_q^2, \quad \Delta_\ell = \xi_0 \mathbb{1} + \xi_1 \mathbf{A}_\ell + \xi_2 \mathbf{A}_\ell^2, \quad (8)$$

where, model-independently, $\zeta_{0,1,2}$ and $\xi_{0,1,2}$ are free parameters expected to be at most of $\mathcal{O}(1)$ and with negligible imaginary components [50], so that we can make the approximations $\Delta_q^\dagger = \Delta_q$ and $\Delta_\ell^\dagger = \Delta_\ell$.

It is convenient to work in the basis where $Y_{d,e}$ are diagonal,

$$Y_d = \text{diag}(y_d, y_s, y_b), \quad Y_e = \text{diag}(y_e, y_\mu, y_\tau), \quad (9)$$

with $y_f = \sqrt{2}m_f/v$, and U_k , D_k , $\tilde{\nu}_{k,L}$, $N_{k,R}$, and E_k refer to the mass eigenstates. In that case,

$$Q_j = \begin{pmatrix} \sum_k (V_{\text{CKM}}^\dagger)_{jk} U_k \\ D_j \end{pmatrix}, \quad L_j = \begin{pmatrix} \sum_k (U_{\text{PMNS}})_{jk} \tilde{\nu}_k \\ E_j \end{pmatrix}, \quad Y_u = V_{\text{CKM}}^\dagger \text{diag}(y_u, y_c, y_t),$$

$$\mathbf{A}_q = V_{\text{CKM}}^\dagger \text{diag}(y_u^2, y_c^2, y_t^2) V_{\text{CKM}}, \quad \mathbf{A}_\ell = \frac{2\mathcal{M}}{v^2} U_{\text{PMNS}} \hat{m}_\nu^{1/2} O O^\dagger \hat{m}_\nu^{1/2} U_{\text{PMNS}}^\dagger. \quad (10)$$

From this point on, we write $\ell_k = E_k$, and so $(\ell_1, \ell_2, \ell_3) = (e, \mu, \tau)$.

Without introducing other new interactions or particles, one then sees that the operators of lowest dimension that are flavor invariant, SM gauge singlet, and of the type that can readily give rise to the NP terms in Eq. (3) are [51]

$$O_1^6 = \overline{Q} \gamma_\eta \Delta_q P_L Q \overline{L} \gamma^\eta \Delta_\ell P_L L, \quad O_2^6 = \overline{Q} \gamma_\eta \tilde{\Delta}_q P_L \tau_a Q \overline{L} \gamma^\eta \tilde{\Delta}_\ell P_L \tau_a L, \quad (11)$$

where $\tilde{\Delta}_q$ and $\tilde{\Delta}_\ell$ are, respectively, of the same form as Δ_q and Δ_ℓ in Eq. (8), but have their own independent coefficients $\tilde{\zeta}_r$ and $\tilde{\xi}_r$, and the index $a = 1, 2, 3$ of the Pauli matrix τ_a is implicitly summed over. The MFV effective Lagrangian of interest is then

$$\mathcal{L}_{\text{eff}}^{\text{MFV}} = \frac{1}{\Lambda^2} (O_1^6 + O_2^6), \quad (12)$$

where the mass scale Λ characterizes the heavy NP underlying these interactions.

From Eq. (12), one could obtain interactions that can account for the $b \rightarrow s\ell^+\ell^-$ anomalies and investigate some of the implications [51] without explicitly addressing the underlying NP. Specifically, among $b \rightarrow (s, d)\ell\bar{\ell}'$ and $s \rightarrow d\ell\bar{\ell}'$ decays with $\ell \neq \ell'$ as well as related processes with neutrinos in the final states, there could be predicted rates which are not far from their experimental results and, therefore, may be testable in near future searches [51].

In the rest of this paper, we concentrate instead on a scenario in which a Z' gauge boson with nonuniversal couplings to SM fermions is responsible for the NP effects on $b \rightarrow s\ell^+\ell^-$. Such a particle exists in myriad models [52]. Since O_2^6 contains charged-currents, only O_1^6 is attributable to the Z' contribution at tree level. It is worth remarking that, although this analysis concerns the Z' gauge boson, the main results are applicable to any new electrically neutral spin-1 particle, which could be composite, having similar flavor-violating couplings.

III. Z' -MEDIATED INTERACTIONS

The renormalizable Lagrangian for the interactions between SM fermions and the Z' boson fulfilling the MFV criterion can take the form

$$\mathcal{L}_{Z'} = -(\bar{Q}\gamma^\eta\Delta_q P_L Q + \bar{L}\gamma^\eta\Delta_\ell P_L L)Z'_\eta, \quad (13)$$

where any overall coupling constant of the Z' has been absorbed into the coefficients $\zeta_{0,1,2}$ and $\xi_{0,1,2}$ in Δ_q and Δ_ℓ , respectively. We also suppose that any mixing between the Z' and SM gauge bosons is negligible and that the Z' mass, $m_{Z'}$, is above the electroweak scale.

From Eq. (13), one can readily derive the MFV Lagrangian, \mathcal{L}_{MFV} , that involves three types of effective four-fermion operators with dimension up to 6. Thus, besides O_1^6 , the additional operators that can appear due to Z' exchange at tree level are given by

$$\begin{aligned} \mathcal{L}_{\text{MFV}} &= \frac{-1}{m_{Z'}^2} (\mathcal{O}^{4q} + \mathcal{O}^{4\ell} + \mathcal{O}^{2q2\ell}), \\ \mathcal{O}^{4q} &= \frac{1}{2} \bar{Q}\gamma_\eta\Delta_q P_L Q \bar{Q}\gamma^\eta\Delta_q P_L Q, \\ \mathcal{O}^{4\ell} &= \frac{1}{2} \bar{L}\gamma_\eta\Delta_\ell P_L L \bar{L}\gamma^\eta\Delta_\ell P_L L, \\ \mathcal{O}^{2q2\ell} &= O_1^6 = \bar{Q}\gamma_\eta\Delta_q P_L Q \bar{L}\gamma^\eta\Delta_\ell P_L L, \end{aligned} \quad (14)$$

where $m_{Z'}$ is taken to be large compared to the energies of the external fermions. With the extra operators to consider, we will need to deal with more constraints than in a model-independent analysis based on the $\bar{Q}Q\bar{L}L$ operators in Eq. (12) alone.

In the following, we discuss the effects of \mathcal{O}^{4q} , \mathcal{O}^{4l} , and \mathcal{O}^{2q2l} in turn and study the restrictions on the elements of $\Delta_{q,\ell}$ from existing data. In view of the recent great interest in the $b \rightarrow s\ell^+\ell^-$ anomalies, we start with a discussion on the interactions involving $\mathcal{O}^{2q2\ell}$.

A. Diquark-Dilepton Interactions

In the presence of $\mathcal{O}^{2q2\ell}$ in \mathcal{L}_{MFV} , the effective interaction responsible for $b \rightarrow s\ell\bar{\ell}'$ is

$$\mathcal{L}_{\text{eff}} \supset \frac{\sqrt{2}\alpha_e\lambda_{sb}G_F}{\pi} C_{\ell\ell'} \bar{s}\gamma^\eta P_L b \bar{\ell}\gamma_\eta P_L \ell', \quad (15)$$

where

$$\lambda_{q'q} = V_{tq'}^* V_{tq}, \quad C_{\ell\ell'} = \delta_{\ell\ell'} C_9^{\text{SM}} + c_{\ell\ell'}, \quad (16)$$

with the approximation $C_{10}^{\text{SM}} = -C_9^{\text{SM}}$. Hence, in terms of the elements of $\Delta_{q,\ell}$

$$\begin{aligned} c_{\ell_j \ell_k} &= \frac{-\pi}{\sqrt{2}\alpha_e G_F m_{Z'}^2} (\zeta_1 y_t^2 + \zeta_2 y_t^4) (\Delta_\ell)_{jk} \\ &\simeq -25.3 \text{ TeV}^2 \frac{(\zeta_1 y_t^2 + \zeta_2 y_t^4) (\Delta_\ell)_{jk}}{m_{Z'}^2}, \end{aligned} \quad (17)$$

where $\alpha_e = 1/133$ has been used and terms with $y_{u,c}$ are dropped. It follows that $|C_{\ell\ell'}| = |C_{\ell'\ell}|$. Analogously, one can write down the corresponding expressions for $b \rightarrow d\ell\bar{\ell}'$ and $s \rightarrow d\ell\bar{\ell}'$.

Subsequent to the recent LHCb finding on R_{K^*} , it has been pointed out that one of the best fits to the $b \rightarrow s\ell^+\ell^-$ data has the NP Wilson coefficients [11]

$$c_{ee} = 0, \quad -1.00 \leq c_{\mu\mu} \leq -0.32 \quad (18)$$

at the 2σ level, which can be interpreted to imply that the Z' boson does not couple to electrons. This is the scenario that we will continue to analyze in this work. Since $c_{ee} \propto (\Delta_\ell)_{11}$, we then have from Eq. (18) the condition $(\Delta_\ell)_{11} = 0$.

The same operator, $\mathcal{O}^{2q2\ell}$, contributes at tree level to $\mu \rightarrow e$ conversion in nuclei which is subject to stringent empirical limits. Nevertheless, as outlined in Appendix A, the $\mathcal{O}^{2q2\ell}$ contribution to this process can be made consistent with its current data by sufficiently reducing the size of the coefficient ζ_0 in Δ_q .

There may also be constraints from collider data. However, given that $(\Delta_\ell)_{11} = 0$, limits implied by LEP measurements on $e^+e^- \rightarrow q\bar{q}$ [53] can be evaded. Moreover, our numerical calculations show that potential restraints from recent LHC results on $pp \rightarrow \mu^+\mu^-$ [54] are not yet realized, as sketched in Appendix A.

B. Four-Quark Interactions

The operator \mathcal{O}^{4q} in \mathcal{L}_{MFV} contributes at tree level to the heavy-light mass difference of neutral B_d (B_s) mesons, $\Delta M_{d(s)}$. Including the SM contribution, we express it as [55]

$$\Delta M_{d(s)} = \Delta M_{d(s)}^{\text{SM}} \left| 1 + \frac{S_d^{Z'}}{S_0(x_t)} \right|, \quad (19)$$

where $S_0(x_t) = 2.35$ for $m_t = 165 \text{ GeV}$ is due to SM loop diagrams and the Z' part is

$$S_d^{Z'} = \frac{4(\Delta_q)_{13(23)}^2 \tilde{r}}{\lambda_{db(sb)}^2 g_{\text{SM}}^2 m_{Z'}^2} = \frac{4(\zeta_1 y_t^2 + \zeta_2 y_t^4)^2 \tilde{r}}{g_{\text{SM}}^2 m_{Z'}^2}, \quad (20)$$

with [55] $g_{\text{SM}}^2 = 1.78 \times 10^{-7} \text{ GeV}^{-2}$ and the QCD factor $\tilde{r} \sim 1$ for $m_{Z'} \sim 1 \text{ TeV}$.

The experimental and SM values of $\Delta M_{d,s}$ are, in units of ps^{-1} ,

$$\begin{aligned} \Delta M_d^{\text{exp}} &= 0.5064 \pm 0.0019 \quad [56], & \Delta M_d^{\text{SM}} &= 0.575_{-0.090}^{+0.093} \quad [57], \\ \Delta M_s^{\text{exp}} &= 17.757 \pm 0.021 \quad [56], & \Delta M_s^{\text{SM}} &= 18.6_{-2.3}^{+2.4} \quad [57], \end{aligned} \quad (21)$$

with updated parameters used in the SM predictions. From these numbers, we can calculate the 2σ ranges

$$0.60 \leq \frac{\Delta M_d^{\text{exp}}}{\Delta M_d^{\text{SM}}} \leq 1.16, \quad 0.71 \leq \frac{\Delta M_s^{\text{exp}}}{\Delta M_s^{\text{SM}}} \leq 1.19 \quad (22)$$

after combining in quadrature the relative errors in the measurements and predictions. The first, and somewhat stronger, of these two constraints then translates into

$$0 \leq \frac{S_d^{Z'}}{S_0(x_t)} = 9.56 \times 10^6 \text{ GeV}^2 \frac{(\zeta_1 y_t^2 + \zeta_2 y_t^4)^2 \tilde{r}}{m_{Z'}^2} \leq 0.16 \quad (23)$$

or, with $\tilde{r} = 1$,

$$\frac{|\zeta_1 y_t^2 + \zeta_2 y_t^4|}{m_{Z'}} \leq \frac{0.13}{\text{TeV}}. \quad (24)$$

This caps the quark part of $\mathbf{c}_{\ell_j \ell_k}$ in Eq. (17).

It is worth noting that the neutral-kaon system can furnish a comparable, but weaker, restraint, as \mathcal{O}^{4q} can modify the SM predictions for the K_L - K_S mass difference ΔM_K and CP -violation parameter ϵ_K . The Z' contribution $M_{12}^{K,Z'} = \lambda_{ds}^2 (\zeta_1 y_t^2 + \zeta_2 y_t^4)^2 \eta_2 \hat{B}_K f_K^2 m_{K^0} \tilde{r} / (6m_{Z'}^2)$, with [55] $\eta_2 = 0.5765 \pm 0.0065$, $\hat{B}_K = 0.767 \pm 0.010$, and $f_K = (156.1 \pm 1.1) \text{ MeV}$, enters via $\Delta M_K = 2 \text{Re}(M_{12}^{K,\text{SM}} + M_{12}^{K,Z'}) + \Delta M_K^{\text{LD}}$ and $\epsilon_K = \text{Im}(M_{12}^{K,\text{SM}} + M_{12}^{K,Z'}) / (\sqrt{2} \Delta M_K^{\text{exp}})$, where ΔM_K^{LD} encodes long-distance effects and $\Delta M_K^{\text{exp}} = (52.89 \pm 0.10) \times 10^{10} / \text{s}$ [39]. Given the potential sizable uncertainties in the ΔM_K calculation [55], we focus on ϵ_K , whose measured and SM values are $|\epsilon_K^{\text{exp}}| = (2.228 \pm 0.011) \times 10^{-3}$ [63] and $|\epsilon_K^{\text{SM}}| = (2.27_{-0.42}^{+0.21}) \times 10^{-3}$ [39]. The 2σ ranges of these numbers then suggest that we can impose $|M_{12}^{K,Z'}| < 5\sqrt{2} \times 10^{-4} \Delta M_K^{\text{exp}}$, which implies $|\zeta_1 y_t^2 + \zeta_2 y_t^4| / m_{Z'} < 0.17 / \text{TeV}$.

C. Four-Lepton Interactions

The $\mathcal{O}^{4\ell}$ operator in \mathcal{L}_{MFV} , induced by the Z' boson at tree level, give rise to various processes that conserve or violate lepton flavors at tree level or 1-loop level. As searches for the flavor-violating decays of charged leptons have yielded the most stringent bounds on some of the interactions of interest, we treat these processes first.

For $\ell_1 \rightarrow \ell_2 \ell_3 \bar{\ell}_4$ and $\ell_1 \rightarrow \ell_2 \gamma$, we employ the relevant formulas from Ref. [58]. Thus, we arrive at the rates

$$\begin{aligned}\Gamma_{\tau \rightarrow ee\bar{\mu}} &= \frac{|(\Delta_\ell)_{12}(\Delta_\ell)_{13}|^2 m_\tau^5}{768 \pi^3 m_{Z'}^4}, & \Gamma_{\tau \rightarrow \mu\mu\bar{e}} &= \frac{|(\Delta_\ell)_{21}(\Delta_\ell)_{23}|^2 m_\tau^5}{768 \pi^3 m_{Z'}^4}, \\ \Gamma_{\tau \rightarrow \mu e\bar{e}} &= \frac{|(\Delta_\ell)_{21}(\Delta_\ell)_{13}|^2 m_\tau^5}{1536 \pi^3 m_{Z'}^4}, & \Gamma_{\tau \rightarrow 3\mu} &= \frac{|(\Delta_\ell)_{22}(\Delta_\ell)_{23}|^2 m_\tau^5}{768 \pi^3 m_{Z'}^4}, \\ \Gamma_{\tau \rightarrow e\mu\bar{\mu}} &= \frac{|(\Delta_\ell)_{22}(\Delta_\ell)_{13} + (\Delta_\ell)_{12}(\Delta_\ell)_{23}|^2 m_\tau^5}{1536 \pi^3 m_{Z'}^4}\end{aligned}\quad (25)$$

from tree-level Z' -exchange diagrams and

$$\begin{aligned}\Gamma_{\mu \rightarrow e\gamma} &= \frac{\alpha_e m_\mu^5}{2304 \pi^4 m_{Z'}^4} |(\Delta_\ell)_{12}(\Delta_\ell)_{22} + (\Delta_\ell)_{13}(\Delta_\ell)_{32}|^2, \\ \Gamma_{\tau \rightarrow e\gamma} &= \frac{\alpha_e m_\tau^5}{2304 \pi^4 m_{Z'}^4} |(\Delta_\ell)_{12}(\Delta_\ell)_{23} + (\Delta_\ell)_{13}(\Delta_\ell)_{33}|^2, \\ \Gamma_{\tau \rightarrow \mu\gamma} &= \frac{\alpha_e m_\tau^5}{2304 \pi^4 m_{Z'}^4} |(\Delta_\ell)_{21}(\Delta_\ell)_{13} + (\Delta_\ell)_{22}(\Delta_\ell)_{23} + (\Delta_\ell)_{23}(\Delta_\ell)_{33}|^2\end{aligned}\quad (26)$$

from Z' -loop diagrams, where we have neglected the final leptons' masses and taken into account the choice $(\Delta_\ell)_{11} = 0$, which also leads to $\Gamma_{\mu \rightarrow 3e} = \Gamma_{\tau \rightarrow 3e} = 0$. The experimental data are [39, 59]

$$\begin{aligned}\mathcal{B}(\tau \rightarrow ee\bar{\mu})_{\text{exp}} &< 1.5 \times 10^{-8}, & \mathcal{B}(\tau \rightarrow \mu\mu\bar{e})_{\text{exp}} &< 1.7 \times 10^{-8}, \\ \mathcal{B}(\tau \rightarrow e\mu\bar{\mu})_{\text{exp}} &< 2.7 \times 10^{-8}, & \mathcal{B}(\tau \rightarrow 3\mu)_{\text{exp}} &< 2.1 \times 10^{-8}, \\ \mathcal{B}(\tau \rightarrow \mu e\bar{e})_{\text{exp}} &< 1.8 \times 10^{-8}, & \mathcal{B}(\mu \rightarrow e\gamma)_{\text{exp}} &< 4.2 \times 10^{-13}, \\ \mathcal{B}(\tau \rightarrow e\gamma)_{\text{exp}} &< 3.3 \times 10^{-8}, & \mathcal{B}(\tau \rightarrow \mu\gamma)_{\text{exp}} &< 4.4 \times 10^{-8},\end{aligned}\quad (27)$$

all at 90% CL. The strictest of the bounds on these decay modes is from $\mathcal{B}(\mu \rightarrow e\gamma)_{\text{exp}}$, which translates into

$$\frac{|(\Delta_\ell)_{12}(\Delta_\ell)_{22} + (\Delta_\ell)_{13}(\Delta_\ell)_{32}|}{m_{Z'}^2} < \frac{5.4 \times 10^{-4}}{\text{TeV}^2}. \quad (28)$$

This indicates that some tuning is needed so that $(\Delta_\ell)_{22} = \mathcal{O}(0.2)$ can be maintained in order to satisfy Eq. (18). The other modes, notably $\tau \rightarrow 3\mu$, can also be important.

Related to $\ell_1 \rightarrow \ell_2 \gamma$ is the Z' contribution to the anomalous magnetic moment of charged lepton ℓ_j ,

$$a_{\ell_j}^{Z'} = \frac{-m_{\ell_j}^2}{12\pi^2 m_{Z'}^2} \sum_k |(\Delta_\ell)_{jk}|^2. \quad (29)$$

With $a_{\ell_j}^{Z'}$ being always negative, due to the purely left-handed Z' couplings, it does not help resolve the discrepancy between a_μ^{SM} and a_μ^{exp} , presently differing by $a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (288 \pm 80) \times 10^{-11}$ [39]. Nevertheless, requiring $|a_{E_j}^{Z'}|$ to be less than the error in this difference does not result in a strict limitation on the Z' couplings.

Another type of low-energy process which can be affected by the Z' is the SM-dominated decay $\ell \rightarrow \ell' \nu \nu'$. Since the neutrinos are unobserved, its rate comes from channels with all possible combinations of neutrino flavors in the final states, namely

$$\Gamma_{\tau \rightarrow \mu \nu \nu'} = \Gamma_{\tau \rightarrow \mu \nu_\mu \nu_\tau} + \Gamma_{\tau \rightarrow \mu \nu_e \nu_\tau} + \Gamma_{\tau \rightarrow \mu \nu_\tau \nu_\tau} + \Gamma_{\tau \rightarrow \mu \nu_e \nu_\mu} + \Gamma_{\tau \rightarrow \mu \nu_\mu \nu_\mu}, \quad (30)$$

where

$$\begin{aligned} \Gamma_{\tau \rightarrow \mu \nu_\mu \nu_\tau} &= \frac{G_F^2 m_\tau^5}{192 \pi^3} (1 - 8\rho_\mu + 8\rho_\mu^3 - \rho_\mu^4 - 12\rho_\mu^2 \ln \rho_\mu) (1 + R_{23})^2, \\ \rho_\ell &= \frac{m_\ell^2}{m_\tau^2}, \quad R_{rs} = \frac{|(\Delta_\ell)_{rs}|^2}{\sqrt{8} G_F m_{Z'}^2}, \end{aligned} \quad (31)$$

and the other partial rates in Eq. (30) can be neglected, without SM contributions and proportional to $|(\Delta_\ell)_{23}^2 (\Delta_\ell)_{rs}^2|$. One could write down an analogous formula for $\Gamma_{\tau \rightarrow e \nu \nu'}$. From the data $\mathcal{B}(\tau \rightarrow e \nu \nu')_{\text{exp}} = (17.82 \pm 0.04)\%$ and $\mathcal{B}(\tau \rightarrow \mu \nu \nu')_{\text{exp}} = (17.39 \pm 0.04)\%$ [39] and the SM predictions $\mathcal{B}(\tau \rightarrow e \nu \nu')_{\text{SM}} = 0.1778 \pm 0.0003$ and $\mathcal{B}(\tau \rightarrow \mu \nu \nu')_{\text{SM}} = 0.1729 \pm 0.0003$ [60], we calculate

$$\frac{\mathcal{B}(\tau \rightarrow e \nu \nu')_{\text{exp}}}{\mathcal{B}(\tau \rightarrow e \nu \nu')_{\text{SM}}} = 1.002 \pm 0.006, \quad \frac{\mathcal{B}(\tau \rightarrow \mu \nu \nu')_{\text{exp}}}{\mathcal{B}(\tau \rightarrow \mu \nu \nu')_{\text{SM}}} = 1.006 \pm 0.006, \quad (32)$$

with 2σ errors. Numerically, we get $(1 + R_{13,23})^2 - 1 < 0.0011$ for the Z' effect represented by our benchmark points, and so it is at least several times smaller than the errors in Eq. (32).

At higher energies, the Z' effects may be probed by LEP experiments on the scattering $e^+ e^- \rightarrow \ell^+ \ell^-$ for $\ell = e, \mu, \tau$. In particular, the lower limits at 95% CL on the effective heavy mass scale derived from fits to their data [53] imply

$$\frac{|(\Delta_\ell)_{11}|}{m_{Z'}} \leq \frac{0.28}{\text{TeV}}, \quad 0 \leq \frac{(\Delta_\ell)_{11}(\Delta_\ell)_{jj} + (\Delta_\ell)_{1j}(\Delta_\ell)_{j1}}{m_{Z'}^2} \leq \frac{0.13}{\text{TeV}^2}, \quad (33)$$

where $j = 2, 3$. The first constraint is automatically satisfied by our preference $(\Delta_\ell)_{11} = 0$, and consequently, since Δ_ℓ is Hermitian, the second one becomes

$$\frac{|(\Delta_\ell)_{1j}|}{m_{Z'}} \leq \frac{0.36}{\text{TeV}}, \quad j = 2, 3. \quad (34)$$

As can be expected, these restrictions turn out to be less important than that in Eq. (28). Although not explicitly addressed in this study, we mention that at 1-loop level the leptonic Z' contributions to Z -pole observables, also measured at LEP [39], are not strongly constrained either, provided that $m_{Z'} > 0.5 \text{ TeV}$.

IV. NUMERICAL ANALYSIS

As discussed in the preceding sections, we deal with the fermionic interactions of the Z' by imposing MFV on both its quark and lepton couplings and, for the latter, by incorporating the type-I seesaw mechanism with 3 heavy right-handed neutrinos. One could perform instead a simpler implementation of leptonic MFV by assuming a minimal field content with only SM fermions plus the dimension-5 Weinberg operator, as was done in Ref. [61]. However, in the type-I seesaw case, there is significantly more freedom to satisfy the various constraints.

Given that the Z' leptonic interactions in Eq. (13) involve $\Delta_\ell = \xi_0 \mathbb{1} + \xi_1 \mathbf{A}_\ell + \xi_2 \mathbf{A}_\ell^2$ with \mathbf{A}_ℓ defined in Eq. (10), to evaluate them we need the values of the elements of U_{PMNS} as well as the coefficients $\xi_{0,1,2}$. Thus, adopting the standard U_{PMNS} parametrization [39], we employ the parameter values quoted in Table I from a recent fit to global neutrino data [62]. The majority of these numbers depend on whether the light neutrinos' masses have a normal ordering (NO), where $m_1 < m_2 < m_3$, or an inverted one (IO), where $m_3 < m_1 < m_2$. As the absolute scale of $m_{1,2,3}$ is not yet established, for definiteness we will pick $m_{1(3)} = 0$ in the NO (IO) case. Although U_{PMNS} may generally also contain Majorana phases, which are still unknown, for simplicity we set them to zero. As for $\xi_{0,1,2}$, one of them is no longer free due to the requisite $(\Delta_\ell)_{11} = 0$ implied by Eq. (18), and so we fix $\xi_0 = -\xi_1 (\mathbf{A}_\ell)_{11} - \xi_2 (\mathbf{A}_\ell^2)_{11}$, but permit the other two to have any real values as long as $|\xi_{1,2}| \leq \mathcal{O}(1)$.

We begin our numerical analysis by looking first at the simplest possibility, which is that the orthogonal matrix O in \mathbf{A}_ℓ is real and consequently $\mathbf{A}_\ell = 2\mathcal{M}U_{\text{PMNS}}\hat{m}_\nu U_{\text{PMNS}}^\dagger/v^2$. In our numerical explorations for this scenario, after fixing $(\zeta_1 y_t^2 + \zeta_2 y_t^4)/m_{Z'} = 0.13/\text{TeV}$ in $\mathbf{c}_{\mu\mu}$ according to Eqs. (17) and (24), we scan the quantities listed in Table I within their 2σ intervals and let $\xi_{1,2}/m_{Z'}$ have any real values between $\pm 1.5/\text{TeV}$. In the NO case, we find that we can attain $-0.46 \lesssim \mathbf{c}_{\mu\mu} \leq -0.32$, which is a portion of the $\mathbf{c}_{\mu\mu}$ range in Eq. (18), but on its upper side, as long as the Dirac CP -violation phase δ in U_{PMNS} lies below its central value in Table I by about 1σ or more. Consequently, although it may be too early to rule out this possibility, it is disfavored. The situation in the IO case is worse, as we are not able to reach the desired values of $\mathbf{c}_{\mu\mu}$ during our scans. It is therefore of interest to consider another choice of \mathbf{A}_ℓ which has a less simple structure.

Parameter	NH	IH
$\sin^2\theta_{12}$	0.306 ± 0.012	0.306 ± 0.012
$\sin^2\theta_{23}$	$0.441^{+0.027}_{-0.021}$	$0.587^{+0.020}_{-0.024}$
$\sin^2\theta_{13}$	0.02166 ± 0.00075	0.02179 ± 0.00076
$\delta/^\circ$	261^{+51}_{-59}	277^{+40}_{-46}
$\Delta m_{21}^2 = m_2^2 - m_1^2$	$(7.50^{+0.19}_{-0.17}) \times 10^{-5} \text{ eV}^2$	$(7.50^{+0.19}_{-0.17}) \times 10^{-5} \text{ eV}^2$
$\Delta m_{3\ell}^2$	$m_3^2 - m_1^2 = (2.524^{+0.039}_{-0.040}) \times 10^{-3} \text{ eV}^2$	$m_3^2 - m_2^2 = (-2.514^{+0.038}_{-0.041}) \times 10^{-3} \text{ eV}^2$

TABLE I: The best-fit values, and their one-sigma errors, of neutrino oscillation parameters from the global analysis in Ref. [62].

A more promising situation is where the O matrix is complex, which leads to

$$A_\ell = \frac{2}{v^2} \mathcal{M} U_{\text{PMNS}} \hat{m}_\nu^{1/2} O O^\dagger \hat{m}_\nu^{1/2} U_{\text{PMNS}}^\dagger, \quad (35)$$

Since we can in general write $O = e^{iR} e^{R'}$ with real antisymmetric matrices R and R' , we have

$$O O^\dagger = e^{2iR}, \quad R = \begin{pmatrix} 0 & r_1 & r_2 \\ -r_1 & 0 & r_3 \\ -r_2 & -r_3 & 0 \end{pmatrix}, \quad (36)$$

where $r_{1,2,3}$ are independent real constants. These extra free parameters prove to be helpful for our purposes. When conducting our scans in this scenario, we let the other parameters fall within their ranges specified in the previous paragraph, whereas $r_{1,2,3}$ are allowed to have any real values.²

With O being complex, during our scans we can obtain $c_{\mu\mu}$ values consistent with Eq. (18), and all the neutrino mixing parameters stay within their 2σ regions, including δ which can fall even inside its 1σ range. To illustrate this, in Fig. 1 we present sample distributions of δ versus $c_{\mu\mu}$ in the NO (magenta) and IO (cyan) cases corresponding, respectively, to 2000 and 3000 benchmark points in the parameter space fulfilling the different constraints described earlier. Evidently, it is easier in the IO scenario to achieve a larger size of $c_{\mu\mu}$ while satisfying the various restrictions.

As expected, the limit from $\mu \rightarrow e\gamma$ searches plays a major constraining role for many of the benchmarks, as can be viewed in Figs. 2 and 3, where we plot the branching fractions of




FIG. 1: Samples of the Dirac CP -violation phase δ in U_{PMNS} versus $c_{\mu\mu}$ corresponding to benchmark points within the allowed parameter space in the NO (magenta) and IO (cyan) cases. The magenta dashed (cyan dotted) lines mark the boundaries of the 2σ region of δ in the NO (IO) case.

² In our numerical analysis, we aim mainly at obtaining viable solutions under our MFV framework with the Z' that can account for the $b \rightarrow s\ell^+\ell^-$ anomalies and looking at some of the implications. As our results demonstrate, there are indeed a substantial amount of points in the Z' parameter space of interest which can accomplish our purposes and are simultaneously compatible with the pertinent constraints. Therefore, in this study, as also in [51], we leave aside concerns about the issue of fine tuning which has been raised in [32].




FIG. 2: Top: samples of the branching fractions of $\mu \rightarrow e\gamma$ and $\tau \rightarrow 3\mu, ee\bar{\mu}, e\mu\bar{\mu}$ divided by their respective experimental upper-limits versus $c_{\mu\mu}$ corresponding to points within the allowed parameter space in the NO cases, as described in the text. Bottom: distributions of some of the pairs of ratios in the top plots.




FIG. 3: Same as Fig. 2, but for the IO case.

$\mu \rightarrow e\gamma$ and $\tau \rightarrow 3\mu, ee\bar{\mu}, e\mu\bar{\mu}$ normalized by their respective experimental bounds, which are quoted in Eq. (27), versus $c_{\mu\mu}$. The $\tau \rightarrow 3\mu$ data can also be important, especially in the NO case, in which $c_{\mu\mu} < -0.46$ is not possible without $\mathcal{B}(\tau \rightarrow 3\mu)$ violating its empirical limit, as can be inferred from the middle plot in Fig. 2. In these figures, we do not display the corresponding ratios for $\tau \rightarrow e\gamma, \mu\gamma, \mu\mu\bar{e}, \mu e\bar{e}$ because they are comparatively less able to reach unity.

V. PREDICTIONS

We notice in Figs. 2 and 3 that there are a considerable amount of points that are not far less than unity, implying that the calculated branching fractions are close to their experimental limits, even within factors of a few. Therefore, these theoretical results and the corresponding $c_{\mu\mu}$ values constitute some of the predictions of our Z' model. They are testable with future quests or detections of these LFV decays and upcoming improved measurements of $b \rightarrow s\ell^+\ell^-$ processes.

In Figs. 2 and 3, we also see that the NO and IO scenarios predict different potential correlations among the branching fractions of these LFV decays which may be confirmed or excluded when they are observed in the future with sufficient precision. To illustrate these possibilities, based on these graphs we present in Figs. 4 and 5 the distributions of several pairs of the ratios $R = \mathcal{B}/\mathcal{B}_{\text{exp}}$ of the calculated branching fractions to their corresponding experimental bounds.

The fact that Eq. (17) also describes LFV couplings implies that they give rise to $b \rightarrow s\ell\bar{\ell}'$ and, analogously, also $b \rightarrow d\ell\bar{\ell}'$ and $s \rightarrow d\ell\bar{\ell}'$, with $\ell \neq \ell'$, all of which strictly do not occur in the SM with massless neutrinos. Accordingly, we have predictions for a number of exclusive $B_{d,s}$ -meson and kaon decays. Using the pertinent formulas given in Ref. [51] with updated CKM parameters [63], we find the maximum $|\mathbf{c}_{\ell\ell'}|$ from our benchmark points to calculate the branching fractions collected in Table II. We observe that a few of the predictions (*e.g.*, the $B \rightarrow K^{(*)}e\mu$, $B \rightarrow \pi e\mu$, and $K_L \rightarrow e\mu$ modes) are within two orders of magnitude from their experimental bounds, especially $K_L \rightarrow e\mu$, and consequently may be probed in near-future searches.

Future measurements of $b \rightarrow s\tau^+\tau^-$ decays, such as $B \rightarrow K^{(*)}\tau^+\tau^-$ and $B_{d,s} \rightarrow \tau^+\tau^-$, which are not yet seen [56], may offer extra tests on the coefficient $\mathbf{c}_{\tau\tau}$. From our benchmark points, we find $-0.63 (-0.85) < \mathbf{c}_{\tau\tau} < +0.80 (-0.11)$ in the NO (IO) of light neutrino masses. This implies that our Z' scenario predicts a modification to the SM expectation by a factor of

$$0.72 (0.64) < \left| 1 + \frac{\mathbf{c}_{\tau\tau}}{C_{9,10}^{\text{SM}}} \right|^2 < 1.42 (0.95). \quad (37)$$




FIG. 4: Sample distributions of correlations between various pairs of the ratios $R = \mathcal{B}/\mathcal{B}_{\text{exp}}$ of the calculated branching fractions to their corresponding experimental bounds in Fig. 2.




FIG. 5: Sample distributions of correlations between various pairs of the ratios $R = \mathcal{B}/\mathcal{B}_{\text{exp}}$ of the calculated branching fractions to their corresponding experimental bounds in Fig. 3.

Hence the Z' contribution may be detectable, and the NO and IO possibilities may be distinguishable, in the acquired data. However, it may be a while before this can be probed because the SM predictions for their branching fractions are of order 10^{-6} or lower [65], whereas experimentally there are still only upper bounds of order 10^{-3} for the different channels [56, 66].

The flavor-violating decays $Z \rightarrow \ell\bar{\ell}'$ also have not yet been observed, but there have been searches for them which resulted in the limits quoted in Table III. These processes can occur in our model because of the flavor-violating Z' -loop modifications to the $Z\ell\bar{\ell}$ vertex and leptonic self-energy diagrams. From the amplitude

$$\mathcal{M}_{Z \rightarrow \ell\bar{\ell}'} = \bar{u}_\ell \not{\epsilon}_Z (L_{\ell\ell'} P_L + R_{\ell\ell'} P_R) v_{\bar{\ell}'}, \quad (38)$$

one obtains the rate

$$\Gamma_{Z \rightarrow \ell\bar{\ell}'} = \frac{|\mathbf{p}_\ell|}{12\pi m_Z^2} \left\{ (|L_{\ell\ell'}|^2 + |R_{\ell\ell'}|^2) \left[m_Z^2 - \frac{m_\ell^2 + m_{\ell'}^2}{2} - \frac{(m_\ell^2 - m_{\ell'}^2)^2}{2m_Z^2} \right] + 6 \text{Re}(L_{\ell\ell'}^* R_{\ell\ell'}) m_{\ell'} m_\ell \right\}, \quad (39)$$

where \mathbf{p}_ℓ is the three-momentum of ℓ in the Z rest-frame. Including the SM and Z' contributions, one has

$$L_{\ell\ell'} = \delta_{\ell\ell'} g_L^{\text{SM}} + L_{\ell\ell'}^{Z'}, \quad R_{\ell\ell'} = \delta_{\ell\ell'} g_R^{\text{SM}}, \quad (40)$$

Decay mode	Branching fractions		
	Measured upper limit at 90% CL [39, 56]	Prediction maximum or range	
		NO	IO
$B \rightarrow Ke^\pm\mu^\mp$	3.8×10^{-8}	2.9×10^{-9}	3.0×10^{-9}
$B \rightarrow K^*e^\pm\mu^\mp$	5.1×10^{-7}	7.8×10^{-9}	7.8×10^{-9}
$B_s \rightarrow e^\pm\mu^\mp$	1.1×10^{-8}	8.6×10^{-12}	9.0×10^{-12}
$B \rightarrow \pi e^\pm\mu^\mp$	9.2×10^{-8}	1.2×10^{-10}	1.3×10^{-10}
$B \rightarrow \rho e^\pm\mu^\mp$	3.2×10^{-6}	3.1×10^{-10}	3.2×10^{-10}
$B^0 \rightarrow e^\pm\mu^\mp$	2.8×10^{-9}	2.6×10^{-13}	2.7×10^{-13}
$B^+ \rightarrow K^+e^\pm\tau^\mp$	3.0×10^{-5}	8.1×10^{-9}	5.9×10^{-9}
$B^+ \rightarrow K^{*+}e^\pm\tau^\mp$	–	1.6×10^{-8}	1.2×10^{-8}
$B_s \rightarrow e^\pm\tau^\mp$	–	8.0×10^{-9}	5.8×10^{-9}
$B^+ \rightarrow \pi^+e^-\tau^+$	2.0×10^{-5}	1.9×10^{-10}	1.4×10^{-10}
$B^+ \rightarrow \rho^+e^\pm\tau^\mp$	–	7.1×10^{-10}	5.2×10^{-10}
$B^0 \rightarrow e^\pm\tau^\mp$	2.8×10^{-5}	2.4×10^{-10}	1.7×10^{-10}
$B^+ \rightarrow K^+\mu^\pm\tau^\mp$	4.8×10^{-5}	$[0.3, 3.1] \times 10^{-9}$	2.6×10^{-9}
$B^+ \rightarrow K^{*+}\mu^\pm\tau^\mp$	4.8×10^{-5}	$[0.7, 6.1] \times 10^{-9}$	5.1×10^{-9}
$B_s \rightarrow \mu^\pm\tau^\mp$	–	$[0.3, 3.1] \times 10^{-9}$	2.6×10^{-9}
$B^+ \rightarrow \pi^+\mu^\pm\tau^\mp$	7.2×10^{-5}	$[0.2, 1.5] \times 10^{-10}$	1.2×10^{-10}
$B^+ \rightarrow \rho^+\mu^\pm\tau^\mp$	7.2×10^{-5}	$[0.3, 2.7] \times 10^{-10}$	2.3×10^{-10}
$B^0 \rightarrow \mu^\pm\tau^\mp$	2.2×10^{-5}	$[1, 9] \times 10^{-11}$	7.7×10^{-11}
$K_L \rightarrow e^\pm\mu^\mp$	4.7×10^{-12}	1.4×10^{-12}	1.5×10^{-12}

TABLE II: The maximum predictions for the branching fractions of exclusive b -meson (kaon) decays involving $e\mu$, $e\tau$, and $\mu\tau$ ($e\mu$) in the final states. The lower end of a prediction is also displayed if exceeding one per mill of its upper end. For comparison, the data are quoted if available. To conform to the experimental reports [64], the $B \rightarrow K^{(*)}e\mu$ prediction is the simple average over the B^+ and B^0 channels, $\mathcal{B}(B \rightarrow K^{(*)}e^\pm\mu^\mp) = (\mathcal{B}(B^+ \rightarrow K^{(*)+}e^\pm\mu^\mp) + \mathcal{B}(B^0 \rightarrow K^{(*)0}e^\pm\mu^\mp))/2$, whereas the $B \rightarrow \pi e\mu$ prediction is from $\mathcal{B}(B \rightarrow \pi e^\pm\mu^\mp) = \mathcal{B}(B^+ \rightarrow \pi^+e^\pm\mu^\mp)/2 + \mathcal{B}(B^0 \rightarrow \pi^0e^\pm\mu^\mp)$ and similarly for $B \rightarrow \rho e^\pm\mu^\mp$. The predictions for $B_s \rightarrow \phi\ell\ell'$ are close to those for $B \rightarrow K^*\ell\ell'$.

where $g_L^{\text{SM}} = g(2s_w^2 - 1)/(2c_w)$ and $g_R^{\text{SM}} = gs_w^2/c_w$ are the SM contributions at tree level, with g being the weak coupling constant, $c_w = \sqrt{1 - s_w^2}$, and s_w^2 the squared sine of the Weinberg angle. For $\ell\ell' = E_k E_l$, the Z' contribution is given by [67, 68]

$$\begin{aligned}
L_{E_k E_l}^{Z'} &= \frac{-\mathcal{F}(\varrho)}{16\pi^2} \sum_o (\Delta_\ell)_{ko} (\Delta_\ell)_{ol}, \quad \varrho = \frac{m_{Z'}^2}{m_Z^2}, \\
\mathcal{F}(\varrho) &= \frac{7}{2} + 2\varrho + 2(1 + \varrho)^2 \text{Li}_2\left(-\frac{1}{\varrho}\right) + (\ln \varrho + i\pi) \left[3 + 2\varrho + 2(1 + \varrho)^2 \ln \frac{\varrho}{1 + \varrho}\right]. \quad (41)
\end{aligned}$$

Numerically, we have checked that for $\ell' = \ell$ the points in the viable Z' parameter space found above produce effects on the Z -pole observables that are well within the 95% CL ranges of their data [39], as long as $m_{Z'} \gtrsim 0.5 \text{ TeV}$. At the same time, for $\ell' \neq \ell$ the Z' contributions to $Z \rightarrow \ell\bar{\ell}'$ may be detectable in the not-too-distant future. In Table III, from the allowed

Decay mode	Branching fractions				
	Measured upper limit at 95% CL [39]	Prediction maximum or range			
		NO		IO	
		0.6 TeV	1 TeV	0.6 TeV	1 TeV
$Z \rightarrow e^\pm \mu^\mp$	7.5×10^{-7}	8.3×10^{-10}	1.8×10^{-11}	8.3×10^{-10}	1.8×10^{-11}
$Z \rightarrow e^\pm \tau^\mp$	9.8×10^{-6}	3.2×10^{-6}	7.0×10^{-8}	4.7×10^{-7}	1.0×10^{-8}
$Z \rightarrow \mu^\pm \tau^\mp$	1.2×10^{-5}	$[0.8, 8.5] \times 10^{-7}$	$[0.2, 1.9] \times 10^{-8}$	8.8×10^{-7}	1.9×10^{-8}

TABLE III: The maximum predictions of the branching fractions of $Z \rightarrow e\mu, e\tau, \mu\tau$ due to loop contributions of the Z' with mass $m_{Z'} = 0.6$ and 1 TeV, compared to the experimental limits. The lower end of a prediction is also displayed if exceeding one per mill of its upper end.

parameter space, we present some predictions of the branching fractions of these LFV decays for $m_{Z'} = 0.5$ and 1 TeV. We see that, although the $e\mu$ mode is likely to be undetectable, the $e\tau$ and $\mu\tau$ predictions can be only a few times below their respective experimental bounds.

VI. CONCLUSIONS

Inspired by the recent hint of lepton flavor nonuniversality in the $B \rightarrow K^* \mu \bar{\mu}$ and $K^* e \bar{e}$ decays, along with several anomalies observed in other $b \rightarrow s \ell \bar{\ell}$ transitions, we have studied within the minimal flavor violation framework whether the parameter space preferred by such data can be consistent with a wider class of observables. Restricting ourselves to new physics operators up to dimension-6, we have shown that the new interactions are chiral and feature a specific relation for the Wilson coefficients in the effective Hamiltonian for $b \rightarrow s \ell \bar{\ell}$ decays: $C_9^{\ell, \text{NP}} = -C_{10}^{\ell, \text{NP}}$. With the hierarchy in quark Yukawa couplings and assuming $\mathcal{O}(1)$ neutrino Yukawa couplings, we have found that only the Δ_q and Δ_ℓ couplings, defined in Eq. (8), can induce flavor-violating interactions.

We have also considered a scenario where the new physics effects on the $b \rightarrow s \ell \bar{\ell}$ decays are mediated by a Z' gauge boson with nonuniversal couplings to SM fermions. In particular, the Z' boson is assumed to have couplings with charged leptons except for the electron. Such new interactions lead to dimension-6 operators with flavor violation that are constrained by the limits or measurements of various observables, of which we find that the B - \bar{B} mixing data are very consequential and the empirical bounds for $\mu \rightarrow e\gamma$ and $\tau \rightarrow 3\mu$ often play the major role in further constraining the allowed parameter space in the model.

Through numerical scans of parameters in the lepton sector for both the normal and inverted orderings of neutrino masses, we have obtained sampling benchmark points that are compatible with the different constraints and made predictions particularly for lepton flavor-violating processes in lepton decays, B -meson and kaon decays, and Z boson decays. The viable parameter space depends highly on the structure of the \mathbf{A}_ℓ matrix constructed from the right-handed neutrinos' Yukawa couplings and on the light neutrinos' mass ordering. With the simplest form of \mathbf{A}_ℓ , only the NO case possesses viable parameter space, albeit marginally.

Adopting a less simple choice of A_ℓ with additional complex phases, we demonstrate that both the NO and IO scenarios have allowed parameter space, with the IO case being preferred, and subsequently predict a number of observables. The upper bounds of our predictions for some of these observables can be further probed by searches or measurements in the near future.

Acknowledgments

This work was supported in part by MOE Academic Excellence Program (Grant No. 105R891505) and NCTS of ROC. C.W.C. was also supported in part by MOST of ROC (Grant No. MOST 104-2628-M-002-014-MY4). X.G.H. was also supported in part by MOST of ROC (Grant No. MOST104-2112-M-002-015-MY3) and in part by NSFC (Grant Nos. 11175115 and 11575111), Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education, and Shanghai Key Laboratory for Particle Physics and Cosmology (SKLPPC) (Grant No. 11DZ2260700) of PRC.

Appendix A: Extra constraints on $\mathcal{O}^{2q2\ell}$

At tree level, the Z' interactions in Eq. (13) contribute to $\mu \rightarrow e$ conversion in nuclei via the operator $\mathcal{O}^{2q2\ell}$ in Eq. (12). To calculate the rate $\mathcal{B}(\mu\mathcal{N} \rightarrow e\mathcal{N})$ of $\mu \rightarrow e$ conversion in nucleus \mathcal{N} , we employ the formulas from Ref. [69]. Thus, we arrive at

$$\mathcal{B}(\mu\mathcal{N} \rightarrow e\mathcal{N}) = \frac{m_\mu^5 |(2g_{uue\mu} + g_{dde\mu})V_{\mathcal{N}}^p + (g_{uue\mu} + 2g_{dde\mu})V_{\mathcal{N}}^n|^2}{\omega_{\text{capt}}^{\mathcal{N}}}, \quad (\text{A1})$$

$$\begin{aligned} g_{uue\mu} &= \frac{(V_{\text{CKM}}^\dagger \Delta_q V_{\text{CKM}})_{11} (\Delta_\ell)_{12}}{m_{Z'}^2} = \frac{(\zeta_0 + \zeta_1 y_u^2 + \zeta_2 y_u^4) (\Delta_\ell)_{12}}{m_{Z'}^2}, \\ g_{dde\mu} &= \frac{(\Delta_q)_{11} (\Delta_\ell)_{12}}{m_{Z'}^2} = \frac{[\zeta_0 + |V_{td}|^2 (\zeta_1 y_t^2 + \zeta_2 y_t^4)] (\Delta_\ell)_{12}}{m_{Z'}^2}, \end{aligned} \quad (\text{A2})$$

where $V_{\mathcal{N}}^{p(n)}$ is an overlap integral for the protons (neutrons) in \mathcal{N} and $\omega_{\text{capt}}^{\mathcal{N}}$ the rate of muon capture in \mathcal{N} . Based on the data on $\mu \rightarrow e$ transition in nuclei [39] and the corresponding $V_{\mathcal{N}}^{p(n)}$ and $\omega_{\text{capt}}^{\mathcal{N}}$ values [69], we find the gold limit $\mathcal{B}(\mu\text{Au} \rightarrow e\text{Au})_{\text{exp}} < 7.0 \times 10^{-13}$ at 90% CL [39] to supply the strictest restraint. Using $V_{\text{Au}}^{p(n)} = 0.0974$ (0.146) and $\omega_{\text{capt}}^{\text{Au}} = 13.07 \times 10^6/\text{s}$ [69], we then extract

$$|g_{uue\mu} + 1.14 g_{dde\mu}| < \frac{2.0 \times 10^{-6}}{\text{TeV}^2}. \quad (\text{A3})$$

Since our scans of the benchmark points within the permitted parameter space in the NO (IO) case result in the bound $|(\Delta_\ell)_{12}|/m_{Z'} < 0.058$ (0.064)/TeV, while $|\zeta_1 y_t^2 + \zeta_2 y_t^4|/m_{Z'} < 0.142/\text{TeV}$ from Eq. (24), and from quark data $y_u^2 \sim 10^{-10}$ and $|V_{td}|^2 \sim 7 \times 10^{-5}$ [39], it is evident that choosing $|\zeta_0|/m_{Z'} \lesssim 8 \times 10^{-6}/\text{TeV}$ in Eq. (A2) will make the Z' contributions compatible with the condition in Eq. (A3).

The recent LHC measurements on $pp \rightarrow \mu^+ \mu^-$ [54] translate into restrictions on NP effects on the partonic reactions $\bar{q}q \rightarrow \mu^+ \mu^-$. The relevant Z' couplings are

$$\begin{aligned} g_{uu\mu\mu} &\simeq \frac{\zeta_0 (\Delta_\ell)_{22}}{m_{Z'}^2}, & g_{dd\mu\mu} &= \frac{[\zeta_0 + |V_{td}|^2 (\zeta_1 y_t^2 + \zeta_2 y_t^4)] (\Delta_\ell)_{22}}{m_{Z'}^2}, \\ g_{cc\mu\mu} &\simeq \frac{(\zeta_0 + \zeta_1 y_c^2) (\Delta_\ell)_{22}}{m_{Z'}^2}, & g_{ss\mu\mu} &= \frac{[\zeta_0 + |V_{ts}|^2 (\zeta_1 y_t^2 + \zeta_2 y_t^4)] (\Delta_\ell)_{22}}{m_{Z'}^2}, \\ g_{bb\mu\mu} &= \frac{[\zeta_0 + |V_{tb}|^2 (\zeta_1 y_t^2 + \zeta_2 y_t^4)] (\Delta_\ell)_{22}}{m_{Z'}^2}. \end{aligned} \quad (\text{A4})$$

The aforementioned scans yield in the NO (IO) case $|(\Delta_\ell)_{22}|/m_{Z'} < 0.14 (0.26)/\text{TeV}$. Then, with $|V_{ts}|^2 \sim 0.0016$, $|V_{tb}|^2 \sim 1$, and $y_c^2 \sim 2 \times 10^{-5}$ [39], as well as the other parameter values specified in the preceding paragraph, we derive, in units of TeV^{-2} ,

$$\begin{aligned} |g_{uu\mu\mu}| &\lesssim 2.1 \times 10^{-6}, & |g_{dd\mu\mu}| &\lesssim 4.8 \times 10^{-6}, & |g_{cc\mu\mu}| &\lesssim 7.3 \times 10^{-6}, & |g_{ss\mu\mu}| &\lesssim 6.2 \times 10^{-5}, \\ |g_{bb\mu\mu}| &\lesssim 0.038. \end{aligned} \quad (\text{A5})$$

Most of these numbers are at least three orders of magnitude below their respective bounds inferred in Ref. [23] from the $pp \rightarrow \mu^+ \mu^-$ data [54], except $-0.38 \lesssim g_{bb\mu\mu}^{\text{exp}} \text{TeV}^2 \lesssim 0.46$, which is still an order of magnitude above its Z' counterpart in Eq. (A5).

-
- [1] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **111**, 191801 (2013) doi:10.1103/PhysRevLett.111.191801 [arXiv:1308.1707 [hep-ex]].
 - [2] R. Aaij *et al.* [LHCb Collaboration], JHEP **1602**, 104 (2016) doi:10.1007/JHEP02(2016)104 [arXiv:1512.04442 [hep-ex]].
 - [3] A. Abdesselam *et al.* [Belle Collaboration], arXiv:1604.04042 [hep-ex].
 - [4] S. Wehle *et al.* [Belle Collaboration], Phys. Rev. Lett. **118**, no. 11, 111801 (2017) doi:10.1103/PhysRevLett.118.111801 [arXiv:1612.05014 [hep-ex]].
 - [5] R. Aaij *et al.* [LHCb Collaboration], JHEP **1307**, 084 (2013) doi:10.1007/JHEP07(2013)084 [arXiv:1305.2168 [hep-ex]].
 - [6] R. Aaij *et al.* [LHCb Collaboration], JHEP **1509**, 179 (2015) doi:10.1007/JHEP09(2015)179 [arXiv:1506.08777 [hep-ex]].
 - [7] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **113**, 151601 (2014) doi:10.1103/PhysRevLett.113.151601 [arXiv:1406.6482 [hep-ex]].
 - [8] R. Aaij *et al.* [LHCb Collaboration], arXiv:1705.05802 [hep-ex].
 - [9] G. D'Amico, M. Nardecchia, P. Panci, F. Sannino, A. Strumia, R. Torre, and A. Urbano, arXiv:1704.05438 [hep-ph].
 - [10] W. Altmannshofer, C. Niehoff, P. Stangl, and D.M. Straub, arXiv:1703.09189 [hep-ph].
 - [11] W. Altmannshofer, P. Stangl, and D.M. Straub, arXiv:1704.05435 [hep-ph].
 - [12] B. Capdevila, A. Crivellin, S. Descotes-Genon, J. Matias and J. Virto, arXiv:1704.05340 [hep-ph].
 - [13] A. Celis, J. Fuentes-Martin, A. Vicente and J. Virto, arXiv:1704.05672 [hep-ph].
 - [14] D. Bećirević and O. Sumensari, arXiv:1704.05835 [hep-ph].
 - [15] Y. Cai, J. Gargalionis, M. A. Schmidt and R. R. Volkas, arXiv:1704.05849 [hep-ph].

- [16] D. Ghosh, arXiv:1704.06240 [hep-ph].
- [17] J. F. Kamenik, Y. Soreq and J. Zupan, arXiv:1704.06005 [hep-ph].
- [18] S. Di Chiara, A. Fowlie, S. Fraser, C. Marzo, L. Marzola, M. Raidal and C. Spethmann, arXiv:1704.06200 [hep-ph].
- [19] A. K. Alok, D. Kumar, J. Kumar and R. Sharma, arXiv:1704.07347 [hep-ph].
- [20] A. K. Alok, B. Bhattacharya, A. Datta, D. Kumar, J. Kumar and D. London, arXiv:1704.07397 [hep-ph].
- [21] W. Wang and S. Zhao, arXiv:1704.08168 [hep-ph].
- [22] R. Alonso, P. Cox, C. Han and T. T. Yanagida, arXiv:1704.08158 [hep-ph].
- [23] A. Greljo and D. Marzocca, arXiv:1704.09015 [hep-ph].
- [24] F. Feruglio, P. Paradisi and A. Pattori, arXiv:1705.00929 [hep-ph].
- [25] C. Bonilla, T. Modak, R. Srivastava and J. W. F. Valle, arXiv:1705.00915 [hep-ph].
- [26] J. Ellis, M. Fairbairn and P. Tunney, arXiv:1705.03447 [hep-ph].
- [27] F. Bishara, U. Haisch and P. F. Monni, arXiv:1705.03465 [hep-ph].
- [28] R. Alonso, P. Cox, C. Han and T. T. Yanagida, arXiv:1705.03858 [hep-ph].
- [29] Y. Tang and Y. L. Wu, arXiv:1705.05643 [hep-ph].
- [30] A. Datta, J. Kumar, J. Liao and D. Marfatia, arXiv:1705.08423 [hep-ph].
- [31] D. Das, C. Hati, G. Kumar and N. Mahajan, arXiv:1705.09188 [hep-ph].
- [32] D. Dinh, L. Merlo, S.T. Petcov, and Vega-Alvarez, arXiv:1705.09284 [hep-ph].
- [33] D. Bardhan, P. Byakti and D. Ghosh, arXiv:1705.09305 [hep-ph].
- [34] M. Bordone, D. Buttazzo, G. Isidori and J. Monnard, arXiv:1705.10729 [hep-ph].
- [35] S. Neshatpour, V. G. Chobanova, T. Hurth, F. Mahmoudi and D. Martinez Santos, arXiv:1705.10730 [hep-ph].
- [36] R.S. Chivukula and H. Georgi, Phys. Lett. B **188**, 99 (1987); L.J. Hall and L. Randall, Phys. Rev. Lett. **65**, 2939 (1990); A.J. Buras, P. Gambino, M. Gorbahn, S. Jager, and L. Silvestrini, Phys. Lett. B **500**, 161 (2001) [hep-ph/0007085]; A.J. Buras, Acta Phys. Polon. B **34**, 5615 (2003) [hep-ph/0310208]; A.L. Kagan, G. Perez, T. Volansky, and J. Zupan, Phys. Rev. D **80**, 076002 (2009) [arXiv:0903.1794 [hep-ph]].
- [37] G. D'Ambrosio, G.F. Giudice, G. Isidori, and A. Strumia, Nucl. Phys. B **645**, 155 (2002) [hep-ph/0207036].
- [38] V. Cirigliano, B. Grinstein, G. Isidori, and M.B. Wise, Nucl. Phys. B **728**, 121 (2005) [hep-ph/0507001].
- [39] C. Patrignani *et al.* [Particle Data Group], Chin. Phys. C **40**, no. 10, 100001 (2016).
- [40] X.G. He, C.J. Lee, J. Tandean, and Y.J. Zheng, Phys. Rev. D **91**, no. 7, 076008 (2015) [arXiv:1411.6612].
- [41] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p. 95; Prog. Theor. Phys. **64**, 1103 (1980); M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam, 1979), p. 315; P. Ramond, arXiv:hep-ph/9809459; S.L. Glashow, in *Proceedings of the 1979 Cargese Summer Institute on Quarks and Leptons*, edited by M. Levy *et al.* (Plenum Press, New York, 1980), p. 687; R.N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980); J. Schechter and J.W.F. Valle, Phys. Rev. D **22**, 2227 (1980); Phys. Rev. D **25**, 774 (1982).
- [42] R. Foot, H. Lew, X.G. He, and G.C. Joshi, Z. Phys. C **44**, 441 (1989).
- [43] M.B. Gavela, T. Hambye, D. Hernandez, and P. Hernandez, JHEP **0909**, 038 (2009)

- [arXiv:0906.1461].
- [44] M. Magg and C. Wetterich, Phys. Lett. B **94**, 61 (1980); T.P. Cheng and L.F. Li, Phys. Rev. D **22**, 2860 (1980); R.N. Mohapatra and G. Senjanovic, Phys. Rev. D **23**, 165 (1981); G. Lazarides, Q. Shafi, and C. Wetterich, Nucl. Phys. B **181**, 287 (1981).
 - [45] G.C. Branco, A.J. Buras, S. Jager, S. Uhlig, and A. Weiler, JHEP **0709**, 004 (2007) [hep-ph/0609067];
 - [46] S. Davidson and F. Palorini, Phys. Lett. B **642**, 72 (2006) [hep-ph/0607329]; A.S. Joshipura, K.M. Patel, and S.K. Vempati, Phys. Lett. B **690**, 289 (2010) [arXiv:0911.5618]; R. Alonso *et al.*, JHEP **1106**, 037 (2011) [arXiv:1103.5461]; D. Aristizabal Sierra, A. Degee, and J.F. Kamenik, JHEP **1207**, 135 (2012) [arXiv:1205.5547]; A. Pilaftsis and D. Teresi, Phys. Rev. D **92**, no. 8, 085016 (2015) [arXiv:1506.08124].
 - [47] X.G. He, C.J. Lee, S.F. Li, and J. Tandean, Phys. Rev. D **89**, 091901 (2014) [arXiv:1401.2615]; JHEP **1408**, 019 (2014) [arXiv:1404.4436].
 - [48] B. Pontecorvo, Sov. Phys. JETP **26** (1968) 984 [Zh. Eksp. Teor. Fiz. **53** (1968) 1717]; Z. Maki, M. Nakagawa, and S. Sakata, Prog. Theor. Phys. **28**, 870 (1962).
 - [49] J.A. Casas and A. Ibarra, Nucl. Phys. B **618**, 171 (2001) [hep-ph/0103065].
 - [50] G. Colangelo, E. Nikolidakis, and C. Smith, Eur. Phys. J. C **59**, 75 (2009) [arXiv:0807.0801 [hep-ph]]; L. Mercolli and C. Smith, Nucl. Phys. B **817**, 1 (2009) [arXiv:0902.1949 [hep-ph]].
 - [51] C.J. Lee and J. Tandean, JHEP **1508**, 123 (2015) [arXiv:1505.04692 [hep-ph]].
 - [52] P. Langacker, Rev. Mod. Phys. **81**, 1199 (2009) [arXiv:0801.1345 [hep-ph]].
 - [53] J. Alcaraz *et al.* [ALEPH and DELPHI and L3 and OPAL Collaborations and LEP Electroweak Working Group], hep-ex/0612034.
 - [54] The ATLAS collaboration [ATLAS Collaboration], ATLAS-CONF-2017-027.
 - [55] A.J. Buras, F. De Fazio, and J. Girrbach, JHEP **1302**, 116 (2013) [arXiv:1211.1896 [hep-ph]].
 - [56] Y. Amhis *et al.*, arXiv:1612.07233 [hep-ex].
 - [57] M. Artuso, G. Borissov, and A. Lenz, Rev. Mod. Phys. **88**, no. 4, 045002 (2016) [arXiv:1511.09466 [hep-ph]]; T. Jubb, M. Kirk, A. Lenz, and G. Tetlalmatzi-Xolocotzi, Nucl. Phys. B **915**, 431 (2017) [arXiv:1603.07770 [hep-ph]].
 - [58] C.W. Chiang, Y.F. Lin, and J. Tandean, JHEP **1111**, 083 (2011) [arXiv:1108.3969 [hep-ph]].
 - [59] A.M. Baldini *et al.* [MEG Collaboration], Eur. Phys. J. C **76**, no. 8, 434 (2016) [arXiv:1605.05081 [hep-ex]].
 - [60] A. Pich, Prog. Part. Nucl. Phys. **75**, 41 (2014) doi:10.1016/j.pnpnp.2013.11.002 [arXiv:1310.7922 [hep-ph]].
 - [61] C.S. Kim, X.B. Yuan, and Y.J. Zheng, Phys. Rev. D **93**, no. 9, 095009 (2016) [arXiv:1602.08107 [hep-ph]].
 - [62] I. Esteban, M.C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, and T. Schwetz, JHEP **1701**, 087 (2017) [arXiv:1611.01514 [hep-ph]]. Online updates available at <http://www.nu-fit.org>.
 - [63] J. Charles, O. Deschamps, S. Descotes-Genon, H. Lacker, A. Menzel, S. Monteil, V. Niess, and J. Ocariz *et al.*, Phys. Rev. D **91**, no. 7, 073007 (2015) [arXiv:1501.05013 [hep-ph]]. Online updates available at <http://ckmfitter.in2p3.fr>.
 - [64] K. W. Edwards *et al.* [CLEO Collaboration], Phys. Rev. D **65**, 111102 (2002) [hep-ex/0204017]; B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. D **73**, 092001 (2006) [hep-ex/0604007]; Phys. Rev. Lett. **99**, 051801 (2007) [hep-ex/0703018].
 - [65] C. Bobeth, M. Gorbahn, T. Hermann, M. Misiak, E. Stamou and M. Steinhauser, Phys. Rev. Lett. **112**, 101801 (2014) doi:10.1103/PhysRevLett.112.101801 [arXiv:1311.0903 [hep-ph]].
 - [66] J.P. Lees *et al.* [BaBar Collaboration], Phys. Rev. Lett. **118**, no. 3, 031802 (2017)

- doi:10.1103/PhysRevLett.118.031802 [arXiv:1605.09637 [hep-ex]]; R. Aaij *et al.* [LHCb Collaboration], arXiv:1703.02508 [hep-ex].
- [67] C.W. Chiang, T. Nomura, and J. Tandean, Phys. Rev. D **87**, 075020 (2013) [arXiv:1302.2894 [hep-ph]].
- [68] C.D. Carone and H. Murayama, Phys. Rev. Lett. **74**, 3122 (1995) [hep-ph/9411256]; Phys. Rev. D **52**, 484 (1995) [hep-ph/9501220]; E. Ma and D.P. Roy, Phys. Rev. D **58**, 095005 (1998) [hep-ph/9806210].
- [69] R. Kitano, M. Koike, and Y. Okada, Phys. Rev. D **66**, 096002 (2002) [Erratum-ibid. D **76**, 059902 (2007)] [hep-ph/0203110].