

Probing the electromagnetic dipole moments of the tau-neutrino in the $U(1)_{B-L}$ model at the ILC and CLIC energies

A. Llamas-Bugarin*,¹ A. Gutiérrez-Rodríguez†,¹ and M. A. Hernández-Ruíz‡²

¹*Facultad de Física, Universidad Autónoma de Zacatecas
Apartado Postal C-580, 98060 Zacatecas, México.*

²*Unidad Académica de Ciencias Químicas, Universidad Autónoma de Zacatecas
Apartado Postal C-585, 98060 Zacatecas, México.*

(Dated: March 12, 2022)

Abstract

In this work we study the sensitivity on the anomalous magnetic and electric dipole moments of the tau-neutrino in the framework of the $SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ electroweak model at future e^+e^- linear colliders as the ILC and CLIC. For our study we consider the process $e^+e^- \rightarrow (Z, Z', \gamma) \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$. For center-of-mass energies of $\sqrt{s} = 1000 - 3000 \text{ GeV}$ and integrated luminosities of $\mathcal{L} = 500 - 2000 \text{ fb}^{-1}$, we derive 95% C.L. limits on the dipole moments $|\mu_{\nu_\tau}(\mu_B)| \leq 6.28 \times 10^{-9}$ and $|d_{\nu_\tau}(ecm)| \leq 1.21 \times 10^{-21}$ improve the existing limits by two or three orders of magnitude. Our study complements other studies on the dipole moments of the tau-neutrino at hadron and e^+e^- colliders.

PACS numbers: 14.60.St, 13.40.Em, 12.15.Mm

Keywords: Non-standard-model neutrinos, Electric and Magnetic Moments, Neutral Currents.

* maria.llamas@fisica.uaz.edu.mx

† alexgu@fisica.uaz.edu.mx

‡ mahernan@uaz.edu.mx

I. INTRODUCTION

In the Standard Model (SM) [1–3] minimally extended with Dirac neutrino masses, the neutrino magnetic moment induced by radiative corrections is unobservably small [4–6],

$$\mu_{\nu_i} = \frac{3m_e G_F}{4\sqrt{2}\pi^2} m_{\nu_i} \simeq 3.1 \times 10^{-19} \left(\frac{m_{\nu_i}}{1 \text{ eV}} \right) \mu_B, \quad (1)$$

where $\mu_B = e/2m_e$ is the Bohr Magneton. Current limits on these magnetic moments are several orders of magnitude larger, so that a magnetic moment close to these limits would indicate a window for probing effects induced by new physics beyond the SM [6]. Similarly, a neutrino electric dipole moment will also point to new physics and will be of relevance in astrophysics and cosmology, as well as terrestrial neutrino experiments [7]. Some bounds on the neutrino magnetic moment are shown in Table I.

TABLE I: Bounds on the neutrino magnetic moment.

Experiment/ Method	Limit	C. L.	Reference
Laboratory experiment Borexino	$\mu_\nu \leq 5.4 \times 10^{-11} \mu_B$	90%	[8]
Laboratory experiment TEXONO	$\mu_\nu < 2.9 \times 10^{-11} \mu_B$	90%	[9]
Cooling rates of white dwarfs	$\mu_\nu \lesssim 10^{-11} \mu_B$	90%	[10]
Cooling rates of red giants	$\mu_\nu \lesssim 3 \times 10^{-12} \mu_B$	90%	[11]
Supernova energy loss	$\mu_\nu \lesssim (1.1 - 2.7) \times 10^{-12} \mu_B$	90%	[12]
Absence of high-energy events in the SN1987A neutrino signal	$\mu_\nu \lesssim 10^{-12} \mu_B$	90%	[13]
Standard model (Dirac mass)	$\mu_\nu \simeq 3.1 \times 10^{-19} (m_\nu/1 \text{ eV}) \mu_B$		[4–6]

In the case of the anomalous magnetic moment of the tau-neutrino, the current best limit on μ_{ν_τ} has been obtained in the Borexino experiment which explores solar neutrinos. Searches for the magnetic moment of the tau-neutrino have also been performed in accelerator experiments. The experiment E872 (DONUT) is based at $\nu_\tau e^-$, $\bar{\nu}_\tau e^-$ elastic scattering. In the CERN experiment WA-066, a limit on μ_{ν_τ} is obtained on an assumed flux of tau-neutrinos in the neutrino beam. The L3 collaboration obtain a limit on the magnetic moment of the tau-neutrino from a sample of e^+e^- annihilation events at the Z resonance.

Experimental limits on the magnetic moment of the tau-neutrino are shown in Table II. Others limits on the magnetic moment of the μ_{ν_τ} are reported in the literature [14–36].

TABLE II: Experimental limits on the magnetic moment of the tau-neutrino.

Experiment	Method	Limit	C. L.	Reference
Borexino	Solar neutrino	$\mu_{\nu_\tau} < 1.9 \times 10^{-10} \mu_B$	90%	[8]
E872 (DONUT)	Accelerator $\nu_\tau e^-, \bar{\nu}_\tau e^-$	$\mu_{\nu_\tau} < 3.9 \times 10^{-7} \mu_B$	90%	[37]
CERN-WA-066	Accelerator	$\mu_{\nu_\tau} < 5.4 \times 10^{-7} \mu_B$	90%	[38]
L3	Accelerator	$\mu_{\nu_\tau} < 3.3 \times 10^{-6} \mu_B$	90%	[39]

The discovery of CP violation in the decays of neutral kaons [40], and later in the decays of neutral B mesons [41] and D^0 [42], shed light on the nature and origin of the violation of this symmetry. The CP violation is one of the open problems of the SM. For this reason, the measurement of large amounts of CP violation can be indicative of signs of new physics. The signs of new physics can be analyzed by investigating the electromagnetic dipole moments of the tau-neutrino such as its magnetic moment (MM) and electric dipole moment (EDM) defined as a source of CP violation.

In the case of the electric dipole moment of the tau-neutrino some theoretical limits are presented in Table III. Others limits on the d_{ν_τ} are reported in the literature [19–27, 29, 30].

TABLE III: Theoretical limits on the electric dipole moment of the electron-neutrino, muon-neutrino and the tau-neutrino.

Particle	Model	Limit	C. L.	Reference
$\nu_{e,\mu}$	Model-independent	$d_{\nu_e,\nu_\mu} < 2 \times 10^{-21} ecm$	95%	[43]
ν_τ	Effective Lagrangian approach	$d_{\nu_\tau} < 5.2 \times 10^{-17} ecm$	95%	[15]
ν_τ	Model-independent	$d_{\nu_\tau} < O(2 \times 10^{-17} ecm)$	95%	[18]
ν_τ	Vector like Multiplets	$d_{\nu_\tau} < O(10^{-18} - 10^{-20} ecm)$	95%	[44]

The $U(1)_{B-L}$ model [45–49] is one of the simplest extensions of the SM with an extra $U(1)$ local gauge symmetry [50], where B and L represent the baryon number and lepton number, respectively. This B-L symmetry plays an important role in various physics scenarios beyond the SM. The features that distinguish the $U(1)_{B-L}$ models from other models are the following: **a)** The gauge $U(1)_{B-L}$ symmetry group is contained in the Grand Unification Theory (GUT) described by a $SO(10)$ group [45]. **b)** The scale of the B-L symmetry breaking is related to the mass scale of the heavy right-handed Majorana neutrino mass terms and provide the well-known see-saw mechanism [51–55] to explain light left-handed neutrino mass. **c)** The B-L symmetry and the scale of its breaking are tightly connected to the baryogenesis mechanism through leptogenesis [6]. **d)** Another distinctive feature of the $U(1)_{B-L}$ models is the possibility of the Z' heavy boson decaying into pairs of heavy neutrinos $\Gamma(Z' \rightarrow \nu_h \bar{\nu}_h)$. The model contains an extra gauge boson Z' corresponding to B-L gauge symmetry and an extra SM singlet scalar (heavy Higgs boson H). These new particles can change the SM phenomenology significantly and lead to interesting signatures at the current and future colliders such as the Large Hadron Collider (LHC) [56, 57], International Linear Collider (ILC) [58–63] and the Compact Linear Collider (CLIC) [64–66].

The B-L model [67, 68] is attractive due to its relatively simple theoretical structure. The crucial test of the model is the detection of the new heavy neutral (Z') gauge boson and the new Higgs boson (H). On the other hand, searches for both the heavy gauge boson (Z') and the additional heavy neutral Higgs boson (H) predicted by the B-L model are presently being conducted at the LHC. In this regard, the additional boson Z' of the B-L model has a mass which is given by the relation $M_{Z'} = 2v'g'_1$ [48, 49, 67, 68]. This boson Z' interacts with the leptons, quarks, heavy neutrinos and light neutrinos with interaction strengths proportional to the B-L gauge coupling g'_1 . The sensitivity limits on the mass $M_{Z'}$ of the boson Z' of the $U(1)_{B-L}$ model derived for the ATLAS and CMS collaborations are of the order of $\mathcal{O}(1.83 - 2.65) TeV$ [69–77]. It is noteworthy that future LHC runs at 13-14 TeV could increase the Z' mass bounds to higher values, or evidence may be found of its existence. Precision studies of the Z' properties will require a new linear collider [78], which will allow us to perform precision studies of the Higgs sector. We refer the readers to Refs. [48, 49, 67, 68, 79–84] for a detailed description of the B-L model.

Our aim in the present paper is to analyze the reaction $e^+e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$ in the framework of the $U(1)_{B-L}$ model and we attribute an anomalous magnetic moment and an electric dipole

moment to a massive tau-neutrino. It is worth mentioning that at higher s , the dominant contribution involves the exchange of the Z, Z' bosons. The dependence on the magnetic moment (μ_{ν_τ}) and the electric dipole moment (d_{ν_τ}) comes from the radiation of the photon observed by the neutrino or antineutrino in the final state. However, in order to improve the limits on the magnetic moment and the electric dipole moment of the tau-neutrino, in our calculation of the process $e^+e^- \rightarrow \nu_\tau\bar{\nu}_\tau\gamma$ we consider the contribution that involves the exchange of a virtual photon. In this case, the dependence on the dipole moments comes from a direct coupling to the virtual photon, and the observed photon is a result of initial-state Bremsstrahlung. The Feynman diagrams which give the most important contribution to the cross section are shown in Fig. 1. This process sets limits on the tau-neutrino MM and EDM. In this paper, we take advantage of this fact to set limits on μ_{ν_τ} and d_{ν_τ} for integrated luminosities of $500 - 2000 \text{ fb}^{-1}$ and center-of-mass energies between $1000 - 3000 \text{ GeV}$, that is to say in the next generation of linear colliders, namely, the International Linear Collider (ILC) [58] and the Compact Linear Collider (CLIC) [64].

The L3 Collaboration [39] evaluated the selection efficiency using detector-simulated $e^+e^- \rightarrow \nu\bar{\nu}\gamma(\gamma)$ events, random trigger events, and large-angle $e^+e^- \rightarrow e^+e^-$ events. From Fig. 1 of Ref. [39] the process $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ with γ emitted in the initial state is the sole background in the $[44.5^0, 135.5^0]$ angular range (white histogram). From the same figure in this angular interval that is $-0.7 < \cos\theta_\gamma < 0.7$ we see that only 6 events were found, this is the real background, not 14 events. In this case a simple method [24, 85, 86] is that at 1σ level (68% *C.L.*) for a null signal the number of observed events should not exceed the fluctuation of the estimated background events: $N = N_B + \sqrt{N_B}$. Of course, this method is good only when N_B is sufficiently large (i.e. when the Poisson distribution can be approximated with a gaussian [24, 85, 86]) but for $N_B > 10$ it is a good approximation. This means that at $1\sigma, 2\sigma, 3\sigma$ level (68%, 90%, 95% *C.L.*) the limits on the non-standard parameters are found replacing the equation for the total number of events expected $N = N_B + \sqrt{N_B}$ in the expression $N = \sigma(\mu_{\nu_\tau}, d_{\nu_\tau})\mathcal{L}$. The distributions of the photon energy and the cosine of its polar angle are consistent with SM predictions.

This paper is organized as follows: In Section II, we present the B-L theoretical model. In Sec. III we present the calculation of the process $e^+e^- \rightarrow \nu_\tau\bar{\nu}_\tau\gamma$ in the context of the B-L model. Finally, we present our results and conclusions in Sect. IV.

II. BRIEF REVIEW OF THE B-L THEORETICAL MODEL

The solid evidence for the non-vanishing neutrino masses has been confirmed by various neutrino oscillation phenomena and indicates the evidence of new physics beyond the SM. In the SM, neutrinos are massless due to the absence of right-handed neutrinos and the exact B-L conservation. The most attractive idea to naturally explain the tiny neutrino masses is the seesaw mechanism [52–54, 87], in which the right-handed (RH) neutrinos singlet under the SM gauge group is introduced. The gauged $U(1)_{B-L}$ model based on the gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ [51, 88] is an elegant and simple extension of the SM in which the RH heavy neutrinos are essential both for anomaly cancelation and preserving gauge invariance. In addition, the mass of RH neutrinos arises associated with the $U(1)_{B-L}$ gauge symmetry breaking. Therefore, the fact that neutrinos are massive indicates that the SM requires extension.

We consider a $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ model, which is one of the simplest extensions of the SM [48, 49, 51, 67, 79–84, 88], where $U(1)_{B-L}$, represents the additional gauge symmetry. The gauge invariant Lagrangian of this model is given by

$$\mathcal{L} = \mathcal{L}_s + \mathcal{L}_{YM} + \mathcal{L}_f + \mathcal{L}_Y, \quad (2)$$

where $\mathcal{L}_s, \mathcal{L}_{YM}, \mathcal{L}_f$ and \mathcal{L}_Y are the scalar, Yang-Mills, fermion and Yukawa sector, respectively.

The model consists of one doublet Φ and one singlet χ and we briefly describe the Lagrangian including the scalar, fermion and gauge sector, respectively. The Lagrangian for the gauge sector is given by [48, 83, 89, 90],

$$\mathcal{L}_g = -\frac{1}{4}B_{\mu\nu}B^{\mu\nu} - \frac{1}{4}W_{\mu\nu}^a W^{a\mu\nu} - \frac{1}{4}Z'_{\mu\nu}Z'^{\mu\nu}, \quad (3)$$

where $W_{\mu\nu}^a, B_{\mu\nu}$ and $Z'_{\mu\nu}$ are the field strength tensors for $SU(2)_L, U(1)_Y$ and $U(1)_{B-L}$, respectively.

The Lagrangian for the scalar sector of the model is

$$\mathcal{L}_s = (D^\mu\Phi)^\dagger(D_\mu\Phi) + (D^\mu\chi)^\dagger(D_\mu\chi) - V(\Phi, \chi), \quad (4)$$

where the potential term is [81],

$$V(\Phi, \chi) = m^2(\Phi^\dagger\Phi) + \mu^2|\chi|^2 + \lambda_1(\Phi^\dagger\Phi)^2 + \lambda_2|\chi|^4 + \lambda_3(\Phi^\dagger\Phi)|\chi|^2, \quad (5)$$

with Φ and χ as the complex scalar Higgs doublet and singlet fields, respectively. The covariant derivative is given by [79–81]

$$D_\mu = \partial_\mu + ig_s t^\alpha G_\mu^\alpha + i[gT^a W_\mu^a + g_1 Y B_\mu + (\tilde{g}Y + g'_1 Y_{B-L})B'_\mu], \quad (6)$$

where g_s , g , g_1 and g'_1 are the $SU(3)_C$, $SU(2)_L$, $U(1)_Y$ and $U(1)_{B-L}$ couplings with t^α , T^a , Y and Y_{B-L} being their respective group generators. The mixing between the two Abelian groups is described by the new coupling \tilde{g} . The electromagnetic charges on the fields are the same as those of the SM and the Y_{B-L} charges for quarks, leptons and the scalar fields are given by: $Y_{B-L}^{\text{quarks}} = 1/3$, $Y_{B-L}^{\text{leptons}} = -1$ with no distinction between generations for ensuring universality, $Y_{B-L}(\Phi) = 0$ and $Y_{B-L}(\chi) = 2$ [48, 49, 79–81] to preserve the gauge invariance of the model, respectively.

An effective coupling and effective charge such as g' and Y' are usually introduced as $g'Y' = \tilde{g}Y + g'_1 Y_{B-L}$ and some specific benchmark models [91, 92] can be recovered by particular choices of both \tilde{g} and g'_1 gauge couplings at a given scale, generally the electroweak scale. For instance, the pure B-L model is obtain by the condition $\tilde{g} = 0$ ($Y' = Y_{B-L}$) which implies the absence of mixing at the electroweak scale. Other benchmark models of the general parameterisation are the Sequential Standar Model (SSM), the $U(1)_R$ model and the $U(1)_\chi$ model. The SSM is reproduced by the condition $g'_1 = 0$ ($Y' = Y$), and the $U(1)_R$ extension is realised by the condition $\tilde{g} = -2g'_1$, while the $SO(10)$ -inspired $U(1)_\chi$ model is described by $\tilde{g} = -\frac{4}{5}g'_1$.

The doublet and singlet scalars are

$$\Phi = \begin{pmatrix} G^\pm \\ \frac{v+\phi^0+iG_Z}{\sqrt{2}} \end{pmatrix}, \quad \chi = \left(\frac{v' + \phi'^0 + iz'}{\sqrt{2}} \right), \quad (7)$$

with G^\pm , G_Z and z' the Goldstone bosons of W^\pm , Z and Z' , respectively, while $v \approx 246 \text{ GeV}$ is the electroweak symmetry breaking scale and v' is the B-L symmetry breaking scale constrained by the electroweak precision measurement data whose value is assumed to be of the order TeV .

After spontaneous symmetry breaking, the two scalar fields can be written as,

$$\Phi = \begin{pmatrix} 0 \\ \frac{v+\phi^0}{\sqrt{2}} \end{pmatrix}, \quad \chi = \frac{v' + \phi'^0}{\sqrt{2}}, \quad (8)$$

with v and v' real and positive.

In Table IV, the interactions of h and H with the gauge bosons and scalar are expressed in terms of the parameters of the B-L model.

To determine the mass spectrum of the gauge bosons, we have to expand the scalar kinetic terms as with the SM. We expect that there exists a massless gauge boson, the photon, while the other gauge bosons become massive. The extension we are studying is in the Abelian sector of the SM gauge group, so that the charged gauge bosons W^\pm will have masses given by their SM expressions related to the $SU(2)_L$ factor only. The other gauge boson masses are not so simple to identify because of mixing. In fact, analogous to the SM, the fields of definite mass are linear combinations of B^μ , W_3^μ and B'^μ , the relation between the neutral gauge bosons (B^μ , W_3^μ and B'^μ) and the corresponding mass eigenstates is given by [67, 68, 79, 80]

$$\begin{pmatrix} B^\mu \\ W_3^\mu \\ B'^\mu \end{pmatrix} = \begin{pmatrix} \cos \theta_W & -\sin \theta_W \cos \theta_{B-L} & \sin \theta_W \sin \theta_{B-L} \\ \sin \theta_W & \cos \theta_W \cos \theta_{B-L} & -\cos \theta_W \sin \theta_{B-L} \\ 0 & \sin \theta_{B-L} & \cos \theta_{B-L} \end{pmatrix} \begin{pmatrix} A^\mu \\ Z^\mu \\ Z'^\mu \end{pmatrix}, \quad (9)$$

with $-\frac{\pi}{4} \leq \theta_{B-L} \leq \frac{\pi}{4}$, such that

$$\tan 2\theta_{B-L} = \frac{2\tilde{g}\sqrt{g^2 + g_1^2}}{\tilde{g}^2 + 16\left(\frac{v'}{v}\right)^2 g_1'^2 - g^2 - g_1^2}, \quad (10)$$

and the mass spectrum of the gauge bosons is given by

$$\begin{aligned} M_\gamma &= 0, \\ M_{W^\pm} &= \frac{1}{2}vg, \\ M_Z &= \frac{v}{2}\sqrt{g^2 + g_1^2} \sqrt{\frac{1}{2}\left(\frac{\tilde{g}^2 + 16\left(\frac{v'}{v}\right)^2 g_1'^2}{g^2 + g_1^2} + 1\right) - \frac{\tilde{g}}{\sin 2\theta_{B-L}\sqrt{g^2 + g_1^2}}}, \\ M_{Z'} &= \frac{v}{2}\sqrt{g^2 + g_1^2} \sqrt{\frac{1}{2}\left(\frac{\tilde{g}^2 + 16\left(\frac{v'}{v}\right)^2 g_1'^2}{g^2 + g_1^2} + 1\right) + \frac{\tilde{g}}{\sin 2\theta_{B-L}\sqrt{g^2 + g_1^2}}}, \end{aligned} \quad (11)$$

where M_Z and M_{W^\pm} are the SM gauge bosons masses and $M_{Z'}$ is the mass of new neutral gauge boson Z' , which strongly depends on v' and g'_1 . For $\tilde{g} = 0$, there is no mixing between the new and SM gauge bosons Z' and Z . In this case, the $U(1)_{B-L}$ model is called the pure or minimal model $U(1)_{B-L}$. In this article we consider the case $\tilde{g} \neq 0$, which is mostly determined by the other gauge couplings g_1 and g'_1 [93–95]. The electroweak precision measurement data can give stringent constraints on the $Z - Z'$ mixing angle θ_{B-L} expressed in Eq. (10) [96].

In the Lagrangian of the $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ model, the terms for the interactions between neutral gauge bosons Z, Z' and a pair of fermions of the SM can be written in the form [33, 48, 49, 97–99]

$$\mathcal{L}_{NC} = \frac{-ig}{\cos\theta_W} \sum_f \bar{f}\gamma^\mu \frac{1}{2}(g_V^f - g_A^f\gamma^5)fZ_\mu + \frac{-ig}{\cos\theta_W} \sum_f \bar{f}\gamma^\mu \frac{1}{2}(g_V^{\prime f} - g_A^{\prime f}\gamma^5)fZ'_\mu. \quad (12)$$

From this Lagrangian we determine the expressions for the new couplings of the Z, Z' bosons with the SM fermions, which are given in Table IV. The couplings g_V^f ($g_V^{\prime f}$) and g_A^f ($g_A^{\prime f}$) depend on the $Z - Z'$ mixing angle θ_{B-L} and the coupling constant g'_1 of the B-L interaction. In these couplings, the current bound on the mixing angle is $|\theta_{B-L}| \leq 10^{-3}$ [24]. In the decoupling limit, when $\theta_{B-L} = 0$ and $g'_1 = 0$, the couplings of the SM are recovered.

III. THE DECAY WIDTHS OF THE Z' BOSON IN THE B-L MODEL

In this section we present the decay widths of the Z' boson [97, 98, 100–104] in the context of the B-L model needed in the calculation of the cross section for the process $e^+e^- \rightarrow \nu_\tau\bar{\nu}_\tau\gamma$. The decay width of the Z' boson to fermions is given by

$$\Gamma(Z' \rightarrow f\bar{f}) = \frac{2G_F}{3\pi\sqrt{2}} N_c M_Z^2 M_{Z'} \sqrt{1 - \frac{4M_f^2}{M_{Z'}^2}} \left[(g_V^{\prime f})^2 \left\{ 1 + 2\left(\frac{M_f^2}{M_{Z'}^2}\right) \right\} + (g_A^{\prime f})^2 \left\{ 1 - 4\left(\frac{M_f^2}{M_{Z'}^2}\right) \right\} \right], \quad (13)$$

where N_c is the color factor ($N_c = 1$ for leptons, $N_c = 3$ for quarks) and the couplings $g_V^{\prime f}$ and $g_A^{\prime f}$ of the Z' boson with the SM fermions are given in Table IV.

The decay width of the Z' boson to heavy neutrinos is

TABLE IV: The new couplings of the Z, Z' bosons with the SM fermions and vector boson, scalar coupling in the B-L model. $g = e/\sin\theta_W$ and θ_{B-L} is the $Z - Z'$ mixing angle.

Particle	Couplings
$f\bar{f}Z$	$g_V^f = T_3^f \cos\theta_{B-L} - 2Q_f \sin^2\theta_W \cos\theta_{B-L} + \frac{2g_1'}{g} \cos\theta_W \sin\theta_{B-L},$ $g_A^f = T_3^f \cos\theta_{B-L}$
$f\bar{f}Z'$	$g_V^{'f} = -T_3^f \sin\theta_{B-L} - 2Q_f \sin^2\theta_W \sin\theta_{B-L} + \frac{2g_1'}{g} \cos\theta_W \cos\theta_{B-L},$ $g_A^{'f} = -T_3^f \sin\theta_{B-L}$
$Z_\mu Z'_\nu h$	$g_{ZZ'h} = 2i[\frac{1}{4}v \cos\alpha f(\theta_{B-L}, g_1') - v' \sin\alpha g(\theta_{B-L}, g_1')]g_{\mu\nu},$ $f(\theta_{B-L}, g_1') = -\sin(2\theta')(g_1^2 + g_2^2 + g_1'^2) - 2\cos(2\theta')g_1\sqrt{g_1^2 + g_2^2},$ $g(\theta_{B-L}, g_1') = \frac{1}{4}\sin(2\theta')g_1'^2$
$Z_\mu Z'_\nu H$	$g_{ZZ'H} = 2i[\frac{1}{4}v \sin\alpha f(\theta_{B-L}, g_1') + v' \cos\alpha g(\theta_{B-L}, g_1')]g_{\mu\nu},$
$W_\mu^-(p_1)W_\nu^+(p_2)Z'_\rho(p_3)$	$g_{W^-W^+Z'} = -ig \cos\theta_W \sin\theta_{B-L}[(p_1 - p_2)_\rho g_{\mu\nu} + (p_2 - p_3)_\mu g_{\nu\rho} + (p_3 - p_1)_\nu g_{\rho\mu}],$

$$\Gamma(Z' \rightarrow \nu_h \bar{\nu}_h) = \frac{g_1'^2}{24\pi} \sin^2\theta_{B-L} M_{Z'} \sqrt{\left(1 - \frac{4M_{\nu_h}^2}{M_{Z'}^2}\right)^3}, \quad (14)$$

where the width given by Eq. (14) implies that the right-handed neutrino must be lighter than half the Z' mass, $M_{\nu_h} < \frac{M_{Z'}}{2}$, and the conditions under which this inequality holds is for coupled heavy neutrinos, i.e. with minor mass less than $\frac{M_{Z'}}{2}$. The possibility of the Z' heavy boson decaying into pairs of heavy neutrinos is certainly one of the most interesting of its features.

The Z' partial decay widths involving vector bosons and the scalar bosons are

$$\Gamma(Z' \rightarrow W^+W^-) = \frac{G_F M_W^2}{24\pi\sqrt{2}} \cos^2\theta_W \sin^2\theta_{B-L} M_{Z'} \left(\frac{M_{Z'}}{M_Z}\right)^4 \sqrt{\left(1 - 4\frac{M_W^2}{M_{Z'}^2}\right)^3} \left[1 + 20\frac{M_W^2}{M_{Z'}^2} + 12\frac{M_W^4}{M_{Z'}^4}\right], \quad (15)$$

$$\Gamma(Z' \rightarrow Zh) = \frac{G_F M_Z^2 M_{Z'}}{24\pi\sqrt{2}} \sqrt{\lambda_h} \left[\lambda_h + 12\frac{M_Z^2}{M_{Z'}^2}\right] \left[f(\theta_{B-L}, g_1') \cos\alpha + g(\theta_{B-L}, g_1') \sin\alpha\right]^2, \quad (16)$$

$$\Gamma(Z' \rightarrow ZH) = \frac{G_F M_Z^2 M_{Z'}}{24\pi\sqrt{2}} \sqrt{\lambda_H} \left[\lambda_H + 12 \frac{M_Z^2}{M_{Z'}^2} \right] \left[f(\theta_{B-L}, g'_1) \sin \alpha - g(\theta_{B-L}, g'_1) \cos \alpha \right]^2, \quad (17)$$

where

$$\begin{aligned} \lambda_{h,H} \left(1, \frac{M_Z^2}{M_{Z'}^2}, \frac{M_{h,H}^2}{M_{Z'}^2} \right) &= 1 + \left(\frac{M_Z^2}{M_{Z'}^2} \right)^2 + \left(\frac{M_{h,H}^2}{M_{Z'}^2} \right)^2 - 2 \left(\frac{M_Z^2}{M_{Z'}^2} \right) - 2 \left(\frac{M_{h,H}^2}{M_{Z'}^2} \right) - 2 \left(\frac{M_Z^2}{M_{Z'}^2} \right) \left(\frac{M_{h,H}^2}{M_{Z'}^2} \right), \\ f(\theta_{B-L}, g'_1) &= \left(1 + \frac{v^2 g_1'^2}{4M_Z^2} \right) \sin(2\theta_{B-L}) + \left(\frac{v g_1'}{M_Z} \right) \cos(2\theta_{B-L}), \\ g(\theta_{B-L}, g'_1) &= \left(\frac{v v'}{4M_Z^2} \right) g_1'^2 \sin(2\theta_{B-L}). \end{aligned} \quad (18)$$

In the B-L model, the heavy gauge boson mass $M_{Z'}$ satisfies the relation $M_{Z'} = 2v'g'_1$ [48, 49, 67, 68, 79, 80], and considering the most recent limit from $\frac{M_{Z'}}{g'_1} \geq 6.9 \text{ TeV}$ [92, 105, 106], it is possible to obtain a direct bound on the B-L breaking scale v' . In our next numerical calculation, we will take $v' = 3.45 \text{ TeV}$, while $\alpha = \frac{\pi}{9}$ for the $h - H$ mixing angle in correspondence with Refs. [48, 56, 57, 107].

IV. THE TOTAL CROSS SECTION

In this section we calculate the total cross section for the reaction $e^+e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$. The respective transition amplitudes are thus given by

$$\begin{aligned} \mathcal{M}_1 &= \frac{-g^2}{4 \cos^2 \theta_W (l^2 - m_\nu^2)} \left[\bar{u}(p_3) \Gamma^\alpha (l + m_\nu) \gamma^\beta (g_V^\nu - g_A^\nu \gamma_5) v(p_4) \right] \\ &\times \frac{(g_{\alpha\beta} - p_\alpha p_\beta / M_Z^2)}{[(p_1 + p_2)^2 - M_Z^2 - iM_Z \Gamma_Z]} \left[\bar{u}(p_2) \gamma^\alpha (g_V^e - g_A^e \gamma_5) v(p_1) \right] \epsilon_\alpha^\lambda, \end{aligned} \quad (19)$$

$$\begin{aligned} \mathcal{M}_2 &= \frac{-g^2}{4 \cos^2 \theta_W (l'^2 - m_\nu^2)} \left[\bar{u}(p_3) \gamma^\beta (g_V^\nu - g_A^\nu \gamma_5) (l' + m_\nu) \Gamma^\alpha v(p_4) \right] \\ &\times \frac{(g_{\alpha\beta} - p_\alpha p_\beta / M_Z^2)}{[(p_1 + p_2)^2 - M_Z^2 - iM_Z \Gamma_Z]} \left[\bar{u}(p_2) \gamma^\alpha (g_V^e - g_A^e \gamma_5) v(p_1) \right] \epsilon_\alpha^\lambda, \end{aligned} \quad (20)$$

$$\begin{aligned} \mathcal{M}_3 &= \frac{-g^2}{4 \cos^2 \theta_W (r^2 - m_\nu^2)} \left[\bar{u}(p_3) \Gamma^\alpha (r + m_\nu) \gamma^\beta (g_V^\nu - g_A^\nu \gamma_5) v(p_4) \right] \\ &\times \frac{(g_{\alpha\beta} - p_\alpha p_\beta / M_{Z'}^2)}{[(p_1 + p_2)^2 - M_{Z'}^2 - iM_{Z'} \Gamma_{Z'}]} \left[\bar{u}(p_2) \gamma^\alpha (g_V^e - g_A^e \gamma_5) v(p_1) \right] \epsilon_\alpha^\lambda, \end{aligned} \quad (21)$$

$$\begin{aligned} \mathcal{M}_4 &= \frac{-g^2}{4 \cos^2 \theta_W (r'^2 - m_\nu^2)} \left[\bar{u}(p_3) \gamma^\beta (g_V^\nu - g_A^\nu \gamma_5) (\not{r}' + m_\nu) \Gamma^\alpha v(p_4) \right] \\ &\times \frac{(g_{\alpha\beta} - p_\alpha p_\beta / M_{Z'}^2)}{\left[(p_1 + p_2)^2 - M_{Z'}^2 - i M_{Z'} \Gamma_{Z'} \right]} \left[\bar{u}(p_2) \gamma^\alpha (g_V^e - g_A^e \gamma_5) v(p_1) \right] \epsilon_\alpha^\lambda, \end{aligned} \quad (22)$$

$$\mathcal{M}_5 = \frac{e^2}{(k'^2 - m_e^2)} \left[\bar{u}(p_3) \Gamma^\alpha v(p_4) \right] \frac{g_{\alpha\beta}}{(p_1 + p_2)^2} \left[\bar{u}(p_2) \gamma^\alpha (\not{k}' + m_e) \gamma^\beta v(p_1) \right] \epsilon_\alpha^\lambda, \quad (23)$$

and

$$\mathcal{M}_6 = \frac{e^2}{(k'^2 - m_e^2)} \left[\bar{u}(p_3) \Gamma^\alpha v(p_4) \right] \frac{g_{\alpha\beta}}{(p_1 + p_2)^2} \left[\bar{u}(p_2) \gamma^\beta (\not{k}' + m_e) \gamma^\alpha v(p_1) \right] \epsilon_\alpha^\lambda, \quad (24)$$

where the most general expression consistent with Lorentz and electromagnetic gauge invariance, for the tau-neutrino electromagnetic vertex may be parameterized in terms of four form factors:

$$\Gamma^\alpha = e F_1(q^2) \gamma^\alpha + \frac{ie}{2m_{\nu_\tau}} F_2(q^2) \sigma^{\alpha\mu} q_\mu + e F_3(q^2) \gamma_5 \sigma^{\alpha\mu} q_\mu + e F_4(q^2) \gamma_5 (\gamma^\mu q^2 - \not{q} q^\mu), \quad (25)$$

where e is the charge of the electron, m_{ν_τ} is the mass of the tau-neutrino, q^μ is the photon momentum, and $F_{1,2,3,4}(q^2)$ are the electromagnetic form factors of the neutrino, corresponding to charge radius, MM, EDM and anapole moment (AM), respectively, at $q^2 = 0$ [15, 108–113], while ϵ_α^λ is the polarization vector of the photon. $l, r(k)$ and $l', r'(k')$ stand for the momentum of the virtual neutrino (electron) and antineutrino (positron) respectively. The form factors corresponding to charge radius and the anapole moment, do not concern us here.

The MM and EDM give a contribution to the total cross section for the process $e^+ e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$ of the form:

$$\begin{aligned} \sigma_{Tot}(e^+ e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma) &= \int \frac{\alpha^2}{96\pi} \left(\kappa^2 \mu_B^2 + d_{\nu_\tau}^2 \right) \\ &\times \left\{ 4 \left[\frac{(g_V^e)^2 + (g_A^e)^2}{x_W^2 (1 - x_W)^2} \right] \left[\frac{((g_V^\nu)^2 + (g_A^\nu)^2)(s - 2\sqrt{s} E_\gamma) + (g_A^\nu)^2 E_\gamma^2 \sin^2 \theta_\gamma}{(s - M_Z^2)^2 + M_Z^2 \Gamma_Z^2} \right] \right. \\ &+ 4 \left[\frac{(g_V^e)^2 + (g_A^e)^2}{x_W^2 (1 - x_W)^2} \right] \left[\frac{((g_V^\nu)^2 + (g_A^\nu)^2)(s - 2\sqrt{s} E_\gamma) + (g_A^\nu)^2 E_\gamma^2 \sin^2 \theta_\gamma}{(s - M_{Z'}^2)^2 + M_{Z'}^2 \Gamma_{Z'}^2} \right] \end{aligned}$$

$$\begin{aligned}
& + 32 \left[\frac{s - 2\sqrt{s}E_\gamma + 2E_\gamma^2 - E_\gamma^2 \sin^2 \theta_\gamma}{sE_\gamma^2 \sin^2 \theta_\gamma} \right] \\
& + 6 \left[\frac{(g_V^e g_V^{\prime e} + g_A^e g_A^{\prime e})}{x_W^2 (1 - x_W)^2} \right] \left[\frac{(s - M_Z^2)(s - M_{Z'}^2) + M_Z M_{Z'} \Gamma_Z \Gamma_{Z'}}{[(s - M_Z^2)^2 + M_Z^2 \Gamma_Z^2][(s - M_{Z'}^2)^2 + M_{Z'}^2 \Gamma_{Z'}^2]} \right] \\
& \times \left[(g_V^e g_V^{\prime e} + g_A^e g_A^{\prime e})(s - 2\sqrt{s}E_\gamma) + (g_A^\nu g_A^{\prime \nu})E_\gamma^2 \sin^2 \theta_\gamma \right] \Big\} E_\gamma dE_\gamma d \cos \theta_\gamma,
\end{aligned} \tag{26}$$

where $x_W \equiv \sin^2 \theta_W$ and E_γ , $\cos \theta_\gamma$ are the energy and the opening angle of the emitted photon.

The expression given in Eq. (26) corresponds to the total cross section with the exchange of the Z, Z', γ bosons. The SM expression for the cross section of the reaction $e^+ e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$ can be obtained in the decoupling limit when $\theta_{B-L} = 0$, $g'_1 = 0$ and $\alpha = 0$. In this case, the terms that depend on θ_{B-L} , g'_1 and α in Eq. (26) are zero and Eq. (26) is reduced to the expression given in Ref. [14] for the standard model minimally extended to include massive Dirac neutrinos.

V. RESULTS AND CONCLUSIONS

In order to evaluate the integral of the total cross section as a function of the parameters of the model, that is to say, μ_{ν_τ} and d_{ν_τ} we require cuts on the photon angle and energy to avoid divergences when the integral is evaluated at the important intervals of each experiment. We integrate over θ_γ from 44.5° to 135.5° and E_γ from 15 GeV to 100 GeV . Using the following values for numerical computation [24]: $\sin^2 \theta_W = 0.23126 \pm 0.00022$, $m_\tau = 1776.82 \pm 0.16 \text{ MeV}$, $m_b = 4.6 \pm 0.18 \text{ GeV}$, $m_t = 172 \pm 0.9 \text{ GeV}$, $M_{W^\pm} = 80.389 \pm 0.023 \text{ GeV}$, $M_Z = 91.1876 \pm 0.0021 \text{ GeV}$, $\Gamma_Z = 2.4952 \pm 0.0023 \text{ GeV}$, $M_h = 125 \pm 0.4 \text{ GeV}$, $M_H = 500 \text{ GeV}$ and considering the most recent limit from [92, 105, 106]:

$$\frac{M_{Z'}}{g'_1} \geq 6.9 \text{ TeV}, \tag{27}$$

it is possible to obtain a direct bound on the B-L breaking scale v' and take $v' = 3.45 \text{ TeV}$ and $\alpha = \frac{\pi}{9}$. In our numerical analysis, we obtain the total cross section $\sigma_{Tot} = \sigma_{Tot}(\mu_{\nu_\tau}, d_{\nu_\tau}, \sqrt{s}, M_{Z'}, g'_1, \theta_{B-L}, \alpha)$. Thus, in our numerical computation, we will assume \sqrt{s} , $M_{Z'}$, g'_1 , θ_{B-L} and α as free parameters.

As was discussed in Refs. [14, 39, 114, 115], $N \approx \sigma_{Tot}(\mu_{\nu_\tau}, d_{\nu_\tau}, \sqrt{s}, M_{Z'}, g'_1, \theta_{B-L}, \alpha)\mathcal{L}$, where $N = N_B + \sqrt{N_B}$ is the total number of $e^+e^- \rightarrow \nu_\tau\bar{\nu}_\tau\gamma$ events expected at $1\sigma, 2\sigma, 3\sigma$ level as is mentioned in the introduction and $\mathcal{L} = 500 - 2000 \text{ fb}^{-1}$ according to the data reported by the ILC and CLIC Refs. [58, 64]. Taking this into consideration, we can obtain a limit for the tau-neutrino magnetic moment with $d_{\nu_\tau} = 0$.

TABLE V: Bounds on the μ_{ν_τ} magnetic moment and d_{ν_τ} electric dipole moment for $\sqrt{s} = 1000, 2000, 3000 \text{ GeV}$ and $\mathcal{L} = 500, 1000, 2000 \text{ fb}^{-1}$ at $1\sigma, 2\sigma$ and 3σ .

$\mathcal{L} = 500, 1000, 2000 \text{ fb}^{-1}$		
$\sqrt{s} = 1000 \text{ GeV}; M_{Z'} = 1000 \text{ GeV}, g'_1 = 0.145$		
C. L.	$ \mu_{\nu_\tau}(\mu_B) $	$ d_{\nu_\tau}(\text{ecm}) $
1σ	$(3.11, 2.20, 1.55) \times 10^{-8}$	$(6.01, 4.25, 3.00) \times 10^{-19}$
2σ	$(3.53, 2.50, 1.76) \times 10^{-8}$	$(6.82, 4.82, 3.41) \times 10^{-19}$
3σ	$(3.91, 2.75, 1.95) \times 10^{-8}$	$(7.55, 5.34, 3.77) \times 10^{-19}$
$\sqrt{s} = 2000 \text{ GeV}; M_{Z'} = 2000 \text{ GeV}, g'_1 = 0.290$		
C. L.	$ \mu_{\nu_\tau}(\mu_B) $	$ d_{\nu_\tau}(\text{ecm}) $
1σ	$(1.51, 1.07) \times 10^{-8}, 7.57 \times 10^{-9}$	$(2.92, 2.06, 1.46) \times 10^{-19}$
2σ	$(1.72, 1.21) \times 10^{-8}, 8.60 \times 10^{-9}$	$(3.31, 2.34, 1.65) \times 10^{-19}$
3σ	$(1.90, 1.34) \times 10^{-8}, 9.52 \times 10^{-9}$	$(3.67, 2.59, 1.83) \times 10^{-19}$
$\sqrt{s} = 3000 \text{ GeV}, M_{Z'} = 3000 \text{ GeV}, g'_1 = 0.435$		
C. L.	$ \mu_{\nu_\tau}(\mu_B) $	$ d_{\nu_\tau}(\text{ecm}) $
1σ	$1.00 \times 10^{-8}, (7.07, 5.00) \times 10^{-9}$	$(1.93, 1.36) \times 10^{-19}, 9.65 \times 10^{-20}$
2σ	$1.13 \times 10^{-8}, (8.03, 5.68) \times 10^{-9}$	$(2.19, 1.55, 1.09) \times 10^{-19}$
3σ	$1.25 \times 10^{-8}, (8.89, 6.28) \times 10^{-9}$	$(2.42, 1.71, 2.21) \times 10^{-19}$

As an indicator of the order of magnitude on the dipole moments, we present the bounds obtained on the μ_{ν_τ} magnetic moment and d_{ν_τ} electric dipole moment in Table V for several center-of-mass energies $\sqrt{s} = 1000, 2000, 3000 \text{ GeV}$, integrated luminosity $\mathcal{L} = 500, 1000, 2000 \text{ fb}^{-1}$ and heavy gauge boson masses $M_{Z'} = 1000, 2000, 3000 \text{ GeV}$ with $g'_1 = 0.145, 0.290, 0.435$ at $1\sigma, 2\sigma$ and 3σ , respectively. It is worth mentioning that the values

reported in Table V for the dipole moments are determined while preserving the relationship between $M_{Z'}$ and g'_1 given in Eq. (27). This relationship will always remain throughout the article. We observed that the results obtained in Table V are better than those reported in the literature [14–18, 26–34, 38, 39].

TABLE VI: Bounds on the μ_{ν_τ} magnetic moment and d_{ν_τ} electric dipole moment for $\sqrt{s} = 1000, 2000, 3000 \text{ GeV}$ and $\mathcal{L} = 500, 1000, 2000 \text{ fb}^{-1}$ at $1\sigma, 2\sigma$ and 3σ .

$\mathcal{L} = 500, 1000, 2000 \text{ fb}^{-1}$		
$\sqrt{s} = 1000 \text{ GeV}$		
C. L.	$ \mu_{\nu_\tau}(\mu_B) $	$ d_{\nu_\tau}(\text{ecm}) $
1σ	$(2.22, 1.57, 1.11) \times 10^{-7}$	$(4.29, 3.03, 2.14) \times 10^{-18}$
2σ	$(2.52, 1.78, 1.26) \times 10^{-7}$	$(4.87, 3.44, 2.43) \times 10^{-18}$
3σ	$(2.79, 1.97, 1.39) \times 10^{-7}$	$(5.39, 3.81, 2.69) \times 10^{-18}$
$\sqrt{s} = 2000 \text{ GeV}$		
C. L.	$ \mu_{\nu_\tau}(\mu_B) $	$ d_{\nu_\tau}(\text{ecm}) $
1σ	$1.08 \times 10^{-7}, (7.64, 5.40) \times 10^{-8}$	$(2.08, 1.47, 1.04) \times 10^{-18}$
2σ	$1.22 \times 10^{-7}, (8.68, 6.14) \times 10^{-8}$	$(2.36, 1.67, 1.18) \times 10^{-18}$
3σ	$1.35 \times 10^{-7}, (9.61, 6.79) \times 10^{-8}$	$(2.62, 1.85, 1.31) \times 10^{-18}$
$\sqrt{s} = 3000 \text{ GeV}$		
C. L.	$ \mu_{\nu_\tau}(\mu_B) $	$ d_{\nu_\tau}(\text{ecm}) $
1σ	$(7.14, 5.05, 3.57) \times 10^{-8}$	$1.37 \times 10^{-18}, (9.74, 6.88) \times 10^{-19}$
2σ	$(8.11, 5.73, 4.05) \times 10^{-8}$	$(1.56, 1.10) \times 10^{-18}, 7.82 \times 10^{-19}$
3σ	$(8.97, 6.34, 4.48) \times 10^{-9}$	$(1.73, 1.22) \times 10^{-18}, 8.65 \times 10^{-19}$

The previous analysis and comments can readily be translated to the EDM of the τ -neutrino with $\mu_{\nu_\tau} = 0$. The resulting limits for the EDM as a function of $\sqrt{s}, M_{Z'}$ and g'_1 are shown in Table V.

In the case of the standard model minimally extended [14], i.e. in the decoupling limit when $\theta_{B-L} = 0, g'_1 = 0$ and $\alpha = 0$, the bounds generated on the dipole moments are given in Table VI. These bounds are weaker than those obtained with the $U(1)_{B-L}$ model.

The vector and axial-vector e^+e^-Z couplings g_V^e and g_A^e which depend on g_1' and θ_{B-L} are given in Table IV. To see the dependence of g_V^e and g_A^e on the parameters of the model we plot the relative correction $\frac{\delta g_V^e}{(g_V^e)_{SM}} = \frac{(g_V^e)_{B-L} - (g_V^e)_{SM}}{(g_V^e)_{SM}}$ and $\frac{\delta g_A^e}{(g_A^e)_{SM}} = \frac{(g_A^e)_{B-L} - (g_A^e)_{SM}}{(g_A^e)_{SM}}$ as a function of (g_1', θ_{B-L}) in Fig. 2. From the top panel, we can see that the absolute value of the relative correction $\frac{\delta g_V^e}{(g_V^e)_{SM}}$ increases when the parameter g_1' increases and is almost independent of the mixing angle θ_{B-L} . However, the absolute value of $\frac{\delta g_V^e}{(g_V^e)_{SM}}$ is in the ranges from 10%–70% in most of the parameter space. In the bottom panel, we present the relative correction $\frac{\delta g_A^e}{(g_A^e)_{SM}}$ as a function of g_1' and θ_{B-L} . Here it is shown that the absolute value of $\frac{\delta g_A^e}{(g_A^e)_{SM}}$ increases when the parameter g_1' increases and is almost independent of the mixing angle θ_{B-L} . For $g_1' = 1$, the absolute value of $\frac{\delta g_A^e}{(g_A^e)_{SM}}$ is in the range of 4%. We find that the relative change in g_V^e is much greater than that for g_A^e for the values of the free parameters g_1' and θ_{B-L} near the endpoints. We conclude that the deviations of the couplings g_V^e and g_A^e from its SM value are relatively large in the parameter space (g_1', θ_{B-L}) .

In Fig. 3 we present the total decay width of the Z' boson as a function of $M_{Z'}$ and the new $U(1)_{B-L}$ gauge coupling g_1' , respectively, with the other parameters held fixed to three different values and $\theta_{B-L} = 10^{-3}$. From the top panel, we see that the total width of the Z' new gauge boson varies from very few to hundreds of GeV over a mass range of $1000 GeV \leq M_{Z'} \leq 3500 GeV$, depending on the value of g_1' , when $g_1' = 0.145, 0.290, 0.435$, respectively. In the case of the bottom panel, a similar behavior is obtained in the range $0 \leq g_1' \leq 1$ and depends on the value $M_{Z'} = 1000, 2000, 3000 GeV$. In both figures a clear dependence is observed on the parameters of the $U(1)_{B-L}$ model.

Figure 4 shows the total cross section for $e^+e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$ as a function of the center-of-mass energy \sqrt{s} and different values representative of the magnetic moment, which are reported in the literature, that is to say, $\mu_{\nu_\tau} = 3.3 \times 10^{-6} \mu_B$ (L3), $5.4 \times 10^{-7} \mu_B$ (BEBC (CERN)), $2.75 \times 10^{-8} \mu_B$ (Table V) with $M_{Z'} = 3000 GeV$ and $g_1' = 0.435$. Starting from a center-of-mass energy of the order of the Z mass, a minimum around $\sqrt{s} \simeq 100 GeV$ occurs due to the SM Z -boson resonance tail on the high energies. For different values of the parameter μ_{ν_τ} the shape of the curves does not change and there is only a shift of these depending on the value of the magnetic moment.

The dependence of the sensitivity limits of the magnetic moment μ_{ν_τ} with respect to the collider luminosity \mathcal{L} for three different values of the center-of-mass energy, $\sqrt{s} = 1000, 2000, 3000 GeV$, heavy gauge boson mass of $M_{Z'} = 1000, 2000, 3000 GeV$ and $g_1' =$

0.145, 0.290, 0.435, respectively, is presented in Fig. 5. The figure clearly shows a strong dependence of μ_{ν_τ} with respect to \mathcal{L} and the parameters of the $U(1)_{B-L}$ model. In addition, the spacing between the curves are broader for larger g'_1 values, as the total width of the Z' boson increases with g'_1 , as shown in figure 3. Finally, in order to see how the total cross section $e^+e^- \rightarrow \nu_\tau\bar{\nu}_\tau\gamma$ change with respect to the dipole moments μ_{ν_τ} and d_{ν_τ} we give a 3D plot as shown in Fig. 6. In this figure we consider $M_{Z'} = 3000 \text{ GeV}$ and $g'_1 = 0.435$ in correspondence with Eq. (27).

It is worth mentioning that by reversing the process, we can obtain specific predictions on the $U(1)_{B-L}$ models from the expression of the scattering cross section of the process $e^+e^- \rightarrow \nu_\tau\bar{\nu}_\tau\gamma$. Predictions about the models can be obtained by using the upper bound on the ν_τ magnetic moment reported in the literature by the L3 Collaboration as an input, which maximize the total cross section, namely $\mu_{\nu_\tau} = 3.3 \times 10^{-6} \mu_B$ (90% C.L.) [39], and using the data obtained by the ALEPH Collaboration $\sigma = (3.09 \pm 0.234) \text{ pb}$ Ref. [116, 117] for the cross section, where the error is statistical.

In conclusion, we have found that the process $e^+e^- \rightarrow \nu_\tau\bar{\nu}_\tau\gamma$ in the context of the standard model minimally extended to include massive Dirac neutrino at the high energies and luminosities expected at the ILC/CLIC colliders can be used to probe for bounds on the magnetic moment μ_{ν_τ} and electric dipole moment d_{ν_τ} . In particular, we can appreciate that the 95% C.L. sensitivity limits expected for the magnetic moment at 1000 – 3000 GeV center-of-mass energies already can provide proof of these bounds of order $10^{-8} - 10^{-9}$, that is to say, 2-3 orders of magnitude better than those reported in the literature, see Table II and refs. [14–24, 26–33, 33, 34]. Our results in Table V compare favorably with the limits obtained by the L3 Collaboration [39], and with other limits reported in the literature [14–18, 26–34, 38, 39].

In the case of the electric dipole moment the 95% C.L. sensitivity limits at 1000 – 3000 GeV center-of-mass energies and integrated luminosities of 2000 fb^{-1} can provide proof of these bounds of order $10^{-19} - 10^{-20}$, that is to say, are improved by 2-3 orders of magnitude than those reported in the literature, see Table III and refs. [19–24, 26, 27, 29, 30].

The above results do not appear outside the realm of detection in future experiment with improved sensitivity. In addition, the analytical and numerical results for the cross section could be of relevance for the scientific community. Further, the results above could have possible astrophysical implications. In this regard, the stellar energy loss rates data

have been used to put constraints on the properties and interaction of light particles [118–121]. In addition, one of the most interesting possibilities to use stars as particle physics laboratories [122, 123] is to study the backreaction of the novel energy loss rates implied by the existence of new low-mass particles such as axions [124, 125], or by non-standard neutrino properties such as magnetic moment and electric dipole moment [10, 11, 126–128]. Our study complements other studies on the dipole moments of the tau-neutrino at hadron and e^+e^- colliders.

Acknowledgments

We acknowledge support from CONACyT, SNI and PROFOCIE (México).

-
- [1] S. L. Glashow, *Nucl. Phys.* **22**, 579 (1961).
 - [2] S. Weinberg, *Phys. Rev. Lett.* **19**, 1264 (1967).
 - [3] A. Salam, in *Elementary Particle Theory*, Ed. N. Svartholm (Almqvist and Wiksell, Stockholm, 1968) 367.
 - [4] K. Fujikawa and R. Shrock, *Phys. Rev. Lett.* **45**, 963 (1980).
 - [5] Robert E. Shrock, *Nucl. Phys.* **B206**, 359 (1982).
 - [6] M. Fukugita and T. Yanagida, *Physics of Neutrinos and Applications to Astrophysics*, (Springer, Berlin, 2003).
 - [7] A. Cisneros, *Astrophys. Space Sci.* **10**, 87 (1971).
 - [8] C. Arpesella, *et al.*, [Borexino Collaboration], *Phys. Rev. Lett.* **101**, 091302 (2008).
 - [9] M. Deniz *et al.* [TEXONO Collaboration], *Phys. Rev.* **D81**, 072001 (2010).
 - [10] S. I. Blinnikov, N. V. Dunina-Barkovskaya, *Mon. Not. R. Astron. Soc.* **266**, 289 (1994).
 - [11] G. G. Raffelt, *Phys. Rept.* **320**, 319 (1999) and references therein.
 - [12] A. V. Kuznetsov, N. V. Mikheev and A. A. Okrugin, arXiv:0907.2905 [hep-ph].
 - [13] R. Barbieri and R. N. Mohapatra, *Phys. Rev. Lett.* **61**, 27 (1988).

- [14] T. M. Gould and I. Z. Rothstein, *Phys. Lett.* **B333**, 545 (1994).
- [15] R. Escribano and E. Massó, *Phys. Lett.* **B395**, 369 (1997).
- [16] H. Grotch and R. Robinet, *Z. Phys.* **C39**, 553 (1988).
- [17] Aytekin Aydemir and Ramazan Sever, *Mod. Phys. Lett.* **A16** 7, 457 (2001).
- [18] Keiichi Akama, Takashi Hattori and Kazuo Katsuura, *Phys. Rev. Lett.* **88**, 201601 (2002).
- [19] A. Gutiérrez-Rodríguez, M. Koksál and A. A. Billur, *Phys. Rev.* **D91**, 093008 (2015).
- [20] A. Gutiérrez-Rodríguez, *Int. J. Theor. Phys.* **54**, 236 (2015).
- [21] A. Gutiérrez-Rodríguez, M. Koksál and A. A. Billur, *Proc. Sci.*, EPS-HEP2015 (2015) 036.
- [22] A. Gutiérrez-Rodríguez, *Advances in High Energy Physics* **2014**, 491252 (2014).
- [23] M. A. Hernández-Ruíz, *et al.*, *Nucl. Phys. Proc. Suppl.* **253-255**, 204 (2014).
- [24] K. A. Olive, *et al.*, [Particle Data Group], *Chin. Phys.* **C38**, 090001 (2014).
- [25] A. Gutiérrez-Rodríguez, *Proceedings 20th International Conference on Particles and Nuclei (PANIC 14)*, Hamburg, Germany, August 24-29, edited by A. Schmidt and C. Sander (DESY, Hamburgo, 2015), p. 683.
- [26] A. Gutiérrez-Rodríguez, *Pramana Journal of Physics* **79**, 903 (2012).
- [27] A. Gutiérrez-Rodríguez, *Eur. Phys. J.* **C71**, 1819 (2011).
- [28] C. Aydın, M. Bayar and N. Kilic, *Chin. Phys.* **C32**, 608 (2008).
- [29] A. Gutiérrez-Rodríguez, *et al.*, *Phys. Rev.* **D74**, 053002 (2006).
- [30] A. Gutiérrez-Rodríguez, *et al.*, *Phys. Rev.* **D69**, 073008 (2004).
- [31] A. Gutiérrez-Rodríguez, *et al.*, *Acta Phys. Slov.* **53**, 293 (2003).
- [32] A. Gutiérrez-Rodríguez, *et al.*, *Rev. Mex. Fís.* **45**, 249 (1999).
- [33] A. Gutiérrez-Rodríguez, *et al.*, *Phys. Rev.* **D58**, 117302 (1998).
- [34] P. Abreu, *et al.*, [DELPHI Collaboration], *Z. Phys.* **C74**, 577 (1997).
- [35] I. Sahin, *Phys. Rev. D* **85**, 033002 (2012).
- [36] I. Sahin and M. Koksál, *J. High Energy Phys.* **03**, 100 (2011).
- [37] R. Schwinhorst, *et al.*, [DONUT Collaboration], *Phys. Lett.* **B513**, 23 (2001).
- [38] A. M. Cooper-Sarkar, *et al.*, [WA66 Collaboration], *Phys. Lett.* **B280**, 153 (1992).
- [39] M. Acciarri *et al.*, [L3 Collaboration], *Phys. Lett.* **B412**, 201 (1997).
- [40] J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turlay, *Phys. Rev. Lett.* **13**, 138 (1964).
- [41] Abe K., *et al.*, *Phys. Rev. Lett.* **87**, 091802 (2001).
- [42] R. Aaij, *et al.*, [LHCb Collaboration], *JHEP* **07**, 041 (2014).

- [43] F. del Aguila and M. Sher, *Phys. Lett.* **B252**, 116 (1990).
- [44] Tarek Ibrahim and Pran Nath, *Phys. Rev.* **D81**, 033007 (2010).
- [45] W. Buchmuller, C. Greub, and P. Minkowski, *Phys. Lett.* **B267**, 395 (1991).
- [46] R. Marshak and R. N. Mohapatra, *Phys. Lett.* **B91**, 222 (1980).
- [47] R. N. Mohapatra and R. Marshak, *Phys. Rev. Lett.* **44**, 1316 (1980).
- [48] S. Khalil, *J. Phys. G: Nucl. Part. Phys.* **G35**, 055001 (2008).
- [49] S. Khalil, *Eur. Phys. J.* **C52**, 625 (2007).
- [50] E. D. Carlson, *Nucl. Phys.* **B286**, 378 (1987).
- [51] R.N. Mohapatra and G. Senjanovic, *Phys. Rev. Lett.* **44**, 912 (1980).
- [52] P. Minkowski, *Phys. Lett.* **B67**, 421 (1977).
- [53] P. Van Nieuwenhuizen and D. Z. Freedman, 341 (1979), Amsterdam, Netherlands: North-Holland.
- [54] T. Yanagida, in Proceedings of the Workshop on the *Baryon Number of the Universe and Unified Theories*, Tsukuba, Japan, 13-14 February, (1979), p. 95.
- [55] M. Gell-Mann, P. Ramond, and R. Slansky print-80-0576 (CERN).
- [56] G. Aad, *et al.*, [ATLAS Collaboration], *Phys. Lett.* **B716**, 1 (2012).
- [57] S. Chatrchyan, *et al.*, [CMS Collaboration], *Phys. Lett.* **B716**, 30 (2012).
- [58] T. Abe, *et al.* [Am. LC Group], arXiv: hep-ex/0106057.
- [59] G. Aarons *et al.*, [ILC Collaboration], arXiv: 0709.1893 [hep-ph].
- [60] J. Brau, *et al.*, [ILC Collaboration], arXiv: 0712.1950 [physics.acc-ph].
- [61] H. Baer, T. Barklow, K. Fujii, *et al.*, *The International Linear Collider, Technical Design Report-Vol. 2: Physics*, arXiv:1306.6352 [hep-ph].
- [62] D. M. Asner, *et al.*, *ILC Higgs White Paper*, arXiv: 1310.0763 [hep-ph].
- [63] *Proceedings of the Workshop e^+e^- Collisions at 500 GeV: The Physics Potential*, Munich-Annecy-Hamburg, ed. P. M. Zerwas, Reports DESY 92-123A, B; 93-123C.
- [64] E. Accomando, *et al.* [CLIC Physics Working Group Collaboration], arXiv: hep-ph/0412251, CERN-2004-005.
- [65] H. Abramowicz, *et al.*, *The CLIC Detector and Physics Study*, arXiv:1307.5288 [hep-ex].
- [66] D. Dannheim, P. Lebrun, L. Linssen *et al.*, arXiv: 1208.1402 [hep-ex].
- [67] L. Basso, *et al.*, *Phys. Rev.* **D80**, 055030 (2009).
- [68] L. Basso, *et al.*, *JHEP* **0910**, 006 (2009).

- [69] G. Aad, *et al.*, [ATLAS Collaboration], *Phys. Rev.* **D90**, 052005 (2014).
- [70] ATLAS collaboration, ATLAS-CONF-2015-070.
- [71] CMS Collaboration, CMS-PAS-EXO-12-023.
- [72] V. Khachatryan, *et al.*, [CMS Collaboration], *JHEP* **1504**, 025 (2015).
- [73] V. Khachatryan, *et al.*, [CMS Collaboration], *Phys. Rev.* **D91**, 052009 (2015).
- [74] S. Chatrchyan, *et al.*, [CMS Collaboration], *JHEP* **1209**, 029 (2012).
- [75] G. Aad, *et al.*, [ATLAS Collaboration], [arXiv:1502.07177].
- [76] ATLAS Collaboration, ATLAS-CONF-2013-017.
- [77] CMS Collaboration, CMS-PAS-EXO-12-061.
- [78] B. C. Allanach, *et al.*, arXiv:hep-ph/0403133.
- [79] L. Basso, *et al.*, *Eur. Phys. J.* **C71**, 1613 (2011).
- [80] L. Basso, S. Moretti and G. M. Pruna, *J. Phys. G: Nucl. Part. Phys.* **G39**, 025004 (2012).
- [81] L. Basso, S. Moretti and G. M. Pruna, *Phys. Rev.* **D82**, 055018 (2010).
- [82] L. Basso, *et al.*, *Eur. Phys. J.* **C71**, 1724 (2011).
- [83] L. Basso, arXiv:1106.4462 [hep-hp].
- [84] Satoshi Iso, Nobuchica Ocada and Yuta Orikasa, *Phys. Rev.* **D80**, 115007 (2009).
- [85] Rick S. Gupta and James D. Wells, arXiv: 1110.0824 [hep-ph].
- [86] G. L. Bayatian *et al.*, (CMS Collaboration), *J. Phys.* **G34**, 995 (2007).
- [87] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, Proceedings of Workshop, Stony Brook, New York, 1979, edited by P. Van Nieuwenhuizen and D. Z. Freedman (North-Holland, Amsterdam, 1979), p 315.
- [88] R. E. Marshak and R. N. Mohapatra, *Phys. Lett.* **B91**, 222 (1980).
- [89] A. Ferroglia, A. Lorca and J. J. van der Bij, *Annals of Physics* **16**, 563 (2007) and references therein.
- [90] T. G. Rizzo, arXiv:hep-ph/0610104 and references therein.
- [91] T. Appelquist, B. A. Dobrescu, and A. R. Hopper, *Phys. Rev.* **D68**, 035012, (2003).
- [92] M. Carena, A. Daleo, B. A. Dobrescu, and T. M. Tait, *Phys. Rev.* **D70**, 093009 (2004).
- [93] P. Bandyopadhyay, E. J. Chun, H. Okada and J. C. Park, *JHEP* **1301**, 079 (2013).
- [94] L. Basso, *Phys. Lett.* **B725**, 322 (2013).
- [95] H. Mansour, N. Bakhet, *J. Open Microphys.* **4**, 37 (2013).
- [96] S. Schael, *et al.*, *Phys. Rept.* **427**, 257 (2006).

- [97] A. Gutiérrez-Rodríguez and M. A. Hernández-Ruíz, *Advances in High Energy Physics* **2015**, 593898 (2015).
- [98] F. Ramírez-Sánchez, *et al.*, *J. Phys. G: Nucl. Part. Phys.* **43**, 095003 (2016).
- [99] Shi-Yu, *et al.*, *Int. J. Theor. Phys.* **55**, 648 (2016).
- [100] A. Leike, *Phys. Rep.* **317**, 143 (1999).
- [101] R. W. Robinett and J. L. Rosner, *Phys. Rev.* **D25**, 3036 (1982).
- [102] V. Barger and K. Whisnant, *Phys. Rev.* **D36**, 3429 (1987).
- [103] Pavel Fileviez Perez and Tao Han, Tong Li, *Phys. Rev.* **D80**, 073015 (2009).
- [104] P. Langacker, *Rev. Mod. Phys.* **81**, 1199 (2009).
- [105] J. Heeck, *Phys. Lett.* **B739**, 256 (2014).
- [106] G. Cacciapaglia, C. Csaki, G. Marandella and A. Strumia, *Phys. Rev.* **D74**, 033011 (2006).
- [107] Lorenzo Basso, Stefano Moretti, and Giovanni Marco Pruna, *Phys. Rev.* **D83**, 055014 (2011).
- [108] P. Vogel and J. Engel, *Phys. Rev.* **D39**, 3378 (1989).
- [109] J. Bernabeu, *et al.*, *Phys. Rev.* **D62**, 113012 (2000).
- [110] J. Bernabeu, *et al.*, *Phys. Rev. Lett.* **89**, 101802 (2000); *Phys. Rev. Lett.* **89**, 229902 (2002).
- [111] M. S. Dvornikov and A. I. Studenikin, *Jour. of Exp. and Theor. Phys.* **99**, 254 (2004).
- [112] C. Giunti and A. Studenikin, *Phys. Atom. Nucl.* **72**, 2089 (2009).
- [113] C. Brogini, C. Giunti, A. Studenikin, *Adv. High Energy Phys.* **2012**, 459526 (2012) and references therein.
- [114] R. M. Barnett *et al.*, *Phys. Rev.* **D54**, 166 (1996).
- [115] G. J. Feldman and R. D. Cousins, *Phys. Rev.* **D57**, 3873 (1998).
- [116] A. Heister, *et al.*, [ALEPH Collaboration], *Eur. Phys. J.* **C28**, 1 (2003).
- [117] G. Taylor, *et al.*, [ALEPH Collaboration], ALEPH 99-049, CONF 99-025, TAMPERE ABSTRACT # 6-429, 30 June 1999.
- [118] D. A. Dicus, *Phys. Rev.* **D6**, 941 (1972).
- [119] Duane A. Dicus, *et al.*, *Astrophys. J.* **210**, 481 (1976).
- [120] Stephen W. Bruenn, *Astrophys. J. Suppl. Ser.* **58**, 771 (1985).
- [121] J. Ellis, K. Olive, *Nucl. Phys.* **B223**, 252 (1983).
- [122] G. G. Raffelt, *Stars as laboratories for fundamental physics: The astrophysics of neutrinos, axions, and other weakly interacting particles* (University of Chicago Press, Chicago, USA, 1996) p. 664 and references therein.

- [123] M. A. Hernández-Ruíz, *et al.*, *Eur. Phys. J.* **A53**, 16 (2017).
- [124] A. Payez, *et al.*, *J. Cosmol. Astropart. Phys.* **02**, 006 (2015).
- [125] T. Fischer, *et al.*, *Phys. Rev. D* **94**, 085012 (2016).
- [126] B. K. Kerimov, *et al.*, *Phys. Lett.* **B274**, 477 (1992) and references therein.
- [127] Alexander Heger, *et al.*, *Astrophys. J.* **696**, 608 (2009).
- [128] A. Llamas-Bugarin, A. Gutiérrez-Rodríguez and M. A. Hernández-Ruíz, work in progress.




FIG. 1: The Feynman diagrams contributing to the process $e^+e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$ (1-4) when the $Z(Z')$ vector bosons are produced on mass-shell and (5, 6) contributions from anomalous neutrino electromagnetic couplings with initial-state radiation.




FIG. 2: Top panel: The relative correction $\frac{\delta g_V^e}{(g_V^e)_{SM}}$ as a function of (g'_1, θ_{B-L}) . Bottom panel: The relative correction $\frac{\delta g_A^e}{(g_A^e)_{SM}}$ as a function of (g'_1, θ_{B-L}) .




FIG. 3: Top panel: Z' width as a function of $M_{Z'}$ for fixed values of g'_1 . Bottom panel: Z' width as a function of g'_1 for fixed values of $M_{Z'}$.




FIG. 4: The curves show the shape for $e^+e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$ as a function of center-of-mass energy and different values of the μ_{ν_τ} magnetic moment.




FIG. 5: Dependence of the sensitivity limits at 95% C. L. for the anomalous magnetic moment for three different values of $M_{Z'}$ and g'_1 in the process $e^+e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$.




FIG. 6: The surface show the shape for the cross section of the process $e^+e^- \rightarrow \nu_\tau \bar{\nu}_\tau \gamma$ as a function of the μ_{ν_τ} magnetic moment and the d_{ν_τ} electric dipole moment.