Proceedings of the Fifth Annual LHCP ATL-PHYS-PROC-2017-064 June 9, 2017

# NEW RESULTS ON COLLECTIVITY WITH ATLAS

KRZYSZTOF W. WOŹNIAK<sup>1</sup>

On behalf of the ATLAS Experiment, Institute of Nuclear Physics, Polish Academy of Sciences Krakow, Poland

#### ABSTRACT

The collective phenomena are observed not only in heavy ion collisions, but also in the proton-nucleus and in high-multiplicity pp collisions. The latest results from this area obtained in ATLAS are presented. In p+Pb collisions the emission source of particles is measured using the HBT method. The analysis of p+Pbdata collected in 2016 provides information on the elliptic flow of charged hadrons and muons. Low multiplicity events from pp, p+Pb and peripheral Pb+Pb collisions are studied with the cumulant methods. A deeper understanding of Pb+Pb collisions is provided by the analysis of longitudinal fluctuations of the collective flow parameters.

#### PRESENTED AT

The Fifth Annual Conference on Large Hadron Collider Physics Shanghai Jiao Tong University, Shanghai, China May 15-20, 2017

 $<sup>^1 \</sup>rm Work$  supported in part by the National Science Centre, Poland, grant 2015/18/M/ST2/00087 and by PL-Grid Infrastructure.




Figure 1: The ratio of exponential radii  $R_{\text{out}}/R_{\text{side}}$  (left) as a function of the pair transverse momentum,  $k_{\text{T}}$ , in four centrality bins [2] and (right) as a function of the angular distance from the second-order event plane  $2(\phi_k - \Psi_2)$  in five bins of elliptic flow vector magnitude  $|\vec{q}_2|$  [3]. The points for different intervals of  $|\vec{q}_2|$  are offset for visibility, as indicated in the legend.

### 1 Introduction

Strong collective effects were first found in heavy ion collisions, where they indicate a creation of the Quark-Gluon Plasma. This phase of matter manifests azimuthal correlations among produced particles, referred to as collective particle flow. However, similar effects are observed in proton-nucleus and even in high-multiplicity proton-proton collisions. The measurements of Pb+Pb, p+Pb and pp collisions at energies available from the Large Hadron Collider (LHC) and performed using the ATLAS detector [1] allow to study collective phenomena in detail. The obtained results are important for understanding the particle production at LHC energies.

### 2 Results

One of the questions, which can be answered by analysing correlations among produced particles is the size of their emission source. The Bose-Einstein correlations of identical bosons, as a function of the difference of their momenta,  $\mathbf{q} = \mathbf{p}^{\mathbf{a}} - \mathbf{p}^{\mathbf{b}}$ , can be parameterized as:

$$C_{\rm BE}(\mathbf{q}) = 1 + e^{\|\mathbf{R}\mathbf{q}\|},\tag{1}$$

where the diagonal elements of the matrix **R** are  $R_{\text{out}}$ ,  $R_{\text{side}}$  and  $R_{\text{long}}$ . These parameters were studied for p+Pb collisions at  $\sqrt{s_{\text{NN}}} = 5.02$  TeV [2] as a function of  $k_{\text{T}}$  (where  $\mathbf{k} = 0.5(\mathbf{p^a} - \mathbf{p^b})$ ) and the number of nucleons participating in collisions,  $N_{\text{part}}$ . Especially interesting is the ratio  $R_{\text{out}}/R_{\text{side}}$  which represents asymmetry of the source. In Figure 1 (left) one can see that this ratio depends on  $k_{\text{T}}$ , which indicates radial expansion of the source of particles [2]. This ratio depends also on the orientation of the correlated pairs (represented by  $\phi_k$ , azimuthal angle of  $k_{\text{T}}$ ) with respect to the event plane,  $\Psi_2$ , as shown in Figure 1 (right). The  $R_{\text{out}}/R_{\text{side}}$  ratio is larger out of plane than in plane ( $\phi_k \approx \Psi_2$ ) as  $R_{\text{out}}$  is enhanced out of plane and  $R_{\text{side}}$  does not change much with  $\phi_k - \Psi_2$  [3].

In the studies of azimuthal correlations among charged particles the Fourier decomposition is used:

$$\frac{\mathrm{d}N_{\mathrm{ch}}}{\mathrm{d}\phi} \sim 1 + 2\sum_{n=1}^{\infty} v_n(p_{\mathrm{T}}, \eta) \cos(n(\phi - \Psi_n)).$$
<sup>(2)</sup>

The flow harmonics  $v_n$  can be obtained also from two-particle correlations. In pp and p+Pb collisions these correlations contain large contributions from other sources (non-flow), which have to be subtracted. ATLAS




Figure 2: The  $v_2$  values, as a function of  $N_{\rm ch}^{\rm rec}$ , obtained (left) in p+Pb collisions at 8.16 TeV and 5.02 TeV and (right) in p+Pb collisions at 8.16 TeV from h-h correlations (circles) and h- $\mu$  correlations (squares) [5].




Figure 3: Multiplicity dependence of (left)  $v_2\{2, |\Delta \eta| > 2\}$ ,  $v_2\{4\}$ ,  $v_2\{6\}$  and  $v_2\{8\}$  for p+Pb collisions at 5.02 TeV and low-multiplicity Pb+Pb collisions at 2.76 TeV and (right)  $v_2\{2, |\Delta \eta| > 2\}$  for pp collisions at 5.02 TeV and 13 TeV, p+Pb collisions at 5.02 TeV and low-multiplicity Pb+Pb collisions at 2.76 TeV [6].

applies a *template method* [4] to remove them. In Figure 2 the elliptic flow  $v_2$  in p+Pb collisions at 5.02 TeV and 8.16 TeV for all hadrons (h-h) and for hadron-muon  $(h-\mu)$  correlations (at 8.16 TeV) is shown as a function of charged-particle multiplicity,  $N_{\rm ch}^{\rm rec}$  [5]. There is no dependence of  $v_2$  on the energy of collisions and a very weak increase with multiplicity. The values of  $v_2^{h-\mu}$  reach only about  $0.6 v_2^{h-h}$ .

The non-flow effects are most pronounced in low-multiplicity events as they usually involve relatively small number of particles in limited kinematical range (resonance decays, jets) and are thus suppressed in multi-particle correlations. Such correlation can be used to calculate cumulants,  $c_n\{2k\}$  [6], closely related to flow harmonics:

$$v_n\{2\} = \sqrt{c_n\{2\}}, \qquad v_n\{4\} = \sqrt[4]{-c_n\{4\}}, \qquad v_n\{6\} = \sqrt[6]{-c_n\{6\}/4}, \qquad v_n\{8\} = \sqrt[8]{-c_n\{8\}/33}.$$
(3)

In the case of  $c_n\{2\}$  a separation of particles in pseudorapidity  $(|\Delta \eta| > 2)$  is required in calculations of  $v_n\{2, |\Delta \eta| > 2\}$  to suppress short-range correlations. In Figure 3 elliptic flow  $v_2$  values obtained for p+Pb, Pb+Pb and pp collisions using different multi-particle cumulants are compared [6]. While the  $v_2\{2k\}$  values are similar (for k=2, 3, 4), those for k = 1 with  $|\Delta \eta| > 2$  requirement are larger. Comparison of




Figure 4: The  $c_2$ {4} cumulants in p+Pb collisions at 5.02 TeV calculated for charged particles with (left)  $0.3 < p_T < 3$  GeV and (right)  $0.5 < p_T < 5$  GeV compared across the three cumulant methods [7].




Figure 5: The  $v_2$ {4} values calculated for charged particles with  $0.3 < p_T < 3$  GeV using the three-subevent method in pp collisions at (left) 5.02 TeV, (middle) 13 TeV and (right) in p+Pb collisions at 5.02 TeV. They are compared to  $v_2$  obtained from a two-particle correlation analysis using a template fit procedure (solid circles) or peripheral subtraction (solid line) to remove non-flow effects [7].




Figure 6: The number of sources inferred from  $v_2\{2\}$  and  $v_2\{4\}$  using Eq. 4 in pp and p+Pb collisions at 13 TeV and 5.02 TeV, respectively, as a function of charged-particle multiplicity,  $N_{\rm ch}^{\rm Sel}$ , obtained by selecting charged particles with (left)  $0.3 < p_{\rm T} < 3$  GeV and (right)  $0.5 < p_{\rm T} < 5$  GeV [7].




Figure 7: (left) The  $v_n$  values as a function of transverse momentum  $p_{\rm T}$ , in (20–30)% centrality interval of Pb+Pb collisions at 5.02 TeV [8]. (right) Nuclear modification factor  $R_{\rm AA}$  as a function of  $p_{\rm T}$  for charged hadrons measured in Pb+Pb collisions at 5.02 TeV for five centrality intervals [9].




Figure 8: (left) The ratio of decorrelation parameter  $F_{n;1}^r$  in Pb+Pb collisions at 2.76 TeV and 5.02 TeV as a function of centrality represented by  $N_{\text{part}}$ . (right) The event-plane twist component  $F_{n;2}^{\text{twi}}$  and asymmetry component  $F_{n;2}^{\text{asy}}$  as a function of  $N_{\text{part}}$  for n=2 and n=3 for Pb+Pb collisions at 5.02 TeV [10].

 $v_2\{2, |\Delta \eta| > 2\}$  for different systems shows an increase with the size of colliding projectiles for events with the same number of produced particles. The  $v_2\{2k\}$  harmonics do not change with the multiplicity and the energy for pp collisions while are increasing with multiplicity for p+Pb and Pb+Pb collisions.

Further suppression of short range correlations can be achieved if in the calculations of correlations, used to obtain cumulants, particles from different ranges of pseudorapidity (i.e. *subevents*) are used [7]. For  $c_2\{4\}$  negative values are expected, as otherwise  $v_2\{4\}$  can not be calculated using Eq. 3. In Figure 4 one can see that positive  $c_2\{4\}$  are obtained at low multiplicities, but are reduced in two- and especially three-subevent cumulant methods. The  $v_2\{4\}$  values from three-subevent method are lower than  $v_2\{2\}$  from peripheral subtraction or template fit method (Figure 5). This difference can be interpreted as a result of event-by-event flow fluctuations, which are closely related to the effective number of sources,  $N_s$ , for particle production:

$$\frac{v_2\{4\}}{v_2\{2\}} = \left(\frac{4}{3+N_s}\right)^{1/4}.$$
(4)

The number of sources shown in Figure 6 is similar for pp and p+Pb collisions and for the latter it increases from 10 to 20 in the full available event-multiplicity range.

For events with large multiplicities, such as measured in Pb+Pb collisions at 5.02 TeV, standard methods of calculation of flow harmonics are sufficiently robust against non-flow effects. In Figure 7 (left)  $v_n$ , for n = 2 - 7, as a function of  $p_T$  in 20-30% centrality interval are shown [8]. Flow harmonics up to  $v_7$  are non-zero. The  $v_2$  is 0.05 even for  $p_T > 20$  GeV. Relatively large  $v_2$  for particles with high transverse momenta means that also very energetic partons are interacting in the QGP. This observation is consistent with the measured suppression of high- $p_T$  charged hadrons as quantified by the nuclear modification factor,  $R_{AA}$  [9]. In Figure 7 (right) one can see that  $R_{AA} < 1$  in the same centrality and  $p_T$  range.

A deeper understanding of flow phenomena may provide study of the dependence of flow fluctuations on position in pseudorapidity using correlators  $r_{n|n;k}$  and  $R_{n,n|n,n}$  (see Ref. [10] for definitions). Assuming that their dependence on pseudorapidity is linear it can be parameterized as:

$$r_{n|n;k} \approx 1 - 4 k F_{n;k}^{\mathrm{r}} \eta$$
 and  $R_{n,n|n,n} \approx 1 - 4 F_{n;n}^{\mathrm{twi}} \eta$ , (5)

where  $F_{n;k}^{\rm r} = F_{n;n}^{\rm asy} + F_{n;n}^{\rm twi}$ ,  $F_{n;n}^{\rm asy}$  and  $F_{n;n}^{\rm twi}$  are decorrelation parameters connected with magnitude (asymmetry) and twist fluctuations, respectively. In Pb+Pb collisions the decorrelation parameters  $F_{n;k}^{\rm r}$  are similar for all centralities but depend on the energy of the collision and are 10-16% larger at 2.76 TeV than at 5.02 TeV (see Figure 8 (left)). On the other hand the magnitude and twist decorrelation parameters, shown in Figure 8 (right), are approximately constant for  $N_{\rm part} > 100$ . In the whole multiplicity range  $F_{n;2}^{\rm asy} \approx F_{n;2}^{\rm twi}$  for the same n [10].

## 3 Conclusions

Studies of correlations among particles produced in different types of collisions available at LHC provide valuable information on properties of their source. In p+Pb collisions the volume from which particles are emitted has an elongated shape and undergoes a radial expansion. The flow of muons originating from b or c quarks is much smaller than that of charged hadrons. Results on flow harmonics obtained using cumulant methods clearly show that the non-flow contributions are very important in low-multiplicity events and need to be properly subtracted. In new detailed studies of Pb+Pb collisions non-zero flow harmonics up to  $v_7$  were measured. Analysis of longitudinal fluctuations of flow harmonics reveals decorrelation effects which are stronger at lower collision energy, but similar when decomposed into magnitude and twist contributions.

#### References

- [1] ATLAS Collaboration, JINST 3 (2008) S08003.
- [2] ATLAS Collaboration, arXiv:1704.01621 [hep-ex], submitted to Phys. Rev. C.
- [3] ATLAS Collaboration, ATLAS-CONF-2017-008, https://cds.cern.ch/record/2244818.
- [4] ATLAS Collaboration, Phys. Rev. Lett. 116 (2016) 172301.
- [5] ATLAS Collaboration, ATLAS-CONF-2017-006, https://cds.cern.ch/record/2244808.
- [6] ATLAS Collaboration, arXiv:1705.04176 [hep-ex], submitted to EPJC.
- [7] ATLAS Collaboration, ATLAS-CONF-2017-002, https://cds.cern.ch/record/2244764.
- [8] ATLAS Collaboration, ATLAS-CONF-2016-105, https://cds.cern.ch/record/2220372.
- [9] ATLAS Collaboration, ATLAS-CONF-2017-012, https://cds.cern.ch/record/2244824.
- [10] ATLAS Collaboration, ATLAS-CONF-2017-003, https://cds.cern.ch/record/2244796.