

ASYMPTOTICS OF QUANTUM $6j$ SYMBOLS

QINGTAO CHEN AND JUN MURAKAMI

ABSTRACT. The asymptotics of the quantum $6j$ symbols corresponding to a hyperbolic tetrahedron is investigated and the first two leading terms are determined for the case that the tetrahedron has an ideal or an ultra-ideal vertex. These terms are given by the volume and the determinant of the Gram matrix of the tetrahedron.

1. INTRODUCTION

The quantum $6j$ symbol was introduced by A. Kirillov and N. Reshetikhin from the representation theory of the quantum group $\mathcal{U}_q(sl_2)$ and was used to reconstruct quantum sl_2 invariants of knots and knotted graphs in [9]. Then V. Turaev and O. Viro constructed a 3-manifold invariant from its tetrahedral decomposition by assigning the quantum $6j$ symbol to each tetrahedron. The quantum $6j$ symbol is a quantum analogue of the classical $6j$ symbol introduced by Wigner, which is a highly symmetric version of Racah's W -coefficients expressing the recoupling coefficient of 3 angular momenta. The relation between the asymptotics of the classical $6j$ symbols and the volume of an Euclidean tetrahedron was conjectured by G. Ponzano and T. Regge [15], and was proved by J. Roberts [17].

Here we give the asymptotics of the quantum $6j$ symbols corresponding to a hyperbolic tetrahedron with at least one ideal or ultra-ideal vertex. Let T be a hyperbolic tetrahedron, v be one of its vertex, a, b, c be the edges of T touching to v and $\theta_a, \theta_b, \theta_c$ be the dihedral angles at a, b, c . Then $\theta_a + \theta_b + \theta_c > \pi$ if v is a usual vertex, $\theta_a + \theta_b + \theta_c = \pi$ if v is an ideal vertex and $\theta_a + \theta_b + \theta_c < \pi$ if v is a ultra-ideal vertex. Let $r \geq 3$ be a positive odd integer and $\xi_r = \exp(2\pi\sqrt{-1}/r)$. A triplet (a, b, c) of half integers is called *admissible* if it satisfies the Clebsch-Gordan condition, i.e. $|a - b| \leq c \leq a + b$ and $a + b + c \in \mathbf{Z}$.

Theorem 1. *Let T be a hyperbolic tetrahedron at least one of whose vertices are ideal or ultra-ideal. Let $\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f$ be dihedral angles of T where a, \dots, f are edges of T given as in Figure 1 and let a_r, b_r, \dots, f_r be sequences of half integers for odd $r \geq 3$ satisfying*

$$(1) \quad \lim_{r \rightarrow \infty} \frac{a_r}{r} = \frac{1}{4} - \frac{\theta_a}{4\pi}, \quad \lim_{r \rightarrow \infty} \frac{b_r}{r} = \frac{1}{4} - \frac{\theta_b}{4\pi}, \quad \dots, \quad \lim_{r \rightarrow \infty} \frac{f_r}{r} = \frac{1}{4} - \frac{\theta_f}{4\pi},$$

where the triplets $(a_r, b_r, e_r), (a_r, c_r, f_r), (b_r, d_r, f_r)$ and (c_r, d_r, e_r) are all admissible. We also assume that $a_r + b_r + e_r \geq \frac{r-3}{2}$ for the three edges a, b, e attached to the same ideal or ultra-ideal vertex. Then

$$(2) \quad \lim_{r \rightarrow \infty} \frac{2\pi}{r} \log \left| \begin{Bmatrix} a_r & b_r & e_r \\ d_r & c_r & f_r \end{Bmatrix}_{q=\xi_r^2}^{RW} \right| = \text{Vol}(T),$$

where $\text{Vol}(T)$ is the hyperbolic volume of T .

By specifying the sequences a_r, b_r, \dots, f_r more precisely, we also have the following refined asymptotics.

Theorem 2. *Let T be a hyperbolic tetrahedron with at least one of whose vertices are ideal or ultra-ideal. Let $\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f$ be dihedral angles of T and $\eta_a = \frac{\pi - \theta_a}{2\pi}, \dots, \eta_f = \frac{\pi - \theta_f}{2\pi}$. Assume that η_a, \dots, η_f are all rational numbers whose denominators and numerators are odd integers. Let r be an*

The first author was supported by the National Center of Competence in Research SwissMAP of the Swiss National Science Foundation, and the second author was supported in part by KAKENHI (25287014, 16H06336, 17K18728).




FIGURE 1. Edges of a tetrahedron and a truncated tetrahedron

odd integer which is a common multiple of the denominators of η_a, \dots, η_f . Then $a_r = \frac{r}{2}\eta_a - \frac{1}{2}, \dots, f_r = \frac{r}{2}\eta_f - \frac{1}{2}$ are integers and

$$(3) \quad \left| \begin{Bmatrix} a_r & b_r & e_r \\ d_r & c_r & f_r \end{Bmatrix}_{q=\xi_r^2}^{RW} \right| \underset{r \rightarrow \infty}{\sim} \frac{\sqrt{2}\pi}{r^{3/2} \sqrt[4]{-\det G}} e^{\frac{r}{2\pi} \text{Vol}(T)},$$

where G is the Gram matrix of T given by

$$G = \begin{pmatrix} 1 & -\cos \theta_a & -\cos \theta_b & -\cos \theta_f \\ -\cos \theta_a & 1 & -\cos \theta_e & -\cos \theta_c \\ -\cos \theta_b & -\cos \theta_e & 1 & -\cos \theta_d \\ -\cos \theta_f & -\cos \theta_c & -\cos \theta_d & 1 \end{pmatrix}.$$

Remark 1. For the classical $6j$ symbol, such asymptotics were conjectured by Wigner, Ponzano and Regge, and proved by J. Roberts [17]. For non-squared q case, the asymptotics of the quantum $6j$ symbol corresponding to a spherical tetrahedron was given by Taylor and Woodward in [18]. Here we use squared $q = \xi_r^2$, then the quantum $6j$ symbol corresponding to a hyperbolic tetrahedron grows exponentially and the leading term is given by the volume of the tetrahedron. This point is quite different from the asymptotics given in the previous works, but our asymptotics also relate to the determinant of the Gram matrix as in the previous works.

The paper is organized as follows. In Section 2, we explain the quantum $6j$ symbol, and prove Theorem 1 in Section 3 that the leading term of the asymptotics of the quantum $6j$ symbol is given by the hyperbolic volume of the corresponding tetrahedron. We prove Theorem 2 in Section 4 by showing that the second leading term of the asymptotics of the quantum $6j$ symbols is given by the determinant of the corresponding Gram matrix. Some part of the proof of Theorem 3 is given in Appendix. In Section 4, a conjecture about the symmetry of the asymptotics of the quantum $6j$ symbols is proposed.

2. QUANTUM $6j$ SYMBOLS

The quantum $6j$ symbol was introduced by Kirillov and Reshetikhin [9] to describe the structure of the tensor representations of the quantum group $\mathcal{U}_q(sl_2)$. Let q be an indeterminate. Finite-dimensional irreducible representations of $\mathcal{U}_q(sl_2)$ are parametrized by non-negative half integers, which is called the spin. For a non-negative half integer a , let V_a denote the corresponding finite-dimensional irreducible representation. It is known that V_c is contained in $V_a \otimes V_b$ if a, b, c satisfied the Clebsch-Gordan condition, which means that $|a - b| \leq c \leq a + b$ and $a + b + c$ is an integer. Let i_{ab}^c denote the inclusion map from V_c to $V_a \otimes V_b$. For six non-negative half integers a, b, c, d, e, f , the quantum $6j$ symbol is defined by the following relation between the following two inclusions of an irreducible representation to a tensor of three irreducible representations $V_c \rightarrow V_d \otimes V_b \otimes V_a$, one through $V_d \otimes V_e$ and another one through $V_f \otimes V_a$.

$$\left(V_c \xrightarrow{i_{de}^c} V_d \otimes V_e \xrightarrow{id \otimes i_{ba}^e} V_d \otimes V_b \otimes V_a \right) =$$

$$\sum_f \left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\}_q \left(V_c \xrightarrow{i_{fa}^c} V_f \otimes V_a \xrightarrow{i_{db}^f \otimes id} V_d \otimes V_b \otimes V_a \right).$$

Here we assume that the triplets (a, b, e) , (a, c, f) , (b, d, f) and (c, d, e) satisfy the Clebsch-Gordan condition. The Racah-Wigner version of the quantum $6j$ symbol is given by

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\}_q^{RW} = \frac{1}{\sqrt{-1}^{2(c+d+2e-a-b)} \sqrt{[2e+1][2f+1]}} \left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\}_q,$$

where $[n] = \frac{q^{n/2} - q^{-n/2}}{q^{1/2} - q^{-1/2}}$. Let $[n]! = \prod_{j=1}^n [j]$. Then this version has the following symmetric formula

$$(4) \quad \left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\}_q^{RW} = \Delta_r(a, b, e) \Delta_r(a, c, f) \Delta_r(b, d, f) \Delta_r(c, d, e) \sum_{z=m}^M \frac{(-1)^z [z+1]!}{\prod_{j=1}^4 [z - S_j]! \prod_{k=1}^3 [T_k - z]!},$$

where

$$(5) \quad \begin{aligned} S_1 &= a + b + e, & S_2 &= a + c + f, & S_3 &= b + d + f, & S_4 &= c + d + e, \\ T_1 &= a + b + c + d, & T_2 &= a + d + e + f, & T_3 &= b + c + e + f, \\ m &= \max(S_1, S_2, S_3, S_4), & M &= \min(T_1, T_2, T_3), \\ \Delta_r(u, v, w) &= \left(\frac{[u+v-w]! [v+w-u]! [w+u-v]!}{[u+v+w+1]!} \right)^{1/2}. \end{aligned}$$

For quantum invariants of knots, its relation to the hyperbolic volume of the knot complement was found by R. Kashaev [8], and reformulated using the colored Jones polynomial in [11]. This relation is called the volume conjecture and proved for a few knots and links. The volume conjecture suggests that there must be a relation between the quantum $6j$ symbol and the volume of a hyperbolic tetrahedron. Using this idea, a volume formula for hyperbolic and spherical tetrahedra was obtained by the second author and M. Yano in [12], and the asymptotics of the quantum $6j$ symbol was studied by F. Costantino [5].

Let

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\}_q^{RW}$$

be the Racah-Wigner version of the quantum $6j$ symbol which is defined for six non-negative half integers a, b, \dots, f with the quantum parameter q . A triplet (a, b, c) of half integers is called *admissible* if it satisfies the Clebsch-Gordan condition, i.e. $|a - b| \leq c \leq a + b$ and $a + b + c \in \mathbf{Z}$.

Theorem 3 (F. Costantino [5]). *Let T be a hyperbolic truncated tetrahedron with dihedral angles $\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f$ at edges a, \dots, f in Figure 1. Let a_r, b_r, \dots, f_r be sequences of non-negative half integers satisfying*

$$2\pi \left| \lim_{r \rightarrow \infty} \frac{a_r}{r} - \frac{1}{2} \right| = \theta_a, \quad 2\pi \left| \lim_{r \rightarrow \infty} \frac{b_r}{r} - \frac{1}{2} \right| = \theta_b, \quad \dots, \quad 2\pi \left| \lim_{r \rightarrow \infty} \frac{f_r}{r} - \frac{1}{2} \right| = \theta_f,$$

and the triplets (a_r, b_r, e_r) , (a_r, c_r, f_r) , (b_r, d_r, f_r) and (c_r, d_r, e_r) are all admissible. Then

$$\lim_{r \rightarrow \infty} \frac{2\pi}{r} \log \left| \text{ev}_r \left\{ \begin{matrix} a_r & b_r & e_r \\ d_r & c_r & f_r \end{matrix} \right\}_{\xi_r}^{RW} \right| = 2 \text{Vol}(T),$$

where $\xi_r = \exp(2\pi\sqrt{-1}/r)$ and $\text{Vol}(T)$ is the hyperbolic volume of T .

Here the operator ev_r removes the pole of the quantum $6j$ symbol at $q = \xi_r$. In this paper, we put $q = \xi_r^2$ instead of ξ_r , then the quantum $6j$ symbol doesn't have a pole at $q = \xi_r^2$ and its actual asymptotics is given by the volume and the determinant of the Gram matrix of the corresponding tetrahedron.

3. RELATION TO THE HYPERBOLIC VOLUME

In this section, we prove Theorem 1 giving a relation between the Racah-Wigner version of the quantum $6j$ symbols and the hyperbolic volume of the corresponding hyperbolic tetrahedra.

Proof of Theorem 1. The idea of proof is similar to the proof of the volume conjecture for figure-eight knot, for example, see [10]. Such an idea is also used in [5] for the quantum $6j$ symbols with $q = \xi_r$, where he used ev_r map. Here we follow the argument in [5] without ev_r map. At the end of proof, the limiting value is compared with the volume formula in [20] based on [12].

From the assumption, $m = \max(S_{r,1}, \dots, S_{r,4}) \geq \frac{r-3}{2}$, $M = \min(T_{r,1}, T_{r,2}, T_{r,3}) < \frac{3r}{4}$ and $\frac{r-1}{2} \leq m+1 \leq z+1 \leq r-1$. For such z , $(-1)^z [z+1]!$ has the same sign since $[k+1] < 0$ for $m+1 \leq k \leq M$. Therefore the terms in the sum of (4) are all real numbers with the same sign, and the limit in (2) is determined by the term having the largest absolute value. Indeed, let

$$(6) \quad \alpha_r(z) = \frac{(-1)^z [z+1]!}{\prod_{j=1}^4 [z - S_{r,j}]! \prod_{k=1}^3 [T_{r,k} - z]!} \Big|_{q=\xi_r^2},$$

where $S_{r,j}, T_{r,j}$ are defined by using a_r, b_r, \dots, f_r as in (5),

$$(7) \quad m_r = \max_{j=1}^4 S_{r,j}, \quad M_r = \min_{k=1}^3 T_{r,k}$$

and let $|\alpha_r(z_0)|$ be the maximum of $\{|\alpha_r(z)| \mid m_r \leq z \leq M_r\}$. Then

$$\lim_{r \rightarrow \infty} \frac{1}{r} \log |\alpha_r(z_0)| < \lim_{r \rightarrow \infty} \frac{1}{r} \log \left| \sum_{z=m}^M \alpha_r(z) \right| < \lim_{r \rightarrow \infty} \frac{1}{r} \log |r \alpha_r(z_0)|.$$

Since $\lim_{r \rightarrow \infty} \frac{1}{r} \log r = 0$, we get $\lim_{r \rightarrow \infty} \frac{1}{r} \log \left| \sum_{z=m}^M \alpha_r(z) \right| = \lim_{r \rightarrow \infty} \frac{1}{r} \log |\alpha_r(z_0)|$.

Now compare

$$\lim_{r \rightarrow \infty} \frac{2\pi}{r} \log |\Delta_r(a_r, b_r, e_r) \Delta_r(a_r, c_r, f_r) \Delta_r(b_r, d_r, f_r) \Delta_r(c_r, d_r, e_r) \alpha_r(z_0)|$$

with the hyperbolic volume of T . This part is the same as the proof of Theorem 1 in [5], which concerns the quantum $6j$ symbol with $q = \xi_r$. Here we assume r to be odd and q to be ξ_r^2 instead of ξ_r , so the proof in [5] works well without applying the operator ev_r to remove the factor $\xi_r^{r/2} - \xi_r^{-r/2}$ in $[S_{r,j} + 1]!$ and $[z + 1]!$. Here we may have factors $\xi_r^{(r \pm 1)/2} - \xi_r^{-(r \pm 1)/2}$ in $[S_{r,j} + 1]!$ and $[z + 1]!$ but we never have the factor $\xi_r^{r/2} - \xi_r^{-r/2}$ since $q = \xi_r^2$ and r is odd. \square

Theorem 1 can suggest the following a bit more general conjecture which is also inspired by the new volume conjecture proposed by the first author and T. Yang [4].

Conjecture 1. Let T be a hyperbolic tetrahedron and let $\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f$ are dihedral angles of T and a_r, b_r, \dots, f_r be sequences of admissible half integers for odd $r \geq 3$ satisfying (1) as in Theorem 1. Then

$$(8) \quad \lim_{r \rightarrow \infty} \frac{2\pi}{r} \log \left| \left\{ \begin{matrix} a_r & b_r & e_r \\ d_r & c_r & f_r \end{matrix} \right\}_{q=\xi_r^2}^{RW} \right| = \text{Vol}(T).$$

4. RELATION TO THE GRAM MATRIX

In this section, we investigate a bit more about the asymptotics of $\alpha_r(z)$ and prove Theorem 2 by using the method to prove the volume conjecture for 5_2 knot given in [13]. Here we use notations $S_{r,j}, T_{r,k}, m_r, M_r$ introduced in (6).

First of all, we express $a_r(z)$ in terms of the dilogarithm function. Let

$$(q)_n = \prod_{k=1}^n (1 - q^k).$$

Then

$$[n]! = q^{\frac{-n(n+1)}{4}} \frac{(q)_n}{2^n (-i)^n \sin^n \frac{2\pi}{r}}$$

and $\alpha_r(z)$, $\Delta_r(a_r, b_r, e_r)$, \dots , $\Delta_r(c_r, d_r, e_r)$ are reformulated as

$$(9) \quad \alpha_r(z) = \frac{\xi_r^{2(d_1+d_2)} (-1)^{z+1} (\xi_r^2)_{z+1}}{2i \sin \frac{2\pi}{r} \prod_{j=1}^4 (\xi_r^2)_{z-S_j} \prod_{k=1}^3 (\xi_r^2)_{T_k-z}},$$

where

$$\begin{aligned} d_1 &= - \left(2a_r + 2b_r + 2c_r + 2d_r + 2e_r + 2f_r + \frac{1}{2} \right) z + \frac{3z^2}{2}, \\ d_2 &= a_r^2 + a_r b_r + b_r^2 + a_r c_r + b_r c_r + c_r^2 + a_r d_r + b_r d_r + c_r d_r + d_r^2 + a_r e_r + \\ &\quad b_r e_r + c_r e_r + d_r e_r + e_r^2 + a_r f_r + b_r f_r + c_r f_r + d_r f_r + e_r f_r + f_r^2 - \frac{1}{2}, \end{aligned}$$

and

$$\begin{aligned} \Delta_r(a_r, b_r, e_r) &= \left(\xi_r^{2d_3(a_r, b_r, e_r)} \frac{-2i \sin \frac{2\pi}{r} (\xi_r^2)_{a_r+b_r-e_r} (\xi_r^2)_{b_r+e_r-a_r} (\xi_r^2)_{e_r+a_r-b_r}}{(\xi_r^2)_{a_r+b_r+e_r+1}} \right)^{1/2}, \\ &\quad \dots, \\ \Delta_r(c_r, d_r, e_r) &= \left(\xi_r^{2d_3(c_r, d_r, e_r)} \frac{-2i \sin \frac{2\pi}{r} (\xi_r^2)_{c_r+d_r-e_r} (\xi_r^2)_{d_r+e_r-c_r} (\xi_r^2)_{e_r+c_r-d_r}}{(\xi_r^2)_{c_r+d_r+e_r+1}} \right)^{1/2} \end{aligned}$$

where

$$d_3(a_r, b_r, e_r) = -\frac{1}{2} (a_r^2 + b_r^2 + e_r^2 - 2a_r b_r - 2a_r e_r - 2b_r e_r - a_r - b_r - e_r - 1).$$

Let φ_r is the function introduced in [7] for $0 \leq t \leq 1$ by the following integral.

$$\varphi_r(t) = \int_{-\infty}^{\infty} \frac{e^{(2t-1)x}}{4x \sinh x \sinh(2x/r)} dx,$$

where the singularity at $x = 0$ is bypassed by a small half circle above the real line.

Lemma 1. *The function φ_r satisfies*

$$(10) \quad (1 - e^{2\pi i a}) = \exp \left(\varphi_r \left(a - \frac{1}{r} \right) - \varphi_r \left(a + \frac{1}{r} \right) \right) \quad \left(\frac{1}{r} \leq a \leq \frac{r-1}{r} \right),$$

$$(11) \quad (\xi_r^2)_n = \exp \left(\varphi_r \left(\frac{1}{r} \right) - \varphi_r \left(\frac{2n+1}{r} \right) \right) \quad \left(0 \leq n \leq \frac{r-1}{2} \right),$$

$$(12) \quad (\xi_r^2)_n = \exp \left(\varphi_r \left(\frac{1}{r} \right) - \varphi_r \left(\frac{2n+1}{r} - 1 \right) + \log 2 \right) \quad \left(\frac{r-1}{2} < n < \frac{2r-1}{2} \right),$$

$$(13) \quad \operatorname{Im} \varphi_r(t) = -\pi i \frac{6r^2 t^2 - 6r^2 t + r^2 - 2}{24r}.$$

Proof. We have

$$\begin{aligned} \varphi_r \left(a - \frac{1}{r} \right) - \varphi_r \left(a + \frac{1}{r} \right) &= \int_{-\infty}^{\infty} \frac{e^{(2a-1-\frac{2}{r})x} - e^{(2a-1+\frac{2}{r})x}}{4x \sinh x \sinh(2x/r)} dx \\ &= - \int_{-\infty}^{\infty} \frac{e^{(2a-1)x}}{4x \sinh x} dx. \end{aligned}$$

The function $-\frac{e^{(2a-1)x}}{2x \sinh x}$ has poles at $x = k\pi i$ ($k \in \mathbb{N}$) in the upper half plane, and the corresponding residues are $-\frac{e^{2a\pi ki}}{2\pi ki}$. The sum of these residues is equal to $\frac{\log(1-e^{2a\pi i})}{2\pi i}$ and this implies (10).

The formula (11) is a product of (10) for $a = 2/r, 4/r, \dots, 2n/r$.

To prove (12), we decompose $(\xi_r^2)_n$ into a product of $(\xi_r^2)_{\frac{r-1}{2}}$ and $\prod_{k=1}^{n-\frac{r-1}{2}} (1 - \xi_r^{2k-1})$. Then $(\xi_r^2)_{\frac{r-1}{2}} = \exp(\varphi_r(1/r) - \varphi_r(1))$ and $\prod_{k=1}^{n-\frac{r-1}{2}} (1 - \xi_r^{2k-1}) = \exp(\varphi_r(0) - \varphi_r(\frac{2n+1}{r} - 1))$. Moreover, $\varphi_r(0) - \varphi_r(1)$ is given by $2\pi i$ times the sum of residues of $-\frac{1}{2x \sinh(2x/r)}$, which is $\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} = \log 2$. Hence we get (12).

The imaginary part of $\varphi_r(t)$ comes from the pole at $x = 0$. For the other part, $\varphi_r(t)$ is defined as an integral of a real function. We define the integral path to avoid $x = 0$ by taking a path in the upper half plane, so by taking a small upper half circle around $x = 0$, the contribution is given by $-\pi i$ times the residue of $\frac{e^{(2t-1)x}}{4x \sinh x \sinh(2x/r)}$ at $x = 0$, which is the right hand side of (13). \square

Now, we reformulate $\alpha_r(z)$ and $\Delta_r(a_r, b_r, e_r)$ as continuous functions by using (11) and (12).

$$(14) \quad \tilde{\alpha}_r(z) = i^r \xi_r^{2d_2} \exp \left(-\pi i z - 6\varphi_r\left(\frac{1}{r}\right) - \varphi_r\left(\frac{2z+3}{r} - 1\right) + \log 2 + \sum_{j=1}^4 \varphi_r\left(\frac{2z-2S_{r,j}+1}{r}\right) \right. \\ \left. + \sum_{j=1}^3 \varphi_r\left(\frac{2T_{r,j}-2z+1}{r}\right) + \frac{2\pi i}{r} (3z^2 - z - 4(a_r + b_r + \dots + f_r)z) \right),$$

and

$$(15) \quad \tilde{\Delta}_r(a_r, b_r, e_r) = \exp \left(\frac{1}{2} \left(2\varphi_r\left(\frac{1}{r}\right) + \varphi_r\left(\frac{2a_r+2b_r+2e_r+3}{r} - 1\right) - \log 2 - \varphi_r\left(\frac{2a_r+2b_r-2e_r+1}{r}\right) - \varphi_r\left(\frac{2b_r+2e_r-2a_r+1}{r}\right) - \varphi_r\left(\frac{2e_r+2a_r-2b_r+1}{r}\right) \right) \right).$$

Then $\tilde{\alpha}_r(z)$ is a positive real function for $m_r < z < M_r$,

$$\alpha_r(z) = \frac{(-1)^{\frac{r+1}{2}}}{2 \sin \frac{2\pi}{r}} \tilde{\alpha}_r(z)$$

for an integer z ($m_r \leq z \leq M_r$) and

$$\Delta_r(a_r, b_r, e_r) = \sqrt{-2i \sin \frac{2\pi}{r}} \xi_r^{d_3(a_r, b_r, e_r)} \tilde{\Delta}_r(a_r, b_r, e_r).$$

Next, we apply the Poisson summation formula to $\sum_{z=m_r}^{M_r} \tilde{\alpha}_r(z)$. Let ε be a small positive real number and $\psi(x)$ be a smooth function satisfying $0 \leq \psi(x) \leq 1$, $\psi(0) = 0$ for $x \in \mathbb{R} \setminus ((2m_r+3)/r + \varepsilon, (2M_r+3)/r - \varepsilon)$, and $\psi(x) = 1$ for $x \in [(2m_r+3)/r + 2\varepsilon, (2M_r+3)/r - 2\varepsilon]$. Note that $(2m_r+3)/r$ and $(2M_r+3)/r$ do not depend on r . Let

$$(16) \quad h_r(z) = \psi\left(\frac{2z+3}{r}\right) \tilde{\alpha}_r(z).$$

By the Poisson summation formula, we have

$$\sum_{m \in \mathbb{Z}} h_r(m) = \sum_{m \in \mathbb{Z}} \hat{h}_r(m)$$

where $\hat{h}_r(m) = \int_{-\infty}^{\infty} e^{-2\pi i m t} h_r(t) dt$. The function $h_r(t)$ is non-negative or non-positive depending on r , and satisfies the following.

Lemma 2. *There are positive constants c and δ' satisfying*

$$(17) \quad \frac{\left| \sum_{m \neq 1} \widehat{h}_r(m) \right|}{\left| \widehat{h}_r(0) \right|} < c e^{-r\delta'}.$$

We also have the following.

Lemma 3. *Let $A_r = \sum_{z=m_r}^{M_r} \widehat{\alpha}_r(z)$ and $B_r = \sum_{z=m_r}^{M_r} h_r(z)$, then there is positive constants c' and δ' not depending on r satisfying*

$$(18) \quad \frac{|A_r - B_r|}{|\widehat{h}_r(0)|} < c' e^{-\delta'}.$$

The proof of the above two lemmas are given in Appendix.

The above two lemmas shows that the main asymptotics of the quantum $6j$ symbol is determined by the asymptotics of $\widehat{h}_r(0)$ with respect to r . Let $\zeta = \frac{2z+3}{r}$, $\sigma_j = \frac{2S_{r,j}+3}{r}$ ($1 \leq j \leq 4$), and $\tau_k = \frac{2T_{r,k}+4}{r}$ ($1 \leq k \leq 3$). Note that σ_j, τ_k do not depend on r since they are determined by η_a, \dots, η_f . Let

$$(19) \quad g_r(\zeta, \eta) = \psi(\zeta) \exp \left(\frac{-\pi i r \zeta}{2} + \frac{3\pi i}{2} - 6\varphi_r\left(\frac{1}{r}\right) - \varphi_r(\zeta - 1) + \log 2 + \sum_{j=1}^4 \varphi_r(\zeta - \sigma_{r,j} + \frac{1}{r}) \right. \\ \left. + \sum_{j=1}^3 \varphi_r(\tau_{r,j} - \zeta) + 2\pi i r \left(\frac{3}{4} \zeta^2 - (\eta_a + \dots + \eta_f) \zeta + \frac{\zeta}{r} \right) \right),$$

where $\eta = (\eta_a, \eta_b, \dots, \eta_f)$. Then the function $h_r(z)$ is equal to $|g_r(\frac{2z+3}{r}, \eta)|$, the argument of $g(\zeta, \eta)$ does not depend on ζ and

$$\left| \widehat{h}_r(0) \right| = \left| \int_{-\infty}^{\infty} h_r(x) dx \right| = \frac{r}{2} \left| \int_{-\infty}^{\infty} g_r(\zeta, \eta) d\zeta \right|.$$

The parameters η_a, \dots, η_f do not depend on r , but the function $\varphi_r(x)$ depends on r . We investigate the asymptotics of

$$(20) \quad \left| r \sin \frac{2\pi}{r} \widetilde{\Delta}\left(\frac{r\eta_a-1}{2}, \frac{r\eta_b-1}{2}, \frac{r\eta_c-1}{2}\right) \dots \widetilde{\Delta}\left(\frac{r\eta_e-1}{2}, \frac{r\eta_d-1}{2}, \frac{r\eta_f-1}{2}\right) \int_{-\infty}^{\infty} g_r(\zeta, \eta) d\zeta \right|.$$

We expand $g_r(\zeta, \eta)$ by using the following relations.

Lemma 4 ([13, Appendix A]). *The asymptotics of φ_r is given as follows.*

$$(21) \quad \varphi_r\left(\frac{1}{r}\right) = \frac{r}{4\pi i} \left(\frac{\pi^2}{6} + \frac{2\pi i}{r} \log \frac{r}{2} - \frac{\pi^2}{r} + O\left(\frac{1}{r^2}\right) \right),$$

$$(22) \quad \varphi_r(t) = \frac{r}{4\pi i} \left(\text{Li}_2(e^{2\pi i t}) + O\left(\frac{1}{r^2}\right) \right).$$

We also have the following expansion for small α since $\frac{d}{dx} \text{Li}_2(x) = -\frac{\log(1-x)}{x}$.

$$(23) \quad \text{Li}_2(e^{2\pi i(p+\alpha)}) = \text{Li}_2(e^{2\pi i p}) - 2\pi i \log(1 - e^{2\pi i p}) \alpha + O(\alpha^2).$$

By reformulating g_r by using (21), (22) and (23), we get

$$\begin{aligned}
 g_r(\zeta, \eta) &= \frac{16\psi(\zeta)}{r^3} \exp\left(\frac{-\pi i r \zeta}{2} + \frac{3\pi i}{2} + \right. \\
 &\quad \left. \frac{r}{4\pi i} \left(-\pi^2 + \frac{6\pi^2}{r} - \text{Li}_2(e^{2\pi i \zeta}) + \sum_{j=1}^4 \text{Li}_2(e^{2\pi i(\zeta - \sigma_j)}) + \sum_{j=1}^3 \text{Li}_2(e^{2\pi i(\tau_j - \zeta)}) \right) + \right. \\
 (24) \quad &\quad \left. 2\pi i r \left(\frac{3}{4} \zeta^2 - (\eta_a + \dots + \eta_f) \zeta + \frac{\zeta}{r} \right) - \frac{1}{2} \sum_{j=1}^4 \log(1 - e^{2\pi i(\zeta - \sigma_j)}) + O\left(\frac{1}{r}\right) \right) \\
 &= \frac{16\psi(\zeta)}{r^3} \frac{-e^{2\pi i \zeta}}{\sqrt{\prod_{j=1}^4 (1 - \frac{e^{2\pi i \zeta}}{e^{2\pi i \sigma_j}})}} \exp\left(\frac{r}{4\pi i} \left(-\pi^2 - 2\pi^2 \zeta - \text{Li}_2(e^{2\pi i \zeta}) + \right. \right. \\
 &\quad \left. \left. \sum_{j=1}^4 \text{Li}_2\left(\frac{e^{2\pi i \zeta}}{e^{2\pi i \sigma_j}}\right) + \sum_{j=1}^3 \text{Li}_2\left(\frac{e^{2\pi i \tau_j}}{e^{2\pi i \zeta}}\right) - 6\pi^2 \zeta^2 + 8\pi^2 (\eta_a + \dots + \eta_f) \zeta \right) + O\left(\frac{1}{r}\right) \right).
 \end{aligned}$$

Moreover, let

$$(25) \quad \delta(\eta_1, \eta_2, \eta_3) = \frac{1}{8\pi i} \left(\text{Li}_2(e^{2\pi i(\eta_1 + \eta_2 + \eta_3)}) - \text{Li}_2\left(\frac{e^{2\pi i(\eta_1 + \eta_2)}}{e^{2\pi i \eta_3}}\right) - \text{Li}_2\left(\frac{e^{2\pi i(\eta_2 + \eta_3)}}{e^{2\pi i \eta_1}}\right) - \text{Li}_2\left(\frac{e^{2\pi i(\eta_3 + \eta_1)}}{e^{2\pi i \eta_2}}\right) \right)$$

for $0 \leq \eta_i \leq 1$ and $1 \leq \eta_1 + \eta_2 + \eta_3 \leq 2$. Then

$$(26) \quad \tilde{\Delta}_r\left(\frac{r\eta_1-1}{2}, \frac{r\eta_2-1}{2}, \frac{r\eta_3-1}{2}\right) = \frac{\sqrt{r}}{2} \exp\left(r\delta(\eta_1, \eta_2, \eta_3) + O\left(\frac{1}{r}\right)\right).$$

Now we apply the stationary phase method to the function $g_r(\zeta, \eta)$. The arguments of

$$\frac{16\psi(\zeta)}{r^3} \frac{-e^{2\pi i \zeta}}{\sqrt{\prod_{j=1}^4 (1 - \frac{e^{2\pi i \zeta}}{e^{2\pi i \sigma_j}})}}$$

does not depend on ζ since $\arg(1 - z) = -(\pi - \arg z)/2$ for $z \in \mathbb{C}$ satisfying $|z| = 1$. Similarly, the argument of

$$\exp\left(\frac{r}{4\pi i} (-2\pi^2 \zeta - \text{Li}_2(e^{2\pi i \zeta}) + \sum_{j=1}^4 \text{Li}_2\left(\frac{e^{2\pi i \zeta}}{e^{2\pi i \sigma_j}}\right) + \sum_{j=1}^3 \text{Li}_2\left(\frac{e^{2\pi i \tau_j}}{e^{2\pi i \zeta}}\right) - 6\pi^2 \zeta^2 + 8\pi^2 (\eta_a + \dots + \eta_f) \zeta)\right)$$

does not depend on ζ since $\text{Re Li}_2(e^{2\pi i x}) = \pi^2((x - \lfloor x \rfloor)^2 - \frac{1}{2})^2 - \frac{\pi^2}{12}$ for $x \in \mathbb{R}$ where $\lfloor x \rfloor$ is the largest integer not exceeding x . Let

$$\begin{aligned}
 F(\zeta, \eta) &= \frac{1}{4\pi i} \left(-2\pi^2 \zeta - \text{Li}_2(e^{2\pi i \zeta}) + \sum_{j=1}^4 \text{Li}_2\left(\frac{e^{2\pi i \zeta}}{e^{2\pi i \sigma_j}}\right) + \right. \\
 &\quad \left. \sum_{j=1}^3 \text{Li}_2\left(\frac{e^{2\pi i \tau_j}}{e^{2\pi i \zeta}}\right) - 6\pi^2 \zeta^2 + 8\pi^2 (\eta_a + \dots + \eta_f) \zeta \right).
 \end{aligned}$$

Let ζ_0 ($m_r \leq \zeta_0 \leq M_r$) be a solution of

$$\frac{\partial}{\partial \zeta} \text{Re} F(\zeta, \eta) = 0.$$

Then the derivative of $\text{Im} F(\zeta, \eta)$ with respect to ζ is 0 and ζ_0 satisfies

$$(27) \quad \frac{\partial}{\partial \zeta} F(\zeta, \eta) = 0.$$

The equation (27) is essentially a quadratic equation and we have only one solution which corresponds to the maximum. If we had two solutions corresponding to maxima, we also should have at least one solution corresponding to a minimum, but the total number of the solutions of (27) is two.

Since $\zeta_0 \in \mathbf{R}$ and the integral path is the real line, we can apply the stationary phase method to $|\int_{-\infty}^{\infty} g_r(\zeta, \eta) d\zeta| = \int_{-\infty}^{\infty} |g_r(\zeta, \eta)| d\zeta$ and we have

$$(28) \quad \left| \int_{-\infty}^{\infty} g_r(\zeta, \eta) d\zeta \right| \underset{r \rightarrow \infty}{\sim} \frac{16}{r^3} \left| \frac{e^{2\pi i \zeta_0}}{\prod_{j=1}^4 \sqrt{1 - \frac{e^{2\pi i \zeta_0}}{e^{2\pi i \sigma_j}}}} \sqrt{\frac{2\pi}{r |F_{\zeta\zeta}(\zeta_0, \eta)|}} \exp(rF(\zeta_0, \eta) + O(\frac{1}{r})) \right|.$$

Lemma 5.

$$\frac{\prod_{j=1}^4 (1 - \frac{e^{2\pi i \zeta_0}}{e^{2\pi i \sigma_j}})}{e^{4\pi i \zeta_0}} F_{ss}(\zeta_0, \eta) = \pm 4\pi \sqrt{-\det G}$$

where G is the gram matrix of T .

Proof. The second derivative of $F(\zeta, \eta)$ at ζ_0 is given by

$$F_{\zeta\zeta}(\zeta_0, \eta) = \pi i \left(-\frac{e^{2\pi i \zeta_0}}{1 - e^{2\pi i \zeta_0}} + \sum_{j=1}^4 \frac{e^{2\pi i \zeta_0}}{e^{2\pi i \sigma_j} - e^{2\pi i \zeta_0}} - \sum_{j=1}^3 \frac{e^{2\pi i \zeta_0}}{e^{2\pi i \tau_j} - e^{2\pi i \zeta_0}} \right).$$

On the other hand,

$$\prod_{j=1}^4 (e^{2\pi i \sigma_j} - e^{2\pi i \zeta_0}) = (1 - e^{2\pi i \zeta_0}) \prod_{j=1}^3 (e^{2\pi i \tau_j} - e^{2\pi i \zeta_0})$$

since $F_{\zeta}(\zeta_0) = 0$. Let

$$(29) \quad a_2 u^2 + a_1 u + a_0 = \left(\prod_{j=1}^4 (e^{2\pi i \sigma_j} - u) - (1 - u) \prod_{j=1}^3 (e^{2\pi i \tau_j} - u) \right) / u.$$

Then $e^{2\pi i \zeta_0}$ is a solution of $a_2 u^2 + a_1 u + a_0 = 0$. Let $u_0 = e^{2\pi i \zeta_0}$ and u'_0 be another solution of (29). Then we have

$$\begin{aligned} & \frac{\prod_{j=1}^4 (e^{2\pi i \sigma_k} - u_0)}{u_0^2} F_{\zeta\zeta}(\zeta_0, \eta) \\ &= \frac{\pi i}{u_0} \left(-\prod_{j=1}^3 (e^{2\pi i \tau_j} - u_0) + \sum_{j=1}^4 \prod_{1 \leq k \leq 4, k \neq j} (e^{2\pi i \sigma_k} - u_0) - \sum_{j=1}^3 (1 - u_0) \prod_{1 \leq k \leq 3, k \neq j} (e^{2\pi i \tau_k} - u_0) \right) \\ &= \pi i (3a_2 u_0 + 2a_1 + a_0 u_0^{-1}) = \pi i (a_2 u_0 - a_0 u_0^{-1}) = a_2 \pi i (u_0 - u'_0) \\ (30) \quad &= \pm \pi i \sqrt{a_1^2 - 4a_0 a_2} = \pm 4\pi \sqrt{-\det G}. \end{aligned}$$

Here we use

$$a_1^2 - 4a_0 a_2 = 16 \det G$$

which is obtained by direct computation. □

Recall that the volume $\text{Vol}(T)$ is given in [12] and [20] as follows.

$$(31) \quad \text{Vol}(T) =$$

$$\left| 2\pi (F(\zeta_0, \eta) + \delta(\eta_a, \eta_b, \eta_e) + \delta(\eta_a, \eta_c, \eta_f) + \delta(\eta_b, \eta_d, \eta_f) + \delta(\eta_c, \eta_d, \eta_e)) \right|.$$

Combining (18), (17), (20), (26), (28), (30) and (31), we get

$$\begin{aligned}
& \left| \begin{Bmatrix} a_r & b_r & e_r \\ d_r & c_r & f_r \end{Bmatrix}_{q=\xi_r^2}^{RW} \right| \sim_{r \rightarrow \infty} \\
& 2\pi \frac{r^2}{16} \frac{16}{r^3} \left| \frac{e^{2\pi i \zeta_0}}{\prod_{j=1}^4 \sqrt{1 - \frac{e^{2\pi i \zeta_0}}{e^{2\pi i \sigma_j}}}} \sqrt{\frac{2\pi}{r |F_{\zeta\zeta}(\zeta_0, \eta)|}} \exp\left(\frac{r}{2\pi} \text{Vol}(T) + O\left(\frac{1}{r}\right)\right) \right| \sim_{r \rightarrow \infty} \\
& \frac{\sqrt{2}\pi}{r^{3/2} \sqrt[4]{-\det G}} e^{\frac{\text{Vol}(T)r}{2\pi} + O\left(\frac{1}{r}\right)}.
\end{aligned}$$

5. SYMMETRY

A widely applicable version of the volume conjecture was introduced by the first author and T. Yang in [4]. The original volume conjecture is for knots and links in S^3 , while this new version works well for the Turaev-Viro invariants [19] and the Witten-Reshetikhin-Turaev invariants [16] of 3-manifolds. Here the Turaev-Viro invariant include the extension to cusped 3-manifolds and hyperbolic 3-manifolds with totally geodesic boundaries (see [2, 4]). Let $\xi_r = \exp(2\pi\sqrt{-1}/r)$ for a positive odd integer $r \geq 3$. Substituting $q = \xi_r^2$ instead of $q = \xi_r$ and doing numerical experiments for some manifolds, they observe that the Turaev-Viro invariant and the Witten-Reshetikhin-Turaev invariant have exponential growth with respect to r and the leading asymptotic is determined by the hyperbolic volume of the manifold.

Conjecture 2 (Chen-Yang [4]). Let M be a hyperbolic manifold and let $\text{Vol}(M)$ be its hyperbolic volume.

- (1) Let M be a 3-manifold which is closed, cusped, or with totally geodesic boundary, and let $TV_r(M)$ be the Turaev-Viro invariant [19, 4, 2] with $q = \xi_r^2$ for odd r . Then

$$\lim_{r \rightarrow \infty, r: \text{odd}} \frac{2\pi}{r} \log |TV_r(M)| = \text{Vol}(M).$$

- (2) Let $WRT_r(M)$ be the Witten-Reshetikhin-Turaev invariant [21, 16, 3] with $q = \xi_r^2$ for odd r . Then

$$\lim_{r \rightarrow \infty, r: \text{odd}} \frac{4\pi\sqrt{-1}}{r} \log WRT_r(M) = \text{CS}(M) + \text{Vol}(M)\sqrt{-1} \pmod{\pi^2 \mathbf{Z}}$$

where $\text{CS}(M)$ is the Chern-Simons invariant of M multiplied by $2\pi^2$.

Conjecture 2 is proved for some cases, one is for 3-manifolds obtained by integral surgeries along the figure-eight knot by T. Ohtsuki in [14], and another one is for the complements of the figure-eight knot and the Borromean rings by R. Detcherry, E. Kalfagianni and T. Yang in [6] and those for fundamental shadow links in [1]. Since the Turaev-Viro invariant is constructed by using the quantum $6j$ symbols, this conjecture suggests that the terms of asymptotic expansion of the quantum $6j$ symbol are all expressed by some geometric data of the corresponding tetrahedra.

We are considering a non-oriented tetrahedron. In this case, we can think of its volume, area of faces, lengths of edge and dihedral angles along edges. But there is no canonical way to assign the signs of these contents. So formulas of geometric contents of a tetrahedron must have certain symmetry for changing the signs of the lengths and angles. For example, the entries of the Gram matrix is given in terms of cosines of the dihedral angles and they don't depend on the signs of dihedral angles. In the construction of the Turaev-Viro invariant, the normalized quantum $6j$ symbol $\left| \begin{smallmatrix} a & b & e \\ d & c & f \end{smallmatrix} \right|$ is assigned to

each tetrahedron instead of $\left\{ \begin{smallmatrix} a & b & e \\ d & c & f \end{smallmatrix} \right\}^{RW}$ where

$$\left| \begin{smallmatrix} a & b & e \\ d & c & f \end{smallmatrix} \right| = \sqrt{-1}^{-2(a+b+c+d+e+f)} \left\{ \begin{smallmatrix} a & b & e \\ d & c & f \end{smallmatrix} \right\}^{RW}.$$

So it is natural to expect certain symmetry about each parameter of $\begin{vmatrix} a & b & e \\ d & c & f \end{vmatrix}$ reflecting the change of the sign of the corresponding dihedral angle. Here we consider such symmetry for the following continuous version of the quantum 6j symbol. Let

$$F_r(\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f) = 2(-1)^{\frac{r+1}{2}} \sin \frac{2\pi}{r} e^{\frac{2\pi i}{r}(d_3(a_r, b_r, e_r) + d_3(a_r, c_r, f_r) + d_3(b_r, d_r, f_r) + d_3(c_r, d_r, e_r))} \times \\ \tilde{\Delta}_r(a_r, b_r, e_r) \tilde{\Delta}_r(a_r, c_r, f_r) \tilde{\Delta}_r(b_r, d_r, f_r) \tilde{\Delta}_r(c_r, d_r, e_r) \int_{-\infty}^{\infty} \psi_r(z) \tilde{\alpha}_r(z) dz,$$

where $a_r = \frac{r(\pi - \theta_a)}{4\pi} - \frac{1}{2}, \dots, f_r = \frac{r(\pi - \theta_f)}{4\pi} - \frac{1}{2}$. For the functions $\tilde{\alpha}_r, \tilde{\Delta}_r, \psi_r$, refer (14), (15) and (16) respectively. According to the argument in Appendix, the asymptotics of $\begin{vmatrix} a_r & b_r & e_r \\ d_r & c_r & f_r \end{vmatrix}$ with respect to r is equal to that of $F_r(\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f)$ where $a_r = \frac{r(\pi - \theta_a)}{4\pi} - \frac{1}{2}, \dots, f_r = \frac{r(\pi - \theta_f)}{4\pi} - \frac{1}{2}$.

Conjecture 3 (Symmetry of asymptotics). Let T be a hyperbolic truncated tetrahedron and $\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f$ be dihedral angles at edges a, \dots, f in Figure 1. Then the asymptotic expansions of $F_r(\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f)$ and $F_r(-\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f)$ with respect to r are expected to agree at all orders of $\frac{1}{r}$.

This conjecture explains the very mysterious big cancellation happening in the Chen-Yang's Volume Conjecture of the Turaev-Viro invariants.

Remark 2. If $a_r, b_r, c_r, d_r, e_r, f_r$ are admissible and $a'_r, b'_r, c_r, d_r, e_r, f'_r$ are also admissible, then we have the following identity proved in [6].

$$\begin{vmatrix} a_r & b_r & e_r \\ d_r & c_r & f_r \end{vmatrix}_{q=\xi_r^2} = \begin{vmatrix} a'_r & b'_r & e_r \\ d_r & c_r & f'_r \end{vmatrix}_{q=\xi_r^2}$$

where $a'_r = r \frac{\pi + \theta_a}{4\pi} - \frac{1}{2}$, i.e. $\theta'_a = -\theta_a$. Applying our Conjecture 3 three times predicts that those 6j-symbols have the same expansion, but in this case the identity shows that they are completely equal.

Remark 3. Let

$$F_r(\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f) \underset{r \rightarrow \infty}{\sim} (-1)^{\frac{r+1}{2}} A e^{\frac{r}{2\pi} B} \left(1 + C_1 \frac{1}{r} + C_2 \frac{1}{r^2} + \dots \right)$$

be the asymptotic expansion of $F_r(\theta_a, \theta_b, \theta_c, \theta_d, \theta_e, \theta_f)$. Then we showed that

$$A = \frac{\sqrt{2}}{r^{3/2} \sqrt[4]{-\det G}}, \quad B = \text{Vol}(T),$$


where $\text{Vol}(T)$ is the volume of the tetrahedron T with dihedral angles $\theta_a, \dots, \theta_f$ and $\det G$ is the Gram matrix of T . The Conjecture 3 means that the coefficients A, B, C_1, C_2, \dots are all symmetric with respect to the switching of the sign of dihedral angles. The volume function $\text{Vol}(T)$ and the Gram matrix $\det G$ are not changed by changing the sign of any dihedral angle, the above conjecture is true up to the second leading term.

We computed the term C_1 . Let $C_1(\theta_a)$ denote the term C_1 with 5 fixed parameters ($\theta_b = \frac{2}{25}\pi, \theta_c = \frac{3}{25}\pi, \theta_d = \frac{4}{25}\pi, \theta_e = \frac{1}{5}\pi, \theta_f = \frac{6}{25}\pi$). We computed $C_1(\theta_a)$ numerically as the following table and graph. It suggests that $C_1(\theta_a) = C_1(-\theta_a)$.

APPENDIX. COMPARISON OF $\hat{h}_r(0)$ AND THE SUM OF $\tilde{\alpha}_r(z)$

Let $\beta_r(x) = \frac{1}{r} \log \tilde{\alpha}_r(\frac{rx-3}{2})$. Then $\beta_r(x)$ is a real analytic function defined on the interval (u, v) with $u = \frac{2m_r+3}{r}$ and $v = \frac{2M_r+3}{r}$ where m_r and M_r are the lower and upper bounds in the line below (6). Note that u and v do not depend on r . Such function $\beta_r(x)$ exists since $\tilde{\alpha}_r(\frac{rx-3}{2})$ is a positive real analytic function.

θ_a	$-\frac{3}{5}\pi$	$-\frac{2}{5}\pi$	$-\frac{1}{5}\pi$	0	$\frac{1}{5}\pi$	$\frac{2}{5}\pi$	$\frac{3}{5}\pi$
$C_1(\theta_a)$	4.85928	2.33391	1.98659	1.90677	1.98659	2.33391	4.85928

TABLE 1. Values of $C_1(\theta_a)$.FIGURE 2. Graph of $C_1(\eta_a)$.

Thanks to Lemma 2, $\beta_r(x)$ has a limiting function $\beta(x) = \lim_{r \rightarrow \infty} \beta_r(x)$, which is a real analytic function which takes its maximum at $x = x_0 \in (u, v)$. Recall that $h_r(z) = \psi_r(\frac{2z+3}{r}) \tilde{\alpha}_r(z)$. Then, by applying the Poisson summation formula, we have

$$\sum_{z \in \mathbf{Z}} h_r(z) = \sum_{m \in \mathbf{Z}} \hat{h}_r(m),$$

and

$$\begin{aligned} \hat{h}_r(m) &= \int_{\mathbf{R}} e^{-2\pi i m z} \psi\left(\frac{2z+3}{r}\right) \tilde{\alpha}_r(z) dz = \int_{\mathbf{R}} e^{-2\pi i m \frac{rx-3}{2}} \psi(x) \tilde{\alpha}_r\left(\frac{rx-3}{2}\right) \frac{r}{2} dx \\ &= (-1)^m \frac{r}{2} \int_{\mathbf{R}} \psi(x) e^{r(\beta_r(x) - \pi i m x)} dx. \end{aligned}$$

Lemma 6. *There are positive constants c'' and δ satisfying*

$$(32) \quad \frac{|\hat{h}_r(1)|}{|\hat{h}_r(0)|} < c'' e^{-r\delta}.$$

for some $\delta > 0$.


Proof. At first, we extend $\beta(x)$ to a complex-valued function around $x = x_0$. Let $\beta_{r,1}(x) = \beta_r(x) - \pi i x$. Choose $\varepsilon' > 0$ so that $\beta(x) - \pi i x$ doesn't have a critical point in the ε' neighborhood $B(x_0, \varepsilon')$ of x_0 in \mathbf{C} and $[u + 2\varepsilon, v - 2\varepsilon] \cap B(x_0, \varepsilon') = (x_0 - \varepsilon', x_0 + \varepsilon')$ where ε is the small positive number introduced in the definition of Ψ in (16). Let $x_0^{(r)}$ be the maximum point of $\beta_r(x)$, then $x_0^{(r)}$ is close to x_0 when r is sufficiently large and

$$\beta_r(x) = \beta_r(x_0^{(r)}) - a_r(x - x_0^{(r)})^2 + O((x - x_0^{(r)})^3)$$

in a small neighborhood of $x_0^{(r)}$. For $x \in \mathbf{C}$ near $x_0^{(r)}$, the contours of $\text{Re } \beta_r(x)$ and $\text{Re } \beta_{r,1}(x)$ are given in the following figure. Choose $x_1 \in (x_0 - \varepsilon', x_0)$ so that the contour $C_{r,1}$ corresponding to the value $\text{Re } \beta_{r,1}(x_1)$ connects x_1 to a real point $x_1^{(r)} \in (x_0, x_0 + \varepsilon)$ in $B(x_0, \varepsilon')$ and for any sufficiently large r as in the figure. Please note that $\text{Re } \beta_{r,1}(x) = \text{Re } \beta_r(x)$ for $x \in \mathbf{R}$. Then, due to $\beta_r(x_1) \rightarrow \beta(x_1)$, $\beta_r(x_0^{(r)}) \rightarrow \beta(x_0)$ when $r \rightarrow \infty$ and $\beta(x_1) \neq \beta(x_0)$, there is $\delta > 0$ such that $\beta_r(x_1) < \beta_r(x_0^{(r)}) - \delta$ for any sufficiently large r .

Now recall that

$$\hat{h}_r(0) = \frac{r}{2} \int_{u+\varepsilon}^{v-\varepsilon} \psi(x) e^{r\beta_r(x)} dx \underset{r \rightarrow \infty}{\sim} \text{const. } r^{-5/2} e^{r\beta(x_0)},$$



$$\begin{aligned} |\widehat{h}_r(1)| &= \frac{r}{2} \left| \int_{u+\epsilon}^{v-\epsilon} \psi(x) e^{r(\beta_r(x) - \pi i x)} dx \right| \leq \frac{r}{2} \int_{C'} \psi(x) e^{r(\text{Re}(\beta_r(x) - \pi i x))} dx \\ &\leq \frac{r}{2} \int_{C'} \psi(x) e^{r(\text{Re } \beta_r(x_1))} dx = \frac{r}{2} \ell(C') e^{r \text{Re } \beta_r(x_1)} \end{aligned}$$

where $C' = [u, x_1] \cup C_{r,1} \cup [x_1^{(r)}, v]$ and $\ell(C')$ is the length of C' . Since $\text{Re } \beta_r(x_1) < \text{Re } \beta_r(x_0) - \delta$, the relation (17) holds for some positive constant c'' . \square

Lemma 7. For $m > 1$, $\widehat{h}(m)$ satisfies

$$(33) \quad |\widehat{h}_r(m)| \leq \frac{D}{m^2} |\widehat{h}_r(1)|.$$

for some constant D not depend on m .

Proof. We have $\text{Re}(\beta_r(z) - \pi m i z) = \text{Re}(\beta_r(z) - \pi i z)$ on $[u, v]$ and $\text{Re}(\beta_r(z) - \pi m i z) \leq \text{Re}(\beta_r(z) - \pi i z)$ on $C_{r,1}$. Now we reformulate $\widehat{h}_r(m)$ as

$$\begin{aligned} \widehat{h}_r(m) &= (-1)^m \frac{r}{2} \int_{C'} e^{-\pi i r m} \psi(x) e^{r \beta_r(x)} dx \\ &= \frac{(-1)^{m+1} r}{2 \pi i m} \int_{C'} e^{-\pi i r m} (\psi'(x) + r \beta_r'(x)) e^{r \beta_r(x)} dx \\ &= \frac{(-1)^{m+1} r}{4 \pi^2 m^2} \int_{C'} e^{-\pi i r m} (\psi''(x) + r \beta_r''(x) + r (\psi'(x) + r \beta_r'(x)) \eta_r'(x)) e^{r \beta_r(x)} dx \end{aligned}$$

then there is a constant d satisfying

$$|e^{-\pi i r m} (\psi''(x) + r \beta_r''(x) + r (\psi'(x) + r \beta_r'(x)) \eta_r'(x))| < d.$$

$$\text{Hence } |\widehat{h}_r(m)| < \frac{dr}{4\pi^2 m^2} |\widehat{h}_r(1)|. \quad \square$$

Proof of Lemma 2. The above properties also holds for $\widehat{h}_r(-1)$ and $\widehat{h}_r(m)$ for negative m . So we have $|\sum_{m \neq 0} \widehat{h}_r(m)| \leq M |\widehat{h}_r(1)|$ for some constant M since $\sum_{m \neq 1} \frac{1}{m^2}$ converges. Hence Lemma 6 implies (17). \square

Now we prove Lemma 3.

Proof of Lemma 3. Let $x_1, x_1^{(r)}$ be points on $[u, v]$ given in the proof of Lemma 6. Then $\Psi(x) = 1$ for $x \in [x_1, x_1^{(r)}]$. Therefore $\widetilde{\alpha}_r(x) - h_r(x) = 0$ on $[x_1, x_1^{(r)}]$. Since $|h_r(x)| \leq |\widetilde{\alpha}_r(x)|$ and $|\widetilde{\alpha}_r(x)|$ is increasing for $x \leq x_1$ and decreasing for $x \geq x_1^{(r)}$. Hence

$$|A_r - B_r| \leq 2r |\widetilde{\alpha}_r(x_1)| = 2r e^{r \beta_r(x_1)} < 2r e^{r(\beta_r(x_0^{(r)}) - \delta)}.$$

By putting $\delta' = \delta$, this implies (18). \square

ACKNOWLEDGEMENTS

The first author would like to thank Francis Bonahon and Nicolai Reshetikhin for their strong interests, valuable discussions, as well as cordial guidance during past years. He also thank Giovanni Felder for his large interests, strong support and enlightening discussions, and Rinat Kashaev, Tomotada Ohtsuki, Shing-Tung Yau for their strong interests, hospitality, encouragement and inspiring discussion. Finally he would like to thank Tian Yang for his proof reading, discussion and many valuable suggestions.

REFERENCES

- [1] Beletti G., Detcherry R., Kalfagianni E., Yang T.: Growth of quantum $6j$ -symbols and applications to the volume conjecture, arXiv:1807.03327.
- [2] Benedetti R., Petronio C.: On Robert's proof of the Turaev-Walker theorem, *J. Knot Theory Ramifications* **5** (1996), 427–439.
- [3] Blanchet C., Habegger N., Masbaum G., Vogel P.: Three-manifold invariants derived from the Kauffman bracket, *Topology* **31** (1992), 685–699.
- [4] Chen Q., Yang T.: 'Volume conjectures for the Reshetikhin-Turaev and the Turaev-Viro invariants', *Quantum Topol.* **9** (2018), 419–460.
- [5] Costantino F.: $6j$ -symbols, hyperbolic structures and the volume conjecture, *Geom. Topol.* **11** (2007), 1831–1854.
- [6] Detcherry R., Kalfagianni E., Yang T.: Turaev-Viro invariants, colored Jones polynomials and volume, *Quantum Topol.* **9** (2018), 775–813.
- [7] Faddeev L.: Discrete Heisenberg-Weyl group and modular group, *Lett. Math. Phys.* **34** (1995), 249–254.
- [8] Kashaev R. M.: The hyperbolic volume of knots from the quantum dilogarithm, *Lett. Math. Phys.* **39** (1997), 269–275.
- [9] Kirillov A. N., Reshetikhin N. Yu.: Representations of the algebra $\mathcal{U}_q(sl(2))$, q -orthogonal polynomials and invariants of links, *Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988)*, Adv. Ser. Math. Phys., Vol. 7, World Sci. Publ., Teaneck, NJ, 1989, pp. 285–339.
- [10] Murakami H.: An introduction to the volume conjecture, *Interactions between hyperbolic geometry, quantum topology and number theory*, Contemp. Math., Vol. 541, Amer. Math. Soc., Providence, RI, 2011, pp. 1–40.
- [11] Murakami H., Murakami J.: The colored Jones polynomials and the simplicial volume of a knot, *Acta Math.* **186** (2001), 85–104.
- [12] Murakami J., Yano M.: On the volume of a hyperbolic and spherical tetrahedron, *Comm. Anal. Geom.* **13** (2005), 379–400.
- [13] Ohtsuki T.: On the asymptotic expansion of the Kashaev invariant of the 5_2 knot, *Quantum Topology* **7** (2016), 669–735.
- [14] Ohtsuki T.: 'On the asymptotic expansion of the quantum $SU(2)$ invariant at $q = \exp(4\pi\sqrt{-1}/N)$ for closed hyperbolic 3-manifolds obtained by integral surgery along the figure-eight knot', *Algebr. Geom. Topol.* **18** (2018), 4187–4274.
- [15] Ponzano G., Regge T.: 'Semiclassical limit of Racah coefficients', *Spectroscopic and group theoretical methods in physics*, Wiley Interscience, 1968, pp. 1–98.
- [16] Reshetikhin N. Yu, Turaev V. G.: 'Invariants of 3-manifolds via link polynomials and quantum groups', *Invent. Math.* **103** (1991), 547–597.
- [17] Roberts J.: Classical $6j$ -symbols and the tetrahedron. *Geom. Topol.* **3** (1999), 21–66.
- [18] Taylor Y. and Woodward C.: $6j$ symbols for $Uq(sl(2))$ and non-Euclidean tetrahedra, *Selecta Math. (N.S.)* **11** (2005), 539–571.
- [19] Turaev V. G. and Viro O.: State sum invariants of 3-manifolds and quantum $6j$ -symbols, *Topology* **4** (1992), 865–902.
- [20] Ushijima A.: A volume formula for generalized hyperbolic tetrahedra, *Non-Euclidean geometries*, Math. Appl. (N. Y.), Vol. 581, Springer, New York, 2006, pp. 249–265.
- [21] Witten E.: Quantum field theory and the Jones polynomial, *Comm. Math. Phys.* **121** (1989), 351–399.

Qingtao Chen, Division of Science, New York University Abu Dhabi, PO Box 129188 Abu Dhabi, United Arab Emirates, chenqtao@nyu.edu

Jun Murakami, Department of Mathematics, Faculty of Science and Engineering, Waseda University, 3-4-1 Ohkubo, Shinjuku-ku, Tokyo 169-8555 Japan, murakami@waseda.jp