

**Erratum: Numerical values of the f^F, f^D and f^S coupling
constants in SU(3) invariant Lagrangian of the interaction of the
vector-meson nonets with $1/2^+$ octet baryons [Phys. Rev. C93,
055208 (2016)]**

Cyril Adamuščin, Erik Bartoš, and Stanislav Dubníčka

Institute of Physics, Slovak Academy of Sciences, Bratislava, Slovak Republic

Anna Z. Dubníčková

Department of Theoretical Physics, Comenius University, Bratislava, Slovak Republic

(Dated: February 26, 2022)

Some aspects of the presented determination of numerical values of the f^F, f^D and f^S coupling constants in the SU(3) invariant interaction Lagrangian

$$\begin{aligned}
L_{VBB} = & \frac{i}{\sqrt{2}} f^F [\bar{B}_\beta^\alpha \gamma_\mu B_\gamma^\beta - \bar{B}_\gamma^\beta \gamma_\mu B_\beta^\alpha] (V^\mu)_\alpha^\gamma + \\
& + \frac{i}{\sqrt{2}} f^D [\bar{B}_\beta^\alpha \gamma_\mu B_\gamma^\beta + \bar{B}_\gamma^\beta \gamma_\mu B_\beta^\alpha] (V^\mu)_\alpha^\gamma + \\
& + \frac{i}{\sqrt{2}} f^S \bar{B}_\beta^\alpha \gamma_\mu B_\alpha^\beta \omega_0^\mu
\end{aligned}$$

of the vector-meson nonet with $1/2^+$ octet baryons in [1] were not correct.

The crucial ingredient in that determination is an application of the $\omega - \phi$ mixing, which has been entered into the procedure in [1] two times. First in a derivation of expressions (51), (52), (55) and (56) for coupling constants of vector-mesons with nucleons from the above-mentioned Lagrangian, and so, also in a derivation of the reverse expressions (66), (67), (68) and (69), and then in a determination of the signs of the universal vector-meson coupling constants f_ρ, f_ω and f_ϕ .

Generally there are in literature the following four different physically acceptable forms of the $\omega - \phi$ mixing

$$\begin{aligned}
1. \quad \phi &= \omega_8 \cos \theta - \omega_0 \sin \theta \\
\omega &= \omega_8 \sin \theta + \omega_0 \cos \theta
\end{aligned} \tag{1}$$

$$\begin{aligned}
2. \quad \phi &= -\omega_8 \cos \theta + \omega_0 \sin \theta \\
\omega &= \omega_8 \sin \theta + \omega_0 \cos \theta
\end{aligned} \tag{2}$$

$$\begin{aligned}
3. \quad \phi &= -\omega_8 \cos \theta + \omega_0 \sin \theta \\
\omega &= -\omega_8 \sin \theta - \omega_0 \cos \theta
\end{aligned} \tag{3}$$

$$\begin{aligned}
4. \quad \phi &= \omega_8 \cos \theta - \omega_0 \sin \theta \\
\omega &= -\omega_8 \sin \theta - \omega_0 \cos \theta
\end{aligned} \tag{4}$$

which manifest themselves in the four different forms of expressions for f^F, f^D, f^S

$$\begin{aligned}
1. \quad f^F &= \frac{1}{2}[f_{\rho NN} + \sqrt{3}(f_{\Phi NN} \cos \theta + f_{\omega NN} \sin \theta)] \\
f^D &= \frac{1}{2}[3f_{\rho NN} - \sqrt{3}(f_{\Phi NN} \cos \theta + f_{\omega NN} \sin \theta)]
\end{aligned} \tag{5}$$

$$\begin{aligned}
f^S &= \sqrt{2}(f_{\omega NN} \cos \theta - f_{\Phi NN} \sin \theta) \\
2. \quad f^F &= \frac{1}{2}[f_{\rho NN} + \sqrt{3}(-f_{\Phi NN} \cos \theta + f_{\omega NN} \sin \theta)] \\
f^D &= \frac{1}{2}[3f_{\rho NN} - \sqrt{3}(-f_{\Phi NN} \cos \theta + f_{\omega NN} \sin \theta)] \\
f^S &= \sqrt{2}(f_{\omega NN} \cos \theta + f_{\Phi NN} \sin \theta)
\end{aligned} \tag{6}$$

$$\begin{aligned}
3. \quad f^F &= \frac{1}{2}[f_{\rho NN} - \sqrt{3}(f_{\Phi NN} \cos \theta + f_{\omega NN} \sin \theta)] \\
f^D &= \frac{1}{2}[3f_{\rho NN} + \sqrt{3}(f_{\Phi NN} \cos \theta + f_{\omega NN} \sin \theta)] \\
f^S &= -\sqrt{2}(f_{\omega NN} \cos \theta - f_{\Phi NN} \sin \theta)
\end{aligned} \tag{7}$$

$$\begin{aligned}
4. \quad f^F &= \frac{1}{2}[f_{\rho NN} + \sqrt{3}(f_{\Phi NN} \cos \theta - f_{\omega NN} \sin \theta)] \\
f^D &= \frac{1}{2}[3f_{\rho NN} - \sqrt{3}(f_{\Phi NN} \cos \theta - f_{\omega NN} \sin \theta)] \\
f^S &= -\sqrt{2}(f_{\omega NN} \cos \theta + f_{\Phi NN} \sin \theta).
\end{aligned} \tag{8}$$

On the other hand an application of the same $\omega - \phi$ mixing configurations (1), (2), (3), (4) leads to the rates of the overthrown values of the universal vector-meson coupling constants with different signs as follows

$$1. \quad \frac{1}{f_\rho} : \frac{1}{f_\omega} : \frac{1}{f_\phi} = \sqrt{3} : \sin \theta : \cos \theta, \tag{9}$$

which can be found e.g. in ref. [2] on p. 52 relation (A)

$$2. \quad \frac{1}{f_\rho} : \frac{1}{f_\omega} : \frac{1}{f_\phi} = \sqrt{3} : \sin \theta : -\cos \theta \tag{10}$$

to be found in ref. [3] on p. 539

$$3. \quad \frac{1}{f_\rho} : \frac{1}{f_\omega} : \frac{1}{f_\phi} = \sqrt{3} : -\sin \theta : -\cos \theta \tag{11}$$

$$4. \quad \frac{1}{f_\rho} : \frac{1}{f_\omega} : \frac{1}{f_\phi} = \sqrt{3} : -\sin \theta : \cos \theta, \tag{12}$$

and this last case can be found e.g. in ref. [4] on p. 446.

All these relations can be explained by the following considerations.

Starting e.g. from the $\omega - \phi$ mixing configuration (1) and substituting explicitly

$$\begin{aligned}\omega_8 &= \frac{1}{\sqrt{6}}(\bar{u}u + \bar{d}d - 2\bar{s}s) \\ \omega_0 &= \frac{1}{\sqrt{3}}(\bar{u}u + \bar{d}d + \bar{s}s)\end{aligned}\tag{13}$$

and for the ideal mixing angle $\theta = 35.3^\circ$

$$\sin \theta = \sqrt{\frac{1}{3}}; \quad \cos \theta = \sqrt{\frac{2}{3}},\tag{14}$$

one obtains

$$\phi = -\bar{s}s\tag{15}$$

and

$$\omega = +\frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d).\tag{16}$$

On the other hand the hadronic electromagnetic (EM) current

$$J_\mu^h = \frac{2}{3}\bar{u}\gamma_\mu u - \frac{1}{3}\bar{d}\gamma_\mu d - \frac{1}{3}\bar{s}\gamma_\mu s,\tag{17}$$

can be formally arranged to the shape

$$J_\mu^h = \frac{1}{2}(\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d) + \frac{1}{6}(\bar{u}\gamma_\mu u + \bar{d}\gamma_\mu d) - \frac{1}{3}(\bar{s}\gamma_\mu s),$$

and because

$$\begin{aligned}\frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d) &= J_\mu^{\rho^0} \\ +\frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu u + \bar{d}\gamma_\mu d) &= J_\mu^\omega \text{ due to the sign "+" in (16)} \\ -\bar{s}\gamma_\mu s &= J_\mu^\phi \text{ due to the sign "-" in (15),}\end{aligned}$$

are the ρ^0 -, ω -, ϕ - meson EM currents, respectively, the hadronic EM current acquires the following form

$$J_\mu^h = \frac{1}{\sqrt{2}}J_\mu^{\rho^0} + \frac{1}{3\sqrt{2}}J_\mu^\omega + \frac{1}{3}J_\mu^\phi.\tag{18}$$

Now, if the results of the Kroll-Lee-Zumino paper [5], that a linear combination of the neutral vector-meson fields $\rho_\mu^0, \omega_\mu, \phi_\mu$

$$J_\mu^h = \frac{m_{\rho^0}^2}{f_\rho}\rho_\mu^0 + \frac{m_\omega^2}{f_\omega}\omega_\mu + \frac{m_\phi^2}{f_\phi}\phi_\mu,\tag{19}$$

with the universal vector-meson coupling constants f_ρ, f_ω, f_ϕ , is proportional by some real constant A to the hadronic EM current (18), are taken into account, considering a dimension

of the Dirac quark fields u, d, s in (18), in the framework of the natural units $\hbar = c = 1$, to be $\text{m}^{\frac{3}{2}}$ and a dimension of the vector-meson fields in (19) to be m^1 , the relations

$$\frac{1}{f_\rho} = A \frac{1}{\sqrt{2}}; \quad \frac{1}{f_\omega} = +A \frac{1}{3\sqrt{2}}; \quad \frac{1}{f_\phi} = +A \frac{1}{3}, \quad (20)$$

are found.

Then from these last relations the rates

$$\begin{aligned} \frac{1}{f_\rho} : \frac{1}{f_\omega} : \frac{1}{f_\phi} &= \frac{1}{\sqrt{2}} : +\frac{1}{3\sqrt{2}} : +\frac{1}{3} \\ &= \frac{\sqrt{6}}{\sqrt{2}} : +\frac{\sqrt{6}}{3\sqrt{2}} : +\frac{\sqrt{6}}{3} \\ &= \sqrt{3} : +\frac{1}{\sqrt{3}} : +\sqrt{\frac{2}{3}} \\ &= \sqrt{3} : +\sin \theta : +\cos \theta, \end{aligned}$$

are obtained, giving the signs of the universal vector-meson coupling constants $+f_\rho, +f_\omega, +f_\phi$.

As a matter of fact the signs of the universal vector-meson coupling constants are specified already from the relations (20), however, a community of physicists prefers the relations (9)-(12), in which the universal vector-meson coupling constants are related to $\sin \theta$ and $\cos \theta$ where the angle θ is determined from the quadratic Gell-Mann-Okubo vector-meson mass formula, which provide more realistic values of the latter to be in fair agreement with experimental evaluations. Therefore we also favor a presentation of the universal vector-meson coupling constants signs in the form (9)-(12).

In a like manner, starting from the $\omega - \phi$ mixing configuration (2), and relations (13) and (14), one obtains

$$\phi = +\bar{s}s \quad (21)$$

and

$$\omega = +\frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d). \quad (22)$$

Then comparing the hadronic EM current to be multiplied by the real constant A

$$J_\mu^h = \frac{1}{\sqrt{2}}J_\mu^{\rho^0} + \frac{1}{3\sqrt{2}}J_\mu^\omega - \frac{1}{3}J_\mu^\phi, \quad (23)$$

where

$$J_\mu^{\rho^0} = \frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d)$$

$$J_\mu^\omega = +\frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu u + \bar{d}\gamma_\mu d) \text{ due to the sign "+" in (22)}$$

$$J_\mu^\phi = +\bar{s}\gamma_\mu s \text{ due to the sign "+" in (21),}$$

with (19), one obtains relations

$$\frac{1}{f_\rho} = A\frac{1}{\sqrt{2}}; \quad \frac{1}{f_\omega} = +A\frac{1}{3\sqrt{2}}; \quad \frac{1}{f_\phi} = -A\frac{1}{3} \quad (24)$$

and from them the rates (10), giving the following signs of universal vector-meson coupling constants $+f_\rho, +f_\omega, -f_\phi$.

Again, starting from the $\omega - \phi$ mixing configuration (3), and relations (13) and (14), one obtains

$$\phi = +\bar{s}s \quad (25)$$

and

$$\omega = -\frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d). \quad (26)$$

Then comparing the hadronic EM current to be multiplied by the real constant A

$$J_\mu^h = \frac{1}{\sqrt{2}}J_\mu^{\rho^0} - \frac{1}{3\sqrt{2}}J_\mu^\omega - \frac{1}{3}J_\mu^\phi, \quad (27)$$

where

$$J_\mu^{\rho^0} = \frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d)$$

$$J_\mu^\omega = -\frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu u + \bar{d}\gamma_\mu d) \text{ due to the sign "-" in (26)}$$

$$J_\mu^\phi = +\bar{s}\gamma_\mu s \text{ due to the sign in (25),}$$

with (19), one obtains relations

$$\frac{1}{f_\rho} = A\frac{1}{\sqrt{2}}; \quad \frac{1}{f_\omega} = -A\frac{1}{3\sqrt{2}}; \quad \frac{1}{f_\phi} = -A\frac{1}{3} \quad (28)$$

and from them the rates (11), giving the following signs of universal vector-meson coupling constants $+f_\rho, -f_\omega, -f_\phi$.

Finally, starting from the $\omega - \phi$ mixing configuration (4), and relations (13) and (14), one obtains

$$\phi = -\bar{s}s \quad (29)$$

and

$$\omega = -\frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d). \quad (30)$$

Then comparing the hadronic EM current to be multiplied by the real constant A

$$J_\mu^h = \frac{1}{\sqrt{2}}J_\mu^{\rho^0} - \frac{1}{3\sqrt{2}}J_\mu^\omega + \frac{1}{3}J_\mu^\phi, \quad (31)$$

where

$$J_\mu^{\rho^0} = \frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d)$$

$$J_\mu^\omega = -\frac{1}{\sqrt{2}}(\bar{u}\gamma_\mu u + \bar{d}\gamma_\mu d) \text{ due to the sign "-" in (30)}$$

$$J_\mu^\phi = -\bar{s}\gamma_\mu s \text{ due to the sign "-" in (29),}$$

with (19), one obtains relations

$$\frac{1}{f_\rho} = A\frac{1}{\sqrt{2}}; \quad \frac{1}{f_\omega} = -A\frac{1}{3\sqrt{2}}; \quad \frac{1}{f_\phi} = +A\frac{1}{3} \quad (32)$$

and from them the rates (12), giving the following signs of universal vector-meson coupling constants $+f_\rho, -f_\omega, +f_\phi$.

The signs of the universal vector-meson coupling constants f_ρ, f_ω, f_ϕ are very important to be known, as the numerical values of these constants are regularly estimated from the experimental values [6] of the vector-meson lepton widths by means of the formula

$$\Gamma(V \rightarrow e^+e^-) = \frac{\alpha^2 m_V}{3} \left(\frac{f_V^2}{4\pi} \right)^{-1}, \quad (33)$$

in which f_V is contained in a quadratic form.

From (5), (6), (7), (8) it seems at first sight that numerical values of the coupling constants f^F, f^D, f^S have to depend on the choice of the $\omega - \phi$ mixing version.

However, if in (5), (6), (7), (8) the coupling constants of vector-mesons with nucleons are determined e.g from $(f_{\rho NN}^{(1)}/f_\rho), (f_{\omega NN}^{(1)}/f_\omega), (f_{\phi NN}^{(1)}/f_\phi)$, to be found in a fitting procedure of all existing data on nucleon EM structure, by means of the signs of f_ρ, f_ω, f_ϕ following from (9), (10), (11), (12), as it is demonstrated above, then from all four different expressions in (5), (6), (7), (8) one obtains the same numerical values for f_1^F, f_1^D, f_1^S as follows

$$f_1^F = 5.414; \quad f_1^D = -1.699; \quad f_1^S = 42.916. \quad (34)$$

By means of a similar procedure one can find also numerical values of all other coupling constants under consideration

$$\begin{aligned} f_2^F &= 7.626; & f_2^D &= 21.088; & f_2^S &= -7.111; \\ f_1^{F'} &= 8.343; & f_1^{D'} &= 12.498; & f_1^{S'} &= -7.858; \\ f_2^{F'} &= -30.450; & f_2^{D'} &= -5.271; & f_2^{S'} &= -18.614. \end{aligned} \quad (35)$$

In the paper [1] for the numerical evaluation of the vector-meson-nucleon coupling constants from the $(f_{\rho NN}/f_\rho), (f_{\omega NN}/f_\omega), (f_{\phi NN}/f_\phi)$ we have applied the signs of the universal vector-meson coupling constants (10) (to be inspired by Close and Cottingham in [3]), which moreover have been combined with expressions for f^F, f^D, f^S

$$\begin{aligned} f^F &= \frac{1}{2} \left[f_{\rho NN} + \sqrt{3} (f_{\phi NN} \cos \theta - f_{\omega NN} \sin \theta) \right] \\ f^D &= \frac{1}{2} \left[3f_{\rho NN} - \sqrt{3} (f_{\phi NN} \cos \theta - f_{\omega NN} \sin \theta) \right] \\ f^S &= \sqrt{2} (f_{\omega NN} \cos \theta + f_{\phi NN} \sin \theta) \end{aligned} \tag{36}$$

to be generated by the physically non-acceptable form of the $\omega - \phi$ mixing

$$\begin{aligned} \phi &= \omega_8 \cos \theta + \omega_0 \sin \theta \\ \omega &= -\omega_8 \sin \theta + \omega_0 \cos \theta \end{aligned} \tag{37}$$

$$\tag{38}$$

used by Gasiorowicz [4] on p.325, leading to the numerical values of f^F, f^D, f^S differing from the correct values presented in this Erratum.

-
- [1] C.Adamuscin, E.Bartos, S.Dubnicka and A.Z.Dubnickova, Phys. Rev. C93, 055208 (2016)
 - [2] J. P.Perez-y-Jorba and F. M.Renard, Phys. Reports 31C, 1 (1977)
 - [3] F.E.Close and W.N.Cottingham: e^+e^- Annihilation in *Electromagnetic Interactions of Hadrons, Vol.2 Editors: A.Donnachie, G.Shaw, Originally published by Plenum Press, New York in 1978*
 - [4] S.Gasiorowicz: Elementary particle physics, John Wiley & Sons, Inc. New York, 1966
 - [5] N. M.Kroll, T. D.Lee and B. Zumino, Phys. Rev. 157, 1376 (1967)
 - [6] C. Patrignani et al.(Particle Data Group), Chin. Phys. C40, 100001 (2016)