

# Optimal modification of the LRT for the equality of two high-dimensional covariance matrices

Qiuyan Zhang, Jiang Hu\* and Zhidong Bai

KLASMOE & School of Mathematics and Statistics  
Northeast Normal University, China.

February 3, 2022

## Abstract

This paper considers the optimal modification of the likelihood ratio test (LRT) for the equality of two high-dimensional covariance matrices. The optimality here means that the modification of LRT cannot be improved anymore in our model settings. It is well-known that the classical log-LRT is not well defined when the dimension is larger than or equal to the sample size. Or even the log-LRT is well-defined, it is usually perceived as a bad statistic in high dimension cases for their low powers under some alternatives. In this paper, we shall argue some goodnesses of the modified log-LRT, and an optimally modified test that works well in cases where the dimension is larger than the sample sizes is proposed. Besides, the test is established under the weakest conditions on the moments and the dimensions of the samples. The asymptotic distribution of the proposed test statistic is also obtained under the null hypotheses. What is more, we also propose a lite version of the modified LRT in the paper. A simulation study and a real data analysis show that the performances of the two proposed statistics are confirmed to be invariant to affine transformations.

*Keywords:* Likelihood ratio test, High-dimensional data, Hypothesis testing, Random matrix theory.

---

\*Corresponding author.

# 1 Introduction

Since the assumption of homogeneity of covariance matrices is needed in many multivariate statistical analyses based on two populations, the equality of two covariance matrices is among the most active hypothesis tests. These tests date back to the work of Wilks (1932), which was followed by a huge amount of literature. Suppose we have  $N := N_1 + N_2$  observations  $\{\mathbf{z}_i^{(l)} \sim N(\boldsymbol{\mu}_l, \Sigma_l), i = 1, \dots, N_l, l = 1, 2\}$  and wish to test the hypothesis

$$H_0 : \Sigma_1 = \Sigma_2 \quad \text{v.s.} \quad H_1 : \Sigma_1 \neq \Sigma_2, \quad (1.1)$$

where  $\boldsymbol{\mu}_l$  and  $\Sigma_l$  are the population mean vectors and covariance matrices of the  $p$ -dimensional vectors  $\mathbf{z}_i^{(l)}$ ,  $l = 1, 2$  respectively. It is natural to first consider the likelihood ratio test (LRT) if it is “applicable”. But when is the LRT applicable to testing the equality of two covariance matrices? The traditional viewpoint is that the LRT is applicable if the sample sizes are both much larger than the dimensions based on the  $\chi^2$  approximation (Wilks, 1946). However, Bai et al. (2009) and Jiang and Yang (2013) showed that when the dimensions are large but smaller than the sample sizes, the traditional  $\chi^2$  approximation of the LRT fails to work well. This problem encouraged us to investigate the conditions under which the high-dimensional LRT is applicable. Additionally, we consider a lite LRT proposed by Pillai and Jayachandran (1967, 1968).

Currently, there are three general types of test procedures for the high-dimensional hypothesis (1.1) that are widely discussed in the literature: (i) Corrected classical LRTs, see, e.g., Bai et al. (2009); Jiang et al. (2012); Jiang and Yang (2013); (ii) Nonparametric methods, see, e.g., Ledoit and Wolf (2002); Li and Chen (2012); Srivastava and Yanagihara (2010); (iii) Maximum element methods, see, e.g., Cai et al. (2013); Cai and Ma (2013). The strengths and weaknesses of these three methods are significant. Nonparametric methods and maximum element methods can address cases where the dimensions are much larger than the sample sizes but are strongly restricted by the structure or eigenvalues of the

population covariance matrices. By contrast, corrected classical LRT requires the dimensions to be smaller than the sample sizes, but there is no assumption on the population covariance matrices. In addition, if the dimensions are fixed, LRT has been shown to be unbiased and uniformly most powerful among affine-transform-invariant tests. Therefore, we focus on the LRT.

In the following, we denote  $\mathbf{X}_{N_1}^{(1)} = (x_{ij}^{(1)})_{p \times N_1}$ , where  $\{x_{ij}^{(1)}\}$  are independent and identically distributed (i.i.d.) random variables with mean zero and variance one. Similarly  $\mathbf{X}_{N_2}^{(2)} = (x_{ij}^{(2)})_{p \times N_2}$ , which is independent with  $\mathbf{X}_{N_1}^{(1)}$ , constitutes another i.i.d. sample with mean zero and variance one. For  $l = 1, 2$ , we assume that the  $N_l$  observations  $\mathbf{z}_j^{(l)}$  satisfy the *linear transformation model*

$$\mathbf{z}_j^{(l)} = \Sigma_l^{1/2} \mathbf{x}_j^{(l)} + \boldsymbol{\mu}_l, \quad (1.2)$$

where  $\mathbf{x}_j^{(l)}$  is the  $j$ th column of  $\mathbf{X}_{N_l}^{(l)}$  and  $\boldsymbol{\mu}_l$  and  $\Sigma_l$  are the population mean vectors and covariance matrices of  $\{\mathbf{z}_j^{(l)}\}$ , respectively. Here,  $\Sigma_l^{1/2}$  can be chosen as any square root of matrix  $\Sigma_l$ . The linear transformation model covers the case where the samples  $\mathbf{z}_i^{(l)}$  are normally distributed, although we are not restricted here. Denote  $n_l = N_l - 1$ ,  $\bar{\mathbf{z}}^{(l)} = \frac{1}{N_l} \sum_{i=1}^{N_l} \mathbf{z}_i^{(l)}$ ,  $\mathring{\mathbf{z}}_i^{(l)} = \mathbf{z}_i^{(l)} - \bar{\mathbf{z}}^{(l)}$  and  $\mathbf{S}_l^{\mathbf{z}} := \mathbf{S}_{n_l}^{\mathbf{z}} = \frac{1}{n_l} \sum_{i=1}^{N_l} \mathring{\mathbf{z}}_i^{(l)} (\mathring{\mathbf{z}}_i^{(l)})'$ . We recall the famous Bartlett corrected LRT statistic  $L$  proposed by Bartlett (1937), which is given by,

$$L = \frac{2}{n_1 + n_2} \log \left( \frac{|\mathbf{S}_1^{\mathbf{z}}|^{\frac{n_1}{2}} \cdot |\mathbf{S}_2^{\mathbf{z}}|^{\frac{n_2}{2}}}{|c_1 \mathbf{S}_1^{\mathbf{z}} + c_2 \mathbf{S}_2^{\mathbf{z}}|^{\frac{n_1+n_2}{2}}} \right).$$

Here, and in the following, we denote  $c_1 = n_1/(n_1 + n_2)$  and  $c_2 = n_2/(n_1 + n_2)$ . Moreover, under the null hypothesis of (1.1) and linear transformation model (1.2), it is not difficult to rewrite this statistic as

$$L = \frac{2}{n_1 + n_2} \log \left( \frac{|\mathbf{S}_1^{\mathbf{x}}|^{\frac{n_1}{2}} \cdot |\mathbf{S}_2^{\mathbf{x}}|^{\frac{n_2}{2}}}{|c_1 \mathbf{S}_1^{\mathbf{x}} + c_2 \mathbf{S}_2^{\mathbf{x}}|^{\frac{n_1+n_2}{2}}} \right),$$

where  $\mathbf{S}_l^{\mathbf{x}} = \frac{1}{n_l} \sum_{i=1}^{N_l} (\mathbf{x}_i^{(l)} - \frac{1}{N_l} \sum_{i=1}^{N_l} \mathbf{x}_i^{(l)}) (\mathbf{x}_i^{(l)} - \frac{1}{N_l} \sum_{i=1}^{N_l} \mathbf{x}_i^{(l)})'$ ,  $l = 1, 2$ . Thus we know that  $L$  is independent with the population means  $\boldsymbol{\mu}_l$  and covariance matrices  $\Sigma_l$  under the null

hypothesis  $H_0$ . In the following, we drop the superscripts of  $\mathbf{S}$  and denote  $\mathbf{S}_1 := \mathbf{S}_1^{\mathbf{x}}$  and  $\mathbf{S}_2 := \mathbf{S}_2^{\mathbf{x}}$  for simplicity. In addition, by simple calculation, we can rewrite

$$L = c_1 \log(c_1^{-1} |\mathbf{B}_n|) + c_2 \log(c_2^{-1} |\mathbf{I}_p - \mathbf{B}_n|),$$

where

$$\mathbf{B}_n = n_1 \mathbf{S}_1 (n_1 \mathbf{S}_1 + n_2 \mathbf{S}_2)^{-1}.$$

We now analyze  $L$ . If  $p > n_1$  or  $p > n_2$ , then  $L$  is undefined because: (1) if  $p \geq n_1 + n_2$ , then matrix  $n_1 \mathbf{S}_1 + n_2 \mathbf{S}_2$  is singular, which makes the inverse of matrix  $n_1 \mathbf{S}_1 + n_2 \mathbf{S}_2$  undefined; and (2) if  $p < n_1 + n_2$ , i.e., the inverse of  $n_1 \mathbf{S}_1 + n_2 \mathbf{S}_2$  is well-defined almost surely (with the fourth moments finite assumption), but  $p > n_1$  or  $p > n_2$ , then at least one of the determinants of  $\mathbf{B}_n$  or  $\mathbf{I}_p - \mathbf{B}_n$  is zero, which makes the logarithm functions undefined. However, from random matrix theory (RMT), we know that if the fourth moment of  $x_{ij}^{(l)}$  exists, matrix  $\mathbf{B}_n$  almost certainly has  $p - n_1$  zero eigenvalues and  $p - n_2$  one eigenvalues according to the condition  $p > n_1$  and  $p > n_2$ , respectively (see Bai et al. (2015)). Therefore, we can naturally redefine the LRT  $L$  by restricting the non-zero and non-one eigenvalues of  $\mathbf{B}_n$ , i.e.,

$$\mathcal{L} = \sum_{\lambda_i^{\mathbf{B}_n} \in (0,1)} [c_1 \log \lambda_i^{\mathbf{B}_n} + c_2 \log(1 - \lambda_i^{\mathbf{B}_n})], \quad (1.3)$$

where  $\lambda_i^{\mathbf{B}_n}$  denotes the  $i$ -th smallest eigenvalue of  $\mathbf{B}_n$ . Therefore, we only need to obtain the asymptotic distributions of the redefined LRT in (1.3), which is addressed in the next section.

This paper also considers the test statistic  $\tilde{L}$ ,

$$\tilde{L} = \log |n_1 \mathbf{S}_1^{\mathbf{z}} (n_1 \mathbf{S}_1^{\mathbf{z}} + n_2 \mathbf{S}_2^{\mathbf{z}})^{-1}|,$$

which was proposed by Pillai and Jayachandran (1967, 1968) and can be viewed as a lite

LRT. Similar to  $\mathcal{L}$ , we redefine  $\tilde{L}$  by

$$\tilde{\mathcal{L}} = \sum_{\lambda_i^{\mathbf{B}_n} \in (0,1)} \log \lambda_i^{\mathbf{B}_n}. \quad (1.4)$$

It is obvious that  $\tilde{\mathcal{L}}$  is monotone for matrix  $\mathbf{B}_n$ ; thus, it should be more powerful than  $\mathcal{L}$ .

The rest of this paper is organized as follows. The main results are presented in Section 2, including the asymptotic normality of  $\mathcal{L}$  and  $\tilde{\mathcal{L}}$  and their optimal properties, which stand for the modification of LRT cannot be improved anymore in our model settings. In Section 3, we present the simulation results for the proposed statistics by comparison with that proposed by Li and Chen (2012) and Cai et al. (2013). In Section 5, we introduce a real data application to demonstrate the application of the proposed tests. All technical details are relegated to the appendix. We note that for the high-dimensional testing problem (1.1), the exact distribution of the test statistic is difficult to obtain when the distributions are free.

## 2 Main results

In this section, we give the asymptotic distributions of the redefined LRT in (1.3) and (1.4). For the application, we also present consistent estimators for the fourth moments of the samples under the null hypothesis. Before presenting the main results, we give some notation and the optimal assumptions. In the following, we denote the indicator function

by  $\delta(\cdot)$ , the natural logarithm function by  $\log(\cdot)$ , convergence in distribution by  $\xrightarrow{D}$ , and

$$\begin{aligned}
y_1 &:= y_{n1} = p/n_1, \quad y_2 := y_{n2} = p/n_2, \quad h := h_n = \sqrt{y_1 + y_2 - y_1 y_2}, \\
c_1 &:= n_1/(n_1 + n_2) = y_2/(y_1 + y_2), \quad c_1 := n_2/(n_1 + n_2) = y_1/(y_1 + y_2), \\
l(y_1, y_2) &= \log(h^{\frac{2c_1 h^2}{y_1 y_2}}) \delta_{y_1 > 1} - \log(y_1^{\frac{c_1(1+y_2)}{y_2}} y_2^{\frac{c_1(1-y_1)}{y_1}}) \delta_{y_1 > 1}, \\
u(y_1, y_2) &= \log\left(\frac{y_1^{c_1}}{h^{c_1}}\right) \delta_{y_1 > 1}, \quad v(y_1, y_2) = \log\left(\frac{y_1^{2c_1}}{h^{2c_1(c_1+2c_2)}}\right) \delta_{y_1 > 1}, \\
\Psi(y_1, y_2) &= c_2 y_1^2 [y_2^4 \delta_{y_2 < 1} + h^2(2y_2^2 - h^2) \delta_{y_2 > 1}] \\
&\quad - c_1 y_2^2 [y_1^3 (y_1 + 2y_2) \delta_{y_1 < 1} + h^2(y_1 + y_2 + y_1 y_2) \delta_{y_1 > 1}].
\end{aligned}$$

We set two assumptions of the sample that will be shown to be optimal for the proposed test statistics.

- (Moments Assumption:)  $\mathbb{E}x_{11}^{(1)} = \mathbb{E}x_{11}^{(2)} = 0, \mathbb{E}(x_{11}^{(1)})^2 = \mathbb{E}(x_{11}^{(2)})^2 = 1, \mathbb{E}(x_{11}^{(1)})^4 = \Delta_1 + 3 < \infty$  and  $\mathbb{E}(x_{11}^{(2)})^4 = \Delta_2 + 3 < \infty$ ;
- (Dimensions Assumption:)  $y_1 \neq 1, y_2 \neq 1$  and  $p/(n_1 + n_2) < 1$ .

We are now ready to present the main results of this paper.

**Theorem 2.1.** *In addition to the Moments Assumption and Dimensions Assumption, we assume that as  $\min\{p, n_1, n_2\} \rightarrow \infty$ ,  $\lim y_1 \notin \{0, 1\}$ ,  $\lim y_2 \notin \{0, 1\}$  and  $\lim p/(n_1 + n_2) \in (0, 1)$ . Then under the null hypothesis, we have*

$$T := \frac{\mathcal{L} - p\ell_n - \mu_n}{\nu_n} \xrightarrow{D} N(0, 1), \quad (2.1)$$

where

$$\ell_n = \log \left( \frac{y_1^{c_2} y_2^{c_1} h^{\frac{2h^2}{y_1 y_2}}}{(y_1 + y_2)^{\frac{(y_1 + y_2)}{y_1 y_2}} |1 - y_1|^{\frac{c_1|1 - y_1|}{y_1}} |1 - y_2|^{\frac{c_2|1 - y_2|}{y_2}}} \right) - l(y_1, y_2) - l(y_2, y_1),$$

$$\begin{aligned}\mu_n = & \log \left[ \frac{(y_1 + y_2)^{\frac{1}{2}} |1 - y_1|^{\frac{c_1}{2}} |1 - y_2|^{\frac{c_2}{2}}}{h} \right] - u(y_1, y_2) - u(y_2, y_1) \\ & + \frac{\Delta_1 \Psi(y_1, y_2)}{2y_1 y_2^2 (y_1 + y_2)^2} + \frac{\Delta_2 \Psi(y_2, y_1)}{2y_2 y_1^2 (y_1 + y_2)^2},\end{aligned}$$

and

$$\begin{aligned}\nu_n^2 = & \log \frac{h^4}{|1 - y_1|^{2c_1^2} |1 - y_2|^{2c_2^2} (y_1 + y_2)^2} + 2[v(y_1, y_2) + v(y_2, y_1) + \log(h^{4c_1 c_2}) \delta_{y_1 > 1} \delta_{y_2 > 1}] \\ & + \frac{(y_1 \Delta_1 + y_2 \Delta_2)}{y_1^2 y_2^2 (y_1 + y_2)^2} [(y_1 - 1) y_2^2 \delta_{y_1 > 1} - (y_2 - 1) y_1^2 \delta_{y_2 > 1}]^2.\end{aligned}$$

According to the above theorem, we can easily conclude the following corollary under normal circumstances:

**Corollary 2.2.** *If  $z_j^{(l)}$ ,  $j = 1 \dots, n_l$ ,  $l = 1, 2$  are normally distributed, then (2.1) in Theorem 2.1 reduces to:*

$$T := \frac{\mathcal{L} - p\ell_n - \mu_n}{\nu_n} \xrightarrow{D} N(0, 1),$$

where

$$\begin{aligned}\ell_n = & \log \left( \frac{y_1^{c_2} y_2^{c_1} h^{\frac{2h^2}{y_1 y_2}}}{(y_1 + y_2)^{\frac{(y_1 + y_2)}{y_1 y_2}} |1 - y_1|^{\frac{c_1 |1 - y_1|}{y_1}} |1 - y_2|^{\frac{c_2 |1 - y_2|}{y_2}}} \right) - l(y_1, y_2) - l(y_2, y_1), \\ \mu_n = & \log \left[ \frac{(y_1 + y_2)^{\frac{1}{2}} |1 - y_1|^{\frac{c_1}{2}} |1 - y_2|^{\frac{c_2}{2}}}{h} \right] - u(y_1, y_2) - u(y_2, y_1),\end{aligned}$$

and

$$\nu_n^2 = \log \frac{h^4}{|1 - y_1|^{2c_1^2} |1 - y_2|^{2c_2^2} (y_1 + y_2)^2} + 2[v(y_1, y_2) + v(y_2, y_1) + \log(h^{4c_1 c_2}) \delta_{y_1 > 1} \delta_{y_2 > 1}].$$

**Remark 2.3.** *If the Dimensions Assumption in Theorem 2.1 is satisfied, the limit condition that  $\lim y_1 \notin \{0, 1\}$ ,  $\lim y_2 \notin \{0, 1\}$  and  $\lim p/(n_1 + n_2) \in (0, 1)$  could be considered to hold,*

because there is no information for the convergence of its dimensions and size for any dataset. In addition, Jiang and Yang (2013) showed that if  $p/n_l \rightarrow 1$ , Theorem 2.1 also holds for normally distributed data. However, if  $y_l$  is near 1, then the variance  $\nu_n$  will be large and the LRT will become unstable, see Figure 1 for illustration.

**Remark 2.4.** When  $y_1 < 1$  and  $y_2 < 1$ , Corollary 2.2 recovers Theorem 4.1 in (Bai et al., 2009) directly.

The optimality of the Dimensions Assumption is clear because of the definitions of  $\ell_n$ ,  $\mu_n$  and  $\nu_n$ . For the Moments Assumption, we only need to consider the fourth moments of the sample. From the variance  $\nu_n$  in Theorem 2.1, we know that its fourth-moment term cannot be removed except  $y_1 < 1$  and  $y_2 < 1$ . However, if  $y_1 < 1$  and  $y_2 < 1$ , it is not difficult to obtain that

$$\Psi(y_2, y_1) = \Psi(y_1, y_2) = -y_1^3 y_2^3,$$


which implies that the fourth-moment term of  $\mu_n$  cannot be removed. Therefore, we conclude that the existence of the fourth moment of the sample is necessary for the modified LRT statistic  $\mathcal{L}$ . However, when  $y_1$  or  $y_2$  is close to 1, the variance  $\nu_n$  will increase rapidly, resulting in poor power. For illustration, we present two 3D figures of  $\mu_n$  and  $\nu_n^2$  with  $\Delta_1 = \Delta_2 = 0$  in Figure 1.

Now, we give the asymptomatic distribution of  $\tilde{\mathcal{L}}$ .

**Theorem 2.5.** In addition to the Moments Assumption and Dimensions Assumption, we assume that as  $\min\{p, n_1, n_2\} \rightarrow \infty$ ,  $\lim y_1 \neq 1$ ,  $\lim y_2 \neq 1$  and  $\lim p/(n_1 + n_2) < 1$ . Then under the null hypothesis, we have

$$\tilde{T} := \frac{\tilde{\mathcal{L}} - p\tilde{\ell}_n - \tilde{\mu}_n}{\tilde{\nu}_n} \xrightarrow{D} N(0, 1),$$





(a) 3D figure of the mean  $\mu_n$ .

(b) 3D figure of the variance  $\nu_n^2$ .

Figure 1: This figure was made using Maple software with  $y_1 \in (0, 2)$  and  $y_2 \in (0, 2)$ . The vertical axes present the values of  $\mu_n$  and  $\nu_n$ , respectively.

where

$$\tilde{\ell}_n = \log \left( \frac{y_2 h^{\frac{2h^2}{y_1 y_2}}}{(y_1 + y_2)^{\frac{(y_1 + y_2)}{y_1 y_2}} |1 - y_1|^{\frac{|1 - y_1|}{y_1}}} \right) - \log \left( \frac{h^{\frac{2h^2}{y_1 y_2}}}{\frac{1 + y_2}{y_1} \frac{1 - y_1}{y_2}} \right) \delta_{y_1 > 1}$$

$$\begin{aligned} \tilde{\mu}_n = & \log \left[ \frac{(y_1 + y_2)^{\frac{1}{2}} |1 - y_1|^{\frac{1}{2}}}{h} \right] - \log \left( \frac{y_1}{h} \right) \delta_{y_1 > 1} \\ & - \frac{\Delta_1 [y_1^3 (y_1 + 2y_2) \delta_{y_1 < 1} + h^2 (y_1 + y_2 + y_1 y_2) \delta_{y_1 > 1}]}{2y_1 (y_1 + y_2)^2} + \frac{\Delta_2 [y_1^4 y_2 \delta_{y_1 < 1} + h^2 y_2 (2y_1^2 - h^2) \delta_{y_1 > 1}]}{2y_1^2 (y_1 + y_2)^2}, \end{aligned}$$

and

$$\tilde{\nu}_n^2 = 2 \log \frac{h^2}{|1 - y_1| (y_1 + y_2)} + 2 \log \left( \frac{y_1^2}{h^2} \right) \delta_{y_1 > 1} + \frac{(y_1 \Delta_1 + y_2 \Delta_2)}{y_1^2 (y_1 + y_2)^2} [y_1^4 \delta_{y_1 < 1} + h^4 \delta_{y_1 > 1}].$$

**Remark 2.6.** Notice that the asymptotic distributions in Theorem 2.1 and Theorem 2.5 are obtained under the null hypothesis, which can only guarantee the Type I errors. For the powers, that is under the alternative hypothesis  $\Sigma_1 \neq \Sigma_2$ , the asymptotic distributions of

statistics  $\mathcal{L}$  and  $\tilde{\mathcal{L}}$  will depend on the eigenvalues of  $\Sigma_1 \Sigma_2^{-1}$ . In this case, if the dimension  $p$  is smaller than either of the two sample sizes, the power functions for  $T$  and  $\tilde{T}$  can be obtained by the CLT of the general Fisher matrices which is derived by Zheng et al. (2017). However, if  $p$  is bigger than both of the two sample sizes, because of the lack of theoretical results about the general Beta matrix  $n_1 \mathbf{S}_1 (n_1 \mathbf{S}_1 + n_2 \mathbf{A}^{1/2} \mathbf{S}_2 \mathbf{A}^{1/2})^{-1}$ , the asymptotic distributions of statistics  $\mathcal{L}$  and  $\tilde{\mathcal{L}}$  are also open problems and will be left for our future work. Here  $\mathbf{A}$  is any non-random symmetric matrix.

If  $\Delta_l \neq 0$ , or more specifically, if the samples are not normally distributed, the estimates for  $\Delta_l$  are necessary for the test application. Thus, we obtain their consistent estimators using the method of moments and random matrix theory. Let

$$\hat{\Delta}_1 = (1 - y)^2 \frac{\sum_{j=1}^{N_1} [(\mathbf{z}_j^{(1)} - \bar{\mathbf{z}}^{(1)})' (c_{11} \mathbf{S}_{1j}^{\mathbf{z}} + c_{12} \mathbf{S}_2^{\mathbf{z}})^{-1} (\mathbf{z}_j^{(1)} - \bar{\mathbf{z}}^{(1)}) - \frac{p}{1-y}]^2}{pN_1} - \frac{2}{1-y} \quad (2.2)$$

$$\hat{\Delta}_2 = (1 - y)^2 \frac{\sum_{j=1}^{N_2} [(\mathbf{z}_j^{(2)} - \bar{\mathbf{z}}^{(2)})' (c_{21} \mathbf{S}_1^{\mathbf{z}} + c_{22} \mathbf{S}_{2j}^{\mathbf{z}})^{-1} (\mathbf{z}_j^{(2)} - \bar{\mathbf{z}}^{(2)}) - \frac{p}{1-y}]^2}{pN_2} - \frac{2}{1-y}, \quad (2.3)$$

where  $y = \frac{p}{n_1 + n_2 - 1}$ ,  $c_{11} = \frac{n_1 - 1}{n_1 + n_2 - 1}$ ,  $c_{12} = \frac{n_2}{n_1 + n_2 - 1}$ ,  $c_{21} = \frac{n_1}{n_1 + n_2 - 1}$ ,  $c_{22} = \frac{n_2 - 1}{n_1 + n_2 - 1}$  and  $\mathbf{S}_{lj}^{\mathbf{z}}$  is the sample covariance matrix by removing the vector  $\mathbf{z}_j^{(l)}$  from the  $l$ -th sample,  $l = 1, 2$ .

**Theorem 2.7.** *Under the same assumptions of Theorem 2.1 and under the null hypothesis, we have the estimators  $\hat{\Delta}_l$ ,  $l = 1, 2$ , defined in (2.2) and (2.3) are weakly consistent and asymptotically unbiased.*

**Remark 2.8.** *Actually we use  $(c_{11} \mathbf{S}_{1j}^{\mathbf{z}} + c_{12} \mathbf{S}_2^{\mathbf{z}})^{-1}$  here is to avoid the case where the sample covariance matrices when the sample sizes are not larger than the dimension. Otherwise, if the dimension  $p$  is smaller than some sample size, say,  $p < n_1$ , we can estimate  $\Delta_1$  by replacing the term  $(c_{11} \mathbf{S}_{1j}^{\mathbf{z}} + c_{12} \mathbf{S}_2^{\mathbf{z}})^{-1}$  and  $y = \frac{p}{n_1 + n_2 - 1}$  in (2.2) by  $(\mathbf{S}_{1j}^{\mathbf{z}})^{-1}$  and  $y = \frac{p}{n_1 - 1}$  respectively.*

The proofs of Theorem 2.1, Theorem 2.5 and Theorem 2.7 are given in the appendix.

### 3 Results of the simulation

In this section, we compare the performance of the statistics proposed in Li and Chen (2012) and Cai et al. (2013) and our modified LRTs  $\mathcal{L}$  and  $\tilde{\mathcal{L}}$  under various settings of sample size and dimensionality. The classical LRT statistic in (Wilks, 1946) was shown to have poor performance for (1.1) by Bai et al. (2009); thus, it will not be considered in this section. Without loss of generality, we assume  $\boldsymbol{\mu}_l = 0$  and set

$$\Sigma_1 = (1 + a/n_1)\Sigma_2,$$

where  $a$  is a constant. The samples are drawn from the following distributions:

Case 1:  $\mathbf{x}^{(1)}$  and  $\mathbf{x}^{(2)}$  are both standard normal distributed and  $\Sigma_2 = \mathbf{I}_p$ ;

Case 2:  $\mathbf{x}^{(1)}$  and  $\mathbf{x}^{(2)}$  are from the uniform distribution  $U(-\sqrt{3}, \sqrt{3})$  and  $\Sigma_2 = \mathbf{I}_p$ ;

Case 3:  $\mathbf{x}^{(1)}$  and  $\mathbf{x}^{(2)}$  are from the uniform distribution  $U(-\sqrt{3}, \sqrt{3})$  and

$$\Sigma_2 = \text{Diag}(p^2, 1, \dots, 1);$$

Case 4:  $\mathbf{x}^{(1)}$  and  $\mathbf{x}^{(2)}$  are from the uniform distribution  $U(-\sqrt{3}, \sqrt{3})$  and

$$\Sigma_2 = (0.5\mathbf{I}_p + 0.5\mathbf{1}_p\mathbf{1}_p').$$

Here,  $\mathbf{1}_p$  represents a  $p$ -dimensional vector with all entries 1. The results are obtained based on 10,000 replicates. In the tables,  $T$  and  $\tilde{T}$  denote the proposed modified LRTs,  $T_{lc}$  denotes the nonparametric test of Li and Chen (2012) and  $T_{clx}$  denotes the maximum element test of Cai et al. (2013).

In the first part of this section, we report the results by assuming the forth moments of the  $\mathbf{x}^{(1)}$  and  $\mathbf{x}^{(2)}$  are known. Tables 1- 4 present the empirical sizes and empirical powers of Cases 1- 4. Additionally, we provide four figures (Figures 2 - 5) to show the divergence of

powers of the four test statistics as the parameter  $a$  increases. The results indicate that the sizes  $T$  and  $\tilde{T}$  perform quite well for all cases. However, in Case 3 and Case 4, the sizes of  $T_{lc}$  and  $T_{clx}$  are not accurate, which reflects the fact that the null distributions of the test statistics  $T_{lc}$  and  $T_{clx}$  are not well approximated by their asymptotic distributions in these cases. For the empirical power, we can conclude that  $T$  and  $\tilde{T}$  are more sensitive than  $T_{lc}$  and  $T_{clx}$  when at least one of the sample sizes is smaller than the dimensions. Otherwise, the LRT  $T$  does not perform as well when the dimensions are large. However, the lite LRT  $\tilde{T}$  is always powerful because of its monotonicity for matrix  $\mathbf{B}_n$ , which coincides with our intuition.

Next we will show the performance of the estimator we proposed in Theorem 2.7. Here we have to explain the reason that why we did not use the estimators in last simulation results. That is because our estimator is based on moment method, and need  $n = n_1 + n_2$  times loop and inverse process for one replication, and we need 10,000 replications for one result, that makes the running times to be terrible. In the simulation, we set  $x_{ij}$  be standard normal distributed and uniform distributed on  $(-\sqrt{3}, \sqrt{3})$  to estimate the  $\Delta_1$  respectively. Under each circumstance we repeat 10,000 times and the results are reported at Tables 5 and 6. From the numerical results, the performance of the estimator is remarkable, especially when the sample size is large. Therefore, we believe that the proposed modified LRTs must be also perform good at the null hypothesis when using the estimators instead of their true values. But, under the alternative, we can not make any arbitrary decision right now because of the less theoretical results about the general Beta matrix.

## 4 An example

For illustration, we apply the proposed test statistics to the daily returns of a selection of stocks issued by companies on Standard & Poor's (S&P) 500. The original data are

$(n_1, n_2, p)$ $y_1 > 1, y_2 > 1$		(25,35,40)			(50,70,80)			(100,140,160)			(200,280,320)		
		size		power	size		power	size		power	size		power
		$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$		0.055	0.951	1	0.053	0.951	1	0.054	0.951	1	0.051	0.946	1
$\tilde{T}$		0.056	0.999	1	0.048	1	1	0.048	1	1	0.049	1	1
$T_{lc}$		0.074	0.804	1	0.057	0.489	0.999	0.054	0.208	0.954	0.054	0.093	0.571
$T_{clx}$		0.082	0.152	0.591	0.057	0.071	0.343	0.048	0.049	0.120	0.042	0.046	0.062
$(n_1, n_2, p)$ $y_1 > 1, y_2 < 1$		(25,35,30)			(50,70,60)			(100,140,120)			(200,280,240)		
		size		power	size		power	size		power	size		power
		$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$		0.059	0.685	0.999	0.057	0.618	0.998	0.054	0.558	0.994	0.052	0.523	0.987
$\tilde{T}$		0.058	0.997	1	0.053	0.999	1	0.050	0.999	1	0.049	1	1
$T_{lc}$		0.063	0.793	1	0.055	0.494	0.999	0.053	0.205	0.952	0.050	0.098	0.574
$T_{clx}$		0.081	0.154	0.621	0.054	0.076	0.380	0.049	0.048	0.147	0.045	0.046	0.068
$(n_1, n_2, p)$ $y_1 < 1, y_2 > 1$		(35,25,30)			(70,50,60)			(140,100,120)			(280,200,240)		
		size		power	size		power	size		power	size		power
		$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$		0.060	0.109	0.075	0.055	0.167	0.282	0.050	0.213	0.515	0.050	0.248	0.645
$\tilde{T}$		0.056	0.956	1	0.052	0.985	1	0.052	0.989	1	0.046	0.994	1
$T_{lc}$		0.068	0.314	0.992	0.059	0.143	0.896	0.054	0.077	0.489	0.051	0.054	0.179
$T_{clx}$		0.077	0.203	0.677	0.054	0.102	0.433	0.049	0.061	0.189	0.042	0.051	0.076
$(n_1, n_2, p)$ $y_1 < 1, y_2 < 1$		(25,35,20)			(50,70,40)			(100,140,80)			(200,280,160)		
		size		power	size		power	size		power	size		power
		$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T_1$		0.060	0.251	0.960	0.056	0.119	0.715	0.050	0.079	0.330	0.049	0.053	0.139
$\tilde{T}$		0.057	0.996	1	0.052	0.999	1	0.049	1	1	0.050	0.999	1
$T_{lc}$		0.069	0.783	1	0.059	0.476	0.999	0.051	0.211	0.948	0.049	0.095	0.563
$T_{clx}$		0.076	0.169	0.679	0.058	0.083	0.438	0.046	0.053	0.180	0.047	0.050	0.072

Table 1: Empirical sizes and empirical powers of the tests  $T$ ,  $\tilde{T}$ ,  $T_{lc}$  and  $T_{clx}$  in Case 1.

These results are based on the 5% significance level.




Figure 2: Graphs of the divergence of the four powers in Case 1.




Figure 3: Graphs of the divergence of the four powers in Case 2.

$(n_1, n_2, p)$ $y_1 > 1, y_2 > 1$	(25,35,40)			(50,70,80)			(100,140,160)			(200,280,320)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.052	0.961	1	0.046	0.953	1	0.046	0.950	1	0.048	0.948	1
$\tilde{T}$	0.054	1	1	0.050	1	1	0.049	1	1	0.048	1	1
$T_{lc}$	0.055	0.815	1	0.056	0.486	0.999	0.050	0.209	0.959	0.050	0.093	0.579
$T_{clx}$	0.191	0.858	1	0.125	0.567	1	0.086	0.207	0.996	0.070	0.095	0.663
$(n_1, n_2, p)$ $y_1 > 1, y_2 < 1$	(25,35,30)			(50,70,60)			(100,140,120)			(200,280,240)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.056	0.701	0.999	0.050	0.610	0.998	0.047	0.563	0.994	0.053	0.517	0.986
$\tilde{T}$	0.057	0.999	1	0.052	1	1	0.051	1	1	0.050	1	1
$T_{lc}$	0.061	0.818	1	0.053	0.481	0.999	0.050	0.205	0.956	0.050	0.093	0.573
$T_{clx}$	0.163	0.855	1	0.112	0.581	1	0.087	0.228	0.997	0.068	0.095	0.679
$(n_1, n_2, p)$ $y_1 < 1, y_2 > 1$	(35,25,30)			(70,50,60)			(140,100,120)			(280,200,240)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.056	0.108	0.071	0.052	0.179	0.272	0.058	0.225	0.503	0.055	0.247	0.654
$\tilde{T}$	0.058	0.977	1	0.053	0.993	1	0.053	0.997	1	0.051	0.998	1
$T_{lc}$	0.058	0.307	0.995	0.050	0.137	0.904	0.052	0.072	0.488	0.049	0.051	0.177
$T_{clx}$	0.169	0.755	1	0.117	0.409	0.998	0.087	0.156	0.858	0.066	0.081	0.303
$(n_1, n_2, p)$ $y_1 < 1, y_2 < 1$	(25,35,20)			(50,70,40)			(100,140,80)			(200,280,160)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.053	0.245	0.976	0.055	0.117	0.724	0.050	0.072	0.330	0.051	0.055	0.134
$\tilde{T}$	0.061	0.999	1	0.056	1	1	0.050	1	1	0.047	1	1
$T_{lc}$	0.052	0.813	1	0.057	0.490	0.999	0.052	0.204	0.957	0.051	0.093	0.568
$T_{clx}$	0.146	0.845	1	0.106	0.583	1	0.078	0.237	0.996	0.064	0.098	0.706

Table 2: Empirical sizes and empirical powers of the tests  $T$ ,  $\tilde{T}$ ,  $T_{lc}$  and  $T_{clx}$  in Case 2. These results are based on the 5% significance level.

$(n_1, n_2, p)$ $y_1 > 1, y_2 > 1$	(25,35,40)			(50,70,80)			(100,140,160)			(200,280,320)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.051	0.960	1	0.046	0.954	1	0.046	0.949	1	0.048	0.949	1
$\tilde{T}$	0.054	1	1	0.050	1	1	0.049	1	1	0.048	1	1
$T_{lc}$	0.020	0.618	0.982	0.017	0.409	0.934	0.016	0.219	0.746	0.016	0.106	0.455
$T_{clx}$	0.191	0.861	1	0.125	0.568	1	0.086	0.210	0.996	0.070	0.095	0.662
$(n_1, n_2, p)$ $y_1 > 1, y_2 < 1$	(25,35,30)			(50,70,60)			(100,140,120)			(200,280,240)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.056	0.702	0.999	0.051	0.610	0.998	0.046	0.564	0.994	0.052	0.517	0.986
$\tilde{T}$	0.058	0.999	1	0.053	1	1	0.052	1	1	0.049	1	1
$T_{lc}$	0.019	0.613	0.983	0.011	0.407	0.934	0.016	0.216	0.755	0.015	0.103	0.458
$T_{clx}$	0.162	0.856	1	0.112	0.579	1	0.087	0.229	0.997	0.068	0.094	0.679
$(n_1, n_2, p)$ $y_1 < 1, y_2 > 1$	(35,25,30)			(70,50,60)			(140,100,120)			(280,200,240)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.057	0.106	0.071	0.052	0.179	0.274	0.057	0.223	0.502	0.054	0.249	0.648
$\tilde{T}$	0.060	0.978	1	0.053	0.992	1	0.053	0.997	1	0.052	0.998	1
$T_{lc}$	0.019	0.313	0.869	0.016	0.185	0.667	0.018	0.098	0.408	0.017	0.056	0.220
$T_{clx}$	0.168	0.757	1	0.117	0.407	0.999	0.088	0.157	0.855	0.067	0.081	0.305
$(n_1, n_2, p)$ $y_1 < 1, y_2 < 1$	(25,35,20)			(50,70,40)			(100,140,80)			(200,280,160)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.053	0.250	0.975	0.054	0.119	0.725	0.050	0.071	0.327	0.052	0.054	0.136
$\tilde{T}$	0.060	0.999	1	0.056	1	1	0.050	1	1	0.048	1	1
$T_{lc}$	0.019	0.629	0.983	0.016	0.403	0.930	0.015	0.212	0.752	0.015	0.113	0.445
$T_{clx}$	0.147	0.847	1	0.106	0.581	1	0.079	0.237	0.996	0.063	0.098	0.706

Table 3: Empirical sizes and empirical powers of the tests  $T$ ,  $\tilde{T}$ ,  $T_{lc}$  and  $T_{clx}$  in Case 3.

These results are based on the 5% significance level.






Figure 4: Graphs of the divergence of the four powers in Case 3.




Figure 5: Graphs of the divergence of the four powers in Case 4.

$(n_1, n_2, p)$ $y_1 > 1, y_2 > 1$	(25,35,40)			(50,70,80)			(100,140,160)			(200,280,320)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.052	0.961	1	0.046	0.954	1	0.045	0.950	1	0.047	0.949	1
$\tilde{T}$	0.052	1	1	0.049	1	1	0.049	1	1	0.048	1	1
$T_{lc}$	0.083	0.505	0.909	0.081	0.348	0.793	0.087	0.240	0.587	0.082	0.179	0.386
$T_{clx}$	0.062	0.207	0.778	0.044	0.076	0.492	0.030	0.032	0.187	0.022	0.020	0.068
$(n_1, n_2, p)$ $y_1 > 1, y_2 < 1$	(25,35,30)			(50,70,60)			(100,140,120)			(200,280,240)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.055	0.701	0.999	0.050	0.611	0.998	0.046	0.565	0.994	0.053	0.518	0.987
$\tilde{T}$	0.057	0.999	1	0.053	1	1	0.052	1	1	0.049	1	1
$T_{lc}$	0.089	0.517	0.918	0.083	0.359	0.786	0.083	0.235	0.596	0.081	0.172	0.395
$T_{clx}$	0.068	0.230	0.818	0.043	0.085	0.539	0.031	0.036	0.221	0.021	0.023	0.078
$(n_1, n_2, p)$ $y_1 < 1, y_2 > 1$	(35,25,30)			(70,50,60)			(140,100,120)			(280,200,240)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.057	0.107	0.069	0.051	0.178	0.276	0.057	0.224	0.506	0.054	0.248	0.649
$\tilde{T}$	0.060	0.979	1	0.054	0.993	1	0.053	0.997	1	0.051	0.998	1
$T_{lc}$	0.084	0.310	0.736	0.083	0.229	0.563	0.081	0.170	0.375	0.084	0.125	0.247
$T_{clx}$	0.065	0.272	0.777	0.042	0.145	0.487	0.029	0.084	0.244	0.021	0.048	0.112
$(n_1, n_2, p)$ $y_1 < 1, y_2 < 1$	(25,35,20)			(50,70,40)			(100,140,80)			(200,280,160)		
	size		power	size		power	size		power	size		power
	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$	$a=0$	$a=10$	$a=20$
$T$	0.052	0.245	0.975	0.055	0.119	0.724	0.050	0.072	0.329	0.051	0.055	0.135
$\tilde{T}$	0.061	0.999	1	0.056	1	1	0.050	1	1	0.048	1	1
$T_{lc}$	0.084	0.518	0.924	0.079	0.362	0.792	0.083	0.249	0.595	0.081	0.172	0.398
$T_{clx}$	0.072	0.273	0.862	0.043	0.107	0.608	0.032	0.044	0.261	0.021	0.025	0.092

Table 4: Empirical sizes and empirical powers of the tests  $T$ ,  $\tilde{T}$ ,  $T_{lc}$  and  $T_{clx}$  in Case 4.

These results are based on the 5% significance level.

(200,280)			(400,560)			(800,1120)		
p	mean	variance	p	mean	variance	p	mean	variance
2	0.0186	0.1368	4	0.0099	0.0441	8	0.0049	0.0160
10	0.0197	0.0642	20	0.0081	0.0260	40	0.0034	0.0116
20	0.0193	0.0563	40	0.0107	0.0252	80	0.0040	0.0116
100	0.0310	0.0732	200	0.0121	0.0338	400	0.0057	0.0166
180	0.0552	0.1189	360	0.0223	0.0555	720	0.0135	0.0261
220	0.0719	0.1634	440	0.0332	0.0736	880	0.0137	0.0355
240	0.0739	0.1887	480	0.0395	0.0870	960	0.0190	0.0424
300	0.1491	0.3568	600	0.0695	0.1598	1200	0.0373	0.0771

Table 5: Numerical results for the estimator  $\hat{\Delta}_1$  of standard normal distribution. Here the true value  $\Delta_1 = 0$ .

(200,280)			(400,560)			(800,1120)		
p	mean	variance	p	mean	variance	p	mean	variance
2	-1.2021	0.0058	4	-1.2020	0.0032	8	-1.2012	0.0015
10	-1.2009	0.0073	20	-1.2002	0.0036	40	-1.2018	0.0019
20	-1.2005	0.0081	40	-1.2001	0.0039	80	-1.1994	0.0020
100	-1.1971	0.0179	200	-1.1977	0.0091	400	-1.2001	0.0044
180	-1.1858	0.0450	360	-1.1892	0.0213	720	-1.1957	0.0105
220	-1.1704	0.0721	440	-1.1894	0.0314	880	-1.1938	0.0159
240	-1.1650	0.0902	480	-1.1829	0.0417	960	-1.1892	0.0204
300	-1.1170	0.2170	600	-1.1537	0.0974	1200	-1.1760	0.0445

Table 6: Numerical results for the estimator  $\hat{\Delta}_1$  of uniform distribution  $(-\sqrt{3}, \sqrt{3})$ . Here the true value  $\Delta_1 = -1.2$ .

	Consumer Discretionary	Consumer Discretionary	Materials	Materials
$(n_1, n_2, p)$	(61,62,71)	(60,61,71)	(60,58,22)	(60,61,22)
$T$	0	0	0	0
$\tilde{T}$	0	$8.1307 * 10^{-8}$	0	0
$T_{lc}$	$9.4786 * 10^{-6}$	0	0	0
$T_{clx}$	$4.7868 * 10^{-3}$	$2.3442 * 10^{-4}$	$1.9239 * 10^{-6}$	$1.3389 * 10^{-3}$

Table 7:  $p$ -values of the test of equality of the two covariance matrices of daily returns from the same sector in different quarters.

the closing prices or the bid/ask average of these stocks for the trading days during 2012 and 2013. This dataset is derived from the Center for Research in Security Prices Daily Stock from Wharton Research Data Services. A common interest is to test whether the covariance matrices of the logarithmic daily returns for some stocks are the same over a period of time. Logarithmic daily returns are commonly used in finance. There are several theoretical and practical advantages of using logarithmic daily returns, including that we can assume that the sequences of logarithmic daily returns are independent of each other, see (Cont, 2001).

According to the Global Industry Classification Standard (GICS), which is an industry taxonomy developed in 1999 by MSCI and S&P for use by the global financial community, we select two sectors, Consumer Discretionary and Materials, which include 71 and 22 stocks, respectively. First, for Consumer Discretionary sector, we test whether the covariance matrices of the logarithmic daily returns of the second quarters (1 April-30 June) of 2012 and 2013 are the same. There were 63 and 64 trading days, respectively, in the second quarters of 2012 and 2013. By using the logarithmic difference transformation on the original data, we obtain the logarithmic daily returns dataset with sample sizes  $n_1 = 61$  and  $n_2 = 62$ . Simultaneously, we use the first quarters (1 January-31 March) data of 2012 and 2013 when applying to Materials sector. That makes the sample sizes be  $n_1 = 60$

and  $n_2 = 58$ . The  $p$  values obtained by applying the four test statistics  $T$ ,  $\tilde{T}$ ,  $T_{lc}$  and  $T_{clx}$  are shown in the first and third columns of Table 7. Next, we use the same procedure to test whether the covariance matrices of the logarithmic daily returns of the first quarter (1 January-31 March) and the second quarter (1 April-30 June) of 2012 are the same. The results are shown in the second and fourth columns of Table 7. The results show that all the  $p$ -values are much smaller than 0.05. Thus, there is strong evidence that the two covariance matrices are different. Therefore, we suggest caution with the assumption that the daily returns are identically distributed.

## 5 Conclusion and discussion

In this paper, we propose two modified LRTs for the equality of two high-dimensional covariance matrices and show the asymptotic distributions under the Moments Assumption and the null hypothesis. Furthermore, we show that the Moments Assumption and Dimensions Assumption are necessary for our results. More specifically, if the samples are Gaussian distributed, our modifications are optimal. We also present the weakly consistent and asymptotic unbiased estimators of  $\Delta_l$  for non-Gaussian distributions under the null hypothesis. According to the simulation results, the performances of these modified LRTs are remarkable under conditions where they are applicable, but the theoretical results for the alternative hypothesis are not considered in this paper because of the lack of random matrix theory. The optimal modification of the LRT under the alternative hypothesis test will be presented in the future.

## Acknowledgement

J. Hu was partially supported by NSFC (No.11771073) and Science and Technology Development foundation of Jilin (No.20160520174JH). Z. D. Bai was partially supported by

## A Appendix

In this appendix, we provide the proofs of Theorem 2.5 and Theorem 2.7. By comparing the definitions of  $\mathcal{L}$  and  $\tilde{\mathcal{L}}$ , it is easy to verify that the proof of Theorem 2.1 can be split into two parts, one of which is the same as the proof of Theorem 2.5 and another of which is an analogous analysis. Hence, we present the proof Theorem 2.5 in this paper.

### A.1 Proof of Theorem 2.5

The main tool used to prove the theorems is the Cauchy integral formula and Theorem 1.6 in Bai et al. (2015), which established the central limit theorem of linear spectral statistics of random matrix  $\mathbf{B}_n$  and is presented below for convenience. Denote  $\alpha_n = n_2/n_1$ ,  $G_n(x) = p(\mathbf{F}^{\mathbf{B}_n}(x) - \mathbf{F}_{y_1, y_2}(x))$  and

$$\mathbf{F}_{y_1, y_2}(x) = \frac{(\alpha_n + 1)\sqrt{(x_r - x)(x - x_l)}}{2\pi y_1 x(1 - x)} \delta_{x \in (x_l, x_r)}$$

$x_l, x_r = \frac{y_2(h \mp y_1)^2}{(y_1 + y_2)^2}$  be the limit spectral distribution of  $\mathbf{B}_n$  with parameters  $\alpha_n, y_1, y_2$ ,

**Lemma A.1** (Theorem 1.6 in Bai et al. (2015)). *In addition to the Moments Assumption and the Dimensions Assumption, we further assume that:*

(1) As  $\min\{p, n_1, n_2\} \rightarrow \infty$ ,  $y_1 \rightarrow \gamma_1 \in (0, 1) \cup (1, \infty)$ ,  $y_2 \rightarrow \gamma_2 \in (0, 1) \cup (1, \infty)$ , and  $\alpha_n \rightarrow \alpha > 0$ .

(2) Let  $f_1, \dots, f_k$  be the analytic functions on an open region containing the interval  $[a_l, a_r]$ , where  $a_l = v^{-1}(1 - \sqrt{\gamma_1})^2$ ,  $a_r = 1 - \alpha v^{-1}(1 - \sqrt{\gamma_2})^2$ , and  $v$  is defined as  $v = (1 + \frac{\gamma_1}{\gamma_2})(1 - \sqrt{\frac{\gamma_1 \gamma_2}{\gamma_1 + \gamma_2}})^2$ .

Then, as  $\min(n_1, n_2, p) \rightarrow \infty$ , the random vector

$$(\int f_i dG_n(x)), \quad i = 1, \dots, k,$$

converges weakly to a Gaussian vector  $(G_{f_1}, \dots, G_{f_k})$  with mean functions

$$\begin{aligned}\mathbb{E}G_{f_i} &= \frac{1}{4\pi i} \oint f_i\left(\frac{z}{\alpha+z}\right) d\log\left(\frac{(1-\gamma_2)m_3^2(z) + 2m_3(z) + 1 - \gamma_1}{(1+m_3(z))^2}\right) \\ &\quad + \frac{\Delta_1}{2\pi i} \oint \gamma_1 f_i\left(\frac{z}{\alpha+z}\right) (1+m_3)^{-3} dm_3(z) \\ &\quad + \frac{\Delta_2}{4\pi i} \oint f_i\left(\frac{z}{\alpha+z}\right) (1-\gamma_2 m_3^2(z)(1+m_3(z))^{-2}) d\log(1-\gamma_2 m_3^2(z)(1+m_3(z))^{-2})\end{aligned}$$

and covariance functions

$$\begin{aligned}\mathbf{Cov}(G_{f_i}, G_{f_j}) &= -\frac{1}{2\pi^2} \oint \oint \frac{f_i\left(\frac{z_1}{\alpha+z_1}\right) f_j\left(\frac{z_2}{\alpha+z_2}\right) dm_3(z_1) dm_3(z_2)}{(m_3(z_1) - m_3(z_2))^2} \\ &\quad - \frac{\gamma_1 \Delta_1 + \gamma_2 \Delta_2}{4\pi^2} \oint \oint \frac{f_i\left(\frac{z_1}{\alpha+z_1}\right) f_j\left(\frac{z_2}{\alpha+z_2}\right) dm_3(z_1) dm_3(z_2)}{(m_3(z_1) + 1)^2 (m_3(z_2) + 1)^2},\end{aligned}$$

where

$$\begin{aligned}m_0(z) &= \frac{(1+\gamma_1)(1-z) - \alpha z(1-\gamma_2) + \sqrt{((1-\gamma_1)(1-z) + \alpha z(1-\gamma_2))^2 - 4\alpha z(1-z)}}{2z(1-z)(\gamma_1(1-z) + \alpha z\gamma_2)} - \frac{1}{z}, \\ m_1(z) &= \frac{\alpha}{(\alpha+z)^2} m_0\left(\frac{z}{\alpha+z}\right) - \frac{1}{\alpha+z}, \quad m_2(z) = -z^{-1}(1-\gamma_1) + \gamma_1 m_1(z), \\ m_{mp}^{\gamma_2}(z) &= \frac{1-\gamma_2-z + \sqrt{(z-1-\gamma_2)^2 - 4\gamma_2}}{2\gamma_2 z}, \quad m_3(z) = \gamma_2 m_{mp}^{\gamma_2}(-m_2(z)) + (m_2(z))^{-1}(1-\gamma_2).\end{aligned}$$

All the above contour integrals can be evaluated on any contour enclosing the interval  $\left[\frac{\alpha c_l}{1-c_l}, \frac{\alpha c_r}{1-c_r}\right]$ .

Note that Lemma A.1 is proved based on the centralized sample covariance matrices, which are constructed by not subtracting the sample mean vector from each sample vector. However, Zheng et al. (2015) showed that the only difference between the two types of sample covariance matrices is normalization by  $p/n_l$  and  $p/N_l$ . It is not difficult to verify that Theorem 2.7 satisfies Lemma A.1 by choosing the kernel function to the logarithmic function. Therefore, the main task of proving Theorem 2.7 is to calculate the three integrals, which will be shown step by step.

*Proof of the limit part  $\tilde{\ell}_n$ .* When calculating the integral

$$p \int \log x \frac{(\alpha_n + 1) \sqrt{(x_r - x)(x - x_l)}}{2\pi y_1 x(1 - x)} \delta_{x \in (x_l, x_r)} dx \quad (\text{A.1})$$

to achieve the limit part  $\tilde{\ell}_n$ , we choose a transformation

$$x = \frac{y_2 |y_1 + h\xi|^2}{(y_1 + y_2)^2}.$$

Clearly, when  $x$  moves from  $\frac{y_2(h-y_1)^2}{(y_1+y_2)^2}$  to  $\frac{y_2(h+y_1)^2}{(y_1+y_2)^2}$  two times,  $\xi$  shifts along a unit circle in the positive direction. Then, we obtain that the integral (A.1) is equal to

$$p \frac{(y_1 + y_2)h^2 i}{4\pi} \oint [\log \frac{y_2 |y_1 + h\xi|^2}{(y_1 + y_2)^2}] \cdot \frac{(\xi^2 - 1)^2}{\xi^3 |y_1 + h\xi|^2 |y_2 - h\xi|^2} d\xi \quad (\text{A.2})$$

when  $y_1 > 1$  and

$$p \frac{(y_1 + y_2)h^2 i}{4\pi} \oint [\log \frac{y_2 |h + y_1 \xi|^2}{(y_1 + y_2)^2}] \cdot \frac{(\xi^2 - 1)^2}{\xi^3 |y_1 + h\xi|^2 |y_2 - h\xi|^2} d\xi \quad (\text{A.3})$$

when  $y_1 < 1$ . Two forms of the integral ensure that the logarithmic function returns a finite number of poles related to  $y_1$  of the integrand. The pole related to  $y_2$  of the integrand is  $h/y_2$  when  $y_2 > 1$ . There is no difference in the integral value before and after the transformation  $\xi = 1/\xi$ , except that the residue point in the unit disc turns into  $y_2/h$ , which is the residue point under the assumption  $y_2 < 1$ . Thus, we assume  $y_2 > 1$  without loss of generality. Then, we obtain that if  $y_1 > 1$ , (A.2) can be rewritten as

$$\begin{aligned} & p \frac{(y_1 + y_2)h^2 i}{4\pi} \oint [\log \frac{y_2(y_1 + h\xi)}{(y_1 + y_2)^2}] \times \frac{(\xi^2 - 1)^2}{\xi^3 |y_1 + h\xi|^2 |y_2 - h\xi|^2} d\xi \\ & + p \frac{(y_1 + y_2)h^2 i}{4\pi} \oint [\log \frac{y_2(y_1 + \frac{h}{\xi})}{(y_1 + y_2)^2}] \times \frac{(\xi^2 - 1)^2}{\xi^3 |y_1 + h\xi|^2 |y_2 - h\xi|^2} d\xi \end{aligned}$$

which is equal to

$$p \frac{(y_1 + y_2)h^2 i}{4\pi} \oint [\log \frac{y_2(y_1 + h\xi)^2}{(y_1 + y_2)^2}] \times \frac{(\xi^2 - 1)^2}{\xi^3 (y_1 + h\xi)(y_1 + \frac{h}{\xi})(y_2 - h\xi)(y_2 - \frac{h}{\xi})} d\xi.$$



Similarly, if  $y_1 < 1$ , we have (A.3) equals

$$p \frac{(y_1 + y_2)h^2 i}{4\pi} \oint [\log \frac{y_2(h + y_1\xi)^2}{(y_1 + y_2)^2}] \times \frac{(\xi^2 - 1)^2}{\xi^3(y_1 + h\xi)(y_1 + \frac{h}{\xi})(y_2 - h\xi)(y_2 - \frac{h}{\xi})} d\xi.$$

According to Cauchy's residue theorem, we find three poles

$$0, \quad -\frac{h}{y_1}, \quad \frac{h}{y_2}$$

under the settings of  $y_1 > 1, y_2 > 1$ . The corresponding residues are

$$\frac{p(y_1 + y_2)}{2y_1y_2} \log \frac{y_1^2 y_2}{(y_1 + y_2)^2}, \quad \frac{p(1 - y_1)}{2y_1} \log \frac{y_2(y_1 - 1)^2}{y_1^2}, \quad \frac{p(1 - y_2)}{2y_2} \log \frac{1}{y_2},$$

respectively. In the same way, under the assumptions  $y_1 < 1, y_2 > 1$ , we obtain three poles

$$0, \quad -\frac{y_1}{h}, \quad \frac{h}{y_2}$$

and three residues

$$\frac{p(y_1 + y_2)}{2y_1y_2} \log \frac{h^2 y_2}{(y_1 + y_2)^2}, \quad \frac{p(1 - y_1)}{2y_1} \log \frac{h^2}{y_2(1 - y_1)^2}, \quad \frac{p(1 - y_2)}{2y_2} \log \frac{h^2}{y_2}.$$

Therefore, by combining the above results and basic calculations, we obtain the limit part (A.1) is

$$\log \left( \frac{y_2 h^{\frac{2h^2}{y_1 y_2}}}{(y_1 + y_2)^{\frac{(y_1 + y_2)}{y_1 y_2}} |1 - y_1|^{\frac{|1 - y_1|}{y_1}}} \right) - \log \left( \frac{h^{\frac{2h^2}{y_1 y_2}}}{y_1^{\frac{1 + y_2}{y_2}} y_2^{\frac{1 - y_1}{y_1}}} \right) \delta_{y_1 > 1},$$

which completes the proof.  $\square$

*Proof of the mean part  $\tilde{\mu}_n$ .* Because  $m_3$  satisfies the equation

$$z = -\frac{m_3(z)(m_3(z) + 1 - y_1)}{(1 - y_2)(m_3(z) + \frac{1}{1 - y_2})},$$

we make an integral conversion  $z = (1 + hr\xi)(1 + \frac{h}{r\xi})/(1 - y_2)^2$ , where  $r$  is a number greater than but close to 1. According to the discussion in the last section, we assume  $y_2 > 1$  without loss of generality. From the equation

$$\frac{(1 + hr\xi)(1 + \frac{h}{r\xi})}{(1 - y_2)^2} = -\frac{m_3(m_3 + 1 - y_1)}{(1 - y_2)m_3 + 1},$$

we obtain that  $m_3 = -(1 + hr\xi)/(1 - y_2)$  or  $m_3 = -(1 + \frac{h}{r\xi})/(1 - y_2)$ . When  $z$  runs in the positive direction around the contour enclosing the interval  $[\frac{\alpha c_l}{1-c_l}, \frac{\alpha c_r}{1-c_r}]$ ,  $m_3$  runs in the opposite direction. Thus, when  $y_2 > 1$ , we choose the outcome  $m_3 = -(1 + \frac{h}{r\xi})/(1 - y_2)$ . Consequently, we have

$$\frac{z}{\alpha + z} = \frac{y_2|1 + hr\xi|^2}{|y_2 + hr\xi|^2}.$$

Therefore, for  $y_1 > 1$ , we get the mean part  $\tilde{\mu}_n$  is equal to

$$\lim_{r \downarrow 1} \frac{1}{4\pi i} \oint_{|\xi|=1} \left( \log \frac{y_2(1 + hr\xi)(1 + \frac{h}{r\xi})}{(y_2 + hr\xi)(y_2 + \frac{h}{r\xi})} \right) \times \left( \frac{1}{\xi - \frac{1}{r}} + \frac{1}{\xi + \frac{1}{r}} - \frac{2}{\xi + \frac{h}{y_2 r}} \right) d\xi \quad (\text{A.4})$$

$$+ \lim_{r \downarrow 1} \frac{\Delta_1}{2\pi i} \oint -y_1 \left( \log \frac{y_2(1 + hr\xi)(1 + \frac{h}{r\xi})}{(y_2 + hr\xi)(y_2 + \frac{h}{r\xi})} \right) \frac{(1 - y_2)^2 h}{y_2^3} \frac{\xi}{(\xi + \frac{h}{y_2 r})^3} d\xi \quad (\text{A.5})$$

$$+ \lim_{r \downarrow 1} \frac{\Delta_2}{4\pi i} \oint \left( \log \frac{y_2(1 + hr\xi)(1 + \frac{h}{r\xi})}{(y_2 + hr\xi)(y_2 + \frac{h}{r\xi})} \right) \frac{(y_2 - 1)(\xi^2 - \frac{h^2}{y_2 r^2})}{y_2(\xi + \frac{h}{y_2 r})^2} \left[ \frac{2\xi}{(\xi^2 - \frac{h^2}{y_2 r^2})} - \frac{2}{\xi + \frac{h}{y_2 r}} \right] d\xi. \quad (\text{A.6})$$

If  $y_1 < 1$ , we only need to change the logarithmic term in the above expression into the following form

$$\log \frac{y_2(1 + hr\xi)(1 + \frac{h}{r\xi})}{(h + y_2 r\xi)(h + \frac{y_2}{r\xi})},$$

and the other terms remain the same. For  $y_1 > 1$ , there are four poles of term (A.4)

$$\frac{1}{r}, \quad -\frac{1}{r}, \quad -\frac{h}{y_2 r}, \quad 0,$$

and four residues

$$\frac{1}{2} \log \frac{y_2(1 + h)}{y_2 + h}, \quad \frac{1}{2} \log \frac{y_2(1 - h)}{y_2 - h}, \quad \frac{1}{2} \log \frac{(y_1 + y_2)^2}{y_2^2 y_1^2}, \quad 0.$$

Thus, by Cauchy's residue theorem, we have

$$(\text{A.4}) = \frac{1}{2} \log \frac{(y_1 + y_2)(y_1 - 1)}{y_1^2}.$$

Analogously,

$$(A.5) = -\Delta_1 \frac{h^2(y_1 + y_2 + y_1 y_2)}{2y_1(y_1 + y_2)^2},$$

and

$$(A.6) = \Delta_2 \frac{h^2 y_2 (2y_1^2 - h^2)}{2y_1^2 (y_1 + y_2)^2}.$$

For  $y_1 < 1$ ,

$$\begin{aligned} & \lim_{r \downarrow 1} \frac{1}{4\pi i} \oint_{|\xi|=1} \left( \log \frac{y_2(1 + hr\xi)(1 + \frac{h}{r\xi})}{(h + y_2 r\xi)(h + \frac{y_2}{r\xi})} \right) \times \left( \frac{1}{\xi - \frac{1}{r}} + \frac{1}{\xi + \frac{1}{r}} - \frac{2}{\xi + \frac{h}{y_2 r}} \right) d\xi \\ &= \lim_{r \downarrow 1} \frac{1}{4\pi i} \oint_{|\xi|=1} \left( \log \frac{y_2(1 + \frac{h}{r\xi})}{(h + \frac{y_2}{r\xi})} \right) \times \left( \frac{1}{\xi - \frac{1}{r}} + \frac{1}{\xi + \frac{1}{r}} - \frac{2}{\xi + \frac{h}{y_2 r}} \right) d\xi \\ &+ \lim_{r \downarrow 1} \frac{1}{4\pi i} \oint_{|\xi|=1} \left( \log \frac{(\xi + hr)}{(h\xi + y_2 r)} \right) \times \frac{1}{\xi} \times \left( \frac{r}{r - \xi} + \frac{r}{r + \xi} - \frac{\frac{2y_2 r}{h}}{\frac{y_2 r}{h} + \xi} \right) d\xi \\ &= \frac{1}{2} \log \frac{(y_1 + y_2)(1 - y_1)}{h^2}, \\ & \lim_{r \downarrow 1} \frac{\Delta_1}{2\pi i} \oint -y_1 \left( \log \frac{y_2(1 + hr\xi)(1 + \frac{h}{r\xi})}{(h + y_2 r\xi)(h + \frac{y_2}{r\xi})} \right) \frac{(1 - y_2)^2 h}{y_2^3} \frac{\xi}{(\xi + \frac{h}{y_2 r})^3} d\xi \\ &= \Delta_1 \frac{y_1^2 (y_1 + 2y_2)}{-2(y_1 + y_2)^2}, \end{aligned}$$

and

$$\lim_{r \downarrow 1} \frac{\Delta_2}{4\pi i} \oint \left( \log \frac{y_2(1 + hr\xi)(1 + \frac{h}{r\xi})}{(h + y_2 r\xi)(h + \frac{y_2}{r\xi})} \right) \frac{(y_2 - 1)(\xi^2 - \frac{h^2}{y_2 r^2})}{y_2(\xi + \frac{h}{y_2 r})^2} \left[ \frac{2\xi}{(\xi^2 - \frac{h^2}{y_2 r^2})} - \frac{2}{\xi + \frac{h}{y_2 r}} \right] d\xi$$

$$= \Delta_2 \frac{y_1^2 y_2}{2(y_1 + y_2)^2}.$$

Thus, by combining the above results, the mean part  $\tilde{\mu}_n$  is

$$\begin{aligned} & \log \left[ \frac{(y_1 + y_2)^{\frac{1}{2}} |1 - y_1|^{\frac{c_1}{2}}}{h} \right] - \log\left(\frac{y_1}{h}\right) \delta_{y_1 > 1} \\ & - \frac{\Delta_1 [y_1^3 (y_1 + 2y_2) \delta_{y_1 < 1} + h^2 (y_1 + y_2 + y_1 y_2) \delta_{y_1 > 1}]}{2y_1 (y_1 + y_2)^2} + \frac{\Delta_2 [y_1^4 y_2 \delta_{y_1 < 1} + h^2 y_2 (2y_1 - h^2) \delta_{y_1 > 1}]}{2y_1^2 (y_1 + y_2)^2}. \end{aligned}$$

This completes the proof. □

*Proof of the variance part.* To calculate the variance part  $\tilde{\nu}_n$

$$- \frac{1}{2\pi^2} \oint \oint \frac{f(\frac{z_1}{\alpha + z_1}) f(\frac{z_2}{\alpha + z_2}) dm_3(z_1) dm_3(z_2)}{(m_3(z_1) - m_3(z_2))^2} \quad (\text{A.7})$$

$$- \frac{y_1 \Delta_1 + y_2 \Delta_2}{4\pi^2} \oint \oint \frac{f(\frac{z_1}{\alpha + z_1}) f(\frac{z_2}{\alpha + z_2}) dm_3(z_1) dm_3(z_2)}{(m_3(z_1) + 1)^2 (m_3(z_2) + 1)^2}, \quad (\text{A.8})$$

we make an analogous integral conversion

$$z_1 = (1 + hr_1 \xi_1) (1 + \frac{h}{r_1 \xi_1}) / (1 - y_2)^2$$

and

$$z_2 = (1 + hr_2 \xi_2) (1 + \frac{h}{r_2 \xi_2}) / (1 - y_2)^2.$$

Therefore, the relationship between  $\xi_l$  and  $m_3(z_l)$ ,  $l = 1, 2$  is

$$m_3(z_1) = -\frac{1 + \frac{h}{r_1 \xi_1}}{(1 - y_2)}, \quad m_3(z_2) = -\frac{1 + \frac{h}{r_2 \xi_2}}{(1 - y_2)}.$$

Without loss of generality, we assume  $r_1 < r_2$ . When  $y_1 > 1, y_2 > 1$ ,

$$- \frac{1}{2\pi^2} \oint \oint \frac{f_i(\frac{z_1}{\alpha + z_1}) f_j(\frac{z_2}{\alpha + z_2}) dm_3(z_1) dm_3(z_2)}{(m_3(z_1) - m_3(z_2))^2} = 2 \lim_{r_2 \downarrow 1} \oint \frac{1}{2\pi i} \left( \log \frac{y_2 (1 + hr_2 \xi_2) (1 + \frac{h}{r_2 \xi_2})}{(y_2 + hr_2 \xi_2) (y_2 + \frac{h}{r_2 \xi_2})} \right) \times$$

$$\left\{ \lim_{r_1 \downarrow 1} \oint \frac{1}{2\pi i} \left( \log \frac{y_2 (1 + hr_1 \xi_1) (1 + \frac{h}{r_1 \xi_1})}{(y_2 + hr_1 \xi_1) (y_2 + \frac{h}{r_1 \xi_1})} \right) \frac{r_1}{(r_1 \xi_1 - r_2 \xi_2)^2} d\xi_1 \right\} r_2 d\xi_2.$$

$$\begin{aligned}
&= 2 \lim_{r_2 \downarrow 1} \oint \frac{1}{2\pi i} \left( \log \frac{y_2(1 + hr_2\xi_2)(1 + \frac{h}{r_2\xi_2})}{(y_2 + hr_2\xi_2)(y_2 + \frac{h}{r_2\xi_2})} \right) \times \\
&\quad \left\{ \lim_{r_1 \downarrow 1} \oint \frac{1}{2\pi i} \left[ \left( \log \frac{y_2(1 + hr_1\xi_1)}{(y_2 + hr_1\xi_1)} \right) \frac{r_1}{(r_1\xi_1 - r_2\xi_2)^2} + \left( \log \frac{(1 + \frac{h}{r_1\xi_1})}{(y_2 + \frac{h}{r_1\xi_1})} \right) \frac{r_1}{r_2^2\xi_2^2(\frac{r_1}{r_2\xi_2} - \xi_1)^2} \right] d\xi_1 \right\} r_2 d\xi_2.
\end{aligned}$$

There is only one pole  $\frac{r_1}{r_2\xi_2}$  in the unit disc of the integration formula with respect to  $\xi_1$ .

Thus, by Cauchy's residue theorem, (A.7) is equal to

$$2 \lim_{r_2 \downarrow 1} \oint \frac{1}{2\pi i} \left( \log \frac{y_2(1 + hr_2\xi_2)(1 + \frac{h}{r_2\xi_2})}{(y_2 + hr_2\xi_2)(y_2 + \frac{h}{r_2\xi_2})} \right) \cdot \frac{1}{\xi_2} \left( \frac{h}{h + r_2\xi_2} - \frac{h}{h + y_2r_2\xi_2} \right) d\xi_2$$

which has three poles

$$0, \quad -\frac{h}{r_2}, \quad -\frac{h}{r_2y_2}.$$

Thus, we finally obtain that

$$(\text{A.7}) = \frac{2y_1^2}{(y_1 + y_2)(y_1 - 1)}.$$

Similarly, if  $y_1 < 1, y_2 > 1$ ,

$$(\text{A.7}) = \frac{2h^2}{(y_1 + y_2)(1 - y_1)}$$

In addition, as when  $y_1 > 1, y_2 > 1$ ,

$$\begin{aligned}
& -\frac{y_1\Delta_1 + y_2\Delta_2}{4\pi^2} \oint \oint \frac{f_i(\frac{z_1}{\alpha+z_1})f_j(\frac{z_2}{\alpha+z_2})dm_3(z_1)dm_3(z_2)}{(m_3(z_1) + 1)^2(m_3(z_2) + 1)^2} \\
&= (y_1\Delta_1 + y_2\Delta_2) \left\{ \lim_{r_2 \downarrow 1} \oint \frac{1}{2\pi i} \left( \log \frac{y_2(1 + hr_2\xi_2)(1 + \frac{h}{r_2\xi_2})}{(y_2 + hr_2\xi_2)(y_2 + \frac{h}{r_2\xi_2})} \right) \frac{(1 - y_2)r_1 h d\xi_1}{(y_2r_1\xi_1 + h)^2} \right\} \\
&\quad \left\{ \lim_{r_1 \downarrow 1} \oint \frac{1}{2\pi i} \left( \log \frac{y_2(1 + hr_1\xi_1)(1 + \frac{h}{r_1\xi_1})}{(y_2 + hr_1\xi_1)(y_2 + \frac{h}{r_1\xi_1})} \right) \frac{(1 - y_2)r_2 h d\xi_2}{(y_2r_2\xi_2 + h)^2} \right\}
\end{aligned}$$

$$= (y_1\Delta_1 + y_2\Delta_2) \cdot \frac{h^4}{y_1^2(y_1 + y_2)^2}$$

and

$$\begin{aligned} & -\frac{y_1\Delta_1 + y_2\Delta_2}{4\pi^2} \oint \oint \frac{f_i(\frac{z_1}{\alpha+z_1})f_j(\frac{z_2}{\alpha+z_2})dm_3(z_1)dm_3(z_2)}{(m_3(z_1) + 1)^2(m_3(z_2) + 1)^2} \\ & = (y_1\Delta_1 + y_2\Delta_2) \cdot \frac{y_1^2}{(y_1 + y_2)^2} \end{aligned}$$

when  $y_1 < 1, y_2 > 1$ . Thus, the variance part  $\tilde{\nu}_n$  is equal to

$$2 \log \frac{h^2}{|1 - y_1|(y_1 + y_2)} + 2 \log(\frac{y_1^2}{h^2})\delta_{y_1 > 1} + \frac{(y_1\Delta_1 + y_2\Delta_2)}{y_1^2(y_1 + y_2)^2} [y_1^4\delta_{y_1 < 1} + h^4\delta_{y_1 > 1}].$$

This completes the proof.  $\square$

## A.2 Proof Theorem 2.7

We now prove that  $\hat{\Delta}_1$  is a weakly consistent and asymptotically unbiased estimator of  $\Delta_1$ .

*Proof.* From the definition of  $\Delta_1$  in (2.2) and Chebyshev's inequality, we only need to prove the following two results: Under the same assumptions in Theorem 2.1 and the null hypothesis,

$$\mathbb{E}\hat{\Delta}_1 \rightarrow \Delta_1 \tag{A.9}$$

and

$$\mathbb{E}(\hat{\Delta}_1 - \Delta_1)^2 \rightarrow 0. \tag{A.10}$$

We first consider (A.9). Without loss of generality, we assume the mean of  $\mathbf{z}_j^{(1)}$  is zero. And following the same truncation steps in (Bai and Silverstein, 2004) we may truncate and re-normalize the random variables as follows

$$|x_{ij}^{(1)}| \leq \eta_n \sqrt{n}, \quad \mathbb{E}x_{ij}^{(1)} = 0, \quad \mathbb{E}(x_{ij}^{(1)})^2 = 1 \quad \text{and} \quad \mathbb{E}(x_{ij}^{(1)})^4 = \Delta_1 + 3 + O(n^{-1}),$$

where  $\eta_n \rightarrow 0$  slowly. In addition, in the following we assume the sample covariance matrix without the minus sample mean, because their difference is only a rank one matrix  $\bar{\mathbf{z}}_1 \bar{\mathbf{z}}_1'$  which will not affect the results. Form the proof of Theorem 1 in (Bai and Yin, 1993), one can conclude that under the conditions of Theorem 2.1, there exists a constant  $M > 0$  such that for any  $k > 0$

$$\mathbb{P}(\|(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\| \geq M) \leq n^{-k},$$

where  $\|\cdot\|$  means the spectral norm of a matrix. Thus, in the sequel, we can assume the smallest eigenvalue of  $c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}}$  is bounded away from zero uniformly. Then we have

$$\begin{aligned} & \mathbb{E}[(\mathbf{z}_j^{(1)} - \bar{\mathbf{z}}^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{z}} + c_{12}\mathbf{S}_2^{\mathbf{z}})^{-1}(\mathbf{z}_j^{(1)} - \bar{\mathbf{z}}^{(1)}) - \frac{p}{1-y}]^2 \\ &= \mathbb{E}[(\mathbf{z}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{z}} + c_{12}\mathbf{S}_2^{\mathbf{z}})^{-1}\mathbf{z}_j^{(1)}]^2 + \frac{p^2}{(1-y)^2} \\ & \quad - \frac{2p}{1-y} \mathbb{E}[(\mathbf{z}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{z}} + c_{12}\mathbf{S}_2^{\mathbf{z}})^{-1}\mathbf{z}_j^{(1)}] + O(p). \end{aligned}$$

Here we use the fact that  $\mathbb{E}[(\bar{\mathbf{z}}^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{z}} + c_{12}\mathbf{S}_2^{\mathbf{z}})^{-1}(\bar{\mathbf{z}}^{(1)})] = O(1)$  and  $\mathbb{E}[(\bar{\mathbf{z}}^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{z}} + c_{12}\mathbf{S}_2^{\mathbf{z}})^{-1}(\mathbf{z}_j^{(1)})] = O(1)$ . These results can be found in (Pan and Zhou, 2011). According to the independence of  $\mathbf{z}_j^{(1)}$  and  $\mathbf{S}_{1j}^{\mathbf{z}}$  and under the null hypothesis, we obtain that

$$\mathbb{E}[(\mathbf{z}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{z}} + c_{12}\mathbf{S}_2^{\mathbf{z}})^{-1}\mathbf{z}_j^{(1)}] = \mathbb{E}[(\mathbf{x}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_j^{(1)}] = \mathbb{E}tr[(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]$$

and

$$\begin{aligned} & \mathbb{E}[(\mathbf{z}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{z}} + c_{12}\mathbf{S}_2^{\mathbf{z}})^{-1}\mathbf{z}_j^{(1)}]^2 = (\Delta_1 - 3)\mathbb{E}tr[(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1} \circ (c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}] \\ & \quad + 2\mathbb{E}tr[(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-2}] + \{\mathbb{E}tr[(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]\}^2. \end{aligned}$$

where  $\circ$  is the Hadamard product. Notice that  $c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}}$  is a sample covariance matrix with dimension  $p$  and sample size  $n_1 + n_2 - 1$ , whose limit spectral distribution is the famous Marcenko-Pastur law with Stieltjes transform

$$s_{mp}(z, \gamma) = \frac{1 - \gamma - z + \sqrt{(z - 1 - \gamma)^2 - 4\gamma}}{2\gamma z}, \quad z \in \mathbb{C}^+. \quad (\text{A.11})$$

Here  $y := p/(n_1 + n_2 - 1) \rightarrow \gamma \in (0, \infty)$  and for any distribution function  $F$ , its Stieltjes transform is defined by

$$s(z) = \int_{-\infty}^{+\infty} \frac{1}{x - z} dF(x),$$

where  $z \in \mathbb{C}^+$ . Thus it is not difficult to check (see Section 3.3.2 in Bai and Silverstein (2010)) that

$$\frac{1}{p} \mathbb{E} \text{tr}[(c_{11} \mathbf{S}_{1j}^{\mathbf{x}} + c_{12} \mathbf{S}_2^{\mathbf{x}})^{-1}] - \lim_{z \downarrow 0} s_{mp}(z, y) \rightarrow 0 \quad (\text{A.12})$$

and

$$\frac{1}{p} \mathbb{E} \text{tr}[(c_{11} \mathbf{S}_{1j}^{\mathbf{x}} + c_{12} \mathbf{S}_2^{\mathbf{x}})^{-2}] - \lim_{z \downarrow 0} \frac{\partial s_{mp}(z, y)}{\partial z} \rightarrow 0.$$

From (A.11), L'Hospital's rule and the fact that  $y < 1$ , we have

$$\lim_{z \downarrow 0} s_{mp}(z, y) - \frac{1}{1 - y} = 0 \quad (\text{A.13})$$

and

$$\lim_{z \downarrow 0} \frac{\partial s_{mp}(z, y)}{\partial z} - \frac{1}{(1 - y)^3} = 0.$$

In addition, applying Lemma 4.3 of Bai et al. (2015), we obtain that

$$\begin{aligned} & \mathbb{E} \text{tr}[(c_{11} \mathbf{S}_{1j}^{\mathbf{x}} + c_{12} \mathbf{S}_2^{\mathbf{x}})^{-1} \circ (c_{11} \mathbf{S}_{1j}^{\mathbf{x}} + c_{12} \mathbf{S}_2^{\mathbf{x}})^{-1}] \\ &= \mathbb{E} \sum_{i=1}^p [(c_{11} \mathbf{S}_{1j}^{\mathbf{x}} + c_{12} \mathbf{S}_2^{\mathbf{x}})^{-1}(i, i)]^2 = p^{-1} \{\mathbb{E} \text{tr}[(c_{11} \mathbf{S}_{1j}^{\mathbf{x}} + c_{12} \mathbf{S}_2^{\mathbf{x}})^{-1}]\}^2 + o(p), \end{aligned}$$

which together with (A.12) and (A.13) implies

$$\mathbb{E} \text{tr}[(c_{11} \mathbf{S}_{1j}^{\mathbf{x}} + c_{12} \mathbf{S}_2^{\mathbf{x}})^{-1} \circ (c_{11} \mathbf{S}_{1j}^{\mathbf{x}} + c_{12} \mathbf{S}_2^{\mathbf{x}})^{-1}] = \frac{p}{(1 - y)^2} + o(p).$$

Thus, combining the above results, we conclude that

$$\mathbb{E}[(\mathbf{z}_j^{(1)} - \bar{\mathbf{z}}^{(1)})'(c_{11} \mathbf{S}_{1j}^{\mathbf{z}} + c_{12} \mathbf{S}_2^{\mathbf{z}})^{-1}(\mathbf{z}_j^{(1)} - \bar{\mathbf{z}}^{(1)}) - \frac{p}{1 - y}]^2 = \frac{p\Delta_1}{(1 - y)^2} + \frac{2p}{(1 - y)^3} + o(p),$$



which complete the proof of (A.9).

For (A.10), by the same argument as above, we have that  $\mathbb{E}(\hat{\Delta}_1 - \Delta_1)^2$  to

$$\begin{aligned} & \mathbb{E}(\hat{\Delta}_1 - \Delta_1)^2 \\ &= \frac{(1-y)^4 \mathbb{E}\{\sum_{j=1}^{N_1} [(\mathbf{x}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_j^{(1)} - \text{tr}(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]^2\}^2}{p^2 N_1^2} \\ & \quad - (\Delta_1 + \frac{2}{1-y})^2 + o(1). \end{aligned}$$

Now, by (2.1) in Bai and Silverstein (2004), we obtain that

$$\frac{\sum_{j=1}^{N_1} \mathbb{E}[(\mathbf{x}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_j^{(1)} - \text{tr}(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]^4}{p^2 N_1^2} = O(\eta_m^4 N_1^{-1}) \rightarrow 0. \quad (\text{A.14})$$

Next we use the fact that

$$\begin{aligned} (c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1} &= (c_{11}\mathbf{S}_{1ji}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1} \\ & \quad - \frac{\frac{1}{n_1+n_2-1}(c_{11}\mathbf{S}_{1ji}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_i^{(1)}(\mathbf{x}_i^{(1)})'(c_{11}\mathbf{S}_{1ji}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}}{1 + \frac{1}{n_1+n_2-1}(\mathbf{x}_i^{(1)})'(c_{11}\mathbf{S}_{1ji}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_i^{(1)}} \end{aligned}$$

and

$$(\mathbf{x}_i^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_i^{(1)} = \frac{(\mathbf{x}_i^{(1)})'(c_{11}\mathbf{S}_{1ji}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_i^{(1)}}{1 + \frac{1}{n_1+n_2-1}(\mathbf{x}_i^{(1)})'(c_{11}\mathbf{S}_{1ji}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_i^{(1)}},$$

where  $\mathbf{S}_{1ji}^{\mathbf{x}}$  is the sample covariance matrix by removing the vector  $\mathbf{x}_j^{(1)}$  and  $\mathbf{x}_i^{(1)}$ . Thus, we get that

$$\begin{aligned} & \sum_{i \neq j}^{N_1} \mathbb{E}\{[(\mathbf{x}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_j^{(1)} - \text{tr}(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]^2 \\ & \quad \times [(\mathbf{x}_i^{(1)})'(c_{11}\mathbf{S}_{1i}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_i^{(1)} - \text{tr}(c_{11}\mathbf{S}_{1i}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]^2\} \\ &= N_1(N_1 - 1) \mathbb{E}\{[(\mathbf{x}_1^{(1)})'(c_{11}\mathbf{S}_{112}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_1^{(1)} - \text{tr}(c_{11}\mathbf{S}_{112}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]^2 \\ & \quad \times [(\mathbf{x}_2^{(1)})'(c_{11}\mathbf{S}_{112}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_2^{(1)} - \text{tr}(c_{11}\mathbf{S}_{112}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]^2\} + O(N_1^2) \\ &= N_1(N_1 - 1) \left( \frac{p\Delta_1}{(1-y)^2} + \frac{2p}{(1-y)^3} \right)^2 + O(N_1^3), \end{aligned}$$

which together with (A.14) implies

$$\frac{(1-y)^4 \mathbb{E}\{\sum_{j=1}^{N_1} [(\mathbf{x}_j^{(1)})'(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}\mathbf{x}_j^{(1)} - \text{tr}(c_{11}\mathbf{S}_{1j}^{\mathbf{x}} + c_{12}\mathbf{S}_2^{\mathbf{x}})^{-1}]^2\}^2}{p^2 N_1^2} \\ = (\Delta_1 + \frac{2}{1-y})^2 + o(1).$$

Then we complete the proof of Theorem 2.7. □

## References

- Z. D. Bai, J. Hu, G. Pan, and W. Zhou. Convergence of the empirical spectral distribution function of Beta matrices. *Bernoulli*, 21(3):1538–1574, 2015.
- Z. D. Bai and J. W. Silverstein. CLT for linear spectral statistics of large-dimensional sample covariance matrices. *The Annals of Probability*, 32(1):553–605, 2004.
- Z. D. Bai and J. W. Silverstein. *Spectral analysis of large dimensional random matrices. Second Edition*. Springer Verlag, 2010.
- Z. D. Bai and Y. Q. Yin. Limit of the smallest eigenvalue of a large dimensional sample covariance matrix. *The Annals of Probability*, 21(3):1275–1294, 1993.
- Z. D. Bai, D. D. Jiang, J. F. Yao, and S. R. Zheng. Corrections to LRT on large-dimensional covariance matrix by RMT. *The Annals of Statistics*, 37(6B):3822–3840, Dec. 2009.
- M. S. Bartlett. Properties of Sufficiency and Statistical Tests. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 160(901):268–282, 1937.
- T. Cai, W. Liu, and Y. Xia. Two-Sample Covariance Matrix Testing and Support Recovery in High-Dimensional and Sparse Settings. *Journal of the American Statistical Association*, 108(501):265–277, 2013.

- T. T. Cai and Z. Ma. Optimal hypothesis testing for high dimensional covariance matrices. *Bernoulli*, 19(5B):2359–2388, 2013.
- L. Chen, D. Paul, R. Prentice, and P. Wang. A regularized Hotelling’s  $T$  (2) test for pathway analysis in proteomic studies. *Journal of the American Statistical Association*, 106(496):1345–1360, 2011.
- R. Cont. Empirical properties of asset returns: stylized facts and statistical issues. *Quantitative Finance*, 1(April):223–236, 2001.
- D. Jiang, T. Jiang, and F. Yang. Likelihood ratio tests for covariance matrices of high-dimensional normal distributions. *Journal of Statistical Planning and Inference*, 142(8):2241–2256, Aug. 2012.
- T. Jiang and F. Yang. Central limit theorems for classical likelihood ratio tests for high-dimensional normal distributions. *The Annals of Statistics*, 41(4):2029–2074, Aug. 2013.
- O. Ledoit and M. Wolf. Some hypothesis tests for the covariance matrix when the dimension is large compared to the sample size. *The Annals of Statistics*, 30(4):1081–1102, 2002.
- J. Li and S. X. Chen. Two sample tests for high-dimensional covariance matrices. *The Annals of Statistics*, 40(2):908–940, Apr. 2012.
- G. Pan and W. Zhou. Central limit theorem for Hotelling’s  $T^2$  statistic under large dimension. *The Annals of Applied Probability*, 21(5):1860–1910, 2011.
- K. C. S. Pillai and K. Jayachandran. Power comparisons of tests of two multivariate hypotheses based on four criteria. *Biometrika*, 54(1):195–210, 1967.
- K. C. S. Pillai and K. Jayachandran. Power comparisons of tests of equality of two covariance matrices based on four criteria. *Biometrika*, 55(2):335–342, 1968.

- M. S. Srivastava and H. Yanagihara. Testing the equality of several covariance matrices with fewer observations than the dimension. *Journal of Multivariate Analysis*, 101(6): 1319–1329, July 2010.
- S. S. Wilks. Certain generalizations in the analysis of variance. *Biometrika*, 24(3):471–494, 1932.
- S. S. Wilks. Sample criteria for testing equality of means, equality of variances, and equality of covariances in a normal multivariate distribution. *The Annals of Mathematical Statistics*, pages 257–281, 1946.
- Q. Y. Zhang, J. Hu and Z. D. Bai. Supplementary material to “Optimal modification of the LRT for the equality of two high-dimensional covariance matrices”.
- S. Zheng, Z. Bai, and J. Yao. Substitution principle for CLT of linear spectral statistics of high-dimensional sample covariance matrices with applications to hypothesis testing. *Annals of Statistics*, 43(2):546–591, 2015.
- S. Zheng, Z. Bai, and J. Yao. CLT for eigenvalue statistics of large-dimensional general Fisher matrices with applications. *Bernoulli*, 23(2):1130–1178, 2017.

Qiuyan Zhang, Jiang Hu and Zhidong Bai

School of Mathematics and Statistics

Northeast Normal University, China.

E-mail: (zhangqy919@nenu.edu.cn, huj156@nenu.edu.cn, baizd@nenu.edu.cn)