

Global algorithms for maximal eigenpair

Mu-Fa Chen

Assisted by Yue-Shuang Li

(Beijing Normal University)

April 29, 2017

Abstract

This paper is a continuation of [5] where an efficient algorithm for computing the maximal eigenpair was introduced first for tridiagonal matrices and then extended to the irreducible matrices with nonnegative off-diagonal elements. This paper introduces two global algorithms for computing the maximal eigenpair in a rather general setup, including even a class of real (with some negative off-diagonal elements) or complex matrices.

2000 *Mathematics Subject Classification*: 60J60, 34L15

Key words and phrases. Maximal eigenpair, shifted inverse iteration, global algorithm.

1 Introduction

To compute the maximal eigenpair of the tridiagonal matrices with positive sub-diagonal elements, an efficient algorithm was introduced [5; §3]. In the tridiagonal case, the construction of the initials for the algorithm is explicit. In some sense, the results are more or less complete (a modified algorithm, Algorithm 17, is included in §4.4). Next, the algorithm was extended to the general case in [5; §4] which is still efficient for tridiagonally dominant matrices. Note that the initial v_0 constructed in [5; §4.2] may not be efficient enough, since the shape of the maximal eigenvector can be rather arbitrary, could be quite far away from v_0 constructed in [5; §4.2]. Thus, we are worrying about the efficiency of the extended algorithm and moreover a global algorithm is still missed in our general setup. This is the aim of this paper. In §3, a part of the off-diagonal elements of the matrices are allowed to be negative. We can even handle with some complex matrices. Let us concentrate on the nonnegative matrices from now on, unless otherwise is stated.

By a shift if necessary, unless otherwise stated, we assume that the given matrix $A = (a_{ij} : 0 \leq i, j \leq N)$ is irreducible and nonnegative: $a_{ij} \geq 0$. We now state our algorithms. To guarantee the convergence of the iterations in the paper, we assume that the matrix is irreducible having positive trace, or equivalently,

$$A^n > 0 \quad \text{for each } n \geq \text{some } n_0. \quad (1)$$

We mention that in the present nonnegative case, the condition having positive trace is not serious, otherwise, simply adopt a shift as mentioned at the beginning of [5].

In what follows, we omit, without mention time by time, the trivial case that $\sum_j a_{ij} \equiv \text{constant } m > 0$. Since then the maximal eigenpair of A becomes $(m, \mathbb{1})$, where $\mathbb{1}$ is the constant function having components 1 everywhere.

Recall that the choice of the initials is quite essential for the Rayleigh Quotient Iteration (RQI), a special shifted inverse iteration. In general, it seems no hope at the moment to have such explicit analytic formulas as used in [5; §3]. Instead, as suggested in many textbooks, one may use other approach to obtain in a numerical way the required initials, say use the power iteration for instance. The last approach is safe, but rather slow as shown at the beginning of [5]. This leads us to come back to the shifted inverse iterations which is a fast cubic algorithm. The ratio of the numbers of iterations for these two algorithms can be thousands. Now, a critical point is to avoid the dangerous pitfalls, i.e., the region $(0, \rho(A))$, where $\rho(A)$ is the maximal eigenvalue of A . The answer is given in part (1) of the next two algorithms. At the moment, we are interesting in the generality and safety, do not take care much about the convergence speed, perhaps, maybe some price we have to pay here. We will see soon what happen in the next section.

Algorithm 1 (Specific Rayleigh quotient iteration) Let $A = (a_{ij})$ be given.

(1) Define column vectors

$$w^{(0)} = (1, 1, \dots, 1)^*, \quad v^{(0)} = w^{(0)} / \sqrt{N+1},$$

and set

$$z^{(0)} = \max_{0 \leq i \leq N} (Aw^{(0)})_i.$$

(2) For given $v := v^{(n-1)}$ and $z := z^{(n-1)}$, let $w := w^{(n)}$ solve the equation

$$(zI - A)w = v. \quad (2)$$

As in step (1), define $v^{(n)} = w / \sqrt{w^* w}$. Next, define

$$x^{(n)} = \min_{0 \leq j \leq N} \frac{(Aw^{(n)})_j}{w_j^{(n)}}, \quad y^{(n)} = \max_{0 \leq j \leq N} \frac{(Aw^{(n)})_j}{w_j^{(n)}}, \quad z^{(n)} = v^{(n)*} A v^{(n)}.$$

(3) If at some $n \geq 1$, $y^{(n)} - x^{(n)} < 10^{-6}$ (or $|z^{(n)} - z^{(n+1)}| < 10^{-6}$)(say!), then stop the computation. At the same time, regard $(z^{(n)}, v^{(n)})$ as an approximation of the maximal eigenpair.

The algorithm was presented in [5; §4.1: Choice I]. The simplest choice v_0 is reasonable in the sense that it enables us to cover the general case. We did not pay enough attention on this algorithm since it looks less efficient. However, as some examples will be illustrated below, this algorithm is actually rather powerful. It is the place to state the main new algorithm of the paper.

Algorithm 2 (Shifted inverse iteration) Everything is the same as in Algorithm 1, except $y^{(n)}$ and $z^{(n)}$ defined in parts (2) and (3) there are exchanged. Moreover, the resulting $z^{(n)}$ (resp. $x^{(n)}$) is decreasing (resp. increasing) in n .

Let us repeat the sequences $z^{(n)}$, $y^{(n)}$ and $x^{(n)}$ defined in Algorithm 2:

$$x^{(n)} = \min_{0 \leq j \leq N} \frac{(Aw^{(n)})_j}{w_j^{(n)}}, \quad y^{(n)} = v^{(n)*} A v^{(n)}, \quad z^{(n)} = \max_{0 \leq j \leq N} \frac{(Aw^{(n)})_j}{w_j^{(n)}}.$$

It is obvious that

$$x^{(n)} \leq y^{(n)} \leq z^{(n)}.$$

In general, Algorithm 1 is a little effective than Algorithm 2, saving one iteration for instance, but in Algorithm 2, each iteration is safe, never failed into the pitfall. This is based on the following dual variational formula.

Proposition 3 [11; Theorem (8)] For a nonnegative irreducible matrix A , the Collatz–Wielandt formula holds:

$$\sup_{x > 0} \min_{i \in E} \frac{(Ax)_i}{x_i} = \rho(A) = \inf_{x > 0} \max_{i \in E} \frac{(Ax)_i}{x_i}.$$

Actually, suppose that we have $w^{(n-1)} > 0$ in Algorithm 2. Then by Proposition 3 and step (2) of Algorithm 2, we have $z^{(n-1)} > \rho(A)$ and then the solution $w^{(n)}$ to the equation (2) should be positive: $w^{(n)} > 0$. Otherwise, if $z^{(n-1)} < \rho(A)$, then the solution $w^{(n)}$ is negative. This is the main reason why we choose such a $z^{(n-1)}$ for each $n \geq 1$ in Algorithm 2 and in the case of $n = 0$ in Algorithm 1 as our shift, avoiding the change of signs. Note that in Algorithm 1 we adopt $y^{(n)}$ at each step $n \geq 1$, hence the solution $w^{(n)}$ changes its sign often. This seems dangerous because $y^{(n)}$ is located in the dangerous region, but up to now, we have not meet trouble. Therefore, it is still regarded as one of our two main algorithms. Certainly, if necessary, you can replace $z^{(k)}$ defined in Algorithm 1 by $z^{(k)} = \max_{0 \leq j \leq N} (Aw^{(k)})_j / w_j^{(k)}$ for some $k = 1, 2, 3$ or so. Or, simply use Algorithm 2 instead.

A careful comparison of Algorithm 1 and the powerful one introduced in [5; §3] is delayed to the Appendix.

An easier way to see the efficiency of Algorithms 1 and 2 is comparing them with the one given in [5; §4.2]. Suppose that we have used three iterations in computing a model using the method introduced in [5; §4.2], this means on the one hand we have solved the linear equations in three times. On the other hand, we have solved three more times in advance to figure out the initials $v^{(0)}$ and $z^{(0)}$ in terms of the triple (ψ, h, μ) . Altogether, we have solved six linear equations. Or in other words, we have used 6 iterations in the computation for the specific model. Thus, Algorithms 1 and 2 should be regarded as efficient one if no more than 6 iterations are used in the computation for the same model. As we will see soon, we are actually in such a successful situation.

To conclude this section, we rewrite Algorithms 1 and 2 to a class of matrices with nonnegative off-diagonal elements and negative diagonal elements: $Q = (q_{ij})$:

$$q_{ij} \geq 0, \quad i \neq j; \quad \sum_{j=0}^N q_{ij} \leq 0, \quad 0 \leq i \leq N.$$

In this case, we are studying the maximal eigenpair of Q , or alternatively, the minimal eigenpair of $-Q$. To which, the next two algorithms are devoted.

Again, the trivial case that $\sum_{j=0}^N q_{ij}$ equals a constant is ignored throughout the paper.

Algorithm 4 (Specific Rayleigh quotient iteration) Let $Q = (q_{ij})$ be given.

(1) Define column vectors

$$w^{(0)} = (1, 1, \dots, 1)^*, \quad v^{(0)} = w^{(0)} / \sqrt{N+1},$$

and set $z^{(0)} = 0$.

(2) For given $v := v^{(n-1)}$ and $z := z^{(n-1)}$, let $w := w^{(n)}$ solve the equation

$$(-Q - zI)w = v. \quad (3)$$

As in step (1), define $v^{(n)} = w / \sqrt{w^* w}$. Next, define

$$x^{(n)} = \min_{0 \leq j \leq N} \frac{((-Q)w^{(n)})_j}{w_j^{(n)}}, \quad y^{(n)} = \max_{0 \leq j \leq N} \frac{((-Q)w^{(n)})_j}{w_j^{(n)}}, \quad z^{(n)} = v^{(n)*}(-Q)v^{(n)}.$$

(3) If at some $n \geq 1$, $y^{(n)} - x^{(n)} < 10^{-6}$ (or $|z^{(n)} - z^{(n+1)}| < 10^{-6}$)(say!), then stop the computation. At the same time, regard $(z^{(n)}, v^{(n)})$ as an approximation of the minimal eigenpair.

Algorithm 5 (Shifted inverse iteration) Everything is the same as in Algorithm 4, except $x^{(n)}$ and $z^{(n)}$ defined in parts (2) and (3) there are exchanged. Moreover, the resulting $z^{(n)}$ (resp. $x^{(n)}$) is increasing (resp. decreasing) in n .

Algorithms 4 and 5 are based on [5; Corollary 12], a corollary of Proposition 3.

2 Examples

To illustrate the power of the algorithms introduced in the last section, we examine some typical examples in this section.

To go to practical computation for concrete models, our readers are urged to prepare enough patience, one may have a large number of iterations since the initials given in part (1) are quite rough.

The efficient application of Algorithm 1 was illustrated by [5; Examples 13–16]. To have a concrete comparison of the present algorithms with the one introduced in [5; §4.2], let us consider a simple example.

Example 6 [5; Example 21] Let

$$Q = \begin{pmatrix} -3 & 2 & 0 & 1 & 0 \\ 4 & -7 & 3 & 0 & 0 \\ 0 & 5 & -5 & 0 & 0 \\ 10 & 0 & 0 & -16 & 6 \\ 0 & 0 & 0 & 11 & -11 - b_4 \end{pmatrix}.$$

Corresponding to different b_4 , the minimal eigenvalue λ_0 of $-Q$ and its approximation are as follows. Here and in what follows, we stop at $z^{(2)}$ once the outputs $z^{(k)} = z^{(2)}$ for every $k \geq 2$.

Table 1. The outputs by Algorithm 1

b_4	$z^{(1)}$	$z^{(2)}$	$z^{(3)} = \lambda_{\min}(-Q)$
0.01	0.000278773	0.000278686	
1	0.0251531	0.0245175	
100	0.191729	0.182822	0.182819
10^4	0.201695	0.195019	0.195015

Table 2. The outputs by Algorithm 2

b_4	$z^{(1)}$	$z^{(2)}$	$z^{(3)} = \lambda_{\min}(-Q)$
0.01	0.000278637	$0.000278686 = \lambda_{\min}(-Q)$	
1	0.0241546	0.0245175	
100	0.168776	0.18275	0.182819
10^4	0.179525	0.194932	0.195015

While the outputs by the algorithm given in [5; §4.2] are the following.

Table 3. The outputs by the algorithm given in [5]

b_4	$z^{(1)}$	$z^{(2)}$	$z^{(3)} = \lambda_{\min}(-Q)$
0.01	0.000278573	0.000278686	
1	0.0236258	0.0245174	0.0245175
100	0.200058	0.182609	0.182819

These tables show that the three algorithms are more or less at the same level of effectiveness. However, the first two are actually more economic since the last one requires an extra work computing the initial v_0 .

Comparing [5; Example 15] with the corrected version of [5; Example 20] and its improvements given in [5; Tables 11, 12] (see the author's homepage), we see that the extended algorithm introduced in [5; §4.2] can be less efficient than Algorithm 1, it has some limitation for general non-symmetrizable (non-symmetric) matrices. We call a matrix $A = (a_{ji})$ is symmetrizable, if there exists a positive measure (μ_i) such that

$$\mu_i a_{ij} = \mu_j a_{ji}, \quad i \neq j.$$

A simple necessary condition for the symmetrizability is

$$a_{ij} > 0 \iff a_{ji} > 0, \quad i \neq j.$$

Refer to [3; Chapter 7] and references within for the solution to the symmetrizability problem.

Let us start at a class of non-symmetrizable matrices which are taken from the so-called single birth Q -matrix (cf. [3] and references within). Define

$$Q = \begin{pmatrix} -1 & 1 & 0 & 0 & \cdots & 0 & 0 \\ a_1 & -a_1-2 & 2 & 0 & \cdots & 0 & 0 \\ a_2 & 0 & -a_2-3 & 3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \cdots & N-2 & 0 \\ a_{N-1} & 0 & 0 & 0 & \cdots & -a_{N-1}-N+1 & N-1 \\ a_N & 0 & 0 & 0 & \cdots & 0 & -a_N-N \end{pmatrix}. \quad (4)$$

For this matrix, we have computed several cases:

$$a_k = 1/(k+1), \quad a_k \equiv 1, \quad a_k = k, \quad a_k = k^2.$$

Among them, the first one is hardest and is hence presented below.

Example 7 Let Q be defined by (4). For different N , the outputs of Algorithm 5 (equivalently, Algorithm 2) are as follows.

Table 4. The outputs for different N by Algorithm 5

N	$z^{(1)}$	$z^{(2)}$	$z^{(3)}$	$z^{(4)}$	$z^{(5)}$	$z^{(6)}$
8	0.276727	0.427307	0.451902	0.452339		
16	0.222132	0.367827	0.399959	0.400910		
32	0.187826	0.329646	0.370364	0.372308	0.372311	
50	0.171657	0.311197	0.357814	0.360776	0.360784	
100	0.152106	0.287996	0.343847	0.349166	0.349197	
500	0.121403	0.247450	0.321751	0.336811	0.337186	
1000	0.111879	0.233257	0.313274	0.334155	0.335009	0.335010
5000	0.0947429	0.205212	0.293025	0.328961	0.332609	0.332635
10^4	0.0888963	0.194859	0.284064	0.326285	0.332113	0.332188

The last line shows that when $N = 10^4$, $\lambda_{\min}(-Q) \approx 0.332188$. If we use the shifted matrix $A = Q + mI$, then $\rho(A) \approx 9999.67$. From which, we get $\lambda_{\min}(-Q) \approx 10^4 + 10^{-4} - 9999.67$. Clearly, the second approach has a less precise output. That is the main difference between Algorithms 1, 2 and 4, 5, even though they are equivalent analytically.

It should be meaningful to have a comparison of the present results with those produced by [5; §4.2]. The outputs listed in the table below come from the algorithm without using δ_1 defined in that section. For the outputs using δ_1 , one more iteration is needed for those N from 16 to 100 listed in the table.

Table 5. The outputs for different N by the algorithm given in [5; §4.2]

N	$z^{(1)}$	$z^{(2)}$	$z^{(3)}$
8	0.450694	0.452338	0.452339
16	0.399520	0.400910	
32	0.371433	0.372311	
64	0.355722	0.355940	
100	0.349501	0.349197	
500	0.340666	0.337185	0.337186
1000	0.340871	0.335003	0.335010
5000	0.347505	0.332536	0.332635
10^4	0.352643	0.331975	0.332188

Clearly, the general algorithm introduced in [5; §4.2] is efficient for this non-symmetrizable model. We have seen that the present algorithms require more iterations than the earlier one, this is reasonable since the computations of the initials are excluded from the last table. Actually, the computations of the last table cost double time than the previous one.

The next example is motivated from the classical branching process. Denote by $(p_k : k \geq 0)$ a given probability measure with $p_1 = 0$. Let

$$Q = \begin{pmatrix} -1 & p_2 & p_3 & p_4 & \cdots & p_{N-1} & \sum_{k \geq N} p_k \\ 2p_0 & -2 & 2p_2 & 2p_3 & \cdots & 2p_{N-2} & 2 \sum_{k \geq N-1} p_k \\ 0 & 3p_0 & -3 & 3p_2 & \cdots & 3p_{N-3} & 3 \sum_{k \geq N-2} p_k \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \vdots & \vdots & \ddots & \ddots & -(N-1) & (N-1)p_2 \\ 0 & 0 & 0 & 0 & \cdots & Np_0 & -Np_0 \end{pmatrix},$$

In the original model, the state 0 is an absorbing one. Here we regard it as a killing boundary. Hence it is ruled out from our state space. Thus, the matrix is defined on $E := \{1, 2, \dots, N\}$. Set $M_1 = \sum_{k \in E} kp_k$. When $N = \infty$, in the subcritical case that $M_1 < 1$, with a little modification at 0, it is known that the process generated by Q is ergodic, and is indeed exponentially ergodic (cf. [8; Theorem 1.4 (iii)]). Hence the exponential convergence rate should be positive. Otherwise, the process is not ergodic and so the convergence rate should be zero.

From now on, fix

$$p_0 = \alpha/2, \quad p_1 = 0, \quad p_2 = (2 - \alpha)/2^2, \quad \dots, \quad p_n = (2 - \alpha)/2^n, \quad \dots, \quad \alpha \in (0, 2).$$

Then $M_1 = 3(2 - \alpha)/2$ and hence we are in the subcritical case iff $\alpha \in (4/3, 2)$.

Example 8 Set $\alpha = 1$. Then the outputs of the approximation for the minimal eigenvalue of $-Q$ by Algorithm 2 (or 5) are as follows.

Table 6. The outputs in the supercritical case

N	$z^{(1)}$	$z^{(2)}$	$z^{(3)}$
8	0.0311491	0.0346044	0.0346310
16	0.00256281	0.00260088	

When $N \geq 50$, $z^{(1)} < 10^{-6}$. Hence, $z^{(n)}$ decays quite quick to zero when $N \rightarrow \infty$ (for $n \geq 2$). This is reasonable since we are now away from the subcritical region.

Example 9 Set $\alpha = 7/4$. We are now in the subcritical case and so the maximal eigenvalue should be positive. We want to know how fast the local maximal eigenvalue becomes stable (i.e., close enough to the converge rate at $N = \infty$). Again, we adopt Algorithm 2 (or 5). Up to $N = 10^4$, the steps of the iterations we need are no more than 6. To fasten the convergence, we adopt a convex combination, as we did several times in [5]. Replace the original $z^{(0)} = \max_{0 \leq j \leq N} (Aw^{(0)})_j$ by

$$z^{(0)} = \xi \min_{0 \leq j \leq N} (Aw^{(0)})_j + (1 - \xi)(v^{(0)})^* Av^{(0)}.$$

In view of the practice on $N = 8$, we make the choice that $\xi = 0.69$. Then the outputs of the approximation of the minimal eigenvalue of $-Q$ for different N are as follows.

Table 7. The outputs in the subcritical case

N	$z^{(1)}$	$z^{(2)}$	$z^{(3)}$	$z^{(4)}$
8	0.637800	0.638153		
16	0.621430	0.625490	0.625539	
50	0.609976	0.624052	0.624997	0.625000
100	0.606948	0.623377	0.624991	0.625000
500	0.604409	0.622116	0.624962	0.625000
1000	0.604082	0.621688	0.624944	0.625000
5000	0.603817	0.620838	0.62489	0.625000
10^4	0.603784	0.620511	0.624861	0.625000

From the above table, we see that for N varies from 8 to 10^4 , in each case, we need at most 4 iterations only. The computation in each case costs no more than one minute. Besides, starting from $N = 50$, the final outputs are all the same: 0.625, which then can be regarded as a very good approximation of the maximal eigenvalue at infinity $N = \infty$.

Hopefully, we have already shown the power of our algorithms.

3 A class of real or complex matrices

This section is out of the scope of [5] which depends heavily on probabilistic idea. Thanks are given to the extended Perron–Frobenius theory ([10]–[12]) which makes this section possible.

First, we consider the real case. The special case that all off-diagonal elements of A are negative has been treated above, using $-Q$ instead of A here. Thus, we are now mainly interested in the case that a part of the off-diagonal elements are negative. Again, we are concentrated in the study of the maximal eigenpair.

Proposition 10 Let A be a real matrix satisfying (1). Then Algorithms 1 and 2 are available.

Proof. By [10; Theorem 2.2], condition (1) implies that the matrix A possesses the strong Perron–Frobenius property. Hence it has the maximal eigenvalue $\rho(A)$ which is simple, positive and corresponds to a positive eigenvector. Besides, by [10; Theorem 2.6], the Collatz–Wielandt formula given in Proposition 3 holds. These facts are enough to use Algorithms 1 and 2. \square

The next simple observation is helpful.

Lemma 11 Condition (1) holds iff

$$A^k > 0 \quad \text{for } k = n_0, n_0 + 1, \dots, 2n_0 - 1.$$

Proof. Given $n \geq n_0$, write

$$n = rn_0 + s$$

for some integer $r \geq 1$ and $s = 0, 1, \dots, n_0 - 1$. If $r = 1$, then the conclusion holds by assumption. Otherwise, let $r \geq 2$. Then express

$$n = (r - 1)n_0 + (n_0 + s).$$

It follows that

$$A^n = (A^{n_0})^{r-1} A^{n_0+s} > 0$$

as required. \square

We now illustrate our algorithms by a simple example.

Example 12 [11; Example (7)] Let

$$A = \begin{pmatrix} -1 & 8 & -1 \\ 8 & 8 & 8 \\ -1 & 8 & 8 \end{pmatrix}.$$

Then

$$A^2 = \begin{pmatrix} 66 & 48 & 57 \\ 48 & 192 & 120 \\ 57 & 120 & 129 \end{pmatrix} > 0, \quad A^3 = \begin{pmatrix} 261 & 1368 & 774 \\ 1368 & 2880 & 2448 \\ 774 & 2448 & 1935 \end{pmatrix} > 0.$$

By Lemma 11, condition (1) holds with $n_0 = 2$. The eigenvalues of A are as follows.

$$17.5124, \quad -7.4675, \quad 4.95513.$$

The corresponding maximal eigenvector is

$$(0.486078, 1.24981, 1)$$

which is positive.

Here are the outputs of our algorithms. Both algorithms are started at $z^{(0)} = 24$.

Table 8. The outputs for a matrix with more negative elements

n	$z^{(n)}$: Algorithm 1	$z^{(n)}$: Algorithm 2
1	17.3772	18.5316
2	17.5124	17.5416
3		17.5124

Next, we turn to study the complex case. Instead of (1), we assume that

$$\operatorname{Re}(A^n) > 0 \quad \text{for } n \geq \text{some } n_0. \quad (5)$$

Certainly, as usual $\operatorname{Re}(A)$ means the real part of a complex matrix A . This condition is based on [12; Theorems 2.3 and 2.2], from which we know that A has the maximal, simple, positive eigenvalue. Then we have a weak extension of the Collatz–Wielandt formula as follows.

Proposition 13 [12; Theorems 2.3 and 2.4] Let $A^k \neq 0$ for each $k \geq 1$ and $\operatorname{Re}(A^n) \geq 0$ for every large enough n . Then we have for each $x > 0$

$$\min_{0 \leq j \leq N} \frac{(\operatorname{Re}(A)x)_j}{x_j} \leq \rho(A) \leq \max_{0 \leq j \leq N} \frac{(\operatorname{Re}(A)x)_j}{x_j}.$$

Since for the complex conjugate \bar{x}^* of x , the quantity \bar{x}^*Ax may still be complex, in view of this, Proposition 13 and the positivity of $\rho(A)$ by (5), it seems not reasonable to use $\bar{x}^*Ax/(\bar{x}^*x)$ as a shift. In this sense, we do not have a modified version of Algorithm 1. Fortunately, Algorithm 2 is still meaningful.

Algorithm 14 (Shifted inverse iteration) Assume (5).

(1) Define column vectors

$$w^{(0)} = (1, 1, \dots, 1)^*, \quad v^{(0)} = w^{(0)} / \sqrt{N+1},$$

and set

$$z^{(0)} = \max_{0 \leq i \leq N} (\operatorname{Re}(A)w^{(0)})_i.$$

(2) For given $v := v^{(n-1)}$ and $z := z^{(n-1)}$, let $w := w^{(n)}$ solve the equation

$$(zI - A)w = v. \quad (6)$$

As in step (1), define $v^{(n)} = w / \sqrt{\bar{w}^* w}$. Next, define

$$z^{(n)} = \max_{0 \leq j \leq N} \frac{(\operatorname{Re}(A)\operatorname{Re}(w^{(n)}))_j}{\operatorname{Re}(w^{(n)})_j}, \quad y^{(n)} = (\bar{v}^{(n)})^* A v^{(n)}.$$

(3) If at some $n \geq 1$, $|y^{(n+1)} - y^{(n)}| < 10^{-6}$ (say!), then stop the computation. At the same time, regard $(y^{(n)}, v^{(n)})$ as an approximation of the maximal eigenpair.

Note that in Algorithm 14, the sequence $\{y^{(n)}\}_{n \geq 0}$, but not $\{z^{(n)}\}_{n \geq 0}$, converges to $\rho(A)$. To illustrate the use of the algorithm, we consider the following example.

Example 15 [12; Example 2.1] Let

$$A = \begin{pmatrix} 0.75 - 1.125i & 0.5882 - 0.1471i & 1.0735 + 1.4191i \\ -0.5 - i & 2.1765 + 0.7059i & 2.1471 - 0.4118i \\ 2.75 - 0.125i & 0.5882 - 0.1471i & -0.9265 + 0.4191i \end{pmatrix},$$

where the coefficients are all accurate, to four decimal digits. Then A has eigenvalues

$$3, \quad -2 - i, \quad 1 + i$$

with maximal eigenvector

$$(0.408237, 0.816507, 0.408237).$$

The outputs of Algorithm 14 are as follows.

Table 9. The outputs for a complex matrix

$y^{(1)}$	$y^{(2)}$	$y^{(3)}$
$3.03949 - 0.0451599i$	$3.00471 - 0.0015769i$	3

4 Appendix

4.1 Proof of the last assertion in Algorithm 2

Proposition 16 The sequence

$$z^{(n)} = \max_{0 \leq j \leq N} \frac{(Aw^{(n)})_j}{w_j^{(n)}} \quad \left(\text{resp. } x^{(n)} = \min_{0 \leq j \leq N} \frac{(Aw^{(n)})_j}{w_j^{(n)}} \right)$$

defined in Algorithm 2 is decreasing (resp. increasing) in n .

Proof. Let $w > 0$ and define

$$\bar{\rho} = \max_{0 \leq j \leq N} \frac{(Aw)_j}{w_j}.$$

Then $(Aw)_j \leq \bar{\rho} w_j$ for every j . That is,

$$(A_z w)_j \leq \bar{\rho}_z w_j \quad \forall j, \quad A_z := A/z, \quad \bar{\rho}_z = \bar{\rho}/z.$$

Since $A_z \geq 0$, it follows that

$$A \sum_{n=0}^{\infty} A_z^n w \leq A \left(w + \bar{\rho}_z \sum_{n=0}^{\infty} A_z^n w \right) \leq \bar{\rho} w + \sum_{n=1}^{\infty} \bar{\rho} A_z^n w = \bar{\rho} \sum_{n=0}^{\infty} A_z^n w.$$

This means that

$$A(I - A_z)^{-1} w \leq \bar{\rho}(I - A_z)^{-1} w$$

since $z > \rho(A)$ by assumption and then $\rho(A_z) < 1$. Hence

$$\max_{0 \leq j \leq N} \frac{(A((I - A_z)^{-1} v))_j}{((I - A_z)^{-1} v)_j} \leq \bar{\rho}, \quad v := w/\sqrt{w^* w}.$$

Regarding $w = w^{(n-1)}$ and $v = v^{(n-1)}$, this gives us

$$z^{(n)} = \max_{0 \leq j \leq N} \frac{(Aw^{(n)})_j}{w_j^{(n)}} \leq \bar{\rho} = \max_{0 \leq j \leq N} \frac{(Aw^{(n-1)})_j}{w_j^{(n-1)}} = z^{(n-1)}.$$

Here we have assumed that $z^{(n-1)} > \rho(A)$, otherwise, the computation should be finished at the step $n - 1$. We have thus proved the assertion on $z^{(n)}$. Dually, we have the assertion on $x^{(n)}$. \square

4.2 Proof of the last assertion in Algorithm 5

Recall the sequence $\{z^{(n)}\}$ used in Algorithm 2 is given in Proposition 16. Denote by $\{\tilde{z}^{(n)}\}$. Then, by the relation of Q and A used in Algorithm 5: $A = Q + mI$, where $m = \max_i \sum_j a_{ij}$. Hence

$$z^{(0)}I - A = -Q - (m - z^{(0)})I.$$

This means not only $\tilde{z}^{(0)} = 0$, but also

$$w^{(1)} = (z^{(0)}I - A)^{-1}v^{(0)} = (-Q - \tilde{z}^{(0)}I)^{-1}v^{(0)} =: \tilde{w}^{(1)},$$

where $\tilde{w}^{(1)}$ is obtained by the first iteration of Algorithm 5. Furthermore, similar to the proof of [5; Corollary 12], we have

$$\tilde{z}^{(1)} = \min_i \frac{(-Q\tilde{w}^{(1)})_i}{\tilde{w}_i^{(1)}} = m - \max_i \frac{(Aw^{(1)})_i}{w_i^{(1)}} = m - z^{(1)}.$$

Recursively, we obtain the required assertion. \square

4.3 Comparison of Algorithms 1 and 4 with the one given in [5; §3]

Since Algorithms 1 and 4 are equivalent, we need only to compare Algorithm 4 with the one given in [5; §3]. The main difference is their initial $(v^{(0)}, z^{(0)})$. Clearly, the initial $v^{(0)}$ used in [5; §3] is finer than the one used in Algorithm 4. Hence, we need only to compare their $z^{(0)}$.

Next, let $v := v^{(0)}$ be the initial vector used in [5; §3]. Denote by w be the solution of the ordinary inverse iteration (that is the first step of Algorithm 4 or equivalently, Algorithm 1):

$$-Qw = v.$$

Then

$$\frac{(-Qw)_j}{w_j} = \frac{v_j}{((-Q)^{-1}v)_j} = \Pi_j(v)^{-1}. \quad (7)$$

Here in the last equality of (7), we have used the first formula in the proof of [5; Proposition 23]. Hence

$$\inf_j \frac{(-Qw)_j}{w_j} = \inf_j \Pi_j(v)^{-1}. \quad (8)$$

The right-hand side of (8) is just δ_1^{-1} used in [5; §3] as its initial $z^{(0)}$. The left-hand side of (8) should be positive, due to the inverse iteration algorithm, it is certainly bigger than 0 used as the initial $z^{(0)}$ in Algorithm 4. In conclusion, both initials used in [5; §3] are better than those used in Algorithm 4. This completes the comparison of Algorithm 4 and the one given in [5; §3].

Naturally, this comparison leads to the next remark.

4.4 Modification of the algorithm defined in [5; §3]

Step 1. By a shift if necessary, we may assume that we are given a matrix Q having the form

$$Q = \begin{pmatrix} -(b_0 + c_0) & b_0 & 0 & 0 & \cdots \\ a_1 & -(a_1 + b_1 + c_1) & b_1 & 0 & \cdots \\ 0 & a_2 & -(a_2 + b_2 + c_2) & b_2 & \cdots \\ \vdots & \vdots & \ddots & \ddots & \ddots \\ 0 & 0 & 0 & a_N & -(a_N + c_N) \end{pmatrix},$$

where $a_i > 0$, $b_i > 0$, $c_i \geq 0$ but $c_i \neq 0$. Note that the maximal eigenvalue of Q is shifted from the original one but the corresponding eigenvector remain the same.

Step 2. Following [5; §3], assume for a moment that some of c_i ($i = 0, 1, \dots, N-1$) is positive. Then, define

$$r_0 = 1 + \frac{c_0}{b_0}, \quad r_n = 1 + \frac{a_n + c_n}{b_n} - \frac{a_n}{b_n r_{n-1}}, \quad 1 \leq n < N,$$

$$h_0 = 1, \quad h_n = h_{n-1} r_{n-1} = \prod_{k=0}^{n-1} r_k, \quad 1 \leq n \leq N,$$

and additionally,

$$h_{N+1} = c_N h_N + a_N (h_N - h_{N-1}).$$

We remark that in the special case that

$$c_0 = \cdots = c_{N-1} = 0,$$

by induction, it is easy to check that

$$r_0 = \cdots = r_{N-1} = 1$$

and hence

$$h_0 = \cdots = h_N = 1.$$

Furthermore, $h_{N+1} = c_N$. Thus, in this special case, we simply ignore the sequence $\{h_k\}$ but replace c_N by b_N . Note that here we use all of the three sequence $\{a_k\}$, $\{b_k\}$ and $\{c_k\}$ given in Q but no extra thing. The role of the sequence $\{h_k\}$ is reducing the former case to the last special one and keep the same spectrum, in terms of the H -transform \tilde{Q} :

$$\tilde{Q} = \text{Diag}(h_i)^{-1} Q \text{Diag}(h_i). \quad (9)$$

The maximal eigenpair $(\rho(Q), g)$ is transformed to $(\rho(\tilde{Q}) = \rho(Q), \text{Diag}(h_i)^{-1} g)$.

Step 3. In view of Step 2 above, it suffices to consider the following matrix

$$Q = \begin{pmatrix} -b_0 & b_0 & 0 & 0 & \cdots \\ a_1 & -(a_1 + b_1) & b_1 & 0 & \cdots \\ 0 & a_2 & -(a_2 + b_2) & b_2 & \cdots \\ \vdots & \vdots & \ddots & \ddots & \ddots \\ 0 & 0 & 0 & a_N & -(a_N + b_N) \end{pmatrix}, \quad (10)$$

where $a_i, b_i > 0$. This step is changed from the original, where everything we are working here is transfer into the original matrix Q rather than the simpler one here. It seems a direct treatment of the present matrix Q is slightly simpler.

Define the sequence (μ_i) as usual:

$$\mu_0 = 1, \quad \mu_n = \mu_{n-1} \frac{b_{n-1}}{a_n} = \frac{b_0 b_1 \cdots b_{n-1}}{a_1 a_2 \cdots a_n}, \quad 1 \leq n \leq N.$$

Next, define

$$\varphi_n = \sum_{k=n}^N \frac{1}{\mu_k b_k}, \quad 0 \leq n \leq N. \quad (11)$$

and

$$\delta_1 = \max_{0 \leq n \leq N} \left[\sqrt{\varphi_n} \sum_{k=0}^n \mu_k \sqrt{\varphi_k} + \frac{1}{\sqrt{\varphi_n}} \sum_{n+1 \leq j \leq N} \mu_j \varphi_j^{3/2} \right]. \quad (12)$$

Having these preparations at hand, we can now start our iterations.

Step 4. As in [5; §3], choose

$$w^{(0)} = \sqrt{\varphi}, \quad v^{(0)} = w^{(0)} / \|w^{(0)}\|_{\mu,2}, \quad z^{(0)} = \delta_1^{-1}, \quad (13)$$

where $\|\cdot\|_{\mu,2}$ denotes the $L^2(\mu)$ -norm. Note that here in the non-symmetric case, the use of the measure (μ_i) cannot be ignored since in this case, we are based on, δ_k for instance, the $L^2(\mu)$ setup.

Step 5. For given $v = v^{(n-1)}$ and $z = z^{(n-1)}$, let $w = w^{(n)}$ solve the linear equation

$$(-Q - zI)w = v \quad (14)$$

and then define $v^{(n)} = w / \|w\|_{\mu,2}$. An explicit solution of this w is now available, refer to [6; Algorithm 3].

Step 6. At the k th ($k \geq 1$) iteration, in addition to the one $(v^{(k)}, -Qv^{(k)})_{\mu,2}$ used in [5; §3], one may also adopt $z^{(k)} = \delta_k^{-1}$:

$$\delta_k = \max_{0 \leq i \leq N} \frac{1}{v_i^{(k)}} \left[\varphi_i \sum_{j=0}^i \mu_j v_j^{(k)} + \sum_{i+1 \leq j \leq N} \mu_j \varphi_j v_j^{(k)} \right]. \quad (15)$$

This is the main new point in the modified algorithm. Since [4; Theorems 2.4 (3), 3.2 (1) and (3.6)], we have

$$\delta_k^{-1} \leq \lambda_{\min}(-Q) \leq (v^{(k)}, -Qv^{(k)})_{\mu,2} \text{ for each } k \text{ and } n.$$

By [5; Proposition 23] and [4; Theorem 3.2 (1)], we have known that the sequence $\{\delta_k^{-1}\}$, deduced in the theorem just cited using the approximating eigenvectors obtained by the ordinary inverse iteration (without shift), is increasing to $\lambda_{\min}(-Q)$. It should be clear that the present sequence $\{\delta_k^{-1}\}$ produced by the advanced shifted inverse iteration should converge to $\lambda_{\min}(-Q)$ more faster. Thus the new $z^{(k)}$ ($k \geq 1$) not only avoids the dangerous region but may also accelerate the convergence of the algorithm. Certainly, the computation of δ_k needs more work than the one of $(v^{(k)}, -Qv^{(k)})_{\mu,2}$.

The use of the quantity (15) is motivated from the remark above on “Comparison of Algorithms 1 and 4 with the one given in [5; §3]”. The formula (15) is a corollary of [4; Theorem 2.4 (3)] which depends on the form (10) of Q . For general Q such the one in Step 1, we do not have an analog of [4; Theorem 2.4 (3)], and so (15) is not applicable in such a general situation.

Step 7. To go back to the original matrix A , denote its maximal eigenpair by $(\rho(A), g)$. Recall that the matrix Q at the beginning is obtained from A by a shift: $Q = A - mI$, $m := \max_i \sum_j a_{ij}$. Let (z, v) be the output from the last iteration in Step 6. Then we have

$$\rho(A) \approx m - z, \quad g \approx \text{Diag}(h_i)v. \quad (16)$$

We now summery the above discussions as a modified algorithm.

Algorithm 17 For tridiagonal matrix, the Step 1–Step 7 above consist a modified algorithm of the one introduced in [5; §3].

We are now ready to study a randomly chosen example, introduced to the author by Tao Tang, to justify the power of our algorithms and also to compare their efficiency.

Example 18 Let

$$A = \begin{pmatrix} 2.334 & 0.9962 & 0 & 0 & 0 & 0 \\ 0.5142 & 2.6725 & 0.1111 & 0 & 0 & 0 \\ 0 & 0.2115 & 2.263 & 0.1405 & 0 & 0 \\ 0 & 0 & 0.8442 & 2.8457 & 0.7595 & 0 \\ 0 & 0 & 0 & 0.2347 & 2.2257 & 0.0781 \\ 0 & 0 & 0 & 0 & 0.9837 & 2.1582 \end{pmatrix}.$$

Then the eigenvalues of A are

$$3.26753, 3.16247, 2.40182, 2.12632, 1.80416, 1.73679.$$

The outputs of our algorithms are given in the table below.

Table 10. Comparison of four algorithms

Algorithm	$z^{(1)}$	$z^{(2)}$	$z^{(3)}$	$z^{(4)}$	$z^{(5)}$
Algorithm 1	3.30193	3.26737	3.26754	3.26753	
Algorithm 2	3.64033	3.32623	3.26937	3.26756	3.26753
Algorithm 17a	3.2618	3.26752	3.26753		
Algorithm 17b	3.27947	3.2685	3.26754	3.26753	

where the algorithms in the last two lines mean that

Algorithm 17a: take $z^{(k)} = (v^{(k)}, -Qv^{(k)})_{\mu,2}$ for each $k \geq 1$.

Algorithm 17b: take $z^{(k)} = \delta_k^{-1}$ defined by (15) for each $k \geq 1$.

Proof. To apply Algorithm 17, take $m = 4.4494$. Then $Q = A - mI$:

$$Q = \begin{pmatrix} -2.1154 & 0.9962 & 0 & 0 & 0 & 0 \\ 0.5142 & -1.7769 & 0.1111 & 0 & 0 & 0 \\ 0 & 0.2115 & -2.1864 & 0.1405 & 0 & 0 \\ 0 & 0 & 0.8442 & -1.6037 & 0.7595 & 0 \\ 0 & 0 & 0 & 0.2347 & -2.2237 & 0.0781 \\ 0 & 0 & 0 & 0 & 0.9837 & -2.2912 \end{pmatrix}.$$

We have $h = (2.12347, 29.3339, 453.284, 924.514, 24961)$. The H -transform of Q becomes

$$\tilde{Q} = \begin{pmatrix} -2.1154 & 2.1154 & 0 & 0 & 0 & 0 \\ 0.242151 & -1.7769 & 1.53475 & 0 & 0 & 0 \\ 0 & 0.0153104 & -2.1864 & 2.17109 & 0 & 0 \\ 0 & 0 & 0.0546316 & -1.6037 & 1.54907 & 0 \\ 0 & 0 & 0 & 0.115072 & -2.2237 & 2.10863 \\ 0 & 0 & 0 & 0 & 0.0364346 & -2.2912 \end{pmatrix}.$$

Then we are ready to use Algorithm 17 for the maximal eigenpair of \tilde{Q} and finally return to the one for A by (16). \square

To explain the word “modified” in detail, we transfer Algorithm 17 to the one presented in [5; §3]. To do so, we keep the notation Q , μ , φ , δ_1 and so on used in [5; §3], but add superscript \sim to those notation used in Steps 3, 4 above. Let $\tilde{\mu} = h^2\mu$ (i.e., $\tilde{\mu}_i = h_i^2\mu_i$). Then, as mentioned in [5; §5], the mapping $f \rightarrow \tilde{f} := f/h$ gives us not only an isometry from $L^2(\mu)$ to $L^2(\tilde{\mu})$ (i.e., $\|f\|_{\mu,2} = \|\tilde{f}\|_{\tilde{\mu},2}$), and then also an isospectrum of Q on $L^2(\mu)$ and \tilde{Q} on $L^2(\tilde{\mu})$:

$$(f, Qf)_\mu = (\tilde{f}, \tilde{Q}\tilde{f})_{\tilde{\mu}}, \quad \|f\|_{\mu,2} = 1.$$

Now, from $L^2(\tilde{\mu})$ to $L^2(\mu)$, we have

$$\tilde{\varphi}_n = \sum_{k=n}^N \frac{1}{\tilde{\mu}_k \tilde{b}_k} \rightarrow \sum_{k=n}^N \frac{1}{h_k h_{k+1} \mu_k b_k} = \varphi_n, \quad 0 \leq n \leq N.$$

Here the transform $\tilde{\mu}_k \tilde{b}_k \rightarrow h_k h_{k+1} \mu_k b_k$ for each $k \leq N-1$ is regular, except the last term in the sum $(\tilde{\mu}_N \tilde{b}_N)^{-1}$, where \tilde{b}_N is actually the element \tilde{c}_N which is obtained from the transform $Q \rightarrow \tilde{Q}$, and h_{N+1} and b_N are specified in [5; §3] to make the unified expression in the second sum. We mention here that h_{N+1} is the original paper [5] should be replaced by

$$h_{N+1} = c_N h_N + a_N (h_N - h_{N-1})$$

since the sequence (c_i) used in [5] and [7] have different sign. Next,

$$\begin{aligned} \tilde{\delta}_1 &= \max_{0 \leq n \leq N} \left[\sqrt{\tilde{\varphi}_n} \sum_{k=0}^n \tilde{\mu}_k \sqrt{\tilde{\varphi}_k} + \frac{1}{\sqrt{\tilde{\varphi}_n}} \sum_{n+1 \leq j \leq N} \tilde{\mu}_j \tilde{\varphi}_j^{3/2} \right] \\ \rightarrow \delta_1 &= \max_{0 \leq n \leq N} \left[\sqrt{\varphi_n} \sum_{k=0}^n \mu_k h_k^2 \sqrt{\varphi_k} + \frac{1}{\sqrt{\varphi_n}} \sum_{n+1 \leq j \leq N} \mu_j h_j^2 \varphi_j^{3/2} \right]. \end{aligned}$$

At the same time,

$$\begin{aligned} (-\tilde{Q} - \tilde{z}I)\tilde{w} &= \tilde{v} \\ \iff (-\text{Diag}(h)^{-1}Q \text{Diag}(h) - \tilde{z}I)\tilde{w} &= \tilde{v} \\ \iff (-Q - \tilde{z}I)\text{Diag}(h)\tilde{w} &= \text{Diag}(h)\tilde{v} \\ \iff (-Q - zI)w &= v. \end{aligned}$$

Here in the last line, \tilde{z} is replaced by z , this is due to the isospectrum: an lower bound of the spectrum of $-\tilde{Q}$ is also the one of $-Q$. The fact that $\text{Diag}(h)\tilde{w} = w$ comes from the definition of our mapping $f \rightarrow \tilde{f}$. Finally, since the isometry, we have $\|w\|_{\mu,2} = \|\tilde{w}\|_{\tilde{\mu},2}$. We have thus deduced the algorithm presented in [5; §3] from the modified one.

4.5 Modification of the algorithm introduced in [5; §4.2]

In parallel to §4.4, we may introduce a modification of the algorithm presented in [5; §4.2]. The main idea is: once we obtain the function h , it can be ignored since we can use the general transform \tilde{Q} defined in (9) instead of the original Q to continue the procedure of the algorithm constructed in [5; §4.2]. Since this modification is only a mimic of the one for tridiagonal matrix (§4.4), something may be lost. For instance, the sequence $\{\delta_k^{-1}\}$ formally defined by (15) may no longer be the lower bound of $\lambda_{\min}(-Q)$, one has to take care in practice.

To conclude this paper, we remark some possible extension of the algorithms given here to a more general setup. For a larger class of Markov generators, the algorithms are meaningful. Actually, the Perron–Frobenius property as well as the the Collatz–Wielandt formula have been generalized by a number of authors. In particular, the part of the Collatz–Wielandt formula used in Algorithm 5 as $z^{(n)}$ was extended by [9; $\psi_2(V)$ in the Theorem].

See also [13; (1.1) i) and §2] and more recently, [1; Theorem 2.1]. Note the difference: we are working on $\lambda_{\min}(-L)$ here rather than $\lambda_{\max}(L)$ in the cited papers.

In the nonlinear case, the shifted inverse iteration (Algorithms 2 or 5) is more essential, actually Algorithm 1 may no longer be applicable since equation (2) often has no real solution. This point is illustrated in [6] where the shift is based on a generalization of (15). In view of [2; Theorem 2.3 and Corollary 2.5], it seems that Algorithm 2 and its variations could be applied to a more general setup.

Acknowledgments The author thanks Ms Yue-Shuang Li for her assistance in computing the large matrices using MatLab, and also pointed out the error on h_{N+1} mentioned in §4.4. The author also acknowledges Mr Xu Zhu for constructing Example 18 which leads us to find out the error just mentioned. Research supported in part by National Natural Science Foundation of China (No. 11626245), the project from the Ministry of Education in China, and the Project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions.

References

- [1] Arapostathis, A., Borkar, V.S. and Kumar, K.S. (2016). *Risk-sensitive control and an abstract Collatz–Wielandt formula*. J. Theor. Probab. 29(4), 1458–1484.
- [2] Chang, K.C. (2014). *Nonlinear extensions of the Perron–Frobenius theorem and the Krein–Rutman theorem*. J. Fixed Point Theory Appl. 15, 433–457.
- [3] Chen, M.F. (2004). *From Markov Chains to Non-Equilibrium Particle Systems*. World Scientific, Singapore, 2nd Ed. (1st Ed., 1992).
- [4] Chen, M.F. (2010). Speed of stability for birth–death processes. Front. Math. China 5(3), 379–515.
- [5] Chen, M.F. (2016). *Efficient initials for computing the maximal eigenpair*. Front. Math. China 11(6): 1379–1418. See also volume 4 in the middle of the author’s homepage:

<http://math0.bnu.edu.cn/~chenmf>

A package based on the paper is available on CRAN now. One may check it through the link:

<https://cran.r-project.org/web/packages/EfficientMaxEigenpair/index.html>

- [6] Chen, M.F. (2017) *Efficient algorithm for principal eigenpair of discrete p-Laplacian*. Preprint.
- [7] Chen, M.F. and Zhang, X. (2014) *Isospectral operators*. Commu Math Stat 2, 17–32.
- [8] Chen, R.R. (1997). *An Extended Class of Time-Continuous Branching Processes*. J. Appl. Probab. 34(1), 14–23
- [9] Donsker, W.D. and Varadhan, S.R.S. (1975). *On a variational formula for the principal eigenvalue for operators with maximum principle*. Proc. Natl. Acad. Sci. 72(3), 780–783.
- [10] Noutsos, D. (2006). *On Perron-Frobenius property of matrices having some negative entries*. Linear Algebra Appl. 412, 132–153.

- [11] Noutsos, D. (2008). *Perron Frobenius theory and some extensions*.
<http://www.pdfdrive.net/perron-frobenius-theory-and-some-extensions-e10082439.html>
- [12] Noutsos, D. and Varga, R.S. (2012). *On the Perron–Frobenius theory for complex matrices*. Linear Algebra and its Applications 437, 1071–1088.
- [13] Sheu, S.J. (1984). *Stochastic control and principal eigenvalue*. Stochastics 11(3–4), 191–211.

Mu-Fa Chen

School of Mathematical Sciences, Beijing Normal University, Laboratory of Mathematics and Complex Systems (Beijing Normal University), Ministry of Education, Beijing 100875, The People's Republic of China.

E-mail: mfchen@bnu.edu.cn

Home page: http://math0.bnu.edu.cn/~chenmf/main_eng.htm