

Production of muons from heavy-flavour hadron decays at forward rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV

Zuman Zhang ^{1,2} for the ALICE Collaboration

¹ Key Laboratory of Quark and Lepton Physics (MOE) and Institute of Particle Physics, CCNU, Wuhan, China

² Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France

Abstract

The measurement of the production of single muons from heavy-flavour hadron decays at forward rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV collected in 2015 is presented as a function of transverse momentum (p_T) and collision centrality. A strong suppression of the yield is observed at high p_T in the most central collisions compared to the binary-scaled expectation from pp collisions at the same energy. The p_T -integrated nuclear modification factor (R_{AA}) as function of the number of participating nucleons indicates an increase of the suppression from peripheral to central collisions. Comparisons with the results for Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV and with transport model predictions are shown. A similar suppression is measured at both $\sqrt{s_{NN}} = 2.76$ TeV and 5.02 TeV. The results available from $p_T = 7$ GeV/c up to $p_T = 20$ GeV/c, will provide new constraints on transport model ingredients and new insights on the understanding of the evolution of the hot and dense matter formed in ultra-relativistic heavy-ion collisions.

Keywords:

LHC, ALICE, ultra-relativistic heavy-ion collisions, open heavy flavour production, nuclear modification factor

1. Introduction

ALICE [1] is the dedicated heavy-ion experiment at the LHC, optimized to investigate the properties of strongly-interacting matter under extreme conditions of temperature and energy density where the formation of the Quark-Gluon Plasma (QGP) is expected. Heavy quarks (charm and beauty) are regarded as efficient probes of the properties of the QGP as they are created on a very short time scale in initial hard parton scattering processes and subsequently interact with the medium. Open heavy flavours are expected to be sensitive to the energy density of the system through the in-medium energy loss of heavy quarks. In the high p_T region, the suppression of the heavy-flavour yields is quantified by means of the nuclear modification factor, R_{AA} :

$$R_{AA}(p_T) \equiv \frac{d^2 N_{AA}/dp_T dy}{\langle T_{AA} \rangle d^2 \sigma_{pp}/dp_T dy} \quad (1)$$

It is defined as the ratio of the particle yield measured in Pb–Pb collisions to the p_T -differential cross section measured in pp collisions scaled by the average nuclear overlap function. Due to the color-charge effect, the radiative energy loss of gluons should be larger than that of quarks. Moreover, heavy-quark energy loss may be reduced with respect to that of light quarks due to the dead-cone effect [2].

ALICE detector and data sample

We focus on the measurement of the nuclear modification factor of muons from open heavy-flavour hadron decays with the ALICE muon spectrometer in the pseudo-rapidity range of $-4 < \eta < -2.5$. The ALICE muon spectrometer consists of a thick

front absorber, a beam shield, a dipole magnet, five tracking stations and two trigger stations behind an iron wall. In addition to the muon spectrometer, for the analysis presented here, the V0 detector comprising two scintillator arrays and the Silicon Pixel Detector (SPD) are used. Collisions were classified according to their centrality, determined from the sum of the amplitudes of the signals in the V0 detector and defined in terms of percentiles of the total hadronic Pb–Pb cross section. The event vertex is reconstructed with the SPD.

The analysis uses the Pb–Pb data sample collected in 2015 at $\sqrt{s_{NN}} = 5.02$ TeV and is based on muon triggered events. The trigger condition consists of a signal in the two V0 arrays (minimum bias trigger) with at least one candidate track with a high p_T threshold of about 4.2 GeV/c in the muon trigger system. The integrated luminosity is $202 \pm 6 \mu\text{b}^{-1}$ in the 0–90% centrality class.


Muons are reconstructed within the acceptance of the muon spectrometer. Then, it is required that the track candidate in the tracking system matches a track reconstructed in the muon trigger system. This condition allows us to reject most of the punch-through hadrons that are stopped in the iron wall. Furthermore, the correlation between the track momentum and the geometrical distance of closest approach to the primary vertex is used in order to remove fake tracks and tracks from beam-gas interactions. The remaining background for $p_T > 2$ GeV/c after these selection cuts consists of muons from decays of primary light hadrons (π and K, main contribution at low and intermediate p_T , called decay muons in the following) and muons from decays of W and Z/ γ^* (main contribution at high p_T).

2. Analysis strategy


$$\frac{dN_{AA}^{\mu \leftarrow HF}}{dp_T} = \frac{dN_{AA}^{incl \mu}}{dp_T} - \frac{dN_{AA}^{\mu \leftarrow K, \pi, W, Z/\gamma^*}}{dp_T} \quad (2)$$

The analysis strategy in Pb–Pb collisions includes the following steps: normalization of muon samples to equivalent number of minimum-bias (MB) events on a run by run basis, acceptance \times efficiency correction and background subtraction (Eq. 2). The acceptance \times efficiency correction is obtained from realistic GEANT3 simulations using p_T and rapidity distributions of muons from heavy-flavour signals by NLO pQCD calculations. The centrality dependence of the tracking efficiency is estimated via an embedding procedure.

The background of π and K decay muons that needs to be subtracted is evaluated by a data-tuned




ALI-PREL-113634



ALI-PREL-113627

Figure 1: p_T -differential R_{AA} of heavy-flavour hadron decay muons at forward rapidity in different centrality classes.



ALI-PREL-113666

Figure 2: R_{AA} of heavy-flavour decay muons as a function of the mean number of participating nucleons for different transverse momentum ranges.

Monte-Carlo cocktail. The fraction of π and K decay muons is about 3% in $7 < p_T < 10$ GeV/c. The contribution of W and Z/γ^* [3] decay muons in Pb–Pb templates is obtained by combining simulations of W and Z/γ^* decay muons in pp, pn, np and nn collisions with POWHEG. The fraction of W and Z/γ^* decay muons increases with p_T and is about 38% (19%) in the 0–10% (60–80%) centrality class at $p_T = 20$ GeV/c.


The study of in-medium effects with the R_{AA} observable requires a pp reference. The latter was obtained from the published p_T -differential cross section of heavy-flavour hadron decay muons at $\sqrt{s} = 7$ TeV measured in $2 < p_T < 12$ GeV/c [4] scaled to $\sqrt{s} = 5.02$ TeV using simulations based on Fixed-Order-Next-to-Leading-Log (FONLL) [5] calculations. The p_T -differential cross section of heavy-flavour decay muons in $p_T > 12$ GeV/c is obtained by scaling FONLL predictions according the ratio between data and FONLL.

The systematic uncertainty on the nuclear modification factor of muons from heavy-flavour hadron decays is affected by the systematic uncertainties of the inclusive muon yield, the background subtraction, the pp reference and the normalization procedure. The final systematic uncertainty on the p_T -differential R_{AA} is obtained by adding in quadrature these contributions. It varies from about 11% ($p_T = 7$ GeV/c) to about 22% ($p_T = 20$ GeV/c) in central collisions.

3. Results


Figure 1 shows the p_T -differential R_{AA} of heavy-flavour decay muons in the p_T interval $7 < p_T < 20$ GeV/c, in the centrality classes 0–10%, 10–20%, 20–40%, 40–60% and 60–80%. The vertical error bars represent the statistical uncertainty. The open boxes include the systematic uncertainty sources except the systematic uncertainty on normalization which is shown as a filled box at $R_{AA} = 1$. A stronger suppression is observed in central collisions compared to peripheral collisions, with no significant p_T dependence within uncertainties. The suppression reaches a factor of about three in the 10% most central collisions.

Figure 2 shows the centrality dependence of R_{AA} of heavy-flavour hadron decay muons which is studied as a function of the average number of participating nucleons ($\langle N_{part} \rangle$). The suppression increases with increasing centrality in the domain $7 < p_T < 20$ GeV/c. The results are compatible with those obtained in Pb–Pb




ALI-PREL-113642

Figure 3: p_T -differential R_{AA} of heavy-flavour hadron decay muons at forward rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ compared with the measurement at $\sqrt{s_{NN}} = 2.76$ TeV.



ALI-PREL-113670



ALI-PREL-113674

Figure 4: R_{AA} of muons from heavy-flavour hadron decays in $2.5 < y < 4$, compared with different model predictions.

collisions at $\sqrt{s_{NN}} = 2.76$ TeV [6], although in a slightly different p_T interval.

Figure 3 presents a comparison of the p_T -differential R_{AA} of heavy-flavour hadron decay muons at forward rapidity in Pb–Pb collisions at 5.02 TeV and 2.76 TeV. A similar suppression is observed at 5.02 TeV and at 2.76 TeV within uncertainties.

Figure 4 presents a comparison of the R_{AA} of muons from heavy-flavour decays in $2.5 < y < 4$ with transport models in different centrality classes. The Vitev model [7] (Fig. 4, top) describes the R_{AA} of heavy-flavour hadron decay muons in central collisions. The TAMU model [8] tends to overestimate the R_{AA} of muons from heavy-flavour hadron decays in central collisions and reproduces the measurement in peripheral collisions. These R_{AA} measurements at $\sqrt{s_{NN}} = 5.02$ TeV provide new constraints on energy loss models.

In addition to final-state effects where in-medium energy loss would be dominant, initial-state effects could influence the R_{AA} measurement. In the kinematic range relevant for the study of the production of muons from heavy-flavour hadron decays, the main expected effects are nuclear modification of parton distribution functions [9], energy loss [10] and k_T broadening via multiple soft scatterings in the initial state [11]. It is worth noting that since the corresponding measured R_{pPb} is consistent with unity over the whole p_T range [12], the suppression observed at high p_T in central Pb–Pb collisions results from final-state effects related to parton energy loss [2] in-medium.

4. Summary

We presented the first measurement of the nuclear modification factor of open heavy flavours with ALICE via single muons measured at forward rapidity ($2.5 < y < 4$) in a wide p_T range ($7 < p_T < 20$ GeV/ c) in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV collected in the LHC Run2. A strong suppression in the 10% most central collisions reaching a factor of about three in $7 < p_T < 12$ GeV/ c is observed. Results are compatible within uncertainties with those obtained at $\sqrt{s_{NN}} = 2.76$ TeV. The measured suppression is due to final-state effects. The results of R_{AA} in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV in various centrality classes may provide important constraints to the models.

Acknowledgement

This work is supported partly by the NSFC 11375071, the National Basic Research Program of

China (2013CB837803) and NSFC IRG11521064 and 11475068.

References

- [1] Aamodt K, et al., ALICE Collaboration, JINST 3 (2008) S08002
- [2] Y. Dokshitzer and D. Kharzeev, Phys. Lett. B 519 (2001) 199
- [3] S. Chatrchyan, et al., CMS Collaboration, Phys. Lett. B 715 (2012) 66–87
- [4] R. Averbeck et al., ALICE Collaboration, Phys. Lett. B 708 (2012) 265
- [5] R. Averbeck et al., ALICE Collaboration, arXiv:1107.3243
- [6] B. Abelev, et al., ALICE Collaboration, Phys. Rev. Lett. 109 (2012) 112301
- [7] R. Sharma and B. W. Zhang, Phys. Rev. C 80 (2009) 054902
- [8] M. He, R. J. Fries, R. Rapp, Phys. Lett. B 735 (2014) 445
- [9] H. Fujii and K. Watanabe, Nucl. Phys. A 915 (2013) 1
- [10] I. Vitev, Phys. Rev. C 75 (2007) 064906
- [11] X. N. Wang, Phys. Rev. C 61 (2000) 064910
- [12] S. Li, Nucl. Phys. A 931 (2014) 546–551