

A new method to study the number of colors in the final-state interactions of hadrons

Ling-Yun Dai¹, Ulf-G. Meißner^{2,1}

¹ *Institut für Kernphysik, Institute for Advanced Simulation and Jülich Center for Hadron Physics, Forschungszentrum Jülich, D-52425 Jülich, Germany*

² *Helmholtz Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany*

Abstract

Based on Chiral Perturbation Theory we introduce the dependence on the number of colors (N_C) for the $\pi\pi \rightarrow \pi\pi$ scattering amplitudes. Those amplitudes are calculated from dispersion relations that respect analyticity and coupled channel unitarity, as well as accurately describing experiment. By varying N_C the trajectories of the poles and residues (the couplings to $\pi\pi$) of the light mesons, the $\sigma/f_0(500)$, $f_0(980)$, $\rho(770)$ and $f_2(1270)$ are investigated. Our results show that the method proposed is a reliable way to study the N_C dependence in hadron-hadron scattering with final-state interactions.

Keywords: Dispersion relations, Partial-wave analysis, Chiral Lagrangian, meson production

The lightest scalar mesons are rather interesting as they have the same quantum numbers as the QCD vacuum. The nature of them is still a mystery [1]-[4]. The phenomenology of these is complicated due to the contribution from important final-state interactions (FSI) [5, 6]. Dispersion relations are the natural way to include FSI, see e.g. [7, 8]. For some of the light mesons, like the σ , κ , their existence has been confirmed [9]-[12] and accurate pole locations and $\pi\pi$ couplings, including also the $\rho(770)$, have been given in Refs. [13, 14, 15]. Concerning the nature of the scalar mesons, there are a cornucopia of models [16]-[35]. Among them tracking the large N_C trajectories of the poles is an effective diagnostics to distinguish ordinary from non-ordinary quark-antiquark structure as considered in [36]-[40]. However, these analyses based on unitarized Chiral Perturbation theory ($U\chi PT$) lack crossing symmetry. Unitarization itself will also generate spurious poles [41] and cuts [42, 40]. In contrast, dispersion relations respect analyticity, but including coupled channel unitarity and the N_C dependence is difficult. Clearly, both analyticity and coupled channel unitarity are critical in the region of the $\bar{K}K$ threshold, close to which the $f_0(980)$ is located. To solve this problem,

we use an Omnès representation based on the phase of the relevant amplitudes, rather than the elastic phase shift [43, 44]. There has been renewed interest in the study of the large N_C limit [45, 46] of the properties of resonances [47, 48, 49]. Weinberg [50] pointed out that resonant tetraquark states could exist due to the contribution of the leading order (LO) ‘connected’ diagrams to the Green functions. Their widths are $O(N_C^{-1})$, as narrow as ordinary mesons. They could be even narrower, with width of $O(N_C^{-2})$, when the flavor of the quarks is combined in different ways [51]. The width could also be wide as $O(1)$, see [52]. There are many other interesting discussions such as [53, 54] and references therein. In this paper we focus on establishing a ‘practical’ way to study the N_C dependence of the scattering amplitudes, built into dispersion relations. Resonances appearing in the intermediate states are also studied.

In this letter we first use dispersive methods to obtain the $\pi\pi$ scattering amplitudes up to 2 GeV. We construct the amplitudes in a model-independent way, which is both analytic and respects coupled channel unitarity. We also recalculate the analytical expressions of the $IJ = 00, 02, 11$ waves in $SU(3)$ Chiral Perturbation Theory (χPT) up to $O(p^4)$. By matching with the χPT amplitudes up to $O(N_C^{-1})$, we introduce the N_C -dependence into the dispersive amplitudes. This N_C dependence is automatically transferred to the high-energy region,

Email addresses: l.dai@fz-juelich.de (Ling-Yun Dai¹),
meissner@hiskp.uni-bonn.de (Ulf-G. Meißner^{2,1})

where the FSI are implemented by a dispersion relation. We give the trajectories of the poles and residues by varying N_C . The behavior of the $\rho(770)$, $f_2(1270)$, $\sigma/f_0(500)$ and $f_0(980)$ show that this is a reliable way to study the number of colors in hadron-hadron scattering. The N_C trajectory of the light scalar mesons supports a mixed structure of hadronic molecule and $\bar{q}q$ components (for a recent review on hadronic molecules, see Ref. [55]), while a tetra-quark component is also possible.

We first present our $IJ = 00, 02, 11$ partial waves of $\pi\pi \rightarrow \pi\pi$ calculated in a model-independent way. We start from:

$$T_J^I(s) = P_J^I(s)\Omega_J^I(s). \quad (1)$$

where $\Omega_J^I(s)$ is the Omnès function [56]:

$$\Omega_J^I(s) = \exp\left(\frac{s}{\pi} \int_{s_{th}}^{\infty} ds' \frac{\varphi_J^I(s')}{s'(s' - s)}\right). \quad (2)$$

with $\varphi_J^I(s)$ the phase of the partial wave amplitude $T_J^I(s)$, which has been given in previous amplitude analysis [43, 44]. This phase is known from experiment up to roughly 2 GeV, while in the higher energy region it is constrained by unitarity. The Omnès function is truncated at $s = 22 \text{ GeV}^2$. The function $P_J^I(s)$ includes the effect of the left-hand cut (l.h.c) and corrections that come from the distant right-hand cut (r.h.c) above 2 GeV. The latter one has a tiny contribution to the region where the light mesons appear. Other information is provided by chiral dynamics that fixes the Adler zero in the S-wave, and the approach to the threshold of the S-, P-, and D-waves in terms of scattering lengths and effective ranges. We therefore parameterize the $P_J^I(s)$ as

$$P_J^I(s) = (s - z_J^I)^{n_J} \sum_{k=1}^n \alpha_{Jk}^I (s - 4M_\pi^2)^{k-1}, \quad (3)$$

where z_J^I is the Adler zero for the S-wave and $4M_\pi^2$ for P- and D-waves. The parameter n_J is 1 for S- and P-waves and 2 for D-waves. The parameters α_i are given in Table 1. The units of the α_k are chosen to ensure that the amplitude $T_J^I(s)$ is dimensionless.

The fit results are shown in Fig. 1. We fit the amplitudes in the region of $s \in [0, 4\text{GeV}^2]$, where the ‘data’ is as follows: χ PT amplitudes at $[0, 4M_\pi^2]$, amplitudes of K-matrix and Roy-like equation at $[4M_\pi^2, 2\text{GeV}^2]$, and experiment data up to 4GeV^2 . The fits are of high quality, even in our ‘prediction’ region where $s \in [-4M_\pi^2, 0]$. In this region the real part of our amplitudes is in good agreement with that of χ PT ($O(p^4)$), and the imaginary part certainly vanishes. Notice that the imaginary part of the χ PT amplitudes is rather small, too.

	T_{11S}^0	T_{11D}^0	T_{11P}^1
α_1	2.4051	0.2972	0.4283
α_2	-1.9451	-0.9354	-0.2976
α_3	1.3473(36)	2.1931(2)	0.6173(16)
α_4	-0.4629(19)	-2.6108(2)	-0.7092(11)
α_5	0.0038(6)	1.6508(1)	0.3774(4)
α_6	0.0307(2)	-0.5679(1)	-0.0909(1)
α_7	-0.0045(1)	0.1004(1)	0.0081(1)
α_8	—	-0.0071(1)	—

Table 1: The fit parameters as given in Eq. (3). The errors are given by MINUIT and notice that the $\alpha_{1,2}$ are fixed by the scattering lengths and slope parameters [44, 57, 58].

There is no l.h.c in our parametrization¹, but its contribution to the shaded region on the complex s -plane, as shown in Fig. 2, are properly implemented, as we fit our amplitudes to the data as well as the amplitudes of Roy equation [10] in the presence of crossing symmetry². We note that the amplitudes on the upper half of s -plane are readily obtainable from the ones on the lower side according to the Schwarz reflection principle. From Fig. 2 we see that our amplitudes are compatible to that of the Roy equation analysis in the complex s -plane. The distribution of contours is in good agreement and moreover, their gradient variations are compatible with each other, as shown by the shading of the color from blue to red. Nevertheless, amplitudes on the edge of the domain and in the region of $\text{Im}s < -0.3 \text{ GeV}^2$ for $T_S^0(s)$ are less consistent with differences ≤ 0.1 . The difference around $\sqrt{s} \approx 1 \text{ GeV}$ of $T_S^0(s)$ are also not ignorable, this is caused by the different treatment of the $K\bar{K}$: ours include the physical K^+K^- and $K^0\bar{K}^0$ [43], while it is treated as $K\bar{K}$ in the isospin basis in the analysis of the Roy equations [10].

With these amplitudes, we can extract the poles and residues on the second sheet. The residue $g_{f\pi\pi}$ and pole s_R on the second Riemann sheet are defined as

$$T^{II}(s) = \frac{g_{f\pi\pi}^2}{s_R - s}. \quad (4)$$

The pole locations and their residues are listed in Ta-

¹It is worth to note that in [13] the mass and width of σ deviates about 15% from the original value when the l.h.c is removed.

²The comparison of the I=1 P-wave amplitudes has already been given in [59]. However, we present it here for convenience. Notice that the D-wave is absent in the Roy equation analysis[13], and the $f_2(1270)$ is quite far away from the l.h.c. We thus do not discuss it here.




Figure 1: The fit of $\pi\pi$ scattering amplitudes are in the left column. The violet and magenta bands are from K-Matrix [43]. The olive and light grey bands in the low energy region are from χ PT [60, 61, 62, 42, 40]. The cyan and green bands are from CFDIV [57]. The CERN-Munich data is from [63]. The absolute values of the amplitudes by varying N_C are in the right column. The black solid, orange dashed, blue dotted and red dash-dotted lines are for $N_C = 3, 6, 12, 120$, respectively.

ble 2. These are very similar from those of previous analyses [13, 15, 43, 64, 65]. For the $f_0(1370)$ and $f_2(1270)$, to find the pole closest to the physical sheet one needs to include the $\pi\pi$, $\bar{K}K$, 4π as coupled channels. The pole obtained here from a single $\pi\pi$ channel is of course, not on the Riemann sheet which is closest to the physical region. Therefore, we do not discuss the $f_0(1370)$ in the next sections. For the $f_2(1270)$, $\pi\pi$ is the dominant decay channel and the pole on the second sheet is not far away from the physical one.

Having analytically calculated the partial wave amplitudes of $IJ = 00, 11, 02$ within one-loop SU(3) χ PT, we can match our dispersive results to these and so fix their N_C dependence. We note several points about this matching: First, though the matching is done in the low-energy region, this N_C dependence is transferred to high-energy region. As the FSI of hadrons in the higher energy region correspond to ‘hadron loop’ corrections, which could be translated into higher order corrections




Figure 2: Comparison of our amplitudes ($IJ = 00, 11$) with the ones from the Roy equation analysis in the domain where the Roy equations work. On the left side there are real and imaginary parts of our amplitudes, and on the right side are those from Roy equations [10].

State	pole locations (MeV)	$g_{f\pi\pi} = g_{f\pi\pi} e^{i\varphi}$	
		$ g_{f\pi\pi} $ (GeV)	φ ($^\circ$)
$\sigma/f_0(500)$	$436.2(12.2) - i260.7(6.8)$	$0.45(0.02)$	$74(2)$
$f_0(980)$	$997.7(1.1) - i21.7(1.9)$	$0.27(0.02)$	$84(3)$
$f_0(1370)$	$1431.6(34.6) - i185.4(22.4)$	$0.78(0.21)$	$-47(18)$
$f_2(1270)$	$1278.3(7.0) - i79.3(17.8)$	$0.50(0.05)$	$0.7(4.3)$
$\rho(770)$	$762.4(3.9) - i68.7(6.3)$	$0.34(0.01)$	$12(3)$

Table 2: The pole locations and residues on the second Riemann sheet.

(quark loops) in large N_C QCD, and are thus suppressed by an extra factor of N_C^{-1} . Of course, this N_C dependence is energy-dependent, see the discussion below. Second, we only calculate the χ PT amplitudes up to $O(p^4)$. Here, the imaginary part of the χ PT amplitudes is given entirely from one-loop integrals and thus both l.h.c and r.h.c are N_C^{-2} , while the real part is N_C^{-1} . For $O(p^6)$, it is still N_C^{-1} for the real part of the amplitudes as given by the contact terms, and again it is N_C^{-2} for the imaginary part. The latter one is either from the 2-loop corrections or from the 1-loop corrections with one insertion of a contact term of $O(p^4)$. This N_C dependence continues when one goes to higher orders. One thus finds that $\text{Im}T/\text{Re}T \sim 1/N_C$ and we define:

$$\varphi(s, N_C) = \arctan \left[\frac{3}{N_C} \tan \varphi(s) \right], \quad (5)$$

where $\varphi(s)$ is the phase. One notices that $\varphi(M_R^2, N_C) = 90^\circ$, where the phase is 90° at M_R with $N_C = 3$. This implies for the low-energy region:

$$\Omega_J^I(s, N_C) = \bar{\Omega}_J^I(s) + O(N_C^{-1}). \quad (6)$$

$\bar{\Omega}_J^I(s)$ is close to 1, where the phase is obtained by $N_C \rightarrow \infty$ and behaves as a step function. Third, we only match up to $O(N_C^{-1})$. In this case, Eq.(6) shows that one only needs to inject the N_C dependence as

$$P_J^I(s, N_C) = \frac{3}{N_C} P_J^I(s). \quad (7)$$

This is compatible with the fact that the real part of the χ PT amplitudes is $O(N_C^{-1})$ while the imaginary part of the χ PT amplitudes is $O(N_C^{-2})$, up to any order. For simplicity we ignore all the complicated higher order N_C dependence.

We test the stability of the amplitudes by varying N_C , shown in Fig. 1. Both the shape and the magnitude of the amplitude of the S wave vary dramatically. The peak of $|T|$ behaves as $O(1)$ for the P and D waves and the width behaves as $O(N_C^{-1})$. This is consistent with that of a Breit-Wigner description, where there is a zero of the real part amplitude ($s = M_R^2$), and the N_C dependence is canceled leading to $T \simeq i/\rho$. Thus, the N_C dependence is properly implemented. The trajectories of the pole locations and their residues $g_{f\pi\pi}$ on the second Riemann sheet are obtained from Eq. (4) and plotted in Fig. 3. To estimate the uncertainties, we choose $\frac{2}{N_C} + \frac{3}{N_C^2}$ and/or $\frac{4}{N_C} - \frac{3}{N_C^2}$ to replace $\frac{3}{N_C}$ in Eqs. (5,7). The variation of poles and residues with these different N_C dependence are collected as the uncertainty. Here we simply assume that the whole contribution of the N_C^{-2}




Figure 3: The trajectories of the pole locations and their residues by varying N_C from 3 to a large number, in steps of $\Delta N_C = 1$. The black filled circle and lines represent poles and residues, respectively. The red dotted lines are linear fits (with the variable $\sqrt{N_C}$ or $1/\sqrt{N_C}$) to the residues obtained, as explained in the text.

corrections is roughly one third of that of N_C^{-1} . The separated N_C^{-2} dependence of each polynomial in Eq. (7) is not discussed, as the complete N_C^{-2} part of the real amplitudes in χ PT is not available and these polynomials are correlated with each other. Only the first two polynomials are undoubtedly fixed due to the scattering lengths and slope parameters. We are aware that the N_C dependence of these polynomials, especially the third or fourth polynomial could have fairly large effects on the uncertainties. One needs a careful analysis on N_C^{-2} corrections for a refined analysis. The uncertainties of the poles/residues are presented as cyan bands in Fig. 3. They are compatible to the black solid line with only

N_C^{-1} dependence³. Next, we discuss the various resonances within the accuracy of our approach.

$\rho(770)$

The pole trajectory of the $\rho(770)$ moves toward the real axis, similar to what was found using $U\chi PT$ [37, 40, 39]. In Fig.3, the red dotted line represents for $|g_{\rho\pi\pi}| \times \sqrt{3/N_C}$ with $|g_{\rho\pi\pi}|$ the residue at $N_C = 3$. The black solid and red dotted lines overlap perfectly. This confirms that the modulus of the residue decreases exactly as $1/\sqrt{N_C}$. Similarly the width decreases exactly as $1/N_C$. Such behavior confirms the widely accepted $\bar{q}q$ structure.

$f_2(1270)$

For the $f_2(1270)$, the trajectory is quite similar to that of the $\rho(770)$. Again this confirms a $\bar{q}q$ structure. In Fig.3, the red dotted line represents $|g_{f_2\pi\pi}| \times \sqrt{3/N_C}$ with $|g_{f_2\pi\pi}|$ the residue at $N_C = 3$. The black solid and red dotted lines are much the same, indicating that the residue decreases exactly as $1/\sqrt{N_C}$. Similarly, the width decreases exactly as $1/N_C$.

$\sigma/f_0(500)$

For $\sigma/f_0(500)$, its mass is roughly $O(1)$ and the width is $O(N_C)$. Mixing with $O(N_C)$ or $O(N_C^{-1})$ can not be excluded. The pole moves far away from the real axis on the complex s -plane when N_C is increased. We note that the third and fourth quadrants in Eq. (3) may be important and their higher N_C^{-2} corrections may distort this trajectory. However, considering that the pole is mainly determined by the unitary cut at $N_C = 3$, and both the l.h.c and distant r.h.c are $O(N_C^{-2})$, and also in the $O(N_C^{-1})$ case the pole moves far away from the real axis (or cuts), it is natural to assume that the $1/N_C$ part of these quadrants dominates the trajectory at large N_C . At present, we can not quantify how much difference these $O(N_C^{-2})$ corrections would generate. To get such a large width it could have a molecular component [51] and/or a tetra-quark component [52]. Notice in [40, 43] the shadow pole in the third sheet suggests a $\bar{q}q$ component. So the σ might be a mixed state including molecule, $\bar{q}q$, etc. The modulus of the residue increases, reaching the peak at $N_C = 5$ rapidly and then decreases, again it increases much slower at the turning point $N_C = 17$. The red dotted line is for $0.47 + 0.013\sqrt{N_C}$, and it is essentially coincident with the black solid line where $N_C \geq 17$. This indicates that the residue should contain components with $O(N_C^{1/2})$ and $O(1)$. Also, the sharp peak implies

³We note that Ref. [67] points out the sub-leading N_C dependence of the LECs may be sizable. In [36, 68], the uncertainty caused by the regularization scale μ is discussed. These uncertainties are larger than that shown in Fig.3, particularly for scalar mesons. Here, we simply absorb all these uncertainties into an overall higher correction of $O(N_C^{-2})$.

that the residue should contain $O(N_C)$ or higher components. A mixture of $O(N_C^{-1/2})$ or lower components is also possible. Moreover, it is of interest to note that around the sharp peak (with $N_C \in [3, 7]$) the real parts of the pole locations in complex s -plane are quite close to the $\pi\pi$ threshold. The curved behavior of the trajectory is consistent with the mixing structure of molecule and $\bar{q}q$, while an extra tetra-quark component is possible.

$f_0(980)$

For the $f_0(980)$, the pole moves to the real axis, slightly below $\bar{K}K$ threshold. It is similar to that of $\rho(770)$ and $f_2(1270)$, implying an $\bar{s}s$ component. In contrast, in Refs [69, 40] the pole moves to the real axis above $\bar{K}K$ threshold and goes onto the fourth Riemann sheet. We may need higher order $1/N_C$ corrections, especially that caused by kaon loops, to obtain a more accurate pole trajectory. The residue increases at first and then decreases, as a ‘curve’. The red dotted line is $1.10/\sqrt{N_C}$, which is coincident with the black solid line where $N_C \geq 10$. This indicates that the residue must have the component of $O(N_C^{-1/2})$. Note it is most likely to be a $\bar{K}K$ molecule in other analysis [2, 40, 43]. The peak around $N_C = 5$ implies also $O(N_C^{1/2})$ or even higher components. This might indicate a possible tetra-quark component. We note that the pole locations, around the sharp peak with $N_C \in [3, 6]$, are close to the $\bar{K}K$ threshold. Our findings support the idea that the $f_0(980)$ is a mixture of $\bar{K}K$ molecular and $\bar{s}s$ components. The relative strengths of these components can not be inferred from the analysis presented here.

In this letter we presented a new method to study the large N_C behaviour of resonances. The $\pi\pi$ scattering amplitudes with FSI are constructed in a model-independent way. We match it with the amplitudes of χPT to give the N_C -dependence of the coefficients and phase in the dispersive approach. This is a reliable way to study the N_C -dependence with final-state interactions. We obtain the trajectories of the poles and residues as N_C changes. Those for both the $\rho(770)$ and the $f_2(1270)$ quantitatively support, as expected, the standard $\bar{q}q$ structure. The N_C trajectory of the light scalar mesons are obtained, especially the coupling of $g_{\sigma\pi\pi}$ has $O(N_C^{1/2})$ component and $g_{f_0(980)\pi\pi}$ has $O(N_C^{-1/2})$ component. They are consistent with each being a combination of $\bar{q}q$ and multi-hadron (molecular) components, while a tetra-quark component can not be excluded. We stress that some of these conclusions might be modified when higher order corrections in the χPT amplitudes are accounted for.

Acknowledgements

We are very grateful to M.R. Pennington for many valuable suggestions and discussions to improve the paper. Special thanks to J. Ruiz de Elvira for his thoughtful and critical reading of the manuscript. Helpful discussions with Z.H. Guo are also acknowledged. This work is supported by the DFG (SFB/TR 110, “Symmetries and the Emergence of Structure in QCD”), by the Chinese Academy of Sciences (CAS) President’s International Fellowship Initiative (PIFI) (Grant No. 2018DM0034) and the VolkswagenStiftung (Grant No. 93562).

References

- [1] M.R. Pennington, *Mod. Phys. Lett.* **A22** 1439 (2007), arXiv: 0705.3314 [hep-ph].
- [2] M.R. Pennington, *AIP Conf. Proc.* **1257** 27 (2010), arXiv: 1003.2549 [hep-ph].
- [3] R.J. Jaffe, *Phys. Rev.* **D 15** 267 (1977), *Phys. Rept.* **409** 1 (2005).
- [4] J. R. Pelaez, *Physics Reports* **658** 1 (2016), arXiv: 1510.00653
- [5] K.L. Au, D. Morgan and M.R. Pennington, *Phys. Rev.* **D 35** 1633 (1987); D. Morgan and M.R. Pennington, *Phys. Rev.* **D 48** 1185 (1993).
- [6] U.-G. Meißner, *Comments Nucl. Part. Phys.* **20**, 119 (1991).
- [7] X.W. Kang, B. Kubis, C. Hanhart and Ulf-G. Meißner, *Phys. Rev.* **D 89** 053015 (2014), arXiv: 1312.1193 [hep-ph].
- [8] Y.H. Chen, J. T. Daub, F.K. Guo, B. Kubis, Ulf-G. Meißner and B.S. Zou, *Phys. Rev.* **D 93** 034030 (2015), arXiv: 1512.03583 [hep-ph].
- [9] Z.G. Xiao and H.Q. Zheng, *Nucl. Phys.* **A695** 273 (2001), arXiv: 0011260 [hep-ph].
- [10] G. Colangelo, J. Gasser and H. Leutwyler, *Nucl. Phys.* **B603**, 125 (2001), arXiv: 0103088 [hep-ph].
- [11] Z.Y. Zhou, G. Y. Qin, P. Zhang, Z. G. Xiao, H. Q. Zheng, and N. Wu, *JHEP* **0502** 043 (2005), arXiv: 0406271 [hep-ph].
- [12] S. Descotes-Genon, B. Moussallam, *Eur. Phys. J.* **C48** 553 (2006), arXiv: [hep-ph/0607133].
- [13] I. Caprini, G. Colangelo and H. Leutwyler, *Phys. Rev. Lett.* **96** 132001 (2006), arXiv: 0512364 [hep-ph].
- [14] P. Büttiker, S. Descotes-Genon and B. Moussallam, *Eur. Phys. J.* **C33** 409 (2004), arXiv:0310283 [hep-ph].
- [15] R. García-Martín, R. Kamiński, J. R. Peláez, and J. Ruiz de Elvira, *Phys. Rev. Lett.* **107** 072001 (2011), arXiv: 1107.1635 [hep-ph].
- [16] R. L. Jaffe, *Phys. Rev.* **D 15**, 267 (1977).
- [17] E. van Beveren, T. A. Rijken, K. Metzger, C. Dullemond, G. Rupp and J. E. Ribeiro, *Z. Phys.* **C30** 615 (1986), arXiv:0710.4067 [hep-ph].
- [18] N. A. Tornqvist and M. Roos, *Phys. Rev. Lett.* **76** (1996) 1575, arXiv: hep-ph/9511210.
- [19] J. A. Oller and E. Oset, *Nucl. Phys. A* **620**, 438 (1997) hep-ph/9702314.
- [20] J. A. Oller, E. Oset and J. R. Pelaez, *Phys. Rev. D* **59**, 074001 (1999) hep-ph/9804209.
- [21] D. Black, A. H. Fariborz, F. Sannino and J. Schechter, *Phys. Rev.* **D 59** 074026 (1999), arXiv: 9808415 [hep-ph]; D. Black, A. H. Fariborz and J. Schechter, *Phys. Rev.* **D 61** 074001 (2000), arXiv: [hep-ph/9907516].
- [22] V. Baru, J. Haidenbauer, C. Hanhart, Yu. Kalashnikova, and A. Kudryavtsev, *Phys. Lett.* **B586** 53 (2004), arXiv: 0308129 [hep-ph].
- [23] N. N. Achasov, *Nucl. Phys.* **A728** 425 (2003), arXiv: 0309118 [hep-ph].
- [24] N. N. Achasov and G. N. Shestakov, *Phys. Rev. Lett.* **99** 072001 (2007), arXiv: 0704.2368 [hep-ph].
- [25] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, *Phys. Rev. Lett.* **93** 212002 (2004), arXiv:0407017 [hep-ph]; *Eur. Phys. J.* **C50**, (2007) 609, arXiv: 0604018 [hep-ph].
- [26] G. ’t Hooft, G. Isidori, L. Maiani, A.D. Polosa, and V. Riquer, *Phys. Lett.* **B662** 424 (2008), arXiv: 0801.2288 [hep-ph].
- [27] Z.Y. Zhou and Z.G. Xiao, *Phys. Rev.* **D 83** 014010 (2011), arXiv: 1007.2072 [hep-ph].
- [28] G. Mennessier, S. Narison, X.G. Wang, *Phys. Lett.* **B688** 59 (2010), arXiv: 1002.1402 [hep-ph]; *Phys. Lett.* **B696** 40 (2011), arXiv: 1009.2773 [hep-ph].
- [29] Z.H. Guo and J. A. Oller, *Phys. Rev.* **D 84** 034005 (2011), arXiv: 1104.2849 [hep-ph].
- [30] Z.H. Guo, J. A. Oller and J. Ruiz de Elvira, *Phys. Lett.* **B712**, 407 (2012), arXiv: 1203.4381 [hep-ph]; *Phys. Rev.* **D 86**, 054006 (2012), arXiv: 1206.4163 [hep-ph].
- [31] J. Nebreda, J. T. Londergan, J. R. Peláez and A. P. Szczepaniak, arXiv: 1403.2790 [hep-ph].
- [32] T. Cohen, F. J. Llanes-Estrada, J. R. Pelaez and J. Ruiz de Elvira, *Phys. Rev.* **D 90** 036003 (2014), arXiv: 1405.4831 [hep-ph].
- [33] Z.-H. Guo, Ulf-G. Meißner, and D.-L. Yao, *Phys. Rev.* **D 92**, 094008 (2015), arXiv: 1507.03123 [hep-ph].
- [34] Z.H. Guo, and J. A. Oller, *Phys. Rev.* **D 93**, 096001 (2016), arXiv: 1508.06400 [hep-ph].
- [35] Raul A. Briceno, Jozef J. Dudek, Robert G. Edwards, David J. Wilson, *Phys. Rev. Lett.* **118**, 022002, (2017). arXiv: 1607.05900 [hep-ph].
- [36] J.R. Pelaez, *Phys. Rev. Lett.* **92** 102001 (2004). arXiv: 0309292 [hep-ph].
- [37] J.R. Pelaez and G. Rios, *Phys. Rev. Lett.* **97** 242002 (2006). arXiv: 0610397 [hep-ph].
- [38] Z.X. Sun *et al.*, *Mod. Phys. Lett.* **A22** 711 (2007). arXiv: 0503195 [hep-ph].
- [39] J.R. Pelaez, M.R. Pennington, J. Ruiz de Elvira and D.J. Wilson, *Phys. Rev.* **D 84** 096006 (2011), arXiv: 1009.6204 [hep-ph].
- [40] L.Y. Dai, X.G. Wang and H.Q. Zheng, *Commun. Theor. Phys.* **57** 841 (2012), arXiv: 1108.1451 [hep-ph]; *Commun. Theor. Phys.* **58** 410 (2012), arXiv: 1206.5481 [hep-ph].
- [41] G. Y. Qin, W. Z. Deng, Z. Xiao and H. Q. Zheng, *Phys. Lett. B* **542** 89 (2002), hep-ph/0205214.
- [42] A. Gomez Nicola, J. R. Pelaez, *Phys. Rev.* **D 65** 054009 (2002). arXiv: 0109056 [hep-ph].
- [43] L.Y. Dai and M.R. Pennington, *Phys. Lett.* **B736** 11 (2014), arXiv: 1403.7514 [hep-ph]; *Phys. Rev.* **D 90** 036004 (2014), arXiv: 1404.7524 [hep-ph].
- [44] L. Y. Dai, V. Mathieu, E. Passemar, M. R. Pennington and A. Szczepaniak, in preparation.
- [45] G. ’t Hooft, *Nucl. Phys.* **B72** 461 (1974).
- [46] G. ’t Hooft, *Nucl. Phys.* **B75** 461 (1974).
- [47] E. Witten, *Nucl. Phys.* **B160** 57 (1979).
- [48] S.R. Coleman, *Aspects of Symmetry*, (Cambridge Univ. Press, Cambridge, England, 1985), pp. 377-378.
- [49] T.D. Cohen, *Phys. Lett.* **B 427**, 348 (1998), arXiv: [hep-ph/9801316].
- [50] S. Weinberg, *Phys. Rev. Lett.* **110** 261601 (2013), arXiv: 1303.0342 [hep-ph].
- [51] M. Knecht and S. Peris, *Phys. Rev.* **D 88**, 036016 (2013), arXiv: 1307.1273 [hep-ph].
- [52] T. Cohen, F. J. Llanes-Estrada, J. R. Pelaez and J. Ruiz de Elvira,

- Phys. Rev. D **90**, 036003 (2014), arXiv: 1405.4831 [hep-ph].
- [53] R.F. Lebed, Phys. Rev. D **88**, 057901 (2013), arXiv: 1308.2657 [hep-ph].
 - [54] T.D. Cohen and R.F. Lebed, Phys. Rev. D **89**, 054018 (2014), arXiv: 1401.1815 [hep-ph]; Phys. Rev. D **90**, 016001 (2014), arXiv: 1403.8090 [hep-ph].
 - [55] F. K. Guo, C. Hanhart, Ulf-G. Meißner, Q. Wang, Q. Zhao and B. S. Zou, Rev. Mod. Phys. **90** 015004 (2018), arXiv: <https://arxiv.org/abs/1705.00141> [hep-ph].
 - [56] R. Omnès, Nuovo Cim. **8**, 316 (1958).
 - [57] R. García- Martín, R. Kamiński, J. R. Peláez, J. Ruiz de Elvira, and F. J. Ynduráin, Phys. Rev. D **83** 074004 (2011), arXiv: 1102.2183 [hep-ph].
 - [58] M.M. Nagels *et al.*, Nucl. Phys. **B147** 189 (1979);
 - [59] L. Y. Dai, X. W. Kang, U. G. Meißner, X. Y. Song and D. L. Yao, Phys. Rev. D **97** 036012 (2018), arXiv: 1712.02119 [hep-ph].
 - [60] J. Gasser, H. Leutwyler, Ann. Phys. (NY) **158** 142 (1984).
 - [61] J. Gasser, H. Leutwyler, Nucl. Phys. **B250** 465 (1985).
 - [62] J. Bijnens, G. Colangelo and J. Gasser, Nucl. Phys. **B427** 427 (1994).
 - [63] B. Hyams, *et al.*, Nucl. Phys. **B64** 134 (1973); G. Grayer, *et al.*, Nucl. Phys. **B75** 189 (1974); B. Hyams, *et al.*, Nucl. Phys. **B100** 205 (1975).
 - [64] M. Döring, Ulf-G. Meißner, E. Oset, A. Rusetsky, Eur. Phys. J. **A47** 139 (2011), arXiv: 1107.3988 [hep-lat]; Eur. Phys. J. **A48** 114 (2012), arXiv: 1205.4838 [hep-lat].
 - [65] C. Patrignani *et al.*, [PDG], Chin. Phys. **C40** 100001 (2016).
 - [66] G. Ecker, J. Gasser, A. Pich, and E. de Rafael, Nucl. Phys. **B321** 311 (1989).
 - [67] T. Ledwig, J. Nieves, A. Pich, E.R. Arriola, and J. Ruiz de Elvira, Phys. Rev. D **90** 114020 (2014), arXiv: 1407.3750 [hep-lat].
 - [68] J. Nieves and E. Ruiz Arriola, Phys. Lett. B **679**, 449 (2009), arXiv:0904.4590 [hep-ph].
 - [69] M. Uehara, arXiv: 0404221 [hep-ph].