

# An inequality for solutions of the Navier-Stokes equations in $\mathbb{R}^n$

T. HAGSTROM,<sup>1</sup> J. LORENZ,<sup>2</sup> J. P. ZINGANO<sup>3</sup> AND P. R. ZINGANO<sup>3</sup>

<sup>1</sup>Department of Mathematics  
Southern Methodist University  
Dallas, TX 75275-0235, USA

<sup>2</sup>Department of Mathematics and Statistics  
University of New Mexico  
Albuquerque, NM 87131-0001, USA

<sup>3</sup>Departamento de Matemática Pura e Aplicada  
Universidade Federal do Rio Grande do Sul  
Porto Alegre, RS 91509-900, Brazil

## Abstract

We obtain a new inequality that holds for general Leray solutions of the incompressible Navier-Stokes equations in  $\mathbb{R}^n$  ( $n \leq 4$ ). This recovers important results previously obtained by other authors regarding the time decay of solution derivatives (of arbitrary order).

**2010 AMS Subject Classification:** 35B40 (primary), 35D30, 35Q30, 76D05

**Keywords:** time decay of solution derivatives, Leray global weak solutions, incompressible Navier-Stokes equations, Schonbek-Wiegner estimates

## 1. Introduction

In this note we derive a fundamental new inequality for general Leray solutions of the incompressible Navier-Stokes equations (in dimension  $n \leq 4$ ), that is, global solutions  $\mathbf{u}(\cdot, t) \in L^\infty((0, \infty), \mathbf{L}_\sigma^2(\mathbb{R}^n)) \cap L^2((0, \infty), \dot{\mathbf{H}}^1(\mathbb{R}^n)) \cap C_w([0, \infty), \mathbf{L}^2(\mathbb{R}^n))$  of the fluid flow system

$$\mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = \nu \Delta \mathbf{u}, \quad (1.1a)$$

$$\nabla \cdot \mathbf{u}(\cdot, t) = 0, \quad (1.1b)$$

$$\mathbf{u}(\cdot, 0) = \mathbf{u}_0 \in \mathbf{L}_\sigma^2(\mathbb{R}^n), \quad (1.1c)$$

that satisfy the generalized energy inequality

$$\| \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^n)}^2 + 2\nu \int_s^t \| D\mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^n)}^2 d\tau \leq \| \mathbf{u}(\cdot, s) \|_{L^2(\mathbb{R}^n)}^2, \quad \forall t \geq s \quad (1.2)$$

for a.e.  $s \geq 0$ , including  $s = 0$ . Such solutions were first constructed by Leray [9, 10] for  $n \leq 3$ , and later by other authors with different methods and more general space dimension, see e.g. [3, 4, 8, 16, 17, 18]. In (1.1) above,  $\nu > 0$  is a given constant,  $\mathbf{u} = \mathbf{u}(x, t)$  and  $p = p(x, t)$  are the unknowns (the flow velocity and pressure, respectively), with condition (1.1c) satisfied in  $\mathbf{L}^2(\mathbb{R}^n)$ , i.e.,  $\| \mathbf{u}(\cdot, t) - \mathbf{u}_0 \|_{L^2(\mathbb{R}^n)} \rightarrow 0$  as  $t \searrow 0$ . In the present work, we always assume  $2 \leq n \leq 4$ .

A well known property of Leray solutions is that they are eventually very regular: there is always some  $t_* \geq 0$  such that one has  $\mathbf{u} \in C^\infty(\mathbb{R}^n \times (t_*, \infty))$  and, moreover,

$$\mathbf{u}(\cdot, t) \in C((t_*, \infty), \mathbf{H}^m(\mathbb{R}^n)), \quad \forall m \geq 0, \quad (1.3)$$

see e.g. [4, 7, 9, 10, 15, 17].<sup>1</sup> It is also well established that  $\lim_{t \rightarrow 0} \| \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^n)} = 0$  and, more generally,

$$\lim_{t \rightarrow 0} t^{m/2} \| D^m \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^n)} = 0 \quad (1.4)$$

for every  $m \geq 1$ , and for all Leray solutions to the system (1.1) [1, 4, 11, 12, 13, 15]. Furthermore, suitable stronger assumptions on the initial data have led to interesting finer estimates for the solutions and their derivatives, see e.g. [6, 12, 15, 19]. An important shortcut for many of these results (including (1.4) and the Schonbek-Wiegner estimates [15]) is provided by the following fundamental inequality recently discovered by the authors, which has eluded previous studies.<sup>2</sup>

**Main Theorem.** *Let  $n \leq 4$ ,  $\mathbf{u}_0 \in \mathbf{L}_\sigma^2(\mathbb{R}^n)$ , and let  $\mathbf{u}(\cdot, t)$  be any particular Leray solution to the Navier-Stokes equations (1.1). Then we have, for every  $\alpha \geq 0$ :*

$$\limsup_{t \rightarrow \infty} t^{\alpha+m/2} \| D^m \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^n)} \leq K(\alpha, m) \nu^{-m/2} \limsup_{t \rightarrow \infty} t^\alpha \| \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^n)} \quad (1.5)$$

for every  $m \geq 1$ , where

$$K(\alpha, m) = \min_{\delta > 0} \left\{ \delta^{-1/2} \prod_{j=0}^m (\alpha + j/2 + \delta)^{1/2} \right\}.$$

<sup>1</sup>It is known that  $t_* = 0$  if  $n = 2$ ,  $t_* \leq \nu^{-5} \| \mathbf{u}_0 \|_{L^2(\mathbb{R}^3)}^4$  if  $n = 3$ ,  $t_* \leq \nu^{-3} \| \mathbf{u}_0 \|_{L^2(\mathbb{R}^4)}^2$  if  $n = 4$ .

<sup>2</sup>For the definition of  $\| \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^n)}$ ,  $\| D^m \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^n)}$  and other similar norms, see (1.6).

In Section 2 we present our original derivation of (1.5), which was based in part on some previous ideas in [2, 5, 6, 20]. Alternative proofs could also be developed (using e.g. Schonbek's Fourier splitting method [13, 14]), but we prefer to follow the very way in which (1.5) was first revealed.

**Notation.** As already shown, boldface letters are used for vector quantities, as in  $\mathbf{u}(x, t) = (u_1(x, t), \dots, u_n(x, t))$ . Also,  $\nabla p \equiv \nabla p(\cdot, t)$  denotes the spatial gradient of  $p(\cdot, t)$ ;  $D_j = \partial/\partial x_j$ ;  $\nabla \cdot \mathbf{u} = D_1 u_1 + \dots + D_n u_n$  is the (spatial) divergence of  $\mathbf{u}(\cdot, t)$ .  $\mathbf{L}_\sigma^2(\mathbb{R}^n)$  denotes the space of solenoidal fields  $\mathbf{v} = (v_1, \dots, v_n) \in \mathbf{L}^2(\mathbb{R}^n) \equiv L^2(\mathbb{R}^n)^n$  with  $\nabla \cdot \mathbf{v} = 0$  in the distributional sense;  $\dot{\mathbf{H}}^1(\mathbb{R}^n) = \dot{H}^1(\mathbb{R}^n)^n$  with  $\dot{H}^1(\mathbb{R}^n)$  being the homogeneous  $L^2$  Sobolev space of order 1;  $\mathbf{H}^m(\mathbb{R}^n) = H^m(\mathbb{R}^n)^n$ , where  $H^m(\mathbb{R}^n)$  is the space of functions  $v \in L^2(\mathbb{R}^n)$  whose  $m$ -th order derivatives are also square integrable.  $C_w(I, \mathbf{L}^2(\mathbb{R}^n))$  denotes the set of mappings from a given interval  $I \subseteq \mathbb{R}$  to  $\mathbf{L}^2(\mathbb{R}^n)$  that are  $L^2$ -weakly continuous at each  $t \in I$ .  $\|\cdot\|_{L^q(\mathbb{R}^n)}$ ,  $1 \leq q \leq \infty$ , are the standard norms of the Lebesgue spaces  $L^q(\mathbb{R}^n)$ , with the vector counterparts

$$\|\mathbf{u}(\cdot, t)\|_{L^q(\mathbb{R}^n)} = \left\{ \sum_{i=1}^n \int_{\mathbb{R}^n} |u_i(x, t)|^q dx \right\}^{1/q} \quad (1.6a)$$

$$\|D^m \mathbf{u}(\cdot, t)\|_{L^q(\mathbb{R}^n)} = \left\{ \sum_{i, j_1, \dots, j_m=1}^n \int_{\mathbb{R}^n} |D_{j_1} \dots D_{j_m} u_i(x, t)|^q dx \right\}^{1/q} \quad (1.6b)$$

if  $1 \leq q < \infty$ ; if  $q = \infty$ , then  $\|\mathbf{u}(\cdot, t)\|_{L^\infty(\mathbb{R}^n)} = \max \{ \|u_i(\cdot, t)\|_{L^\infty(\mathbb{R}^n)} : 1 \leq i \leq n \}$ ,  $\|D\mathbf{u}(\cdot, t)\|_{L^\infty(\mathbb{R}^n)} = \max \{ \|D_j u_i(\cdot, t)\|_{L^\infty(\mathbb{R}^n)} : 1 \leq i, j \leq n \}$ , and so forth.

## 2. Proof of (1.5)

The derivation of (1.5) below takes advantage of the regularity property (1.3) and proceeds by induction in  $m$ . It combines standard techniques (energy inequalities and related interpolation estimates) with well known properties of Leray solutions (namely, that  $\|\mathbf{u}(\cdot, t)\|_{L^2(\mathbb{R}^n)} \rightarrow 0$  (as  $t \rightarrow \infty$ ), or that

$$\lim_{t \rightarrow \infty} \|D\mathbf{u}(\cdot, t)\|_{L^2(\mathbb{R}^n)} = 0, \quad (2.1)$$

which are easy to obtain directly). As the proofs for  $n = 2, 3, 4$  are entirely similar, we will present the details for one case only — say,  $n = 4$ . Let then  $\mathbf{u}(\cdot, t)$  be any given Leray solution to (1.1), in  $\mathbb{R}^4$ , such that we have, for some  $\alpha \geq 0$ ,

$$\limsup_{t \rightarrow \infty} t^\alpha \|\mathbf{u}(\cdot, t)\|_{L^2(\mathbb{R}^4)} =: \lambda_0(\alpha) < \infty. \quad (2.2)$$

Let  $\delta > 0$ ,  $0 < \epsilon < 2$  be given, and let  $t_*$  be the solution's regularity time as defined in (1.3). Recalling the basic estimate

$$\|u\|_{L^4(\mathbb{R}^4)} \leq \|Du\|_{L^2(\mathbb{R}^4)}, \quad (2.3)$$

from which we get

$$\|D^\ell u\|_{L^4(\mathbb{R}^4)} \|D^{m-\ell} u\|_{L^4(\mathbb{R}^4)} \leq \|Du\|_{L^2(\mathbb{R}^4)} \|D^{m+1} u\|_{L^2(\mathbb{R}^4)} \quad (2.4)$$

for arbitrary  $m \geq 0$ ,  $0 \leq \ell \leq m$ , we may proceed along the lines of [2, 20] as follows. Taking the dot product of (1.1a) with  $(t - t_0)^{2\alpha+\delta} \mathbf{u}(x, t)$  and integrating the result on  $\mathbb{R}^4 \times [t_0, t]$ , for  $t \geq t_0 > t_*$ , we obtain, because of (1.1b),

$$\begin{aligned} (t - t_0)^{2\alpha+\delta} \|\mathbf{u}(\cdot, t)\|_{L^2(\mathbb{R}^4)}^2 + 2\nu \int_{t_0}^t (\tau - t_0)^{2\alpha+\delta} \|D\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)}^2 d\tau \\ = (2\alpha + \delta) \int_{t_0}^t (\tau - t_0)^{2\alpha+\delta-1} \|\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)}^2 d\tau \end{aligned}$$

for  $t \geq t_0 \geq t_*$ . This promptly gives, by (2.2), that

$$\int_{t_0}^t (\tau - t_0)^{2\alpha+\delta} \|D\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)}^2 d\tau \leq \frac{1}{2\nu} \frac{2\alpha + \delta}{\delta} (\lambda_0(\alpha) + \epsilon)^2 (t - t_0)^\delta \quad (2.5)$$

for all  $t \geq t_0$  (choosing  $t_0 \geq t_*$  sufficiently large). Next, for  $m = 1$ , we similarly have

$$\begin{aligned} (t - t_0)^{2\alpha+1+\delta} \|D\mathbf{u}(\cdot, t)\|_{L^2(\mathbb{R}^4)}^2 + 2\nu \int_{t_0}^t (\tau - t_0)^{2\alpha+1+\delta} \|D^2\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)}^2 d\tau \\ \leq (2\alpha + 1 + \delta) \int_{t_0}^t (\tau - t_0)^{2\alpha+\delta} \|D\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)}^2 d\tau + \\ K_1 \int_{t_0}^t (\tau - t_0)^{2\alpha+1+\delta} \|\mathbf{u}(\cdot, \tau)\|_{L^4(\mathbb{R}^4)} \|D\mathbf{u}(\cdot, \tau)\|_{L^4(\mathbb{R}^4)} \|D^2\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)} d\tau \end{aligned}$$

(where  $K_1 = 8\sqrt{2}$ ), which gives, by (2.3):

$$\begin{aligned} (t - t_0)^{2\alpha+1+\delta} \|D\mathbf{u}(\cdot, t)\|_{L^2(\mathbb{R}^4)}^2 + 2\nu \int_{t_0}^t (\tau - t_0)^{2\alpha+1+\delta} \|D^2\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)}^2 d\tau \\ \leq (2\alpha + 1 + \delta) \int_{t_0}^t (\tau - t_0)^{2\alpha+\delta} \|D\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)}^2 d\tau + \\ K_1 \int_{t_0}^t (\tau - t_0)^{2\alpha+1+\delta} \|D\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)} \|D^2\mathbf{u}(\cdot, \tau)\|_{L^2(\mathbb{R}^4)}^2 d\tau \end{aligned}$$

for  $t \geq t_0$ . By (2.1) and (2.5), we then get (increasing  $t_0$  if necessary):

$$(t - t_0)^{2\alpha+1} \| D \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^4)}^2 \leq \frac{1}{2\nu} (2\alpha + 1 + \delta) \frac{2\alpha + \delta}{\delta} (\lambda_0(\alpha) + \epsilon)^2 \quad (2.6a)$$

and

$$\int_{t_0}^t (\tau - t_0)^{2\alpha+1+\delta} \| D^2 \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)}^2 d\tau \leq \frac{(2\alpha + 1 + \delta)(2\alpha + \delta)}{\delta \cdot [(2 - \epsilon)\nu]^2} (\lambda_0(\alpha) + \epsilon)^2 (t - t_0)^\delta \quad (2.6b)$$

for all  $t \geq t_0$ . Proceeding in this way ( $m = 2, 3, \dots$ ) we obtain at the  $m$ th step

$$\begin{aligned} & (t - t_0)^{2\alpha+m+\delta} \| D^m \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^4)}^2 + 2\nu \int_{t_0}^t (\tau - t_0)^{2\alpha+m+\delta} \| D^{m+1} \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)}^2 d\tau \\ & \leq (2\alpha + m + \delta) \int_{t_0}^t (\tau - t_0)^{2\alpha+m-1+\delta} \| D^m \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)}^2 d\tau + \\ & K_m \int_{t_0}^t (\tau - t_0)^{2\alpha+m+\delta} \| D^{m+1} \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)} \sum_{\ell=0}^{[m/2]} \| D^\ell \mathbf{u}(\cdot, \tau) \|_{L^4(\mathbb{R}^4)} \| D^{m-\ell} \mathbf{u}(\cdot, \tau) \|_{L^4} d\tau \end{aligned}$$

for  $t \geq t_0$ , and some constant  $K_m > 0$ , where  $[m/2]$  denotes the integer part of  $m/2$ . This gives, by (2.4):

$$\begin{aligned} & (t - t_0)^{2\alpha+m+\delta} \| D^m \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^4)}^2 + 2\nu \int_{t_0}^t (\tau - t_0)^{2\alpha+m+\delta} \| D^{m+1} \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)}^2 d\tau \\ & \leq (2\alpha + m + \delta) \int_{t_0}^t (\tau - t_0)^{2\alpha+m-1+\delta} \| D^m \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)}^2 d\tau + \\ & \left(1 + \left\lceil \frac{m}{2} \right\rceil\right) \cdot K_m \int_{t_0}^t (\tau - t_0)^{2\alpha+m+\delta} \| D \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)} \| D^{m+1} \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)}^2 d\tau. \end{aligned} \quad (2.7)$$

At this stage, we would already know from the previous steps that

$$(t - t_0)^{2\alpha+k} \| D^k \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^4)}^2 \leq \frac{1}{\delta \cdot [(2 - \epsilon)\nu]^k} \left\{ \prod_{j=0}^k (2\alpha + j + \delta) \right\} (\lambda_0(\alpha) + \epsilon)^2 \quad (2.8a)$$

and

$$\begin{aligned} \int_{t_0}^t (\tau - t_0)^{2\alpha+k+\delta} \| D^{k+1} \mathbf{u}(\cdot, \tau) \|_{L^2(\mathbb{R}^4)}^2 d\tau & \leq \frac{\delta^{-1}}{[(2 - \epsilon)\nu]^{k+1}} \left\{ \prod_{j=0}^k (2\alpha + j + \delta) \right\} \\ & \times (\lambda_0(\alpha) + \epsilon)^2 \cdot (t - t_0)^\delta \end{aligned} \quad (2.8b)$$

for all  $t \geq t_0$ , and each  $0 \leq k < m$ . By (2.1) and (2.7), and increasing  $t_0$  if necessary, we would then obtain (2.8) for  $k = m$  as well, completing the induction step.

The argument above established that, for each  $m \geq 1$ , we have

$$(t - t_0)^{2\alpha+m} \| D^m \mathbf{u}(\cdot, t) \|_{L^2(\mathbb{R}^4)}^2 \leq \frac{1}{\delta \cdot [(2 - \epsilon)\nu]^m} \left\{ \prod_{j=0}^m (2\alpha + j + \delta) \right\} (\lambda_0(\alpha) + \epsilon)^2$$

for all  $t$  sufficiently large. Since  $\delta > 0$ ,  $0 < \epsilon < 2$  are arbitrary, this gives the result.

### Acknowledgements.

Work of the first author was supported by NSF Grant DMS-1418871, and that of the last author by CAPES Grant # 88881.067966/2014-01. Any opinions, findings, and conclusions or recommendations are those of the authors and do not necessarily reflect the views of the National Science Foundation or CAPES.

## References

- [1] J. BENAMEUR AND R. SELMI, *Long time decay to the Leray solution of the two-dimensional Navier-Stokes equations*, Bull. London Math. Soc. **44** (2012), 1001-1019.
- [2] P. BRAZ E SILVA, L. SCHÜTZ AND P. R. ZINGANO, *On some energy inequalities and supnorm estimates for advection-diffusion equations in  $\mathbb{R}^n$* , Nonl. Anal. **93** (2013), 90-96.
- [3] H. FUJITA AND T. KATO, *On the Navier-Stokes initial value problem*, Arch. Rat. Mech. Anal. **16** (1964), 269-315.
- [4] T. KATO, *Strong  $L^p$ -solutions of the Navier-Stokes equations in  $\mathbb{R}^m$ , with applications to weak solutions*, Math. Z. **187** (1984), 471-480.
- [5] H.-O. KREISS, T. HAGSTROM, J. LORENZ AND P. R. ZINGANO, *Decay in time of the solutions of the Navier-Stokes equations for incompressible flows*, unpublished note, University of New Mexico, Albuquerque, NM, 2002.
- [6] H.-O. KREISS, T. HAGSTROM, J. LORENZ AND P. R. ZINGANO, *Decay in time of incompressible flows*, J. Math. Fluid Mech. **5** (2003), 231-244.
- [7] H.-O. KREISS AND J. LORENZ, *Initial-boundary value problems and the Navier-Stokes equations*, Academic Press, New York, 1989. (Reprinted in the series SIAM Classics in Applied Mathematics, Vol. 47, 2004.)
- [8] O. A. LADYZHENSKAYA, *The Mathematical Theory of Viscous Incompressible Flow* (2nd ed.), Gordon and Breach, New York, 1969.

- [9] J. LERAY, *Étude de diverses équations intégrales non linéaires et de quelques problèmes que pose l'hydrodynamique*, J. Math. Pures Appl. **12** (1933), 1-82.
- [10] J. LERAY, *Essai sur le mouvement d'un fluide visqueux emplissant l'espace*, Acta Math. **63** (1934), 193-248.
- [11] K. MASUDA, *Weak solutions of the Navier-Stokes equations*, Tôhoku Math. Journal **36** (1984), 623-646.
- [12] M. OLIVER AND E. S. TITI, *Remark on the rate of decay of higher order derivatives for solutions to the Navier-Stokes equations in  $\mathbb{R}^n$* , J. Funct. Anal. **172** (2000), 1-18.
- [13] M. E. SCHONBEK,  *$L^2$  decay for weak solutions of the Navier-Stokes equations*, Arch. Rat. Mech. Anal. **88** (1985), 209-222.
- [14] M. E. SCHONBEK, *The Fourier splitting method*, in: P. Concus and K. Lancaster (Eds.), *Advances in Geometric Analysis and Continuum Mechanics*, International Press, Cambridge, 1995, pp. 269-274.
- [15] M. E. SCHONBEK AND M. WIEGNER, *On the decay of higher-order norms of the solutions of Navier-Stokes equations*, Proc. Roy. Soc. Edinburgh **126A** (1996), 677-685.
- [16] M. SHINBROT AND S. KANIEL, *The initial value problem for the Navier-Stokes equations*, Arch. Rat. Mech. Anal. **21** (1966), 270-285.
- [17] H. SOHR, *The Navier-Stokes Equations*, Birkhäuser, Basel, 2001.
- [18] R. TEMAM, *Navier-Stokes Equations: theory and numerical analysis* (2nd ed.), AMS/Chelsea, Providence, 1984.
- [19] M. WIEGNER, *Decay results for weak solutions of the Navier-Stokes equations on  $\mathbb{R}^n$* , J. London Math. Soc. **35** (1987), 303-313.
- [20] P. R. ZINGANO, *Nonlinear  $L^2$  stability under large disturbances*, J. Comp. Appl. Math. **103** (1999), 207-219.