

# Exact Monte Carlo likelihood-based inference for jump-diffusion processes

F. B. Gonçalves<sup>a,1</sup>, K. Łatuszyński<sup>b,c</sup>, G. O. Roberts<sup>b</sup>

<sup>a</sup> Universidade Federal de Minas Gerais, Brazil

<sup>b</sup> University of Warwick, UK

<sup>c</sup> Alan Turing Institute

## Abstract

Statistical inference for discretely observed jump-diffusion processes is a complex problem which motivates new methodological challenges. Thus existing approaches invariably resort to time-discretisations which inevitably lead to approximations in inference. In this paper, we give the first general collection of methodologies for exact (in this context meaning discretisation-free) likelihood-based inference for discretely observed jump-diffusions. We shall introduce both frequentist and Bayesian approaches, illustrating the methodology through simulated and real examples.

*Key Words:* Retrospective sampling, MCEM, MCMC, Poisson estimator, Barker's algorithm.

## 1 Introduction

Jump-diffusion processes are used in a variety of applications in several scientific areas, especially economics (see Ball and Roma, 1993; Duffie et al., 2000; Runggaldier, 2003; Eraker et al., 2003; Eraker, 2004; Johannes, 2004; Barndorff-Nielsen and Shephard, 2004; Feng and Linetsky, 2008; Kennedy et al., 2009; Cont and Tankov, 2004). Other applications can be found, for example, in physics (see Chudley and Elliott, 1961), biomedicine (see Grenander and Miller, 1994) and object recognition (see Srivastava et al., 2002). Jump-diffusions are natural extensions to diffusions, allowing for discrete discontinuities in trajectories which are otherwise described by diffusion dynamics, thus offering additional flexibility in modelling phenomena which exhibit sudden large jumps.

Inference for jump-diffusions is a challenging problem because transition densities are almost always intractable. This problem is typically overcome using approximations based on time-discretisations which typically lead to systematic biases which are difficult to quantify (see Detemple et al., 2006). Therefore, the problem requires new approaches which circumvent the need for approximation, and we shall call them *exact* solutions. Whereas efficient exact solutions have already been proposed for the context where there is no jump component (see Beskos et al., 2006b, 2009; Sermaidis et al., 2013), the problem for jump-diffusions is extremely under-developed.

This paper provides a general suite of methodologies for exact likelihood-based inference for discretely observed jump-diffusions, featuring both maximum likelihood and Bayesian approaches. The only source of inaccuracy is Monte Carlo error which can be controlled using standard Monte Carlo techniques. Some of our approaches directly use an algorithm that performs exact simulation of a class of jump-diffusion bridges proposed in Gonçalves and Roberts (2014) and called the *Jump Exact algorithm* (JBEA). However our most general approaches involve the construction of significant extensions of JBEA.

---

<sup>1</sup>Address: Av. Antônio Carlos, 6627 - DEST, ICEX, UFMG - 31270-901, Belo Horizonte, Minas Gerais, Brazil.  
E-mail: fbgoncalves@est.ufmg.br

First we propose two methodologies (MCEM and MCMC) for the case where the drift and the jump-rate are uniformly bounded and JBEA can be directly applied. Our methodology collapses to that of Beskos et al. (2006b) in the case where no jump component is present. We also propose two methods for the general case where both the drift and the jump-rate may be unbounded. One of these is an importance sampling (IS) adaptation of the first MCEM method we propose providing improved Monte Carlo variance properties. The second method introduces a new perspective for solving the inference problem in an exact framework using an infinite-dimensional Markov chain Monte Carlo (MCMC) algorithm based on Barker’s transitions (Barker, 1965) and novel simulation techniques.

Our exact approach can be applied to a wide class of univariate models which allow non-linear state-dependent drift and diffusion coefficients, and state-dependent and time-inhomogeneous jump rate and jump size distribution. The methodology extends in principle to certain multivariate jump-diffusions, although we do not explore this direction here.

Whilst the focus on this paper is clearly on exact inference for jump-diffusions, the use of exact Barker’s transitions via a *two coin* algorithm which we will introduce appears to be of generic interest for any context with intractable accept/reject ratios. Barker’s method is rarely used, as it is known to be uniformly dominated by the much more well-known Metropolis-Hastings accept/reject formula. However, Metropolis-Hastings never beats Barker by a factor greater than 2 in Peskun order sense (Łatuszyński and Roberts, 2013), and crucially, in contrast to the Metropolis-Hastings, the particular form of the accept/reject formula in the Barker case permits the construction of the two coin procedure.

This paper is organised as follows. The remainder of Section 1 provides a literature review and formally defines the class of jump-diffusion processes to be considered. Section 2 revisits the algorithm for the exact simulation of jump-diffusion bridges, JBEA, proposed in Gonçalves and Roberts (2014). Section 3 presents two inference methods directly based on JBEA and applicable when the drift and jump rates are uniformly bounded. Section 4 presents two other methods which make indirect use of JBEA and do not require the boundedness assumption. In Section 5 the methods are applied to simulated data sets to investigate their efficiency. Finally, two real data sets concerning the exchange rate GBP/USD and the S&P500 index are analysed in Section 6.

## 1.1 Literature review on approximate methods

Existing solutions for the inference problem that we consider are based on approximations which, in turn, rely on *path discretisation* and/or *data augmentation* strategies. These solutions follow basically three main directions: considering alternative estimators to the *maximum likelihood estimator* (MLE); using numerical approximations to the unknown likelihood function; and estimating an approximation to the likelihood by using Monte Carlo methods. Moreover, several of the methods assume state-independence of various components of the model.

Alternative estimators can be found in Duffie and Singleton (1993) and Duffie and Glynn (2004). Numerical approximations can be found in Lo (1988), Aït-Sahalia and Yu (2006) and Filipović et al. (2013). Particle filter-based Monte Carlo methods can be found in Johannes et al. (2002) and Johannes et al. (2009). Golightly (2009) proposes a refinement where the particles are propagated via MCMC. The author proposes a Metropolis-Hastings algorithm generalising the Durham and Gallant bridge (Durham and Gallant (2002)) for the jump-diffusion case.

Finally, an exact methodology that performs unbiased Monte Carlo estimation of the transition density is proposed in Giesecke and Schwenkler (2019) building on the work of Pedersen (1995), Beskos et al. (2009), Chen (2009) and Casella and Roberts (2010). This work provides methodology for exact MLE methods although suffers from two main drawbacks. Firstly, the results that assure finite variance of the transition density estimator and consistency and asymptotic normality of the simulated MLE require strong assumptions that are typically hard to check and often not met in realistic situations. Secondly, even when these assumptions are met, the method can be very inefficient due to high importance sampling weight variances. Our methodology will circumvent these difficulties

through the use of rejection sampling, importance sampling with provably finite variances, and a novel Barker’s MCMC methodology for Bayesian inference.

## 1.2 The jump-diffusion model

Formally, a jump-diffusion is the stochastic process  $V := \{V_s : 0 \leq s \leq t\}$  that solves the SDE:

$$dV_s = b(V_{s-}, s)ds + \sigma(V_{s-}, s)dW_s + \int_E g_1(z, V_{s-})m(dz, ds), \quad V_0 = v_0, \quad (1)$$

where  $b, \sigma : \mathbb{R} \rightarrow \mathbb{R}$  and  $g_1 : E \times \mathbb{R} \rightarrow \mathbb{R}$  are assumed to satisfy the regularity conditions (locally Lipschitz, with a linear growth bound) to guarantee a unique weak solution (see Platen and Bruti-Liberati, 2010, Section 1.9).  $W_s$  is a Brownian motion and  $m(dz, ds)$  is a random counting measure on the product space  $E \times [0, t]$ , for  $E \subseteq \mathbb{R}$ , with associated intensity measure  $\lambda_m$ . We assume that  $\lambda_m$  is absolutely continuous with respect to Lebesgue measure on  $E \times [0, t]$  and Markov dependent on  $V$ :

$$\lambda_m(dz, ds; V_{s-}) = \lambda_m(z, s; V_{s-})dzds = \lambda_1(s, V_{s-})f_Z(z; s)dzds, \quad (2)$$

where, for any  $v \in \mathbb{R}$ ,  $\lambda_1(\cdot, v)$  is a non-negative real valued function on  $[0, t]$  and for any  $s \in \mathbb{R}^+$ ,  $f_Z(\cdot; s)$  is a standard density function with support  $E$ . According to (1) and (2), between any two jumps, the process  $V$  behaves as a homogeneous diffusion process with drift  $b$  and diffusion coefficient  $\sigma$ . The jump times follow a Markov point process on  $[0, t]$  with intensity function (jump rate)  $\lambda_1(s; V_{s-})$ . A random variable  $Z_j$  with density  $f_Z(z; t_j)$  is associated to each of the  $N$  jump times and, along with the state of the process, determines the size of the jump  $g_1(Z_j, V_{t_j-})$  at time  $t_j$ .

## 2 Exact simulation of jump-diffusion bridges

In this section, we briefly present the JBEA algorithm to perform exact simulation of jump-diffusion bridges proposed in Gonçalves and Roberts (2014). JBEA simulates a finite representation from the exact probability law of a class of univariate jump-diffusion bridges.

JBEA can only be directly applied to processes with unit diffusion coefficient. This is a genuine restriction in the multivariate case where not all diffusions can be reduced to this case. On the other hand, in the one-dimensional case, we can always apply the Lamperti transform to obtain a process as required (assuming the diffusion coefficient is continuously differentiable). Thus we set  $X_s = \eta(V_s)$  with the Lamperti transform:

$$\eta(v) := \int_{v^*}^v \frac{1}{\sigma(u)}du, \quad (3)$$

where  $v^*$  is some arbitrary element of the state space of  $V$ .

The transformed process  $X := \{X_s : 0 \leq s \leq t\}$  is a one-dimensional jump-diffusion solving the SDE:

$$dX_s = \alpha(X_{s-})ds + dW_s + \int_E g(z, X_{s-})m(dz, ds), \quad X_0 = \eta(v_0) = x, \quad (4)$$

where

$$\alpha(u) = \frac{b\{\eta^{-1}(u)\}}{\sigma\{\eta^{-1}(u)\}} - \frac{\sigma'\{\eta^{-1}(u)\}}{2}, \quad g(z, u) = \eta(\eta^{-1}(u) + g_1(z, \eta^{-1}(u))) - u \text{ and}$$

$$\lambda_m(dz, ds; X_{s-}) = \lambda_m(z, s; X_{s-})dzds = \lambda_1(s, \eta^{-1}(X_{s-}))f_Z(z; s)dzds = \lambda(s, X_{s-})f_Z(z; s)dzds.$$

For the time interval  $[0, t]$ , we define  $N$  as the number of jumps,  $t_j$  as the jump times and  $J_j$  as the jump size at  $t_j$ ,  $1 \leq j \leq N$ . We also set  $J = \sum_{j=1}^N J_j$  and use  $\mathbf{J}$  as shorthand notation for the collection of jumps  $\{J_1, \dots, J_N\}$  and their corresponding jump times  $\{t_1, \dots, t_N\}$  and, where appropriate, also for the processes consisting of just the jumps. Now let  $f_g(\cdot; t_j, X_{t_j})$  be the pdf or pmf of the jump size at  $t_j$  induced by  $g$ . We shall refer to this as the jump size density. We shall often abbreviate notation where no ambiguity arises, e.g.  $\lambda(s)$ ,  $\lambda(j)$  and  $f_g(\cdot; j)$ .

Let  $\mathbb{P}$  be the probability measure induced by the solution  $X$  of (4) and define  $\tilde{\mathbb{P}}$  to be the law of the jump-diffusion with law  $\mathbb{P}$  given  $X_t = y$ . Now let  $\tilde{\mathbb{D}}$  be the law of the process constructed as the sum of an independent jump process, with jump rate  $\lambda_0$  and jump size density  $f_0$ , and a Brownian bridge  $B^*$  that starts in  $x$  at time 0 and finishes in  $y - J$  at time  $t$ . Note that  $J$  is independent of the ending point  $y$  and it is  $B^*$  that guarantees that the trajectories will end in  $y$ . To simulate a process with law  $\tilde{\mathbb{D}}$  we first simulate the jump process  $\mathbf{J}$  and then the Brownian bridge  $B^*$ , which depends on  $J$ .

We now define a probability measure  $\tilde{\mathbb{F}}$ , called the *Lipschitz proposal*, which differs from  $\tilde{\mathbb{D}}$  only in the distribution of  $(N, \mathbf{J})$  - the conditional distributions of  $\tilde{\mathbb{F}}$  and  $\tilde{\mathbb{D}}$  given  $(N, \mathbf{J})$  are identical. Define  $f_{\tilde{\mathbb{D}}}, f_{\tilde{\mathbb{F}}}$  as the joint density of  $(N, J_{t_1}, \dots, J_{t_N})$  under  $\tilde{\mathbb{D}}$  and  $\tilde{\mathbb{F}}$  respectively. We thus set

$$f_{\tilde{\mathbb{F}}} \propto f_{\tilde{\mathbb{D}}} \exp(c_0 \sum_{j=1}^N |J_j|). \quad (5)$$

Details on how to simulate from  $f_{\tilde{\mathbb{F}}}$  can be found in Gonçalves and Roberts (2014). Our aim is to simulate exact skeletons of the jump-diffusion with probability law  $\tilde{\mathbb{P}}$  using rejection sampling from  $\tilde{\mathbb{F}}$ . The complete list of assumptions is presented in Appendix A.

The algorithm proceeds by proposing  $B \sim \tilde{\mathbb{F}}$  and accepting with probability proportional to the Radon-Nikodym derivative  $\frac{d\tilde{\mathbb{P}}}{d\tilde{\mathbb{F}}}(B)$  and is given by  $a(B) := p_1(B)p_2(B)p_3(B)$ , where

$$\begin{aligned} p_1(B) &= \exp \left\{ -\frac{1}{2t}(y - J - x)^2 \right\}, \quad p_2(B) = \prod_{j=1}^N \left[ \frac{\lambda(j)f_g(J_j; j)}{\lambda_0 f_0(J_j)} \exp \{ -\Delta A(j) - c_0 |J_j| \} \right], \\ p_3(B) &= \exp \left\{ -\int_0^t \phi(s, B_s) - m ds \right\}; \\ \phi(s, B_s) &= \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, B_{s-}), \quad m = \inf_{u \in \mathcal{X}, 0 \leq s \leq t} \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, u), \end{aligned}$$

with  $\mathcal{X}$  being the state space of  $X$ ,  $A(u) := \int_0^u \alpha(y) dy$  and  $\Delta A(j) := A(X_{t_j}) - A(X_{t_{j-}})$ .

Suppose that  $r$  is an upper bound on  $\phi(s, B_s) - m$ ,  $0 \leq s \leq t, B_s \in \mathcal{X}$ . In order to evaluate the acceptance probability of the algorithm we need to simulate three independent events of probabilities  $p_1, p_2$  and  $p_3$ , respectively. The first two are achieved by simulating independent Bernoulli r.v.'s (since  $p_1$  and  $p_2$  are known). Finally note that  $p_3$  is equal to the probability that a Poisson process (PP) with rate  $r$  on  $[0, t] \times [0, 1]$  produces no points below the curve  $(\phi(s, B_s) - m)/r$ . This event can be evaluated by unveiling the proposal only at the time instances of the points from the PP, which makes it feasible to decide whether or not to accept the proposal. We call this the *Poisson Coin* algorithm.

JBEA returns a skeleton of the jump-diffusion bridge with law  $\tilde{\mathbb{P}}$ . Any further points may be simulated from the Brownian bridge  $B$ . Further details on the algorithm can be found in Gonçalves and Roberts (2014), including extensions, simplifications, details on the simulation steps and practical strategies to optimise its computational cost.

### 3 Inference via JBEA

We now present two algorithms based on JBEA to perform exact inference for discretely observed jump-diffusion processes. The first one is a Monte Carlo EM algorithm to find the MLE and the second one is a MCMC algorithm to sample from the posterior distribution of the parameters.

### 3.1 A Monte Carlo EM algorithm

In this algorithm, JBEA is used to obtain a Monte Carlo estimate of the expectation on the E-step. The exactness of the method is obtained by combining the exactness feature of JBEA with some auxiliary variable techniques. We present two versions of the algorithm to cover the cases where the diffusion coefficient does or does not depend on unknown parameters.

#### 3.1.1 The case where the diffusion coefficient is known

Consider  $V := \{V_s : 0 \leq s \leq T\}$  a one-dimensional jump-diffusion process, with probability measure  $\mathbb{S}$ , solving the SDE in (1). Suppose functions  $b$ ,  $g_1$  and  $\lambda_1$  depend on unknown parameter set  $\theta$ . Suppose we observe  $V$  at  $(n + 1)$  time instances  $0 = t_0, \dots, t_n = T$  and want to estimate the parameter vector  $\theta$  based on these observations. Now let  $X$  be the stochastic process with probability measure  $\mathbb{P}$  that solves the SDE in (4), obtained by applying the Lamperti transform  $\eta$  to  $V$ . Naturally, functions  $\alpha$ ,  $\lambda$  and  $f_g$  depend on  $\theta$ . Note, however, that, since the diffusion coefficient of  $V$  does not depend on  $\theta$ , nor does  $\eta$ , which means that  $X$  is observed.

To obtain the likelihood we need the finite-dimensional distributions of  $X$ , which are typically unavailable. We can, however, obtain the augmented or full likelihood function, ie that is obtained from observing the entire jump-diffusion trajectory in  $[0, T]$ . The augmented likelihood is obtained by writing down the Radon-Nikodym derivative of  $\mathbb{P}$  w.r.t. any measure  $\mathbb{Q}$ , such that  $\mathbb{P} \ll \mathbb{Q}$  where  $\mathbb{Q}$  does not depend on  $\theta$  (see Gonçalves and Franklin, 2019). This is the natural environment to apply the *EM algorithm*. However, since  $X_{mis}$  is an infinite-dimensional random variable, we have to be particularly careful in constructing a convergent algorithm.

In our context, we start by separating  $X$  in two parts:  $x_{obs} = \{x_0, \dots, x_n\}$ , the transformed observations, and  $X_{mis}$ , the unobserved part of the process  $X$ . In order to write down the likelihood of a complete path we choose  $\mathbb{Q}$  to be the measure of a jump-diffusion which is the sum of a Brownian motion and a jump process with jump rate 1 and jump size density  $f$  which does not depend on  $\theta$  and such that  $f_g \ll f$ . Since  $\mathbb{Q}$  does not depend on  $\theta$  and  $\mathbb{P} \ll \mathbb{Q}$ , the Radon-Nikodym derivative  $\frac{d\mathbb{P}}{d\mathbb{Q}}$  is a valid likelihood function for  $\theta$ , given a complete path of  $X$  in  $[0, T]$ . Defining  $y = X_T$ , Theorem 2 from Gonçalves and Roberts (2014) implies that the complete log-likelihood function is given by

$$\begin{aligned} l(X|\theta) &= A(y;\theta) - A(x;\theta) - \sum_{j=1}^N (\Delta A(j;\theta)) - \int_0^T \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, X_{s-}; \theta) ds \\ &\quad + \sum_{j=1}^N \log(\lambda(j;\theta)) + \log(f_g(J_j; j, \theta)) + \kappa, \end{aligned} \quad (6)$$

where  $\kappa$  is a constant with respect to  $\theta$  and so can be neglected by the EM algorithm.

Since the expectation of (6) cannot be evaluated analytically, we rely on Monte Carlo methods to obtain an unbiased and strongly consistent estimator of it.

Given the auxiliary variable  $U = (U_1, \dots, U_n)$ , where the  $U_i$ 's are mutually independent with  $U_i \sim Unif(t_{i-1}, t_i)$  and independent of  $X$ , we have that:

$$\begin{aligned} \mathbb{E}_{X_{mis}, U|x_{obs}, \theta'} (l(X|\theta)) &= A(y;\theta) - A(x;\theta) - \mathbb{E} \left[ \sum_{j=1}^N (\Delta A(j;\theta); \theta) \right] + \mathbb{E}(\kappa) \\ &\quad - \mathbb{E} \left[ \sum_{i=1}^n \Delta t_i \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (U_i, X_{U_i-}; \theta) \right] + \mathbb{E} \left[ \sum_{j=1}^N \log(\lambda(j;\theta)) + \log(f_g(J_j; j, \theta)) \right]. \end{aligned} \quad (7)$$

The MCEM algorithm proceeds by maximising the MC estimate of (7) w.r.t.  $\theta$ , based on an iid sample from  $(X_{mis}, U)|x_{obs}, \theta'$ . Samples are obtained by performing JBEA between all the consecutive pair of observations.

Thus a single iteration of the algorithm inputs  $\theta'$  and outputs the maximising  $\theta$  value. The algorithm iterates this procedure until the output values are convergent according to the desired accuracy. It is well documented (see for example Fort and Moulines, 2003, and references therein) that the number of Monte Carlo samples should increase with the EM iterations in order to overcome Monte Carlo error. Finally, the maximisation step may require numerical methods. For well behaved likelihoods, standard numerical optimisation algorithms should work well. For example, the quasi-Newton method BFGS (see, for example, Fletcher, 1987, Section 3.2) is used in the example presented in Section 5.1.

### 3.1.2 The case where the diffusion coefficient is unknown

We now focus on the case where the diffusion coefficient of  $V$  depends on unknown parameters. A very important result in jump-diffusion theory, which is particularly important for inference is that a complete path of a jump-diffusion can be used to perfectly estimate  $\sigma(V_{t-}; \theta)$ . The result states that (Protter, 2004, II.6):

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \left( V_{t_i/n} - V_{t_{(i-1)/n}} \right)^2 = \int_0^T \sigma^2(V_{s-}; \theta) ds + \sum_{j=1}^N (\Delta V_{t_j})^2. \quad (8)$$

The computational implication of this result is that we cannot construct an EM algorithm as in the previous section because there is a perfect correlation between  $\sigma$  and the missing path as described in (8) (see Meng, 1993). This problem is also relevant in MCMC algorithms, where it was first encountered in the context of inference for diffusions (see Roberts and Stramer, 2001; Elerian, 1999).

We propose a solution for this problem which combines together and then substantially generalises Roberts and Stramer (2001) and Beskos et al. (2006b). The method consists of a suitable transformation of the missing data that breaks the dependence between the missing data and the parameters, when conditional on the observed values. In our infinite-dimensional context, this problem is equivalent to finding a reparameterisation of the missing data so that the dominating measure is independent of the parameters. We construct this reparameterisation in two easily interpreted transformations.

The first transformation takes  $X = \eta(V; \theta)$ , with  $\{x(\theta) = x_0(\theta), x_1(\theta), \dots, x_n(\theta) = y(\theta)\}$  being the transformed observations, which now depend on  $\theta$ . Let  $N_i$  be the number of jumps in  $(t_{i-1}, t_i)$  and  $(t_{i_1}, \dots, t_{i_{N_i}})$  be the jump times. Defining  $t_{i_0} = t_{i-1}$  and  $t_{i_{N_i+1}} = t_i$ , we obtain the second level of path transformation  $\{X_s \rightarrow \dot{X}_s; s \in (t_{i-1}, t_i) \setminus \{t_{i_1}, \dots, t_{i_{N_i}}\}\}$ :

$$\dot{X}_s := X_s - \left( 1 - \frac{s - t_{i_{j-1}}}{t_{i_j} - t_{i_{j-1}}} \right) X_{i_{j-1}} - \left( \frac{s - t_{i_{j-1}}}{t_{i_j} - t_{i_{j-1}}} \right) X_{i_j-}; \quad s \in (t_{i_{j-1}}, t_{i_j}), \quad (9)$$

where  $X_{i_j} = X_{t_{i_j}}$ ,  $X_{i_0} = x_{t_{i-1}}(\theta)$  and  $X_{i_{N_i+1}-} = x_{t_i}(\theta)$ .

Note that  $\dot{X}$  is a collection of diffusion bridges starting and ending in 0. Its dynamics depend on  $\theta$  and are typically intractable; nevertheless it is easy to simulate  $\dot{X}$  at any time  $s$ , conditionally on  $V_{obs}$  and a specific value of  $\theta$ , by firstly computing  $x_{t_{i-1}}(\theta)$  and  $x_{t_i}(\theta)$ , then simulating  $X$  via JBEA and, finally, applying the transformation in (9). The inverse transformation of  $\dot{X}_s \rightarrow X_s$  is given by

$$X_s = \varphi(\dot{X}_s; \theta) = \dot{X}_s + \left( 1 - \frac{s - t_{i_{j-1}}}{t_{i_j} - t_{i_{j-1}}} \right) X_{i_{j-1}} + \left( \frac{s - t_{i_{j-1}}}{t_{i_j} - t_{i_{j-1}}} \right) X_{i_j-}; \quad s \in (t_{i_{j-1}}, t_{i_j}) \quad (10)$$

The data augmentation scheme is now based on  $V_{mis} = (\mathbf{J}, X_J, \dot{X})$ , where  $X_J$  is  $X$  at the jump times. We define  $V_{com} = \{V_{obs}, V_{mis}\}$ , with  $V_{obs} = \mathbf{v} = \{v_{t_0}, \dots, v_{t_n}\}$ .

We obtain a likelihood function by writing the joint law of  $V_{obs}$  and  $V_{mis}$  with respect to the law of  $\mathbb{L}^{n+N} \otimes \mathbb{J}_0 \otimes \mathbb{W}_0$ , where  $\mathbb{L}^n$  is the  $n$ -dimensional Lebesgue measure,  $\mathbb{J}_0$  is the law of a jump process with unit jump rate and a parameter-free state-independent jump size distribution that preserves the required absolute continuity, and  $\mathbb{W}_0$  is the collection of standard Brownian bridges between the merged observation and jump times.

**Lemma 1.** *The likelihood of the complete data  $V_{com}$  is given by*

$$\begin{aligned}
L(V_{com}|\theta) &\propto \exp \left\{ A(x_n(\theta); \theta) - A(x_0(\theta); \theta) - \sum_{j=1}^N \Delta A(j; \theta) \right\} \\
&\times \exp \left\{ - \int_0^T \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, \varphi(\dot{X}_s; \theta); \theta) ds \right\} \prod_{j=1}^N [\lambda(j; \theta) f_g(J_j; j, \theta)] \\
&\times \prod_{i=1}^n [\eta'(V_{t_i}; \theta) \pi_{\mathbb{Q}}(X_{J_i}, x_i(\theta) | \mathbf{J}_i, x_{i-1}(\theta))], \tag{11}
\end{aligned}$$

where  $\pi_{\mathbb{Q}}(X_{J_i}, x_i(\theta) | \mathbf{J}_i, x_{i-1}(\theta)) \propto f_N(X_{i_1-}; x_{i-1}(\theta), t_{i_1} - t_{i-1}) f_N(x_i(\theta); X_{i_{N_i}}, t_i - t_{i_{N_i}})$  and  $\eta'(u; \theta)$  is the derivative of  $\eta$  w.r.t.  $u$ .

**Proof.** See Appendix C.

The algorithm is now analogous to the case where  $\sigma$  is known. Care must be exercised to keep track of all terms that depend on  $\theta$ , for example,  $\varphi(\cdot; \theta)$  and the  $x_{t_i}(\theta)$ 's.

### 3.2 A Markov chain Monte Carlo approach

We now present a Bayesian solution for the inference problem with parameter-dependent diffusion coefficient by constructing a Markov chain with stationary distribution given by the exact joint posterior distribution of  $\theta$  and the components output by JBEA.

For  $i = 1, \dots, n$ , define  $(\mathbf{J}_i^*, B_{J_i}, B_i, \Phi_i^*)$  as the random elements of the proposal for JBEA in  $[t_{i-1}, t_i]$ .  $\mathbf{J}_i^*$  is the jump process, which includes the jump times and sizes,  $B_{J_i}$  is the process at the jump times,  $B_i$  is the collection of standard Brownian bridges between these times and  $\Phi_i^*$  is the Poisson process with rate  $r_i(\theta)$  on  $[t_{i-1}, t_i] \times [0, 1]$  used in the accept/reject step of the algorithm. Defining  $I_i$  as the indicator that the proposal is accepted as a realisation of  $\tilde{\mathbb{P}}$ , we have that  $(\mathbf{J}_i^*, B_{J_i}, B_i, \Phi_i^* | I_i = 1) = (\mathbf{J}_i, X_{J_i}, \dot{X}_i, \Phi_i)$ , where  $(\mathbf{J}_i, X_{J_i}, \dot{X}_i)$  is  $(\mathbf{J}, X_J, \dot{X})$  in  $(t_{i-1}, t_i)$ . Now let  $S(i) := S(J_i, X_{J_i}, \dot{X}_i)$  be the corresponding output skeleton and  $\mathbf{S}$  be the union of all skeletons. Finally, let  $\pi(\theta)$  be the prior density of  $\theta$ .

It is likely that we are primarily interested in the posterior distribution  $\pi(\theta | \mathbf{v})$  of  $\theta$  although, depending on the application, we might also be interested in  $\pi(\mathbf{J}, X_J, \dot{X} | \mathbf{v})$ , which can be obtained from the joint posterior  $\pi(\theta, \mathbf{S} | \mathbf{v})$ . We sample from this distribution via Gibbs sampling by alternately sampling from

$$(\mathbf{S} | \theta, \mathbf{v}) \quad \text{and} \quad (\theta | \mathbf{S}, \mathbf{v}).$$

The skeletons  $S(i)$  are conditionally independent given  $\theta$  and are simulated via JBEA. The conditional density  $\pi(\theta | \mathbf{S}, \mathbf{v})$  is given in Theorem 1, which is based on Lemma 4 from Appendix B.

**Theorem 1.** *The full conditional distribution of  $\theta$  is given by:*

$$\begin{aligned} \pi(\theta|S, \mathbf{v}) &\propto \pi(\theta) \exp\{A(x_n(\theta); \theta) - A(x_0(\theta); \theta)\} \\ &\prod_{i=1}^n \left[ \exp\left\{-\Delta t_i(r_i + m_i)(\theta) - \sum_{j=1}^{N_i} (\Delta A(i_j; \theta))\right\} [r_i(\theta)]^{\kappa_i} \prod_{j=1}^{N_i} \lambda(i_j; \theta) f_g(J_{i_j}; i_j, \theta) \right. \\ &\left. \pi_{\mathbb{Q}}(X_{J_i}, x_i(\theta)|\mathbf{J}_i, x_{i-1}(\theta)) \prod_{l=1}^{\kappa_i} \left(1 - \frac{\phi(\psi_l, \dot{X}_{\psi_l}; \theta) - m_i(\theta)}{r_i(\theta)}\right) |\eta'(v_{t_i}; \theta)| \right], \end{aligned} \quad (12)$$

where

$$\phi(s, \dot{X}_s; \theta) = \left(\frac{\alpha^2 + \alpha'}{2} + \lambda\right) (s, \varphi(\dot{X}_s; \theta); \theta), \quad (13)$$

$$m_i(\theta) = \inf_{t_{i-1} \leq s \leq t_i} \phi(s, \dot{X}_s; \theta), \quad (14)$$

$$r_i(\theta) = \sup_{t_{i-1} \leq s \leq t_i} \phi(s, \dot{X}_s; \theta) - m_i(\theta), \quad (15)$$

**Proof.** *See Appendix C.*

It may or may not be possible to simulate directly from (12). If it is not possible, we use a Metropolis-Hastings step. It may be also convenient to sample  $\theta$  in blocks as the full conditionals of parameters in the jump rate and jump size distribution will typically be easy to simulate from given that conjugated priors are used. This way the MH step is only used for parameters in the drift and diffusion coefficient.

## 4 Inference when JBEA is not feasible

The two algorithms proposed in Section 3 require JBEA to be directly applied, that is, there must exist a dominating measure  $\tilde{\mathbb{F}}$  (or  $\tilde{\mathbb{D}}$ , see Section 2) from which we can simulate from and that leads to an uniformly bounded RN derivative of  $\tilde{\mathbb{P}}$  w.r.t.  $\tilde{\mathbb{F}}$ . Finding such a measure  $\tilde{\mathbb{F}}$  may be hard when the drift  $\alpha$  and/or the jump rate is unbounded.

We present two algorithms to perform exact inference which do not make direct use of JBEA, and, therefore, can be applied to a more general class of jump-diffusion models, including those with unbounded drift and/or jump rate. Formally, the two algorithms only require assumptions (a), (c), (e) and (i) from Appendix A to be satisfied.

The first approach introduces an MCEM algorithm that follows similar lines to that previously presented in Subsection 3.1 but uses an importance sampling estimate for the E-step. The second approach introduces an infinite-dimensional MCMC algorithm which is quite different from that in Subsection 3.2.

### 4.1 Importance Sampling MCEM

For simplicity, we present here the algorithm for the case the diffusion coefficient is parameter-free. The algorithm for the general case can be obtained by combining results from Subsection 3.1.2 with this.

The algorithm requires that we can find a dominating measure  $\tilde{\mathbb{F}}$  from which we can simulate and w.r.t. which  $\tilde{\mathbb{P}}$  is absolutely continuous. This option outperforms the algorithm from Subsection 3.1, when the latter is feasible, in the sense that the IS estimator has smaller variance than the rejection sampling one, given that the proposal measure from which the samples are drawn are the same.



On the other hand, the IS estimator has a higher computational cost, given the extra variables and calculations involved.

Defining  $X_{miss}$  as the missing paths of  $X$  and  $X_{com} = \{V_{obs}, X_{mis}\}$ , the expectation of the complete log-likelihood can be written as:

$$\mathbb{E}_{X_{mis}^{\tilde{\mathbb{P}}}|x_{obs},\theta'}[l(X_{com};\theta)] = \mathbb{E}_{X_{mis}^{\tilde{\mathbb{P}}}|x_{obs},\theta'} \left[ \frac{d\tilde{\mathbb{P}}}{d\mathbb{F}}(X_{mis};\theta')l(X_{com};\theta) \right]. \quad (16)$$

Defining  $w = \frac{d\tilde{\mathbb{P}}}{d\mathbb{F}}(X_{mis};\theta')$ , Lemma 5 (see Appendix B) combined with the Markov property gives

$w = \prod_{i=1}^n w_i$ , where

$$\begin{aligned} w_i &= \exp \left\{ - \int_{t_{i-1}}^{t_i} \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, X_{s-}; \theta') ds \right\} \exp \left\{ - \sum_{j=1}^{N_i} \Delta A(i_j; \theta) \right\} \times \kappa_{1i}(X, \theta') \\ &\times \prod_{j=1}^{N_i} \frac{\lambda(i_j; \theta') f_g(J_{i_j}; i_j, \theta')}{\lambda_0(\theta') f_i(J_{i_j}; \theta')} \exp \left\{ - \frac{1}{2\Delta t_i} (\Delta x_i - J_i)^2 \right\} \kappa_{2i}(\theta'), \end{aligned} \quad (17)$$

where  $\kappa_{2i}(\theta')$  is a constant with respect to  $X$ ,  $\kappa_{1i}(X; \theta) = \frac{d\tilde{\mathbb{P}}}{d\mathbb{F}}(X_{mis}; \theta)$ , for  $X$  in  $[t_{i-1}, t_i]$ , and  $\Delta x_i = x_i(\theta) - x_{i-1}(\theta)$ . Furthermore, we also have that  $l(X_{com}; \theta) = \sum_{i=1}^n l_i(\theta)$ , where

$$\begin{aligned} l_i(\theta) &= A(x_{i-1}; \theta) - A(x_i; \theta) - \sum_{j=1}^{N_i} (\Delta A(i_j; \theta)) - \int_{t_{i-1}}^{t_i} \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, X_{s-}; \theta) ds \\ &+ \sum_{j=1}^{N_i} \log(\lambda(i_j; \theta)) + \log(f_g(J_{i_j}; i_j, \theta)). \end{aligned} \quad (18)$$

The first natural choice of an estimator for (16) is

$$E_1(\theta) = \frac{1}{M} \sum_{l=1}^M \sum_{i=1}^n \left( \prod_{k=1}^n w_k^{(l)} \right) l_i^{(l)}(\theta), \quad (19)$$

where  $M$  is the number of Monte Carlo samples.

Nevertheless, an improved estimator can be devised based on the following result.

**Proposition 1.**

$$\mathbb{E}_{X_{mis}^{\tilde{\mathbb{P}}}|x_{obs},\theta'} [l(\theta)] = \sum_{i=1}^n \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [w_i l_i(\theta)]. \quad (20)$$

**Proof.** See Appendix C.

This result leads to the following estimator

$$E_2(\theta) = \frac{1}{M} \sum_{l=1}^M \sum_{i=1}^n \left( w_i^{(l)} l_i^{(l)}(\theta) \right). \quad (21)$$

Estimator  $E_1$  is a global IS estimator whilst estimator  $E_2$  is a local one. The latter is expected to have a much better behavior than the former. In fact, we have the following result.

**Proposition 2.**

$$\text{Var}[E_2] \leq \text{Var}[E_1]. \quad (22)$$

**Proof.** See Appendix C.

In order to devise an exact algorithm we need to avoid the calculation of the integrals in the expressions of  $w_i$  and  $l_i(\theta)$ . This is achieved by firstly defining the auxiliary variables:  $K = (K_1, \dots, K_n)$ ,  $K_i \sim \text{Poi}(\mu_i \Delta t_i)$ ,  $U = (U_1, \dots, U_n)$ ,  $U_i = (U_{i1}, \dots, U_{iK_i})$ ,  $U_{il} \sim \mathcal{U}(t_{i-1}, t_i)$ ,  $\dot{U} = (\dot{U}_1, \dots, \dot{U}_n)$ ,  $\dot{U}_i \sim \mathcal{U}(t_{i-1}, t_i)$ ,  $\forall l = 1, \dots, K_i$ ,  $\forall i = 1, \dots, n$ , where all the components of  $K$ ,  $U$  and  $\dot{U}$  are independent. Now, using the same strategy from Subsection 3.1.1 and the ideas from the Poisson estimator (see Beskos et al., 2006b), we have that

$$\mathbb{E}_{X_{m_{is}}^{\tilde{\mathbb{F}}}|x_{obs}, \theta'}[w_i l_i(\theta)] = \mathbb{E}_{X_{m_{is}}^{\tilde{\mathbb{F}}}, K, U, \dot{U}|x_{obs}, \theta'} \left[ f_{0i} e^{(\mu_i - b_i) \Delta t_i} \mu_i^{-K_i} \prod_{l=1}^{K_i} (b_i - f_{1i}) (f_{2i}(x) - \Delta t_i f_{3i}) \right] \quad (23)$$

$$\begin{aligned} f_{0i} &= \kappa_{1i}(X; \theta') \prod_{j=1}^{N_i} \frac{\lambda(i_j; \theta') f_g(J_{ij}; i_j, \theta')}{\lambda_i(\theta') f_i(J_{ij}; \theta')} \exp \left\{ - \sum_{j=1}^{N_i} (\Delta A(i_j; \theta')) - \frac{1}{2 \Delta t_i} (\Delta x_i - J_i)^2 \right\} \kappa_{2i}; \\ f_{1i} &= \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (U_{il}, X_{U_{il}-}; \theta'); \quad f_{3i} = \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (\dot{U}_i, X_{\dot{U}_i-}; \theta); \\ f_{2i} &= A(x_{i-1}; \theta) - A(x_i; \theta) - \sum_{j=1}^{N(i)} (\Delta A(i_j; \theta)) + \sum_{j=1}^{N_i} \log(\lambda(i_j; \theta)) + \log(f_g(J_{ij}; i_j, \theta)). \end{aligned} \quad (24)$$

We now need to simulate  $K$ ,  $U$ ,  $\dot{U}$ ,  $X_{\dot{U}_i}$ 's,  $X_{U_{il}}$ 's and jump times and sizes to obtain the IS estimate of the expectations on the rhs of (20). Naturally, the choices of  $b_i$  and  $\mu_i$  are closely related to the efficiency of the algorithm. Based on the results from Beskos et al. (2006b) and Fearnhead et al. (2008) we recommend  $b_i = \bar{U}_i$  and  $\mu_i = \bar{U}_i - \bar{L}_i$ , where  $\bar{L}_i \leq f_{1i} \leq \bar{U}_i$ .

The bounds  $\bar{U}_i$  and  $\bar{L}_i$  are obtained by sampling layers for the standard Brownian bridges from  $\tilde{\mathbb{F}}$ . For  $j = 1, \dots, N_i + 1$ , let  $\dot{u}_{ij}$  and  $\dot{l}_{ij}$  be upper and lower bounds for the standard BB in  $(t_{i,j-1}, t_{i,j})$  obtained from the layers simulation, we make

$$\bar{U}_i = \sup_j \sup \{f_{1i}; U_{il} \in (t_{i,j-1}, t_{i,j}), X_{U_{il}-} \in \bar{H}_{ij}\}; \quad (25)$$

$$\bar{L}_i = \inf_j \inf \{f_{1i}; U_{il} \in (t_{i,j-1}, t_{i,j}), X_{U_{il}-} \in \bar{H}_{ij}\}, \quad (26)$$

where  $\bar{H}_{ij} = [\min\{X_{i,j-1}, X_{i,j-}\} + \dot{l}_{ij}, \max\{X_{i,j-1}, X_{i,j-}\} + \dot{u}_{ij}]$ . Details about the layers simulation are given in Appendix E of the supplementary material.

We now have a new estimator

$$\dot{E}_2(\theta) = \frac{1}{m} \sum_{l=1}^m \sum_{i=1}^n \left( \dot{w}_i^{(l)} \dot{l}_i^{(l)}(\theta) \right), \quad (27)$$

where  $\dot{w}_i$  and  $\dot{l}_i(\theta)$  are obtained by rewriting  $w_i$  and  $l_i(\theta)$  using the result in (23).

Note that Proposition 2 does not guarantee any properties of the estimator related to finite variance, in particular, that the weights have finite variance. This condition does not assure that the estimator also has finite variance (it does whenever the target function is bounded), but weights with infinite variance will typically lead to infinite variance estimators. In fact, we have the following result.

**Lemma 2.** Let  $b_i = \bar{U}_i$  and  $\mu_i = \bar{U}_i - \bar{L}_i$ . Then, the weights of the IS estimator have finite variance if the two conditions below are satisfied:

(a)  $f_{1i}$  is uniformly bounded below,  $\forall i$ ;

(b)  $\mathbb{E} [(f_{0i})^2] < \infty, \forall i$ .

**Proof.** For  $f_{1i} \geq l_i \in \mathbb{R}$ ,  $\mathbb{E} [\dot{w}_i^2] \leq \mathbb{E} [(f_{0i})^2 \exp(-2l_i \Delta t_i)] = \exp(-2l_i \Delta t_i) \mathbb{E} [(f_{0i})^2]$ .

Condition (b) from Lemma 2 is true when JBEA is feasible (the drift and jump-rate are bounded) but is hard to check in other cases. Furthermore, we cannot ignore the constants  $\kappa_{2i}$ , which depend on the (unknown) transition density, and potentially unknown constants from  $\kappa_{1i}(X; \theta)$ . Nevertheless, we can modify  $\dot{E}_2$  in a way that the constants can be ignored and also get a nice property regarding the variance of the estimator. The fact that  $\mathbb{E}[\dot{w}_i] = 1$  suggests the following estimator:

$$\dot{E}_3(\theta) = \sum_{i=1}^n \sum_{j=1}^m \left( \tilde{w}_i^{(j)} \dot{l}_i^{(j)}(\theta) \right), \quad (28)$$

where  $\tilde{w}_i^{(j)} = \frac{\dot{w}_i^{(j)}}{\sum_{j=1}^m \dot{w}_i^{(j)}}$ . Estimator  $\dot{E}_3$  is biased but has some nice properties as stated in the following lemma.

**Lemma 3.**  $\dot{E}_3$  is a strongly consistent estimator for the expectation from the E-step and has finite variance whenever  $\dot{l}_i$  has finite variance under  $(\tilde{\mathbb{F}} \otimes \mathbb{A})$ , where  $\mathbb{A}$  is the joint probability measure of  $(K, U, \dot{U})$ .

**Proof.** See Appendix.

Note that this lemma is valid, in particular, when  $\tilde{\mathbb{F}} \equiv \tilde{\mathbb{D}}$ . In fact, this will typically be the case when JBEA is not feasible. Moreover, although  $\dot{E}_3$  is biased, the lemma implies that it is asymptotically unbiased and it is expected that the number of Monte Carlo samples  $M$  is typically large enough to make the bias negligible.

Finally, we also construct an estimator for the covariance matrix of the MLE, which is presented in Appendix D of the supplementary material.

## 4.2 An infinite-dimensional Barker's MCMC

Because JBEA is not feasible, we are unable to find a finite-dimensional representation of the jump-diffusion bridges for which we know and can simulate from the full conditional of the parameters given this representation. Instead, the complete bridges  $V_{mis}$  (as define in Subsection 3.1.2) need to be considered. Our MCMC algorithm iterates between the following sampling steps:

$$(V_{mis} | \theta, \mathbf{v}) \quad \text{and} \quad (\theta | V_{mis}, \mathbf{v}). \quad (29)$$

The first step is complicated by the lack of JBEA and the need to somehow store the entire missing paths  $V_{mis}$ . Nevertheless, we can find a solution without recouring to approximations.

Since direct simulation from the full conditional distributions is infeasible, both  $V_{mis}$  and  $\theta$  are sampled using an independence and a random walk accept-reject procedure, respectively, involving the alternative Barker's step (Barker, 1965). To implement the method, we only need to unveil  $V_{mis}$  at a finite collection of time points.

The main contribution of this algorithm when compared to the one from Subsection 3.2 is that it may be applied to a quite general class of jump-diffusion processes, including processes with unbounded drift and/or unbounded jump rate.

### 4.2.1 Sampling the missing paths

The missing paths are sampled from  $(V_{mis}|\theta, \mathbf{v})$  via Barker's step. The Barker's algorithm was proposed in Barker (1965) and works similarly to the MH algorithm. A proposal is simulated from an arbitrary distribution and accepted according to an acceptance probability that preserves detailed balance. The probability to go from  $x$  to  $y$  when proposing from  $q(x, y)$  is given by

$$\frac{\pi^*(y)q(y, x)}{\pi^*(x)q(x, y) + \pi^*(y)q(y, x)}, \quad (30)$$

where  $\pi^*$  is the invariant distribution of the chain. Barker's method is not as popular as the prolific Metropolis-Hastings algorithm since it is easy to demonstrate that Barker is uniformly dominated by MH in terms of its convergence properties (see Peskun, 1973). However it is also easy to see that Barker is no worse than twice as slow as Metropolis-Hastings and its convergence properties are broadly comparable (Łatuszyński and Roberts, 2013). For us, the smoothness of the Barker's acceptance probability function turns out to be crucial in obtaining a feasible algorithm.

For a given interval  $(t_{i-1}, t_i)$ , let  $X$  be the current state of the chain and a new proposal  $B$  is drawn from  $\mathbb{D}$ . We use  $\mathbb{D}$  as our proposal distribution and so it is convenient to use it as the dominating measure. Lemma 5 (see Appendix B) shows that the acceptance probability can be written as

$$\alpha_X = \frac{\frac{d\tilde{\mathbb{P}}}{d\mathbb{D}}(B)}{\frac{d\tilde{\mathbb{P}}}{d\mathbb{D}}(X) + \frac{d\tilde{\mathbb{P}}}{d\mathbb{D}}(B)} = \frac{s_k p_k}{s_{k-1} p_{k-1} + s_k p_k}, \quad (31)$$

where

$$s_{k-1} = \exp \left\{ - \sum_{j=1}^{N_i} (\Delta A(i_j; \theta)) - \mathcal{I}_{a_i}(\theta) - \frac{1}{2\Delta t_i} (\Delta x_i - J_i)^2 \right\} \prod_{j=1}^{N_i} \frac{\lambda(i_j; \theta) f_g(J_{i_j}; i_j, \theta)}{\lambda_0(\theta) f(J_{i_j}; \theta)}, \quad (32)$$

$$p_{k-1} = \exp \left\{ - \int_{t_{i-1}}^{t_i} \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s; X_{s-}; \theta) - a_i(s; \theta) ds \right\}, \quad (33)$$

for  $\mathcal{I}_{a_i} = \int_{t_{i-1}}^{t_i} a_i(s; \theta) ds$  and  $a_i(s; \theta)$  is any lower bound satisfying

$$a_i(s; \theta) \leq \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, X_{s-}; \theta). \quad (34)$$

Additionally,  $s_k$  and  $p_k$  are obtained by replacing  $X$  by  $B$  in (32)-(34). The choice of  $a_i(s; \theta)$  has direct impact on the efficiency of the algorithm, as it is discussed in Subsection 4.2.2.

To perform the accept/reject step of the algorithm, we need to simulate a *Bernoulli*( $\alpha_X$ ) random variable. Most importantly, we have to use only a finite-dimensional representation of the missing paths to do it. That is achieved by the *two-coin algorithm* given in the following proposition.

**Proposition 3.** *Suppose we want to simulate a Bernoulli( $\alpha_X$ ) random variable, where  $s_k$  and  $s_{k-1}$  are known positive numbers and it is possible to simulate events of unknown probabilities given by  $p_k$  and  $p_{k-1}$ . The following algorithm outputs an exact draw of this Bernoulli random variable:*

## Two-Coin algorithm

1. Sample  $C_1$  from  $\{0, 1\}$ , where  $P(C_1 = 1) = \frac{s_k}{s_{k-1} + s_k}$ ;
2. if  $C_1 = 1$ , sample  $C_2$  from  $\{0, 1\}$ , where  $P(C_2 = 1) = p_k$ ;
  - if  $C_2 = 1$ , output 1;
  - if  $C_2 = 0$ , go back to 1;
3. if  $C_1 = 0$ , sample  $C_2$  from  $\{0, 1\}$ , where  $P(C_2 = 1) = p_{k-1}$ ;
  - if  $C_2 = 1$ , output 0;
  - if  $C_2 = 0$ , go back to 1.

**Proof.** Let  $q$  be the probability that there is no output in one trial of  $C_1$  and  $C_2$ , that is  $q = \frac{s_k(1 - p_k) + s_{k-1}(1 - p_{k-1})}{s_k + s_{k-1}}$ . Then, the probability that the algorithm outputs 1 is  $\frac{s_k}{s_k + s_{k-1}} p_k \sum_{i=0}^{\infty} q^i = \frac{s_k p_k}{s_k p_k + s_{k-1} p_{k-1}}$ . Analogous calculations for the probability that the algorithm outputs 0 complete the proof.

In order to simulate events of probability  $p_k$  and  $p_{k-1}$  we use the Poisson Coin described in Section 2. The algorithm stores a finite-dimensional representation of the missing path. It is crucial though to be able to simulate further points, given this skeleton, on the  $\theta$  step of the Gibbs sampler, as we discuss in Subsection 4.2.2 below.

### 4.2.2 Sampling the parameters

Typically, the full conditional distribution of the parameters will depend on the unknown integral in (38). In order to minimise the complexity of the algorithm, the parameters should firstly be separated into two blocks: the first -  $\theta_1$ , consisting of those parameters whose full conditionals depend on the integral in (38), and the second block -  $\theta_2$ , consisting of the remaining parameters. Parameters in  $\theta_2$  are sampled as in an ordinary tractable MCMC - they may be broken into smaller blocks, sampled directly from the full conditional or via MH steps, conveniently. Parameters in  $\theta_1$  are sampled via Barker's step, which may be performed separately for sub-blocks. The full conditional distribution of any block  $\theta_b$  of  $\theta$  is given by

$$\pi(\theta_b | V_{mis}, \mathbf{v}) \propto \pi(V_{mis}, \mathbf{v} | \theta, \cdot) \pi(\theta), \quad (35)$$

where  $\pi(V_{mis}, \mathbf{v} | \theta, \cdot)$  is given by Lemma 1.

Proposals are drawn from a symmetric random walk -  $\theta_b^{(k)} = \theta_b^{(k-1)} + \epsilon$ , where  $\epsilon$  is a r.v. symmetric around 0 with an appropriately tuned covariance matrix  $\Sigma$  and a new value  $\theta_b^{(k)}$  is accepted with probability:

$$\alpha_{\theta_b} = \frac{\pi(\theta_b^{(k)} | V_{mis}, \mathbf{v})}{\pi(\theta_b^{(k-1)} | V_{mis}, \mathbf{v}) + \pi(\theta_b^{(k)} | V_{mis}, \mathbf{v})} = \frac{s_k p_k}{s_{k-1} p_{k-1} + s_k p_k}, \quad (36)$$

where

$$s_k = \exp \left\{ A(x_n(\theta_b^{(k)}); \theta_b^{(k)}) - A(x_0(\theta_b^{(k)}); \theta_b^{(k)}) - \sum_{i=1}^n \mathcal{I}_{a_i}(\theta_b^{(k)}) - \sum_{j=1}^N \Delta A(i_j; \theta_b^{(k)}) \right\} \\ \times \prod_{j=1}^N \left[ \lambda(j; \theta_b^{(k)}) f_g(J_j; j, \theta_b^{(k)}) \right] \prod_{i=1}^n \left[ \eta'(V_i; \theta_b^{(k)}) \pi_{\mathbb{Q}} \left( X_{J_i}, x_i(\theta_b^{(k)}) | \mathbf{J}_i, x_{i-1}(\theta_b^{(k)}) \right) \right] \pi(\theta_b^{(k)}), \quad (37)$$

$$p_k = \exp \left\{ - \int_0^T \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, \varphi(\dot{X}_s; \theta_b^{(k)}); \theta_b^{(k)}) - a(s; \theta_b^{(k)}) ds \right\}, \quad (38)$$

$$a(s; \theta_b^{(k)}) = a_i(s; \theta_b^{(k)}), \text{ for } s \in [t_{i-1}, t_i]. \quad (39)$$

The expressions for  $s_{k-1}$  and  $p_{k-1}$  are obtained by replacing  $\theta_b^{(k)}$  by  $\theta_b^{(k-1)}$  in (37)-(39).

The efficiency of this sampling step depends on the efficiency of the two coin algorithm which ultimately relies on the probabilities of success of  $C_2$ ,  $p_{k-1}$  and  $p_k$ . The smaller these probabilities are, the higher is the expected number of trials per iteration. Moreover, at every trial, the missing paths have to be unveiled at extra time points, which also increases computational cost. The optimisation of  $p_{k-1}$  and  $p_k$  is related to the optimisation of the lower bounds  $a_i$ . An efficient solution is presented in Appendix F of the supplementary material.

We simulate the second coin piecewise by simulating a sequence of “sub-coins” with probability

$$p_{k,i_j} = \exp \left\{ - \int_{t_{i_j-1}}^{t_{i_j}} \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, \varphi(\dot{X}_s; \theta_b^{(k)}); \theta_b^{(k)}) - a_i(s; \theta_b^{(k)}) ds \right\}. \quad (40)$$

If we get 0 at some point, the two-coin algorithm is restarted.

In order to simulate each sub-coin  $p_{k,i_j}$ , we apply the Poisson Coin algorithm by simulating a Poisson process with rate  $r_{ij}(\theta_b^{(k)})$ , as defined in Appendix F of the supplementary material. The algorithms to simulate layers and bridges given layers are presented in Appendix E of the supplementary material.

The overall MCMC algorithm is the following.

#### Barker’s MCMC for jump-diffusions

1. Provide initial values for all the parameters  $\theta$ ;
2. make  $k=1$ ;
3. sample  $V_{mis}^{(k)}$  via Barker’s by proposing from the measure  $\tilde{\mathbb{D}}$  defined in Section 2 (with BB layers) and accepting with probability  $\alpha_X$  in (31) using the Two-Coin algorithm;
4. sample each block of  $\theta_1^{(k)}$  via Barker’s by proposing from a symmetric random walk and accepting with probability  $\alpha_\theta$  in (36) using the Two-Coin algorithm;
5. sample the remaining parameters  $\theta_2^{(k)}$  directly from their full conditionals or via Gaussian random walk MH;
6. to continue running the chain, make  $k = k + 1$  and GOTO 3, otherwise, STOP.

### 4.2.3 Improving the MCMC algorithm

The efficiency of the proposed MCMC algorithm in terms of convergence and computing cost depends on many factors. Here we shall discuss strategies for improvements.

In order to avoid numerical problems when computing the probability of  $C_1$  in the Two-Coin algorithm, compute  $\dot{s}_i = \log(s_i)$ , for  $i = k - 1, k$  and then  $(1 + \exp(\dot{s}_{k-1} - \dot{s}_k))^{-1}$ .

A way to improve the mixing of the chain is to add an extra step to the Gibbs sampler to sample the missing (continuous) paths between observation and jump times using the EA algorithm (see Beskos et al., 2006a). We only require the condition that  $(\alpha^2 + \alpha')(u)$  is bounded below for all  $u$  in the state space of  $X$ . EA performs rejection sampling by proposing from a Brownian bridge and accepting with probability given by  $\exp \left\{ - \int_{t_{i-1}}^{t_i} \left( \frac{\alpha^2 + \alpha'}{2} \right) (X_s) - l ds \right\}$ , where  $l = \inf_u \left( \frac{\alpha^2 + \alpha'}{2} \right) (u)$ .

Since this algorithm will output an exact draw of the missing paths between observation and jump times, it guarantees that these bridges are updated at every iteration of the chain at least once. This strategy ought to improve the mixing of the chain considerably. Furthermore, it eliminates the

problem of accumulating too many bridge points along the iterations of the chain due to rejections on the Barker's step. It is recommended to add one EA step after each Barker's step - for  $X$  or  $\theta$ -blocks. The example of Section 5.2 implements this extra update step.

Further improvement in the chain mixing could be obtained by modifying the Barker's step for  $V_{mis}$ , which is the only step where the jump process is updated. One idea is to perform multiple steps of this type. Another simple and virtually costless strategy is to adopt a pilot analysis to tune the jump rate based on the average number of jumps in each interval. Moreover, if one has reasonable choices for initial values of the parameters it may be a good idea to warm up the chain in the first iterations before start updating the parameters.

In order to increase the success probability of  $C_2$  in the Two-Coin algorithm we need to improve the lower bound  $a(s, \theta)$ , which can be done by tightening the lower and upper bounds for the missing paths in  $V_{mis}$ . That is achievable by performing what we call the *layer refinement algorithm* which, instead of simulating layers for the standard bridge between times  $t_{i_j}$  (observation or jump times), simulates layers for shorter intervals. For a constant  $m \in \mathbb{N}$  and a time interval  $(t_{i_{j-1}}, t_{i_j})$ , we first simulate the standard BB at times  $t_{i_{j-1}} + k\Delta t_{i_j}/m$ , for  $k = 1, \dots, m-1$ , and then simulate layers for each of the sub-intervals obtained. Naturally, the refined layers will provide tighter bounds for the standard bridge and, consequently, for the  $X$  path. In particular, for a standard bridge of length  $t$ , the range is  $\mathcal{O}(\sqrt{t})$ . One may also set different refinement levels along the intervals of consecutive observations. It is typically straightforward to identify the relation of the Two-coin cost with the values of  $X$ , so the level of refinement may be set as a function of the extreme observations of each interval. The bounds for the  $X$  path are obtained by the formulation provided in Appendix F of the supplementary material. A cheaper version of the EA step described above consists in updating the missing paths between observation, jump and refinement points, conditional on the paths' bounds.

The layer refinement algorithm suggests that the algorithm that simulates only the first coin  $C_1$  may be seen as a discretised method and it is the second coin that guarantees the exactness of the algorithm. In that case, the larger  $m$  is the smaller the error due to the discretisation.

Two other strategies may increase the probability of the second coins. The first one is to break  $\theta_1$  into smaller blocks (the extreme case being one parameter per block). This may help in the sense that a smaller block may simplify the function inside the integral in (38) and allow for a more efficient lower bound  $a(s; \cdot)$ . The second strategy is to divide the numerator and denominator of the Barker's acceptance probability by  $p_{k-1}$  or  $p_k$ . This will make the probability of one of the second coins equals 1 and possibly increase the probability of the other one. In order to choose between  $p_{k-1}$  and  $p_k$  one has to look at the resulting ratio of  $p$ 's and recognise which choice has a ratio of the form  $s_0 \int_0^T \phi_2(X_s) ds$ , where  $s_0$  is computable and  $\phi_2$  is a non-negative function. This choice may vary among the chain's iterations, depending on the proposed value. If this strategy is adopted, the success probability of one of the  $C_2$  coins will be 1 and the other one will typically get smaller as the size of the random walk step increases. For this reason, it may be wise to truncate the random walk distribution (if it is Gaussian) between say  $\pm 3.5$  or  $\pm 4$  standard deviations to avoid the algorithm from collapsing after an average number of  $1/\epsilon$  iterations -  $\epsilon$  being the probability of proposing extreme values. This would have very little effect on the algorithms's convergence properties. Another reasonable strategy is to use a uniform random walk. Another reasonable strategy is to perform multiple updates using smaller variance proposals. The multiple proposals ought to compensate the slower mixing but with considerable gains in computational cost.

### 4.3 Practical implementation

Identifiability is a particular problem when dealing with jump-diffusions. It is crucial to have enough information to distinguish well between continuous and jump variation. Practical strategies to tackle the problem include fixing some of the parameters at reasonable values, which is not always easy, or using informative priors under a Bayesian approach. A general idea that should always be considered

is that of admitting the least possible number of jumps necessary to get a good fit so that the jumps only occur when a pure diffusion process is not flexible enough to model the phenomenon of interest. This not only mitigates the identifiability problem but also favors the interpretability of jumps and parameters and, finally, ought to improve computational efficiency.

Computational cost is another important issue when dealing with the algorithms proposed in this paper. Since the MCEM and MCMC algorithms from Section 3 are considerably cheaper than the algorithms presented in section 4, we may consider some practical strategies to allow the use of the former ones. For example, we may occasionally truncate the jump rate.

We can also reduce the computational cost of JBEA by making its proposal as close as possible to the target. This is also a good idea when using the algorithms from Section 4 - it would reduce the variance of the weights in the ISMCEM algorithm and increase the success probability of  $C_2$  in the MCMC one. The idea is basically to make the proposal jump process (jump rate and jump size distribution) depend on time and/or on the extremes of the interval whenever the target jump process is time and/or state dependent. The time dependence can always be mimicked from the target and the state replaced by a function of the extremes, for example, the mean.

## 5 Simulated examples

In this section we present results from some simulated examples. Firstly, we present an example with bounded drift and bounded jump-rate to apply the algorithms from Section 3. Secondly, the algorithms from Section 4 are applied to an unbounded drift example.

### 5.1 An introductory example

We consider the following model:

$$\begin{aligned} dV_s &= -\tanh(V_s - \delta)ds + \sigma dW_t + dJ_s, & V_0 = v, \quad s \in [0, 1000] \\ \lambda_1(s; V_{s-}) &= \lambda(1 - \tanh^2(V_s - \delta)), \\ g_1(Z_j, V_{t_j-}) &= Z_j \sim \mathcal{N}(\mu, \tau^2), \end{aligned} \tag{41}$$

with  $(\delta, \sigma^2, \lambda, \mu, \tau^2) = (0, 1, 0.1, 2, 0.35^2)$ . Note that this parameter configuration is bounded to cause identifiability problems as potential variations due to  $\sigma$  and jumps overlap, in a certain sense. Results from the MCEM algorithm of Section 3.1.2 and the MCMC algorithm of Section 3.2 are presented in Tables 1 and 2, respectively. The M-step from the MCEM algorithm is performed numerically via the quasi-Newton method BFGS (see, for example, Fletcher, 1987, Section 3.2).

Table 1: Iterations from the MCEM algorithm. 1000 Monte Carlo samples are used in all iterations and the M-step is performed via . Results obtained with other initial conditions (away from the real values) suggest the presence of local modes.

Iteration	$\delta$	$\sigma^2$	$\lambda$	$\mu$	$\tau^2$
0	0.5	1.5	0.2	2	0.25
1	0.149	1.299	0.104	1.771	0.212
5	0.078	1.128	0.096	1.685	0.197
20	0.023	1.031	0.128	1.770	0.203
100	0.005	0.996	0.133	1.848	0.175
200	0.008	0.996	0.128	1.889	0.146
500	0.019	1.001	0.120	1.942	0.100
real	0	1	0.1	2	0.1225

The MCMC algorithm also outputs a sample from the posterior distribution of the jump process. Some interesting posterior statistics can be obtained from this distribution. Figure 1 shows an example.

We also run the algorithms fixing parameters  $\mu$  and  $\tau^2$  at their real values. Results reinforce the issues concerning identifiability. Table 3 shows the results obtained for the two algorithms.



Table 2: Posterior statistics from the MCMC output. Uniform improper priors are adopted for all parameters. The prior of  $\tau^2$  had to be truncated to be above 0.008 to avoid getting trapped in small values. The chain runs for 500k iterations. The parameters are jointly sampled using an adaptive Gaussian random walk MH step which had a 0.31 acceptance rate. Trace plots suggest convergence has been achieved.

	$\delta$	$\sigma^2$	$\lambda$	$\mu$	$\tau^2$
Mean	-0.025	0.993	0.179	1.69	0.291
Median	-0.021	0.991	0.164	1.67	0.275
Mode	-0.023	0.987	0.135	1.54 / 1.66	0.025 / 0.280
St. Dev.	0.086	0.068	0.080	0.30	0.180
real	0	1	0.1	2	0.1225




Figure 1: Posterior probability of jump between each pair of consecutive observations. Solid circles represent intervals where a jump really exists and have size proportional to the size of the jump.

Table 3: Results for the case where  $\mu$  and  $\tau^2$  are fixed at their real values. The specifications of the two algorithms are the same as in the previous run. The output of the MCEM algorithm corresponds to iteration 55, for which it has clearly converged.

	$\delta$	$\sigma^2$	$\lambda$
MCEM	0.024	1.004	0.113
MCMC			
Mean	0.021	1.009	0.117
Median	0.022	1.007	0.115
Mode	0.024	1.004	0.113
St. Dev.	0.064	0.062	0.028
95% Cred. Int.	(-0.107,0.146)	(0.895,1.137)	(0.066,0.177)
real	0	1	0.1

## 5.2 An unbounded drift example: the Ornstein Uhlenbeck process

We now present a simulated example where the drift is unbounded and the methodology from Section 4 is applied. We consider the following model:

$$\begin{aligned}
 dV_s &= -\rho(V_s - \mu)ds + dW_t + dJ_s, & V_0 &= v, & s &\in [0, 500] \\
 \lambda_1(s; V_{s-}) &= \lambda, \\
 g_1(Z_j, V_{t_j-}) &= Z_j \sim \text{Exp}(\theta),
 \end{aligned} \tag{42}$$

with  $(\rho, \mu, \lambda, \theta) = (1, 0, 0.07, 1)$ . Identifiability problems are likely to occur due to the ambiguity involving variation of the continuous part and small jumps from the Exponential distribution. Results from the MCEM algorithm of Section 4.1 are presented in Table 4. The M-step is performed analytically. An algorithm based on estimator  $E_1$  for fixed  $\lambda$  was also implemented and the Monte Carlo variance was too high even for  $8 \times 10^5$  samples.

The convergence of the MCEM algorithm is slow, caused by irregularities of the likelihood function

Table 4: Iterations from the MCEM algorithm using estimator  $\hat{E}_3$ .

Iteration	$\rho$	$\mu$	$\lambda$	$\theta$	MC samples ( $\times 10^3$ )
0	1.5	-0.5	0.5	1.5	
1	1.459	-0.231	0.651	1.245	20
10	1.185	-0.105	0.605	1.667	50
100	1.053	0.021	0.281	1.486	50
200	1.005	0.099	0.094	0.926	50
300	0.989	0.117	0.066	0.801	50
312	0.996	0.114	0.067	0.748	600
316	1.001	0.114	0.066	0.760	600
real	1	0	0.07	1	

related to parameter identifiability. If we fix the jump rate at its real value, we get the estimates  $\hat{\rho} = 1.019$ ,  $\hat{\mu} = 0.112$  and  $\hat{\theta} = 0.780$  (after just 7 iterations) and the estimated covariance matrix

$$\begin{pmatrix} 0.0025367 & 0.00022448 & -0.0011035 \\ 0.00022448 & 0.0015771 & 0.00059542 \\ -0.0011035 & 0.00059542 & 0.038611 \end{pmatrix}$$

For the MCMC algorithm of Section 4.2, an EA step between observation, jump and refinement points is performed after each Barker's step. The chain is warmed-up for  $5k$  iterations before the parameters start to be updated. The layer refinement idea described in Subsection 4.2.3 is applied for  $m = 4$ . The parameter vector is broken into four individual blocks. Parameters from the jump process  $(\lambda, \theta)$  are sampled directly from their full conditionals and the other two parameters are sampled via Two-Coin Barker's. We adopt uniform prior distributions for  $\mu$  and  $\rho$  and the following informative priors:  $\lambda \sim Exp(50)$  and  $\theta \sim Gamma(7, 6)$ . As it is explained in section 4.3, these informative priors play a crucial role in the identifiability of model.

We set uniform random walk proposals for  $\mu$  and  $\rho$  having a higher acceptance rate than optimal but perform six alternate updates of each one on each iteration of the Gibbs sampling. Also, three consecutive updates of  $X$  are performed on each iteration. Trace plots indicate good convergence. Posterior statistics are presented in Table 5. The chosen prior distributions seem to correct the irregularities of the likelihood and lead to reasonably good results.

Table 5: Posterior statistics from the MCMC output.

	$\rho$	$\mu$	$\lambda$	$\theta$
Mean	1.010	0.086	0.050	0.725
Median	1.010	0.086	0.046	0.709
Mode	1.013	0.086	0.039	0.667
St. Dev.	0.0719	0.051	0.025	0.189
95% Cred. Int.	(0.872,1.152)	(-0.014,0.188)	(0.013,0.111)	(0.396,1.131)
real	1	0	0.07	1

## 6 Application

### 6.1 Exchange rate USD $\times$ GBP

We consider the exchange rate between USD and GBP. The USD suffered a considerable depreciation during the 2008 world economic crisis. A few months later it had a moderate recovery and oscillated between 1.45 and 1.7 until beginning of 2016. We will use daily data from May 21, 2009, which was right after the recovery, up to March 27, 2013. This constitutes 1201 data points and is shown in Figure 2 in the log-scale. We choose to fit a scaled Brownian motion to model the continuous part. We shall adopt the MCMC procedure of Section 3.2.

In order to specify the jump component of our model, we look at the graph of the differences between consecutive observations. We obtain an empirical estimate of the number of jumps which

suggests that a piecewise constant jump rate should be enough to reasonably accommodate the data. We also choose the jump size distribution to be a mixture of a positive and a negative gamma distribution to take probability mass away from zero and avoid identifiability problems. The chosen model for the log-rate  $V$  is given by:

$$\begin{aligned} dV_s &= bdW_s + dJ_s, \quad V_0 = v, \quad s \in [0, 1201] \\ \lambda_1(s; V_{s-}) &= \lambda_i, \quad \text{for } s \in A_i, \quad i = 1, \dots, 5 \\ g_1(Z_j, V_{t_j-}) &= Z_j \sim p\mathcal{G}(8, 1000) + (1 - p)(-\mathcal{G}(8, 1000)), \end{aligned} \quad (43)$$

for  $A_1 = [0, 370]$ ,  $A_2 = (370, 825]$ ,  $A_3 = (825, 1000]$ ,  $A_4 = (1000, 1120]$  and  $A_5 = (1120, 1201]$ .

We try to avoid identifiability problems by minimising the number of jumps and maximising  $b$ . This is done by setting informative priors to  $\lambda_1$  and  $\lambda_2$  and by adopting the jump size distribution in (43). We also specify the model in a way that between two consecutive observations there can only be positive jumps if the process goes up and only negative jumps if the process goes down. This is done considering the corresponding distribution from the mixture. Parameters are assumed to be mutually independent with marginal priors:  $b^2, \lambda_3, \lambda_4, \lambda_5 \sim \mathcal{U}(0, \infty)$ ,  $\lambda_1, \lambda_2 \sim \mathcal{G}(1, 50)$ ,  $p \sim \mathcal{U}(0, 1)$ .

It is enough to simulate only jump times and sizes in JBEA to derive the full conditional distributions of the parameters, which all have closed forms. Note however that, although this is a fairly simple model, exact inference is only feasible due to the JBEA algorithm.

We start the chain in values  $b^2 = 0.002^2$ ,  $\lambda_1 = 0.40$ ,  $\lambda_2 = 0.28$ ,  $\lambda_3 = 0.20$ ,  $\lambda_4 = 0.07$ ,  $\lambda_5 = 0.22$ ,  $p = 0.5$ , and run  $50k$  iterations. Standard diagnostics suggest that convergence is rapidly attained. Table 6 shows the posterior statistics of the parameters.

Table 6: Posterior statistics of the parameters for the last  $40k$  iterations.

	$b$	$\lambda_1$	$\lambda_2$	$\lambda_3$	$\lambda_4$	$\lambda_5$	$p$
Mean	0.0025	0.488	0.323	0.259	0.059	0.291	0.492
Median	0.0025	0.487	0.323	0.257	0.053	0.286	0.492
Std Dev	0.0001	0.040	0.030	0.048	0.037	0.071	0.027

## 6.2 S&P500

We now apply the methodology from Section 3.1.2 to fit the Pareto-Beta Jump-Diffusion (PBJD) to daily data from the S&P500 index. The PBJD model was proposed in Ramezani and Zeng (1998) to model stock price behavior. It allows for up and down jumps using a mixture jump size distribution to account for good and bad news. The model is the following:

$$\begin{aligned} dV_s &= \mu V_s ds + \sigma V_s dW_s + dJ_s, \quad V_0 = v \\ \lambda_1(s; V_{s-}) &= \lambda \end{aligned} \quad (44)$$

$$\begin{aligned} g_1(Z_j, V_{t_j-}) &= (Z_j - 1)V_{t_j-} \\ Z_j &\sim p\text{Pareto}(\eta_u) + (1 - p)\text{Beta}(\eta_d, 1), \end{aligned} \quad (45)$$

where the Pareto distribution takes values in  $(1, \infty)$ .

We consider daily data from 03/Jan/2000 to 31/Dec/2013 which consists of 3532 observations. This period incorporates the WTC 09/11 episode in 2001 and the 2008 economic crisis. The data is shown in Figure 2. The two periods mentioned are clear in the graph and it seems reasonable to assume  $\mu = 0$ , which considerably simplifies the algorithm. We apply the MCEM algorithm from Section 3.1. Initial values are chosen through an empirical analysis of the data. Results are presented in Table 7.

The estimated covariance matrix of the MLE returned a negative variance for parameter  $p$ . In order to resolve this we use a different parametrisation to compute this estimate. We make  $\lambda_u = p\lambda$  and  $\lambda_d = (1 - p)\lambda$  - the rates of up and down jumps, respectively.

Table 7: Maximum likelihood estimate and asymptotic 95% confidence interval for the model parameters.

	$\sigma$	$\lambda$	$p$	$\lambda_u$	$\lambda_d$	$\eta_u$	$\eta_d$
MLE	0.00577	0.841	0.532	0.447	0.394	144.5	125.2
C.I. 95%	(0.00497 , 0.00656)	-	-	(0.342 , 0.552)	(0.291 , 0.497)	(124.8 , 164.2)	(104.9 , 145.5)




Figure 2: Left: USD $\times$ GBP log-rate, source: <http://www.Investing.com>. Right: S&P500 index, source: <http://research.stlouisfed.org>

In this paper, we looked at the challenging problem of likelihood-based statistical inference for discretely observed jump-diffusion processes. Our main contribution is to provide for the first time, a comprehensive suite of methodologies for both frequentist and Bayesian *exact* inference for discretely observed jump-diffusion processes.

Our simplest methods are based on the JBEA algorithm for jump-diffusion bridge simulation and are presented in Section 3. However more general methods are required for the case of unbounded drift and jump-rate for the (transformed) process. To this end, we develop a more general approach in Section 4 which involve the construction of novel importance sampling and Barker MCMC algorithms. Both these approaches are useful: the JBEA approach is far more computationally efficient, while the methods presented Section 4 are more generally applicable.

We also address some important identifiability issues related to the inference problem and some practical implementation strategies. In particular, we propose (see supplementary material) an algorithm to simulate upper and lower bounds for a Brownian bridge and to simulated the bridge given these bounds and possibly other bridge points. We also concluded that prior distributions play a crucial role to solve identifiability problems. The Barker MCMC introduced in Section 4.2 seems also of independent interest.

The methods were empirically tested in some simulated examples and performed robustly and efficiently. The examples also addressed key identifiability issues. Finally, jump-diffusion models were used to fit some financial data.

Although beyond the scope of this paper, there is no intrinsic reason why this approach cannot be developed for many multi-dimensional contexts, at least for relatively small-dimensional models. However note that the exact simulation methodology that is used here, is fundamentally limited to

reducible, gradient drift diffusions. Moreover the computational overheads associated with moderately high-dimensional problems are likely to be prohibitive.

Although we give the first general efficient methodology for exact likelihood-based inference for discretely observed jump-diffusions in this paper, we also acknowledge the restrictions and complexity involved in the methodology, which reflects the complexity of the inference problem. Most importantly though, we hope our work will stimulate further work on our ambitious aim to solve the problem exactly.

## Acknowledgements

Flávio Gonçalves would like to thank FAPEMIG, CNPq and the University of Warwick for financial support. Krzysztof Łatuszyński is supported by the Royal Society through the Royal Society University Research Fellowship. Gareth Roberts is supported by the EPSRC grants: *ilike* (EP/K014463/1), CoSInES (EP/R034710/1) and Bayes for Health (EP/R018561/1).

## References

- Aït-Sahalia, Y. and Yu, J. (2006). Saddlepoint approximations for continuous-time Markov processes. *Journal of Econometrics*, 134:507–551.
- Ball, C. A. and Roma, A. (1993). A jump diffusion model for the European monetary system. *Journal of International Money and Finance*, 12:475–492.
- Barker, A. A. (1965). Monte Carlo calculations of the radial distribution functions for a proton-electron plasma. *Australian Journal of Physics*, 18:119–133.
- Barndorff-Nielsen, O. E. and Shephard, N. (2004). Power and bipower variation with stochastic volatility and jumps (with discussion). *Journal of Financial Econometrics*, 2:1–48.
- Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2006a). Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098.
- Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2008). A new factorisation of diffusion measure and sample path reconstruction. *Methodology and Computing in Applied Probability*, 10(1):85–104.
- Beskos, A., Papaspiliopoulos, O., and Roberts, G. O. (2009). Monte carlo maximum likelihood estimation for discretely observed diffusion processes. *The Annals of Statistics*, 37:223–245.
- Beskos, A., Papaspiliopoulos, O., Roberts, G. O., and Fearnhead, P. (2006b). Exact and computationally efficient likelihood-based inference for discretely observed diffusion processes (with discussion). *Journal of the Royal Statistical Society. Series B*, 68(3):333–382.
- Casella, B. and Roberts, G. O. (2010). Exact simulation of jump-diffusion processes with Monte Carlo applications. *Methodology and Computing in Applied Probability*, 12(3).
- Chen, N. (2009). Localization and exact simulation of Brownian motion driven stochastic differential equations. Working paper, Chinese University of Hong Kong.
- Chudley, C. T. and Elliott, R. J. (1961). Neutron scattering from a liquid on a jump diffusion model. *Proceedings of the Physical Society*, 77(2):353–361.
- Cont, R. and Tankov, P. (2004). *Financial modelling with jump processes*. Chapman & Hall/CRC Financial Mathematics Series. Chapman & Hall/CRC, Boca Raton, FL.

- Detemple, J., Garcia, R., and Rindisbacher, M. (2006). Asymptotic properties of Monte Carlo estimators of diffusion processes. *Journal of Econometrics*, 134:1–68.
- Duffie, D. and Glynn, P. (2004). Estimation of continuous-time Markov processes sampled at random time intervals. *Econometrica*, 72:1773–1808.
- Duffie, D., Pan, J., and Singleton, K. (2000). Transform analysis and asset pricing for affine jump-diffusions. *Econometrica*, 68(6):1343–1376.
- Duffie, D. and Singleton, K. J. (1993). Simulated moments estimation of Markov models of asset prices. *Econometrica*, 61:929–952.
- Durham, G. B. and Gallant, R. A. (2002). Numerical techniques for maximum likelihood estimation of continuous-time diffusion processes. *Journal of Business and Economic Statistics*, 20:279–316.
- Elerian, O. (1999). *Simulation estimation of continuous time series models with applications to finance*. PhD thesis, Nuffield College, Oxford.
- Eraker, B. (2004). Do stock prices and volatility jump? reconciling evidence from spot and option prices. *The Journal of Finance*, 59(3):1367–1404.
- Eraker, B., Johannes, M. S., and Polson, N. G. (2003). The impact of jumps in volatility and returns. *The Journal of Finance*, 58(3):1269–1300.
- Fearnhead, P., Papaspiliopoulos, O., and Roberts, G. O. (2008). Particle filters for partially observed diffusions. *Journal of the Royal Statistical Society. Series B*, 70(4):755–777.
- Feng, L. and Linetsky, V. (2008). Pricing options in jump-diffusion models: An extrapolation approach. *Operations Research*, 56(2):304–325.
- Filipović, D., Mayerhofer, E., and Schneider, P. (2013). Density approximations for multivariate affine jump-diffusion processes. *Journal of Econometrics*, 176:93–111.
- Fletcher, R. (1987). *Practical Methods of Optimization*. John Wiley & Sons, New York, 2nd edition.
- Fort, G. and Moulines, E. (2003). Convergence of the Monte Carlo expectation maximization for curved exponential families. *The Annals of Statistics*, 31(4):1220–1259.
- Giesecke, K. and Schwenkler, G. (2019). Simulated likelihood estimators for discretely observed jump-diffusions. *Journal of Econometrics*, 213(2):297–320.
- Golightly, A. (2009). Bayesian filtering for jump-diffusions with application to stochastic volatility. *J. Comput. Graph. Statist.*, 18(2):384–400.
- Gonçalves, F. B. and Franklin, P. (2019). On the definition of likelihood function. *arXiv:1906.10733*.
- Gonçalves, F. B. and Roberts, G. O. (2014). Exact simulation problems for jump-diffusions. *Methodology and Computing in Applied Probability*, 16:907–930.
- Grenander, U. and Miller, M. I. (1994). Representations of knowledge in complex systems. *Journal of the Royal Statistical Society. Series B*, 56(4):549–603.
- Johannes, M. (2004). The statistical and economic role of jumps in continuous-time interest rate models. *The Journal of Finance*, 50(1):227–260.
- Johannes, M. S., Polson, N. G., and Stroud, J. R. (2002). Nonlinear filtering of stochastic differential equations with jumps. Available at SSRN: <http://ssrn.com/abstract=334601> or doi:10.2139/ssrn.334601.

- Johannes, M. S., Polson, N. G., and Stroud, J. R. (2009). Optimal filtering of jump diffusions: Extracting latent states from asset prices. *Review of Financial Studies*, 22(7):2759–2799.
- Kennedy, J. S., Forsyth, P. A., and Vetzal, K. R. (2009). Dynamic hedging under jump diffusion with transaction costs. *Operations Research*, 57(3):541–559.
- Łatuszyński, K. G. and Roberts, G. O. (2013). CLTs and asymptotic variance of time sampled Markov chains. *Methodology and Computing in Applied Probability*, 15:237–247.
- Lo, A. W. (1988). Maximum likelihood estimation of generalized Itô processes with discretely sampled data. *Econometric Theory*, 4(2):231–247.
- Meng, X. Rubin, D. B. (1993). Maximum likelihood estimation via the ECM algorithm: A general framework. *Biometrika*, 80:267–278.
- Oakes, D. (1999). Direct calculation of the information matrix via the EM algorithm. *Journal of the Royal Statistical Society. Series B*, 61:479–482.
- Pedersen, A. (1995). A new approach to maximum likelihood estimation for stochastic differential equations based on discrete observations. *Scandinavian Journal of Statistics*, 22(1):55–71.
- Peskun, P. H. (1973). Optimum Monte Carlo sampling using Markov chains. *Biometrika*, 60:607–612.
- Platen, E. and Bruti-Liberati, N. (2010). *Numerical Solution of Stochastic Differential Equations with Jumps in Finance*. Springer, Berlin.
- Pollock, M., Johansen, A. M., and Roberts, G. O. (2016). On the exact and  $\epsilon$ -strong simulation of (jump) diffusions. *Bernoulli*, 22:794–856.
- Protter, P. E. (2004). *Stochastic integrations and differential equations*. Springer, 2nd edition.
- Ramezani, C. and Zeng, Y. (1998). Maximum likelihood estimation os asymmetric jump-diffusion processes: Aplication to security prices. Working Paper, Department of Mathematics and Statistics, University of Missouri.
- Roberts, G. O. and Stramer, O. (2001). On inference for partially observed nonlinear diffusion models using the Metropolis-Hastings algorithm. *Biometrika*, 88:603–621.
- Runggaldier, W. J. (2003). *Handbook of heavy tailed distributions in finance*, chapter Jump-Diffusion Models, pages 169–209. Handbooks in Finance, Book 1. Elsevier/North-Holland.
- Sermaidis, G., Papaspiliopoulos, O., Roberts, G. O., Beskos, A., and Fearnhead, P. (2013). Markov chain Monte Carlo for exact inference for diffusions. *Scandinavian Journal of Statistics*, 40:294–321.
- Srivastava, A., Grenander, U., Jensen, G. R., and Miller, M. I. (2002). Jump-diffusion markov processes on orthogonal groups for object recognition. *Journal of Statistical Planning and Inference*, 103(1-2):15–37.

## Appendix A - assumptions of JBEA

The following assumptions are required in order to simulate jump-diffusion bridges exactly using JBEA (presented in Section 2).

- (a)  $\alpha(\cdot)$  is continuously differentiable;

- (b)  $(\alpha^2 + \alpha')(\cdot)$  is bounded below;
- (c)  $\tilde{\mathbb{P}} \ll \tilde{\mathbb{D}}$  and the Radon-Nikodym derivative of  $\tilde{\mathbb{P}}$  w.r.t.  $\tilde{\mathbb{D}}$  is given by (47);
- (d)  $\lambda(s, u)$  is uniformly bounded for all  $u$  in the state space of  $V$  and all  $s \in [0, t]$ ;
- (e) the probability measure induced by the density  $f_g(\cdot; j)$  is absolutely continuous w.r.t. the probability measure induced by the density  $f_0(\cdot)$ , for all  $j$ ;
- (f)  $\frac{f_g(u; j)}{f_0(u)}$  is uniformly bounded for all  $u \in E_g$  (support of  $f_g$ ) and all  $j$ ;
- (g)  $\lambda_0$  is chosen such that  $\frac{\lambda(j)f_g(u; j)}{\lambda_0 f_0(u)} \leq 1$ ,  $\forall u \in E_g$  and  $\forall j$ ;
- (h)  $A(u) := \int_0^u \alpha(y) dy$  is Lipschitz ( $\alpha(\cdot)$  is uniformly bounded).
- (i)  $f_{\tilde{\mathbb{P}}}$  is integrable and we can simulate from it.

## Appendix B - Important results

**Lemma 4.** Consider any two fixed points  $x_{i-1}(\theta)$  and  $x_i(\theta)$ . Let  $U_1$  and  $U_2$  be the  $\mathcal{U}(0, 1)$  r.v.'s which are used in JBEA to simulate events of probabilities  $p_1(B)$  and  $p_2(B)$ , respectively, and let  $\kappa_i \sim Po\{r_i(\theta)\Delta t_i\}$  be the number of points from the Poisson process  $\Phi_i = \{\Psi, \Upsilon\}$ . Let also  $\lambda_i(\theta)$  and  $f_i(\cdot; \theta)$  be the jump rate and jump size density in  $[t_{i-1}, t_i]$ , under  $\tilde{\mathbb{D}}$ . Then, the conditional density of  $(\mathbf{J}_i, X_{J_i}, \dot{X}_i, \Phi_i, U_1, U_2 | \theta, x_{i-1}(\theta), x_i(\theta))$  with respect to the measure  $\mathbb{J}_{0,i} + \otimes \mathbb{L}^{N_i+1} \otimes \mathbb{W}_{0,i} \otimes \Phi_i^+ \otimes \mathbb{L}^2$ , (where  $(\mathbb{J}_{0,i}, \mathbb{W}_{0,i})$  is  $(\mathbb{J}_0, \mathbb{W}_0)$  in  $(t_{i-1}, t_i)$ ,  $\mathbb{L}^n$  is the  $(n+N)$ -dimensional Lebesgue measure,  $\mathbb{J}_0$  is the law of a jump process with unit jump rate and a parameter-free state-independent jump size distribution that preserves the required absolute continuity,  $\mathbb{W}_0$  is the collection of standard Brownian bridges between the merged observation and jump times and  $\Phi^+$  is the measure of a unit rate Poisson process on  $[t_{i-1}, t_i] \times [0, 1]$ ) is

$$\begin{aligned}
& \pi \left( \mathbf{J}_i, X_{J_i}, \dot{X}_i, \Phi_i | \theta, x_{i-1}(\theta), x_i(\theta) \right) = \\
& \exp \left\{ -(\Delta t_i) r_i(\theta) \right\} [r_i(\theta)]^{\kappa_i} f_{\tilde{\mathbb{P}}}(N_i, J_{i_1}, \dots, J_{i_{N_i}}) \pi_{\mathbb{Q}}(X_{J_i} | \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta); \theta) \\
& \mathbb{1} \left[ U_1 \leq \exp \left\{ -\frac{1}{2\Delta t_i} (\Delta x_i(\theta) - J_i)^2 \right\} \right] \\
& \mathbb{1} \left[ U_2 \leq \prod_{j=1}^{N_i} \frac{\lambda(i_j; \theta) f_g(J_{i_j}; i_j, \theta)}{\lambda_i(\theta) f_i(J_{i_j}; \theta)} \exp \left\{ -\sum_{j=1}^{N_i} (\Delta A(i_j)) + \kappa_0(J_{i_j}; \theta) \right\} \right] \\
& \prod_{l=1}^{\kappa_i} \mathbb{1} \left[ \frac{1}{r_i(\theta)} \phi \left( \varphi(\dot{X}_{\psi_l}) \right) < v_l \right] \frac{1}{a(x_{i-1}(\theta), x_i(\theta); \theta)} \kappa_1, \tag{46}
\end{aligned}$$

where  $\kappa_1$  is a constant that does not depend on  $\theta$ . Furthermore,

$$f_{\tilde{\mathbb{P}}}(N_i, J_{i_1}, \dots, J_{i_{N_i}}) = \frac{1}{c_i(\theta)} f_{\tilde{\mathbb{D}}}(N_i, J_{i_1}, \dots, J_{i_{N_i}}) \exp \left\{ \sum_{j=1}^{N_i} \kappa_0(J_{i_j}; \theta) \right\},$$

$\kappa_0(J_{i_j}; \theta) = c_0(\theta) |J_{i_j}|$ , if using the Lipschitz proposal (see Section 2),

$\kappa_0(J_{i_j}; \theta) = c_2(\theta)$ , if using the Vanilla proposal (see Gonçalves and Roberts, 2014),

$a(x_{i-1}(\theta), x_i(\theta); \theta)$  is the acceptance probability of JBEA and  $\pi_{\mathbb{Q}}(X_{J_i} | \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta); \theta)$  is the density of  $(X_{J_i} | \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta); \theta)$  under  $\mathbb{Q}$ .

**Proof.** The proof of this lemma is analogous to the proof of Lemma 3 from Beskos et al. (2006b).



**Lemma 5.** Let  $\mathbb{D}$  be as defined in Section 2 having jump rate and jump size density given by  $\lambda_i$  and  $f_i$ , respectively, and let  $p_\theta$  be the transition density under  $\mathbb{P}$ . Then

$$\frac{d\tilde{\mathbb{P}}}{d\tilde{\mathbb{D}}}(\mathbf{J}_i, X_{J_i}, \dot{X}_i; \theta) = \mathcal{G}_i \frac{f_N(\Delta x_i - J_i; 0, \Delta t_i)}{p_\theta(x_{i-1}(\theta), x_i(\theta); \Delta t_i)}, \quad (47)$$

and

$$\frac{d\tilde{\mathbb{P}}}{d\tilde{\mathbb{F}}}(\mathbf{J}_i, X_{J_i}, \dot{X}_i; \theta) = \frac{d\tilde{\mathbb{P}}}{d\tilde{\mathbb{D}}}(\mathbf{J}_i, X_{J_i}, \dot{X}_i; \theta) \frac{f_{\tilde{\mathbb{D}}}}{f_{\tilde{\mathbb{F}}}}(N_i, J_{i_1}, \dots, J_{i_{N_i}}), \quad (48)$$

where

$$\begin{aligned} \mathcal{G}_i &= \exp \left\{ A(x_i(\theta); \theta) - A(x_{i-1}(\theta); \theta) - \int_{t_{i-1}}^{t_i} \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, \dot{X}_s; \theta) ds + \Delta t_i \lambda_i(\theta) \right\} \\ &\times \exp \left\{ - \sum_{j=1}^{N_i} (\Delta A(i_j; \theta)) \right\} \prod_{j=1}^{N_i} \frac{\lambda(j; \theta) f_g(J_{i_j}; j, \theta)}{\lambda_i(\theta) f_i(J_{i_j}; \theta)}. \end{aligned} \quad (49)$$

**Proof.** See Lemma 2 and equation (22) from Gonçalves and Roberts (2014).

## Appendix C - proofs

### Proof of Lemma 1

Define  $s_\theta(a, b; \Delta t_i)$  as the transition density of  $V$  going from  $a$  to  $b$  in a time interval  $[t_{i-1}, t_i]$ , and  $p_\theta(a, b; \Delta t_i)$  as the same density for  $X$ . We shall find the density  $\pi(V_{obs}, X_J, J, \dot{X})$  of the law of  $V_{com}$  with respect to the product measure  $\mathbb{L}^n \otimes \mathbb{L}^N \otimes \mathbb{J}_0 \otimes \mathbb{W}_0$ . We have

$$\pi(V_{obs}, V_{mis} | \theta) = \left[ \prod_{i=1}^n s_\theta(V_{t_{i-1}}, V_{t_i}; \Delta t_i) \right] \pi(V_{mis} | V_{obs}, \theta).$$

Furthermore,

$$s_\theta(V_{t_{i-1}}, V_{t_i}; \Delta t_i) = p_\theta(x_{t_{i-1}}(\theta), x_{t_i}(\theta); \Delta t_i) |\eta'(V_{t_i}; \theta)|,$$

$$\begin{aligned} \pi(V_{mis} | V_{obs}, \theta) &= \prod_{i=1}^n \pi(X_{J_i}, \mathbf{J}_i, \dot{X}_i | x_{i-1}(\theta), x_i(\theta), \theta) \\ &= \prod_{i=1}^n \frac{d\mathbb{P}}{d(\mathbb{L}^{N_i} \otimes \mathbb{J}_0 \otimes \mathbb{W}_0)}(X_{J_i}, \mathbf{J}_i, \dot{X}_i | x_{i-1}(\theta), x_i(\theta), \theta) \\ &= \frac{d\mathbb{P}}{d\mathbb{W}_0}(\dot{X}_i | X_{J_i}, \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta), \theta) \frac{d\mathbb{P}}{d(\mathbb{L}^{N_i} \otimes \mathbb{J}_0)}(X_{J_i}, \mathbf{J}_i | x_{i-1}(\theta), x_i(\theta), \theta). \end{aligned} \quad (50)$$

Now note that

$$\frac{d\mathbb{P}}{d\mathbb{W}_0}(\dot{X}_i | X_{J_i}, \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta), \theta) = \frac{d\mathbb{P}}{d\mathbb{Q}}(\varphi(\dot{X}_i) | X_{J_i}, \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta), \theta), \quad (51)$$

where  $\mathbb{Q}$  is the measure of a jump-diffusion starting in  $x_{i-1}(\theta)$  which is the sum of a BM and a jump process with measure  $\mathbb{J}_0$ ,

$$\begin{aligned} \frac{d\mathbb{P}}{d(\mathbb{L}^{N_i} \otimes \mathbb{J}_0)}(X_{J_i}, \mathbf{J}_i | x_{i-1}(\theta), x_i(\theta), \theta) &= \frac{d\mathbb{P}}{d\mathbb{Q}}(X_{J_i}, \mathbf{J}_i | x_{i-1}(\theta), x_i(\theta), \theta) \\ &= \frac{d\mathbb{Q}}{d(\mathbb{L}^{N_i} \otimes \mathbb{J}_0)}(X_{J_i}, \mathbf{J}_i | x_{i-1}(\theta), x_i(\theta), \theta), \end{aligned} \quad (52)$$

$$\frac{d\mathbb{P}}{d\mathbb{Q}} \left( \varphi(\dot{X}_i) | X_{J_i}, \mathbf{J}_i, \cdot \right) \frac{d\mathbb{P}}{d\mathbb{Q}} (X_{J_i}, \mathbf{J}_i | \cdot) = \frac{d\mathbb{P}}{d\mathbb{Q}} \left( X_{J_i}, \mathbf{J}_i, \varphi(\dot{X}_i) | \cdot \right), \quad (53)$$

$$\begin{aligned} \frac{d\mathbb{Q}}{d(\mathbb{L}^{N_i} \otimes \mathbb{J}_0)} (X_{J_i}, \mathbf{J}_i | x_{i-1}(\theta), x_i(\theta), \theta) &= \frac{d\mathbb{Q}}{d\mathbb{J}_0} (\mathbf{J}_i | x_{i-1}(\theta), x_i(\theta), \theta) \frac{d\mathbb{Q}}{d\mathbb{L}^{N_i}} (X_{J_i} | \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta), \theta) \\ &= \frac{d\mathbb{Q}}{d\mathbb{L}^{N_i}} (X_{J_i} | \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta), \theta) = \pi_{\mathbb{Q}} (X_{J_i} | \mathbf{J}_i; \theta). \end{aligned} \quad (54)$$

Putting (50), (51), (52), (53) and (54) together and using the result in (47) and (49) leads to

$$\pi(V_{obs}, V_{mis} | \theta) = \prod_{i=1}^n \mathcal{G}_i f_N(\Delta x_i(\theta) - J_i; 0, \Delta t_i) |\eta'(V_{t_i}; \theta)| \pi_{\mathbb{Q}} (X_{J_i} | \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta)). \quad (55)$$

Finally, note that

$f_N(\Delta x_i - J_i; 0, \Delta t_i) \pi_{\mathbb{Q}} (X_{J_i} | \mathbf{J}_i, x_{i-1}(\theta), x_i(\theta)) \propto \pi_{\mathbb{Q}} (X_{J_i}, x_i(\theta) | \mathbf{J}_i, x_{i-1}(\theta))$ , which establishes the result of the Lemma. □

## Proof of Theorem 1

We use the following decomposition:

$$\pi(\theta, \mathbf{J}_i, X_{J_i}, \dot{X}_i, \Phi_i | \mathbf{v}) = \pi(\theta | \mathbf{v}) \pi(\mathbf{J}_i, X_{J_i}, \dot{X}_i, \Phi_i | \theta, \mathbf{v}) = \pi(\theta | \mathbf{v}) \prod_{i=1}^n \pi(\mathbf{J}_i, X_{J_i}, \dot{X}_i, \Phi_i | \theta, \mathbf{v}), \quad (56)$$

where the second equality is obtained by the Markov property. Furthermore,

$$\pi(\theta | \mathbf{v}) \propto \pi(\theta) \prod_{i=1}^n p_{\theta}(x_{i-1}(\theta), x_i(\theta); \Delta t_i) |\eta'(V_{t_i}; \theta)|. \quad (57)$$

The next step is to find an equality for the transition density  $p_{\theta}(x_{i-1}(\theta), x_i(\theta); \Delta t_i)$ . Taking the expectation with respect to  $\tilde{\mathbb{F}}$  on both sides of (48), we get

$$p_{\theta}(x_{i-1}(\theta), x_i(\theta); \Delta t_i) = \mathbb{E}_{\tilde{\mathbb{F}}} \left[ \mathcal{G} f_N(\Delta x_i(\theta) - J_i; 0, \Delta t_i) \frac{f_{\tilde{\mathbb{D}}}}{f_{\tilde{\mathbb{F}}}}(N_i, J_{i_1}, \dots, J_{i_{N_i}}) \right]. \quad (58)$$

Since

$$\begin{aligned} a(x_{i-1}(\theta), x_i(\theta); \theta) &= \mathbb{E}_{\tilde{\mathbb{F}}} \left[ \exp \left\{ - \int_{t_{i-1}}^{t_i} \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, \varphi(\dot{X}_s); \theta) - m_i(\theta) ds \right. \right. \\ &\quad \left. \left. - \sum_{j=1}^{N_i} (\Delta A(i_j; \theta) + \kappa_0(J_{i_j}; \theta)) \right\} \prod_{j=1}^{N_i} \frac{\lambda(j; \theta) f_g(J_{i_j}; j, \theta)}{\lambda_i(\theta) f_i(J_{i_j}; \theta)} \exp \left\{ \frac{-1}{2\Delta t_i} (\Delta x_i(\theta) - J_i)^2 \right\} \right], \end{aligned} \quad (59)$$

we have that

$$\begin{aligned} p_{\theta}(x_{i-1}(\theta), x_i(\theta); \Delta t_i) &\propto \exp \{ A(x_i(\theta); \theta) - A(x_{i-1}(\theta); \theta) - \Delta t_i (m_i(\theta) - \lambda_i(\theta)) \} \\ &\quad c_i(\theta) a(x_{i-1}(\theta), x_i(\theta); \theta). \end{aligned} \quad (60)$$

The result is obtained by replacing (60) in (57) and then (46) and (57) in (56), and integrating  $U_1$ ,  $U_2$  and the  $v_l$ 's out. □

## Proof of Proposition 1

We have that

$$\begin{aligned}
\mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [l(\theta)] &= \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} \left[ \sum_{i=1}^n l_i(\theta) \right] = \sum_{i=1}^n \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [l_i(\theta)] \\
&= \sum_{i=1}^n \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [w^* l_i(\theta)] = \sum_{i=1}^n \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} \left[ \prod_{l=1}^n w_l^* l_i(\theta) \right] \\
&= \sum_{i=1}^n \prod_{l \neq i} \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [w_l] \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [w_i^* l_i(\theta)] = \sum_{i=1}^n \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [w_i l_i(\theta)],
\end{aligned}$$

where row two to three is justified by the Markov property and row three by the fact that  $\mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [w_i] = 1, \forall i$ .

□

## Proof of Proposition 2

Firstly, note that  $Var[E_1] = \frac{1}{M} Var \left[ \sum_{i=1}^n \left( \prod_{k=1}^n w_k \right) l_i(\theta) \right]$  and  $Var[E_2] = \frac{1}{M} Var \left[ \sum_{i=1}^n w_i l_i(\theta) \right]$ . Moreover,  $w_i \perp w_j, \forall i \neq j$  and  $\mathbb{E}[w_i] = 1$ .

Define  $A_i = \prod_{k \neq i} w_k$  and  $A_{ij} = \prod_{k \neq i, j} w_k$ . We have

$$\begin{aligned}
M \cdot Var[E_1] &= Var \left[ \sum_{i=1}^n A_i w_i l_i \right] = \mathbb{E} \left[ \left( \sum_{i=1}^n A_i w_i l_i \right)^2 \right] - \left( \mathbb{E} \left[ \sum_{i=1}^n A_i w_i l_i \right] \right)^2 \\
&= \sum_{i=1}^n \mathbb{E}[A_i^2] \mathbb{E}[(w_i l_i)^2] + 2 \sum_{i < j} \mathbb{E}[A_{ij}] \mathbb{E}[w_i l_i] \mathbb{E}[w_j l_j] - \left( \sum_{i=1}^n \mathbb{E}[A_i w_i l_i] \right)^2 \\
&\geq \sum_{i=1}^n \mathbb{E}[(w_i l_i)^2] + 2 \sum_{i < j} \mathbb{E}[w_i l_i] \mathbb{E}[w_j l_j] - \left( \sum_{i=1}^n \mathbb{E}[w_i l_i] \right)^2 \\
&= \mathbb{E} \left[ \left( \sum_{i=1}^n w_i l_i \right)^2 \right] - \left( \mathbb{E} \left[ \sum_{i=1}^n w_i l_i \right] \right)^2 = M \cdot Var[E_2]
\end{aligned} \tag{61}$$

## Proof of Lemma 3

To prove that the estimator is strongly consistent, note that

$$\begin{aligned}
Y_M^{(i1)} &= \frac{1}{M} \sum_{j=1}^M \left( \dot{w}_i^{(j)} \dot{l}_i^{(j)}(\theta) \right) \rightarrow \mathbb{E}_{X_{mis}^{\mathbb{F}}|x_{obs},\theta'} [l_i(\theta)] := a_i, \quad \text{a.s. as } M \rightarrow \infty, \text{ by SLLN,} \\
Y_M^{(i2)} &= \sum_{j=1}^M \left( \dot{w}_i^{(j)} \right) \rightarrow 1 := b_i, \quad \text{a.s. as } M \rightarrow \infty, \text{ by SLLN.}
\end{aligned}$$

Since  $Y_M^{(i1)}/Y_M^{(i2)}$  is a a.s. continuous function of  $(Y_M^{(i1)}, Y_M^{(i2)})$ , the result is established by the Convergence of Transformations Theorem.

To prove the finite variance part, we write  $\tilde{l}_i$  instead of  $\tilde{l}_i(\theta)$  and note that  $\tilde{w}_i^{(j)} < 1$  *a.s.* and

$$\begin{aligned} \text{Var}[E_3] &= \text{Var} \left[ \sum_{i=1}^n \sum_{j=1}^M \left( \tilde{w}_i^{(j)} \tilde{l}_i^{(j)} \right) \right] = \sum_{i=1}^n \text{Var} \left[ \sum_{j=1}^M \left( \tilde{w}_i^{(j)} \tilde{l}_i^{(j)} \right) \right] \\ &= \sum_{i=1}^n \left( \sum_{j=1}^M \text{Var} \left[ \tilde{w}_i^{(j)} \tilde{l}_i^{(j)} \right] + 2 \sum_{1 \leq j_1 < j_2 \leq M} \text{Cov} \left( \tilde{w}_i^{(j_1)} \tilde{l}_i^{(j_1)}, \tilde{w}_i^{(j_2)} \tilde{l}_i^{(j_2)} \right) \right). \end{aligned}$$

Therefore,

$$\text{Var}[E_2] < \infty \Leftrightarrow \text{Var} \left[ \tilde{w}_i^{(j)} \tilde{l}_i^{(j)} \right] < \infty \text{ and } \text{Cov} \left( \tilde{w}_i^{(j_1)} \tilde{l}_i^{(j_1)}, \tilde{w}_i^{(j_2)} \tilde{l}_i^{(j_2)} \right) < \infty, \forall i, j, j_1, j_2.$$

Finally,

$$\begin{aligned} \mathbb{E} \left[ \left( \tilde{w}_i^{(j)} \tilde{l}_i^{(j)} \right)^2 \right] &\leq \mathbb{E} \left[ \left( \tilde{l}_i^{(j)} \right)^2 \right], \forall i, j, \\ \text{Cov} \left( \tilde{w}_i^{(j_1)} \tilde{l}_i^{(j_1)}, \tilde{w}_i^{(j_2)} \tilde{l}_i^{(j_2)} \right) &= \mathbb{E} \left[ \tilde{w}_i^{(j_1)} \tilde{w}_i^{(j_2)} \tilde{l}_i^{(j_1)} \tilde{l}_i^{(j_2)} \right] - \mathbb{E} \left[ \tilde{w}_i^{(j_1)} \tilde{l}_i^{(j_1)} \right] \mathbb{E} \left[ \tilde{w}_i^{(j_2)} \tilde{l}_i^{(j_2)} \right] \\ &\leq \mathbb{E} \left[ \left| \tilde{l}_i^{(j_1)} \tilde{l}_i^{(j_2)} \right| \right] + \mathbb{E} \left[ \left| \tilde{l}_i^{(j_1)} \right| \right] \mathbb{E} \left[ \left| \tilde{l}_i^{(j_2)} \right| \right] = 2 \left( \mathbb{E} \left[ \left| \tilde{l}_i^{(j)} \right| \right] \right)^2 \leq 2 \mathbb{E} \left[ \left( \tilde{l}_i^{(j)} \right)^2 \right] \forall i, j, j_1, j_2. \end{aligned}$$

□

## Appendix D - Estimator for the covariance matrix of the MLE

The covariance matrix of the MLE is approximated by the inverse of the observed information matrix  $I$ , that is

$$I^{-1} = \left[ -\frac{\partial^2}{\partial \theta^2} l(X_{obs}; \theta) \right]^{-1}, \quad (62)$$

given that the required regularity conditions are satisfied. Now Oakes' identity (see Oakes, 1999) can be rewritten to give

$$\begin{aligned} \frac{\partial^2}{\partial \theta^2} l(X_{obs}; \theta) &= \mathbb{E}_{X_{mis}^{\tilde{\mathbb{P}}}|x_{obs}, \theta'} \left[ \frac{\partial^2}{\partial \theta^2} l(X_{com}; \theta) \right] + \mathbb{E}_{X_{mis}^{\tilde{\mathbb{P}}}|x_{obs}, \theta'} \left[ \left( \frac{\partial}{\partial \theta} l(X_{com}; \theta) \right)^2 \right] \\ &\quad - \left[ \mathbb{E}_{X_{mis}^{\tilde{\mathbb{P}}}|x_{obs}, \theta'} \left( \frac{\partial}{\partial \theta} l(X_{com}; \theta) \right) \right]^2. \end{aligned}$$

We can use the general result from Proposition 1 to obtain

$$\frac{\partial^2}{\partial \theta^2} l(X_{obs}; \theta) \approx (M_1 + M_2 + M_3 - M_4) \Big|_{\theta=\hat{\theta}}, \quad (63)$$

where  $\hat{\theta}$  is the MLE of  $\theta$  and each  $M$  is a matrix with the  $[r, c]$ -th entry given by:

$$\begin{aligned} M_1[r, c] &= \sum_{i=1}^n \sum_{j=1}^M \tilde{w}_i^{(j)} \frac{\partial^2}{\partial \theta_r \partial \theta_c} \tilde{l}_i^{(j)}; \quad M_2[r, c] = \sum_{i=1}^n \mathbb{E}_{X_{mis}^{\tilde{\mathbb{P}}}|x_{obs}, \theta'} \left[ \frac{\partial}{\partial \theta_r} \tilde{l}_i \frac{\partial}{\partial \theta_c} \tilde{l}_i \right]; \\ M_3[r, c] &= \sum_{i_1=1}^n \sum_{i_2 \neq i_1} \left( \sum_{j=1}^M \tilde{w}_i^{(j)} \frac{\partial}{\partial \theta_r} \tilde{l}_i^{(j)} \right) \left( \sum_{j=1}^M \tilde{w}_i^{(j)} \frac{\partial}{\partial \theta_c} \tilde{l}_i^{(j)} \right); \\ M_4[r, c] &= \left( \sum_{i=1}^n \sum_{j=1}^M \tilde{w}_i^{(j)} \frac{\partial}{\partial \theta_r} \tilde{l}_i^{(j)} \right) \left( \sum_{i=1}^n \sum_{j=1}^M \tilde{w}_i^{(j)} \frac{\partial}{\partial \theta_c} \tilde{l}_i^{(j)} \right). \end{aligned} \quad (64)$$

To obtain the equations above we use the Leibniz Integral Rule which allows us to exchange the partial derivative w.r.t.  $\theta$  and the integral in  $ds$  - that is because  $X(s)$  is *càdlàg*. To deal with  $M_2$ , note that double integrals will emerge from the product and we use the following identity to obtain the MC estimator:

$$\int_{t_{i-1}}^{t_i} \int_{t_{i-1}}^{t_i} f_1^*(X(s_1), \theta) f_2^*(X(s_2), \theta) ds_1 ds_2 = (\Delta t_i)^2 \mathbb{E}_{\dot{U}_{i1}, \dot{U}_{i2}} \left[ f_1^*(X(\dot{U}_{i1}), \theta) f_2^*(X(\dot{U}_{i2}), \theta) \right], \quad (65)$$

where  $\dot{U}_{i1}$  and  $\dot{U}_{i2}$  are independent and uniformly distributed in  $(t_{i-1}, t_i)$ .

Finally, the same algorithm may be used to estimate the covariance matrix of the MLE when using the algorithms from Section 3.1. In that case, we have  $\tilde{w}_i^{(j)} = 1/M$ , which leads to further simplifications in the formule.

## Appendix E - The layered Brownian bridge

The MCEM and MCMC algorithms presented in Section 4 require the simulation of lower and upper bounds for a collection of Brownian bridges. This can be done by simulating the layered Brownian bridge which basically samples layers that contains the supremum and infimum of the bridges. The algorithm to construct and simulate the layered Brownian bridge (layers and bridge points given the layers) was introduced in Beskos et al. (2008).

Equally important, the algorithms from Section 4.2 also require the simulation of extra points of the bridge given the layers and other points previously simulated. This is not a trivial task and no explicit algorithm to do so is presented in the original paper. For that reason we present here an overview of the layered Brownian bridge algorithm from Beskos et al. (2008) and devise an algorithm to perform the simulation of the extra points.

In fact, we restrict ourselves to the simulation of standard bridges (starting and ending in 0) as this is necessary when there are unknown parameters in the diffusion coefficient and also offers a more efficient (tighter bounds) solution in any case. We also restrict ourselves to the symmetric layer case as this is much simpler and there is no real advantage in using asymmetric layers.

Suppose, without loss of generality, a standard Brownian bridge  $W$  in  $[0, t]$ . Let  $\{b_i\}_{i \geq 1}$  be an increasing sequence of positive real numbers with  $b_0 = 0$  and define the following events:

$$\begin{aligned} U_i &= \left\{ \sup_{0 \leq s \leq t} W_s \in [b_{i-1}, b_i] \right\} \cap \left\{ \inf_{0 \leq s \leq t} W_s > -b_i \right\}, \\ L_i &= \left\{ \inf_{0 \leq s \leq t} W_s \in [-b_i, -b_{i-1}] \right\} \cap \left\{ \sup_{0 \leq s \leq t} W_s < b_i \right\}, \\ D_i &= U_i \cup L_i, \quad i \geq 1. \end{aligned}$$

Define the random variable  $I = I(W)$  such that  $\{I = i\} = D_i$  and note that  $\{I = i\}$  implies that  $\{-b_i < W_s < +b_i, \forall s \in [0, t]\}$ . Beskos et al. (2008) show that

$$F(i) := P(I \leq i) = \gamma(T - t_0; 0, 0, b_i), \quad (66)$$

for  $i \geq 1$ , where

$$\begin{aligned} \gamma(s, u_1, u_2, K) &= 1 - \sum_{j=1}^{\infty} \{ \sigma_j(s, u_1, u_2, K) - \tau_j(s, u_1, u_2, K) \}, \\ \sigma_j(s, u_1, u_2, K) &= \bar{\sigma}_j(s, u_1, u_2, K) + \bar{\sigma}_j(s, -u_1, -u_2, K), \\ \tau_j(s, u_1, u_2, K) &= \bar{\tau}_j(s, u_1, u_2, K) + \bar{\tau}_j(s, -u_1, -u_2, K), \end{aligned}$$

$$\bar{\sigma}_j(s, u_1, u_2, K) = \exp \left\{ -\frac{2}{s} [2Kj - (K + u_1)][2Kj - (K + u_2)] \right\},$$

$$\bar{\tau}_j(s, u_1, u_2, K) = \exp \left\{ -\frac{2j}{s} [4K^2j + 2K(u_1 - u_2)] \right\}.$$

We sample  $I$  by using the inverse cdf method and, although we cannot evaluate  $F(i)$  exactly, we can find bounds for it and apply the alternating series method as follows.

Define  $S_{2j}^{(i)} = 1 - \sum_{l=1}^{j-1} (\sigma_l - \tau_l) - \sigma_j$  and  $S_{2j-1}^{(i)} = 1 - \sum_{l=1}^j (\sigma_l - \tau_l)$ , for  $j \geq 1$ , with  $\sigma$  and  $\tau$  defined for  $I = i$ . It can be shown that

$$0 < S_2^{(i)} < S_4^{(i)} < S_6^{(i)} < \dots < F(i) < \dots < S_5^{(i)} < S_3^{(i)} < S_1^{(i)}, \quad (67)$$

with  $S_{2j}^{(i)} \uparrow F(i)$  and  $S_{2j-1}^{(i)} \downarrow F(i)$ . Defining  $S_j^{(0)} = 0, \forall j \geq 1$ , if  $u \sim U(0, 1)$ , we have the following result:

$$I = i, \text{ if } S_{2j-1}^{(i-1)} < u < S_{2j}^{(i)}, \text{ for some } j \geq 1. \quad (68)$$

This way, we can use the following algorithm to simulate  $I$ :

#### Algorithm to simulate $I$

- (a) Simulate  $u \sim U(0, 1)$ ;
- (b) make  $i = 1$ ;
- (c) make  $j = 1$ ;
- (d) compute  $S_{2j-1}^{(i)}$  and  $S_{2j}^{(i)}$ ;
  - (a) if  $u < S_{2j}^{(i)}$ , output  $I = i$ ;
  - (b) if  $u > S_{2j-1}^{(i)}$ , make  $i = i + 1, j = 1$  and GOTO 4;
  - (c) else, make  $j = j + 1$  and GOTO 4.

Now that we have the algorithm to sample the layers, we shall move on to the algorithm to simulate from  $W_s | I = i$ . In the MCMC algorithm from Section 4.2, this simulation will be required at different stages of the algorithm and may fall into one of the two cases: *i*) no points from the bridge have yet been simulated; *ii*) some points of the bridge have already been simulated; which in turn fall into two more cases: *a*)  $I = 1$ ; *b*)  $I \geq 2$ . A nice solution for the case *ia* (and implicitly for case *iiia*) is presented in Beskos et al. (2008). The authors also present an algorithm for *ib*, but not for *iib*. Case *iib* is more complex and requires some work to devise an efficient solution. We propose the following general strategy, which shall solve the problem in any of the four cases described above.

The first step after  $I$  is simulated is to simulate the extreme (minimum or maximum) that reaches the most external layer - note that  $I = i$  implies that either the minimum is in  $(-b_i, -b_{i-1})$  or the maximum is in  $(b_{i-1}, b_i)$ . This is done via rejection sampling with proposal

$$\mathbb{P}_{D_I} := \frac{1}{2} BB_{\overline{M}_I} + \frac{1}{2} BB_{\underline{M}_I}, \quad (69)$$

where  $BB_{\overline{M}_I}$  is the Brownian bridge  $W$  conditional on  $\overline{M}_I := \max_{0 \leq s \leq t} \{W_s\} \in [b_{I-1}, b_I]$  and  $BB_{\underline{M}_I}$  is  $W$  conditional on  $\underline{M}_I := \min_{0 \leq s \leq t} \{W_s\} \in [-b_I, -b_{I-1}]$ .

To simulate from this proposal, we first choose between  $BB_{\overline{M}_I}$  and  $BB_{\underline{M}_I}$  with probability 1/2 each. If we choose the first (second) one, we have to simulate the maximum (minimum) of  $W$  given that it is in the given interval.

Let  $m$  be the minimum and  $M$  be the maximum of  $W$  in  $[0, t]$ . In order to simulate the minimum conditional on being in an interval  $[m_1, m_2]$ , we need to compute the distribution function  $F_m$  of this minimum, which is:

$$F_m(w) = \exp\left\{\frac{-2w^2}{t}\right\}. \quad (70)$$

Let  $U_1 \sim U[F(m_1), F(m_2)]$  and  $E = -\log(U_1)$ , then

$$m = \frac{\left(-\sqrt{2tE}\right)}{2}. \quad (71)$$

In order to simulate the last time instant  $t_m$  where this minimum is attached, set  $c_1 = \frac{m^2}{2t}$  and let  $U_2 \sim U(0, 1)$ ,  $I_1 \sim IGau(1, 2c_1)$  and  $I_2 \sim 1/IGau(1, 2c_1)$  and define  $V = \mathbb{I}[U_2 < 1/2] \cdot I_1 + \mathbb{I}[U_2 \geq 1/2] \cdot I_2$ . Finally,

$$t_m = \frac{t}{(1 + V)}. \quad (72)$$

To simulate the maximum  $M$  conditional on being in an interval  $[M_1, M_2]$ , we use the symmetry of Brownian bridge by simulating a minimum  $m_0$ , conditional on being in the interval  $[-M_2, -M_1]$ , and its location  $t_{m_0}$  and making

$$M = -m_0 \text{ and } t_M = t - t_{m_0}. \quad (73)$$

Let  $\mathbb{B}_{D_I}$  be the measure of the layered Brownian bridge, we accept the proposal w.p.

$$\frac{d\mathbb{B}_{D_I}}{d\mathbb{P}_{D_I}}(W) \propto \frac{\mathbb{I}[D_I]}{1 + \mathbb{I}[U_I \cap L_I]} \leq 1. \quad (74)$$

In order to compute this probability we need to simulate the two Bernoulli r.v.'s in (74). The first one (in the numerator) is equal to 1 w.p.

$$\delta(t_m, 0, -m, b_I - m) \times \delta(t - t_m, 0, -m, b_I - m) \text{ or} \quad (75)$$

$$\delta(t_M, 0, M, M + b_I) \times \delta(t - t_M, 0, M, M + b_I). \quad (76)$$

and the second one (in the denominator) is 1 w.p. 1 if  $I = 1$  and is 0 w.p.

$$\delta(t_m, 0, -m, b_{I-1} - m; b_I - m) \times \delta(t - t_m, 0, -m, b_{I-1} - m; b_I - m) \text{ or} \quad (77)$$

$$\delta(t_M, 0, M, M + b_{I-1}; M + b_I) \times \delta(t - t_M, 0, M, M + b_{I-1}; M + b_I), \quad (78)$$

if  $I > 1$ , where

$$\delta(s, 0, u_2, K) = 1 - \frac{1}{u_2} \sum_{j=1}^{\infty} \{\zeta_j(s, u_2, K) - \xi_j(s, u_2, K)\} \quad (79)$$

$$\delta(s, 0, u_2, K; L) = \frac{u_2 - \sum_{j=1}^{\infty} \{\zeta_j(s, u_2, K) - \xi_j(s, u_2, K)\}}{u_2 - \sum_{j=1}^{\infty} \{\zeta_j(s, u_2, L) - \xi_j(s, u_2, L)\}} \quad (80)$$

$$\xi_j(s, u_2, K) = (2Kj + u_2) \exp\{-2Kj(Kj + u_2)/s\} \quad (81)$$

$$\zeta_j(s, u_2, K) = \xi_j(s, -u_2, K). \quad (82)$$

The alternating series method can be used for the events involving the functions  $\delta$  if  $3K^2 > s$ . Otherwise, it can be used after the first  $\lceil \sqrt{t + K^2}/(2K) \rceil$  pairs  $S_{2j-1}$  and  $S_{2j}$  have been computed.

This follows from the result in Pollock et al. (2016) (Corollary 5) that states that the sequence of bounds for the probabilities in (75) and (77) are Cauchy sequences after the first  $\lceil \sqrt{t + K^2}/(2K) \rceil$  pairs are computed.

Finally note that the acceptance probability is 0 if the Bernoulli r.v. in the numerator is 0. If this r.v. is 1, the acceptance probability is 1 if the Bernoulli r.v. in the denominator is 0 and 1/2 if it is 1.

The algorithm above outputs the following variables under the measure  $\mathbb{B}\mathbb{B}_{D_I}: (m^*, t^*) = (m, t_m)$  or  $(M, t_M)$ ;  $\mathbb{I}[U_I \cap L_I]$ . We choose to keep only  $(m^*, t^*)$  which is an exact draw from  $\pi_{\mathbb{B}\mathbb{B}_{D_I}}(m^*, t^*)$  - the density of  $(m^*, t^*)$  under  $\mathbb{B}\mathbb{B}_{D_I}$ .

We now proceed to the simulation of the bridge points given the information we have -  $\mathcal{I}_{m^*, I}$ : either the minimum and its location and an upper bound for  $W$  or the maximum and its location and a lower bound for  $W$ . This is done via rejection sampling by proposing from  $\mathbb{B}\mathbb{B}|(m^*, t^*)$  and accepting the proposal if  $\mathbb{I}[D_I] = 1$ .

The law  $\mathbb{B}\mathbb{B}|(m^*, t^*)$  is a three-dimensional Bessel bridge and is constructed as follows. Firstly, denote by  $R(\delta) = \{R_s(\delta); 0 \leq s \leq 1\}$  a three-dimensional Bessel bridge of unit length from 0 to  $\delta \leq 0$ , i.e.

$$R_s(\delta) = \sqrt{(\delta s + W_s^{(1)})^2 + (W_s^{(2)})^2 + (W_s^{(3)})^2}, \quad s \in [0, 1], \quad (83)$$

where the  $W_s^{(\cdot)}$ 's are three independent standard Brownian bridges in  $[0, 1]$ . Then, defining  $W^*$  to be  $W|(m, t_m)$ , we have that

$$\{W_s^*; 0 \leq s \leq t_m\} = \sqrt{t_m} R_{(t_m-s)/(t_m)}(\delta_1) + m, \quad (84)$$

$$\{W_s^*; t_m \leq s \leq t\} = \sqrt{t - t_m} R_{(s-t_m)/(t-t_m)}(\delta_2) + m, \quad (85)$$

where  $\delta_1 = \frac{-m}{\sqrt{t_m}}$  and  $\delta_2 = \frac{-m}{\sqrt{t - t_m}}$ .

If we now define  $W^*$  to be  $W|(M, t_M)$ , we have that

$$\{W_s^*; 0 \leq s \leq t_M\} = M - \sqrt{t_M} R_{(t_M-s)/(t_M)}(\delta_1), \quad (86)$$

$$\{W_s^*; t_M \leq s \leq t\} = M - \sqrt{t - t_M} R_{(s-t_M)/(t-t_M)}(\delta_2), \quad (87)$$

where  $\delta_1 = \frac{M}{\sqrt{t_M}}$  and  $\delta_2 = \frac{M}{\sqrt{t - t_M}}$ .

This construction implies that  $\mathbb{B}\mathbb{B}|\mathcal{I}_{m^*, I}$  is a Markov process and, therefore, the proposed rejection sampling algorithm can be performed interval-wise w.r.t. the sub-intervals defined by the bridge points already simulated. Furthermore, in order to simulate the proposal, all the points from the three standard Brownian bridges  $W_s^{(\cdot)}$  need to be stored along the way.

Finally, once the proposal is simulated in a given sub-interval, the acceptance indicator is simulated via the alternating series method. Suppose that  $\mathbb{B}\mathbb{B}|\mathcal{I}_{m^*, I}$  is to be simulated at locations  $(s_1, \dots, s_n)$  in a sub-interval  $(s_0, s_{n+1})$ . Note that  $s_0$  and  $s_{n+1}$  may be one of the extremes of  $(0, t)$  or even  $t^*$ . We have that

$$P(D_I) = \prod_{k=1}^{n+1} \gamma \left( s_k - s_{k-1}; |m^* - W_{s_{k-1}}^*|, |m^* - W_{s_k}^*|, |m^*| + b_I \right), \quad (88)$$

where

$$\delta(s, u_1, u_2, K) = \frac{\gamma(s, u_1 - K/2, u_2 - K/2, K/2)}{1 - \exp\{-2u_1 u_2 / s\}}.$$

The choice of the sequence  $b_1, b_2, \dots$ , specially  $b_1$ , is of great concern. On one hand, if  $b_1$  is a relatively high value, we will get  $I = 1$  more often, which improves the computational time for sampling from  $W$ . On the other hand, a high value of  $b_1$  leads to conservative bounds for the function  $\phi$  in the two steps of the Barker's MCMC which in turn leads to an inefficient two-coin algorithm. For an interval of length  $t$ , we suggest  $\sqrt{t}/2 \leq b_1 \leq \sqrt{t}$ .



## Appendix F - Obtaining efficient lower bounds $a(s; \theta)$

Let  $\dot{X}_{ij}$  be the standard BB in  $s \in [t_{i_{j-1}}, t_{ij}]$  (simulated from  $\tilde{\mathbb{D}}$ ) and define  $\dot{l}_{ij}$  and  $\dot{u}_{ij}$  as the lower and upper bounds for  $\dot{X}_{ij}$ , obtained from the layer simulation. Now define  $x_{ij} := x_{ij}(\theta) = \min\{X_{i_{j-1}}, X_{i_j-}\}$ ,  $y_{ij} := y_{ij}(\theta) = \max\{X_{i_{j-1}}, X_{i_j-}\}$ ,

$$l_{ij}(s; \theta) = \begin{cases} \frac{(y_{ij}-x_{ij})}{\Delta t_{ij}}(s - t_{i_{j-1}}) + (x_{ij} + \dot{l}_{ij}), & \text{if } x_{ij} = X_{i_{j-1}} \\ -\frac{(y_{ij}-x_{ij})}{\Delta t_{ij}}(s - t_{ij}) + (x_{ij} + \dot{l}_{ij}), & \text{if } x_{ij} = X_{i_j}, \end{cases} \quad (89)$$

and

$$u_{ij}(s; \theta) = \begin{cases} \frac{(y_{ij}-x_{ij})}{\Delta t_{ij}}(s - t_{i_{j-1}}) + (x_{ij} + \dot{u}_{ij}), & \text{if } x_{ij} = X_{i_{j-1}} \\ -\frac{(y_{ij}-x_{ij})}{\Delta t_{ij}}(s - t_{ij}) + (x_{ij} + \dot{u}_{ij}), & \text{if } x_{ij} = X_{i_j}, \end{cases} \quad (90)$$

where  $\Delta t_{ij} = t_{ij} - t_{i_{j-1}}$ . This implies that  $X_s(\theta) = \varphi(\dot{X}_s; \theta) \in [l_{ij}(s; \theta), u_{ij}(s; \theta)] = H_{ij}(s; \theta)$ ,  $\forall s \in [t_{i_{j-1}}, t_{ij}]$ .

Now define  $\phi(\dot{X}_s, s; \theta) = \left( \frac{\alpha^2 + \alpha'}{2} + \lambda \right) (s, \varphi(\dot{X}_s; \theta); \theta)$  and

$$l_{\phi_{ij}}(s; \theta) = \inf_{\varphi(u; \theta) \in H_{ij}(s)} \phi(u, s; \theta) \text{ and } L_{ij}(\theta) = \inf_{s \in [t_{i_{j-1}}, t_{ij}]} l_{\phi_{ij}}(s; \theta), \quad (91)$$

$$u_{\phi_{ij}}(s; \theta) = \sup_{\varphi(u; \theta) \in H_{ij}(s)} \phi(u, s; \theta) \text{ and } U_{ij}(\theta) = \sup_{s \in [t_{i_{j-1}}, t_{ij}]} u_{\phi_{ij}}(s; \theta), \quad (92)$$

If we can compute  $\int_{t_{i_{j-1}}}^{t_{ij}} l_{\phi_{ij}}(\dot{X}_s, s; \theta) ds$ , we make  $a(s; \theta) = l_{\phi_{ij}}(s; \theta)$ , for  $s \in [t_{i_{j-1}}, t_{ij}]$ , otherwise  $a(s; \theta) = L_{ij}(\theta)$ . Finally, we define  $r_{ij}(\theta) = U_{ij}(\theta) - L_{ij}(\theta)$ .