## Short Distance Modification of the Quantum Virial Theorem

Qin Zhao<sup>1</sup>, Mir Faizal<sup>2,3</sup>, Zaid Zaz<sup>4,5</sup>

Department of Physics, National University of Singapore,
 2 Science Drive 3, Singapore
 Irving K. Barber School of Arts and Sciences,
 University of British Columbia,
 Kelowna, British Columbia, V1V 1V7, Canada.
 Department of Physics and Astronomy,
 University of Lethbridge,
 Lethbridge, Alberta, T1K 3M4, Canada.
 Theoritical Physics Division, Department of Physics,
 National Institute of Technology,
 Srinagar, Kashmir-190006, India

<sup>5</sup> Department of Electronics and Communication Engineering, University of Kashmir, Srinagar, Kashmir-190006, India

## Abstract

In this letter, we will analyse the deformation of a semi-classical gravitational system from minimal measurable length scale. In the semi-classical approximation, the gravitational field will be analysed as a classical field, and the matter fields will be treated quantum mechanically. Thus, using this approximation, this system will be represented by a deformation of Schrödinger-Newton equation by the generalised uncertainty principle (GUP). We will analyse the effects of this GUP deformed Schrödinger-Newton equation on the behaviour of such a semi-classical gravitational system. As the quantum mechanical virial theorem can be obtained using the Schrödinger-Newton equation, a short distance modification of the Schrödinger-Newton equation will also result in a short distance modification of the quantum mechanical virial theorem.

Even though there have been various attempts to formulate a quantum theory of gravity, there are serious problems with obtaining a fully quantum theory of gravity. This has motivated the study of semi-classical quantum gravity, and in such approaches the gravitational field is treated classically and the matter fields are treated quantum mechanically. Thus, in this approximation, the gravitational field is a classical field, but it is sourced by quantum mechanical matter fields,  $G_{\mu\nu} = 8\pi G \langle \psi | \hat{T}_{\mu\nu} | \psi \rangle / c^2$ , where  $G_{\mu\nu}$  is the Einstein tensor for a classical space-time,  $\hat{T}_{\mu\nu}$  is the operator for the energy-momentum tensor, and  $|\psi\rangle$  is the wave-function of the matter fields. Furthermore, the Newtonian gravity has been observed to be the correct approximation to general relativity till the smallest length scale to which general relativity has been tested [1]. Thus, it is possible to use the Newtonian gravitational field in this approximation, and describe this system using a Schrödinger-Newton equation [2]-[4],

$$i\hbar \frac{\partial \psi}{\partial t} = H\psi = \frac{1}{2m}\hat{P}^2\psi + m\Phi(\mathbf{R}, t)\psi,$$
 (1)

where H is the Hamiltonian operator and  $\Phi(\mathbf{R},t)$  is the Newtonian potential, and the mass density of the system is  $m|\psi(\mathbf{R},t)|^2 = \rho(\mathbf{R},t)$ . It has been demonstrated that such a Schrödinger-Newton equation can be obtained from a non-relativistic limit of self-gravitating Klein-Gordon and Dirac fields [5]. The Schrödinger-Newton equation has been used to study the semi-classical behaviour of various interesting gravitational systems [6]-[10]. It has been proposed that such a Schrödinger-Newton equation can be tested experimentally using an optomechanical systems [11]-[12]. However, it has also been demonstrated that optomechanical systems can be used to test the deformation of a quantum mechanical system by the generalised uncertainty principle [13]. Thus, it becomes important to study the deformation of Schrödinger-Newton equation by generalised uncertainty principle (GUP), as this deformation can be tested experimentally using optomechanical systems. In fact, as the general relativity has not been tested below 0.4mm [1], it is possible that GUP can produce interesting short distance modification for a gravitational system. This modification can alter the behaviour of the system and can be detected using optomechanical systems. Thus, it is very interesting to analyse the GUP deformation of Schrödinger-Newton equation, and so such deformation of Schrödinger-Newton equation has been thoroughly studied [14]-[17].

Furthermore, there is another reason why such a the GUP deformation of Schrödinger-Newton equation would be interesting. The GUP deformation of the Schrödinger-Newton equation can be viewed as an intermediate step between the full quantum theory of gravity and semi-classical quan-

tum gravity. This is because a universal feature of almost all approaches to quantum gravity is the existence of a minimum measurable length scale. It is postulated that physical measurements below this scale are not possible. In string theory, which is one of the most popular approaches to quantum gravity, the smallest scale at which space-time can be probed is the string length scale. This is because strings are the smallest probes that exist in perturbative string theory, and so it is not possible to probe space-time below string length scale [18]-[22]. Even in loop quantum gravity, the existence of a minimum length has a very interesting consequence as it turns the big bang into a big bounce [23]. Moreover arguments from black hole physics also suggest that any theory of quantum gravity must be equipped with a minimum length scale [24]-[25]. This is due to the fact that the energy required to probe any region of space below the Plank length is greater than the energy required to create a mini black hole in that region of space. Thus, by incorporating an intrinsic minimal length scale in the Schrödinger-Newton equation, certain features of the quantum gravity can be analysed beyond the purely semi-classical approximation.

This minimal length can be incorporated into such a system by studding the GUP deforming of the Schrödinger-Newton equation [14]-[17]. The GUP deformation of quantum mechanical systems is motivated from the fact that usual Heisenberg uncertainty principle is not consistent with the existence of a minimum length scale in the system. Thus, the Heisenberg uncertainty principle has to be generalised to a generalised uncertainty principle (GUP) to incorporate the existence of such a minimum measurable length scale [26]-[31]. It is known that the uncertainty principle is related to the Heisenberg algebra, and so any modification of the uncertainty principle will deform the Heisenberg algebra [32]-[35]. The deformation of the Heisenberg algebra will in turn modify the representation of the momentum operator in a position basis [36]-[38]. These effects can be measured from the deformation of the Heisenberg algebra to [36]-[38]

$$[\hat{X}_i, \hat{P}_j] = i\hbar(\delta_{ij} + \beta \delta_{ij}\hat{P}^2 + 2\beta \hat{P}_i\hat{P}_j). \tag{2}$$

It can be demonstrated using the Jacobi identity that the other commutators for this system do not get deformed [39]-[41]

$$[\hat{X}_i, \hat{X}_j] = 0 = [\hat{P}_i, \hat{P}_j].$$
 (3)

It may be noted that this deformation occurs due to the existence of an extended structure (minimum length) in space-time, and hence it also deforms the standard Poisson brackets as  $\{x, p\} = 1 + \beta p^2$  [16]. This is the minimal

extension of the Heisenberg algebra, and to analyse it further, we define

$$\hat{X}_i \psi(\mathbf{x}) = \hat{x}_i \psi(\mathbf{x}) 
\hat{P}_i \psi(\mathbf{x}) = \hat{p}_i (1 + \beta \hat{p}^2) \psi(\mathbf{x}),$$
(4)

where  $\hat{x}_i$  and  $\hat{p}_j = -i\hbar \frac{\partial}{\partial \hat{x}_j}$  satisfy the canonical commutation relations  $[\hat{x}_i, \hat{p}_j] =$  $i\hbar\delta_{ij}$ . Then, it is easy to show that in the first order of  $\beta$ , (2) is satisfied by this deformed momentum operator (here we neglect terms of order  $\beta^2$ and higher). Physically, we interpret  $\hat{p}_i$  as the momentum in the IR limit and  $P_j$  as momentum in the UV limit of the theory. It is worth mentioning here that the minimum measurable length has to exist at least at the Planck scale. However, it can be argued that the minimum measurable length can exist at a much greater scale than the Planck scale. If the generalised uncertainty principle deforms all field theoretical systems in the same way, then the bound on the existence of the minimum measurable length scale is the electroweak scale [39]. Even this can have low energy consequences. In fact, such corrections have been obtained for various physical phenomena such as Lamb shift and Landau levels [40]. Now we can assume that there is a minimum measurable length scale in the background space-time, at a scale much larger than Planck scale. This will deform the Heisenberg uncertainty principle to GUP which will have low energy effects.

Now by substituting  $\hat{X}_i$ ,  $\hat{P}_i$  by  $x_i, p_i$ , we can rewrite the Schrödinger-Newton equation (1) as (to the leading order in  $\beta$ )

$$i\hbar\frac{\partial\psi}{\partial t} = H\psi = \frac{1}{2m}\hat{p}^2\psi + \frac{\beta}{m}\hat{p}^4\psi + m\Phi(\mathbf{r}, t)\psi. \tag{5}$$

We would like to point out that the GUP deformation of such a system has already been studied [14]-[17]. However, we shall use the GUP deformation of the quantum mechanical virial theorem for such a Schrödinger-Newton equation. The quantum mechanical virial theory has been recently obtained using a Schrödinger-Newton equation, and it has been demonstrated that such a quantum mechanical virial theory can have interesting cosmological consequences [42]. So, in this letter, we will analyse the GUP deformation of such a quantum mechanical virial theory using a GUP deformed Schrödinger-Newton equation. Now the moment of inertia for this system is

 $I = \frac{1}{2}m \int d\mathbf{r} r^2 |\psi|^2$ , and so we can write

$$\dot{I} = -\frac{im}{2\hbar} \int d\mathbf{r} r^2 (\psi^* H \psi - \psi H \psi^*) 
= -\frac{im}{2\hbar} \int d\mathbf{r} \psi^* [r^2, H] \psi 
= \frac{i\hbar}{4} \int d\mathbf{r} \psi^* [r^2, \nabla^2 - 2\beta \hbar^2 \nabla^4] \psi,$$
(6)

noting that  $[r^2, \Phi] = 0$ . Using the derivative along time direction, we can obtain

$$\ddot{I} = \frac{1}{4} \int d\mathbf{r} \psi^*[[r^2, \nabla^2 - 2\beta\hbar^2 \nabla^4], H] \psi. \tag{7}$$

Using this result, it can be demonstrated that the following result holds for this system,

$$[[r^{2}, \nabla^{2}], -\frac{\hbar^{2}}{2m} \nabla^{2} + m\Phi] = -4m(\mathbf{r} \cdot \nabla\Phi) - \frac{4\hbar^{2}}{m} \nabla^{2},$$
(8)  

$$[[r^{2}, \nabla^{2}], \frac{\hbar^{4}\beta}{m} \nabla^{4}] + [[r^{2}, -2\beta\hbar^{2}\nabla^{4}], -\frac{\hbar^{2}}{2m} \nabla^{2}]$$

$$= \frac{2\hbar^{4}\beta}{m} [[r^{2}, \nabla^{2}], \nabla^{4}]$$

$$= \frac{2\hbar^{4}\beta}{m} [r^{2}\nabla^{6} - \nabla^{2}r^{2}\nabla^{4} - \nabla^{4}r^{2}\nabla^{2} + \nabla^{6}r^{2}]$$

$$= \frac{2\hbar^{4}\beta}{m} * 16\nabla^{4},$$
(9)

where we have  $\nabla^2 r^2 \nabla^4 = 6\nabla^4 + 4\mathbf{r} \cdot \nabla \nabla^4 + r^2 \nabla^6$ , and  $\nabla^4 r^2 \nabla^2 = 20\nabla^4 + 8\mathbf{r} \cdot \nabla \nabla^4 + r^2 \nabla^6$ , along with  $\nabla^6 r^2 = 42\nabla^4 + 12\mathbf{r} \cdot \nabla \nabla^4 + r^2 \nabla^6$ . We also obtain

$$[[r^{2}, -2\beta\hbar^{2}\nabla^{4}], m\Phi] = 20\hbar^{2}\beta m(\nabla^{2}\Phi) + 8\hbar^{2}\beta m(\mathbf{r} \cdot \nabla\nabla^{2}\Phi) + 8\hbar^{2}\beta m(\nabla^{2}\Phi)\mathbf{r} \cdot \nabla + 8\hbar^{2}\beta m(\mathbf{r} \cdot \nabla\Phi)\nabla^{2} + 16\hbar^{2}\beta m(\mathbf{r} \cdot \nabla)\nabla\Phi \cdot \nabla.$$
(10)

Then, we can expand  $\ddot{I}$  to

$$\ddot{I} = -\int d\mathbf{r}\psi^{*}(m\psi(\mathbf{r}\cdot\nabla\Phi) + \frac{\hbar^{2}}{m}\nabla^{2}\psi) + \int d\mathbf{r}\psi^{*}\frac{8\hbar^{4}\beta}{m}\nabla^{4}\psi$$

$$+ \int d\mathbf{r}5\psi^{*}\hbar^{2}\beta m(\nabla^{2}\Phi)\psi + \int d\mathbf{r}2\hbar^{2}\beta m\psi^{*}(\mathbf{r}\cdot\nabla\nabla^{2}\Phi)\psi$$

$$+ \int d\mathbf{r}2\hbar^{2}\beta m\psi^{*}(\nabla^{2}\Phi)\mathbf{r}\cdot\nabla\psi + \int d\mathbf{r}2\hbar^{2}\beta m\psi^{*}(\mathbf{r}\cdot\nabla\Phi)\nabla^{2}\psi$$

$$+ \int d\mathbf{r}\psi^{*}4\hbar^{2}\beta m(\mathbf{r}\cdot\nabla)(\nabla\Phi\cdot\nabla\psi).$$
(11)

These results hold for any potential as we have not used the explicit form for the gravitational field. However, now we will use an explicit form for the gravitational field, and analyse this deformation of the Schrödinger-Newton equation from GUP. It may be noted that if the GUP deforms all field theoretical systems equally, then the experimental bound on the GUP has to be of the order of electroweak scale. However, there is no reason for the GUP to deform all field theoretical systems by the same amount, and thus we can assume that the value of the deformation parameter  $\beta$  varies between different field theoretical systems. In this letter, we will analyse the GUP deformation of a gravitational system, and assume that the generalised uncertainty acts differently for gravitational systems. As gravity has not been tested below 0.4mm [1], such effects can be detected in future optomechanical experiments. So, now we explicitly write the form for  $\Phi$ , and analyse the GUP deformation of the Schrödinger-Newton equation

$$\Phi = -G \int \frac{d\mathbf{r}' \rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}.$$
 (12)

Now we can use

$$\nabla \Phi = -G \int \frac{d\mathbf{r}' \rho(\mathbf{r}')(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3}, \qquad \nabla^2 \Phi = 0.$$
 (13)

Therefore, the terms containing  $\nabla^2 \Phi$  vanish<sup>1</sup>. Furthermore, it is also useful to take the wave-function in the form  $\psi = \sqrt{\rho/m} \exp(i\theta)$  with  $\theta$  real, and  $\mathbf{v} = \frac{\hbar \nabla \theta}{m}$ ,

$$\ddot{I} = -\int d\mathbf{r}\rho(\mathbf{r}\cdot\nabla\Phi) + \int d\mathbf{r}\rho\mathbf{v}^{2} + \frac{\hbar^{2}}{m^{2}}\int d\mathbf{r}|\nabla\sqrt{\rho}|^{2}$$

$$+ \int d\mathbf{r}\frac{8\hbar^{4}\beta}{m}(\nabla^{2}\sqrt{\frac{\rho}{m}} - \sqrt{\frac{\rho}{m}}\nabla\theta\cdot\nabla\theta)^{2}$$

$$+ \int d\mathbf{r}\frac{8\hbar^{4}\beta}{m}(\nabla\sqrt{\frac{\rho}{m}}\cdot\nabla\theta + \sqrt{\frac{\rho}{m}}\nabla^{2}\theta)^{2}$$

$$+ \int d\mathbf{r}2\hbar^{2}\beta m(\mathbf{r}\cdot\nabla\Phi)\psi^{*}\nabla^{2}\psi$$

$$+ \int d\mathbf{r}\psi^{*}4\hbar^{2}\beta m(\mathbf{r}\cdot\nabla)(\nabla\Phi\cdot\nabla\psi).$$
(14)

This is the quantum virial theorem for the gravitational field deformed by the generalised uncertainty principle. It may be noted here we have taken

Here, we assume that we do not consider the singularity around r = 0.

the parameter  $\beta$  bounded by short distance measurement of gravitational potential [1]. In the limit case, where  $\beta = 0$ , then we just obtain the quantum virial theorem [42]

$$\ddot{I}_{\beta=0} = -\int d\mathbf{r}\rho(\mathbf{r}\cdot\nabla\Phi) + \int d\mathbf{r}\rho v^2 + \frac{\hbar^2}{m^2} \int d\mathbf{r}|\nabla\sqrt{\rho}|^2$$

$$= W + 2K + 2Q,$$
(15)

where  $K = \frac{1}{2} \int d\mathbf{r} \rho \mathbf{v}^2$  is the kinetic energy, and Q is a new term and it is called as the quantum energy [42]. The potential of the system W is given by

$$W = -\int d\mathbf{r}\rho(\mathbf{r}\cdot\nabla\Phi)$$

$$= -G\int d\mathbf{r}d\mathbf{r}'\frac{\rho(\mathbf{r})\rho(\mathbf{r}')\mathbf{r}\cdot(\mathbf{r}-\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|^{3}}$$

$$= -\frac{G}{2}\int d\mathbf{r}\frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|}.$$
(16)

Then, in a steady state where  $\ddot{I} = 0$  and K >= 0, we have [42]

$$\frac{Q}{|W|} \le \frac{1}{2}.\tag{17}$$

It is easy to realise that when the phase of the wave-function is position-independent and then  $\nabla \theta = 0$ , one can obtain  $Q/|W| = \frac{1}{2}$ , which is the classical limit.

However, when we consider the contribution from the  $\beta$  terms, Q/|W| will receive corrections as follows,

$$\ddot{I}_{\nabla\theta=0} = W + 2Q + \int d\mathbf{r} \frac{8\hbar^2 \beta}{m} (\nabla^2 \sqrt{\frac{\rho}{m}})^2 
+ \int d\mathbf{r} 2\hbar^2 \beta m (\mathbf{r} \cdot \nabla \Phi) \sqrt{\frac{\rho}{m}} \nabla^2 \sqrt{\frac{\rho}{m}} 
+ \int d\mathbf{r} \psi^* 4\hbar^2 \beta m (\mathbf{r} \cdot \nabla) (\nabla \Phi \cdot \nabla \psi).$$
(18)

Since  $(\nabla^2 \sqrt{\frac{p}{m}})^2 \ge 0$ , it is also possible to obtain  $Q/|W| \le \frac{1}{2}$  for the position-independent wave-function. Thus, we have been able to study the effect of generalised uncertainty principle on quantum mechanical virial theorem. In this paper, we proposed that GUP deformation can be important for gravitational fields, if the GUP parameter varies between different field theories. As

the gravitational field has not been measured at very short distances, such a deformation can be detected in near future experiments. We have analysed the effect of such a deformation on quantum mechanical virial theorem [42], and demonstrated that this deformation will have interesting consequences for such a quantum mechanical virial theorem.

It may be noted various other forms of GUP have been proposed using a variety of physical motivations. A different deformation of the Heisenberg algebra has also been motivated from the doubly special relativity [43]-[45]. The doubly special relativity is a theory in which besides the speed of light, Planck energy is also a universal constant. Doubly special relativity has in turn been motivated from deformations of the energy-momentum dispersion relation, and such deformations occur in various approaches to quantum gravity, such as the discrete space-time [66], the spontaneous symmetry breaking of Lorentz invariance in string field theory [47], space-time foam models [48], spin-network in loop quantum gravity [49], non-commutative geometry [50], and Horava-Lifshitz gravity [51]-[53]. It is possible to combine the deformation of the Heisenberg algebra which is dependent on quadratic powers of momentum with the deformation of the Heisenberg algebra produced by the doubly special relativity [54]-[56]. The coordinate representation of the momentum operator for such a deformed Heisenberg algebra which is produced by the combination of both these deformations contains linear powers of the momentum operator. This approach results in non-local fractional derivative contributions in any dimension beyond the simple one dimensional case. However, these fractional derivative contributions can be analysed using the harmonic extension of function [57]-[58]. Another deformation of Heisenberg algebra containing inverse powers of momentum operator has also been proposed [59]. This deformation of the Heisenberg algebra is motivated by non-local quantum mechanics. It will be interesting to study the effects of these different forms of deformations of the Heisenberg algebra on small scale quantum systems with a gravitational potential.

It may be noted that the GUP deformation of quantum mechanics contains higher derivative terms, and such therms can produce Ostrogradsky ghost [60]-[61]. These ghost states can in turn make the theory non-unitary, and such theories can even contain negative norm states. However, it is possible to use different formalism to analyse such higher derivative states. It is possible to remove the negative norm states by tracing over a certain field configuration in the final state [62]-[63]. Thus, it is possible to get rid of the negative norm states from such a higher derivative theory, but the theory would still be non-unitary. It is possible to use the Lee-Wick boundary conditions and obtain a unitary theory [64]-[67]. Thus, it is possible to analyse this deformed system using an Lee-Wick field corresponding to a more mas-

sive mode. Then Lee-Wick boundary conditions can be imposed, and this could make the theory unitary. We would like to point out that such boundary conditions violate causality at microscopic scales, but such a violation does not occur at macroscopic scale [64]-[67]. The relation between Lee-Wick field theories, nonanticommutative deformation of supersymmetric and GUP deformation has already been studied [68]- [70]. Thus, it is interesting to analyse this system using Lee-Wick boundary conditions. It is interesting to note that negative norm states occur in usual gauge theories, but such negative states are removed using the Kugo-Ojima criterion which is based on the BRST symmetry [71]- [72]. Such a criterion has also been used to remove the negative norm states in higher derivative theories [73]-[75]. In fact, various superselection rules have been proposed to remove these ghost states from higher derivative theories [76]-[82]. Even though we do not address such questions in this paper, it is important to note that such higher derivative terms for GUP deformed systems can be consistently analysed. It would be interesting to analyse such terms using some of these approaches. It may be noted that usually such terms are analysed in the framework of effective field theory, and such higher derivative terms occur [83]-[87] in the derivative expansion of effective field theory. This is because these theories have a minimum measurable length scale associated with them, and so this is the scale that has to be integrated out in effective field theories [88]. However, it would be also interesting to analyse these higher derivative terms using other approaches.

## Acknowledgements

The work of Q.Z. is supported by NUS Tier 1 FRC Grant R-144-000-360-112.

## References

- [1] S. Q. Yang, B. F. Zhan, Q. L. Wang, C. G. Shao, L. C. Tu, W. H. Tan and J. Luo, Test of the Gravitational Inverse Square Law at Millimeter Ranges, Phys. Rev. Lett. **108**, 081101 (2012)
- [2] H. Yang, H. Miao, D. s. Lee, B. Helou and Y. Chen, "Macroscopic Quantum Mechanics in a classical spacetime," Phys. Rev. Lett. 110, no. 17, 170401 (2013)

- [3] I. M. Moroz, R. Penrose and P. Tod, "Spherically symmetric solutions of the Schrödinger-Newton equations," Class. Quant. Grav. 15, 2733 (1998)
- [4] J. R. van Meter, "Schrödinger-Newton 'collapse' of the wave function," Class. Quant. Grav. 28, 215013 (2011)
- [5] D. Giulini and A. Grossardt, "The Schrödinger-Newton equation as non-relativistic limit of self-gravitating Klein-Gordon and Dirac fields," Class. Quant. Grav. 29, 215010 (2012)
- [6] D. Giulini and A. Grossardt, "Gravitationally induced inhibitions of dispersion according to the Schrödinger-Newton Equation," Class. Quant. Grav. 28, 195026 (2011)
  - S. Bera, R. Mohan and T. P. Singh, "Stochastic modification of the Schrödinger-Newton equation," Phys. Rev. D **92**, no. 2, 025054 (2015)
- [7] A. Großardt, "Approximations for the free evolution of self-gravitating quantum particles," Phys. Rev. A 94, no. 2, 022101 (2016)
- [8] G. Manfredi, "The Schrödinger-Newton equations beyond Newton," Gen. Rel. Grav. 47, no. 2, 1 (2015)
- [9] D. Giulini and A. Grossardt, "Gravitationally induced inhibitions of dispersion according to a modified Schrödinger-Newton equation for a homogeneous-sphere potential," Class. Quant. Grav. 30, 155018 (2013)
- [10] G. Manfredi, P. A. Hervieux and F. Haas, "Lagrangian approach to the time-dependent Schrödinger-Newton equations," Class. Quant. Grav. 30, 075006 (2013)
- [11] C. C. Gan, C. M. Savage and S. Z. Scully, "Optomechanical tests of a Schrödinger-Newton equation for gravitational quantum mechanics," Phys. Rev. D 93, no. 12, 124049 (2016)
- [12] A. Großardt, J. Bateman, H. Ulbricht and A. Bassi, "Optomechanical test of the Schrödinger-Newton equation," Phys. Rev. D 93, no. 9, 096003 (2016)
- [13] I. Pikovski, M. R. Vanner, M. Aspelmeyer, M. S. Kim and C. Brukner, "Probing Planck-scale physics with quantum optics," Nature Phys. 8, 393 (2012)

- [14] S. Deb, S. Das and E. C. Vagenas, "Discreteness of Space from GUP in a Weak Gravitational Field," Phys. Lett. B 755, 17 (2016)
- [15] A. F. Ali and A. Tawfik, "Modified Newton's Law of Gravitation Due to Minimal Length in Quantum Gravity," Adv. High Energy Phys. 2013, 126528 (2013)
- [16] S. Benczik, L. N. Chang, D. Minic, N. Okamura, S. Rayyan and T. Takeuchi, "Short distance versus long distance physics: The Classical limit of the minimal length uncertainty relation," Phys. Rev. D 66, 026003 (2002)
- [17] P. Pedram, "A Higher Order GUP with Minimal Length Uncertainty and Maximal Momentum II: Applications," Phys. Lett. B 718, 638 (2012)
- [18] D. Amati, M. Ciafaloni and G. Veneziano, Can Space-Time Be Probed Below the String Size?, Phys. Lett. B 216, 41 (1989).
- [19] A. Kempf, G. Mangano and R. B. Mann, Hilbert space representation of the minimal length uncertainty relation, Phys. Rev. D **52**, 1108 (1995)
- [20] L. N. Chang, D. Minic, N. Okamura and T. Takeuchi, Exact solution of the harmonic oscillator in arbitrary dimensions with minimal length uncertainty relations, Phys. Rev. D 65, 125027 (2002)
- [21] L. N. Chang, D. Minic, N. Okamura and T. Takeuchi, The Effect of the minimal length uncertainty relation on the density of states and the cosmological constant problem, Phys. Rev. D 65, 125028 (2002)
- [22] S. Benczik, L. N. Chang, D. Minic, N. Okamura, S. Rayyan and T. Takeuchi, Short distance versus long distance physics: The Classical limit of the minimal length uncertainty relation, Phys. Rev. D 66, 026003 (2002)
- [23] P. Dzierzak, J. Jezierski, P. Malkiewicz and W. Piechocki, The minimum length problem of loop quantum cosmology, Acta Phys. Polon. B 41, 717 (2010)
- [24] M. Maggiore, A generalised uncertainty principle in quantum gravity, Phys. Lett. B **304**, 65 (1993)
- [25] M. I. Park, The generalised Uncertainty Principle in (A)dS Space and the Modification of Hawking Temperature from the Minimal Length, Phys. Lett. B 659, 698 (2008)

- [26] M. Maggiore, Quantum groups, gravity and the generalised uncertainty principle, Phys. Rev. D 49, 5182 (1994)
- [27] M. Maggiore, A generalised uncertainty principle in quantum gravity, Phys. Lett. B **304**, 65 (1993)
- [28] S. Park, B. H. Woo, M. Jung, E. J. Jang and W. S. Chung, Step potential problem and harmonic oscillator problem in the minimum length quantum mechanics, Mod. Phys. Lett. A **30**, no. 20, 1550096 (2015)
- [29] M. Maggiore, The Algebraic structure of the generalised uncertainty principle, Phys. Lett. B **319**, 83 (1993)
- [30] L. J. Garay, Quantum gravity and minimum length, Int. J. Mod. Phys. A 10, 145 (1995)
- [31] F. Scardigli, generalised uncertainty principle in quantum gravity from micro-black hole Gedanken experiment, Phys. Lett. B **452**, 39 (199
- [32] C. Bambi and F. R. Urban, Natural extension of the Generalised Uncertainty Principle, Class. Quant. Grav. 25, 095006 (2008)
- [33] D. Gao and M. Zhan, Constraining the generalised uncertainty principle with cold atoms, Phys. Rev. A **94**, no. 1, 013607 (2016)
- [34] F. Brau, Minimal Length Uncertainty Relation and Hydrogen Atom, J.Phys.A32:7691-7696,1999
- [35] K. Nozari, and B. Fazlpour, Chaos, Solitons and Fractals, 34, 224 (2007)
- [36] K. Nozari, Some aspects of Planck scale quantum optics, Phys. Lett. B 629, 41 (2005)
- [37] K. Nozari and T. Azizi, Some aspects of minimal length quantum mechanics, Gen. Rel. Grav. 38, 735 (2006)
- [38] P. Pedram, A class of GUP solutions in deformed quantum mechanics, Int. J. Mod. Phys. D 19, 2003 (2010)
- [39] S. Das and E. C. Vagenas, Universality of Quantum Gravity Corrections, Phys. Rev. Lett. **101**, 221301 (2008)
- [40] A. F. Ali, S. Das and E. C. Vagenas, A proposal for testing Quantum Gravity in the lab, Phys. Rev. D 84, 044013 (2011)

- [41] A. Kempf, Non-point-like Particles in Harmonic Oscillators, J. Phys. A **30**, 2093 (1997)
- [42] L. Hui, J. P. Ostriker, S. Tremaine and E. Witten, On the hypothesis that cosmological dark matter is composed of ultra-light bosons, arXiv:1610.08297 [astro-ph.CO]
- [43] J. Magueijo and L. Smolin, Lorentz invariance with an invariant energy scale, Phys. Rev. Lett. 88, 190403 (2002)
- [44] J. Magueijo and L. Smolin, String theories with deformed energy momentum relations, and a possible nontachyonic bosonic string, Phys. Rev. D 71, 026010 (2005)
- [45] J. L. Cortes and J. Gamboa, Quantum uncertainty in doubly special relativity, Phys. Rev. D 71, 065015 (2005)
- [46] G. 't Hooft, Quantization of point particles in (2+1)-dimensional gravity and space-time discreteness, Class. Quant. Grav. 13, 1023 (1996).
- [47] V. A. Kostelecky and S. Samuel, Spontaneous Breaking of Lorentz Symmetry in String Theory, Phys. Rev. D 39, 683 (1989)
- [48] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos and S. Sarkar, Tests of quantum gravity from observations of gamma-ray bursts, Nature **393**, 763 (1998).
- [49] R. Gambini and J. Pullin, Nonstandard optics from quantum spacetime, Phys.Rev. D59 (1999) 124021
- [50] S. M. Carroll, J. A. Harvey, V. A. Kostelecky, C. D. Lane and T. Okamoto, Noncommutative field theory and Lorentz violation, Phys. Rev. Lett. 87, 141601 (2001).
- [51] P. Horava, Quantum Gravity at a Lifshitz Point, Phys. Rev. D 79, 084008 (2009)
- [52] P. Horava, Spectral Dimension of the Universe in Quantum Gravity at a Lifshitz Point, Phys. Rev. Lett. **102**, 161301 (2009)
- [53] P. Horava, Membranes at Quantum Criticality, JHEP **0903**, 020 (2009)
- [54] A. F. Ali, S. Das and E. C. Vagenas, Discreteness of Space from the generalised Uncertainty Principle, Phys. Lett. B **678**, 497 (2009)

- [55] S. Das, E. C. Vagenas and A. F. Ali, Discreteness of Space from GUP II: Relativistic Wave Equations, Phys. Lett. B **690**, 407 (2010)
- [56] P. Pedram, On the Modification of Hamiltonians' Spectrum in Gravitational quantum Mechanics, Europhys. Lett. **89**, 50008 (2010)
- [57] M. Faizal, Consequences of Deformation of the Heisenberg Algebra, Int. J. Geom. Meth. Mod. Phys. 12, no. 02, 1550022 (2014)
- [58] S. Pramanik, M. Faizal, M. Moussa and A. F. Ali, Path integral quantization corresponding to the deformed Heisenberg algebra, Annals Phys. **362**, 24 (2015)
- [59] S. Masood, M. Faizal, Z. Zaz, A. F. Ali, J. Raza and M. B. Shah, The most general form of deformation of the Heisenberg algebra from the generalised uncertainty principle, Phys. Lett. B **763**, 218 (2016)
- [60] M. V. Ostrogradsky, "Memoire sur lesequations differentielles relatives aux problemes des isoperimetres" Mem. Acad. St. Petersbourg VI 4, 385 (1850)
- [61] R. P. Woodard, "Avoiding dark energy with 1/r modifications of gravity," Lect. Notes Phys. 720, 403 (2007)
- [62] S. W. Hawking and T. Hertog, "Living with ghosts," Phys. Rev. D 65, 103515 (2002)
- [63] M. Fontanini and M. Trodden, "Tackling Higher Derivative Ghosts with the Euclidean Path Integral," Phys. Rev. D 83, 103518 (2011)
- [64] T. D. Lee and G. C. Wick, "Negative Metric and the Unitarity of the S Matrix," Nucl. Phys. B 9, 209 (1969)
- [65] T. D. Lee and G. C. Wick, "Finite Theory of Quantum Electrodynamics," Phys. Rev. D 2, 1033 (1970)
- [66] T. D. Lee and G. C. Wick, "Questions of Lorentz Invariance in Field Theories With Indefinite Metric," Phys. Rev. D 3, 1046 (1971)
- [67] B. Grinstein, D. O'Connell and M. B. Wise, "The Lee-Wick standard model," Phys. Rev. D 77, 025012 (2008)
- [68] I. Cho and O. K. Kwon, "generalised Lee-Wick Formulation from Higher Derivative Field Theories," Phys. Rev. D 82, 025013 (2010)

- [69] M. Dias, A. F. Ferrari, C. A. Palechor and C. R. Senise, "A relation between deformed superspace and Lee–Wick higher-derivative theories," J. Phys. A 48, no. 27, 275403 (2015)
- [70] M. Faizal, "Supersymmetry breaking as a new source for the generalised uncertainty principle," Phys. Lett. B **757**, 244 (2016)
- [71] C. Becchi, A. Rouet and R. Stora, "Renormalization of Gauge Theories," Annals Phys. 98, 287 (1976)
- [72] I. V. Tyutin, "Gauge Invariance in Field Theory and Statistical Physics in Operator Formalism," arXiv:0812.0580 [hep-th]
- [73] V. O. Rivelles, "Triviality of higher derivative theories," Phys. Lett. B 577, 137 (2003)
- [74] K. Andrzejewski, J. Gonera, P. Machalski and K. Bolonek-Lasoń, "On the triviality of higher-derivative theories," Phys. Lett. B 706, 427 (2012)
- [75] M. Faizal and M. Khan, "A Superspace Formulation of The BV Action for Higher Derivative Theories," Eur. Phys. J. C 71, 1603 (2011)
- [76] M. Mintchev, "Quantization In Indefinite Metric," J. Phys. A 13, 1841 (1980)
- [77] E. S. Fradkin and A. A. Tseytlin, "Renormalizable asymptotically free quantum theory of gravity," Nucl. Phys. B **201**, 469 (1982)
- [78] N. H. Barth and S. M. Christensen, "Quantizing Fourth Order Gravity Theories. 1. The Functional Integral," Phys. Rev. D 28, 1876 (1983)
- [79] U. Moschella, "Infrared Singularities And Breaking Of The Poincaré Group. The Massless Dipole Field," J. Math. Phys. **31**, 2480 (1990)
- [80] M. Asorey, J. L. Lopez and I. L. Shapiro, "Some remarks on high derivative quantum gravity," Int. J. Mod. Phys. A 12, 5711 (1997)
- [81] T. C. Cheng, P. M. Ho and M. C. Yeh, "Perturbative approach to higher derivative and nonlocal theories," Nucl. Phys. B **625**, 151 (2002)
- [82] A. Accioly, A. Azeredo and H. Mukai, "Propagator, tree-level unitarity and effective nonrelativistic potential for higher-derivative gravity theories in D dimensions," J. Math. Phys. 43, 473 (2002)

- [83] C. P. Burgess, "Introduction to Effective Field Theory," Ann. Rev. Nucl. Part. Sci. 57, 329 (2007)
- [84] A. Awad and M. Faizal, "Higher Derivative Terms in Three Dimensional Supersymmetric Theories," JHEP **1510**, 002 (2015)
- [85] T. R. Morris, "Derivative expansion of the exact renormalization group," Phys. Lett. B **329**, 241 (1994)
- [86] L. Y. Hung, "String Creation, D-branes and Effective Field Theory," JHEP 0804, 057 (2008)
- [87] T. Banks, "TASI Lectures on Holographic Space-Time, SUSY and Gravitational Effective Field Theory," arXiv:1007.4001 [hep-th].
- [88] M. Faizal, A. F. Ali and A. Nassar, generalised Uncertainty Principle as a Consequence of the Effective Field Theory, Phys. Lett. B **765** 238 (2017), doi.org/10.1016/j.physletb.2016.11.054.