

Gini estimation under infinite variance

Andrea Fontanari - Delft University of Technology and CWI
 Nassim Nicholas Taleb - Tandon School of Engineering, NYU
 Pasquale Cirillo - Delft University of Technology

Abstract—We study different approaches to the estimation of the Gini index in presence of a fat-tailed data generator process, i.e. one in the stable distribution class with finite mean but infinite variance (i.e. with tail index $\alpha \in (1, 2)$). We show that, in such a case, the Gini coefficient can not be reliably estimated using conventional nonparametric methods, because of a downward bias that emerges in case of fat tails.

We start by discussing how the nonparametric estimator of the Gini index undergoes a phase transition in the symmetry structure of its asymptotic distribution, as the data distribution shifts from the domain of attraction of a light-tailed distribution to that of a fat-tailed one, especially in the case of infinite variance. We show how the nonparametric Gini bias increases with lower values of α .

We then prove that maximum likelihood estimation outperforms nonparametric methods, requiring a much smaller sample size to reach efficiency.

Finally, for fat-tailed data, we provide a simple correction mechanism to the small sample bias of the nonparametric estimator based on the distance between the mode and the mean of its asymptotic distribution.

I. INTRODUCTION

Wealth inequality studies represent a field of economics and statistics exposed to fat-tailed data generator processes, often with infinite variance. This is not at all surprising if we recall that the prototype of fat-tailed distributions, the Pareto, has been proposed for the first time to model household incomes [11].

However, the fat-tailedness of data can be problematic in the context of statistical estimation, as efficiency (and partially consistency) does not hold.

The scope of this work is, accordingly, to show how fat tails affect the estimation of one of the most celebrated measure of income inequality: the Gini index.

The literature concerning the estimation of the Gini index is wide and comprehensive (e.g. [15] for a review). However, strangely enough, almost no attention has been paid to its behavior in presence of fat tails, and this is curious if we consider that the Gini index itself can be seen as a measure of fat-tailedness [6], [7] and variability [2]. The most common method for the estimation of the the Gini index is nonparametric: one just computes the index from the empirical distribution of the available data (see Equation (5) below). But, as we show in this paper, this estimator suffers from a downward bias, when we deal with fat-tailed data.

Therefore our goal is to close this gap by deriving the limiting distribution of the nonparametric Gini estimator in presence of fat tails, and to analyze possible strategies to reduce biases.

We show how the maximum likelihood approach, despite the risk of model misspecification, needs much fewer observations to reach efficiency compared to a nonparametric one.¹

By fat-tailed data we indicate those data generated by a positive random variable X , whose distribution function $F(x)$ is regularly varying of order α . This means that for the corresponding survival function $\bar{F}(x) := 1 - F(x)$ the following holds:

$$\lim_{x \rightarrow \infty} x^\alpha \bar{F}(x) = L(x), \quad (1)$$

where $L(x)$ is a slowly varying function such that $\lim_{x \rightarrow \infty} \frac{L(cx)}{L(x)} = 1$ with $c > 0$ a constant, and where $\alpha > 0$ is called the tail exponent. Regularly varying and fat-tailed are thus synonyms.

Regularly varying distribution functions define a large class of random variables whose properties have been extensively studied in the context of extreme value theory [1], [4], when dealing with the probabilistic behavior of maxima and minima. In fact it is known that, if $(X_i)_{1 \leq i \leq n}$ are i.i.d. observations with a c.d.f. $F(x)$ in the regularly varying class, as defined in Equation (1), then their data generator process falls into the maximum domain of attraction of a Fréchet distribution of parameter ρ , in symbols $X \in MDA(\Phi(\rho))$. This means that, for the partial maximum $M_n = \max(X_1, \dots, X_n)$, one has

$$P\left(a_n^{-1}(M_n - b_n) \leq x\right) \xrightarrow{d} \Phi(\rho) = e^{-x^{-\rho}}, \quad \rho > 0, \quad (2)$$

with $a_n > 0$ and $b_n \in \mathbb{R}$ two normalizing constants [4]. The Fréchet distribution is one of the limiting distributions for extremes in extreme value theory, together with the Gumbel and the Weibull. It represents the fat-tailed and unbounded limiting case. The relationship between regularly varying random variables and the Fréchet class thus allows us to deal with a very large family of random variables, and to show that the Gini index is highly influenced by maxima, i.e. extreme wealth, as intuition also suggests [6], [7].

¹We observe a similar bias affecting the nonparametric measurement of quantile contributions (of the type the top 1% owns $x\%$ of the total wealth)[13]. This paper extends the problem to the more widespread Gini coefficient, and goes deeper by making links with the limit theorems.

The existence (finiteness) of the moments of a fat-tailed random variable depends on the tail exponent α , in fact

$$\begin{aligned} E(X^\delta) &< \infty \text{ if } \delta < \alpha \\ E(X^\delta) &= \infty \text{ if } \delta > \alpha. \end{aligned} \quad (3)$$

In the reminder of this work, we restrict our focus on data generator processes with finite mean and infinite variance, therefore, according to Equation (3), we limit our attention to the class of regularly varying distribution with tail index $\alpha \in (1, 2)$.

Table I and Figure 1 present numerically and graphically our story and already suggest its conclusion. Table I compares the Gini index obtained by the nonparametric estimator (Equation (5)) and the one obtained via maximum likelihood estimation (MLE) of the tail index (Subsection II-B).

As the first column shows, the convergence of the nonparametric estimator to the true value $g = 0.8333$ is extremely slow and monotonically increasing when the data distribution has an infinite variance. This suggests an issue not only in the tail structure of the distribution of the nonparametric estimator but also in its symmetry.

TABLE I: Comparison of nonparametric Gini to ML estimator, assuming tail $\alpha = 1.1$

n (number of obs.)	nonparametric		MLE		Error Ratio
	Mean	Bias	Mean	Bias	
10^3	0.711	-0.122	0.8333	0	1.4
10^4	0.750	-0.083	0.8333	0	3
10^5	0.775	-0.058	0.8333	0	6.6
10^6	0.790	-0.043	0.8333	0	156
10^7	0.802	-0.033	0.8333	0	$> 10^5$

Very interestingly, Figure 1 provides evidence that the limiting distribution of the nonparametric Gini index loses its properties of normality and symmetry [5] shifting to a fatter-tailed and skewed limit, as the distribution of the data enters an infinite variance domain. As we prove in Section II this is exactly what happens. When the data generator process is in the domain of attraction of a fat-tailed distribution, the asymptotic distribution of the Gini index moves away from Gaussianity towards a totally skewed to the right α -stable limit.

This change of behavior is the major responsible for the main problem of the nonparametric estimators for the Gini index: a downwards bias, for almost every sample size, when data are fat-tailed. Such a result, however, also suggests a solution to improve the quality of the nonparametric estimator: the idea is to correct for the skewness of the distribution of the nonparametric estimator in order to place its mode on the true value of the Gini. This correction, we show, improves the consistency and the bias of the estimator, reducing the risk of underestimating the Gini index (with all the

possible consequences in terms of economic and social policies [7])

The paper is organized as follows. In Section II we derive the asymptotic distribution of the sample Gini index when the data distribution has infinite variance. We provide an example with Pareto distributed data and we compare the quality of the limit distribution using the maximum likelihood and the nonparametric estimators. In Section III we propose a simple correction mechanism based on the mode-mean distance of the asymptotic distribution of the nonparametric estimator, to correct for the bias that is present in small samples. Section IV closes the paper.

II. ASYMPTOTICS AND PRE-ASYMPTOTICS OF THE NONPARAMETRIC ESTIMATOR

We derive the asymptotic distribution for the nonparametric estimator of the Gini index when the data generator process is fat-tailed with finite mean but infinite variance.

The so-called stochastic representation of the Gini, denoted by g , is

$$g = \frac{1}{2} \frac{\mathbb{E}(|X' - X''|)}{\mu}. \quad (4)$$

where g is the Gini index and X' and X'' are i.i.d. copies of a random variable X with c.d.f. $F(x) \in [c, \infty)$, $c > 0$ and finite mean $\mathbb{E}[X] = \mu$. The quantity $\mathbb{E}(|X' - X''|)$ is known as the "Gini mean difference" (GMD) [15].

The Gini index is thus the mean expected deviation between any two independent realizations of a random variable X , scaled by twice its mean [3], in order to obtain a quantity that lies in the interval $[0, 1]$.

What we call the nonparametric estimator of the Gini index of a sample $(X_i)_{1 \leq i \leq n}$ is defined as

$$G^{NP}(X_n) = \frac{\sum_{1 \leq i < j \leq n} |X_i - X_j|}{(n-1) \sum_{i=1}^n X_i}, \quad (5)$$

which can also be expressed as

$$G^{NP}(X_n) = \frac{\sum_{i=1}^n (2(\frac{i-1}{n-1} - 1)X_{(i)})}{\sum_{i=1}^n X_{(i)}} = \frac{\frac{1}{n} \sum_{i=1}^n Z_{(i)}}{\frac{1}{n} \sum_{i=1}^n X_i}, \quad (6)$$

where $X_{(1)}, X_{(2)}, \dots, X_{(n)}$ are the ordered statistics of X_1, \dots, X_n , such that: $X_{(1)} < X_{(2)} < \dots < X_{(n)}$.

The asymptotic normality of (6) under the finite variance assumption has been shown already by different authors [7], [15]. The result directly follows from the properties of the U-statistics and the L-estimators involved in the formulation of Equation (6).

A. The α -stable asymptotic limit of the Gini index

We first introduce some notation for what concerns α -stable distributions, as we need them to study the asymptotic limit of the Gini index.




Fig. 1: Empirical distribution for Gini nonparametric estimator for Type I Pareto distribution with different tail index (results has been centered to ease comparison)

A random variable X is distributed accordingly to an α -stable distribution if:

$$X \sim S(\alpha, \beta, \gamma, \delta)$$

where $\alpha \in (0, 2)$ is the tail parameter, $\beta \in [-1, 1]$ is the skewness, $\gamma \in \mathbb{R}^+$ is the scale and $\delta \in \mathbb{R}$ is the location [10]. It is interesting to notice that there is a correspondence between the α parameter of an α -stable random variable, and the α of a regularly varying random variable as per Equation (1). As discussed in [5], [10], a regularly varying random variable of order α is essentially α -stable, with the same tail coefficient. This is why we do not make any distinction in the use of α as parameter.

The standardized α -stable random variable is expressed as

$$Z_{\alpha, \beta} \sim S(\alpha, \beta, 1, 0) \quad (7)$$

α -stable distributions are a subclass of infinitely divisible distributions: thanks to their closure under convolution, they can be used to describe the limiting behavior of (rescaled) partials sums, $S_n = \sum_{i=1}^n X_i$, in the general Central Limit Theorem (CLT) setting [5]. In particular for $\alpha = 2$ we obtain the normal distribution as special case, which is known to be the limit distribution for the most classical CLTs.

In what follows we indicate that a random variable is in the domain of attraction of an α -stable distribution, by writing $X \in DA(S_\alpha)$.

Given the possible confusion that can arise in the parametrization of α -stable distributions, we clarify that we are here referring to the one in [12], also known as the S1 parametrization [10]. The characteristic function defining this type of α -stable is

$$E(e^{itX}) = e^{-\gamma^\alpha |t|^\alpha (1 - i\beta \text{sign}(t)) \tan(\frac{\pi\alpha}{2}) + i\delta t}, \alpha \neq 1$$

$$E(e^{itX}) = e^{-\gamma |t| (1 + i\beta \frac{2}{\pi} \text{sign}(t)) \ln |t| + i\delta t}, \alpha = 1$$

Since we are dealing with finite mean but infinite variance distributions, we restrict ourselves to $\alpha \in (1, 2)$, like for the case of fat-tailed random variables (as said the two α 's coincide). Recall that, for $\alpha \in (1, 2)$, the expected value of an α -stable random variable X is equal to the location parameter δ , i.e. $E(X) = \delta$. For more details, we suggest [10], [12].

We are now ready to study the asymptotic distribution of the Gini index estimator, as presented in (6), when the data generator process is fat-tailed with infinite variance.

Consider a sample made up of $(X_i)_{1 \leq i \leq n}$ i.i.d. observations with a c.d.f. $F(x)$ in the regularly varying class, as defined in Equation (1), with tail index $\rho \in (1, 2)$. This corresponds to considering a sample whose data generator process is in the domain of attraction of a Fréchet distribution with $\rho \in (1, 2)$, according to Equation (2).

The result is divided into two theorems, Theorem II.1 takes care of the limiting distribution of the "Gini Mean Difference" (GMD) (the numerator in Equation (5)), while Theorem II.2 completes the proof for the whole Gini index.

Theorem II.1. Consider a sequence $(X_i)_{1 \leq i \leq n}$ of i.i.d random variables from a distribution X on $[c, +\infty)$ $c > 0$, such that F is in the maximum domain of attraction of an Fréchet random variable, $X \in \text{MDA}(\Phi(\rho))$, with $\rho \in (1, 2)$. The sample Gini mean deviation (GMD) $\frac{\sum_{i=1}^n Z(i)}{n} = \text{GMD}_n$ satisfies the following limit in distribution.

$$\frac{n^{-\frac{1}{\rho}} (\sum_{i=1}^n Z(i) - n\theta)}{L_0(n)} \xrightarrow{d} Z_{\rho, 1}, \quad (8)$$

where $\theta = E[Z_{(i)}]$, $L_0(n)$ is a slowly varying function such that Equation (9) holds, and $Z_{\rho,1}$ is a standardized α -stable random variable defined as in Equation (7).

Proof. Theorem 3.1 (ii) in [8] proves the existence of a weak limit for the GMD prior to the existence of right scaling sequences for the sequence of i.i.d. random variables $Z_i = (2F(X_i) - 1)X_i$, where $F(X)$ is the integral probability transform of $X \sim F(x)$. Therefore what is left to prove is the characterization of scaling sequences and the limiting distribution.

Recall that by assumption $X \in MDA(\Phi(\rho))$. A standard result in extreme value theory [4] characterizes the tail of the distribution in the Fréchet domain as: $P(|X| > x) \sim L(x)x^{-\rho}$, where $L(x)$ is a slowly varying function. Given our restriction on the tail exponent $\rho \in (1, 2)$ this is the characterization for distributions in the domain of attraction of an α -stable distribution [5].

Therefore $X \in MDA(\Phi(\rho))$ is equivalent to $X \in DA(S_\rho)$ with $\rho \in (1, 2)$. This result enables us to use a CLT argument for the convergence of the sum in the estimator.

However, we first need to prove that the r.v. $Z \in DA(S_\alpha)$ as well. i.e. $P(|Z| > z) \sim L(z)z^{-\rho}$, with $\rho \in (1, 2)$ and $L(z)$ slowly varying.

Now notice that

$$P(|\tilde{Z}| > z) \leq P(|Z| > z) \leq P(2X > z),$$

where $\tilde{Z} = (2U - 1)X$ with $U \sim \text{Uniform}[0, 1]$ and $U \perp X$. The first bound holds because of positive dependence between X and $F(X)$ and can be proven rigorously by noting that $2UX \leq 2F(X)X$ by the rearrangement inequality [9]. The upper bound is trivial.

By assumption $P(2X > z) \sim 2^\rho L(z)z^{-\rho}$ (note that $L(\frac{z}{2}) \sim L(z)$ by definition of slowly varying function). In order to show that $\tilde{Z} \in DA(S_\alpha)$, we use the Breiman Theorem which ensure the stability of the Fréchet class under product, as long as the second random variable is not too fat-tailed [14].

To apply the theorem we re-write $P(|\tilde{Z}| > z)$ as:

$$P(|\tilde{Z}| > z) = P(\tilde{Z} > z) + P(-\tilde{Z} > z) = P(\tilde{U}X > z) + P(-\tilde{U}X > z),$$

where $\tilde{U} \sim \text{Uniform}[-1, 1]$ and $\tilde{U} \perp X$.

We focus on $P(\tilde{U}X > z)$ since for $P(-\tilde{U}X > z)$ the procedure is the same.

We have

$$P(\tilde{U}X > z) = P(\tilde{U}X > z | \tilde{U} > 0)P(\tilde{U} > 0) + P(\tilde{U}X > z | \tilde{U} \leq 0)P(\tilde{U} \leq 0),$$

for $z \rightarrow +\infty$. Now, we have that $P(\tilde{U}X > z | \tilde{U} \leq 0) \rightarrow 0$, while applying Breiman Theorem $P(\tilde{U}X > z | \tilde{U} > 0)$ becomes

$$P(\tilde{U}X > z | \tilde{U} > 0) \rightarrow E((\tilde{U})^\rho | U > 0)P(X > z)P(U > 0).$$

Therefore

$$P(|\tilde{Z}| > z) \rightarrow \frac{1}{2}E((\tilde{U})^\rho | U > 0)P(X > z) + \frac{1}{2}E((-\tilde{U})^\rho | U \leq 0)P(X > z).$$

From this

$$\begin{aligned} P(|\tilde{Z}| > z) &\rightarrow \frac{1}{2}P(X > z)[E((\tilde{U})^\rho | U > 0) \\ &\quad + E((-\tilde{U})^\rho | U \leq 0)] \\ &= \frac{2^\rho}{1 - \rho}P(X > z) \sim \frac{2^\rho}{1 - \rho}L(z)z^{-\rho}. \end{aligned}$$

We can then conclude that by the squeezing theorem

$$P(|Z| > z) \sim L(z)z^{-\rho},$$

as $z \rightarrow \infty$. Therefore $Z \in DA(S_\alpha)$ with $\alpha = \rho$.

We are now ready to invoke the generalized Central Limit Theorem [4] for the sequence Z_i , i.e.

$$\frac{\sum_{i=1}^n Z_i - n\theta}{L_0(n)n^{\frac{1}{\rho}}} \xrightarrow{d} Z_{\rho,\beta},$$

with $\theta = E(Z_i)$ and $L_0(n) = c_n n^{-\frac{1}{\rho}}$, where c_n is a sequence which must satisfy

$$\lim_{n \rightarrow \infty} \frac{nL(c_n)}{c_n^\rho} = \frac{1 - \rho}{\Gamma(2 - \rho) \cos(\frac{\pi\rho}{2})} = C_\rho, \quad (9)$$

with $Z_{\rho,\beta}$ a standardized α -stable r.v.

The skewness parameter β is such that

$$\frac{P(Z > z)}{P(|Z| > z)} \rightarrow \frac{1 + \beta}{2}.$$

Recalling that, by construction, $Z \in [-c, +\infty)$, the above expression reduces to

$$\frac{P(Z > z)}{P(Z > z) + P(-Z > z)} \rightarrow \frac{P(Z > z)}{P(Z > z)} = 1 \rightarrow \frac{1 + \beta}{2}, \quad (10)$$

therefore $\beta = 1$.

Hence by applying result (ii) of Theorem 3.1 in [8] we have the existence of the same limiting α -stable distribution also for the ordered version $Z_{(i)}$. \square

Theorem II.2. Given the same assumptions of Theorem II.1, the estimated Gini index $G^{NP}(X_n)$ satisfies the following limit in distribution:

$$n^{\frac{\rho-1}{\rho}} \left(\frac{G^{NP}(X_n) - g}{L_0(n)} \right) \xrightarrow{d} Q, \quad (11)$$

where $g = \mathbb{E}(G^{NP}(X_n))$ and Q is an α -stable random variable $S(\rho, 1, \frac{1}{\mu}, 0)$.

Proof. In Theorem II.1 we proved that $\sum Z_{(i)} \xrightarrow{d} Z_{\rho,\beta}$ if

$\sum Z_i \xrightarrow{d} Z_{\rho,\beta}$. Recall that by Equation (6) the Gini index is given by $\frac{\sum Z_{(i)}}{\sum X_i}$. Therefore it is sufficient to show that $\frac{\sum Z_i}{\sum X_i} \xrightarrow{d} \Lambda$ to prove that $\frac{\sum Z_{(i)}}{\sum X_i} \xrightarrow{d} \Lambda$. We achieve this using a Slutsky type argument.

Call Y_n the sequence $\frac{\sum_{i=1}^n Z_{(i)} - n\theta}{n^{\frac{1}{\rho}} L_0(n)}$. By Theorem II.1 we have that $Y_n \xrightarrow{d} Z_{\rho,1}$. Thanks to the Weak Law of Large Numbers [5] we also have that $m_n = \frac{\sum X_i}{n} \xrightarrow{p} \mu$. By Slutsky Theorem $\frac{Y_n}{m_n} \xrightarrow{d} \frac{1}{\mu} Z_{\rho,1}$.

What is left to prove is that also (11) converges in distribution to the same limit.

A well known theorem in probability theory [5] states that if a sequence $W_n \xrightarrow{d} \Lambda$ and $W_n - V_n = o_p(1)$ with V_n another sequence, then $V_n \xrightarrow{d} \Lambda$.

Take $W_n = \frac{Y_n}{m_n}$ and V_n the sequence defined by $V_n = n^{\frac{\rho-1}{\rho}} \left(\frac{G^{NP}(X_n) - g}{L_0(n)} \right)$, we show that

$$\frac{Y_n}{m_n} - V_n \xrightarrow{p} 0, \quad (12)$$

which reduces to showing that

$$n^{\frac{\rho-1}{\rho}} \theta \left(\frac{n}{\sum X_i} - \frac{1}{\mu} \right) \xrightarrow{p} 0. \quad (13)$$

Thanks to the continuous mapping theorem $\frac{n}{\sum X_i} \xrightarrow{p} \frac{1}{\mu}$, in particular $\frac{n}{\sum X_i} - \frac{1}{\mu} \rightarrow o_p(n^{-1})$.

Therefore Equation (13) goes to zero as $n \rightarrow \infty$ given that $\rho \in (1, 2)$ by assumption.

We conclude the proof by noting that an α -stable random variable is closed under scaling by a constant [12]. In particular this parameter changes as follows: the tail parameter is unchanged and equal to ρ , the skewness parameter $\beta = 1$ and the scale parameter $\gamma = \frac{1}{\mu}$. \square

In view of equation (10), in case of fat tails with $\alpha \in (1, 2)$, the asymptotic distribution of the Gini index estimator is always right-skewed regardless the distribution of the data generator process.

Comparing this result with the case of a finite-variance data generator process (leading to a Gaussian limit distribution), we can see how the limiting distribution of the estimator undergoes a phase transition in its skewness when variance becomes infinite, thus shifting from a symmetric Gaussian to a totally skewed α -stable. Therefore fat-tailed data not only induce a fatter-tail limit but they also change the shape. As a consequence the estimator, whose asymptotic consistency is still guaranteed, even in the fat-tailed case [8], will approach its true value more slowly and from below. Evidence for these behaviors are given in Table I.

B. The Maximum Likelihood estimator

Theorem II.2 shows that the nonparametric estimator for the Gini index is not the best option when dealing with infinite variance distributions, due to the skewness

and the fatness of the limit. A way out is to look for estimators that still preserve asymptotic normality under fat tails. In general this is not possible in view of the α -stable Central Limit Theorem to which any nonparametric estimator will eventually fall into. However, a solution is to use parametric techniques.

Theorem II.3 shows how, once a parametric family for the data generator process has been identified, it is possible to estimate the Gini index via MLE. The so-obtained estimator will not only be asymptotically normal but also asymptotically efficient. In Theorem II.3 we deal with random variables whose distribution belong to the large and flexible exponential family [7].

Theorem II.3. Let $X \sim F_\theta$ such that F_θ is a distribution belonging to the exponential family, i.e. whose density is

$$f_\theta(x) = h(x)e^{(\eta(\theta)T(x) - A(\theta))}$$

with $\theta \in \mathbb{R}$, and where $T(x)$, $\eta(\theta)$, $h(x)$, $A(\theta)$ are known functions. Then the Gini index obtained by plugging-in the maximum likelihood estimator of θ , $G^{ML}(X_n)_\theta$, is asymptotically normal and efficient. Namely:

$$\sqrt{n}(G^{ML}(X_n)_\theta - g_\theta) \xrightarrow{d} N(0, g_\theta^2 I^{-1}(\theta)), \quad (14)$$

where $g'_\theta = \frac{dg_\theta}{d\theta}$ and $I(\theta)$ is the Fisher Information.

Proof. The result follows easily from the asymptotic efficiency of the maximum likelihood estimators of the exponential family and the invariance principle of MLE. In particular the validity of the invariance principle for the Gini index is granted by continuity and monotonicity of g_θ with respect to θ .

The asymptotic variance is then obtained by application of the delta-method. \square

C. A Paretian illustration

We provide an illustration of the above results using a Pareto type I distribution as distribution for the data. Recall that the Pareto type I has the following density:

$$f(x) = \rho x_m^\rho x^{-\rho-1}, x > x_m \quad (15)$$

It is therefore clear that its survival function $\bar{F}(x)$ belongs to the regularly varying class with tail parameter $\alpha = \rho$. We can therefore apply the results of Section II to obtain the following Corollary.

Corollary 1. Let X_1, \dots, X_n be a sequence of i.i.d. observations with Pareto distribution with tail parameter $\rho \in (1, 2)$. Then the nonparametric Gini estimator has the following limit:

$$G^{NP}(X_n) - g \sim S\left(\rho, \frac{d}{n^{\frac{1}{\rho}}} \frac{(\rho-1)}{\rho}, 0, 1\right). \quad (16)$$

Proof. The results is a mere application of Theorem II.2, recalling that a Pareto distribution is in the domain of attraction of α -stable random variables with slowly varying function $L(x) = 1$. The sequence c_n to satisfy Equation (9) becomes $c_n = n^{\frac{1}{\rho}} C_\rho^{-1}$, therefore we have $L_0(n) = C_\rho^{-1}$ independent of n . For convenience we set $C_\rho^{-1} = d$. Additionally the mean of the distribution is also a function of ρ , that is $\mu = \frac{\rho}{\rho-1}$. \square

Corollary 2. *Let the sample X_1, \dots, X_n be distributed as in Corollary 1, let G_θ^{ML} be the MLE for the Gini index as defined in Theorem II.3. In particular $G_\rho^{ML} = \frac{1}{2\rho^{ML}-1}$. Then the asymptotic distribution of G_ρ^{ML} is*

$$G_\rho^{ML}(X_n) - g \sim N\left(0, \frac{4\rho^2}{n(2\rho-1)^4}\right). \quad (17)$$

Proof. The result follows from the fact that the Pareto distribution (with known minimum value x_m) belongs to the exponential family and therefore satisfies the regularity conditions necessary for asymptotic normality and efficiency of the maximum likelihood estimator. Recall also that the Fisher information for a Pareto distribution is $\frac{1}{\rho^2}$. \square

Now that we have worked out both asymptotic distributions, we can compare the quality of the convergence in the MLE and in the nonparametric cases.

In particular, we can approximate the distribution of the deviations from the true value of the Gini index for finite sample sizes, by using Equation (16) and Equation (17). Figure 2 shows how the noise around the mean of the two different types of estimators is distributed and how these distributions change as the number of observations increases. In particular, to ease the comparison between the maximum likelihood and the nonparametric estimators, we have fixed the number of observation in the MLE case and let them vary in the nonparametric one. We perform this study for different types of tail indices showing how big the impact is on the consistency of the estimator. It is worth to point out that, as the tail index goes towards 1 (the threshold value for a infinite mean), the mode of the distribution of the nonparametric estimator shifts farther and farther away from the mean of the distribution (centered on 0 by definition). This effect is responsible for the small sample bias observed in applications. Such a phenomenon is not present in the MLE case, thanks of the normality of the limit for every value of the tail parameter.

We can make our argument more rigorous by assessing the number of observations needed for the nonparametric estimator to be as good as the MLE one. The

following concentration measure is taken as quantitative measure for the expression "as good as":

$$\int \mathbb{1}_{|x|>c} dP_i = P_i(|X| > c), \quad (18)$$

with P_i , $i \in S, N$ being the distribution of each estimator as in Equations (17) and (16). Naturally $\mathbb{1}_A$ is the indicator function.

More precisely, we wish to compare for a fixed number of observations in (17), how many observations are required to reach the same value of the concentration measure in (16).

The problem can be rephrased in the following way. Consider the function

$$r(c, n) = \frac{P_S^n(|X| > c)}{P_N(|X| > c)}.$$

We wish to find \tilde{n} such that $r(c, \tilde{n}) = 1$ for fixed c .

Table II displays the results for different thresholds and tail parameters. In particular, we can see how, for a sample size of $n = 100$, the MLE estimator outperforms the nonparametric one. A much bigger amount of observations is indeed needed to obtain similar tail probabilities for the nonparametric estimator.

One interesting thing to notice is that the number of observations needed to match the tail probabilities does not vary uniformly with the threshold. However this is expected, since as the threshold goes to infinity or to zero, the tail probabilities are the same for every number of n . Therefore, given the unimodality of the distributions, we expect that there will be a threshold maximizing the number of observations needed to match the tail probabilities, while for all the other levels the number of observations will be smaller.

Figure 3 additionally shows some examples on how the equivalence of the tail probabilities is reached for different thresholds c and different tails index.

We conclude that, when in presence of fat-tailed data with infinite variance, a plug-in MLE based estimator should be preferred with respect to the nonparametric one.

TABLE II: The optimal number of observations \tilde{n} needed in order to match tail probabilities of the asymptotic MLE distribution with fixed $n = 100$.

α	Threshold c :			
	0.005	0.01	0.015	0.02
1.8	321	1242	2827	2244
1.5	844	836	2925	23036
1.2	402200	194372	111809	73888

III. SMALL SAMPLES CORRECTION

Theorem II.2 can be also used to provide a correction for the bias of the nonparametric estimator for small samples size.




Fig. 2: Comparisons between the MLE and the nonparametric asymptotic distribution for different values of the tail index α . The number of observations for MLE is fixed to $n = 100$. Note that, despite all the distributions have mean zero, the mode of the nonparametric one is different.

The key idea is to recognize that, for unimodal distributions, most observations come from around the mode. In symmetric distributions the mode and the mean coincide and therefore most observations will be close to the mean value as well. For skewed distributions, conversely, this is not the case. In particular, for right-skewed continuous unimodal distributions the mode is lower than the mean. Therefore, given that the distribution of the nonparametric Gini index is right-skewed, we expect that the realized (i.e. observed) value of the Gini index will be usually lower than the true value placed at the mean level. We can quantify this difference (i.e. the bias) by looking at the distance between the mode and the mean of its distribution, and once this distance is known, we can correct our Gini estimate by adding it.

Formally, we want to derive a corrected nonparametric estimator $G^C(X_n)$ such that

$$G^C(X_n) = G^{NP}(X_n) + ||m(G^{NP}(X_n)) - E(G^{NP}(X_n))||, \quad (19)$$

where $||m(G^{NP}(X_n)) - E(G^{NP}(X_n))||$ is the distance between the mode and the mean of the distribution of the nonparametric Gini estimator $G^{NP}(X_n)$.

Performing the type of correction described in Equation (19) is equivalent to shifting the distribution of $G^{NP}(X_n)$ in order to place its mode on the true value of the Gini index.

Ideally, we would like to measure this mode-mean distance $||m(G^{NP}(X_n)) - E(G^{NP}(X_n))||$ on the exact distribution of the Gini index to get the most accurate correction. However, the finite distribution is not always easily derivable and it requires assumptions on the parametric structure of the data generator process. Therefore we propose to use the limiting distribution for the nonparametric Gini obtained in Section II to approximate the finite sample distribution, and to estimate the mode-mean distance on it. This procedure allows for more freedom in the modeling assumptions and potentially decreases the number of parameters to be estimated, given that the limiting distribution only depends on the tail index of the data and the mean, which can be usually




Fig. 3: Speed of convergence of the probability ratio $r(c, n) = \frac{P_S^n(|X| > c)}{P_N(|X| > c)}$ as n grows. Note that MLE observations are fixed to $n = 100$

assumed to be a function of the tail index itself, as in the Pareto case i.e. $\mu = \frac{\rho}{\rho-1}$.

In particular, by exploiting the location-scale property of α -stable distributions and Equation (11), we approximate the distribution of $G^{NP}(X_n)$ for finite samples by

$$G^{NP}(X_n) \sim S(\rho, 1, \gamma(n), g), \quad (20)$$

where $\gamma(n) = \frac{1}{n^{\frac{\rho-1}{\rho}}} \frac{L_0(n)}{\mu}$ is the scale parameter of the limiting distribution.

As a consequence,

$$\begin{aligned} \|m(G^{NP}(X_n)) - E(G^{NP}(X_n))\| &\approx \|m(\rho, \gamma(n)) + g - g\| \\ &= \|m(\rho, \gamma(n))\|, \end{aligned}$$

where $m(\rho, \gamma(n))$ is the mode function of the α -stable distribution in Equation (20).

This means that, in order to obtain the correction term, the knowledge of the true Gini index is not necessary, view that $m(\rho, \gamma(n))$ does not depend on g .

We then compute the correction term as

$$\zeta(\rho, \gamma(n)) = \arg \max_x f(x), \quad (21)$$

where $f(x)$ is the numerical density² of the associated α -stable distribution in Equation (20) but centered in 0. Recalling that α -stable distributions are unimodal continuous distributions we conclude that $\zeta(\rho, \gamma(n)) = \arg \max_x f(x) = m(\rho, \gamma(n))$.

The corrected nonparametric estimator is thus

$$G^C(X_n) = G^{NP}(X_n) + \zeta(\rho, \gamma(n)), \quad (22)$$

whose asymptotic distribution is

$$G^C(X_n) \sim S(\rho, 1, \gamma(n), g + \zeta(\rho, \gamma(n))). \quad (23)$$

Note that the correction term $\zeta(\rho, \gamma(n))$ is a function of the tail index ρ and is connected to the sample size n by the scale parameter $\gamma(n)$ of the associated limiting distribution. It is important to notice that $\zeta(\rho, \gamma(n))$ is decreasing in n , and that $\lim_{n \rightarrow \infty} \zeta(\rho, \gamma(n)) \rightarrow 0$.

²Note also that for α -stable distributions the mode is not available in closed form, however it can be computed numerically by optimizing the numerical density [10].




Fig. 4: Comparisons between the corrected nonparametric estimator (in red) and the usual nonparametric estimator (black). For small sample sizes the corrected one clearly improves the quality of the estimation.

This happens because, as n increases, the distribution described in Equation (20) becomes more and more centered around its mean value, pushing to zero the distance between the mode and the mean. This ensures the asymptotic equivalence of the corrected estimator and the nonparametric one. Just observe that

$$\begin{aligned} \lim_{n \rightarrow \infty} |\bar{G}(X_n)^C - G^{NP}(X_n)| \\ = \lim_{n \rightarrow \infty} |G^{NP}(X_n) + \zeta(\rho, \gamma(n)) - G^{NP}(X_n)| \\ = \lim_{n \rightarrow \infty} |\zeta(\rho, \gamma(n))| \rightarrow 0. \end{aligned}$$

Naturally, because of the correction, $G^C(X_n)$ will always behave better in small samples.

Please also consider that, from Equation (23) the distribution of the corrected estimator has now mean $g + \zeta(\rho, \gamma(n))$, which converges to the true Gini g as $n \rightarrow \infty$, but its mode does not appear in g . Therefore the estimator is putting most of its probability mass on values close to the true Gini value.

In general, the quality of this correction depends on

the distance between the exact distribution of $G^{NP}(X_n)$ and its α -stable limit; the closer the two are to each other, the better the approximation.

From what we have written so far, it is clear that the correction term depends on the tail index of the data, and possibly also on their mean. These parameters, if not assumed to be known a priori, must be estimated. Therefore the additional uncertainty due to the estimation will reflect as well on the quality of the correction.

We conclude this Section by discussing the effect of the correction procedure with a simple example. We simulate 1000 Paretian samples of increasing size, from $n = 10$ to $n = 2000$, and for each sample size we compute both the original nonparametric estimator $G^{NP}(X_n)$ and the corrected $G^C(X_n)$. We repeat the experiment for different ρ 's.

Figure 4 presents our results. It is clear that the corrected estimators always perform as good as or better than the uncorrected one. In the case of sample size with $n \leq 500$, the gain is quite remarkable.

As expected, the difference between the estimators decreases with the sample size, as the correction term

is decreasing both in n and in the tail index ρ . Notice that, when the value of the tail index is equal to 2, we obtain the symmetric Gaussian distribution and the two estimators will coincide, given that the nonparametric estimator is no longer biased.

IV. CONCLUSIONS

In this paper we address the issue of asymptotic behavior of the Gini index nonparametric estimator in presence of a distribution with infinite variance, an issue that has been curiously ignored by the literature. The central mistake in the nonparametric methods largely used is to believe that asymptotic consistency translates into equivalent pre-asymptotic properties.

We show that a parametric approach provides better asymptotic results thanks to the properties of maximum likelihood estimation. Hence we strongly suggest that, if the collected data are suspected to be fat-tailed, parametric methods should be preferred.

If a fully parametric approach can not be used, we propose a simple correction mechanism for the nonparametric estimator based on the distance between the mode and the mean of its asymptotic distribution. Even if the correction works nicely, we suggest caution in its use for the additional uncertainty deriving from the estimation of the correction term.

REFERENCES

- [1] L. De Haan, A. Ferreira, *Extreme value theory: an introduction*, Springer (2007).
- [2] I. Eliazar, I. M. Sokolov, *Maximization of statistical heterogeneity. From Shannon's entropy to Gini's index*, Physica A 389 (2010) 3023-3038.
- [3] I. Eliazar, I.M. Sokolov, *Measuring statistical evenness: A panoramic overview*, Physica A 391 (2012) 1323-1353.
- [4] P. Embrechts, C. Kluppelberg, T. Mikosch, *Modelling Extremal Events for Insurance and Finance*, Springer (2003).
- [5] W. Feller, *An introduction to probability theory and its applications*, Vol. 2, Wiley 2008.
- [6] A. Fontanari, P. Cirillo, C.W. Oosterlee, *From Concentration Profiles to Concentration Maps. New Tools for the Study of Loss Distributions*, SSRN (2017) 2850949.
- [7] C. Kleiber, S.Kotz, *Statistical Size Distributions in Economics and Actuarial Sciences*, Wiley (2003).
- [8] D. Li, M.B. Rao, R.J.Tomkins, *The law of the iterated logarithm and central limit theorem for L-statistics*, Journal of Multivariate Analysis 78 (2001) 191-217.
- [9] G.H. Hardy, J.E. Littlewood, G. Pólya, *Inequalities*, Cambridge University Press (1952).
- [10] J.P. Nolan, *Parameterizations and modes of stable distributions*, Statistics and Probability Letters 38.2 (1998) 187-195.
- [11] V. Pareto, *La courbe de la répartition de la richesse* (1896), Reprinted in Rivista di Politica Economica 87 (1997) 647-700.
- [12] G. Samorodnitsky, M. S. Taqqu, *Stable non-Gaussian random processes: stochastic models with infinite variance*, Vol. 1, CRC Press (1994).
- [13] N.N. Taleb, R. Douady, *On the super-additivity and estimation biases of quantile contributions*, Physica A: Statistical Mechanics and its Applications 429 (2015) 252-260.
- [14] Y. Yang, S. Hu, T. Wu, *The tail probability of the product of depended random variables from max-domains of attraction*, Statistics and Probability Letters 81 (2011) 1876-1882.
- [15] S.Yitzhaki, E. Schechtman, *The Gini Methodology: A primer on a statistical methodology*, Springer (2012).