

Establishing low-lying doubly charmed baryons

Hua-Xing Chen¹, Qiang Mao^{1,2}, Wei Chen^{3,*}, Xiang Liu^{4,5,†} and Shi-Lin Zhu^{6,7,8‡}

¹*School of Physics and Beijing Key Laboratory of Advanced Nuclear Materials and Physics, Beihang University, Beijing 100191, China*

²*Department of Electrical and Electronic Engineering, Suzhou University, Suzhou 234000, China*

³*School of Physics, Sun Yat-Sen University, Guangzhou 510275, China*

⁴*School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China*

⁵*Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China*

⁶*School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China*

⁷*Collaborative Innovation Center of Quantum Matter, Beijing 100871, China*

⁸*Center of High Energy Physics, Peking University, Beijing 100871, China*

We systematically study the S -wave doubly charmed baryons using the method of QCD sum rules. Our results suggest that the Ξ_{cc}^{++} recently observed by LHCb can be well identified as the S -wave Ξ_{cc} state of $J^P = 1/2^+$. We study its relevant Ω_{cc} state, whose mass is predicted to be around 3.7 GeV. We also systematically study the P -wave doubly charmed baryons, whose masses are predicted to be around 4.1 GeV. Especially, there can be several excited doubly charmed baryons in this energy region, and we suggest to search for them in order to study the fine structure of the strong interaction.

PACS numbers: 14.20.Lq, 12.38.Lg

Keywords: doubly charmed baryons, excited charmed baryons, QCD sum rule

Introduction.— Fifteen years ago, the SELEX Collaboration reported an observation of the doubly charmed baryon Ξ_{cc}^+ in the $\Xi_{cc}^+ \rightarrow \Lambda_c^+ K^- \pi^+$ process [1], and determined its mass to be 3518.9 ± 0.9 MeV [2]. This is the only experimental evidence for the doubly charmed baryons, but all the other experiments did not confirm this signal [3], until the recent LHCb experiment [4], which observed the Ξ_{cc}^{++} in the $\Lambda_c^+ K^- \pi^+ \pi^+$ invariant mass spectrum. However, its mass was measured to be $3621.40 \pm 0.72 \pm 0.27 \pm 0.14$ MeV by LHCb, which value is significantly larger than that determined by SELEX. Note that the Ξ_{cc}^{++} and Ξ_{cc}^+ are isospin partners, and their mass difference should be only a few MeV.

Besides the recent LHCb experiment, there have been many other experiments in recent years [2, 5, 6], which consequently observed many excited heavy baryons, and gradually make the heavy baryons as well as heavy mesons an ideal platform to study the fine structure of the strong interaction. We refer to the review [7] for more discussions on this point. The doubly charmed baryons have been extensively studied using various theoretical methods, such as various quark models [8], the bag model [9], QCD sum rules [10], lattice QCD simulation [11], and others (for a incomplete list of works see Refs. [12]). Again, we refer to Ref. [7] for a brief review on these studies. More relevant discussions can be found in Refs. [13].

In this letter we systematically study the S -wave and P -wave doubly charmed baryons using the method of QCD sum rules [14]. We construct all the local S -wave doubly charmed baryon fields by investigating two configurations: one contains a $[cc]$ diquark [15] together with a light quark, and the other contains a $[cq]$ diquark together with another charm quark. These two configurations can be related by using the Fierz transformation (as long as local fields are used). After carefully examining these relations, we find a doubly charmed baryon field of $J^P = 1/2^+$, where all the three quark fields in-

side are at the ground-state. We use this field mixed with a few other component to perform QCD sum rule analyses, and find that the Ξ_{cc}^{++} recently observed by LHCb [4] can be well identified as the S -wave Ξ_{cc} state of $J^P = 1/2^+$. We also study its relevant S -wave Ω_{cc} state, whose mass is predicted to be around 3.7 GeV.

Following the same approach, we systematically study the P -wave doubly charmed baryons, whose masses are predicted to be around 4.1 GeV. Moreover, our results suggest that there can be several excited doubly charmed baryons in this energy region, similar to our previous studies on the excited singly heavy baryons [16], where we also found that there can be several singly excited charmed baryons. Recently, the LHCb experiment observed as many as five excited Ω_c states [6]. This may also happen for the excited doubly charmed baryons, so we suggest to search for them in the future LHCb and BelleII experiments. We believe this would greatly help to understand the fine structure of the strong interaction, and consequently improve our understanding of the quantum world.

Constructions of S -wave doubly charmed baryon fields.— As the first step, we discuss how we systematically construct the local S -wave doubly charmed baryon fields. They can only have the antisymmetric color configuration $[\epsilon^{abc} q_a c_b c_c]$, where the subscripts $a \cdots c$ are color indices, q represents an up, down or strange quark, and c represents a charm quark. The other structures, including flavor and spin/orbital/total angular momenta, can be generally described by using either

$$\mathcal{B}_1(x) = \epsilon^{abc} \left(c_a^T(x) C \Gamma_1 c_b(x) \right) \Gamma_2 q_c(x), \quad (1)$$

or

$$\mathcal{B}_2(x) = \epsilon^{abc} \left(q_a^T(x) C \Gamma_3 c_b(x) \right) \Gamma_4 c_c(x). \quad (2)$$

Here, $C = i\gamma_2\gamma_0$ is the charge-conjugation operator, the superscript T represents the transpose of the Dirac indices only, and

the matrices $\Gamma_{1\dots 4}$ are Dirac matrices describing the Lorentz structure. The other configuration

$$\mathcal{B}_3(x) = \epsilon^{abc} \left(c_a^T(x) C \Gamma_5 q_b(x) \right) \Gamma_6 c_c(x), \quad (3)$$

can be transformed to be $\mathcal{B}_2(x)$.

The former $\mathcal{B}_1(x)$ contains a $[cc]$ diquark together with a light quark, where we can clearly see the orbital structure between the two charm quarks; while the latter $\mathcal{B}_2(x)$ contains a $[cq]$ diquark together with another charm quark, where we can clearly see the orbital structure between the charm and light quarks inside the $[cq]$ diquark. In the present study we use local fields, so these two configurations can be related by using the Fierz transformation.

The first configuration $\mathcal{B}_1(x)$ can be easily constructed, because we can directly apply the Pauli principle to the two identical charm quarks contained in the $[cc]$ diquark. Following the method used in Ref. [17], we systematically construct all the possible fields, and find altogether two independent Dirac fields (without any free Lorentz index) of the spin-parity $J^P = 1/2^+$, two independent Rarita-Schwinger fields (with one free Lorentz index) of the pure spin-parity $J^P = 3/2^+$, and one tensor field (with two free antisymmetric Lorentz indices) of the same $J^P = 3/2^+$:

$$\eta_1(x) = \epsilon^{abc} \left(c_a^T(x) C \gamma_\mu c_b(x) \right) \gamma^\mu \gamma_5 q_c(x), \quad (4)$$

$$\eta_2(x) = \epsilon^{abc} \left(c_a^T(x) C \sigma_{\mu\nu} c_b(x) \right) \sigma^{\mu\nu} \gamma_5 q_c(x), \quad (5)$$

$$\eta_{3\alpha}(x) = \Gamma_{\alpha\mu} \epsilon^{abc} \left(c_a^T(x) C \gamma^\mu c_b(x) \right) q_c(x), \quad (6)$$

$$\eta_{4\alpha}(x) = \Gamma_{\alpha\mu} \times \left(\epsilon^{abc} \left(c_a^T(x) C \sigma^{\mu\nu} c_b(x) \right) \gamma_\nu q_c(x) \right. \\ \left. + \epsilon^{abc} \left(c_a^T(x) C \sigma^{\mu\nu} \gamma_5 c_b(x) \right) \gamma_\nu \gamma_5 q_c(x) \right), \quad (7)$$

$$\eta_{5\alpha_1\alpha_2}(x) = \Gamma_{\alpha_1\alpha_2\mu\nu} \times \left(\epsilon^{abc} \left(c_a^T(x) C \sigma^{\mu\nu} c_b(x) \right) \gamma_5 q_c(x) \right. \\ \left. + \epsilon^{abc} \left(c_a^T(x) C \sigma^{\mu\nu} \gamma_5 c_b(x) \right) q_c(x) \right), \quad (8)$$

where $\Gamma_{\alpha\mu}$ and $\Gamma_{\alpha_1\alpha_2\mu\nu}$ are the two projection operators:

$$\Gamma_{\mu\nu} = g_{\mu\nu} - \frac{1}{4} \gamma_\mu \gamma_\nu, \quad (9)$$

$$\Gamma_{\mu\nu\alpha\beta} = g_{\mu\alpha} g_{\nu\beta} - \frac{1}{2} g_{\nu\beta} \gamma_\mu \gamma_\alpha + \frac{1}{2} g_{\mu\beta} \gamma_\nu \gamma_\alpha + \frac{1}{6} \sigma_{\mu\nu} \sigma_{\alpha\beta}. \quad (10)$$

Among them, $\eta_{1,3\alpha}(x)$ contain the S -wave $[cc]$ diquark

$$\epsilon^{abc} c_a^T(x) C \gamma_\mu c_b(x) \quad [{}^{2S+1}L_J = {}^3S_1], \quad (11)$$

while the other three contain excited $[cc]$ diquarks. We can further identify:

$$\eta_1(x) : s_{[cc]} = 1, l_{[cc]} = 0, j_{[cc]} = 1, J = 1/2,$$

$$\eta_{3\alpha}(x) : s_{[cc]} = 1, l_{[cc]} = 0, j_{[cc]} = 1, J = 3/2.$$

Here, $s_{[cc]}$, $l_{[cc]}$ and $j_{[cc]}$ are the spin, orbital and total angular momenta of the $[cc]$ diquark, and J is the total angular momentum of the doubly charmed baryon.

The second configuration $\mathcal{B}_2(x)$ can not be so easily constructed. Still following the method used in Ref. [17], we systematically construct all the possible fields, and find that there are five non-vanishing Dirac fields of $J^P = 1/2^+$:

$$\eta'_1(x) = \epsilon^{abc} \left(q_a^T(x) C \gamma_5 c_b(x) \right) c_c(x), \quad (12)$$

$$\eta'_2(x) = \epsilon^{abc} \left(q_a^T(x) C \gamma_\mu c_b(x) \right) \gamma^\mu \gamma_5 c_c(x), \quad (13)$$

$$\eta'_6(x) = \epsilon^{abc} \left(q_a^T(x) C c_b(x) \right) \gamma_5 c_c(x), \quad (14)$$

$$\eta'_7(x) = \epsilon^{abc} \left(q_a^T(x) C \gamma_\mu \gamma_5 c_b(x) \right) \gamma^\mu c_c(x), \quad (15)$$

$$\eta'_8(x) = \epsilon^{abc} \left(q_a^T(x) C \sigma_{\mu\nu} c_b(x) \right) \sigma^{\mu\nu} \gamma_5 c_c(x). \quad (16)$$

However, only two of them are independent, and we can use the Fierz transformation to relate them to $\eta_1(x)$ and $\eta_2(x)$:

$$\left(\eta'_1 \ \eta'_2 \ \eta'_6 \ \eta'_7 \ \eta'_8 \right) = \left(\eta_1 \ \eta_2 \right) \times \begin{pmatrix} -\frac{1}{4} & -\frac{1}{2} & \frac{1}{4} & \frac{1}{2} & 0 \\ -\frac{1}{8} & 0 & -\frac{1}{8} & 0 & -\frac{1}{2} \end{pmatrix}. \quad (17)$$

Among them, $\eta'_{1,2}(x)$ contain the S -wave $[cq]$ diquarks:

$$\epsilon^{abc} q_a^T(x) C \gamma_5 c_b(x) \quad [{}^1S_0], \quad (18)$$

$$\epsilon^{abc} q_a^T(x) C \gamma_\mu c_b(x) \quad [{}^3S_1], \quad (19)$$

while the other three contain excited $[cq]$ diquarks. We can further identify

$$\eta'_1(x) : s_{[cq]} = 0, l_{[cq]} = 0, j_{[cq]} = 0, J = 1/2,$$

$$\eta'_2(x) : s_{[cq]} = 1, l_{[cq]} = 0, j_{[cq]} = 1, J = 1/2.$$

Here, $s_{[cq]}$, $l_{[cq]}$ and $j_{[cq]}$ are the spin, orbital and total angular momenta of the $[cq]$ diquark. Especially, from Eq. (17) we have the relation $\eta_1 = -2\eta'_2$, making this field interesting because all the three quark fields are at the ground-state.

Similarly, we find two independent Rarita-Schwinger fields and one tensor field, all of which have $J^P = 3/2^+$:

$$\eta'_{3\alpha}(x) = \Gamma_{\alpha\mu} \epsilon^{abc} \left(q_a^T(x) C \gamma^\mu c_b(x) \right) c_c(x), \quad (20)$$

$$\eta'_{4\alpha}(x) = \Gamma_{\alpha\mu} \times \left(\epsilon^{abc} \left(q_a^T(x) C \sigma^{\mu\nu} c_b(x) \right) \gamma_\nu c_c(x) \right. \\ \left. + \epsilon^{abc} \left(q_a^T(x) C \sigma^{\mu\nu} \gamma_5 c_b(x) \right) \gamma_\nu \gamma_5 c_c(x) \right), \quad (21)$$

$$\eta'_{5\alpha_1\alpha_2}(x) = \Gamma_{\alpha_1\alpha_2\mu\nu} \times \left(\epsilon^{abc} \left(q_a^T(x) C \sigma^{\mu\nu} c_b(x) \right) \gamma_5 c_c(x) \right. \\ \left. + \epsilon^{abc} \left(q_a^T(x) C \sigma^{\mu\nu} \gamma_5 c_b(x) \right) c_c(x) \right). \quad (22)$$

The former two fields $\eta'_{3\alpha,4\alpha}(x)$ can be related to $\eta_{3\alpha,4\alpha}(x)$ by using the Fierz transformation, and $\eta'_{5\alpha_1\alpha_2}(x)$ can also be related to $\eta_{5\alpha_1\alpha_2}(x)$. Only the first one $\eta'_{3\alpha}(x)$ contains the S -wave $[cq]$ diquark, while the other two contain excited diquarks. We can further identify

$$\eta'_{3\alpha}(x) : s_{[cq]} = 1, l_{[cq]} = 0, j_{[cq]} = 1, J = 3/2.$$

Interpretations of the Ξ_{cc}^{++} .— In the present study we shall use $\eta_1 = -2\eta'_2$, η'_1 , $\eta_{3\alpha}$ and $\eta'_{3\alpha}$ to perform QCD sum rule analyses in order to study the S -wave doubly charmed baryons of $J^P =$

$1/2^+$ and $J^P = 3/2^+$. We shall pay special attention to the field $\eta_1(x) = -2\eta'_2(x)$. Besides these single fields, we shall also study their mixing

$$\eta_1^M(\theta_1, x) = \cos \theta_1 \times \eta_1(x) + \sin \theta_1 \times \eta'_1(x), \quad (23)$$

$$\eta_{2\alpha}^M(\theta_2, x) = \cos \theta_2 \times \eta_{3\alpha}(x) + \sin \theta_2 \times \eta'_{3\alpha}(x). \quad (24)$$

These fields can couple to the doubly charmed baryons \mathcal{B} through

$$\langle 0|\eta|\mathcal{B}\rangle = f_{\mathcal{B}}u(p), \quad (25)$$

Then the two-point correlation functions can be written as:

$$\Pi(q^2) = i \int d^4x e^{iq \cdot x} \langle 0|T[\eta(x)\bar{\eta}(0)]|0\rangle = (\not{q} + M_{\mathcal{B}})\Pi(q^2).$$

In the present study we use the terms proportional to $M_{\mathcal{B}}$ to perform numerical analyses. Following Refs. [18] we obtain $M_{\mathcal{B}}$ through:

$$M_{\mathcal{B}}^2(s_0, M_B) = \frac{\int_{s_<}^{s_0} e^{-s/M_B^2} \rho(s) ds}{\int_{s_<}^{s_0} e^{-s/M_B^2} \rho(s) ds}, \quad (26)$$

where $s_<$ is the physical threshold, and $\rho(s)$ is the QCD spectral density which we evaluate up to dimension ten:

$$\rho(s) = \rho^{pert}(s) + \rho^{\langle \bar{q}q \rangle}(s) + \rho^{\langle GG \rangle}(s) + \rho^{\langle \bar{q}Gq \rangle}(s) + \rho^{\langle \bar{q}q \rangle \langle GG \rangle}(s) + \rho^{\langle \bar{q}Gq \rangle \langle GG \rangle}(s), \quad (27)$$

where $\rho^{pert}(s)$ is the perturbative term; $\rho^{\langle \bar{q}q \rangle}(s)$, $\rho^{\langle GG \rangle}(s)$, $\rho^{\langle \bar{q}Gq \rangle}(s)$, $\rho^{\langle \bar{q}q \rangle \langle GG \rangle}(s)$, and $\rho^{\langle \bar{q}Gq \rangle \langle GG \rangle}(s)$ are the terms containing the quark condensate $\langle \bar{q}q \rangle$, the gluon condensate $\langle g_s^2 GG \rangle$, the quark-gluon mixed condensate $\langle g_s \bar{q} \sigma Gq \rangle$, and their combinations $\langle \bar{q}q \rangle \langle g_s^2 GG \rangle$ and $\langle g_s \bar{q} \sigma Gq \rangle \langle g_s^2 GG \rangle$, respectively. We find that the leading perturbative term ($D = 0$) and the next-to-leading quark condensate $\langle \bar{q}q \rangle$ ($D = 3$) are important. The results of these spectral densities are too lengthy, so we list them in the supplementary file ‘‘OPE.nb’’. We use the values listed in Refs. [18] for these condensates and the charm quark mass (see also Refs. [19]).

There are two free parameters in Eq. (26): the Borel mass M_B and the threshold value s_0 . In order to obtain reliable QCD sum rule results, we require that the s_0 dependence and the M_B dependence of the mass prediction be weak. Beside this, we also need to carefully examine: a) the convergence of the QCD spectral density $\rho(s)$ through:

$$\frac{\Pi_8}{\Pi_{all}} \equiv \frac{\int_{s_<}^{\infty} e^{-s/M_B^2} \times \rho^{\langle \bar{q}q \rangle \langle GG \rangle}(s) \times ds}{\int_{s_<}^{\infty} e^{-s/M_B^2} \times \rho(s) \times ds}, \quad (28)$$

$$\frac{\Pi_{10}}{\Pi_{all}} \equiv \frac{\int_{s_<}^{\infty} e^{-s/M_B^2} \times \rho^{\langle \bar{q}Gq \rangle \langle GG \rangle}(s) \times ds}{\int_{s_<}^{\infty} e^{-s/M_B^2} \times \rho(s) \times ds}, \quad (29)$$

and b) the pole contribution defined as:

$$PC \equiv \frac{\int_{s_<}^{s_0} e^{-s/M_B^2} \times \rho(s) \times ds}{\int_{s_<}^{\infty} e^{-s/M_B^2} \times \rho(s) \times ds}. \quad (30)$$

Firstly, we study the Ξ_{cc} by replacing $q \rightarrow u/d$. The masses obtained using η'_1 and $\eta_1^M(\theta_1 = -11^\circ)$ are shown in Fig. 1 as functions of the threshold value s_0 and Borel mass M_B , and those obtained using $\eta'_{3\alpha}$ and $\eta_{2\alpha}^M(\theta_2 = 6^\circ)$ are shown in Fig. 2. We choose the working regions of s_0 and M_B to be $22 \text{ GeV}^2 < s_0 < 28 \text{ GeV}^2$ and $3.2 \text{ GeV}^2 < M_B^2 < 3.8 \text{ GeV}^2$, and find that the mass curves are quite stable inside these shady regions. We also list them in Table I together with Π_8/Π_{all} , Π_{10}/Π_{all} and PC defined in Eqs. (28-30). We find that the pole contributions are sufficiently large and the OPE spectral densities have good convergence inside these regions, suggesting that our sum rule results are reliable.




FIG. 1: Variations of $M_{\eta'_1, 1/2^+}$ (dashed curves) and $M_{\eta_1^M(\theta_1=-11^\circ), 1/2^+}$ (solid curves) with respect to the threshold value s_0 (left) and the Borel mass M_B (right).




FIG. 2: Variations of $M_{\eta'_{3\alpha}, 3/2^+}$ (dashed curves) and $M_{\eta_{2\alpha}^M(\theta_2=6^\circ), 3/2^+}$ (solid curves) with respect to the threshold value s_0 (left) and the Borel mass M_B (right).

Our results are:

- The spectral densities extracted from $\eta_1(x) = -2\eta'_2(x)$ and $\eta_{3\alpha}(x)$ are not complete ($\rho_{\eta_1}^{pert}(s) = \rho_{\eta_1}^{\langle \bar{q}Gq \rangle}(s) = 0$ and $\rho_{\eta_{3\alpha}}^{pert}(s) = \rho_{\eta_{3\alpha}}^{\langle \bar{q}Gq \rangle}(s) = 0$), so we can not use them to obtain reliable sum rule results.
- The masses extracted from $\eta'_1(x)$ and $\eta'_{3\alpha}(x)$ are both around 4.0 GeV, significantly larger than the mass of the Ξ_{cc} measured in the SELEX and LHCb experiments.
- We carefully fine-tune θ_1 to be -11° , and use $\eta_1^M(\theta_1 = -11^\circ)$ to perform sum rule analyses. The mass is extracted to be

$$M_{\eta_1^M(\theta_1=-11^\circ), \Xi_{cc}(1/2^+)} = 3.58_{-0.16}^{+0.15} \text{ GeV}, \quad (31)$$

where the central value is obtained by choosing $s_0 = 25 \text{ GeV}^2$ and $M_B^2 = 3.5 \text{ GeV}^2$, and the uncertainties come

TABLE I: Masses and decay constants of the S -wave charmed baryons.

Fields	Baryons	s_0 (GeV)	M_B^2 (GeV ²)	Pole	Convergence		Mass (GeV)	f (GeV ³)
					Π_8/Π_{All}	$\Pi_{10}/\Pi_{\text{All}}$		
$\eta_1 = -2\eta'_2$	Ξ_{cc}	—	—	—	—	—	—	—
	Ω_{cc}	—	—	—	—	—	—	—
η'_1	Ξ_{cc}	22 – 28	3.2 – 3.8	> 72%	< 1%	< 1%	$3.94^{+0.14}_{-0.15}$	$0.12^{+0.03}_{-0.03}$
	Ω_{cc}	22 – 28	3.2 – 3.8	> 72%	< 1%	< 1%	$3.98^{+0.12}_{-0.16}$	$0.13^{+0.02}_{-0.03}$
$\eta_{3\alpha}$	Ξ_{cc}	—	—	—	—	—	—	—
	Ω_{cc}	—	—	—	—	—	—	—
$\eta'_{3\alpha}$	Ξ_{cc}	22 – 28	3.2 – 3.8	> 73%	< 1%	< 1%	$3.95^{+0.13}_{-0.15}$	$0.11^{+0.01}_{-0.03}$
	Ω_{cc}	22 – 28	3.2 – 3.8	> 73%	< 1%	< 1%	$3.97^{+0.12}_{-0.17}$	$0.11^{+0.02}_{-0.03}$
$\eta_1^M(\theta_1 = -11^\circ)$	Ξ_{cc}	22 – 28	3.2 – 3.8	> 87%	< 1%	< 1%	$3.58^{+0.15}_{-0.16}$	$0.15^{+0.02}_{-0.03}$
	Ω_{cc}	22 – 28	3.2 – 3.8	> 84%	< 1%	< 1%	$3.70^{+0.13}_{-0.15}$	$0.17^{+0.02}_{-0.03}$
$\eta_2^M(\theta_2 = 6^\circ)$	Ξ_{cc}	22 – 28	3.2 – 3.8	> 90%	< 1%	< 1%	$3.58^{+0.14}_{-0.10}$	$0.061^{+0.006}_{-0.009}$
	Ω_{cc}	22 – 28	3.2 – 3.8	> 86%	< 1%	< 1%	$3.69^{+0.11}_{-0.15}$	$0.074^{+0.008}_{-0.011}$

from $\theta_1 (= -11 \pm 5^\circ)$, s_0 , M_B , and various quark masses and condensates [18]. Hence, the Ξ_{cc}^{++} recently observed by LHCb [4] can be well identified as the S -wave Ξ_{cc} state of $J^P = 1/2^+$.

- Similarly, we fine-tune θ_2 to be 6° , and use $\eta_2^M(\theta_2 = 6^\circ)$ to perform sum rule analyses:

$$M_{\eta_{2\alpha}^M(\theta_2=6^\circ), \Xi_{cc}(3/2^+)} = 3.58^{+0.14}_{-0.10} \text{ GeV}, \quad (32)$$

where the central value is again obtained by choosing $s_0 = 25 \text{ GeV}^2$ and $M_B^2 = 3.5 \text{ GeV}^2$. Hence, the Ξ_{cc}^{++} may also be identified as the S -wave Ξ_{cc} state of $J^P = 3/2^+$. However, the LHCb experiment preferentially retains longer-lived Ξ_{cc}^{++} candidates [4], disfavoring this interpretation because the Ξ_{cc} of $J^P = 3/2^+$ probably has a much shorter lifetime due to its radiative decays.

- Similarly, we use $\eta_1^M(\theta_1 = -11^\circ)$ and $\eta_2^M(\theta_2 = 6^\circ)$ to study the Ω_{cc} by replacing $q \rightarrow s$. The masses are extracted to be

$$M_{\eta_1^M(\theta_1=-11^\circ), \Omega_{cc}(1/2^+)} = 3.70^{+0.13}_{-0.15} \text{ GeV}, \quad (33)$$

$$M_{\eta_{2\alpha}^M(\theta_2=6^\circ), \Omega_{cc}(3/2^+)} = 3.69^{+0.11}_{-0.15} \text{ GeV}. \quad (34)$$

Again, the central values are obtained by choosing $s_0 = 25 \text{ GeV}^2$ and $M_B^2 = 3.5 \text{ GeV}^2$.

Our study on P -wave doubly charmed baryons.— The local P -wave doubly baryon fields are much more complicated than the S -wave ones. We follow Refs. [16] and construct them using the S -wave diquark fields defined in Eqs. (11, 18-19) as well as the following P -wave diquark fields:

$$\epsilon^{abc} [\mathcal{D}^\mu q_a^T(x)] C \gamma_5 c_b(x) [^1 P_1], \quad (35)$$

$$\epsilon^{abc} [\mathcal{D}^\mu q_a^T(x)] C \gamma^\nu c_b(x) [^3 P_0]/[^3 P_1]/[^3 P_2], \quad (36)$$

$$\epsilon^{abc} [\mathcal{D}^\mu c_a^T(x)] C \gamma^\nu c_b(x) [^3 P_0]/[^3 P_1]/[^3 P_2], \quad (37)$$

Here, $D^\mu = \partial^\mu + igA^\mu$ is the gauge-covariant derivative. There are altogether four configurations:

- Type 1- $[Dq-c]c$: we construct three fields of $J^P = 1/2^-$, three of $J^P = 3/2^-$, and one of $J^P = 5/2^-$.
- Type 2- $[q-c]Dc$: we construct three fields of $J^P = 1/2^-$, three of $J^P = 3/2^-$, and one of $J^P = 5/2^-$.
- Type 3- $[c-c]Dq$: we construct two fields of $J^P = 1/2^-$, two of $J^P = 3/2^-$, and one of $J^P = 5/2^-$.
- Type 4- $[Dc-c]q$: we construct one field of $J^P = 1/2^-$ and one of $J^P = 3/2^-$.

We note that: a) these exist more non-vanishing and independent fields containing other diquark fields, and b) there are some relations among these configurations due to the Fierz transformation.

We use all these P -wave doubly baryon fields to perform QCD sum rule analyses and systematically studied the P -wave doubly charmed baryons of $J^P = 1/2^-$, $3/2^-$ and $5/2^-$. We note that we have only used the single fields but have not investigated their mixing. We find that both type 1 ($[Dq-c]c$) and type 2 ($[q-c]Dc$) can lead to reasonable sum rule analyses, and our results suggest that the masses of the P -wave doubly charmed baryons (both Ξ_{cc} and Ω_{cc}) are around 4.0-4.2 GeV. Moreover, our results suggest that there can be several excited doubly charmed baryons in this energy region. We shall detailly discuss these results in our further work.

Summary and discussions.— We have systematically studied the S -wave and P -wave doubly charmed baryons using the method of QCD sum rules. We construct all the local S -wave doubly charmed baryon fields, and find one field of $J^P = 1/2^+$ where all the three quark fields inside are at the ground-state (verified by the Fierz transformation). We use this field mixed with a few other component to perform QCD sum rule analyses, and find that the Ξ_{cc}^{++} recently observed by LHCb [4]

can be well identified as the S -wave Ξ_{cc} state of $J^P = 1/2^+$. We have also studied its relevant S -wave Ω_{cc} state, whose mass is predicted to be around 3.7 GeV. We suggest to search for them in the future LHCb and BelleII experiments in the Cabibbo-favored weak decay channels, such as $\Xi_c^{(\prime)(*)0} D^{(*)+}$, $\Xi_c^{(\prime)(*)+} \bar{K}^{(*)0}$, and $\Omega_c^{(*)0} \pi^+(\rho^+)$, etc.

Following the same approach, we have systematically studied the P -wave doubly charmed baryons, whose masses are predicted to be around 4.1 GeV. Our results suggest that there can be several excited doubly charmed baryons in this energy region, similar to our previous studies on the excited singly heavy baryons [16], where we also found that there can be several singly charmed baryons. Recalling that the LHCb experiment observed as many as five excited Ω_c states [6], we also suggest to search for these excited doubly charmed baryons in the future LHCb and BelleII experiments in similar Cabibbo-favored weak decay channels ($\Xi_{cc} \rightarrow \Sigma^{(*)} D^{(*)}$, $\Lambda D^{(*)}$, $\Sigma_c^{(*)} K^{(*)}$, $\Lambda_c K^{(*)}$, $\Xi_c^{(\prime)(*)} \pi(\rho)$ and $\Omega_{cc} \rightarrow \Xi_c^{(\prime)(*)} D^{(*)}$, $\Xi_c^{(\prime)(*)} K^{(*)}$, $\Omega_c^{(*)} \pi(\rho)$, etc.) in order to study the fine structure of the strong interaction.

Acknowledgments

We thank Fu-Sheng Yu for helpful discussions. This project is supported by the National Natural Science Foundation of China under Grants No. 11475015, No. 11375024, No. 11222547, No. 11175073, No. 11575008, and No. 11621131001; the 973 program; the Ministry of Education of China (SRFDP under Grant No. 20120211110002 and the Fundamental Research Funds for the Central Universities); the National Program for Support of Top-notch Youth Professionals.

* Electronic address: chenwei29@mail.sysu.edu.cn

† Electronic address: xiangliu@lzu.edu.cn

‡ Electronic address: zhusi@pku.edu.cn

- [1] M. Mattson *et al.* [SELEX Collaboration], Phys. Rev. Lett. **89**, 112001 (2002).
- [2] C. Patrignani *et al.* [Particle Data Group], Chin. Phys. C **40**, 100001 (2016).
- [3] R. Chistov *et al.* [Belle Collaboration], Phys. Rev. Lett. **97**, 162001 (2006); B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. D **74**, 011103 (2006); S. P. Ratti, Nucl. Phys. Proc. Suppl. **115**, 33 (2003).
- [4] R. Aaij *et al.* [LHCb Collaboration], arXiv:1707.01621 [hep-ex].
- [5] J. Yelton *et al.* [Belle Collaboration], Phys. Rev. D **94**, 052011 (2016); Y. Kato *et al.* [Belle Collaboration], Phys. Rev. D **94**, 032002 (2016); R. Aaij *et al.* [LHCb Collaboration], JHEP **1705**, 030 (2017).
- [6] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **118**, 182001 (2017).
- [7] H. X. Chen, W. Chen, X. Liu, Y. R. Liu and S. L. Zhu, Rept. Prog. Phys. **80**, 076201 (2017).
- [8] D. Ebert, R. N. Faustov, V. O. Galkin and A. P. Martynenko, Phys. Rev. D **66**, 014008 (2002); V. V. Kiselev, A. K. Likhoded, O. N. Pakhomova and V. A. Saleev, Phys. Rev. D **66**, 034030 (2002); C. Albertus, E. Hernandez, J. Nieves and J. M. Verde-Velasco, Eur. Phys. J. A **32**, 183 (2007), Erratum: [Eur. Phys. J. A **36**, 119 (2008)]; A. Valcarce, H. Garcilazo and J. Vijande, Eur. Phys. J. A **37**, 217 (2008); A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Körner and V. E. Lyubovitskij, Phys. Rev. D **80**, 034025 (2009); T. Branz, A. Faessler, T. Gutsche, M. A. Ivanov, J. G. Körner, V. E. Lyubovitskij and B. Oehl, Phys. Rev. D **81**, 114036 (2010); M. Karliner and J. L. Rosner, Phys. Rev. D **90**, 094007 (2014); Z. Shah and A. K. Rai, Eur. Phys. J. C **77**, 129 (2017); F. S. Yu, H. Y. Jiang, R. H. Li, C. D. Lu, W. Wang and Z. X. Zhao, arXiv:1703.09086 [hep-ph].
- [9] S. Fleck and J. M. Richard, Prog. Theor. Phys. **82**, 760 (1989); D. H. He, K. Qian, Y. B. Ding, X. Q. Li and P. N. Shen, Phys. Rev. D **70**, 094004 (2004).
- [10] E. Bagan, M. Chabab and S. Narison, Phys. Lett. B **306**, 350 (1993); J. R. Zhang and M. Q. Huang, Phys. Rev. D **78**, 094007 (2008); Z. G. Wang, Eur. Phys. J. C **68**, 459 (2010); S. Narison and R. Albuquerque, Phys. Lett. B **694**, 217 (2011); T. M. Aliev, K. Azizi and M. Savci, Nucl. Phys. A **895**, 59 (2012); T. M. Aliev, K. Azizi and M. Savci, Phys. Lett. B **715**, 149 (2012).
- [11] J. M. Flynn *et al.* [UKQCD Collaboration], JHEP **0307**, 066 (2003); T. W. Chiu and T. H. Hsieh, Nucl. Phys. A **755**, 471 (2005); L. Liu, H. W. Lin, K. Orginos and A. Walker-Loud, Phys. Rev. D **81**, 094505 (2010); Y. Namekawa *et al.* [PACS-CS Collaboration], Phys. Rev. D **87**, 094512 (2013).
- [12] J. G. Körner, M. Krämer and D. Pirjol, Prog. Part. Nucl. Phys. **33**, 787 (1994); R. Roncaglia, D. B. Lichtenberg and E. Predazzi, Phys. Rev. D **52**, 1722 (1995); N. Brambilla, A. Vairo and T. Rosch, Phys. Rev. D **72**, 034021 (2005); C. H. Chang, C. F. Qiao, J. X. Wang and X. G. Wu, Phys. Rev. D **73**, 094022 (2006); C. H. Chang, J. X. Wang and X. G. Wu, Comput. Phys. Commun. **177**, 467 (2007); C. H. Chang, T. Li, X. Q. Li and Y. M. Wang, Commun. Theor. Phys. **49**, 993 (2008); X. Y. Wang and X. G. Wu, Comput. Phys. Commun. **184**, 1070 (2013); Z. F. Sun, Z. W. Liu, X. Liu and S. L. Zhu, Phys. Rev. D **91**, 094030 (2015); K. W. Wei, B. Chen and X. H. Guo, Phys. Rev. D **92**, 076008 (2015); M. Karliner and J. L. Rosner, arXiv:1706.06961 [hep-ph].
- [13] H. Y. Cheng, Front. Phys. (Beijing) **10**, 101406 (2015); H. X. Chen, W. Chen, X. Liu and S. L. Zhu, Phys. Rept. **639**, 1 (2016); F. K. Guo, C. Hanhart, U. G. Meissner, Q. Wang, Q. Zhao and B. S. Zou, arXiv:1705.00141 [hep-ph].
- [14] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 385 (1979); L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rept. **127**, 1 (1985); M. Nielsen, F. S. Navarra and S. H. Lee, Phys. Rept. **497**, 41 (2010).
- [15] R. L. Jaffe, Phys. Rept. **409**, 1 (2005).
- [16] X. Liu, H. X. Chen, Y. R. Liu, A. Hosaka and S. L. Zhu, Phys. Rev. D **77**, 014031 (2008); H. X. Chen, W. Chen, Q. Mao, A. Hosaka, X. Liu and S. L. Zhu, Phys. Rev. D **91**, 054034 (2015); Q. Mao, H. X. Chen, W. Chen, A. Hosaka, X. Liu and S. L. Zhu, Phys. Rev. D **92**, 114007 (2015); H. X. Chen, Q. Mao, A. Hosaka, X. Liu and S. L. Zhu, Phys. Rev. D **94**, 114016 (2016).
- [17] H. X. Chen, V. Dmitrasinovic, A. Hosaka, K. Nagata and S. L. Zhu, Phys. Rev. D **78**, 054021 (2008).
- [18] H. X. Chen, W. Chen, X. Liu, T. G. Steele and S. L. Zhu, Phys. Rev. Lett. **115**, 172001 (2015).
- [19] K. C. Yang, W. Y. P. Hwang, E. M. Henley and L. S. Kisslinger, Phys. Rev. D **47**, 3001 (1993); M. Eidmüller and M. Jamin,

Phys. Lett. B **498**, 203 (2001); S. Narison, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **17**, 1 (2002); V. Gimenez, V. Lubicz, F. Mescia, V. Porretti and J. Reyes, Eur. Phys. J. C **41**, 535 (2005); B. L. Ioffe and K. N. Zyblyuk, Eur. Phys. J. C **27**, 229 (2003); A. A. Ovchinnikov and A. A. Pivovarov, Sov.

J. Nucl. Phys. **48**, 721 (1988) [Yad. Fiz. **48**, 1135 (1988)]; P. Colangelo and A. Khodjamirian, “*At the Frontier of Particle Physics/Handbook of QCD*” (World Scientific, Singapore, 2001), Volume 3, 1495.