

Asymptotic Theory for the Maximum of an Increasing Sequence of Parametric Functions

Jonathan B. Hill *
University of North Carolina

March 12, 2022

Abstract

Hill and Motegi (2017) present a new general asymptotic theory for the maximum of a random array $\{\mathcal{X}_n(i) : 1 \leq i \leq \mathcal{L}\}_{n \geq 1}$, where each $\mathcal{X}_n(i)$ is assumed to converge in probability as $n \rightarrow \infty$. The array dimension \mathcal{L} is allowed to increase with the sample size n . Existing extreme value theory arguments focus on observed data $\mathcal{X}_n(i)$, and require a well defined limit law for $\max_{1 \leq i \leq \mathcal{L}} |\mathcal{X}_n(i)|$ by restricting dependence across i . The high dimensional central limit theory literature presumes approximability by a Gaussian law, and also restricts attention to observed data. Hill and Motegi (2017) do not require $\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)|$ to have a well defined limit nor be approximable by a Gaussian random variable, and we do not make any assumptions about dependence across i . We apply the theory to filtered data when the variable of interest $\mathcal{X}_n(i, \theta_0)$ is not observed, but its sample counterpart $\mathcal{X}_n(i, \hat{\theta}_n)$ is observed where $\hat{\theta}_n$ estimates θ_0 . The main results are illustrated by looking at unit root tests for a high dimensional random variable, and a residuals white noise test.

Keywords : maximum of multivariate mean, high dimensionality, non-Gaussian approximation.

MSC2010 subject classifications : 62E17, 62F40, 62M10, 05D10.

1 Introduction

Consider an array of random variables on a probability measure space $(\Omega, \mathcal{F}, \mathcal{P})$:

$$\{\mathcal{X}_n(i), \mathcal{Y}_n(i) : 1 \leq i \leq \mathcal{I}_n\}_{n \geq 1},$$

where $\{\mathcal{I}_n\}_{n \geq 1}$ is a sequence of positive integers, $\mathcal{I}_n \rightarrow \infty$ as $n \rightarrow \infty$. Under the assumption that $\mathcal{X}_n(i) \xrightarrow{P} 0$ as $n \rightarrow \infty$ for each i , Hill and Motegi (2017) prove for some sequence of positive

*Department of Economics, University of North Carolina at Chapel Hill; jbhill@email.unc.edu.

integers $\{\mathcal{L}_n\}$ that satisfies $\mathcal{L}_n \rightarrow \infty$:

$$\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| \xrightarrow{p} 0. \quad (1)$$

By operating on (1), for any two arrays of \mathcal{F} -measurable random variables $\{\mathcal{X}_n(i), \mathcal{Y}_n(i) : 1 \leq i \leq \mathcal{I}_n\}_{n \geq 1}$, if $\mathcal{X}_n(i) - \mathcal{Y}_n(i) \xrightarrow{p} 0$ for each i Hill and Motegi (2017) then show that the difference in maxima satisfy:

$$\left| \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{Y}_n(i)| \right| \xrightarrow{p} 0. \quad (2)$$

Examples are a sample mean $\mathcal{X}_n(i) = 1/n \sum_{t=1}^n x_t(i)$ or standardized means $\mathcal{X}_n(i) = 1/\sqrt{n} \sum_{t=1}^n x_t(i)$ and $\mathcal{Y}_n(i) = 1/\sqrt{n} \sum_{t=1}^n y_t(i)$, where $\{x_t(i), y_t(i)\}_{t=1}^n$ are the sample paths of some processes on $(\Omega, \mathcal{F}, \mathcal{P})$. This has been studied extensively in the Gaussian approximation and high dimensional Gaussian central limit theory literatures, discussed below. At the highest level of generality we work with maxima of positive values in order to exploit convenient inequalities.

The focus of the present paper is to extend the key ideas of Hill and Motegi (2017) to filtered residuals, and to apply the result to a unit root test and a white noise test. The authors do not impose any restrictions on dependence in $(\mathcal{X}_n(i), \mathcal{Y}_n(i))$ across coordinates i , nor do we require $(\mathcal{X}_n(i), \mathcal{Y}_n(i))$ to belong to a specific domain of attraction. For example, in the normalized mean case we neither require either $(\mathcal{X}_n(i), \mathcal{Y}_n(i))$ to be Gaussian nor converge in law to Gaussian random variables. The generality arises from a new result for convergence of high dimensional arrays which does not rely on probabilistic properties, although naturally lends itself to probabilistic applications.

The approach differs from standard weak convergence methods when applied to an array of standardized means $\{\mathcal{X}_n(i) : 1 \leq i \leq \mathcal{I}_n\}_{n \geq 1}$, e.g. $\mathcal{X}_n(i) = 1/\sqrt{n} \sum_{t=1}^n x_t(i)$. Weak convergence for $\{\mathcal{X}_n(i) : i \in N\}$ in the broad sense of Hoffmann-Jorgensen (1984, 1991) is a potential option. It is known that such weak convergence to a Gaussian limit, with a version that has uniformly bounded and uniformly continuous sample paths, is equivalent to pointwise convergence and the existence of a pseudo metric d on N such that (N, d) is a totally bounded pseudo metric

space and a stochastic equicontinuity property based on d holds. See Dudley (1978, 1984) and Pollard (1990, Chapters 9-10). If d is the Euclidean distance, for example, then (N, d) is not totally bounded. Hill and Motegi (2017) take a different approach that completely sidesteps the approach of Hoffmann-Jorgensen (1984, 1991) by first demonstrating pointwise convergence $\mathcal{X}_n(i) \xrightarrow{d} \mathcal{X}(i)$. They then work with a probability construction for the array $\{\mathcal{X}_n(i) - \mathcal{X}(i) : 1 \leq i \leq \mathcal{I}_n\}_{n \geq 1}$ and apply our general result for convergence of arrays to be able to show $|\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}(i)|| \xrightarrow{P} 0$.

The primary tools used here date in some form to seminal theory developed by Ramsey (1930) and its implications for monotone subsequences and convergence, cf. the Erdős-Szekeres theorem (Erdős and Szekeres, 1935). Hill and Motegi (2017) significantly augment a convergence result for non-stochastic arrays presented in Boehme and Rosenfeld (1974, Lemma 1) in key ways. The latter claim that if an array $\{\mathcal{A}_{k,n} : 1 \leq k \leq \mathcal{I}_n\}_{n \geq 1}$, where $\mathcal{I}_n \rightarrow \infty$ as $n \rightarrow \infty$, lies in a first countable topological space, and $\lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \mathcal{A}_{k,n} = 0$, then $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_l), n_l} = 0$ for some infinite subsequence $\{n_l\}_{l=1}^{\infty}$ of positive integers, and some mapping $\mathcal{L}(n_l) \rightarrow \infty$. (Recall that any metric space is a first countable topological space.) In order to use the result for maxima, we extend $\{n_l\}_{l=1}^{\infty}$ to \mathbb{N} and therefore achieve $\lim_{n \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n), n} = 0$ in Lemma 2.1. We require additional assumptions that lends itself to deriving (1) and (2). A practical application includes when $\mathcal{A}_{k,n} = \int_0^1 P(1 - e^{-\max_{1 \leq i \leq k} |\mathcal{X}_n(i)|} > \epsilon) d\epsilon$, the foundation for showing $\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| \xrightarrow{P} 0$ in the proof of Theorem 2.2 below.

Chernozhukov, Chetverikov, and Kato (2013a, 2016) work with normalized sample means $1/\sqrt{n} \sum_{t=1}^n x_t(i)$ where $\{x_t(i)\}_{t=1}^n$ are independent, zero mean and square integrable random variables. They bypass extreme value theoretic arguments and therefore do not need to restrict dependence across i . This is accomplished by developing new tools for deriving Gaussian approximations based on Slepian and Sudakov-Fernique methods. They prove the impressive result that for some $(K, \zeta) > 0$, and $\mathcal{L}_n \rightarrow \infty$ with $\mathcal{L}_n = O(e^{o(n^c)})$ for some $c > 0$:

$$\mathcal{A}_{\mathcal{L}_n, n} \equiv \sup_{c \geq 0} \left| P \left(\max_{1 \leq i \leq \mathcal{L}_n} \left| \frac{1}{\sqrt{n}} \sum_{t=1}^n x_t(i) \right| \leq c \right) - P \left(\max_{1 \leq i \leq \mathcal{L}_n} \left| \frac{1}{\sqrt{n}} \sum_{t=1}^n y_t(i) \right| \leq c \right) \right| \leq K n^{-\zeta} \quad (3)$$

where $y_t(i)$ are zero mean normally distributed with covariance function $E[y_t(i)y_t(j)] = 1/n \sum_{t=1}^n E[x_t(i)x_t(j)]$.

The literature on such Gaussian couplings has a substantial history, where generally $\mathcal{L}_n = O(n^c)$ for some c . See, for example, Pollard (2002, Chapter 10) for a general review, see Yurinskii (1977) for a seminal result, and see Dudley and Philipp (1983, Lemma 2.12) and Le Cam (1988). See also Portnoy (1985, 1986), Götze (1991) and Chernozhukov, Chetverikov, and Kato (2013b, Appendix L) for theory and references on high dimensional Gaussian central limit theory.

In the above literature, Gaussianity is key, which therefore neglects non-standard asymptotics, including heavy tailed data or non-stationary data. It also does not apply in general when working with filtered data: in this case an intermediate step is required linking two sequences in which neither may be Gaussian. As such, it does not include cases where a filter leads to non-standard asymptotics, including with some parameters are weakly or non-identified and when a parameter boundary value occurs (e.g. Andrews, 1999, Andrews and Cheng, 2012).

Consider, for example, random functions $x_t(i, \theta_0(i))$ where $\theta_0(i)$ is not observed but estimable, write $\mathcal{X}_n(i, \theta_0(i)) \equiv 1/\sqrt{n} \sum_{t=1}^n x_t(i, \theta_0(i))$, and let $\hat{\theta}_n(i)$ be a plug-in estimator for $\theta_0(i)$. Write $\mathcal{X}_n(i) \equiv \mathcal{X}_n(i, \hat{\theta}_n(i))$. In order to gain inference on $\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)|$ we first require an asymptotic expansion $\mathcal{X}_n(i) = \mathcal{Y}_n(i) + o_p(1)$ for some process $\{\mathcal{Y}_n(i)\}$ that ostensibly depends on $\theta_0(i)$ and pre-asymptotic properties of $\hat{\theta}_n(i)$, leading to $|\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{Y}_n(i)|| \xrightarrow{p} 0$ as in (2). Even if $\mathcal{Y}_n(i)$ is *asymptotically* normal, it need not be Gaussian, hence the Gaussian approximation literature does not apply. Of course, $\mathcal{Y}_n(i)$ need not be asymptotically normal for (2) to apply. In Section 4 we illustrate non-standard asymptotics by allowing for unit root non-stationarity, and we treat a white noise test for an unobserved regression error term to illustrate the use of a filter.

The maximum of an increasing sequence of normalized sample covariances $\hat{\gamma}_n(i) \equiv 1/n \sum_{t=1}^n x_t x_{t-i}$ has been studied at least since Berman (1964) and Hannan (1974). See Jirak (2011) and Xiao and Wu (2014) for recent theory and references. In this literature x_t is assumed observed, the exact asymptotic distribution form of a suitably normalized $\sqrt{n} \max_{1 \leq i \leq \mathcal{L}_n} |\hat{\gamma}_n(i) - E[x_t x_{t-i}]|$ is sought, and underlying assumptions ensure $\sqrt{n}(\hat{\gamma}_n(i) - E[x_t x_{t-i}])$ converges in finite dimensional dis-

tributions to a Gaussian law $\mathcal{Z}(i)$. In turn, $\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{Z}(i)|$ must converge in law to a well defined random variable, which requires asymptotic independence $E[\mathcal{Z}(i)\mathcal{Z}(j)] \rightarrow 0$ as $|i - j| \rightarrow \infty$. See, e.g., Leadbetter, Lindgren, and Rootzén (1983, Chapter 6), Hüsler (1986, 1993), Hombler and McCormick (1995) and Falk, Hüsler, and Reiss (2011, Chapter 9). We sidestep extreme value theoretic arguments, dependence across i is unrestricted, and residuals are allowed, hence x_t need not be observed in practice. See Section 4.

We cannot generally provide an upper bound on the divergence rate $\mathcal{L}_n \rightarrow \infty$, similar to ones in Chernozhukov, Chetverikov, and Kato (2013a, 2016) in the Gaussian coupling literature, and Xiao and Wu (2014) in the extreme value theory literature. This is an unavoidable cost for our (i) basing probabilistic statements like (2) on a general array convergence result that itself does not make use of probabilistic properties of (Gaussian) random variables; and (ii) allowing for filtered data and therefore requiring asymptotic linkages between maxima that do not involve Gaussian processes. Chernozhukov, Chetverikov, and Kato (2013a, 2015, 2016), however, appear to have the sharpest result and most general bound on $\mathcal{L}_n \rightarrow \infty$ for Gaussian approximations (for independent data).

We do not treat a bootstrap theory, for example for the sample mean maximum $\max_{1 \leq i \leq \mathcal{L}_n} |1/\sqrt{n} \sum_{t=1}^n x_t(i)|$, because by using our main results any existing bootstrap theory will hold under its specified pointwise assumptions. See Hill and Motegi (2017).

The remaining sections are organized as follows. In Section 2 we present the main results (1) and (2) of Hill and Motegi (2017). These are used in Section 3 for convergence of maxima when filtered residuals are used. Examples are provided in Section 4, and concluding remarks are left for Section 5.

In the following $|\cdot|$ and $\|\cdot\|$ are respectively the l_1 - and l_2 -matrix norms.

2 Main Results

All random variables are assumed to exist on a complete measure space, and probabilities where applicable are outer probability measures. See, e.g., Dudley (1984, p. 101) and Pollard (1984,

Appendix C). See Dudley and Philipp (1983) for theory that sidesteps measurability challenges specifically for normalized means.

The first result due to Hill and Motegi (2017) concerns convergence of arrays. It is based on, and extends, a result from Boehme and Rosenfeld (1974, Lemma 1). They work in first countable spaces. Because we require some structure on the space we work in, and because any metric space is first countable, we simply work in (\mathbb{R}, d) with metric d for ease of notation. All proofs are placed in the appendix.

Lemma 2.1 *Assume the array $\{\mathcal{A}_{k,n} : 1 \leq k \leq \mathcal{I}_n\}_{n \geq 1}$ lies (\mathbb{R}, d) , where $\{\mathcal{I}_n\}_{n \geq 1}$ is a sequence of positive integers, $\mathcal{I}_n \rightarrow \infty$ as $n \rightarrow \infty$. Let $\lim_{n \rightarrow \infty} \mathcal{A}_{k,n} = 0$ for each fixed k , and $\mathcal{A}_{k,n} \leq \mathcal{A}_{k+1,n}$ for each n and all k . Then $\lim_{n \rightarrow \infty} \mathcal{A}_{\mathcal{L}_n,n} = 0$ for some sequence $\{\mathcal{L}_n\}$ of positive integers, $\mathcal{L}_n \rightarrow \infty$ and $\mathcal{L}_n \leq \mathcal{I}_n$, that is not unique.*

Remark 1 Boehme and Rosenfeld (1974) only require $\lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \mathcal{A}_{k,n} = 0$, and do not impose monotonicity, thus $\lim_{n \rightarrow \infty} \mathcal{A}_{k,n} = 0 \forall k$ and $\mathcal{A}_{k,n} \leq \mathcal{A}_{k+1,n}$ are stronger assumptions. The first property, however, is key towards proving $\lim_{n \rightarrow \infty} \mathcal{A}_{\mathcal{L}_n,n} = 0$, rather than merely a subsequence $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}_{n_l},n_l} = 0$ as in Boehme and Rosenfeld (1974). Monotonicity is used to identify \mathcal{L}_n as a function only of n based on using a multiple subsequence argument. The maximum over a subsequence of positive values satisfies monotonicity, and $\lim_{n \rightarrow \infty} \mathcal{A}_{k,n} = 0 \forall k$ holds when applied to pointwise probability convergence problems discussed in the sequel.

Remark 2 Monotonicity implies $\{\mathcal{L}_n\}$ is not unique since $\lim_{n \rightarrow \infty} \mathcal{A}_{\hat{\mathcal{L}}_n,n} = 0$ for any sequence $\{\hat{\mathcal{L}}_n\}$ of positive integers with $\hat{\mathcal{L}}_n \rightarrow \infty$ and $\limsup_{n \rightarrow \infty} \{\hat{\mathcal{L}}_n/\mathcal{L}_n\} < 1$.

The next result uses Lemma 2.1 as the basis for deriving (1) and (2).

Theorem 2.2 *Let $\{\mathcal{X}_n(i), \mathcal{Y}_n(i) : 1 \leq i \leq \mathcal{I}_n\}_{n \geq 1}$ be arrays of random variables, where $\{\mathcal{I}_n\}_{n \geq 1}$ is a sequence of positive integers, $\mathcal{I}_n \rightarrow \infty$ as $n \rightarrow \infty$.*

a. If $\mathcal{X}_n(i) \xrightarrow{p} 0$ for each fixed i , then $\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| \xrightarrow{p} 0$ for some sequence $\{\mathcal{L}_n\}$ of positive integers with $\mathcal{L}_n \rightarrow \infty$, that is not unique.

b. If $\mathcal{X}_n(i) - \mathcal{Y}_n(i) \xrightarrow{p} 0$ for each fixed i then for some sequence $\{\mathcal{L}_n\}$ of positive integers with $\mathcal{L}_n \rightarrow \infty$, that is not unique:

$$\left| \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{Y}_n(i)| \right| \leq \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i) - \mathcal{Y}_n(i)| \xrightarrow{p} 0.$$

Remark 3 A similar result exists under *almost sure* convergence, although a different argument is required. See the supplemental material Hill (2017, Appendix B).

Remark 4 The method of proof for Lemma 2.1 shows the *existence* of such a sequence $\{\mathcal{L}_n\}$ and therefore cannot provide an upper bound on the rate $\mathcal{L}_n \rightarrow \infty$. This seems unavoidable since we are working with general array convergence rather than, for example, the specific attributes of Gaussian probability tails. The payoff is that such generality ultimately permits non-Gaussian couplings, a data filter, and arbitrary dependence across i , as we treat in turn below.

3 Asymptotics for Maxima Based on Filtered Data

We now work with a parametric array $\{\mathcal{X}_n(i, \theta_0(i)) : 1 \leq i \leq \mathcal{L}\}_{n \geq 1}$, where $\theta_0(i)$ is an unknown but estimable parameter in \mathbb{R}^k that may depend on i . Let $\hat{\theta}_n(i)$ be an estimator of $\theta_0(i)$. We assume $\mathcal{X}_n(i, \theta_0(i))$ is unobserved while $\mathcal{X}_n(i, \hat{\theta}_n(i))$ is observed. Our leading example is a sample serial correlation coefficient for time series regression model errors (see Section 4). Another is a sample mean of an observed time series scaled by its conditional variance, e.g. GARCH residuals.

Our primary goal is to prove under fairly general conditions that for some stochastic process $\{\mathcal{Z}_n(i)\}_{i \in \mathbb{N}}$, and some non-unique $\{\mathcal{L}_n\}$, $\mathcal{L}_n \rightarrow \infty$:

$$\left| \max_{1 \leq i \leq \mathcal{L}_n} \left| \mathcal{X}_n(i, \hat{\theta}_n(i)) \right| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{Z}_n(i)| \right| \xrightarrow{p} 0. \quad (4)$$

In view of Theorem 2.2, it suffices to prove $\mathcal{X}_n(i, \hat{\theta}_n(i)) - \mathcal{Z}_n(i) \xrightarrow{p} 0$ for each fixed i . Sufficient conditions follow. These are not the most general possible, but give a reasonably general environment to work in.

Assumption 1

a. $\mathcal{X}_n(i, \theta)$ is continuous and differentiable in $\theta \in \Theta(i)$, where $\Theta(i)$ is a compact subset of \mathbb{R}^k , $k \in \mathbb{N}$.

b. $\theta_0(i) \in \Theta(i)$.

c. There exists a continuous non-stochastic function $\mathcal{D}(i, \cdot) : \Theta(i) \rightarrow \mathbb{R}^k$, and a compact neighborhood $\mathcal{N}_0(i)$ of $\theta_0(i)$ with positive Lebesgue measure such that

$$\sup_{\theta \in \mathcal{N}_0(i)} \left| \frac{1}{\sqrt{n}} \frac{\partial}{\partial \theta} \mathcal{X}_n(i, \theta) - \mathcal{D}(i, \theta) \right| \xrightarrow{p} 0.$$

d. There exists stochastic processes $\{\mathcal{S}_n(i), \mathcal{M}_n(i) : i \in \mathbb{N}\}$ such that $\mathcal{X}_n(i, \theta_0(i)) = \mathcal{S}_n(i) + o_p(1)$ and $\sqrt{n}(\hat{\theta}_n(i) - \theta_0(i)) = \mathcal{M}_n(i) + o_p(1)$ for each i . Moreover, there exists non-degenerate stochastic processes $\{\mathcal{S}(i), \mathcal{M}(i) : i \in \mathbb{N}\}$ such that $(\mathcal{S}_n(i), \mathcal{M}_n(i)) \xrightarrow{d} (\mathcal{S}(i), \mathcal{M}(i))$ for each i .

Remark 5 Uniform convergence (c) holds when $(\partial/\partial\theta)\mathcal{X}_n(i, \theta)/\sqrt{n} \xrightarrow{p} \mathcal{D}(i, \theta)$ for each $\theta \in \mathcal{N}_0(i)$, and a stochastic equicontinuity condition holds. The latter holds, for example, when $\mathcal{X}_n(i, \theta)$ is twice continuously differentiable and the envelope

$$E \left[\frac{1}{\sqrt{n}} \sup_{\theta \in \mathcal{N}_0(i)} \left| \frac{\partial^2}{\partial \theta_i \partial \theta_j} \mathcal{X}_n(i, \theta) \right| \right] = O(1).$$

See, for example, Newey (1991) and Andrews (1992).

Remark 6 Joint convergence (d) holds, for example, when $\mathcal{X}_n(i, \cdot) = 1/\sqrt{n} \sum_{t=1}^n x_t(i)$ where $x_t(i)$ has a zero mean and is square integrable, $\sqrt{n}(\hat{\theta}_n(i) - \theta_0(i)) = 1/\sqrt{n} \sum_{t=1}^n m_t(i) + o_p(1)$ for some zero mean square integrable random variables $m_t(i)$, and $\{x_t(i), m_t(i)\}$ satisfy suitable moment and dependence properties. An example concerning a residuals white noise test is provided in Section 4.

Remark 7 Conditions (b) and (d) allow for non-standard cases. One example is when $\theta_0(i)$ lies on the boundary of $\Theta(i)$ (e.g. Andrews, 1999), and another is when a component of $\theta_0(i)$ is weakly

or non-identified (e.g. Andrews and Cheng, 2012). In each case $\sqrt{n}(\hat{\theta}_n(i) - \theta_0(i)) = \mathcal{M}_n(i) + o_p(1) \xrightarrow{d} \mathcal{M}(i)$ is non-Gaussian.

Theorem 3.1 *Let Assumption 1 hold, and write $\mathcal{D}(i) \equiv \mathcal{D}(i, \theta_0)$ where $\mathcal{D}(i, \theta)$ is defined in Assumption 1.c. For some sequence $\{\mathcal{L}_n\}$ of positive integers that is not unique, with $\mathcal{L}_n \rightarrow \infty$: $|\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i, \hat{\theta}_n(i))| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{S}_n(i) + \mathcal{D}(i)' \mathcal{M}_n(i)|| \xrightarrow{p} 0$ and $|\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{S}_n(i) + \mathcal{D}(i)' \mathcal{M}_n(i)| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{S}(i) + \mathcal{D}(i)' \mathcal{M}(i)|| \xrightarrow{p} 0$, hence $|\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i, \hat{\theta}_n(i))| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{S}(i) + \mathcal{D}(i)' \mathcal{M}(i)|| \xrightarrow{p} 0$.*

4 Illustrations

We now consider maximum statistics that involve unit root test statistics and a residuals white noise test statistic. Both require a non-Gaussian approximation theory, demonstrating the unique applicability of Theorem 2.2.

4.1 Unit Root Tests

Consider unit root tests over a set of processes $\{y_t(i) : i \in \mathbb{N}\}$. Suppose $y_t(i) = \phi_0(i)y_{t-1}(i) + \epsilon_t(i)$, $|\phi_0(i)| \leq 1$, for $i \in \mathbb{N}$. We assume $\{\epsilon_t(i) : i \in \mathbb{N}\}$ lies in a probability measure space $(\Omega, \sigma(\cup_{t \in \mathbb{N}} \mathcal{F}_t), \mathcal{P})$, where $\{\mathcal{F}_t\}_{t \in \mathbb{N}}$ is a sequence of σ -felds. Assume $\epsilon_t(i)$ is \mathcal{F}_t -measurable for each i . Write $\mathcal{F}_s^t \equiv \sigma(\cup_{\tau=s}^t \mathcal{F}_\tau)$. Zero mean $\epsilon_t(i)$ is stationary, and $E|\epsilon_t(i)|^r < \infty$ for some $r > 2$. Moreover, α -mixing coefficients $\alpha_h \equiv \sup_{\mathcal{A} \in \mathcal{F}_{-\infty}^{t-h}, \mathcal{B} \in \mathcal{F}_t^\infty} |\mathcal{P}(\mathcal{A} \cap \mathcal{B}) - \mathcal{P}(\mathcal{A})\mathcal{P}(\mathcal{B})|$ satisfy $\alpha_h = O(h^{r/(r-2)}/\ln(h))$. Thus, $\epsilon_t(i)$ and any measurable function of $\{\epsilon_t(i) : 1 \leq i \leq k\}$ has coefficients α_h . Let $\hat{\phi}_n(i)$ be the least squares estimator.

We want to test the hypothesis that all processes have a unit root $H_0 : \phi_0(i) = 1$ for all $i \in \mathbb{N}$. The proposed test statistic is

$$\mathcal{T}_n(k) \equiv n \max_{1 \leq i \leq k} \left| \hat{\phi}_n(i) - 1 \right|.$$

Define $\sigma^2(i) \equiv \lim_{n \rightarrow \infty} E[(1/\sqrt{n} \sum_{t=1}^n \epsilon_t(i))^2]$ and $\sigma_\epsilon^2(i) \equiv E[\epsilon_t^2(i)]$, and define

$$\mathcal{T}(i) \equiv \frac{1}{2} \left\{ \mathcal{W}(i, 1)^2 - \sigma_\epsilon^2(i)/\sigma^2(i) \right\} / \int_0^1 \mathcal{W}(i, \nu)^2 d\nu$$

where $\{\mathcal{W}(\cdot, \nu) : [0, 1]\}$ are standard Wiener processes. $\mathcal{T}(i)$ is the well known limit law for the least squares estimator when there is a unit root and $\epsilon_t(i)$ is possibly dependent: $n(\hat{\phi}_n(i) - 1) \xrightarrow{d} \mathcal{T}(i)$ (Phillips, 1987, Theorem 3.1). If $\epsilon_t(i)$ is iid then $\sigma_\epsilon^2(i)/\sigma^2(i) = 1$, cf. White (1958).

Apply the mapping theorem to yield $n|\hat{\phi}_n(i) - 1| \xrightarrow{d} |\mathcal{T}(i)|$ for each i under H_0 . Therefore $n|\hat{\phi}_n(i) - 1| - |\mathcal{T}(i)| = o_p(1)$ for each i under H_0 . Now invoke Theorem 2.2 to deduce $|\mathcal{T}_n(\mathcal{L}_n) - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{T}(i)|| \xrightarrow{p} 0$ under H_0 for some non-unique sequence of positive integers $\{\mathcal{L}_n\}$, $\mathcal{L}_n \rightarrow \infty$. Conversely, if $|\phi_0(i^*)| < 1$ for some $i^* \in \mathbb{N}$, then such $n(\phi_0(i^*) - 1) \rightarrow -\infty$ will eventually dominate $\mathcal{T}_n(k) = \max_{1 \leq i \leq k} |n(\hat{\phi}_n(i) - \phi_0(i)) - n(\phi_0(i) - 1)|$ for any $k \geq i^*$. Therefore $\mathcal{T}_n(\mathcal{L}_n) \xrightarrow{p} \infty$ for any sequence of positive integers $\{\mathcal{L}_n\}$, $\mathcal{L}_n \rightarrow \infty$.

The limit law $\mathcal{T}(i)$ is not pivotal since it contains nuisance parameters $\sigma_\epsilon^2(i)$ and $\sigma^2(i)$. Let $\hat{\sigma}_{n,\epsilon}^2(i)$ and $\hat{\sigma}_n^2(i)$ be consistent estimators of $\sigma_\epsilon^2(i)$ and $\sigma^2(i)$ respectively (see Phillips, 1987, Section 4), and define

$$\tilde{\mathcal{T}}_n(k) \equiv \max_{1 \leq i \leq k} \left| n \left(\hat{\phi}_n(i) - \frac{\frac{1}{n} \hat{\sigma}_n^2(i) - \hat{\sigma}_{n,\epsilon}^2(i)}{1/n^2 \sum_{t=2}^n y_{t-1}^2(i)} - 1 \right) \right| = \max_{1 \leq i \leq k} |\tilde{\mathcal{T}}_{n,i}|,$$

where $\tilde{\mathcal{T}}_{n,i}$ is implicitly defined. The term $\tilde{\mathcal{T}}_{n,i}$ was proposed in Phillips (1987, Section 5) as an adjustment that leads to a pivotal asymptotic law.

Denote $\tilde{\mathcal{T}}(i) \equiv (1/2)(\mathcal{W}(i, 1)^2 - 1) / \int_0^1 \mathcal{W}(i, \nu)^2 d\nu$, the now classic limit law for the least squares estimator when there is a unit root and iid error (White, 1958). Theorem 5.1 in Phillips (1987) implies $\tilde{\mathcal{T}}_{n,i} = \tilde{\mathcal{T}}(i) + o_p(1)$. Now use Theorem 2.2 to yield $|\tilde{\mathcal{T}}_n(\mathcal{L}_n) - \max_{i \in \mathcal{I}(\mathcal{L}_n)} \tilde{\mathcal{T}}(i)| \xrightarrow{p} 0$ under H_0 for some non-unique sequence of positive integers $\{\mathcal{L}_n\}$, $\mathcal{L}_n \rightarrow \infty$.

4.2 Residuals White Noise Test

Consider a process $\{y_t\}$ modeled as an AR(p) for finite $p \geq 1$,

$$y_t = c_0 + \sum_{i=1}^p \phi_{0,i} y_{t-i} + \epsilon_t = \theta'_0 x_t + \epsilon_t$$

where $\theta_0 \equiv [c_0, \phi'_0]'$, $x_t \equiv [1, y_{t-1}, \dots, y_{t-p}]'$, and $1 - \sum_{i=1}^p \phi_{0,i} z^i$ has roots outside the unit circle. The AR model is assumed to be pseudo true in the sense that θ_0 is the unique point in the interior of compact Θ that satisfies $E[\epsilon_t x_t] = 0$. Assume $E[y_t^2] > 0$ and $E|y_t|^r < \infty$ for some $r > 4$. Define σ -fields $\mathcal{F}_t \equiv \sigma(y_\tau : \tau \leq t)$ and $\mathcal{F}_s^t \equiv \sigma(y_\tau : s \leq \tau \leq t)$, and assume $\mathcal{F}_{t-1} \subset \mathcal{F}_t \forall t$. Assume y_t is stationary α -mixing with coefficients $\alpha_h \equiv \sup_{\mathcal{A} \subset \mathcal{F}_{-\infty}^{t-m}, \mathcal{B} \subset \mathcal{F}_t^\infty} |P(\mathcal{A} \cap \mathcal{B}) - P(\mathcal{A})P(\mathcal{B})| = O(h^{r/(r-2)}/\ln(h))$. Sufficient conditions for the strong mixing property in linear processes are presented in Gorodetskii (1977) and Withers (1981), amongst others.

We want to test $H_0 : E[\epsilon_t \epsilon_{t-h}] = 0 \forall h \geq 1$. Let $\hat{\theta}_n \equiv [\hat{c}_n, \hat{\phi}'_n]$ be the least squares estimator of θ_0 , and define an error function, and the residual sample serial correlation at lag $h \geq 1$:

$$\epsilon_t(\theta) \equiv y_t - \theta' x_t \quad \text{and} \quad \mathcal{X}_n(i, \theta) \equiv \sqrt{n} \frac{1/n \sum_{t=1+h}^n \epsilon_t(\theta) \epsilon_{t-h}(\theta)}{1/n \sum_{t=1}^n \epsilon_t^2(\theta)}.$$

A valid test can be based on the maximum absolute correlation, $\max_{1 \leq h \leq \mathcal{L}_n} |\mathcal{X}_n(i, \hat{\theta}_n)|$, cf. Hill and Motegi (2017). We only provide a proof linking $\max_{1 \leq h \leq \mathcal{L}_n} |\mathcal{X}_n(i, \hat{\theta}_n)|$ to the maximum of a process that depends on ϵ_t and properties of the plug-in estimator, based on an asymptotic expansion. See Hill and Motegi (2017, Section 2, Theorem 2.5) for an asymptotically valid dependent wild bootstrap based on the expansion.

The following is based on arguments in Hill and Motegi (2017, Lemma 2.1).¹ The proof relies on a standard expansion, and Theorem 2.2. We therefore prove the claim in the supplemental

¹As in the present example, Hill and Motegi (2017) work with a residuals sample correlation in a general parametric regression model setting. Since the form of $\mathcal{X}_n(i, \theta)$ is therefore known, they do not need to verify conditions like Assumption 1, and instead deliver an expansion using a more direct proof.

material Hill (2017). Define

$$\mathcal{D}(h) = -E[\epsilon_t x_{t-h}] - E[\epsilon_{t-h} x_t] \text{ and } z_t(h) \equiv \frac{1}{E[\epsilon_t^2]} \epsilon_t \epsilon_{t-h} - \mathcal{D}(h)' (E[x_t x_t'])^{-1} x_t \epsilon_t$$

$$w_t(\lambda) = \lambda_1 \frac{1}{E[\epsilon_t^2]} \epsilon_{t-h} + \lambda_2 (E[x_t x_t'])^{-1} x_t \text{ for arbitrary } (\lambda_1, \lambda_2) \in \mathbb{R}, \lambda_1^2 + \lambda_2^2 = 1.$$

Theorem 4.1

a. Let H_0 hold and assume $\lim_{n \rightarrow \infty} E(1/\sqrt{n} \sum_{t=1}^n w_t(\lambda) \epsilon_t)^2 > 0$. Moreover, assume for any asymptotic draw $\{y_t, \epsilon_t\}_{t=1}^\infty$ that $\inf_{\theta \in \Theta} |\epsilon_t(\theta)| \geq \iota$ a.s. for all $t \in \mathbb{N}/S$ where S is finite, and for some non-random $\iota > 0$. Assumption 1 applies, and therefore the conclusions of Theorem 3.1 hold. In particular, for some non-unique sequence of positive integers $\{\mathcal{L}_n\}$, $\mathcal{L}_n \rightarrow \infty$:

$$\left| \max_{1 \leq h \leq \mathcal{L}_n} |\mathcal{X}_n(i, \hat{\theta}_n)| - \max_{1 \leq h \leq \mathcal{L}_n} \left| \frac{1}{\sqrt{n}} \sum_{t=1+h}^n z_t(h) \right| \right| \xrightarrow{p} 0. \quad (5)$$

Moreover, $\max_{1 \leq h \leq \mathcal{L}_n} \|1/\sqrt{n} \sum_{t=1+h}^n z_t(h)\| - \max_{1 \leq h \leq \mathcal{L}_n} \|\mathcal{Z}(h)\| \xrightarrow{p} 0$ where $\{\mathcal{Z}(h) : h \in \mathbb{N}\}$ is a zero mean Gaussian process with covariance kernel $E[\mathcal{Z}(h)\mathcal{Z}(\tilde{h})] = \lim_{n \rightarrow \infty} 1/n \sum_{s,t=1}^n E[z_s(h)z_t(\tilde{h})]$ and variance $E[\mathcal{Z}(h)^2] < \infty$.

b. If H_0 is false then $\max_{1 \leq h \leq \mathcal{L}_n} |\mathcal{X}_n(i, \hat{\theta}_n)| \xrightarrow{p} \infty$ for any sequence of positive integers $\{\mathcal{L}_n\}$, $\mathcal{L}_n \rightarrow \infty$.

Remark 8 The bound $\lim_{n \rightarrow \infty} E(1/\sqrt{n} \sum_{t=1}^n w_t(\lambda) \epsilon_t)^2 > 0$ ensures a non-degenerate limit theory for a key joint process arising in a sample correlation first order expansion. The limit is finite by the mixing property and $E|w_t(\lambda) \epsilon_t|^{r/2} < \infty$ where $r > 4$ by assumed L_r -boundedness (Ibragimov, 1975, Theorem 1.7). The assumption $\inf_{\theta \in \Theta} |\epsilon_t(\theta)| \geq \iota > 0$ a.s. for all $t \in \mathbb{N}/S$ and finite S expedites the expansion proof. This is mild since $\mathcal{F}_{t-1} \subset \mathcal{F}_t \forall t$ implies $\inf_{\theta \in \Theta} |\epsilon_t(\theta)| > 0$ a.s. $\forall t$ (see the proof of Theorem 4.1).

Remark 9 The result significantly augments known results in the max-correlation literature by permitting residuals, and without restricting dependence in the limit process $\{\mathcal{Z}(h)\}$. Existing

extreme value theory works with observed data, and imposes conditions that ensure $\sqrt{n}(\hat{\gamma}_n(i) - E[x_t x_{t-i}])$ converges in finite dimensional distributions to a Gaussian law $\mathcal{Z}(i)$, and $\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{Z}(i)|$ converges in law to a well defined random variable (e.g. Xiao and Wu, 2014). The latter requires asymptotic independence $E[\mathcal{Z}(i)\mathcal{Z}(j)] \rightarrow 0$ as $|i - j| \rightarrow \infty$, cf. Leadbetter, Lindgren, and Rootzén (1983, Chapter 6) and Hüsler (1986, 1993). The high dimensional central limit theory literature can tackle $\max_{1 \leq h \leq \mathcal{L}_n} |1/\sqrt{n} \sum_{t=1+h}^n z_t(h)|$, e.g. Chernozhukov, Chetverikov, and Kato (2013a), but not the intermediate step (5) since $z_t(h)$ is generally not Gaussian.

5 Conclusion

We provide a general result for convergence of arrays that permits a new theory for the maximum of an increasing sequence of random variables. When linking the maximum of two random variables $\mathcal{X}_n(i)$ and $\mathcal{Y}_n(i)$, unlike the extreme value theory and high dimensional Gaussian central limit theory literatures, we do not require normality or even asymptotic normality of $\mathcal{Y}_n(i)$. This permits new results for maxima, covering heavy tailed data, non-stationary data, and filtered data where the filter may lead to non-standard asymptotics. Two illustrations are provided covering unit root tests and a residual white noise test, both of which appear to be new. A shortcoming of our general approach, based ultimately on Ramsey (1930) theory and its implications for array convergence, is that we cannot bound the allowed array dimension \mathcal{L}_n as $n \rightarrow \infty$. This runs contrary to the max-correlation literature (e.g. Xiao and Wu, 2014), and the Gaussian coupling literature, most recently punctuated by Chernozhukov, Chetverikov, and Kato (2013a, 2016), where the best known bounds on \mathcal{L}_n are available.

A Appendix: Proofs

Proof of Lemma 2.1. We prove in Step 1 that $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_l), n_l} = 0$ for some sequence of positive integers $\{n_l\}_{l=1}^{\infty}$, $n_l < n_{l+1} \forall l$, and some mapping $\mathcal{L}(n_l) \leq \mathcal{L}(n_{l+1})$, $\mathcal{L}(n_l) \rightarrow \infty$ and $n_l \rightarrow \infty$ as $l \rightarrow \infty$. We use that result in Step 2 to prove the claim.

Step 1. We now prove $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_l), n_l} = 0$. By assumption $\{\mathcal{A}_{k,n} : 1 \leq k \leq \mathcal{I}_n\}_{n \geq 1}$ lies in

(\mathbb{R}, d) , which is a first countable topological space, and $\lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \mathcal{A}_{k,n} = 0$. Therefore, by Lemma 1 in Boehme and Rosenfeld (1974) there exists a sequence of positive integers $\{\mathcal{L}_i\}_{i=1}^\infty$, $\mathcal{L}_i \rightarrow \infty$ as $i \rightarrow \infty$, and an integer mapping $n(\mathcal{L}) \rightarrow \infty$ as $\mathcal{L} \rightarrow \infty$ such that $\lim_{i \rightarrow \infty} \mathcal{A}_{\mathcal{L}_i, n(\mathcal{L}_i)} = 0$. The relation $n(\mathcal{L}) \rightarrow \infty$ as $\mathcal{L} \rightarrow \infty$ holds by construction of the array $\{\mathcal{A}_{k,n} : 1 \leq k \leq \mathcal{I}_n\}_{n \geq 1}$ with $\mathcal{I}_n \rightarrow \infty$ as $n \rightarrow \infty$.

We can always assume monotonicity: $\mathcal{L}_i \leq \mathcal{L}_{i+1} \forall i$. Simply note that $\lim_{i \rightarrow \infty} \mathcal{A}_{\mathcal{L}_i, n(\mathcal{L}_i)} = 0$ implies $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}_{i_l}, n(\mathcal{L}_{i_l})} = 0$ for every infinite subsequence $\{i_l\}_{l \geq 1}$ of $\{i\}_{i \geq 1}$. Since $\mathcal{L}_i \rightarrow \infty$ as $i \rightarrow \infty$, we can find a subsequence $\{i_l^*\}_{l \geq 1}$ such that $i_l^* \leq i_{l+1}^*$ and $\mathcal{L}_{i_l^*} \leq \mathcal{L}_{i_{l+1}^*}$ for each l . This follows from the monotone subsequence theorem, which itself follows from Ramsey's (1930) theorem, cf. Erdős and Szekeres (1935) and Burkill and Mirsky (1973). Now define $\mathcal{L}_l^* \equiv \mathcal{L}_{i_l^*}$, hence $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}_l^*, n(\mathcal{L}_l^*)} = 0$ where $\mathcal{L}_l^* \leq \mathcal{L}_{l+1}^*$ and $\mathcal{L}_l^* \rightarrow \infty$ as $l \rightarrow \infty$.

Now let $\{n_i\}_{i=1}^\infty$ and $\{\mathcal{L}(n_i)\}_{i=1}^\infty$ be any sequences satisfying $n_i = n(\mathcal{L}_i)$ and $\mathcal{L}(n_i) = \mathcal{L}_i$. Hence by the above argument $\mathcal{L}(n_i) \leq \mathcal{L}(n_i + 1)$, $\mathcal{L}(n_i) \rightarrow \infty$ and $n_i \rightarrow \infty$, such that $\lim_{i \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_i), n_i} = 0$. Note that $\lim_{i \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_i), n_i} = 0$ if and only if $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_{i_l}), n_{i_l}} = 0$ for every subsequence $\{n_{i_l}\}_{l=1}^\infty$ of $\{n_i\}_{i=1}^\infty$. Since $n_i \rightarrow \infty$ as $i \rightarrow \infty$, by the monotone subsequence theorem there exists a strictly monotonically increasing subsequence $\{n_{i_l}\}_{l=1}^\infty$. Therefore, as required $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_l), n_l} = 0$ for some sequence of positive integers $\{n_l\}_{l=1}^\infty$, $n_l < n_{l+1} \forall l$, and $\mathcal{L}(n_l) \leq \mathcal{L}(n_{l+1})$, $\mathcal{L}(n_l) \rightarrow \infty$ and $n_l \rightarrow \infty$ as $l \rightarrow \infty$.

Step 2. By assumption $\lim_{n \rightarrow \infty} \mathcal{A}_{k,n} = 0 \forall k$. Therefore:

$$\lim_{s \rightarrow \infty} \mathcal{A}_{k, n_s} = 0 \text{ for every } k \text{ and every infinite subsequence } \{n_s\}_{s \geq 1}. \quad (\text{A.1})$$

Now repeat the Step 1 argument for each $\{\mathcal{A}_{k, n_s}\}_{s \geq 1}$: there exists a strictly monotonically increasing subsequence of positive integers $\{n_{s_l}\}_{l \geq 1}$ and some integer mapping $\mathcal{L}_s(n_{s_l})$ that may depend on s , with $n_{s_l} \rightarrow \infty$ and $\mathcal{L}_s(n_{s_l}) \rightarrow \infty$ as $l \rightarrow \infty \forall s$, such that $\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}_s(n_{s_l}), n_{s_l}} = 0 \forall s$. As above, we may take $\mathcal{L}_s(\cdot)$ to be monotonic: $\mathcal{L}_s(\tilde{n}) \leq \mathcal{L}_s(\tilde{n} + 1) \forall \tilde{n}$.

Since monotonic $\mathcal{L}_s(\tilde{n}) \rightarrow \infty$ as $\tilde{n} \rightarrow \infty \forall s$, there exists an integer mapping $\mathcal{L}(\cdot)$ such that $\mathcal{L}(n) \rightarrow \infty$ as $n \rightarrow \infty$ and for each s , $\limsup_{n \rightarrow \infty} \{\mathcal{L}(n)/\mathcal{L}_s(n)\} < 1$. By monotonicity $\mathcal{A}_{k,n} \leq$

$\mathcal{A}_{k+1,n}$ this mapping satisfies

$$\lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_{s_l}), n_{s_l}} \leq \lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}_s(n_{s_l}), n_{s_l}} = 0 \quad \forall s. \quad (\text{A.2})$$

Notice $\mathcal{L}(\cdot)$ is not unique: for any $\mathcal{L}(\cdot)$ that satisfies (A.2) there exists $\tilde{\mathcal{L}}(n) \rightarrow \infty$ such that $\limsup_{n \rightarrow \infty} \tilde{\mathcal{L}}(n)/\mathcal{L}(n) < 1$, hence by monotonicity $\lim_{l \rightarrow \infty} \mathcal{A}_{\tilde{\mathcal{L}}(n_{s_l}), n_{s_l}} \leq \lim_{l \rightarrow \infty} \mathcal{A}_{\mathcal{L}(n_{s_l}), n_{s_l}} = 0$.

Now write $\mathcal{B}_n \equiv \mathcal{A}_{\mathcal{L}(n), n}$. By a direct implication of (A.1) and (A.2), for every subsequence $\{\mathcal{B}_{n_s}\}_{s \geq 1}$ there exists a further subsequence $\{\mathcal{B}_{n_{s_l}}\}_{l \geq 1}$ that converges $\lim_{l \rightarrow \infty} \mathcal{B}_{n_{s_l}} = 0$. Therefore $\lim_{n \rightarrow \infty} \mathcal{B}_n = 0$ (see Royden, 1988, p. 39). This proves $\lim_{n \rightarrow \infty} \mathcal{A}_{\mathcal{L}_n, n} = 0$ with $\mathcal{L}_n = \mathcal{L}(n)$ as required. *QED*.

Proof of Theorem 2.2.

Claim (a). By assumption each $\mathcal{X}_n(i) \xrightarrow{p} 0$ and therefore $\max_{1 \leq i \leq k} |\mathcal{X}_n(i)| \xrightarrow{p} 0$. Define $\mathcal{A}_{k,n} \equiv 1 - \exp\{-\max_{1 \leq i \leq k} |\mathcal{X}_n(i)|\}$ and $\mathcal{P}_{k,n} \equiv \int_0^\infty P(\mathcal{A}_{k,n} > \epsilon) d\epsilon$. By construction $\mathcal{A}_{k,n} \in [0, 1]$ *a.s.* $\forall k$. Lebesgue's dominated convergence theorem, and $\mathcal{A}_{k,n} \xrightarrow{p} 0$, therefore yield for each k :

$$\lim_{n \rightarrow \infty} \mathcal{P}_{k,n} = \lim_{n \rightarrow \infty} \int_0^\infty P(\mathcal{A}_{k,n} > \epsilon) d\epsilon = \lim_{n \rightarrow \infty} \int_0^1 P(\mathcal{A}_{k,n} > \epsilon) d\epsilon = \int_0^1 \lim_{n \rightarrow \infty} P(\mathcal{A}_{k,n} > \epsilon) d\epsilon = 0.$$

Now apply Lemma 2.1 to $\mathcal{P}_{k,n}$ to deduce that there exists a positive integer sequence $\{\mathcal{L}_n\}$ that is not unique, $\mathcal{L}_n \rightarrow \infty$ and $\mathcal{L}_n = o(n)$, such that $\lim_{n \rightarrow \infty} \mathcal{P}_{\mathcal{L}_n, n} = \lim_{n \rightarrow \infty} \int_0^1 P(\mathcal{A}_{\mathcal{L}_n, n} > \epsilon) d\epsilon = 0$. Therefore, by construction $E[\mathcal{A}_{\mathcal{L}_n, n}] = \int_0^1 P(\mathcal{A}_{\mathcal{L}_n, n} > \epsilon) d\epsilon \rightarrow 0$. Hence $\mathcal{A}_{\mathcal{L}_n, n} \xrightarrow{p} 0$ by Markov's inequality, which yields $\max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| \xrightarrow{p} 0$ as claimed.

The sequence $\{\mathcal{L}_n\}$ is not unique for either of the following reasons: (i) the probability limit is asymptotic hence we can always change \mathcal{L}_n for finitely many n ; and (ii) by monotonicity of $\max_{1 \leq i \leq k} |\mathcal{X}_n(i)|$ any other $\{\mathring{\mathcal{L}}_n\}$ that satisfies $\mathring{\mathcal{L}}_n \rightarrow \infty$ and $\limsup_{n \rightarrow \infty} \{\mathring{\mathcal{L}}_n/\mathcal{L}_n\} < 1$ satisfies $\max_{1 \leq i \leq \mathring{\mathcal{L}}_n} |\mathcal{X}_n(i)| \leq \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| \xrightarrow{p} 0$ as $n \rightarrow \infty$.

Claim (b). Apply the triangle inequality twice to yield both $\max_{1 \leq i \leq k} |\mathcal{X}_n(i)| \leq \max_{1 \leq i \leq k} |\mathcal{Y}_n(i)| + \max_{1 \leq i \leq k} |\mathcal{X}_n(i) - \mathcal{Y}_n(i)|$ and $\max_{1 \leq i \leq k} |\mathcal{Y}_n(i)| \leq \max_{1 \leq i \leq k} |\mathcal{X}_n(i)| + \max_{1 \leq i \leq k} |\mathcal{X}_n(i) - \mathcal{Y}_n(i)|$,

hence

$$\left| \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i)| - \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{Y}_n(i)| \right| \leq \max_{1 \leq i \leq \mathcal{L}_n} |\mathcal{X}_n(i) - \mathcal{Y}_n(i)|$$

Now apply (a) to $\mathcal{X}_n(i) - \mathcal{Y}_n(i)$ to yield the desired result. \mathcal{QED} .

Proof of Theorem 3.1. By the mean value theorem, there exists $\theta_n^*(i)$, $\|\theta_n^*(i) - \theta_0(i)\| \leq \|\hat{\theta}_n(i) - \theta_0(i)\|$, such that:

$$\mathcal{X}_n(i, \hat{\theta}_n(i)) = \mathcal{X}_n(i, \theta_0(i)) + \frac{1}{\sqrt{n}} \frac{\partial}{\partial \theta'} \mathcal{X}_n(i, \theta_n^*(i)) \sqrt{n} (\hat{\theta}_n(i) - \theta_0(i)).$$

Under Assumption 1.d, $\sqrt{n}(\hat{\theta}_n(i) - \theta_0(i)) = \mathcal{M}_n(i) + o_p(1) = O_p(1)$, hence $\|\theta_n^*(i) - \theta_0(i)\| \leq \|\hat{\theta}_n(i) - \theta_0(i)\| \xrightarrow{p} 0$. Therefore $\theta_n^*(i)$ lies in any compact neighborhood $\mathcal{N}_0(i)$ of $\theta_0(i)$ with positive Lebesgue measure asymptotically with probability approaching one. Now use Assumption 1.c and continuity of $\mathcal{D}(i, \cdot)$ to yield $n^{-1/2}(\partial/\partial \theta) \mathcal{X}_n(i, \theta_n^*(i)) \xrightarrow{p} \mathcal{D}(i) \equiv \mathcal{D}(i, \theta_0)$. Further, by Assumption 1.d $\mathcal{X}_n(i, \theta_0(i)) = \mathcal{S}_n(i) + o_p(1)$. We may therefore write

$$\mathcal{X}_n(i, \hat{\theta}_n(i)) = \mathcal{S}_n(i) + \mathcal{D}(i)' \mathcal{M}_n(i) + o_p(1). \tag{A.3}$$

Apply Assumption 1,d and the continuous mapping theorem to deduce $\mathcal{S}_n(i) + \mathcal{D}(i) \mathcal{M}_n(i) \xrightarrow{d} \mathcal{S}(i) + \mathcal{D}(i)' \mathcal{M}(i)$. By the definition of convergence in distribution, we may write for each i :

$$\mathcal{S}_n(i) + \mathcal{D}(i)' \mathcal{M}_n(i) = \mathcal{S}(i) + \mathcal{D}(i)' \mathcal{M}(i) + o_p(1). \tag{A.4}$$

The claim now follows from (A.3) and (A.4), and two applications of Theorem 2.2. \mathcal{QED} .

References

- ANDREWS, D. W. K. (1992): “Generic Uniform Convergence,” *Econometric Theory*, 8, 241–257.
- (1999): “Estimation When a Parameter is on a Boundary,” *Econometrica*, 67, 1341–1383.
- ANDREWS, D. W. K., AND X. CHENG (2012): “Estimation and Inference with Weak, Semi-Strong and Strong Identification,” *Econometrica*, 80, 2153–2211.
- BERMAN, S. M. (1964): “Limit theorems for the maximum term in stationary sequences,” *Annals of Mathematical Statistics*, 35, 502–516.

- BOEHME, T. K., AND M. ROSENFELD (1974): “An Example of Two Compact Hausdorff Frechet Spaces Whose Product is not Frechet,” *Journal of the London Mathematical Society*, 8, 339–344.
- BURKILL, H., AND L. MIRSKY (1973): “Monotonicity,” *Journal of Multivariate Analysis and Applications*, 35, 502–516.
- CHERNOZHUKOV, V., D. CHETVERIKOV, AND K. KATO (2013a): “Gaussian Approximations and Multiplier Bootstrap for Maxima of Sums of High-Dimensional Random Vectors,” *Annals of Statistics*, 41, 2786–2819.
- (2013b): “Supplemental Material for ‘Gaussian Approximations and Multiplier Bootstrap for Maxima of Sums of High-Dimensional Random Vectors’,” Massachusetts Institute of Technology.
- (2015): “Comparison and Anti-Concentration Bounds for Maxima of Gaussian Random Vectors,” *Probability Theory and Related Fields*, 162, 47–70.
- (2016): “Central Limit Theorems and Bootstrap in High Dimensions,” *Annals of Probability*, forthcoming.
- DUDLEY, R. M. (1978): “Central Limit Theorems for Empirical Measures,” *Annals of Probability*, 6, 899–929.
- (1984): “A Course on Empirical Processes,” in *Ecole Eté de Probabilités de St. Flour, Lecture Notes in Math*, vol. 1097, pp. 1–142. Springer, New York.
- DUDLEY, R. M., AND W. PHILIPP (1983): “Invariance Principles for Sums of Banach Space Valued Random Elements and Empirical Processes,” *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 62, 509–552.
- ERDÖS, P., AND G. SZEKERES (1935): “A combinatorial problem in geometry,” *Compositio Mathematica*, 2, 463–470.
- FALK, M., J. HÜSLER, AND R.-D. REISS (2011): *Law of Small Numbers: Extremes and Rare Events*. Birkhäuser, Berlin, 3 edn.
- GORODETSKII, V. V. (1977): “On the Strong Mixing Property for Linear Processes,” *Theory of Probability and its Applications*, 22, 411–413.
- GÖTZE, F. (1991): “On the Rate of Convergence in the Multivariate Central Limit Theorem,” *Annals of Probability*, 19, 724–739.
- HANNAN, E. J. (1974): “The Uniform Convergence of Autocovariances,” *Annals of Statistics*, 2, 803–806.
- HILL, J. B. (2017): “Supplemental Material for ‘Asymptotic Theory for the Maximum of an Increasing Sequence of Parametric Functions’,” .
- HILL, J. B., AND K. MOTEGI (2017): “A Max-Correlation White Noise Test for Weakly Dependent Time Series,” Discussion paper, University of North Carolina - Chapel Hill.
- HOFFMANN-JORGENSEN, J. (1984): “Convergence of Stochastic Processes on Polish Spaces,”

- mimeo.
- (1991): “Convergence of Stochastic Processes on Polish Spaces,” in *Various Publications Series 39*. Aarhus Universitet, Aarhus, Denmark, mimeo.
- HOMBLE, P., AND P. MCCORMICK (1995): “Weak Limit Results for the Extremes of a Class of Shot Noise Processes,” *Journal of Applied Probability*, 32, 707–726.
- HÜSLER, J. (1986): “Extreme Values of Non-Stationary Random Sequences,” *Journal of Applied Probability*, 23, 937–950.
- (1993): “Extremes: Limit Results for Univariate and Multivariate Nonstationary Sequences,” in *Extreme Value Theory and Applications*, ed. by J. Galambos, J. Lechner, and E. Simiu, pp. 283–204. Proceedings on the Conference on Extreme Value Theorem and Applications, Volume I, Gaithersburg, Maryland, Kluwer Academic Publishers.
- IBRAGIMOV, I. A. (1975): “A Note on the Central Limit Theorems for Dependent Random Variables,” *Theory of Probability and its Applications*, 20, 135–141.
- JIRAK, M. (2011): “On the Maximum of Covariance Estimators,” *Journal of Multivariate Analysis*, 102, 1032–1046.
- LE CAM, L. (1988): “On the Prohorov distance between the empirical process and the associated Gaussian bridge,” Discussion paper, University of California - Berkely, Dept. of Statistics, UC-Berkeley.
- LEADBETTER, M. R., G. LINDGREN, AND H. ROOTZÉN (1983): *Extremes and Related Properties of Random Sequences and Processes*, Springer Series in Statistics. Springer-Verlag, New York.
- NEWBY, W. K. (1991): “Uniform Convergence in Probability and Stochastic Equicontinuity,” *Econometrica*, 59, 1161–1167.
- PHILLIPS, P. C. B. (1987): “Time Series with a Unit Root,” *Econometrica*, 55, 277–301.
- POLLARD, D. (1984): *Convergence of Stochastic Processes*. Springer, New York.
- (1990): “Empirical Processes: Theory and Applications,” in *NSF-CBMS Regional Conference Series in Probability and Statistics*, vol. II. Institute of Mathematical Statistics and the American Statistical Association.
- (2002): *A Users Guide to Measure Theoretic Probability*. *Cambridge Series in Statistical and Probabilistic Mathematics 8*. Cambridge Univ. Press, Cambridge.
- PORTNOY, S. (1985): “Asymptotic Behavior of M Estimators of p Regression Parameters when p^2/n is Large; II,” *Annals of Statistics*, 13, 1403–1417.
- (1986): “On the Central Limit Theorem in R^p when $p \rightarrow \infty$,” *Probability Theory and Related Fields*, 73, 571–583.
- RAMSEY, F. P. (1930): “On a Problem of Formal Logic,” *Proceedings of the London Mathematical Society*, 30, 264–286.

- ROYDEN, H. L. (1988): *Real Analysis*. Prentise Hall, 3rd edn.
- WHITE, J. S. (1958): “The Limiting Distribution of the Serial Correlation Coefficient in the Explosive Case,” *Annals of Mathematical Statistics*, 29, 1188–1197.
- WITHERS, C. S. (1981): “Conditions for Linear Processes to be Strong Mixing,” *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 57, 477–480.
- XIAO, H., AND W. B. WU (2014): “Portmanteau Test and Simultaneous Inference for Serial Covariances,” *Statistica Sinica*, 24, 577–599.
- YURINSKII, V. V. (1977): “On the Error of the Gaussian Approximation for Convolutions,” *Theory of Probability and its Applications*, 22, 236–247.