

D-wave heavy baryons of the $SU(3)$ flavor $\mathbf{6}_F$

Qiang Mao^{1,2}, Hua-Xing Chen^{2,*}, Atsushi Hosaka^{3,4,†}, Xiang Liu^{5,6,‡} and Shi-Lin Zhu^{7,8,§}

¹*Department of Electrical and Electronic Engineering, Suzhou University, Suzhou 234000, China*

²*School of Physics and Beijing Key Laboratory of Advanced Nuclear Materials and Physics, Beihang University, Beijing 100191, China*

³*Research Center for Nuclear Physics, Osaka University, Ibaraki 567-0047, Japan*

⁴*Advanced Science Research Center, Japan Atomic Energy Agency, Tokai, Ibaraki, 319-1195 Japan*

⁵*School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China*

⁶*Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China*

⁷*School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China*

⁸*Collaborative Innovation Center of Quantum Matter, Beijing 100871, China*

⁹*Center of High Energy Physics, Peking University, Beijing 100871, China*

We use the method of QCD sum rules to study the D -wave charmed and bottom baryons of the $SU(3)$ flavor $\mathbf{6}_F$, and calculate their masses up to the order $\mathcal{O}(1/m_Q)$ with m_Q the heavy quark mass. Our results suggest that the $\Xi_c(3123)$ can be well interpreted as a D -wave $\Xi'_c(\mathbf{6}_F)$ state, and it probably has a partner state close to it. Our results also suggest that there may exist as many as four D -wave Ω_c states in the energy region $3.3 \sim 3.5$ GeV, and we propose to search for them in the further LHCb and BelleII experiments.

PACS numbers: 14.20.Lq, 12.38.Lg, 12.39.Hg

Keywords: excited heavy baryons, QCD sum rule, heavy quark effective theory

I. INTRODUCTION

In the past years important experimental progresses have been made in the field of heavy baryons [1], and we refer to reviews [2–7] for their recent progress. Especially, the LHCb experiment observed as many as five excited Ω_c states at the same time [8], which is probably related to the fine structure of the strong interaction. Some of these excited Ω_c states can be interpreted as P -wave states [9], inspiring us to further study the D -wave heavy baryons. Actually, in Ref. [10] we have systematically studied the D -wave heavy baryons of the $SU(3)$ flavor $\mathbf{\bar{3}}_F$ using the method of QCD sum rules [11, 12] in the framework of heavy quark effective theory (HQET) [13–15], and in the present study we shall follow the same approach to study the $SU(3)$ flavor $\mathbf{6}_F$ ones, including the D -wave Ω_c states.

There have been lots of heavy baryons observed in various experiments [1, 16–18]. Among them, the $\Xi_c(3123)$ observed by the BaBar Collaboration [19] (but not seen in the following Belle experiment [20]) is a good candidate of the D -wave $\Xi'_c(\mathbf{6}_F)$ state, and has been investigated using many phenomenological methods/models, including various quark models [21–26], the Faddeev method [27], the Regge trajectory [28], the relativistic flux tube model [29], the heavy hadron chiral perturbation theory [30], and QCD sum rules [31, 32], etc. More discussions can be found in Refs. [33–37], and we again refer to reviews [2–7] for the recent progress.

We have studied the heavy baryons using the method of QCD sum rules within HQET [9, 10, 38, 40], and in the present study we shall further study the D -wave charmed baryons of the $SU(3)$ flavor $\mathbf{6}_F$. We shall take the $\mathcal{O}(1/m_Q)$ corrections (m_Q is the heavy quark mass) into account during our QCD sum rule analyses, and extract the chromomagnetic splitting within the same baryon multiplet. More discussions on heavy mesons and baryons containing a single heavy quark can be found in Refs. [43–57].

This paper is organized as follows. In Sec. II, we briefly introduce how we use the D -wave heavy baryon interpolating fields to perform QCD sum rule analyses, and we refer interested readers to consult the discussion in Refs. [9, 10, 38, 40] for details. Then we perform numerical analyses in Sec. III, and offer a short summary in Sec. IV.

II. QCD SUM RULE ANALYSES

In Ref. [10] we have systematically constructed all the D -wave heavy baryon interpolating fields, but just studied the $SU(3)$ flavor $\mathbf{\bar{3}}_F$ ones using the method of QCD sum rules in the framework of heavy quark effective theory (HQET). In this paper we use the same method to study the rest ones of the $SU(3)$ flavor $\mathbf{6}_F$, as briefly shown in Fig. 1. We refer interested readers to consult the discussion in Ref. [10] for details.

Here we briefly explain our notations: $J_{j,P,F;j_l,s_l,\rho-\lambda}^{\alpha_1 \cdots \alpha_{j-1/2}}$ denotes the D -wave heavy baryon field belonging to the baryon multiplet $[F, j_l, s_l, \rho - \lambda]$, where j , P , and F denote its total angular momentum, parity and $SU(3)$ flavor representation ($\mathbf{\bar{3}}_F$ or $\mathbf{6}_F$); j_l and s_l are the total an-

*Electronic address: hxchen@buaa.edu.cn

†Electronic address: hosaka@rcnp.osaka-u.ac.jp

‡Electronic address: xiangliu@lzu.edu.cn

§Electronic address: zhusl@pku.edu.cn

$$\begin{aligned}
& [\rho\rho] \quad l_\rho = 2 \text{ (S)}, \quad l_\lambda = 0, \quad \bar{\mathbf{3}}_C \text{ (A)} \\
& \quad \bar{\mathbf{3}}_F \text{ (A): } L = 2 \otimes s_l = 0 \text{ (A)} \\
& \quad \mathbf{6}_F \text{ (S): } L = 2 \otimes s_l = 1 \text{ (S)} \\
& \quad \left\{ \begin{array}{l} j_l = 1: \Sigma_{c1} \left(\frac{1}{2}^+, \frac{3}{2}^+ \right) \quad \Xi_{c1}' \left(\frac{1}{2}^+, \frac{3}{2}^+ \right) \quad \Omega_{c1} \left(\frac{1}{2}^+, \frac{3}{2}^+ \right) \quad (\rho\rho\text{-b}) [\mathbf{6}_F, 1, 1, \rho\rho] \\ j_l = 2: \Sigma_{c2} \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad \Xi_{c2}' \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad \Omega_{c2} \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad (\rho\rho\text{-c}) [\mathbf{6}_F, 2, 1, \rho\rho] \\ j_l = 3: \Sigma_{c3} \left(\frac{5}{2}^+, \frac{7}{2}^+ \right) \quad \Xi_{c3}' \left(\frac{5}{2}^+, \frac{7}{2}^+ \right) \quad \Omega_{c3} \left(\frac{5}{2}^+, \frac{7}{2}^+ \right) \quad (\rho\rho\text{-d}) [\mathbf{6}_F, 3, 1, \rho\rho] \end{array} \right. \\
& [\lambda\lambda] \quad l_\rho = 0 \text{ (S)}, \quad l_\lambda = 2, \quad \bar{\mathbf{3}}_C \text{ (A)} \\
& \quad \bar{\mathbf{3}}_F \text{ (A): } L = 2 \otimes s_l = 0 \text{ (A)} \\
& \quad \mathbf{6}_F \text{ (S): } L = 2 \otimes s_l = 1 \text{ (S)} \\
& \quad \left\{ \begin{array}{l} j_l = 1: \Sigma_{c1} \left(\frac{1}{2}^+, \frac{3}{2}^+ \right) \quad \Xi_{c1}' \left(\frac{1}{2}^+, \frac{3}{2}^+ \right) \quad \Omega_{c1} \left(\frac{1}{2}^+, \frac{3}{2}^+ \right) \quad (\lambda\lambda\text{-b}) [\mathbf{6}_F, 1, 1, \lambda\lambda] \\ j_l = 2: \Sigma_{c2} \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad \Xi_{c2}' \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad \Omega_{c2} \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad (\lambda\lambda\text{-c}) [\mathbf{6}_F, 2, 1, \lambda\lambda] \\ j_l = 3: \Sigma_{c3} \left(\frac{5}{2}^+, \frac{7}{2}^+ \right) \quad \Xi_{c3}' \left(\frac{5}{2}^+, \frac{7}{2}^+ \right) \quad \Omega_{c3} \left(\frac{5}{2}^+, \frac{7}{2}^+ \right) \quad (\lambda\lambda\text{-d}) [\mathbf{6}_F, 3, 1, \lambda\lambda] \end{array} \right. \\
& [\rho\lambda] \quad l_\rho = 1 \text{ (A)}, \quad l_\lambda = 1, \quad \bar{\mathbf{3}}_C \text{ (A)} \\
& \quad \mathbf{6}_F \text{ (S): } L = 2 \otimes s_l = 0 \text{ (A)} \quad \text{---} \quad j_l = 2: \Sigma_{c2} \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad \Xi_{c2}' \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad \Omega_{c2} \left(\frac{3}{2}^+, \frac{5}{2}^+ \right) \quad (\rho\lambda\text{-a}) [\mathbf{6}_F, 2, 0, \rho\lambda] \\
& \quad \bar{\mathbf{3}}_F \text{ (A): } L = 2 \otimes s_l = 1 \text{ (S)} \\
& \quad L = 0/1 : \text{ we do not study these two cases in this paper.}
\end{aligned}$$

FIG. 1: The notations for the D -wave charmed baryons of the $SU(3)$ flavor $\mathbf{6}_F$. See Fig. 1 of Ref. [10] and discussions therein for details.

gular momentum and spin angular momentum of its light components, respectively; there are three types ($\rho - \lambda$): $\rho\rho$ -type ($l_\rho = 2$ and $l_\lambda = 0$), $\lambda\lambda$ -type ($l_\rho = 0$ and $l_\lambda = 2$) and $\rho\lambda$ -type ($l_\rho = 1$ and $l_\lambda = 1$), where we use l_ρ to denote the orbital angular momentum between the two light quarks, and l_λ to denote the orbital angular momentum between the heavy quark and the two-light-quark system. These parameters satisfy $L = l_\lambda \otimes l_\rho = 2$ (note that we only investigate the D -wave heavy baryons in the present study), $j_l = L \otimes s_l$ and $j = j_l \otimes s_Q = |j_l \pm 1/2|$, with $s_Q = 1/2$ the spin of the heavy quark.

The explicit forms of $J_{j,P,F,j_l,s_l,\rho-\lambda}^{\alpha_1 \cdots \alpha_{j-1/2}}$ have been given in Eqs. (2–19) of Ref. [10]. We use them to perform QCD sum rule analyses by assuming their coupling to the state $|j, P, F, j_l, s_l, \rho - \lambda\rangle$ to be

$$\begin{aligned}
\langle 0 | J_{j,P,F,j_l,s_l,\rho-\lambda}^{\alpha_1 \cdots \alpha_{j-1/2}} | j, P, F, j_l, s_l, \rho - \lambda \rangle \\
= f_{F,j_l,s_l,\rho-\lambda} u^{\alpha_1 \cdots \alpha_{j-1/2}}.
\end{aligned} \tag{1}$$

Again, we recommend interested readers to consult Refs. [9, 10, 38–42] for details, but simply list here the equation to evaluate the mass of the heavy baryon belonging to the multiplet $[F, j_l, s_l, \rho - \lambda]$:

$$m_{j,P,F,j_l,s_l,\rho-\lambda} = m_Q + \bar{\Lambda}_{F,j_l,s_l,\rho-\lambda} + \delta m_{j,P,F,j_l,s_l,\rho-\lambda}. \tag{2}$$

Here m_Q is the heavy quark mass; $\bar{\Lambda}_{F,j_l,s_l,\rho-\lambda} = \bar{\Lambda}_{|j_l-1/2|,P,F,j_l,s_l,\rho-\lambda} = \bar{\Lambda}_{j_l+1/2,P,F,j_l,s_l,\rho-\lambda}$ is the sum rule result obtained at the leading order; $\delta m_{j,P,F,j_l,s_l,\rho-\lambda}$

is the sum rule result obtained at the $\mathcal{O}(1/m_Q)$ order:

$$\begin{aligned}
\delta m_{j,P,F,j_l,s_l,\rho-\lambda} \\
= -\frac{1}{4m_Q} (K_{F,j_l,s_l,\rho-\lambda} + d_M C_{mag} \Sigma_{F,j_l,s_l,\rho-\lambda}),
\end{aligned} \tag{3}$$

where $C_{mag}(m_Q/\mu) = [\alpha_s(m_Q)/\alpha_s(\mu)]^{3/\beta_0}$ with $\beta_0 = 11 - 2n_f/3$, and the coefficient $d_M \equiv d_{j,j_l}$ is defined to be

$$\begin{aligned}
d_{j_l-1/2,j_l} &= 2j_l + 2, \\
d_{j_l+1/2,j_l} &= -2j_l,
\end{aligned} \tag{4}$$

which is directly related to the baryon mass splitting within the same multiplet.

In Ref. [10] we have systematically studied the D -wave heavy baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ using the method of QCD sum rules with HQET. In this paper we similarly study the $SU(3)$ flavor $\mathbf{6}_F$ ones, and calculate the analytical formulae for $\bar{\Lambda}_{F,j_l,s_l,\rho-\lambda}$, $K_{F,j_l,s_l,\rho-\lambda}$ and $\Sigma_{F,j_l,s_l,\rho-\lambda}$. There are altogether seven heavy baryon multiplets of the $SU(3)$ flavor $\mathbf{6}_F$ as shown in Fig. 1, i.e., $[\mathbf{6}_F, 1, 1, \rho\rho]$, $[\mathbf{6}_F, 2, 1, \rho\rho]$, $[\mathbf{6}_F, 3, 1, \rho\rho]$, $[\mathbf{6}_F, 1, 1, \lambda\lambda]$, $[\mathbf{6}_F, 2, 1, \lambda\lambda]$, $[\mathbf{6}_F, 3, 1, \lambda\lambda]$ and $[\mathbf{6}_F, 2, 0, \rho\lambda]$. Hence, there can be as many as two $j^P = 1/2^+$ D -wave excited Σ_c states, five $j^P = 3/2^+$ ones, five $j^P = 5/2^+$ ones, and two $j^P = 7/2^+$ ones. The numbers of excited Ξ'_c and Ω_c states are the same. Recalling that the LHCb experiment observed as many as five excited Ω_c states [8], some of these D -wave heavy baryons may be observed experimentally at the same time. Theoretically, in the present

study we can only use five baryon multiplets to perform sum rule analyses, because in Ref. [10] we failed to construct the currents belonging to the other two multiplets $[\mathbf{6}_F, 2, 1, \rho\rho]$ and $[\mathbf{6}_F, 2, 1, \lambda\lambda]$.

As an example, we use the charmed baryon multiplet $[\Sigma_c(\mathbf{6}_F), 1, 1, \rho\rho]$ to perform QCD sum rule analyses. It contains two charmed baryons, $\Sigma_c(1/2^+)$ and $\Sigma_c(3/2^+)$, and the relevant interpolating field is

$$\begin{aligned} J_{1/2^+, \Sigma_c, 1, 1, \rho\rho}(x) &= \epsilon_{abc} \left([\mathcal{D}_{\mu_1}^t \mathcal{D}_{\mu_2}^t u^{aT}(x)] \mathbb{C} \gamma_{\mu_3}^t d^b(x) \right. \\ &\quad - 2[\mathcal{D}_{\mu_1}^t u^{aT}(x)] \mathbb{C} \gamma_{\mu_3}^t [\mathcal{D}_{\mu_2}^t d^b(x)] \\ &\quad \left. + u^{aT}(x) \mathbb{C} \gamma_{\mu_3}^t [\mathcal{D}_{\mu_1}^t \mathcal{D}_{\mu_2}^t d^b(x)] \right) \\ &\quad \times \left(g_t^{\mu_1\mu_3} g_t^{\mu_2\mu_4} + g_t^{\mu_2\mu_3} g_t^{\mu_1\mu_4} \right) \times \gamma_{\mu_4}^t \gamma_5 h_v^c(x). \end{aligned} \quad (5)$$

We can use this current to perform QCD sum rule analyses, and calculate $\bar{\Lambda}_{\Sigma_c, 1, 1, \rho\rho}$, $K_{\Sigma_c, 1, 1, \rho\rho}$ and $\Sigma_{\Sigma_c, 1, 1, \rho\rho}$:

$$\Pi_{\Sigma_c, 1, 1, \rho\rho} = f_{\Sigma_c, 1, 1, \rho\rho}^2 e^{-2\bar{\Lambda}_{\Sigma_c, 1, 1, \rho\rho}/T} \quad (6)$$

$$= \int_0^{\omega_c} \left[\frac{11}{80640\pi^4} \omega^9 - \frac{19\langle g_s^2 GG \rangle}{3072\pi^4} \omega^5 \right] e^{-\omega/T} d\omega, \quad (7)$$

$$f_{\Sigma_c, 1, 1, \rho\rho}^2 K_{\Sigma_c, 1, 1, \rho\rho} e^{-2\bar{\Lambda}_{\Sigma_c, 1, 1, \rho\rho}/T} = \int_0^{\omega_c} \left[-\frac{59}{2217600\pi^4} \omega^{11} + \frac{299\langle g_s^2 GG \rangle}{161280\pi^4} \omega^7 \right] e^{-\omega/T} d\omega, \quad (8)$$

$$f_{\Sigma_c, 1, 1, \rho\rho}^2 \Sigma_{\Sigma_c, 1, 1, \rho\rho} e^{-2\bar{\Lambda}_{\Sigma_c, 1, 1, \rho\rho}/T} = \int_0^{\omega_c} \left[\frac{37\langle g_s^2 GG \rangle}{322560\pi^4} \omega^7 \right] e^{-\omega/T} d\omega.$$

Sum rules for Ξ'_c and Ω_c belonging to the same multiplet, $[\mathbf{6}_F, 1, 1, \rho\rho]$, as well as sum rules for the other four multiplets, $[\mathbf{6}_F, 3, 1, \rho\rho]$, $[\mathbf{6}_F, 1, 1, \lambda\lambda]$, $[\mathbf{6}_F, 3, 1, \lambda\lambda]$ and $[\mathbf{6}_F, 2, 0, \rho\lambda]$, are listed in Appendix. A. In the next section we shall use these equations to perform numerical analyses.

III. NUMERICAL ANALYSES

We use the following values for various condensates and parameters to perform numerical analyses [1, 58–65]:

$$\begin{aligned} \langle \bar{q}q \rangle &= -(0.24 \pm 0.01)^3 \text{ GeV}^3, \\ \langle \bar{s}s \rangle &= 0.8 \times \langle \bar{q}q \rangle, \\ \langle g_s^2 GG \rangle &= (0.48 \pm 0.14) \text{ GeV}^4, \\ \langle g_s \bar{q}\sigma G q \rangle &= M_0^2 \times \langle \bar{q}q \rangle, \\ \langle g_s \bar{s}\sigma G s \rangle &= M_0^2 \times \langle \bar{s}s \rangle, \\ m_s &= 0.125 \text{ GeV}, \\ M_0^2 &= 0.8 \text{ GeV}^2. \end{aligned} \quad (9)$$

Besides them, we also need the *charm* quark mass, for which we use the PDG value $m_c = 1.275 \pm 0.025 \text{ GeV}$ [1] in the $\overline{\text{MS}}$ scheme.

There are two free parameters in Eqs. (6–8): the Borel mass T and the threshold value ω_c . We follow Ref. [10] and use three criteria to constrain them:

1. The first criterion requires that the high-order corrections should be less than 10%:

$$\text{Convergence (CVG)} \equiv \left| \frac{\Pi_{F, j_l, s_l, \rho-\lambda}^{\text{high-order}}(\infty, T)}{\Pi_{F, j_l, s_l, \rho-\lambda}(\infty, T)} \right| \leq 10\%, \quad (10)$$

where $\Pi_{F, j_l, s_l, \rho-\lambda}^{\text{high-order}}(\omega_c, T)$ denotes the high-order corrections, for example,

$$\Pi_{\Sigma_c, 1, 1, \rho\rho}^{\text{high-order}}(\omega_c, T) = \int_0^{\omega_c} \left[-\frac{19\langle g_s^2 GG \rangle}{3072\pi^4} \omega^5 \right] e^{-\omega/T} d\omega. \quad (11)$$

2. The second criterion requires that the pole contribution (PC) should be larger than 10%:

$$\text{Pole Contribution (PC)} \equiv \frac{\Pi_{F, j_l, s_l, \rho-\lambda}(\omega_c, T)}{\Pi_{F, j_l, s_l, \rho-\lambda}(\infty, T)} \geq 10\%. \quad (12)$$

We can use the first and second criteria together to arrive at an interval $T_{\min} < T < T_{\max}$ for a fixed threshold value ω_c .

3. The third criterion requires that the dependence of $m_{j, P, F, j_l, s_l, \rho-\lambda}$ on the threshold value ω_c should be weak.

Still use the baryon multiplet $[\Sigma_c(\mathbf{6}_F), 1, 1, \rho\rho]$ as an example. Firstly, when we take $\omega_c = 3.2 \text{ GeV}$, the Borel window can be evaluated to be $0.425 \text{ GeV} < T < 0.487 \text{ GeV}$, and the following numerical results can be obtained:

$$\begin{aligned} \bar{\Lambda}_{\Sigma_c, 1, 1, \rho\rho} &= 1.425 \text{ GeV}, \\ K_{\Sigma_c, 1, 1, \rho\rho} &= -1.372 \text{ GeV}^2, \\ \Sigma_{\Sigma_c, 1, 1, \rho\rho} &= 0.0091 \text{ GeV}^2. \end{aligned} \quad (13)$$

Their variations are shown in Fig. 2 with respect to the Borel mass T . Inserting them into Eqs. (2–3), we can further obtain

$$\begin{aligned} m_{\Sigma_c(1/2^+)} &= 2.96 \text{ GeV}, \\ m_{\Sigma_c(3/2^+)} &= 2.97 \text{ GeV}, \\ \Delta m_{[\Sigma_c, 1, 1, \rho\rho]} &= 11 \text{ MeV}, \end{aligned} \quad (14)$$

where $m_{\Sigma_c(1/2^+)}$ and $m_{\Sigma_c(3/2^+)}$ are the masses of the $\Sigma_c(1/2^+)$ and $\Sigma_c(3/2^+)$ belonging to the baryon multiplet $[\Sigma_c(\mathbf{6}_F), 1, 1, \rho\rho]$, and $\Delta m_{[\Sigma_c, 1, 1, \rho\rho]}$ is their mass splitting. We show the variation of $m_{\Sigma_c(1/2^+)}$ with respect to the Borel mass T in the right panel of Fig. 3.

Secondly, we change the threshold value ω_c and redo the above process. We show the variation of $m_{\Sigma_c(1/2^+)}$ with respect to the threshold value ω_c in the left panel of Fig. 3. There are non-vanishing Borel windows as long




FIG. 2: Variations of $\bar{\Lambda}_{\Sigma_c,1,1,\rho\rho}$ (left), $K_{\Sigma_c,1,1,\rho\rho}$ (middle), and $\Sigma_{\Sigma_c,1,1,\rho\rho}$ (right) with respect to the Borel mass T , calculated using the charmed baryon doublet $[\Sigma_c(\mathbf{6}_F), 1, 1, \rho\rho]$. The short-dashed, solid, and long-dashed curves are obtained by fixing $\omega_c = 3.0, 3.2$, and 3.4 GeV, respectively.




FIG. 3: Variations of $m_{\Sigma_c(1/2^+)}$ with respect to the threshold value ω_c (left) and the Borel mass T (right), calculated using the charmed baryon doublet $[\Sigma_c(\mathbf{6}_F), 1, 1, \rho\rho]$. The shady band in the left panel is obtained by changing T inside Borel windows. There exist non-vanishing working regions of T as long as $\omega_c \geq 3.0$ GeV, while the results for $\omega_c < 3.0$ GeV are also shown, for which cases we choose the Borel mass T when the PC defined in Eq. (12) is around 10%. In the right panel, the short-dashed, solid and long-dashed curves are obtained by setting $\omega_c = 3.0, 3.2$ and 3.4 GeV, respectively.

as $\omega_c \geq 3.0$ GeV, and the ω_c dependence is weak and acceptable in the region $3.0 \text{ GeV} < \omega_c < 3.4 \text{ GeV}$. The results for $\omega_c \leq 3.0$ GeV are also shown, for which cases we choose the Borel mass T when the PC defined in Eq. (12) is around 10%.

Finally, we choose our working regions to be $3.0 \text{ GeV} < \omega_c < 3.4 \text{ GeV}$ and $0.425 \text{ GeV} < T < 0.487 \text{ GeV}$, and obtain the following numerical results for the baryon multiplet $[\Sigma_c(\mathbf{6}_F), 1, 1, \rho\rho]$:

$$\begin{aligned} m_{\Sigma_c(1/2^+)} &= 2.96_{-0.12}^{+0.17} \text{ GeV}, \\ m_{\Sigma_c(3/2^+)} &= 2.97_{-0.13}^{+0.18} \text{ GeV}, \\ \Delta m_{[\Sigma_c,1,1,\rho\rho]} &= 11_{-9}^{+17} \text{ MeV}, \end{aligned} \quad (15)$$

where the central values correspond to $\omega_c = 3.2$ GeV and $T = 0.456$ GeV, and the uncertainties come from the threshold value ω_c , the Borel mass T , the charm quark mass m_c , and various quark and gluon condensates. We note that there are large theoretical uncertainties in our mass predictions, but the mass splitting within the same doublet is produced quite well with much less theoretical uncertainty, because it does not depend much on the charm quark mass [9, 10, 38, 40].

To study the charmed baryon multiplets $[\Xi'_c(\mathbf{6}_F), 1, 1, \rho\rho]$ and $[\Omega_c(\mathbf{6}_F), 1, 1, \rho\rho]$, we fine-tune the threshold value ω_c to be

$$\begin{aligned} \omega_c([\Xi'_c, 1, 1, \rho\rho]) &\approx 3.7 \text{ GeV}, \\ \omega_c([\Omega_c, 1, 1, \rho\rho]) &\approx 4.2 \text{ GeV}, \end{aligned} \quad (16)$$

so that

$$\begin{aligned} \omega_c([\Omega_c, 1, 1, \rho\rho]) - \omega_c([\Xi'_c, 1, 1, \rho\rho]) &\approx \\ \omega_c([\Xi'_c, 1, 1, \rho\rho]) - \omega_c([\Sigma_c, 1, 1, \rho\rho]) &\approx 0.5 \text{ GeV}, \end{aligned} \quad (17)$$

which value is the same as those used in our previous studies on excited heavy baryons [9, 10, 40].

After fixing the threshold value $\omega_c([\Xi'_c, 1, 1, \rho\rho])$ to be around 3.7 GeV, we can evaluate our working regions to be $3.5 \text{ GeV} < \omega_c < 3.9 \text{ GeV}$ and $0.395 \text{ GeV} < T < 0.574 \text{ GeV}$, and obtain the following numerical results for the charmed baryon multiplet $[\Xi'_c(\mathbf{6}_F), 1, 1, \rho\rho]$:

$$\begin{aligned} m_{\Xi'_c(1/2^+)} &= 3.02_{-0.21}^{+0.15} \text{ GeV}, \\ m_{\Xi'_c(3/2^+)} &= 3.03_{-0.21}^{+0.15} \text{ GeV}, \\ \Delta m_{[\Xi'_c,1,1,\rho\rho]} &= 7_{-6}^{+8} \text{ MeV}, \end{aligned} \quad (18)$$




FIG. 4: Variations of $m_{\Xi'_c(1/2^+)}$ (left) and $m_{\Omega_c(1/2^+)}$ (right) with respect to the Borel mass T , calculated using the charmed baryon multiplets $[\Xi'_c(\mathbf{6}_F), 1, 1, \rho\rho]$ and $[\Omega_c(\mathbf{6}_F), 1, 1, \rho\rho]$. In the left figure, the short-dashed, solid and long-dashed curves are obtained by setting $\omega_c = 3.5, 3.7$ and 3.9 GeV, respectively. In the right figure, the short-dashed, solid and long-dashed curves are obtained by setting $\omega_c = 4.0, 4.2$ and 4.4 GeV, respectively.

where the central values correspond to $\omega_c = 3.7$ GeV and $T = 0.485$ GeV. We show the variation of $m_{\Xi'_c(1/2^+)}$ with respect to the Borel mass T in the left panel of Fig. 4, where these curves are stable inside the Borel window $0.395 \text{ GeV} < T < 0.574 \text{ GeV}$. The masses of $\Xi'_c(1/2^+)$ and $\Xi'_c(3/2^+)$ are both consistent with the mass of the $\Xi_c(3123)$ observed by the BaBar Collaboration [19]:

$$m_{\Xi_c(3123)}^{\text{exp}} = 3122.9 \pm 1.3 \pm 0.3 \text{ MeV}, \quad (19)$$

which supports it to be a D -wave Ξ'_c state of $J^P = 1/2^+$ or $3/2^+$.

Similarly, after fixing $\omega_c([\Omega_c, 1, 1, \rho\rho])$ to be around 4.2 GeV, we can evaluate our working regions to be $4.0 \text{ GeV} < \omega_c < 4.4 \text{ GeV}$ and $0.421 \text{ GeV} < T < 0.655 \text{ GeV}$, and obtain the following numerical results for the charmed baryon multiplet $[\Omega_c(\mathbf{6}_F), 1, 1, \rho\rho]$:

$$\begin{aligned} m_{\Omega_c(1/2^+)} &= 3.29^{+0.17}_{-0.25} \text{ GeV}, \\ m_{\Omega_c(3/2^+)} &= 3.29^{+0.16}_{-0.25} \text{ GeV}, \\ \Delta m_{[\Omega_c, 1, 1, \rho\rho]} &= 5^{+5}_{-4} \text{ MeV}, \end{aligned} \quad (20)$$

where the central values correspond to $\omega_c = 4.2$ GeV and $T = 0.538$ GeV. We show the variation of $m_{\Omega_c(1/2^+)}$ with respect to the Borel mass T in the right panel of Fig. 4, where these curves are stable inside the Borel window $0.421 \text{ GeV} < T < 0.655 \text{ GeV}$.

We list the above results in Table I. Following the same procedures, we study the charmed baryon multiplets, $[\mathbf{6}_F, 3, 1, \lambda\lambda]$, $[\mathbf{6}_F, 2, 0, \rho\lambda]$, $[\mathbf{6}_F, 3, 1, \rho\rho]$, $[\mathbf{6}_F, 1, 1, \lambda\lambda]$, and obtain:

1. The baryon multiplet $[\mathbf{6}_F, 3, 1, \lambda\lambda]$ contains $\Sigma_c(5/2^+, 7/2^+)$, $\Xi'_c(5/2^+, 7/2^+)$, and $\Omega_c(5/2^+, 7/2^+)$. We use them to perform sum rule analyses, and obtain:

$$\begin{aligned} m_{[\Sigma_c, 3, 1, \lambda\lambda]} &\sim 3.32^{+0.64}_{-0.24} \text{ GeV}, \\ m_{[\Xi'_c, 3, 1, \lambda\lambda]} &\sim 3.39^{+0.40}_{-0.20} \text{ GeV}, \\ m_{[\Omega_c, 3, 1, \lambda\lambda]} &\sim 3.49^{+0.30}_{-0.19} \text{ GeV}. \end{aligned} \quad (21)$$

These values are also listed in Table I, and their variations are shown in Fig. 5 with respect to the threshold value ω_c . The masses of $\Xi'_c(5/2^+, 7/2^+)$ are not far from the mass of the $\Xi_c(3123)$ [19], suggesting that the $\Xi_c(3123)$ may also be interpreted as a D -wave Ξ'_c state of $J^P = 5/2^+$ or $7/2^+$.

We note that we can only evaluate their average masses $\frac{1}{14}(6m_{\Sigma_c(5/2^+)} + 8m_{\Sigma_c(7/2^+)})$, $\frac{1}{14}(6m_{\Xi'_c(5/2^+)} + 8m_{\Xi'_c(7/2^+)})$ and $\frac{1}{14}(6m_{\Omega_c(5/2^+)} + 8m_{\Omega_c(7/2^+)})$, because their mass splittings related to the chromomagnetic interaction ($\Sigma_{\mathbf{6}_F, 3, 1, \lambda\lambda}$) are difficult to be evaluated. More discussions can be found in Ref. [10].

2. The baryon multiplet $[\mathbf{6}_F, 2, 0, \rho\lambda]$ contains $\Sigma_c(3/2^+, 5/2^+)$, $\Xi'_c(3/2^+, 5/2^+)$, and $\Omega_c(3/2^+, 5/2^+)$. We use them to perform sum rule analyses, and obtain:

$$\begin{aligned} m_{\Sigma_c(3/2^+)} &= 3.02^{+0.26}_{-0.16} \text{ GeV}, \\ m_{\Sigma_c(5/2^+)} &= 3.04^{+0.28}_{-0.17} \text{ GeV}, \\ \Delta m_{[\Sigma_c, 2, 0, \rho\lambda]} &= 16^{+26}_{-14} \text{ MeV}, \\ m_{\Xi'_c(3/2^+)} &= 3.50^{+0.15}_{-0.13} \text{ GeV}, \\ m_{\Xi'_c(5/2^+)} &= 3.51^{+0.15}_{-0.13} \text{ GeV}, \\ \Delta m_{[\Xi'_c, 2, 0, \rho\lambda]} &= 11^{+12}_{-9} \text{ MeV}, \\ m_{\Omega_c(3/2^+)} &= 3.67^{+0.14}_{-0.14} \text{ GeV}, \\ m_{\Omega_c(5/2^+)} &= 3.68^{+0.15}_{-0.14} \text{ GeV}, \\ \Delta m_{[\Omega_c, 2, 0, \rho\lambda]} &= 9^{+9}_{-8} \text{ MeV}. \end{aligned} \quad (22)$$

These values are also listed in Table I, and variations of $m_{\Sigma_c(3/2^+)}$, $m_{\Xi'_c(3/2^+)}$ and $m_{\Omega_c(3/2^+)}$ are shown in Fig. 6 with respect to the threshold value ω_c . However, the mass differences among $\Sigma_c(3/2^+, 5/2^+)$, $\Xi'_c(3/2^+, 5/2^+)$ and

TABLE I: Masses of the D -wave charmed baryons of the $SU(3)$ flavor $\mathbf{6}_F$, obtained using the charmed baryon multiplets $[\mathbf{6}_F, 1, 1, \rho\rho]$, $[\mathbf{6}_F, 3, 1, \lambda\lambda]$ and $[\mathbf{6}_F, 2, 0, \rho\lambda]$. For the charmed baryon multiplet $[\mathbf{6}_F, 3, 1, \lambda\lambda]$ containing $\Sigma_c(5/2^+, 7/2^+)$, $\Xi'_c(5/2^+, 7/2^+)$, and $\Omega_c(5/2^+, 7/2^+)$, we can only evaluate their average masses, i.e., $\frac{1}{14}(6m_{\Sigma_c(5/2^+)} + 8m_{\Sigma_c(7/2^+)})$, $\frac{1}{14}(6m_{\Xi'_c(5/2^+)} + 8m_{\Xi'_c(7/2^+)})$ and $\frac{1}{14}(6m_{\Omega_c(5/2^+)} + 8m_{\Omega_c(7/2^+)})$, as discussed in Ref. [10]. For the baryon multiplet $[\mathbf{6}_F, 2, 0, \rho\lambda]$ containing $\Sigma_c(3/2^+, 5/2^+)$, $\Xi'_c(3/2^+, 5/2^+)$, and $\Omega_c(3/2^+, 5/2^+)$, the mass differences among Σ_c , Ξ'_c , and Ω_c seem a bit large, so we do not use them to draw conclusions.

Multiplets	B	ω_c (GeV)	Working region (GeV)	$\bar{\Lambda}$ (GeV)	f (GeV ⁵)	K (GeV ²)	Σ (GeV ²)	Baryons (j^P)	Mass (GeV)	Difference (MeV)
$[\mathbf{6}_F, 1, 1, \rho\rho]$	Σ_c	3.2	$0.425 < T < 0.487$	1.425	0.079	-1.372	0.0091	$\Sigma_c(1/2^+)$	$2.96^{+0.17}_{-0.12}$	11^{+17}_{-9}
								$\Sigma_c(3/2^+)$	$2.97^{+0.18}_{-0.13}$	
	Ξ'_c	3.7	$0.395 < T < 0.574$	1.561	0.146	-0.961	0.0057	$\Xi'_c(1/2^+)$	$3.02^{+0.15}_{-0.21}$	7^{+8}_{-6}
								$\Xi'_c(3/2^+)$	$3.03^{+0.15}_{-0.21}$	
	Ω_c	4.2	$0.421 < T < 0.655$	1.761	0.274	-1.302	0.0042	$\Omega_c(1/2^+)$	$3.29^{+0.17}_{-0.25}$	5^{+5}_{-4}
								$\Omega_c(3/2^+)$	$3.29^{+0.16}_{-0.25}$	
$[\mathbf{6}_F, 3, 1, \lambda\lambda]$	Σ_c	3.1	$0.445 < T < 0.459$	1.432	0.014	-3.139	-	$\Sigma_c(5/2^+)$	$3.32^{+0.64}_{-0.24}$	-
								$\Sigma_c(7/2^+)$		
	Ξ'_c	3.3	$0.458 < T < 0.485$	1.509	0.018	-3.064	-	$\Xi'_c(5/2^+)$	$3.39^{+0.40}_{-0.20}$	-
								$\Xi'_c(7/2^+)$		
	Ω_c	3.5	$0.490 < T < 0.509$	1.612	0.024	-3.085	-	$\Omega_c(5/2^+)$	$3.49^{+0.30}_{-0.19}$	-
								$\Omega_c(7/2^+)$		
$[\mathbf{6}_F, 2, 0, \rho\lambda]$	Σ_c	3.0	$0.398 < T < 0.457$	1.336	0.022	-2.156	0.0082	$\Sigma_c(3/2^+)$	$3.02^{+0.26}_{-0.16}$	16^{+26}_{-14}
								$\Sigma_c(5/2^+)$	$3.04^{+0.28}_{-0.17}$	
	Ξ'_c	3.7	$0.505 < T < 0.543$	1.793	0.076	-2.214	0.0055	$\Xi'_c(3/2^+)$	$3.50^{+0.15}_{-0.13}$	11^{+12}_{-9}
								$\Xi'_c(5/2^+)$	$3.51^{+0.15}_{-0.13}$	
	Ω_c	4.0	$0.561 < T < 0.575$	1.986	0.115	-2.121	0.0046	$\Omega_c(3/2^+)$	$3.67^{+0.14}_{-0.14}$	9^{+9}_{-8}
								$\Omega_c(5/2^+)$	$3.68^{+0.15}_{-0.14}$	

$\Omega_c(3/2^+, 5/2^+)$ seem a bit large, and we shall not use them to draw conclusions.

3. We use the baryon multiplet $[\mathbf{6}_F, 3, 1, \rho\rho]$ to perform sum rule analyses, but there do not exist Borel windows when $\omega_c \leq 4.0$ GeV. We also use the baryon multiplet $[\mathbf{6}_F, 1, 1, \lambda\lambda]$ to perform sum rule analyses, but the obtained results depend significantly on the threshold value ω_c . Hence, these results can not be well interpreted, and we shall not use them to draw conclusions.

IV. SUMMARY

In this paper we used the method of QCD sum rules within HQET to study the D -wave charmed baryons of the $SU(3)$ flavor $\mathbf{6}_F$, and calculated their masses up to the order $\mathcal{O}(1/m_Q)$. We investigated five charmed

baryon doublets, $[\mathbf{6}_F, 1, 1, \rho\rho]$, $[\mathbf{6}_F, 3, 1, \rho\rho]$, $[\mathbf{6}_F, 1, 1, \lambda\lambda]$, $[\mathbf{6}_F, 3, 1, \lambda\lambda]$ and $[\mathbf{6}_F, 2, 0, \rho\lambda]$ (we note that in Ref. [10] we failed to construct the currents belonging to the other two multiplets, $[\mathbf{6}_F, 2, 1, \rho\rho]$ and $[\mathbf{6}_F, 2, 1, \lambda\lambda]$). The results are summarized in Table I for the charmed baryon multiplets $[\mathbf{6}_F, 1, 1, \rho\rho]$, $[\mathbf{6}_F, 3, 1, \lambda\lambda]$, and $[\mathbf{6}_F, 2, 0, \rho\lambda]$, and those obtained using the former two multiplets ($[\mathbf{6}_F, 1, 1, \rho\rho]$ and $[\mathbf{6}_F, 3, 1, \lambda\lambda]$) are reasonable/better. We note that there are large theoretical uncertainties in our mass predictions, but the mass splittings within the same doublet are produced quite well with much less theoretical uncertainty.

Our results suggest that the $\Xi_c(3123)$ observed by the BaBar Collaboration [19] can be well interpreted as a D -wave Ξ'_c state, while its quantum numbers can not be determined. It may belong to either the charmed baryon doublet $[\Xi'_c(\mathbf{6}_F), 1, 1, \rho\rho]$ or $[\Xi'_c(\mathbf{6}_F), 3, 1, \lambda\lambda]$, but in both cases there would be a partner state close to it. Our results also suggest that there may exist as many as four




FIG. 5: Variations of $m_{[\Sigma_c(3,1,\lambda\lambda)]}$ (left), $m_{[\Xi'_c(3,1,\lambda\lambda)]}$ (middle) and $m_{[\Omega_c(3,1,\lambda\lambda)]}$ (right) with respect to the threshold value ω_c , calculated using the charmed baryon doublet $[6_F, 3, 1, \lambda\lambda]$. The shady bands in these figures are obtained by changing T inside Borel windows. There exist Borel windows as long as $\omega_c([\Sigma_c(3,1,\lambda\lambda)] \geq 3.1$ GeV (left), $\omega_c([\Xi'_c(3,1,\lambda\lambda)] \geq 3.2$ GeV (left) and $\omega_c([\Omega_c(3,1,\lambda\lambda)] \geq 3.5$ GeV (right). Accordingly, we choose ω_c to be around 3.1, 3.3, and 3.5 GeV in the left, middle and right panels, respectively.




FIG. 6: Variations of $m_{\Sigma_c(3/2^+)}$ (left), $m_{\Xi'_c(3/2^+)}$ (middle) and $m_{\Omega_c(5/2^+)}$ (right) with respect to the threshold value ω_c , calculated using the charmed baryon multiplet $[6_F, 2, 0, \rho\lambda]$. The shady bands in these figures are obtained by changing T inside Borel windows. There exist Borel windows as long as $\omega_c([\Sigma_c(2,0,\rho\lambda)] \geq 2.8$ GeV (left), $\omega_c([\Xi'_c(2,0,\rho\lambda)] \geq 3.6$ GeV (left) and $\omega_c([\Omega_c(2,0,\rho\lambda)] \geq 4.0$ GeV (right). Accordingly, we choose ω_c to be around 3.0, 3.7, and 4.0 GeV in the left, middle and right panels, respectively.

D -wave Ω_c states in the energy region $3.3 \sim 3.5$ GeV. They are the D -wave Ω_c states of $J^P = 1/2^+, 3/2^+, 5/2^+$, and $7/2^+$; the former two belong to the charmed baryon multiplet $[6_F, 1, 1, \rho\rho]$, and the latter two belong to $[6_F, 3, 1, \lambda\lambda]$. Recalling that the LHCb experiment observed as many as five excited Ω_c states [8] at the same time, we propose to search for these D -wave Ω_c states in the further LHCb and BelleII experiments in order to study the fine structure of the strong interaction. We note that the doubly charmed baryon Ξ_{cc}^{++} was recently discovered by the LHCb Collaboration [67], which can also be an idea platform to study the fine structure of the strong interaction [68].

Following the same procedures, we have investigated the D -wave bottom baryons of the $SU(3)$ flavor 6_F . The pole mass of the bottom quark $m_b = 4.78 \pm 0.06$ GeV [1] is used, and the obtained results are listed in Table II. We also suggest to search for them in further experiments. To end this paper, we note that not only masses but also decay and production properties are useful to understand the heavy baryons [66], and J-PARC is planning an experimental project of such studies [69]. In near future, the joint efforts from experimentalists and theorists will

be helpful to identify more and more charmed and bottom baryons.

ACKNOWLEDGMENTS

We thank Xian-Hui Zhong and Cheng-Ping Shen for useful discussions. This project is supported by the National Natural Science Foundation of China under Grants No. 11722540, No. 11475015, No. 11375024, No. 11222547, No. 11175073, No. 11575008, and No. 11621131001; the 973 program; the Ministry of Education of China (SRFDP under Grant No. 20120211110002 and the Fundamental Research Funds for the Central Universities); the Key Natural Science Research Program of Anhui Educational Committee (Grant No. KJ2016A774); the National Program for Support of Top-notch Youth Professionals. A.H. is supported by Grants-in-Aid for Scientific Research (Grants No. JP17K05441(C)).

TABLE II: Masses of the D -wave bottom baryons of the $SU(3)$ flavor $\mathbf{6}_F$, obtained using the bottom baryon multiplets $[\mathbf{6}_F, 1, 1, \rho\rho]$, $[\mathbf{6}_F, 3, 1, \lambda\lambda]$ and $[\mathbf{6}_F, 2, 0, \rho\lambda]$.

Multiplets	B	ω_c (GeV)	Working region (GeV)	$\bar{\Lambda}$ (GeV)	f (GeV ⁵)	K (GeV ²)	Σ (GeV ²)	Baryons (J^P)	Mass (GeV)	Difference (MeV)
$[\mathbf{6}_F, 1, 1, \rho\rho]$	Σ_b	3.2	$0.425 < T < 0.487$	1.425	0.079	-1.372	0.0091	$\Sigma_b(1/2^+)$	$6.28^{+0.18}_{-0.12}$	2^{+4}_{-2}
								$\Sigma_b(3/2^+)$	$6.28^{+0.18}_{-0.12}$	
	Ξ'_b	3.7	$0.395 < T < 0.574$	1.561	0.146	-0.961	0.0057	$\Xi'_b(1/2^+)$	$6.39^{+0.11}_{-0.14}$	1^{+2}_{-1}
								$\Xi'_b(3/2^+)$	$6.39^{+0.11}_{-0.14}$	
	Ω_b	4.2	$0.421 < T < 0.655$	1.761	0.274	-1.302	0.0042	$\Omega_b(1/2^+)$	$6.61^{+0.12}_{-0.16}$	1^{+1}_{-1}
								$\Omega_b(3/2^+)$	$6.61^{+0.12}_{-0.16}$	
$[\mathbf{6}_F, 3, 1, \lambda\lambda]$	Σ_b	3.1	$0.445 < T < 0.459$	1.432	0.014	-3.139	-	$\Sigma_b(5/2^+)$	$6.38^{+0.43}_{-0.17}$	-
								$\Sigma_b(7/2^+)$		
	Ξ'_b	3.3	$0.458 < T < 0.485$	1.509	0.018	-3.064	-	$\Xi'_b(5/2^+)$	$6.45^{+0.28}_{-0.15}$	-
								$\Xi'_b(7/2^+)$		
	Ω_b	3.5	$0.490 < T < 0.509$	1.612	0.024	-3.085	-	$\Omega_b(5/2^+)$	$6.55^{+0.22}_{-0.14}$	-
								$\Omega_b(7/2^+)$		
$[\mathbf{6}_F, 2, 0, \rho\lambda]$	Σ_b	3.0	$0.398 < T < 0.457$	1.336	0.022	-2.156	0.0082	$\Sigma_b(3/2^+)$	$6.23^{+0.19}_{-0.13}$	3^{+6}_{-3}
								$\Sigma_b(5/2^+)$	$6.23^{+0.20}_{-0.13}$	
	Ξ'_b	3.7	$0.505 < T < 0.543$	1.793	0.076	-2.214	0.0055	$\Xi'_b(3/2^+)$	$6.69^{+0.11}_{-0.09}$	2^{+3}_{-2}
								$\Xi'_b(5/2^+)$	$6.69^{+0.09}_{-0.09}$	
	Ω_b	4.0	$0.561 < T < 0.575$	1.986	0.115	-2.121	0.0046	$\Omega_b(3/2^+)$	$6.88^{+0.09}_{-0.08}$	2^{+2}_{-2}
								$\Omega_b(5/2^+)$	$6.88^{+0.09}_{-0.08}$	

Appendix A: Other Sum Rules

In this appendix we list the sum rules for other currents with different quark contents.

-
- [1] C. Patrignani *et al.* [Particle Data Group], Review of Particle Physics, Chin. Phys. C **40**, no. 10, 100001 (2016).
[2] H. X. Chen, W. Chen, X. Liu, Y. R. Liu and S. L. Zhu, A review of the open charm and open bottom systems, Rept. Prog. Phys. **80**, 076201 (2017).
[3] H. Y. Cheng, Charmed baryons circa 2015, Front. Phys. (Beijing) **10**, 101406 (2015).
[4] V. Crede and W. Roberts, Progress towards understanding baryon resonances, Rept. Prog. Phys. **76**, 076301 (2013).
[5] E. Klempt and J. M. Richard, Baryon spectroscopy, Rev. Mod. Phys. **82**, 1095 (2010).
[6] S. Bianco, F. L. Fabbri, D. Benson and I. Bigi, A Cicerone for the physics of charm, Riv. Nuovo Cim. **26N7**, 1 (2003).
[7] J. G. Körner, M. Kramer and D. Pirjol, Heavy baryons, Prog. Part. Nucl. Phys. **33**, 787 (1994).
[8] R. Aaij *et al.* [LHCb Collaboration], Observation of five new narrow Ω_c^0 states decaying to $\Xi_c^+ K^-$, Phys. Rev. Lett. **118**, 182001 (2017).
[9] H. X. Chen, W. Chen, Q. Mao, A. Hosaka, X. Liu and S. L. Zhu, P-wave charmed baryons from QCD sum rules, Phys. Rev. D **91**, 054034 (2015).
[10] H. X. Chen, Q. Mao, A. Hosaka, X. Liu and S. L. Zhu, D-wave charmed and bottomed baryons from QCD sum rules, Phys. Rev. D **94**, 114016 (2016).
[11] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, QCD and Resonance Physics: Theoretical Foundations, Nucl. Phys. B **147**, 385 (1979).
[12] L. J. Reinders, H. Rubinstein and S. Yazaki, Hadron Properties From QCD Sum Rules, Phys. Rept. **127**, 1 (1985).

$$\Pi_{\Xi_c', 1, 1, \rho\rho} = f_{\Xi_c', 1, 1, \rho\rho}^2 e^{-2\bar{\Lambda}_{\Xi_c', 1, 1, \rho\rho}/T} \quad (A1)$$

$$= \int_{2m_s}^{\omega_c} \left[\frac{11}{80640\pi^4} \omega^9 - \frac{51m_s^2}{8960\pi^4} \omega^7 - \frac{m_s \langle \bar{q}q \rangle}{16\pi^2} \omega^5 + \frac{3m_s \langle \bar{s}s \rangle}{40\pi^2} \omega^5 \right. \\ \left. - \frac{19 \langle g_s^2 GG \rangle}{3072\pi^4} \omega^5 + \frac{33m_s^2 \langle g_s^2 GG \rangle}{512\pi^4} \omega^3 - \frac{m_s \langle g_s^2 GG \rangle \langle \bar{q}q \rangle}{16\pi^2} \omega - \frac{31m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{128\pi^2} \omega \right] e^{-\omega/T} d\omega, \quad (A2)$$

$$f_{\Xi_c', 1, 1, \rho\rho}^2 K_{\Xi_c', 1, 1, \rho\rho} e^{-2\bar{\Lambda}_{\Xi_c', 1, 1, \rho\rho}/T} \\ = \int_{2m_s}^{\omega_c} \left[-\frac{59}{2217600\pi^4} \omega^{11} + \frac{29m_s^2}{17920\pi^4} \omega^9 + \frac{m_s \langle \bar{q}q \rangle}{32\pi^2} \omega^7 - \frac{11m_s \langle \bar{s}s \rangle}{224\pi^2} \omega^7 + \frac{m_s \langle g_s \bar{q} \sigma G q \rangle}{4\pi^2} \omega^5 \right. \\ \left. + \frac{69 \langle g_s^2 GG \rangle}{35840\pi^4} \omega^7 - \frac{2527m_s^2 \langle g_s^2 GG \rangle}{61440\pi^4} \omega^5 - \frac{35m_s \langle g_s^2 GG \rangle \langle \bar{q}q \rangle}{192\pi^2} \omega^3 + \frac{839m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{2304\pi^2} \omega^3 \right. \\ \left. - \frac{19m_s \langle g_s^2 GG \rangle \langle g_s \bar{q} \sigma G q \rangle}{96\pi^2} \omega \right] e^{-\omega/T} d\omega, \quad (A3)$$

$$f_{\Xi_c', 1, 1, \rho\rho}^2 \Sigma_{\Xi_c', 1, 1, \rho\rho} e^{-2\bar{\Lambda}_{\Xi_c', 1, 1, \rho\rho}/T} \\ = \int_{2m_s}^{\omega_c} \left[\frac{37 \langle g_s^2 GG \rangle}{322560\pi^4} \omega^7 - \frac{7m_s^2 \langle g_s^2 GG \rangle}{3840\pi^4} \omega^5 + \frac{19m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{1152\pi^2} \omega^3 \right] e^{-\omega/T} d\omega. \quad (A4)$$

$$\Pi_{\Omega_c, 1, 1, \rho\rho} = f_{\Omega_c, 1, 1, \rho\rho}^2 e^{-2\bar{\Lambda}_{\Omega_c, 1, 1, \rho\rho}/T} \quad (A5)$$

$$= \int_{4m_s}^{\omega_c} \left[\frac{11}{80640\pi^4} \omega^9 - \frac{9m_s^2}{1120\pi^4} \omega^7 + \frac{9m_s^4}{80\pi^4} \omega^5 + \frac{m_s \langle \bar{s}s \rangle}{40\pi^2} \omega^5 - \frac{7m_s^3 \langle \bar{s}s \rangle}{8\pi^2} \omega^3 \right. \\ \left. - \frac{19 \langle g_s^2 GG \rangle}{3072\pi^4} \omega^5 + \frac{37m_s^2 \langle g_s^2 GG \rangle}{256\pi^4} \omega^3 - \frac{3m_s^4 \langle g_s^2 GG \rangle}{64\pi^4} \omega - \frac{39m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{64\pi^2} \omega \right] e^{-\omega/T} d\omega,$$

$$f_{\Omega_c, 1, 1, \rho\rho}^2 K_{\Omega_c, 1, 1, \rho\rho} e^{-2\bar{\Lambda}_{\Omega_c, 1, 1, \rho\rho}/T} \\ = \int_{4m_s}^{\omega_c} \left[-\frac{59}{2217600\pi^4} \omega^{11} + \frac{7m_s^2}{2880\pi^4} \omega^9 - \frac{7m_s^4}{160\pi^4} \omega^7 - \frac{m_s \langle \bar{s}s \rangle}{28\pi^2} \omega^7 + \frac{15m_s^3 \langle \bar{s}s \rangle}{16\pi^2} \omega^5 + \frac{m_s \langle g_s \bar{s} \sigma G s \rangle}{2\pi^2} \omega^5 \right. \\ \left. + \frac{23 \langle g_s^2 GG \rangle}{11520\pi^4} \omega^7 - \frac{2153m_s^2 \langle g_s^2 GG \rangle}{30720\pi^4} \omega^5 + \frac{23m_s^4 \langle g_s^2 GG \rangle}{64\pi^4} \omega^3 \right. \\ \left. + \frac{149m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{384\pi^2} \omega^3 - \frac{239m_s^3 \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{192\pi^2} \omega + \frac{17m_s \langle g_s^2 GG \rangle \langle g_s \bar{s} \sigma G s \rangle}{48\pi^2} \omega \right] e^{-\omega/T} d\omega, \quad (A6)$$

$$f_{\Omega_c, 1, 1, \rho\rho}^2 \Sigma_{\Omega_c, 1, 1, \rho\rho} e^{-2\bar{\Lambda}_{\Omega_c, 1, 1, \rho\rho}/T} \\ = \int_{4m_s}^{\omega_c} \left[\frac{37 \langle g_s^2 GG \rangle}{322560\pi^4} \omega^7 - \frac{7m_s^2 \langle g_s^2 GG \rangle}{1920\pi^4} \omega^5 + \frac{19m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{576\pi^2} \omega^3 \right] e^{-\omega/T} d\omega.$$

-
- [13] B. Grinstein, The Static Quark Effective Theory, Nucl. Phys. B **339**, 253 (1990).
- [14] E. Eichten and B. R. Hill, An Effective Field Theory for the Calculation of Matrix Elements Involving Heavy Quarks, Phys. Lett. B **234**, 511 (1990).
- [15] A. F. Falk, H. Georgi, B. Grinstein and M. B. Wise, Heavy Meson Form-factors From QCD, Nucl. Phys. B **343**, 1 (1990).
- [16] J. Yelton *et al.* [Belle Collaboration], Study of Excited Ξ_c States Decaying into Ξ_c^0 and Ξ_c^+ Baryons, Phys. Rev. D **94**, 052011 (2016).
- [17] Y. Kato *et al.* [Belle Collaboration], Studies of charmed strange baryons in the ΛD final state at Belle, Phys. Rev. D **94**, 032002 (2016).
- [18] R. Aaij *et al.* [LHCb Collaboration], Study of the $D^0 p$ amplitude in $\Lambda_b^0 \rightarrow D^0 p \pi^-$ decays, JHEP **1705**, 030 (2017).
- [19] B. Aubert *et al.* [BaBar Collaboration], A Study of Excited Charm-Strange Baryons with Evidence for new Baryons $\Xi_c(3055)^+$ and $\Xi_c(3123)^+$, Phys. Rev. D **77**, 012002 (2008).
- [20] Y. Kato *et al.* [Belle Collaboration], Search for doubly charmed baryons and study of charmed strange baryons at Belle, Phys. Rev. D **89**, 052003 (2014).
- [21] D. Ebert, R. N. Faustov and V. O. Galkin, Masses of excited heavy baryons in the relativistic quark model, Phys. Lett. B **659**, 612 (2008).
- [22] X. H. Zhong and Q. Zhao, Charmed baryon strong decays in a chiral quark model, Phys. Rev. D **77**, 074008 (2008).
- [23] D. Ebert, R. N. Faustov and V. O. Galkin, Spectroscopy and Regge trajectories of heavy baryons in the relativistic quark-diquark picture, Phys. Rev. D **84**, 014025 (2011).

$$\Pi_{\Sigma_c,3,1,\lambda\lambda} = f_{\Sigma_c,3,1,\lambda\lambda}^2 e^{-2\bar{\Lambda}_{\Sigma_c,3,1,\lambda\lambda}/T} = \int_0^{\omega_c} \left[\frac{1}{193536\pi^4} \omega^9 - \frac{13\langle g_s^2 GG \rangle}{46080\pi^4} \omega^5 \right] e^{-\omega/T} d\omega, \quad (A7)$$

$$f_{\Sigma_c,3,1,\lambda\lambda}^2 K_{\Sigma_c,3,1,\lambda\lambda} e^{-2\bar{\Lambda}_{\Sigma_c,3,1,\lambda\lambda}/T} = \int_0^{\omega_c} \left[-\frac{41}{21288960\pi^4} \omega^{11} + \frac{353\langle g_s^2 GG \rangle}{3870720\pi^4} \omega^7 \right] e^{-\omega/T} d\omega. \quad (A8)$$

$$\Pi_{\Xi'_c,3,1,\lambda\lambda} = f_{\Xi'_c,3,1,\lambda\lambda}^2 e^{-2\bar{\Lambda}_{\Xi'_c,3,1,\lambda\lambda}/T} \quad (A9)$$

$$= \int_{2m_s}^{\omega_c} \left[\frac{1}{193536\pi^4} \omega^9 - \frac{m_s^2}{4480\pi^4} \omega^7 - \frac{m_s \langle \bar{q}q \rangle}{480\pi^2} \omega^5 + \frac{m_s \langle \bar{s}s \rangle}{320\pi^2} \omega^5 \right. \\ \left. - \frac{13\langle g_s^2 GG \rangle}{46080\pi^4} \omega^5 + \frac{5m_s^2 \langle g_s^2 GG \rangle}{3072\pi^4} \omega^3 - \frac{m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{256\pi^2} \omega \right] e^{-\omega/T} d\omega, \\ f_{\Xi'_c,3,1,\lambda\lambda}^2 K_{\Xi'_c,3,1,\lambda\lambda} e^{-2\bar{\Lambda}_{\Xi'_c,3,1,\lambda\lambda}/T} \quad (A10)$$

$$= \int_{2m_s}^{\omega_c} \left[-\frac{41}{21288960\pi^4} \omega^{11} + \frac{17m_s^2}{161280\pi^4} \omega^9 + \frac{m_s \langle \bar{q}q \rangle}{960\pi^2} \omega^7 - \frac{9m_s \langle \bar{s}s \rangle}{4480\pi^2} \omega^7 \right. \\ \left. + \frac{353\langle g_s^2 GG \rangle}{3870720\pi^4} \omega^7 - \frac{151m_s^2 \langle g_s^2 GG \rangle}{122880\pi^4} \omega^5 - \frac{5m_s \langle g_s^2 GG \rangle \langle \bar{q}q \rangle}{1728\pi^2} \omega^3 + \frac{31m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{4608\pi^2} \omega^3 \right] e^{-\omega/T} d\omega.$$

$$\Pi_{\Omega_c,3,1,\lambda\lambda} = f_{\Omega_c,3,1,\lambda\lambda}^2 e^{-2\bar{\Lambda}_{\Omega_c,3,1,\lambda\lambda}/T} \quad (A11)$$

$$= \int_{4m_s}^{\omega_c} \left[\frac{1}{193536\pi^4} \omega^9 - \frac{3m_s^2}{8960\pi^4} \omega^7 + \frac{3m_s^4}{640\pi^4} \omega^5 + \frac{m_s \langle \bar{s}s \rangle}{480\pi^2} \omega^5 - \frac{m_s^3 \langle \bar{s}s \rangle}{24\pi^2} \omega^3 \right. \\ \left. - \frac{13\langle g_s^2 GG \rangle}{46080\pi^4} \omega^5 + \frac{5m_s^2 \langle g_s^2 GG \rangle}{1536\pi^4} \omega^3 - \frac{m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{128\pi^2} \omega \right] e^{-\omega/T} d\omega,$$

$$f_{\Omega_c,3,1,\lambda\lambda}^2 K_{\Omega_c,3,1,\lambda\lambda} e^{-2\bar{\Lambda}_{\Omega_c,3,1,\lambda\lambda}/T} \quad (A12)$$

$$= \int_{4m_s}^{\omega_c} \left[-\frac{41}{21288960\pi^4} \omega^{11} + \frac{3m_s^2}{17920\pi^4} \omega^9 - \frac{27m_s^4}{8960\pi^4} \omega^7 - \frac{13m_s \langle \bar{s}s \rangle}{6720\pi^2} \omega^7 + \frac{m_s^3 \langle \bar{s}s \rangle}{24\pi^2} \omega^5 \right. \\ \left. + \frac{353\langle g_s^2 GG \rangle}{3870720\pi^4} \omega^7 - \frac{131m_s^2 \langle g_s^2 GG \rangle}{61440\pi^4} \omega^5 + \frac{7m_s^4 \langle g_s^2 GG \rangle}{1152\pi^4} \omega^3 \right. \\ \left. + \frac{53m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{6912\pi^2} \omega^3 - \frac{m_s^3 \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{64\pi^2} \omega \right] e^{-\omega/T} d\omega.$$

-
- [24] L. H. Liu, L. Y. Xiao and X. H. Zhong, Charm-strange baryon strong decays in a chiral quark model, Phys. Rev. D **86**, 034024 (2012).
- [25] Z. Shah, K. Thakkar, A. Kumar Rai and P. C. Vinodkumar, Excited State Mass spectra of Singly Charmed Baryons, Eur. Phys. J. A **52**, 313 (2016).
- [26] Z. Zhao, D. D. Ye and A. Zhang, Nature of charmed strange baryons $\Xi_c(3055)$ and $\Xi_c(3080)$, Phys. Rev. D **94**, 114020 (2016).
- [27] A. Valcarce, H. Garcilazo and J. Vijande, Towards an understanding of heavy baryon spectroscopy, Eur. Phys. J. A **37**, 217 (2008).
- [28] X. H. Guo, K. W. Wei and X. H. Wu, Some mass relations for mesons and baryons in Regge phenomenology, Phys. Rev. D **78**, 056005 (2008).
- [29] B. Chen, K. W. Wei and A. Zhang, Assignments of Λ_Q and Ξ_Q baryons in the heavy quark-light diquark picture, Eur. Phys. J. A **51**, 82 (2015).
- [30] H. Y. Cheng and C. K. Chua, Strong Decays of Charmed Baryons in Heavy Hadron Chiral Perturbation Theory: An Update, Phys. Rev. D **92**, 074014 (2015).
- [31] J. R. Zhang and M. Q. Huang, Heavy baryon spectroscopy in QCD, Phys. Rev. D **78**, 094015 (2008).
- [32] Z. G. Wang, Analysis of the $\Lambda_c(2860)$, $\Lambda_c(2880)$, $\Xi_c(3055)$ and $\Xi_c(3080)$ as D-wave baryon states with QCD sum rules, arXiv:1705.07745 [hep-ph].
- [33] H. Y. Cheng and C. K. Chua, Strong Decays of Charmed Baryons in Heavy Hadron Chiral Perturbation Theory, Phys. Rev. D **75**, 014006 (2007).
- [34] A. Selem and F. Wilczek, Hadron systematics and emergent diquarks, hep-ph/0602128.
- [35] H. Garcilazo, J. Vijande and A. Valcarce, Faddeev study of heavy baryon spectroscopy, J. Phys. G **34**, 961 (2007).
- [36] S. M. Gerasyuta and E. E. Matskevich, Charmed (70, 1⁻) baryon multiplet, Int. J. Mod. Phys. E **17**, 585 (2008).
- [37] Z. H. Guo and J. A. Oller, Probabilistic interpretation of compositeness relation for resonances, Phys. Rev. D **93**, 096001 (2016).
- [38] X. Liu, H. X. Chen, Y. R. Liu, A. Hosaka and S. L. Zhu, Bottom baryons, Phys. Rev. D **77**, 014031 (2008).

$$\Pi_{\Sigma_c, 2, 0, \rho\lambda} = f_{\Sigma_c, 2, 0, \rho\lambda}^2 e^{-2\bar{\Lambda}_{\Sigma_c, 2, 0, \rho\lambda}/T} = \int_0^{\omega_c} \left[\frac{1}{48384\pi^4} \omega^9 - \frac{5\langle g_s^2 GG \rangle}{6912\pi^4} \omega^5 \right] e^{-\omega/T} d\omega, \quad (\text{A13})$$

$$f_{\Sigma_c, 2, 0, \rho\lambda}^2 K_{\Sigma_c, 2, 0, \rho\lambda} e^{-2\bar{\Lambda}_{\Sigma_c, 2, 0, \rho\lambda}/T} = \int_0^{\omega_c} \left[-\frac{1}{168960\pi^4} \omega^{11} + \frac{227\langle g_s^2 GG \rangle}{1451520\pi^4} \omega^7 \right] e^{-\omega/T} d\omega, \quad (\text{A14})$$

$$f_{\Sigma_c, 2, 0, \rho\lambda}^2 \Sigma_{\Sigma_c, 2, 0, \rho\lambda} e^{-2\bar{\Lambda}_{\Sigma_c, 2, 0, \rho\lambda}/T} = \int_0^{\omega_c} \left[\frac{\langle g_s^2 GG \rangle}{48384\pi^4} \omega^7 \right] e^{-\omega/T} d\omega. \quad (\text{A15})$$

$$\Pi_{\Xi'_c, 2, 0, \rho\lambda} = f_{\Xi'_c, 2, 0, \rho\lambda}^2 e^{-2\bar{\Lambda}_{\Xi'_c, 2, 0, \rho\lambda}/T} \quad (\text{A16})$$

$$= \int_{2m_s}^{\omega_c} \left[\frac{1}{48384\pi^4} \omega^9 - \frac{m_s^2}{1008\pi^4} \omega^7 - \frac{m_s \langle \bar{q}q \rangle}{72\pi^2} \omega^5 + \frac{m_s \langle \bar{s}s \rangle}{48\pi^2} \omega^5 - \frac{5m_s \langle g_s \bar{q}\sigma Gq \rangle}{72\pi^2} \omega^3 \right. \\ \left. - \frac{5\langle g_s^2 GG \rangle}{6912\pi^4} \omega^5 + \frac{5m_s^2 \langle g_s^2 GG \rangle}{432\pi^4} \omega^3 - \frac{5m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{108\pi^2} \omega \right] e^{-\omega/T} d\omega, \quad (\text{A17})$$

$$f_{\Xi'_c, 2, 0, \rho\lambda}^2 K_{\Xi'_c, 2, 0, \rho\lambda} e^{-2\bar{\Lambda}_{\Xi'_c, 2, 0, \rho\lambda}/T} \\ = \int_{2m_s}^{\omega_c} \left[-\frac{1}{168960\pi^4} \omega^{11} + \frac{137m_s^2}{362880\pi^4} \omega^9 + \frac{37m_s \langle \bar{q}q \rangle}{5040\pi^2} \omega^7 - \frac{17m_s \langle \bar{s}s \rangle}{1344\pi^2} \omega^7 + \frac{m_s \langle g_s \bar{q}\sigma Gq \rangle}{15\pi^2} \omega^5 \right. \\ \left. + \frac{227\langle g_s^2 GG \rangle}{1451520\pi^4} \omega^7 - \frac{3419m_s^2 \langle g_s^2 GG \rangle}{552960\pi^4} \omega^5 - \frac{13m_s \langle g_s^2 GG \rangle \langle \bar{q}q \rangle}{648\pi^2} \omega^3 + \frac{461m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{10368\pi^2} \omega^3 \right. \\ \left. - \frac{11m_s \langle g_s^2 GG \rangle \langle g_s \bar{q}\sigma Gq \rangle}{1728\pi^2} \omega \right] e^{-\omega/T} d\omega, \quad (\text{A18})$$

$$f_{\Xi'_c, 2, 0, \rho\lambda}^2 \Sigma_{\Xi'_c, 2, 0, \rho\lambda} e^{-2\bar{\Lambda}_{\Xi'_c, 2, 0, \rho\lambda}/T} \\ = \int_{2m_s}^{\omega_c} \left[\frac{\langle g_s^2 GG \rangle}{48384\pi^4} \omega^7 - \frac{m_s^2 \langle g_s^2 GG \rangle}{2304\pi^4} \omega^5 + \frac{5m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{864\pi^2} \omega^3 \right] e^{-\omega/T} d\omega. \quad (\text{A19})$$

$$\Pi_{\Omega_c, 2, 0, \rho\lambda} = f_{\Omega_c, 2, 0, \rho\lambda}^2 e^{-2\bar{\Lambda}_{\Omega_c, 2, 0, \rho\lambda}/T} \quad (\text{A19})$$

$$= \int_{4m_s}^{\omega_c} \left[\frac{1}{48384\pi^4} \omega^9 - \frac{m_s^2}{672\pi^4} \omega^7 + \frac{m_s^4}{48\pi^4} \omega^5 + \frac{m_s \langle \bar{s}s \rangle}{72\pi^2} \omega^5 - \frac{5m_s^3 \langle \bar{s}s \rangle}{18\pi^2} \omega^3 - \frac{5m_s \langle g_s \bar{s}\sigma Gs \rangle}{36\pi^2} \omega^3 \right. \\ \left. - \frac{5\langle g_s^2 GG \rangle}{6912\pi^4} \omega^5 + \frac{5m_s^2 \langle g_s^2 GG \rangle}{216\pi^4} \omega^3 - \frac{5m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{54\pi^2} \omega \right] e^{-\omega/T} d\omega, \quad (\text{A20})$$

$$f_{\Omega_c, 2, 0, \rho\lambda}^2 K_{\Omega_c, 2, 0, \rho\lambda} e^{-2\bar{\Lambda}_{\Omega_c, 2, 0, \rho\lambda}/T} \\ = \int_{4m_s}^{\omega_c} \left[-\frac{1}{168960\pi^4} \omega^{11} + \frac{211m_s^2}{362880\pi^4} \omega^9 - \frac{211m_s^4}{20160\pi^4} \omega^7 - \frac{107m_s \langle \bar{s}s \rangle}{10080\pi^2} \omega^7 + \frac{181m_s^3 \langle \bar{s}s \rangle}{720\pi^2} \omega^5 + \frac{2m_s \langle g_s \bar{s}\sigma Gs \rangle}{15\pi^2} \omega^5 \right. \\ \left. + \frac{227\langle g_s^2 GG \rangle}{1451520\pi^4} \omega^7 - \frac{2839m_s^2 \langle g_s^2 GG \rangle}{276480\pi^4} \omega^5 + \frac{427m_s^4 \langle g_s^2 GG \rangle}{13824\pi^4} \omega^3 \right. \\ \left. + \frac{253m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{5184\pi^2} \omega^3 - \frac{263m_s^3 \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{3456\pi^2} \omega - \frac{11m_s \langle g_s^2 GG \rangle \langle g_s \bar{s}\sigma Gs \rangle}{864\pi^2} \omega \right] e^{-\omega/T} d\omega, \quad (\text{A21})$$

$$f_{\Omega_c, 2, 0, \rho\lambda}^2 \Sigma_{\Omega_c, 2, 0, \rho\lambda} e^{-2\bar{\Lambda}_{\Omega_c, 2, 0, \rho\lambda}/T} \\ = \int_{4m_s}^{\omega_c} \left[\frac{\langle g_s^2 GG \rangle}{48384\pi^4} \omega^7 - \frac{m_s^2 \langle g_s^2 GG \rangle}{1152\pi^4} \omega^5 + \frac{5m_s \langle g_s^2 GG \rangle \langle \bar{s}s \rangle}{432\pi^2} \omega^3 \right] e^{-\omega/T} d\omega. \quad (\text{A21})$$

- [39] H. X. Chen, V. Dmitrasinovic, A. Hosaka, K. Nagata and S. L. Zhu, Chiral Properties of Baryon Fields with Flavor SU(3) Symmetry, Phys. Rev. D **78**, 054021 (2008).
 [40] Q. Mao, H. X. Chen, W. Chen, A. Hosaka, X. Liu and S. L. Zhu, QCD sum rule calculation for P-wave bottom baryons, Phys. Rev. D **92**, 114007 (2015).

- [41] H. X. Chen, W. Chen, X. Liu, T. G. Steele and S. L. Zhu, Towards exotic hidden-charm pentaquarks in QCD, Phys. Rev. Lett. **115**, 172001 (2015).
 [42] H. X. Chen, E. L. Cui, W. Chen, X. Liu, T. G. Steele and S. L. Zhu, QCD sum rule study of hidden-charm pentaquarks, Eur. Phys. J. C **76**, 572 (2016).

- [43] E. Bagan, P. Ball, V. M. Braun and H. G. Dosch, QCD sum rules in the effective heavy quark theory, Phys. Lett. B **278**, 457 (1992).
- [44] M. Neubert, Heavy meson form-factors from QCD sum rules, Phys. Rev. D **45**, 2451 (1992).
- [45] D. J. Broadhurst and A. G. Grozin, Operator product expansion in static quark effective field theory: Large perturbative correction, Phys. Lett. B **274**, 421 (1992).
- [46] P. Ball and V. M. Braun, Next-to-leading order corrections to meson masses in the heavy quark effective theory, Phys. Rev. D **49**, 2472 (1994).
- [47] T. Huang and C. W. Luo, Light quark dependence of the Isgur-Wise function from QCD sum rules, Phys. Rev. D **50**, 5775 (1994).
- [48] Y. B. Dai, C. S. Huang, M. Q. Huang and C. Liu, QCD sum rules for masses of excited heavy mesons, Phys. Lett. B **390**, 350 (1997).
- [49] P. Colangelo, F. De Fazio and N. Paver, Universal $\tau_{1/2}(y)$ Isgur-Wise function at the next-to-leading order in QCD sum rules, Phys. Rev. D **58**, 116005 (1998).
- [50] S. Groote, J. G. Körner and O. I. Yakovlev, QCD sum rules for heavy baryons at next-to-leading order in α_s , Phys. Rev. D **55**, 3016 (1997).
- [51] S. L. Zhu, Strong and electromagnetic decays of p wave heavy baryons Λ_{c1} , Λ_{c1}^* , Phys. Rev. D **61**, 114019 (2000).
- [52] J. P. Lee, C. Liu and H. S. Song, QCD sum rule analysis of excited Λ_c mass parameter, Phys. Lett. B **476**, 303 (2000).
- [53] C. S. Huang, A. I. Zhang and S. L. Zhu, Excited heavy baryon masses in HQET QCD sum rules, Phys. Lett. B **492**, 288 (2000).
- [54] D. W. Wang and M. Q. Huang, Excited heavy baryon masses to order Λ_{QCD}/m_Q from QCD sum rules, Phys. Rev. D **68**, 034019 (2003).
- [55] F. O. Duraes and M. Nielsen, QCD sum rules study of Ξ_c and Ξ_b baryons, Phys. Lett. B **658**, 40 (2007).
- [56] D. Zhou, E. L. Cui, H. X. Chen, L. S. Geng, X. Liu and S. L. Zhu, The D-wave heavy-light mesons from QCD sum rules, Phys. Rev. D **90**, 114035 (2014).
- [57] D. Zhou, H. X. Chen, L. S. Geng, X. Liu and S. L. Zhu, F-wave heavy-light meson spectroscopy in QCD sum rules and heavy quark effective theory, Phys. Rev. D **92**, 114015 (2015).
- [58] K. C. Yang, W. Y. P. Hwang, E. M. Henley and L. S. Kisslinger, QCD sum rules and neutron proton mass difference, Phys. Rev. D **47**, 3001 (1993).
- [59] W. Y. P. Hwang and K. C. Yang, Phys. Rev. D **49**, 460 (1994).
- [60] S. Narison, QCD as a theory of hadrons (from partons to confinement), Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **17**, 1 (2002).
- [61] V. Gimenez, V. Lubicz, F. Mescia, V. Porretti and J. Reyes, Operator product expansion and quark condensate from lattice QCD in coordinate space, Eur. Phys. J. C **41**, 535 (2005).
- [62] M. Jamin, Flavour-symmetry breaking of the quark condensate and chiral corrections to the Gell-Mann-Oakes-Renner relation, Phys. Lett. B **538**, 71 (2002).
- [63] B. L. Ioffe and K. N. Zyblyuk, Gluon condensate in charmonium sum rules with 3-loop corrections, Eur. Phys. J. C **27**, 229 (2003).
- [64] A. A. Ovchinnikov and A. A. Pivovarov, QCD Sum Rule Calculation Of The Quark Gluon Condensate, Sov. J. Nucl. Phys. **48**, 721 (1988) [Yad. Fiz. **48**, 1135 (1988)].
- [65] P. Colangelo and A. Khodjamirian, “*At the Frontier of Particle Physics/Handbook of QCD*” (World Scientific, Singapore, 2001), Volume 3, 1495.
- [66] H. X. Chen, Q. Mao, W. Chen, A. Hosaka, X. Liu and S. L. Zhu, Decay properties of P -wave charmed baryons from light-cone QCD sum rules, Phys. Rev. D **95**, 094008 (2017).
- [67] R. Aaij *et al.* [LHCb Collaboration], Observation of the doubly charmed baryon Ξ_{cc}^{++} , Phys. Rev. Lett. **119**, 112001 (2017).
- [68] H. X. Chen, Q. Mao, W. Chen, X. Liu and S. L. Zhu, Establishing low-lying doubly charmed baryons, Phys. Rev. D **96**, 031501 (2017).
- [69] Charmed Baryon Spectroscopy via the (π^-, D^{*-}) reaction (2012). (Available at: http://www.j-parc.jp/researcher/Hadron/en/Proposal_e.html#1 J-PARC P50 proposal.