

Supergravity Model of Inflation and Explaining IceCube HESE Data via PeV Dark Matter Decay

Girish Kumar Chakravarty,^{1,*} Najimuddin Khan,^{2,†} and Subhendra Mohanty^{3,‡}

¹*Centre for Theoretical Physics, Jamia Millia Islamia, New Delhi, India*

²*School of Physical Sciences, Indian Association for the Cultivation of Science, Kolkata, India*

³*Theoretical Physics Division, Physical Research Laboratory, Ahmedabad, India*

Abstract

We construct a unified model of inflation and PeV dark matter with an appropriate choice of no-scale Kähler potential, superpotential and gauge kinetic function in terms of MSSM fields and hidden sector Polonyi field. The model is consistent with the CMB observations and can explain the PeV neutrino flux observed at IceCube HESE. A Starobinsky like Higgs-sneutrino plateau inflation is obtained from the D -term SUGRA potential while F -term being subdominant during inflation. To get PeV dark matter, SUSY breaking at PeV scale is achieved through Polonyi field. This sets the scale for soft SUSY breaking parameters $m_0, m_{\frac{1}{2}}, A_0$ at the GUT scale in terms of the parameters of the model. The low energy particles spectrum is obtained by running the RGEs. We show that the ~ 125 GeV higgs and the gauge coupling unification can be obtained in this model. The 6 PeV bino-type dark matter is a subdominant fraction ($\sim 11\%$) of the relic density and its decay gives the PeV scale neutrino flux observed at IceCube by appropriately choosing the couplings of the R -parity violating operators. Also we find that there is a degeneracy in scalar field parameters (γ, β) and coupling ζ value in producing the correct amplitude of CMB power spectrum. However the value of parameter $\tan(\beta) = 1.8$, which is tightly fixed from the requirement of PeV scale SUSY breaking, removes the degeneracy in the values of the scalar field parameters to provide a unique solution for inflation. In this way it brings the explanation for dark matter, PeV neutrinos and inflation within the same framework.

* Email: girish@ctp-jamia.res.in

† Email: khanphysics.123@gmail.com

‡ Email: mohanty@prl.res.in

I. INTRODUCTION

The 125 GeV Higgs boson found at the Large Hadron Collider (LHC) [1–3] completes the spectrum of the Standard Model but raises the question about what protects the mass of the higgs at the electroweak scale despite quantum corrections (the gauge hierarchy problem). Supersymmetry [4–6] has been widely accepted as the natural symmetry argument for protecting the Higgs and other scalar masses against radiative corrections and in addition provides the unification of the gauge couplings at the GUT scale and WIMP dark matter. The idea of naturalness in SUSY [7, 8] requires that to explain the higgs mass, loop corrections should not be too large compared to the tree level and should not rely on the cancellation of large corrections from different loop corrections. This puts upper bounds on the masses of squarks, gluons and gauginos for natural SUSY [8]. The searches for SUSY particles at LHC by ATLAS [9] and CMS [10] at 13 TeV and with 15 fb^{-1} of data do not find the SUSY partners and rule out the simplest models of natural SUSY [11, 12]. The idea of supersymmetry for explaining the Higgs mass without fine tuning can be abandoned while still retaining some of the positive features like coupling constant unification and WIMP dark matter in Split-SUSY [13] scenarios where squarks and gluinos are heavy evading the LHC bounds and the electro-weakinos are of the TeV scale providing the WIMP dark matter and coupling constant unification. With the tight bounds from the direct detection experiments [14, 15] the bino-higgsino dark matter of a few 100 GeV is getting increasingly difficult [16–25]. The evidence that there is need for new physics at PeV scale comes from the IceCube’s High Energy Starting Events (HESE) observations [26–29] of PeV energy neutrinos. The non-observation of neutrinos with deposited energy between 0.4-1 PeV and at Glashow resonance energies has called for an extra source of PeV energy neutrinos, and a popular scenario [30–35] is the decay of PeV scale dark matter with a sizable branching to neutrinos and with a lifetime of $\sim 10^{15} \text{ sec}$ [36] and $\sim 10^{28} \text{ sec}$ [37–50]. Leptoquarks have been used to explain IceCube PeV events in Refs. [51–56]. A PeV scale supersymmetry model with gauge coupling unification, light higgs (with fine tuning) and PeV dark matter was introduced in [57, 58]. It was found [39] that in order to obtain the large decay time of $\sim 10^{28} \text{ sec}$ as required from the observed IceCube neutrino flux, dimension 6 operators suppressed by the GUT scale had to be introduced in the superpotential.

A different motivation for Supersymmetry is its usefulness in inflation. The low upper bound on the tensor-to-scalar ratio by Planck [59–61] and BICEP2/Keck Array (BK15) [62, 63] rules out the standard particle physics models with quartic and quadratic potentials. One surviving model is the Starobinsky $R + R^2$ model which predicts a very low tensor-to-scalar ratio of $\sim 10^{-3}$. It was shown by Ellis et al [64] that by choosing the Kähler potential of the no-scale form one can achieve Starobinsky type plateau inflation in a simple Wess-Zumino model. No-scale model of inflation from F -term has been constructed in SO(10) [65, 66], SU(5) [67], NMSSM [68] and

MSSM [69, 70] models. Inflation models in Supergravity with F -term scalar potential were earlier considered in [71–75] and with a D -term scalar potential in [76–79]. The Higgs-sneutrino inflation along the D -flat directions in MSSM has also been studied in [80–86].

In this paper, we construct a D -term inflation model with no-scale Kähler potential, MSSM superpotential and an appropriate choice of gauge kinetic function in MSSM fields, which gives a Starobinsky like Higgs-sneutrino plateau inflation favored by observations. The SUSY breaking scale is a few PeV which provides a PeV scale bino as dark matter. A fraction of thermal relic density is obtained by turning on a small R -parity violation to give a decaying dark matter whose present density and neutrino flux at IceCube is tuned by choosing the R -parity violating couplings. Supersymmetry breaking is achieved by a hidden sector Polonyi field which takes a non-zero vev at the end of inflation. The gravitino mass is a few PeV which sets the SUSY breaking scale. The mSUGRA model has only five free parameters including soft SUSY breaking parameters. These are the common scalar mass m_0 , the common gaugino mass $m_{\frac{1}{2}}$, the common trilinear coupling parameter A_0 , the ratio of Higgs field $vevs$ $\tan\beta$ and the sign of mass parameter μ , all are given at the gauge coupling unification scale. The spectra and the couplings of sparticles at the electroweak symmetry breaking scale are generated by renormalization group equations (RGEs) of the above soft breaking masses and the coupling parameters. The sparticle spectrum at the low energy scale was generated using the publicly available softwares **FlexibleSUSY** [87, 88], **SARAH** [89, 90] and **SPheno** [91, 92] with the mSUGRA input parameters set m_0 , $m_{\frac{1}{2}}$, A_0 , $\tan\beta$ and $\text{sign}(\mu)$. In our analysis, we have used **SARAH** to generate model files for mSUGRA and relic density of LSP has been calculated using **micrOMEGAs** [93, 94].

R -parity conserving SUSY models include a stable, massive weakly interacting particle (WIPM) and the lightest supersymmetric particle (LSP), namely, neutralino which can be considered as a viable dark matter candidate. With the additional very tiny R -Parity violating (RPV) terms, LSP can decay to SM particles [56, 95–97], however it does not change the effective (co)annihilation cross-section appreciably in the Boltzmann equation [98–100]. In the context of mSUGRA model, the analysis of neutralino dark matter have been carried out in the Refs. [101–113], where SUSY breaking occurs in a hidden sector, which is communicated to the observable sector via gravitational interactions.

The paper is organized as follows. In Section II, we display the MSSM model and the specific form of a Kähler potential and gauge kinetic function which gives the Starobinsky like Higgs-sneutrino plateau inflation. The R -parity violating terms contribute to the F -term potential and to the decay of the dark matter for IceCube. We fix the parameter in the gauge kinetic function and show that a D -term inflation model consistent with the CMB observations is obtained. In Section III, we describe the SUSY breaking mechanism from the Polonyi field and calculate the soft breaking parameters m_0 , $m_{\frac{1}{2}}$, A_0 and gravitino mass $m_{\frac{3}{2}}$ as a function of parameters of the

Polonyi potential. In Section IV, we give the phenomenological consequences of the PeV scale SUSY model. We show that the coupling constant unification can take place and we obtain ~ 125 GeV Higgs mass by fine tuning. The relic density of the bino LSP is overdense after thermal decoupling but due to slow decay from the R -parity violation a fraction of relic density remains in the present epoch. A different R -parity operator is responsible for the decay of the PeV dark matter into neutrinos. We fix the parameters of the R -parity violating couplings to give the correct flux of the PeV scale neutrino events seen at IceCube. In Section V, we summarize the main results and give our conclusions.

II. D -TERM MSUGRA MODEL OF INFLATION

We consider the model with MSSM matter fields and R -parity violation and choose a Kähler potential $K(\phi_i, \phi_i^*, Z, Z^*)$ and gauge kinetic function $f_{ab}(\phi_i, Z)$ with the aim of getting a plateau inflation favored by observations. We choose K and f_{ab} of the form

$$K = -3 \ln \left[1 - \frac{1}{3} \left(H_u^\dagger H_u + H_d^\dagger H_d + L^\dagger L + Q^\dagger Q + \tilde{e}_R^* \tilde{e}_R + \tilde{u}_R^* \tilde{u}_R + \tilde{d}_R^* \tilde{d}_R \right) \right] + ZZ^* + \frac{\alpha}{2} (ZZ^*)^2, \quad (2.1)$$

$$f_{ab} = \frac{e^{-\kappa Z}}{1 + \zeta H_u \cdot H_d} \delta_{ab} \quad (2.2)$$

respectively. And the MSSM superpotential $W(\phi_i)$

$$W = \mu H_u \cdot H_d - Y_d Q \cdot H_d \tilde{d}_R + Y_u Q \cdot H_u \tilde{u}_R - Y_e L \cdot H_d \tilde{e}_R + \mu_z^2 Z + \mu_{zz} Z^3 \quad (2.3)$$

where,

$$H_u = \begin{pmatrix} \phi_u^+ \\ \phi_u^0 \end{pmatrix}, \quad H_d = \begin{pmatrix} \phi_d^0 \\ \phi_d^- \end{pmatrix}, \quad L = \begin{pmatrix} \phi_\nu \\ \phi_e \end{pmatrix}, \quad Q = \begin{pmatrix} u_L \\ d_L \end{pmatrix}, \quad (2.4)$$

and field Z is the hidden sector Polonyi which is introduced to break supersymmetry. The Polonyi field Z is associated with the fluctuations in the overall size of the compactified dimensions which has to be strongly stabilized at SUSY-breaking for the successful implementation of inflation in supergravity. The parameters α , κ , μ_z , μ_{zz} and ζ are coupling constants of the model to be fixed from SUSY breaking and inflation. The other fields bear their standard meanings. Also the above potentials are in the $M_p = (8\pi G)^{-1} = 1$ unit and shall use the same convention through out the analysis of this model.

In addition to the superpotential given in Eq. (2.3), we also consider the R -parity violating interaction terms

$$W_{int} = \lambda_{ijk} L_i L_j e_{Rk}^c + \lambda'_{ijk} L_i Q_j d_{Rk}^c + \frac{1}{2} \lambda''_{ijk} u_{Ri}^c d_{Rj}^c d_{Rk}^c \quad (2.5)$$

which will play role in explaining observed DM relic density and PeV neutrino flux at IceCube.

In supergravity, the scalar potential depends upon the Kähler function $G(\phi_i, \phi_i^*)$ given in terms of superpotential $W(\phi_i)$ and Kähler potential $K(\phi_i, \phi_i^*)$ as $G(\phi_i, \phi_i^*) \equiv K(\phi_i, \phi_i^*) + \ln W(\phi_i) + \ln W^*(\phi_i^*)$, where ϕ_i are the chiral scalar superfields. In $\mathcal{D} = 4$, $\mathcal{N} = 1$ supergravity, the total tree-level supergravity scalar potential is given as the sum of F -term and D -term potentials which are given by

$$V_F = e^G \left[\frac{\partial G}{\partial \phi^i} K_{j*}^i \frac{\partial G}{\partial \phi_j^*} - 3 \right] \quad (2.6)$$

and

$$V_D = \frac{1}{2} [\text{Re } f_{ab}]^{-1} D^a D^b, \quad (2.7)$$

respectively, where $D^a = -g \frac{\partial G}{\partial \phi_k} (\tau^a)_k^l \phi_l$ and g is the gauge coupling constant corresponding to each gauge group and τ^a are corresponding generators. For $SU(2)_L$ symmetry $\tau^a = \sigma^a/2$, where σ^a are Pauli matrices. And the $U(1)_Y$ hypercharges of the fields $H_u, H_d, L, Q, \tilde{e}_R, \tilde{d}_R, \tilde{u}_R$ given in (2.3) are $Y = (\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{6}, 1, \frac{1}{3}, -\frac{2}{3})$ respectively. The quantity f_{ab} is related to the kinetic energy of the gauge fields and is a holomorphic function of superfields ϕ_i . The kinetic term of the scalar fields is given by

$$\mathcal{L}_{KE} = K_i^{j*} \partial_\mu \phi^i \partial^\mu \phi_j^*, \quad (2.8)$$

where K_{j*}^i is the inverse of the Kähler metric $K_{j*}^i \equiv \partial^2 K / \partial \phi^i \partial \phi_j^*$. We assume that during inflation, SUSY is unbroken and the hidden sector field is subdominant compared to inflaton field so that we can safely assume $Z = 0$. Also, for charge conservation, we assume that during inflation the charged fields take zero vev. The D -term and F -term potentials are obtained as

$$V_D = \frac{9}{8} (g_1^2 + g_2^2) \frac{(|\phi_d|^2 - |\phi_u|^2 + |\phi_\nu|^2)^2 (-1 + \zeta |\phi_d|^2)^2}{(-3 + |\phi_d|^2 + |\phi_u|^2 + |\phi_\nu|^2)^2}, \quad (2.9)$$

$$V_F = \frac{3\mu^2 (3|\phi_u|^2 + |\phi_d|^2 (3 - |\phi_u|^2))}{(-3 + |\phi_d|^2 + |\phi_u|^2 + |\phi_\nu|^2)^2}. \quad (2.10)$$

It can be seen that the above expressions reduces to MSSM D -term and F -term potentials in the small field limit after the end of inflation as the terms coming from Kähler potential and gauge kinetic function are Planck suppressed. Now, for simplification, we parametrize the neutral component fields as

$$\phi_u^0 = \phi \sin[\beta], \quad \phi_d^0 = \phi \cos[\beta], \quad \phi_\nu = \gamma \phi. \quad (2.11)$$

For the above parametrization, the kinetic term turns out to be

$$\mathcal{L}_{KE} = \frac{9(1 + \gamma^2)}{(-3 + (1 + \gamma^2)|\phi|^2)^2} |\partial_\mu \phi|^2. \quad (2.12)$$

To obtain the canonical kinetic term for the inflaton field and to better understand the inflation potential, we redefine the field ϕ to χ_c via

$$\phi = \frac{\sqrt{3}}{\sqrt{1 + \gamma^2}} \tanh\left(\frac{\chi_c}{\sqrt{3}}\right). \quad (2.13)$$

For the above field redefinition, the kinetic term becomes

$$\mathcal{L}_{KE} = \text{sech}^2\left(\frac{2\text{Im}[\chi_c]}{\sqrt{3}}\right) |\partial_\mu \chi_c|^2, \quad (2.14)$$

therefore, if the imaginary part of the field χ_c is zero, we obtain the canonical kinetic term in real part of field $\chi_c = \chi$ (say). As the field ϕ is a linear combination of the Higgs and sneutrino field, so is the inflaton field χ , we can call this model a Higgs-sneutrino inflation model. The D -term potential (2.9) in the canonical inflaton field χ becomes

$$V_D = \frac{1}{2} \lambda^2 (g_1^2 + g_2^2) \tanh^4\left(\frac{\chi}{\sqrt{6}}\right), \quad (2.15)$$

where $\lambda = \frac{3}{2} \frac{\gamma^2 + \cos(2\beta)}{1 + \gamma^2}$ and we have made a specific choice $\zeta = \frac{\gamma^2 + 1}{3 \sin(\beta) \cos(\beta)}$ which is critical to obtain a plateau behavior potential at large field values which can fix inflationary observables. Since at the GUT scale, the mass parameter $\mu \sim 0$ and SU(2) gauge couplings are $g_1 = g_2 = 0.62$, the D -term potential dominates over F -term potential during inflation when the field values are at the Planck scale. With the canonical kinetic term and scalar potential obtained in canonical inflaton field χ , the theory is now in the Einstein frame. Therefore, we can use the standard Einstein frame relations to estimate the inflationary observables, namely, amplitude of the curvature perturbation $\Delta_{\mathcal{R}}^2$, scalar spectral index n_s and its running α_s , and tensor-to-scalar ratio r , given by

$$\Delta_{\mathcal{R}}^2 = \frac{1}{24\pi^2} \frac{V_D}{\epsilon}, \quad (2.16)$$

$$n_s = 1 - 6\epsilon + 2\eta, \quad (2.17)$$

$$\alpha_s \equiv \frac{dn_s}{d \ln k} = 16\epsilon\eta - 24\epsilon^2 - 2\xi, \quad (2.18)$$

$$r = 16\epsilon, \quad (2.19)$$

respectively. Here ϵ , η and ξ are the slow-roll parameters, given by

$$\epsilon = \frac{1}{2} \left(\frac{V'_D}{V_D} \right)^2, \quad \eta = \frac{V''_D}{V_D}, \quad \xi = \frac{V'_D V'''_D}{V_D^2}. \quad (2.20)$$




FIG. 1. The D -term inflation potential for $\tan(\beta) = 1.8$, $g_1 = g_2 = 0.62$ is shown. During inflation as the inflaton rolls down from $\chi_i \simeq 7$ to $\chi_e \simeq 1.9$, the potential along the Z direction stays nearly flat implying inflaton χ being much heavier than the Polonyi field Z . For $Z \gtrsim 1$, the potential becomes very steep where slow-roll inflation cannot be achieved.




FIG. 2. After the end of inflation as χ approaches minimum near zero, Polonyi field Z settles down to minimum at $Z_{min} \simeq \pm 0.144$. The positive Polonyi potential is obtained for the parameter values $\alpha = 20$, $\mu_z \simeq 2.43 \times 10^{-6}$, $\mu_{zz} \simeq 2.858 \times 10^{-11}$, $\kappa = 0.56$, which are fixed from the requirement of getting PeV scale soft SUSY breaking parameters.

In order to have flat Universe as observed, the universe must expand at least by more than 60 e -folds during inflation. The displacement in the inflaton field during inflation is $\Delta\chi \equiv \chi_i - \chi_f$.

The field value χ_i is at the onset of inflation, when observable CMB modes starts leaving the horizon, can be determined using the following relation

$$N = \int_{\chi_f}^{\chi_i} \frac{V_D}{V'_D} d\chi, \quad (2.21)$$

once we fix the field value χ_f using the condition $\epsilon(\chi_f) = 1$ which corresponds to end of inflation.

From the Planck-2018 CMB temperature anisotropy data in combination with the EE measurement at low multipoles, we have the scalar amplitude, the spectral index and its running as $\ln(10^{10}\Delta_{\mathcal{R}}^2) = 3.044 \pm 0.014$, $n_s = 0.9649 \pm 0.0042$, $\alpha_s = -0.0045 \pm 0.0067$, respectively, at (68% CL, PlanckTT,TE,EE+lowE+lensing) [59–61]. Also the Planck-2018 data combined with BK15 CMB polarization data put an upper bound on tensor-to-scalar ratio $r_{0.002} < 0.056$ (95% CL, PlanckTT,TE,EE+lowE+lensing+BK15) [62, 63]. Armed with the theoretical and observational results, we perform the numerical analysis of the models. From the condition of end of inflation $\epsilon = 1$, we find $\chi_e \simeq 1.927$. Therefore requiring $N \simeq 60$ e -folds expansion during inflation, we obtain $\chi_i \simeq 7.03$. At $\chi_i \simeq 7.03$, CMB observables are estimated to be $r \simeq 0.0035$, $n_s \simeq 0.965$ and $\alpha_s \simeq -6 \times 10^{-4}$, consistent with the observations. The (n_s, r) predictions are similar to Starobinsky inflation. For $\tan(\beta) = 1.8$ and $g_1 = g_2 = 0.62$, the observed CMB amplitude can be obtained

for $\lambda \simeq 1.703 \times 10^{-5}$. From the specific case considered in (2.15) for successful inflation, we have sneutrino field parameter $\gamma \simeq 0.727$, and coupling ζ of the Planck suppressed operator in the gauge kinetic function (2.2) $\zeta \simeq 1.2$. The evolution of the D -term inflaton potential (2.15) is shown in Fig. 1. During inflation as the inflaton rolls down from $\chi_i \simeq 7$ to $\chi_e \simeq 1.9$, the potential along the Z direction stays nearly flat implying χ being much heavier than Z . During slow-roll phase, the mass of inflaton varies from $m_{\chi_i} \simeq 2.1 \times 10^{13}$ GeV to $m_{\chi_e} \simeq 9.3 \times 10^{12}$ GeV, whereas the mass parameter associated with the Polonyi field Z varies from $m_{Z_i} \simeq 6.1 \times 10^{10}$ GeV to $m_{Z_e} \simeq 2.1 \times 10^8$ GeV. Hence, $m_Z^2 \ll m_\chi^2$, our assumption that Z is subdominant during inflation and contributes insignificantly to the supergravity inflation potential.

In the next Section we will see that the value of $\tan(\beta) = 1.8$ used to fix CMB amplitude is absolutely critical in order to get PeV dark matter whose decay explain PeV neutrino events observed at IceCube, and reproducing low energy particles mass spectrum as shown in Table II. It is important to mention here that the scalar field parameter values $\tan(\beta) = 1.8$ and $\gamma \simeq 0.727$ is not a unique set of value to obtain the observed CMB amplitude. Instead it can be obtained for all the pair of values of (γ, β) which satisfy the relation $\frac{3}{2} \frac{\gamma^2 + \cos(2\beta)}{1 + \gamma^2} = \lambda \simeq 1.703 \times 10^{-5}$ defined in (2.15). However, $\tan(\beta) = 1.8$ fixed from the requirement of PeV scale SUSY breaking removes this degeneracy in (γ, β) in obtaining the correct CMB amplitude and, at the same time, it brings the successful explanation for inflation, dark matter relic density and PeV neutrino events at IceCube within the same framework.

III. SUSY BREAKING

We assumed that during inflation all the fields including the hidden sector field Z are subdominant compared to inflaton field and supersymmetry is unbroken at the time of inflation. Once the inflation ends, the scalar fields effectively become vanishing and the soft mass terms are generated via SUSY breaking as the field Z settles down to a finite minimum of the Polonyi potential. The Polonyi field Z is associated with the fluctuations in the overall size of the compactified dimensions. For the successful implementation of inflation in supergravity the strong stabilization of the Polonyi field is required which can be achieved via appropriate choice of Kähler potential, gauge kinetic function and superpotential in Z . This also allows for a solution of the cosmological Polonyi problem [114, 115] (which is a special case of cosmological moduli problem [116, 117]) associated with the problem of Cosmological Nucleosynthesis. Technically, cosmological Polonyi problem is evaded if Z is heavy $m_Z \sim \mathcal{O}(100 - 1000)$ TeV and the Polonyi mass is much larger than the gravitino mass, *i.e* $m_Z^2 \gg m_{3/2}^2$ [70, 118, 119]. We will see that this problem does not occur in this model.

The appropriately chosen potentials $K(Z, Z^*)$ and $W(Z)$ are shown in eq.(2.1) and eq.(2.3),

and the gauge kinetic function in eq.(2.2). The kinetic term for Polonyi field Z is obtained as $(1 + 2\alpha|Z|^2)|\partial_\mu Z|^2$ and the Polonyi potential is obtained as

$$V(Z, Z^*) = e^{|Z|^2 + \frac{\alpha}{2}|Z|^4} |Z|^2 |\mu_{zz} Z^2 + \mu_z^2|^2 \left[-3 + \frac{1}{1 + 2\alpha|Z|^2} \left| \frac{1}{Z} + Z^* + \alpha Z Z^{*2} + \frac{2\mu_{zz} Z}{\mu_{zz} Z^2 + \mu_z^2} \right|^2 \right]. \quad (3.1)$$

We take the hidden sector Polonyi field Z to be real. The parameters α , μ_z and μ_{zz} are fixed from the requirement that Z which breaks supersymmetry acquire a minima where the gravitino mass is $\sim \mathcal{O}(\text{PeV})$ scale and the Polony potential is positive $V_F(Z) \gtrsim 0$ with a field minimum at $Z = Z_{min}$. We find gravitino mass to be

$$m_{\frac{3}{2}}^2 = e^G = e^{Z^2 + \frac{\alpha}{2}Z^4} (\mu_{zz} Z^3 + \mu_z^2 Z)^2, \quad (3.2)$$

and the scalar masses, using $m_\phi^2 = \frac{\partial_\phi \partial_{\phi^*} (V_F(\phi, \phi^*, Z))}{\partial_\phi \partial_{\phi^*} K}$ evaluated at $\phi = 0$, $Z = Z_{min}$, are obtained as

$$m_{\phi_i}^2 = m_{\frac{3}{2}}^2 \left[-2 + \frac{1}{1 + 2\alpha Z^2} \left(\frac{1}{Z} + Z + \alpha Z^3 + \frac{2\mu_{zz} Z}{\mu_{zz} Z^2 + \mu_z^2} \right)^2 \right], \quad (3.3)$$

$$m_{\phi_{(u,d)}}^2 = m_{\phi_i}^2 + \frac{\mu^2}{(\mu_{zz} Z^3 + \mu_z^2 Z)^2} m_{\frac{3}{2}}^2 \quad (3.4)$$

where $\phi_i = \phi_\nu, \phi_e, \phi_u^+, \phi_d^-, \tilde{u}_L, \tilde{d}_L, \tilde{u}_R, \tilde{d}_R, \tilde{e}_R$. At the GUT energy scale, $\mu \sim 0$, all the scalar masses are equal $m_{\phi_{(u,d)}}^2 = m_{\phi_i}^2$.

Now we calculate the coefficients of the soft SUSY breaking terms which arise from the Kähler potential (2.1) and superpotential (2.3). The effective potential of the observable scalar sector consists of soft mass terms which give scalar masses as given by Eq.s (3.3)-(3.4), and trilinear and bilinear soft SUSY breaking terms given by [120–122]

$$\frac{1}{3} A_{ijk} \phi^i \phi^j \phi^k + \frac{1}{2} B_{ij} \phi^i \phi^j + \text{h.c.},$$

here the coefficients A_{ijk} and B_{ij} given by

$$A_{ijk} = \left[\frac{\partial_{Z^*} \hat{W}^* + \hat{W}^* \partial_{Z^*} \hat{K}}{\partial_{Z^*} \partial_Z \hat{K}} \partial_Z e^{\hat{K}} \right] \tilde{Y}_{ijk}, \quad (3.5a)$$

$$B_{ij} = \left[\frac{\partial_{Z^*} \hat{W}^* + \hat{W}^* \partial_{Z^*} \hat{K}}{\partial_{Z^*} \partial_Z \hat{K}} \partial_Z e^{\hat{K}} - m_{\frac{3}{2}} e^{\hat{K}/2} \right] \tilde{\mu}_{ij}, \quad (3.5b)$$

From Eqs. (2.1) and (2.3), we have $\hat{K}(Z, Z^*) = ZZ^* + \frac{\alpha}{2}(ZZ^*)^2$ and $\hat{W}(Z) = \mu_z^2 Z + \mu_{zz} Z^3$. The coefficients of normalised masses $\tilde{\mu}_{ij}$ and Yukawa couplings \tilde{Y}_{ijk} in the trilinear (3.5a) and bilinear (3.5b) terms, respectively, are obtained as

$$A_0 = e^{Z^2 + \frac{\alpha}{2}Z^4} \frac{Z(1 + Z^2\alpha) [\mu_{zz}Z^2(3 + Z^2 + Z^4\alpha) + (1 + Z^2 + Z^4\alpha)\mu_z^2]}{1 + 2\alpha Z^2}, \quad (3.6)$$

$$B_0 = A_0 - e^{Z^2 + \frac{\alpha}{2}Z^4} (\mu_{zz}Z^3 + \mu_z^2 Z). \quad (3.7)$$

We drop the B_{ij} contribution to the scalar masses by taking corresponding $\tilde{\mu}_{ij}$ small. Also from the fermionic part of the SUGRA Lagrangian, the soft gaugino masses can be obtained as [122–124]

$$m_{\frac{1}{2}} = \frac{1}{2} [\text{Re } f_{ab}]^{-1} e^{G/2} \partial_I f_{ab} \hat{K}^{IJ*} G_{J*}. \quad (3.8)$$

For the choice of gauge kinetic function $f_{ab} = e^{-\kappa Z} \delta_{ab}$ when $\phi \approx 0$ after inflation, we obtain the gaugino mass

$$m_{\frac{1}{2}} = \frac{\kappa}{2} m_{\frac{3}{2}} (1 + 2\alpha Z^2) \left(Z + \alpha Z^3 + \frac{\mu_z^2 + 3\mu_{zz}Z^2}{\mu_z^2 Z + \mu_{zz}Z^3} \right). \quad (3.9)$$

We show the Polonyi potential (3.1) in Fig.2 for the parameters $\alpha = 20$, $\mu_z \simeq 2.43 \times 10^{-6}$, $\mu_{zz} \simeq 2.858 \times 10^{-11}$, $\kappa = 0.56$ has a minimum at $Z_{min} \simeq \pm 0.144$. These parameter values are fixed at the GUT scale from the requirement to achieve the soft SUSY breaking parameters

$$A_0 = -2.2 \text{ PeV}, \quad m_0 = m_\phi = 14 \text{ PeV}, \quad m_{\frac{1}{2}} \simeq 10 \text{ PeV}.$$

with the specific choice of $\tan\beta = 1.8$ and sign of $\mu > 0$ which gives the Higgs mass ~ 125 GeV at electroweak scale satisfying all experimental (LHC etc) and theoretical constraints (stability, unitarity etc). The gravitino mass and the mass of the Polonyi field comes out to be $m_{\frac{3}{2}} \simeq 2.32$ PeV and $m_z \simeq 31.8$ PeV, respectively, which implies $m_Z^2 \gg m_{3/2}^2$ and therefore the $\mathcal{O}(\text{PeV})$ scale oscillations of the Polonyi field near its minimum decay much before the Big Bang Nucleosynthesis. This leads to strong stabilization of Z and therefore cosmological Polonyi problem does not occur in this model.

Knowing the soft breaking parameters at a high energy scale does not tell us anything about the phenomenology we could observe in the experiments such as LHC, direct detection of DM, IceCube, etc. We need to find these parameters at the low energy. In general all parameters appearing in the supersymmetric Lagrangian evolve with RGEs. These RGEs are the inter-mediator between the unified theory at GUT scale and the low-energy masses and couplings, which strictly depend on the boundary conditions. We apply RGEs to calculate low-energy masses and different branching ratios for the above mentioned set of mSUGRA parameters. As the RGEs are coupled differential equations, which cannot be solved analytically. Also the low-energy phenomenology is complicated due to mixing angles and dependence of couplings on the high-scale parameters, therefore one has to rely on numerical techniques to solve RGEs. We calculate all the variables as allowed by the present experimental data. There are various programs publicly available such as **SARAH**, **SPheno**,

Suspect which can generate two-loop RGEs and calculate the mass spectrum and the couplings at low-energy. For this work, we use **SARAH-4.11.0** [89, 90] to generate the RGEs and other input files for **SPheno-4.0.2** [91, 92] which generate the mass spectrum, couplings, branching ratios and decay widths of supersymmetric particles. To study the neutralino as a dark matter candidate we link the output files of **SPheno** and model files from **SARAH** to **micrOMEGAs-3.6.8** [93, 94] to calculate the number density for a PeV neutralino dark matter. In the next section we discuss the consequences of a PeV dark matter.

IV. BINO DOMINATED DM IN MSUGRA MODEL

The neutralinos χ_i ($i=1,2,3,4$) are the physical superpositions of two gauginos namely bino \tilde{B} and wino \tilde{W}_3 , and two Higgsinos \tilde{H}_{u0} and \tilde{H}_{d0} . The neutralino mass matrix is given by,

$$M_N = \begin{pmatrix} M_1 & 0 & -M_Z c_b s_\theta & M_Z s_b s_\theta \\ 0 & M_2 & M_Z c_b c_\theta & -M_Z s_b c_\theta \\ -M_Z c_b s_\theta & M_Z c_b c_\theta & 0 & -\mu \\ M_Z s_b s_\theta & -M_Z s_b c_\theta & -\mu & 0 \end{pmatrix}$$

where $c_b \equiv \cos\beta$, $c_\theta \equiv \cos\theta_W$ and θ_W is the Weinberg mixing angle. M_1 and M_2 are the bino and wino mass parameter at the EWSB scale. The lightest eigenvalue of the above matrix and the corresponding mass eigenstate has good chance of being the LSP. The Higgsinos dominantly contribute in LSP for $|\mu| < M_{1,2}$, whereas for $|\mu| > M_{1,2}$, the LSP can be determined by bino and wino. The lightest neutralino becomes bino-like when $M_2 > M_1$. In mSUGRA model, M_1 and M_2 are equal due to the universality of the gaugino masses at the GUT scale where the value of the gauge coupling constants g_1 , g_2 and g_3 become equal, see Fig. 3. However at low energy $M_2 \approx \frac{5g_2^2(M_Z)}{3g_1^2(M_Z)} M_1$ [110]. This implies that the LSP is mostly bino-like.

A. Relic density and PeV excess at IceCube

In this analysis we present our mSUGRA model with the aim to realize a PeV scale dark matter candidate, which can also explain the IceCube HESE events. IceCube [26–29] recently reported their observation of high-energy neutrinos in the range 30 TeV – 2 PeV. The observation of neutrinos are isotropic in arrival directions. No particular pattern has been identified in arrival times. This implies that the source is not local but broadly distributed. This superheavy neutrino (LSP) might be the dark matter, distributed in the Galactic halo. In this work we find that the thermally produced LSP can serve as a viable PeV dark matter candidate satisfying the dark




FIG. 3. Solid lines represent the gauge coupling evolution for the standard model. Dotted lines represent the gauge coupling unification for our benchmark points (Table I).

SUGRA parameters				RPV couplings	DM mass in PeV	Decay time in (secs), Explains IceCube- HESE events
Masses in PeV			$\tan\beta$			
m_0	$m_{\frac{1}{2}}$	A_0			λ_{121}	
14	10	-2.2	1.8	2.82×10^{-28}	6.3	8.90×10^{28}

TABLE I. A list of the Benchmark points (BP) as used in our analysis. Using these BPs, we have reproduced the Higgs mass at ~ 125 GeV and explain the relic density and the PeV neutrino events seen at the IceCube experiment.

matter relic density in the right ball park which also explains the IceCube excess at PeV. However, an annihilating PeV scale dark matter candidate poses two serious problems.

(1). To maintain the correct relic density using the well-known thermal freeze-out mechanism requires a very large annihilation cross-section, which violates the unitarity bound. The s-wave

SUSY Fields		Masses in PeV			
$\tilde{d}_{iL,R}$ ($i = d, s, b$)	$M_{\tilde{d}_{iL,R}} \approx (15.75, 16.78)_d (16.78, 16.78)_s (17.38, 17.38)_b$				
$\tilde{u}_{iL,R}$ ($i = u, c, t$)	$M_{\tilde{u}_{iL,R}} \approx (13.30, 15.75)_u (16.88, 16.88)_c (17.38, 17.38)_t$				
$\tilde{e}_{iL,R}$ ($i = e, \mu, \tau$)	$M_{\tilde{e}_{iL,R}} \approx (14.36, 14.36)_e (14.36, 14.86)_\mu (14.86, 14.86)_\tau$				
$\tilde{\nu}_i$ ($i = e, \mu, \tau$)	$M_{\tilde{\nu}_i} \approx (14.86)_e (14.86)_\mu (14.86)_\tau$				
h	$M_h \approx 126$ [GeV]				
H	$M_H \approx 19.48$				
A	$M_A \approx 19.48$				
H^-	$M_{H^-} \approx 19.47$				
\tilde{g}_i ($i = 1 \dots 8$)	$M_{\tilde{g}_i} \approx 14.28$				
$\tilde{\chi}^0$ ($i = 1 \dots 4$)	$M_{\tilde{\chi}_1}(DM) \approx 6.3$ $M_{\tilde{\chi}_2} \approx 7.22$ $M_{\tilde{\chi}_3} \approx 7.22$ $M_{\tilde{\chi}_4} \approx 9.22$				
$\tilde{\chi}^-$	$M_{\tilde{\chi}_1^-} \approx 7.22$ $M_{\tilde{\chi}_2^-} \approx 9.21$				

TABLE II. Particle mass spectrum after EWSB for the benchmark point given in Table I.

annihilation cross-section of dark matter with a mass M_{DM} is limited by unitarity as [125],

$$\langle \sigma v \rangle \lesssim \frac{4\pi}{M_{DM}^2 v}. \quad (4.1)$$

The unitarity bound restrict the dark matter mass below 300 TeV [125]. Also, in order to satisfy the unitarity bound, the PeV scale dark matter model produces an overabundance $\mathcal{O}(10^8)$ (depends on the model) of the dark matter.

(2). The decay time of a particle is in general inversely proportional to its mass. So a PeV scale particle is generically too short-lived to be a dark matter candidate. One has to use fine-tuning [126, 127] to stabilize the dark matter as lifetime of the DM particles has to be at least larger than the age of the Universe [29, 128].

Generally, thermal dark matter freeze-out depends on the remaining dark matter in chemical and thermal equilibrium with the SM bath, which leads to depletion of dark matter through Boltzmann suppression [129, 130]. We consider the possibility that the dark matter can also decay out of equilibrium to the SM particles via R -parity violation. In the presence of constant s-wave effective annihilation cross-section and dark matter decay, the Boltzmann equation is given by,

$$\frac{dn}{dt} = -3Hn - \langle \sigma v \rangle (n^2 - n_{eq}^2) - n \sum_i \frac{1}{\tau_{i,DM}}. \quad (4.2)$$

Here, we assume that the decay rates of super partner of SM particles other than LSP are much faster than the rate of the expansion of the Universe, so that all the particles present at the beginning of the Universe have decayed into the lightest neutralino before the freeze-out. Therefore

the density of the lightest neutralino n is the sum of the density of all SUSY particles. $\tau_{i,DM}$ is the decay time for the i^{th} process. In non-relativistic case $T < M_{DM}$, the equilibrium number density n_{eq} is given by the classical Maxwell-Boltzmann distribution,

$$n_{eq} = \left(\frac{M_{DM} T}{2\pi} \right)^{3/2} \text{Exp} \left(-\frac{M_{DM}}{T} \right). \quad (4.3)$$

The entropy density of the Universe at temperature T is $s = g_* T^3 \left(\frac{2\pi^2}{45} \right)$, where parameter g_* is the effective degrees of freedom. We use the relation of entropy density and the expansion rate of the Universe $\frac{ds}{dt} + 3Hs = 0$, in the eqn. 4.2, to obtain

$$\frac{dY}{dt} = s \langle \sigma v \rangle (Y^2 - Y_{eq}^2) - Y \sum_i \frac{1}{\tau_{i,DM}}, \quad (4.4)$$

where, yield Y is defined as $Y \equiv \frac{n}{s}$. We solve the above equation for the decaying LSP with large decay time. Before freeze-out annihilation term in eqn. 4.4 dominates over the exponential decay term due to very large decay time of LSP. Therefore, integrating eqn. 4.4 between the times $t = 0$ (or $T = \infty$) the start of the Universe and $t = t_f$ (or $T = T_f$) the freeze-out time, the yield $Y(t_f)$ comes out to be inversely proportional to the thermally averaged effective annihilation cross section $\langle \sigma v \rangle$ of the dark matter [131]. However after freeze-out, the annihilation of LSP is no longer large enough and decay term becomes dominant. Therefore, we neglect the annihilation term compared to the decay term, and integrate the remaining equation between the freeze-out time t_f and the present age of the Universe τ_U , which, for $t_U \gg t_f$, give the yield today as

$$Y(T_0) = Y(T_f) \text{Exp} \left(-\sum_i \frac{t_U}{\tau_{i,DM}} \right) \quad (4.5)$$

which implies that the number density of LSP reduces with time. Therefore, the relic abundance of LSP in the present Universe can be written as,

$$\Omega = \frac{s_0}{\rho_c} M_{DM} Y(T_0) = \frac{s_0}{\rho_c} M_{DM} Y(T_f) \text{Exp} \left(-\sum_i \frac{t_U}{\tau_{i,DM}} \right) \quad (4.6)$$

where $s_0 \sim 2890 \text{ cm}^{-3}$ is the current entropy density and $\rho_c \sim 1.05 \times 10^{-5} h^2 \text{ GeV cm}^{-3}$ is critical density of the present Universe, $h = 0.72$ is a dimensionless parameter defined through the Hubble parameter $H_0 = 100 h \text{ km sec}^{-1} \text{ Mpc}^{-1}$. This implies that the relic density is related to the annihilation cross-section and the decay life time as

$$\Omega h^2 \simeq \frac{9.62 \times 10^{-28} [\text{cm}^3 \text{sec}^{-1}]}{\langle \sigma v \rangle} \text{Exp} \left(-\sum_i \frac{t_U}{\tau_{i,DM}} \right) \quad (4.7)$$

Using particles spectrum (see table II) in micrOMEGAs, we get

$$\langle\sigma v\rangle = 6.78 \times 10^{-26} \text{ cm}^3\text{sec}^{-1}. \quad (4.8)$$

The main contribution to the DM effective annihilation cross-section comes from the co-annihilation channel (see Fig. 4) $\chi_1\chi_1^\pm \rightarrow ZW^\pm$ which is consistent with unitarity bound [125] and gives the bino-type DM density $\Omega h^2 \simeq 0.0142 \text{Exp}\left(-\sum_i \frac{t_U}{\tau_{i,DM}}\right)$. The joint results of CMB observations from WMAP and Planck Collaboration [132] give $\Omega h^2 = 0.1198 \pm 0.0026$. It has been shown that if a subdominant fraction of DM is decaying [36, 133], then it can resolve the conflict between σ_8 and H_0 which exists in Λ CDM model. Berezhiani et. al. have shown that if $\sim 5\%$ of the dark matter has a decay lifetime of 1.6×10^{16} seconds resolves the $\sigma_8 - H_0$ conflict [133]. For the PeV DM Anchordoqui et. al. have shown that $\sim 5\%$ of DM of mass scale 76 PeV and lifetime 6×10^{15} seconds is required for explaining the IceCube events [36]. In this paper, we assume the lifetime of DM decay to leptons via the R -parity violation LLe^c operator is 8.9×10^{28} seconds which gives 11% (i.e. $\Omega h^2 \simeq 0.0142$) of the DM relic density and is consistent with the structure formation. The neutrino flux is obtained by this decay channel to give the PeV neutrino flux required to explain the IceCube's HESE data [26–29]. We will explain this result in the subsection IV B in more detail.




FIG. 4. Co-annihilation diagram of lightest neutralino χ_1 (DM) with the charginos χ_1^\pm .

The presence of R -parity violating couplings 2.5 can explain the high energy neutrino events at IceCube. We take only one non-zero dimensionless trilinear R -parity violating couplings namely λ_{121} coefficient of $L_e L_\mu e^c$. λ_{121} helps in producing the neutrino flux through the decay of the Neutralino (see Fig. 5). We find the decay width for the dark matter decaying into $e^+\mu^-\nu_e$ and $e^-\mu^+\nu_e$ [134] as

$$\Gamma(\chi_1 \rightarrow e^+\mu^-\nu_e) = \frac{\kappa_{\chi_1 e^+ \bar{e}^-}^2 \lambda_{121}^2 M_{\chi_1}^5}{(8\pi)^3 M_{\bar{e}^-}^4} \quad \text{and} \quad \Gamma(\chi_1 \rightarrow e^-\mu^+\nu_e) = \frac{\kappa_{\chi_1 \mu^+ \bar{\mu}^-}^2 \lambda_{121}^2 M_{\chi_1}^5}{(8\pi)^3 M_{\bar{\mu}^-}^4} \quad (4.9)$$

where coupling $\kappa_{\chi_1 e^+ \bar{e}^-} = \kappa_{\chi_1 \mu^+ \bar{\mu}^-} \sim \sqrt{2}g_2$ [4, 5], $M_{\chi_1} \simeq 6.3$ PeV and $M_{\bar{e}^-, \bar{\mu}^-} \simeq 14.36$ PeV (see the Table II). For the lepton number violating coupling $\lambda_{121} \approx 2.82 \times 10^{-28}$, we get the decay time for this channel $\tau_{\nu,DM} \approx 8.9 \times 10^{28}$ sec. It is to be noted that we need very tiny λ 's to explain the IceCube HESE data. It indeed has the fine-tuning issue and the dynamical explanation lies somewhere else. To explain IceCube HESE data in terms of decaying dark matter, fine-tuning actually seems to be the most "natural" option [56, 95–97].




FIG. 5. Decay diagrams of neutralino via lepton number violating coupling which produces the Ice-cube flux.

B. Fitting the IceCube data

In this subsection, we fit the flux of neutrinos observed at IceCube from the decay of the neutralino LSP. The total contribution to the neutrino flux from the atmospheric background and the astrophysical sources along with the galactic (G) DM halo and extragalactic (EG) DM is given by,

$$\frac{d\Phi_{tot}}{dE_\nu} = \frac{d\Phi_{Atm}}{dE_\nu} + \frac{d\Phi_{astro}}{dE_\nu} + \frac{d\Phi_G}{dE_\nu} + \frac{d\Phi_{EG}}{dE_\nu} \quad (4.10)$$

Following the analysis as in Ref. [39] for our mSUGRA model parameters, we compute the number of neutrino events as a function of deposited energy

$$N_{bin} = T_E \int_{E_{min}^{bin}(E_{dep})}^{E_{max}^{bin}(E_{dep})} A(E_\nu) \frac{d\Phi_{tot}}{dE_\nu} dE_{dep}(E_\nu), \quad (4.11)$$

where $T_E = 2635$ is the total exposure time, E_{dep} is the deposited energy in the laboratory frame and $A(E_\nu)$ is the neutrino effective area for particular lepton flavor [135]. We sum over all the neutrino flavors. Due to low statistical data points at IceCube, it is acceptable to assume that the two energies coincide, i.e., $E_{dep} \simeq E_\nu$ [41].

We have extracted the atmospheric background data from Ref. [136] and added it to the data from astrophysical sources which come mainly due to the decay of highly energetic pions. The source of these pions could be known sources like active galactic nuclei or the Supernova remnants. We use the power-law flux $E_\nu^2 \frac{d\Phi_{astro}}{dE_\nu} = \Phi_0 (E_\nu)^{-\gamma}$ with $\Phi_0 = 2.2 \times 10^{-8} \text{ cm}^{-2} \text{ sec}^{-1} \text{ sr}^{-1}$ and $\gamma = 0.8$ [41] to fit the data from the astrophysical sources. Both the galactic and the extragalactic neutrino flux contributions are important to explain the excess neutrino flux around 1-2 PeV [39–43, 53, 135]. We take NFW (Navarro, Frenk, and White) DM density profile to account for the galactic dark matter contribution, where the neutrino energy spectrum $\frac{dN}{dE_\nu}$ evaluated at our model parameter




FIG. 6. Normalized neutrino events distribution is shown. The contributions to PeV neutrino flux from the galactic and the extragalactic DM decay is shown by the green region. The total flux from all the sources is shown by the blue line.

values has been used [137]. We present the neutrino events distribution in Fig. 6. The contribution to the neutrino flux from the DM decay can explain the PeV excess at IceCube.

V. CONCLUSIONS

In this paper we have constructed a Starobinsky like Higgs-sneutrino model of plateau inflation from supergravity D -term in MSSM fields, and 6 PeV bino-type dark matter gives the observed flux of PeV neutrino events at IceCube HESE. A subdominant fraction ($\sim 11\%$) of the DM relic density and its decay to neutrinos is obtained by choosing the couplings of the R -parity violating operators. The SUSY breaking is obtained by Polonyi field which sets the scale of soft SUSY parameters m_0 , $m_{\frac{1}{2}}$, A_0 and $m_{\frac{3}{2}}$ as a function of the parameters of the model. By running the RGEs, we show the low energy (PeV scale) spectrum of SUSY particles and show that the model can accommodate the ~ 125 GeV Higgs. The SUSY spectrum so obtained modifies the RGEs above the PeV scale and gives the coupling constant unification at the GUT scale, however such a gauge coupling unification can be achieved for a large range of SUSY-breaking scales. We find that there is a degeneracy in the values of inflation model parameters (γ, β, ζ) in predicting the correct CMB amplitude. However $\tan(\beta) = 1.8$, fixed from SUSY breaking, removes this degeneracy and provide a unique solution for inflation, and brings the explanation for dark matter, PeV neutrinos and inflation within the same model setup.

DATA AVAILABILITY

The data used to support the findings of this study are available from the corresponding author upon request.

CONFLICTS OF INTEREST

The author declares that there are no conflicts of interest regarding the publication of this paper.

ACKNOWLEDGEMENTS

G.K.C. would like to thank University Grants Commission, Govt. of India to provide financial support via Dr. D. S. Kothari Postdoctoral Fellowship (Grant No.: BSR/PH/2017-18/0026). N. K. acknowledges private communications with Florian Staub.

-
- [1] ATLAS collaboration, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett.* **B716** (2012) 1–29, [1207.7214].
 - [2] P. P. Giardino, K. Kannike, I. Masina, M. Raidal and A. Strumia, *The universal Higgs fit*, *JHEP* **05** (2014) 046, [1303.3570].
 - [3] CMS collaboration, S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett.* **B716** (2012) 30–61, [1207.7235].
 - [4] M. Drees, R. Godbole and P. Roy, *Theory and phenomenology of sparticles: An account of four-dimensional $N=1$ supersymmetry in high energy physics*. 2004.
 - [5] S. P. Martin, *A Supersymmetry primer*, [hep-ph/9709356](#).
 - [6] H. Baer and X. Tata, *Weak Scale Supersymmetry: From Superfields to Scattering Events*. 2006.
 - [7] J. R. Ellis, K. Enqvist, D. V. Nanopoulos and F. Zwirner, *Observables in Low-Energy Superstring Models*, *Mod. Phys. Lett.* **A1** (1986) 57.
 - [8] R. Barbieri and G. F. Giudice, *Upper Bounds on Supersymmetric Particle Masses*, *Nucl. Phys.* **B306** (1988) 63–76.
 - [9] ATLAS COLLABORATION collaboration, *Atlas collaboration*, <https://twiki.cern.ch/twiki/bin/view/atlaspublic>, .

- [10] CMS COLLABORATION collaboration, *Cms collaboration*,
<http://cms-results.web.cern.ch/cms-results/public-results/publications/>, .
- [11] M. R. Buckley, D. Feld, S. Macaluso, A. Monteux and D. Shih, *Cornering Natural SUSY at LHC Run II and Beyond*, 1610.08059.
- [12] J. L. Feng, *Naturalness and the Status of Supersymmetry*, *Ann. Rev. Nucl. Part. Sci.* **63** (2013) 351–382, [1302.6587].
- [13] G. F. Giudice and A. Romanino, *Split supersymmetry*, *Nucl. Phys.* **B699** (2004) 65–89, [hep-ph/0406088].
- [14] XENON collaboration, E. Aprile et al., *First Dark Matter Search Results from the XENON1T Experiment*, 1705.06655.
- [15] LUX collaboration, D. S. Akerib et al., *Results from a search for dark matter in the complete LUX exposure*, *Phys. Rev. Lett.* **118** (2017) 021303, [1608.07648].
- [16] GAMBIT collaboration, P. Athron et al., *Global fits of GUT-scale SUSY models with GAMBIT*, 1705.07935.
- [17] M. Abdughani, L. Wu and J. M. Yang, *The mixed bino-higgsino dark matter in natural SUSY confronted with XENON1T/PandaX and LHC data*, 1705.09164.
- [18] B. Zhu, R. Ding and T. Li, *Higgs Mass and Muon $g - 2$ Anomaly in MSSM with Gauge-Gravity hybrid Mediation*, 1610.09840.
- [19] M. Peiro and S. Robles, *Low-mass neutralino dark matter in supergravity scenarios: phenomenology and naturalness*, *JCAP* **1705** (2017) 010, [1612.00460].
- [20] F. Wang, W. Wang and J. M. Yang, *Reconcile muon $g-2$ anomaly with LHC data in SUGRA with generalized gravity mediation*, *JHEP* **06** (2015) 079, [1504.00505].
- [21] M. Chakraborti, U. Chattopadhyay and S. Poddar, *How light a higgsino or a wino dark matter can become in a compressed scenario of MSSM*, 1702.03954.
- [22] W. Abdallah and S. Khalil, *MSSM Dark Matter in Light of Higgs and LUX Results*, *Adv. High Energy Phys.* **2016** (2016) 5687463, [1509.07031].
- [23] M. Badziak, M. Olechowski and P. Szczerbiak, *Is well-tempered neutralino in MSSM still alive after 2016 LUX results?*, *Phys. Lett.* **B770** (2017) 226–235, [1701.05869].
- [24] J. Cao, Y. He, L. Shang, W. Su, P. Wu and Y. Zhang, *Strong constraints of LUX-2016 results on the natural NMSSM*, *JHEP* **10** (2016) 136, [1609.00204].
- [25] A. Basirnia, S. Macaluso and D. Shih, *Dark Matter and the Higgs in Natural SUSY*, *JHEP* **03** (2017) 073, [1605.08442].
- [26] A. Schneider, *Characterization of the Astrophysical Diffuse Neutrino Flux with IceCube High-Energy Starting Events*, *PoS ICRC2019*, 1004 (2020) doi:10.22323/1.358.1004 [arXiv:1907.11266 [astro-ph.HE]].

- [27] ICECUBE collaboration, M. G. Aartsen et al., *Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector*, *Science* **342** (2013) 1242856, [1311.5238].
- [28] ICECUBE collaboration, M. G. Aartsen et al., *First observation of PeV-energy neutrinos with IceCube*, *Phys. Rev. Lett.* **111** (2013) 021103, [1304.5356].
- [29] ICECUBE collaboration, M. G. Aartsen et al., *Observation of High-Energy Astrophysical Neutrinos in Three Years of IceCube Data*, *Phys. Rev. Lett.* **113** (2014) 101101, [1405.5303].
- [30] K. Murase, *On the Origin of High-Energy Cosmic Neutrinos*, *AIP Conf. Proc.* **1666** (2015) 040006, [1410.3680].
- [31] B. Feldstein, A. Kusenko, S. Matsumoto and T. T. Yanagida, *Phys. Rev. D* **88**, no.1, 015004 (2013) doi:10.1103/PhysRevD.88.015004 [arXiv:1303.7320 [hep-ph]].
- [32] Y. Sui and P. Bhupal Dev, *JCAP* **07**, 020 (2018) doi:10.1088/1475-7516/2018/07/020 [arXiv:1804.04919 [hep-ph]].
- [33] M. Kachelriess, O. Kalashev and M. Kuznetsov, *Phys. Rev. D* **98**, no.8, 083016 (2018) doi:10.1103/PhysRevD.98.083016 [arXiv:1805.04500 [astro-ph.HE]].
- [34] A. Bhattacharya, A. Esmaili, S. Palomares-Ruiz and I. Sarcevic, *JCAP* **05**, 051 (2019) doi:10.1088/1475-7516/2019/05/051 [arXiv:1903.12623 [hep-ph]].
- [35] M. Pandey, D. Majumdar and A. Halder, [arXiv:1909.06839 [hep-ph]].
- [36] L. A. Anchordoqui, V. Barger, H. Goldberg, X. Huang, D. Marfatia, L. H. M. da Silva et al., *IceCube neutrinos, decaying dark matter, and the Hubble constant*, *Phys. Rev.* **D92** (2015) 061301, [1506.08788].
- [37] ICECUBE collaboration, R. Abbasi et al., *Search for dark matter from the Galactic halo with the IceCube Neutrino Telescope*, *Phys. Rev.* **D84** (2011) 022004, [1101.3349].
- [38] M. Chianese and A. Merle, *A Consistent Theory of Decaying Dark Matter Connecting IceCube to the Sesame Street*, *JCAP* **1704** (2017) 017, [1607.05283].
- [39] S. B. Roland, B. Shakya and J. D. Wells, *PeV neutrinos and a 3.5 keV x-ray line from a PeV-scale supersymmetric neutrino sector*, *Phys. Rev.* **D92** (2015) 095018, [1506.08195].
- [40] D. Borah, A. Dasgupta, U. K. Dey, S. Patra and G. Tomar, *Multi-component Fermionic Dark Matter and IceCube PeV scale Neutrinos in Left-Right Model with Gauge Unification*, **1704.04138**.
- [41] C. El Aisati, M. Gustafsson and T. Hambye, *New Search for Monochromatic Neutrinos from Dark Matter Decay*, *Phys. Rev.* **D92** (2015) 123515, [1506.02657].
- [42] B. Shakya, S. B. Roland and J. D. Wells, *Hints of a PeV Scale Supersymmetric Neutrino Sector: Neutrino Masses, Dark Matter, keV and PeV Anomalies*, in *Proceedings, Meeting of the APS Division of Particles and Fields (DPF 2015): Ann Arbor, Michigan, USA, 4-8 Aug 2015*, 2015, 1510.08540, <https://inspirehep.net/record/1401341/files/arXiv:1510.08540.pdf>.
- [43] P. S. B. Dev, D. Kazanas, R. N. Mohapatra, V. L. Teplitz and Y. Zhang, *Heavy right-handed neutrino dark matter and PeV neutrinos at IceCube*, *JCAP* **1608** (2016) 034, [1606.04517].

- [44] Y. Bai, R. Lu and J. Salvado, *Geometric Compatibility of IceCube TeV-PeV Neutrino Excess and its Galactic Dark Matter Origin*, *JHEP* **01** (2016) 161, [1311.5864].
- [45] M. Ahlers, Y. Bai, V. Barger and R. Lu, *Galactic neutrinos in the TeV to PeV range*, *Phys. Rev. D* **93** (2016) 013009, [1505.03156].
- [46] N. Hiroshima, R. Kitano, K. Kohri and K. Murase, *High-energy Neutrinos from Multi-body Decaying Dark Matter*, 1705.04419.
- [47] A. Esmaili and P. D. Serpico, *Are IceCube neutrinos unveiling PeV-scale decaying dark matter?*, *JCAP* **1311** (2013) 054, [1308.1105].
- [48] A. Esmaili and P. D. Serpico, *Gamma-ray bounds from EAS detectors and heavy decaying dark matter constraints*, *JCAP* **1510** (2015) 014, [1505.06486].
- [49] A. Bhattacharya, A. Esmaili, S. Palomares-Ruiz and I. Sarcevic, *Probing decaying heavy dark matter with the 4-year IceCube HESE data*, *JCAP* **1707** (2017) 027, [1706.05746].
- [50] S. M. Boucenna, M. Chianese, G. Mangano, G. Miele, S. Morisi, O. Pisanti et al., *Decaying Leptophilic Dark Matter at IceCube*, *JCAP* **1512** (2015) 055, [1507.01000].
- [51] B. Dutta, Y. Gao, T. Li, C. Rott and L. E. Strigari, *Leptoquark implication from the CMS and IceCube experiments*, *Phys. Rev. D* **91** (2015) 125015, [1505.00028].
- [52] U. K. Dey and S. Mohanty, *Constraints on Leptoquark Models from IceCube Data*, *JHEP* **04** (2016) 187, [1505.01037].
- [53] N. Mileo, A. de la Puente and A. Szyrkman, *Implications of a Electroweak Triplet Scalar Leptoquark on the Ultra-High Energy Neutrino Events at IceCube*, *JHEP* **11** (2016) 124, [1608.02529].
- [54] V. Barger, E. Basso, Y. Gao and W.-Y. Keung, *Neutrino signals in IceCube from weak production of top and charm quarks*, *Phys. Rev. D* **95** (2017) 093002, [1611.00773].
- [55] B. Chauhan, B. Kindra and A. Narang, *A Leptoquark explanation for $(g - 2)_\mu$, R_K , R_{K^*} and, IceCube PeV events*, 1706.04598.
- [56] L. A. Anchordoqui, C. A. Garcia Canal, H. Goldberg, D. Gomez Dumm and F. Halzen, *Phys. Rev. D* **74**, 125021 (2006) doi:10.1103/PhysRevD.74.125021 [arXiv:hep-ph/0609214 [hep-ph]].
- [57] J. D. Wells, *PeV-scale supersymmetry*, *Phys. Rev. D* **71** (2005) 015013, [hep-ph/0411041].
- [58] S. A. R. Ellis and J. D. Wells, *High-scale Supersymmetry, the Higgs Mass and Gauge Unification*, 1706.00013.
- [59] Y. Akrami et al. [Planck], [arXiv:1807.06211 [astro-ph.CO]].
- [60] PLANCK collaboration, P. A. R. Ade et al., *Planck 2015 results. XIII. Cosmological parameters*, 1502.01589.
- [61] PLANCK collaboration, P. A. R. Ade et al., *Planck 2015 results. XX. Constraints on inflation*, 1502.02114.

- [62] P. Ade *et al.* [BICEP2 and Keck Array], *Phys. Rev. Lett.* **121**, 221301 (2018) doi:10.1103/PhysRevLett.121.221301 [arXiv:1810.05216 [astro-ph.CO]].
- [63] BICEP2, KECK ARRAY collaboration, P. A. R. Ade et al., *Improved Constraints on Cosmology and Foregrounds from BICEP2 and Keck Array Cosmic Microwave Background Data with Inclusion of 95 GHz Band*, *Phys. Rev. Lett.* **116** (2016) 031302, [1510.09217].
- [64] J. Ellis, D. V. Nanopoulos and K. A. Olive, *No-Scale Supergravity Realization of the Starobinsky Model of Inflation*, *Phys. Rev. Lett.* **111** (2013) 111301, [1305.1247].
- [65] I. Garg and S. Mohanty, *No scale SUGRA SO(10) derived Starobinsky Model of Inflation*, *Phys. Lett.* **B751** (2015) 7–11, [1504.07725].
- [66] J. Ellis, M. A. G. Garcia, N. Nagata, D. V. Nanopoulos and K. A. Olive, “Starobinsky-Like Inflation and Neutrino Masses in a No-Scale SO(10) Model,” *JCAP* **1611**, no. 11, 018 (2016) doi:10.1088/1475-7516/2016/11/018 [arXiv:1609.05849 [hep-ph]].
- [67] J. Ellis, H. J. He and Z. Z. Xianyu, “New Higgs Inflation in a No-Scale Supersymmetric SU(5) GUT,” *Phys. Rev. D* **91**, no. 2, 021302 (2015) doi:10.1103/PhysRevD.91.021302 [arXiv:1411.5537 [hep-ph]].
- [68] I. Garg and S. Mohanty, “No-scale SUGRA Inflation and Type-I seesaw,” arXiv:1711.01979 [hep-ph].
- [69] G. K. Chakravarty, G. Gupta, G. Lambiase and S. Mohanty, *Plateau Inflation in SUGRA-MSSM*, *Phys. Lett.* **B760** (2016) 263–266, [1604.02556].
- [70] G. K. Chakravarty, U. K. Dey, G. Lambiase and S. Mohanty, *Plateau Inflation in R-parity Violating MSSM*, *Phys. Lett.* **B763** (2016) 501–506, [1607.06904].
- [71] L. Alvarez-Gaume, C. Gomez and R. Jimenez, *Minimal Inflation*, *Phys. Lett.* **B690** (2010) 68–72, [1001.0010].
- [72] L. Alvarez-Gaume, C. Gomez and R. Jimenez, *A Minimal Inflation Scenario*, *JCAP* **1103** (2011) 027, [1101.4948].
- [73] G. K. Chakravarty and S. Mohanty, *Power law Starobinsky model of inflation from no-scale SUGRA*, *Phys. Lett.* **B746** (2015) 242–247, [1405.1321].
- [74] S. Ferrara, R. Kallosh and A. Linde, *Cosmology with Nilpotent Superfields*, *JHEP* **10** (2014) 143, [1408.4096].
- [75] G. K. Chakravarty, S. Das, G. Lambiase and S. Mohanty, *Dilaton assisted two-field inflation from no-scale supergravity*, *Phys. Rev.* **D94** (2016) 023521, [1511.03121].
- [76] S. Ferrara, R. Kallosh, A. Linde and M. Porrati, *Minimal Supergravity Models of Inflation*, *Phys. Rev.* **D88** (2013) 085038, [1307.7696].
- [77] F. Farakos, A. Kehagias and A. Riotto, *On the Starobinsky Model of Inflation from Supergravity*, *Nucl. Phys.* **B876** (2013) 187–200, [1307.1137].

- [78] S. Ferrara, P. Fre and A. S. Sorin, *On the Gauged $K\tilde{A}d\tilde{h}$ ler Isometry in Minimal Supergravity Models of Inflation*, *Fortsch. Phys.* **62** (2014) 277–349, [1401.1201].
- [79] K. Nakayama, K. Saikawa, T. Terada and M. Yamaguchi, “Structure of $K\tilde{A}d\tilde{h}$ ler potential for D-term inflationary attractor models,” *JHEP* **1605**, 067 (2016) doi:10.1007/JHEP05(2016)067 [arXiv:1603.02557 [hep-th]].
- [80] R. Deen, B. A. Ovrut and A. Purves, *Supersymmetric Sneutrino-Higgs Inflation*, 1606.00431.
- [81] K. Nakayama, F. Takahashi and T. T. Yanagida, *Chaotic Inflation with Right-handed Sneutrinos after Planck*, *Phys. Lett. B* **730** (2014) 24–29, [1311.4253].
- [82] C. S. Aulakh and I. Garg, *Supersymmetric Seesaw Inflation*, *Phys. Rev. D* **86** (2012) 065001, [1201.0519].
- [83] Y. G. Kim, H. M. Lee and W.-I. Park, *The μ problem and sneutrino inflation*, *JHEP* **08** (2011) 126, [1107.1113].
- [84] C. Pallis and N. Toumbas, *Non-Minimal Sneutrino Inflation, Peccei-Quinn Phase Transition and non-Thermal Leptogenesis*, *JCAP* **1102** (2011) 019, [1101.0325].
- [85] S. Antusch, M. Bastero-Gil, J. P. Baumann, K. Dutta, S. F. King and P. M. Kostka, *Gauge Non-Singlet Inflation in SUSY GUTs*, *JHEP* **08** (2010) 100, [1003.3233].
- [86] R. Allahverdi, K. Enqvist, J. Garcia-Bellido and A. Mazumdar, *Gauge invariant MSSM inflaton*, *Phys. Rev. Lett.* **97** (2006) 191304, [hep-ph/0605035].
- [87] P. Athron, J.-h. Park, D. Stücker and A. Voigt, *Flexible SUSY $\tilde{A}\tilde{A}\tilde{A}\tilde{A}$ spectrum generator generator for supersymmetric models*, *Comput. Phys. Commun.* **190** (2015) 139–172, [1406.2319].
- [88] P. Athron, J.-h. Park, T. Steudtner, D. Stücker and A. Voigt, *Precise Higgs mass calculations in (non-)minimal supersymmetry at both high and low scales*, *JHEP* **01** (2017) 079, [1609.00371].
- [89] F. Staub, *Exploring new models in all detail with SARAH*, *Adv. High Energy Phys.* **2015** (2015) 840780, [1503.04200].
- [90] F. Staub and W. Porod, *Improved predictions for intermediate and heavy Supersymmetry in the MSSM and beyond*, *Eur. Phys. J. C* **77** (2017) 338, [1703.03267].
- [91] W. Porod, *SPheno, a program for calculating supersymmetric spectra, SUSY particle decays and SUSY particle production at e^+e^- colliders*, *Comput. Phys. Commun.* **153** (2003) 275–315, [hep-ph/0301101].
- [92] W. Porod and F. Staub, *SPheno 3.1: Extensions including flavour, CP-phases and models beyond the MSSM*, *Comput. Phys. Commun.* **183** (2012) 2458–2469, [1104.1573].
- [93] G. Belanger, F. Boudjema, P. Brun, A. Pukhov, S. Rosier-Lees, P. Salati et al., *Indirect search for dark matter with micrOMEGAs2.4*, *Comput. Phys. Commun.* **182** (2011) 842–856, [1004.1092].

- [94] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, *micrOMEGAs3: A program for calculating dark matter observables*, *Comput. Phys. Commun.* **185** (2014) 960–985, [1305.0237].
- [95] M. Carena, D. Choudhury, S. Lola and C. Quigg, *Phys. Rev. D* **58**, 095003 (1998) doi:10.1103/PhysRevD.58.095003 [arXiv:hep-ph/9804380 [hep-ph]].
- [96] P. S. B. Dev, D. K. Ghosh and W. Rodejohann, *Phys. Lett. B* **762**, 116–123 (2016) doi:10.1016/j.physletb.2016.08.066 [arXiv:1605.09743 [hep-ph]].
- [97] D. BeĀĎireviĀĜ, B. Panes, O. Sumensari and R. Zukanovich Funchal, *JHEP* **06**, 032 (2018) doi:10.1007/JHEP06(2018)032 [arXiv:1803.10112 [hep-ph]].
- [98] S. Ando and K. Ishiwata, *Constraints on decaying dark matter from the extragalactic gamma-ray background*, *JCAP* **1505** (2015) 024, [1502.02007].
- [99] S. Shirai, F. Takahashi and T. T. Yanagida, *R-violating Decay of Wino Dark Matter and electron/positron Excesses in the PAMELA/Fermi Experiments*, *Phys. Lett.* **B680** (2009) 485–488, [0905.0388].
- [100] K. Hamaguchi, T. Moroi and K. Nakayama, *AMS-02 Antiprotons from Annihilating or Decaying Dark Matter*, *Phys. Lett.* **B747** (2015) 523–528, [1504.05937].
- [101] H. Baer, A. Mustafayev, E.-K. Park and X. Tata, *Collider signals and neutralino dark matter detection in relic-density-consistent models without universality*, *JHEP* **05** (2008) 058, [0802.3384].
- [102] S. Akula, M. Liu, P. Nath and G. Peim, *Naturalness, Supersymmetry and Implications for LHC and Dark Matter*, *Phys. Lett.* **B709** (2012) 192–199, [1111.4589].
- [103] N. Bhattacharyya, A. Choudhury and A. Datta, *Low mass neutralino dark matter in mSUGRA and more general models in the light of LHC data*, *Phys. Rev.* **D84** (2011) 095006, [1107.1997].
- [104] H. Baer, V. Barger and A. Mustafayev, *Neutralino dark matter in mSUGRA/CMSSM with a 125 GeV light Higgs scalar*, *JHEP* **05** (2012) 091, [1202.4038].
- [105] P. S. Bhupal Dev, S. Mondal, B. Mukhopadhyaya and S. Roy, *Phenomenology of Light Sneutrino Dark Matter in cMSSM/mSUGRA with Inverse Seesaw*, *JHEP* **09** (2012) 110, [1207.6542].
- [106] B. Kaufman, P. Nath, B. D. Nelson and A. B. Spisak, *Light Stops and Observation of Supersymmetry at LHC RUN-II*, *Phys. Rev.* **D92** (2015) 095021, [1509.02530].
- [107] D. Hu, T. Li, A. Lux, J. A. Maxin and D. V. Nanopoulos, *General No-Scale Supergravity: An \mathcal{F} -SU(5) Tale*, *Phys. Lett.* **B771** (2017) 264–270, [1703.02917].
- [108] H. Baer, C. Balazs, A. Belyaev, T. Krupovnickas and X. Tata, *Updated reach of the CERN LHC and constraints from relic density, $b \rightarrow s\gamma$ and $a(\mu)$ in the mSUGRA model*, *JHEP* **06** (2003) 054, [hep-ph/0304303].
- [109] H. P. Nilles, *Supersymmetry, Supergravity and Particle Physics*, *Phys. Rept.* **110** (1984) 1–162.
- [110] I. Niessen, *Supersymmetric Phenomenology in the mSUGRA Parameter Space*, Ph.D. thesis, Nijmegen U., IMAPP, 2008. 0809.1748.

- [111] U. Chattopadhyay, A. Corsetti and P. Nath, *WMAP constraints, SUSY dark matter and implications for the direct detection of SUSY*, *Phys. Rev.* **D68** (2003) 035005, [[hep-ph/0303201](#)].
- [112] K. L. Chan, U. Chattopadhyay and P. Nath, *Naturalness, weak scale supersymmetry and the prospect for the observation of supersymmetry at the Tevatron and at the CERN LHC*, *Phys. Rev.* **D58** (1998) 096004, [[hep-ph/9710473](#)].
- [113] H. Baer, V. Barger and M. Savoy, *Supergravity gauge theories strike back: There is no crisis for SUSY but a new collider may be required for discovery*, *Phys. Scripta* **90** (2015) 068003, [[1502.04127](#)].
- [114] G. Coughlan, W. Fischler, E. W. Kolb, S. Raby and G. G. Ross, *Phys. Lett. B* **131**, 59-64 (1983) doi:10.1016/0370-2693(83)91091-2
- [115] J. R. Ellis, D. V. Nanopoulos and M. Quiros, *Phys. Lett. B* **174**, 176-182 (1986) doi:10.1016/0370-2693(86)90736-7
- [116] T. Banks, D. B. Kaplan and A. E. Nelson, *Phys. Rev. D* **49**, 779-787 (1994) doi:10.1103/PhysRevD.49.779 [[arXiv:hep-ph/9308292](#) [[hep-ph](#)]].
- [117] B. de Carlos, J. Casas, F. Quevedo and E. Roulet, *Phys. Lett. B* **318**, 447-456 (1993) doi:10.1016/0370-2693(93)91538-X [[arXiv:hep-ph/9308325](#) [[hep-ph](#)]].
- [118] A. Linde, Y. Mambrini and K. A. Olive, *Phys. Rev. D* **85**, 066005 (2012) doi:10.1103/PhysRevD.85.066005 [[arXiv:1111.1465](#) [[hep-th](#)]].
- [119] E. Dudas, A. Linde, Y. Mambrini, A. Mustafayev and K. A. Olive, *Eur. Phys. J. C* **73**, no.1, 2268 (2013) doi:10.1140/epjc/s10052-012-2268-7 [[arXiv:1209.0499](#) [[hep-ph](#)]].
- [120] S. K. Soni and H. A. Weldon, *Analysis of the Supersymmetry Breaking Induced by $N=1$ Supergravity Theories*, *Phys. Lett.* **B126** (1983) 215–219.
- [121] V. S. Kaplunovsky and J. Louis, *Model independent analysis of soft terms in effective supergravity and in string theory*, *Phys. Lett.* **B306** (1993) 269–275, [[hep-th/9303040](#)].
- [122] E. Dudas and S. K. Vempati, *Large D -terms, hierarchical soft spectra and moduli stabilisation*, *Nucl. Phys.* **B727** (2005) 139–162, [[hep-th/0506172](#)].
- [123] A. Brignole, L. E. Ibanez and C. Munoz, *Towards a theory of soft terms for the supersymmetric Standard Model*, *Nucl. Phys.* **B422** (1994) 125–171, [[hep-ph/9308271](#)].
- [124] A. Brignole, L. E. Ibanez and C. Munoz, *Soft supersymmetry breaking terms from supergravity and superstring models*, *Adv. Ser. Direct. High Energy Phys.* **18** (1998) 125–148, [[hep-ph/9707209](#)].
- [125] K. Griest and M. Kamionkowski, *Unitarity Limits on the Mass and Radius of Dark Matter Particles*, *Phys. Rev. Lett.* **64** (1990) 615.
- [126] C. Rott, K. Kohri and S. C. Park, *Superheavy dark matter and IceCube neutrino signals: Bounds on decaying dark matter*, *Phys. Rev.* **D92** (2015) 023529, [[1408.4575](#)].
- [127] M. Re Fiorentin, V. Niro and N. Fornengo, *A consistent model for leptogenesis, dark matter and the IceCube signal*, *JHEP* **11** (2016) 022, [[1606.04445](#)].

- [128] B. Audren, J. Lesgourgues, G. Mangano, P. D. Serpico and T. Tram, *Strongest model-independent bound on the lifetime of Dark Matter*, *JCAP* **1412** (2014) 028, [1407.2418].
- [129] K. Garrett and G. Duda, *Dark Matter: A Primer*, *Adv. Astron.* **2011** (2011) 968283, [1006.2483].
- [130] H. Murayama, *Physics Beyond the Standard Model and Dark Matter*, in *Les Houches Summer School - Session 86: Particle Physics and Cosmology: The Fabric of Spacetime Les Houches, France, July 31-August 25, 2006*, 2007, 0704.2276,
<https://inspirehep.net/record/748786/files/arXiv:0704.2276.pdf>.
- [131] E. W. Kolb and M. S. Turner, *The Early Universe*. 1989.
- [132] PLANCK collaboration, P. A. R. Ade et al., *Planck 2013 results. XVI. Cosmological parameters*, *Astron. Astrophys.* **571** (2014) A16, [1303.5076].
- [133] Z. Berezhiani, A. D. Dolgov, and I. I. Tkachev, “Reconciling Planck results with low redshift astronomical measurements,” *Phys. Rev.* **D92** no. 6, (2015) 061303, [arXiv:1505.03644](#) [astro-ph.CO].
- [134] E. A. Baltz and P. Gondolo, *Neutralino decay rates with explicit R-parity violation*, *Phys. Rev.* **D57** (1998) 2969–2973, [hep-ph/9709445].
- [135] S. Palomares-Ruiz, A. C. Vincent and O. Mena, *Spectral analysis of the high-energy IceCube neutrinos*, *Phys. Rev.* **D91** (2015) 103008, [1502.02649].
- [136] ICECUBE collaboration, M. G. Aartsen et al., *Atmospheric and astrophysical neutrinos above 1 TeV interacting in IceCube*, *Phys. Rev.* **D91** (2015) 022001, [1410.1749].
- [137] A. Esmaili, A. Ibarra and O. L. G. Peres, *Probing the stability of superheavy dark matter particles with high-energy neutrinos*, *JCAP* **1211** (2012) 034, [1205.5281].