

Indication for Double Parton Scatterings in W +Prompt J/ψ Production at the LHC

Jean-Philippe Lansberg^a, Hua-Sheng Shao^{b,c}, Nodoka Yamanaka^{a,d}

^a*IPNO, CNRS-IN2P3, Univ. Paris-Sud, Université Paris-Saclay, 91406 Orsay Cedex, France*

^b*Sorbonne Universités, UPMC Univ. Paris 06, UMR 7589, LPTHE, F-75005 Paris, France*

^c*CNRS, UMR 7589, LPTHE, F-75005 Paris, France*

^d*THES Research Group, RIKEN, Wako, Saitama 351-0198, Japan*

Abstract

We re-analyse the associated production of a prompt J/ψ and a W boson in pp collisions at the LHC following the results of the ATLAS Collaboration. We perform the first study of the Single-Parton-Scattering (SPS) contributions at the Next-to-Leading Order (NLO) in α_s in the Colour-Evaporation Model (CEM), an approach based on the quark-hadron-duality. Our study provides clear indications for Double-Parton-Scattering (DPS) contributions, in particular at low transverse momenta, since our SPS CEM evaluation, which can be viewed as a conservative upper limit of the SPS yields, falls short compared to the ATLAS experimental data by 3.1 standard deviations. We also determine a finite allowed region for σ_{eff} , inversely proportional to the size of the DPS yields, corresponding to the otherwise opposed hypotheses, namely our NLO CEM evaluation and the LO direct Colour-Singlet (CS) Model contribution. In both cases, the resulting DPS yields are significantly larger than that initially assumed by ATLAS based on jet-related analyses but is consistent with their observed raw-yield azimuthal distribution and with their prompt $J/\psi + J/\psi$ and $Z + \text{prompt } J/\psi$ data.

1. Introduction.

The simultaneous production of vector bosons and quarkonia at high-energy colliders is a very useful observable to study perturbative and nonperturbative aspects of Quantum Chromodynamics (QCD). It also provides original means to search for new physics beyond the standard model via the Higgs sector, as illustrated by the pioneering study of CDF Collaboration [1, 2]. Recently, the final states $J/\psi + W$ [3] and $J/\psi + Z$ [4] were observed by the ATLAS Collaboration. Similar processes with bottomonia have however not been observed yet¹.

Besides the associated production with a vector boson, quarkonium-pair production has been the object of a number of recent experimental studies at the LHC and the Tevatron [14–18] –30 years after the pioneering analyses of NA3 [19, 20]. At small rapidity separations ($\Delta y_{\psi\psi}$), none of them exhibit any tension with the SPS CS contributions (*i.e.* the LO in v^2 of Non Relativistic QCD (NRQCD) [21], known up to NLO accuracy [22–24]) whereas they point at a significant DPS contributions for increasing $\Delta y_{\psi\psi}$ – in accordance with previously studied observables [25–31]. Di- J/ψ hadroproduction has in fact been the object of a large number of theoretical works [22–24, 32–45]. The case of $\Upsilon + J/\psi$ production, measured by the D0 collaboration [46], is slightly different as it seems highly dominated by DPS contributions. We refer to [47] for a complete and up-to-date theory discussion of this reaction. Similar conclusions, pointing at a dominant DPS yield, were made by LHCb for

J/ψ +charm [48] and Υ +charm [49] production with measured cross sections significantly larger than the SPS expectations [50–52]. We however note that it is not clear whether the magnitude of all these DPS yields fit in a coherent picture with a *universal* σ_{eff} which would be inversely proportional to the probability of a second parton scattering.

In the case of $J/\psi + Z$ production, the yield observed by ATLAS happens to be up to nearly one order of magnitude larger than the theory evaluations from NRQCD (with CS and/or Colour-Octet (CO) –higher order in v^2 – contributions) [53, 54]. A natural explanation for such a gap would be –like for quarkonium pairs at large $\Delta y_{\psi\psi}$ – associated production through DPS but for the fact that the azimuthal distribution of the observed events shows a significant back-to-back peak pointing at a large SPS. In a recent paper [55], we have solved this apparent conflict by invoking a misinterpretation of this raw-yield azimuthal distribution when it results from two different sources with very different transverse-momentum distributions in a detector like ATLAS with a strongly transverse-momentum-dependent acceptance. In a further study [56], we have shown that the production of a Z with a non-prompt J/ψ (*i.e.* from a b -hadron decay), as reported by ATLAS in [4], is well accounted by the SPS predictions with a limited impact of DPSs.

In the case of $J/\psi + W$ production, the observed yield of ATLAS [3] also ends up to be nearly one order of magnitude larger than the recent theory SPS evaluations from NRQCD (with CS and/or CO contributions) [57, 58] (see [59, 60] for earlier studies). Just like the $J/\psi + Z$ case, the ATLAS raw-yield azimuthal distribution seems to exhibit a non-trivial structure hinting at the presence of SPS events.

In this context, we have decided to have another look at

¹Other reactions of specific interest include the production of a quarkonium + a photon. It was proposed to constrain the quarkonium-production mechanisms [5–9], to study the proton gluon content [10, 11] or to probe the H^0 coupling to the heavy-quarks [12, 13].

prompt- $J/\psi + W$ production at the LHC, in particular relying on an analysis of the SPS yield at NLO under the assumption of quark-hadron duality [the Colour-Evaporation Model (CEM)] [61, 62] as in our earlier study [55]. As we demonstrated then, the CEM provides a conservative upper limit to the SPS yield which we use to draw definite conclusions about the importance of the DPSs.

This paper is organised as follows. In Section 2, we briefly explain our NLO CEM SPS computation which, as an upper theory limit for the SPS, allows us to claim for a clear indication of the contributions of DPSs at a 3 σ level. In Section 3, we extract the DPS yield and determine σ_{eff} with its uncertainty range. We then show that the azimuthal distribution of the prompt $J/\psi + W$ events collected by ATLAS remains compatible with a yield highly dominated by DPSs. Section 4 is devoted to our conclusions and outlook.

2. Our NLO evaluation of the SPS yield

In this work, we thus focus on the production of a W with a prompt J/ψ (which does not result from b -hadron decays) at the LHC in the kinematical region accessible by the ATLAS detector (see Table 1).

W boson selection ²	
$P_T^{\mu^\pm} > 25 \text{ GeV}, E_T > 20 \text{ GeV}, M_T(W) > 40 \text{ GeV}, \eta_{\mu^\pm} < 2.4$	
J/ ψ selection	
ATLAS fiducial [3]	ATLAS inclusive [3]
$8.5 < P_T^{J/\psi} < 30 \text{ GeV}$	$8.5 < P_T^{J/\psi} < 30 \text{ GeV}$
$ y_{J/\psi} < 2.1$	$ y_{J/\psi} < 2.1$
$P_T^{\text{leading } \mu^\pm} > 4.0 \text{ GeV}$	
$ \eta^{\text{leading } \mu^\pm} < 2.5$	
either $\left(P_T^{\text{sub-leading } \mu^\pm} > 2.5 \text{ GeV} \right)$	
$1.3 \leq \eta^{\text{sub-leading } \mu^\pm} < 2.5$	
or $\left(P_T^{\text{sub-leading } \mu^\pm} > 3.5 \text{ GeV} \right)$	
$ \eta^{\text{sub-leading } \mu^\pm} < 1.3$	

Table 1: Phase-space definition of the measured fiducial/inclusive production cross-section following the geometrical acceptance of the ATLAS detector.

As said above, the SPS contributions have been studied in the past within NRQCD, considering either CS [58] or CO [57] channels. The latter CO evaluation at NLO accuracy in α_s is however perfectible as it relies on artificially low scales ignoring the emission of a W boson and with NRQCD Long Distance Matrix Elements (LDMEs) which are significantly larger than the NLO LDMEs compatible with the LHC yield and polarisation data (see *e.g.* [63]). That being said, since the direct CS yield does not depend on free tunable parameters, it provides a strict *lower* limit on the SPS $J/\psi + W$ production cross section.

² $M_T^W \equiv 2P_T^{\mu^\pm} E_T (1 - \cos(\phi^\mu - \phi^{\nu\mu}))$ not to be confused with $m_T^2 = m^2 + P_T^2$.

On the contrary, the CEM most certainly sets an *upper* limit on the SPS yield. Indeed, the CEM tends to overshoot the experimental data for *single* quarkonium production at large $P_T^{J/\psi}$ [64–66]³. This follows from the precocious dominance of the gluon-fragmentation topologies whose strength is therefore too high at large $P_T^{J/\psi}$. For $J/\psi + W$, the dominant CEM channel is quark-antiquark annihilation with the emission of a W and an off-shell gluon fragmenting into a quarkonium. As such, the CEM likely results in an overestimation of the prompt $J/\psi + W$ cross section. It can thus be considered as a conservative upper limit of the theoretical prediction of the SPS yield. In other words, any set of NRQCD LDMEs which would be compatible with the yield predicted here would severely overshoot the very precise single- J/ψ data. Let us also add that, due to the simplicity of the model, the CEM avoids the complexities encountered in NLO analyses of NRQCD, where a large theoretical uncertainty in any case remains. Overall, we will consider that the difference between both (direct CSM & CEM) represents our best evaluation of the theoretical uncertainty on the SPS cross section.

In the CEM, one considers the integral of the cross section for $Q\bar{Q}$ pair production with an invariant-mass between the quark mass threshold $2m_Q$ and that of open-heavy-flavour hadrons $2m_H$, where the hadronisation of heavy quarks into a quarkonium is likely. To obtain the cross section to the hadron state (Q), this integral is multiplied by a phenomenological factor for the probability of the quark states to hadronise into a given quarkonium state. In short, we would consider

$$\sigma_Q^{(\text{N})\text{LO}, \text{direct prompt}} = \mathcal{P}_Q^{(\text{N})\text{LO}, \text{direct prompt}} \int_{2m_Q}^{2m_H} \frac{d\sigma_{Q\bar{Q}}^{(\text{N})\text{LO}}}{dm_{Q\bar{Q}}} dm_{Q\bar{Q}}. \quad (1)$$

\mathcal{P}_Q can be paralleled to the LDMEs in NRQCD. With the approach just described, the direct or prompt yields result from the same computation but with a different overall factor. In practice, we use $\mathcal{P}_{J/\psi}^{\text{LO}, \text{prompt}} = 0.014 \pm 0.001$ and $\mathcal{P}^{\text{NLO}, \text{prompt}} = 0.009 \pm 0.0004$ which we have fit in [55] on the differential cross section for the production of single J/ψ as measured by ATLAS [71] using $m_c = 1.27 \text{ GeV}$ (see [72] for the mass choice). We refer to [73] for the first NLO CEM fits using the P_T -integrated cross sections.

In view of Eq. (1), computing $J/\psi + W$ production at NLO can be performed with modern tools of automated NLO frameworks, with some slight tunings. In practice, we have used MADGRAPH5_AMC@NLO [74]⁴.

The hard scattering process to be considered is⁵ $ij \rightarrow c\bar{c} + W^\pm + k$ with i, j and k standing for g, q or \bar{q} . As what regards the parton distribution function (PDF), we have used

³To cure this issue, different mechanisms [67–70] were proposed but are not the object of a consensus. In our recent study [55], we have shown that the NLO corrections to the $P_T^{J/\psi}$ spectrum reduce the scale sensitivity but confirm the issue.

⁴We stress that, for the CEM, there is no need to use a specific automated tool like MADONIA [75] and HELAC-ONIA [76, 77] which are by the way currently not able to treat loop corrections.

⁵Unlike the CSM case, channels such as $sg \rightarrow W^- J/\psi c$ negligibly contribute to the CEM yield and can safely be neglected.

Experiment [3]	LO CEM SPS	NLO CEM SPS	LO direct CSM SPS [58]	DPS [3]
$12.6 \pm 3.2 \pm 0.9^{+4.1}_{-2.5}$	$0.44^{+0.15}_{-0.10}$	$0.78^{+0.21}_{-0.19}$	0.3 ± 0.1	$72 \text{ mb}/\sigma_{\text{eff}}$

Table 2: Comparison of the experimental inclusive data for the normalised ratio $\frac{d\sigma(pp \rightarrow W^\pm + J/\psi)}{dy_{J/\psi}} \frac{\text{BR}(J/\psi \rightarrow \mu^+ \mu^-)}{\sigma(pp \rightarrow W^\pm)}$ with our results of calculations in the CEM. We also show the normalised LO CS direct yield [58] and the DPS yield ratios extracted from [3] as a function of σ_{eff} . The unit is in 10^{-7} . The theoretical uncertainty for the (N)LO SPS is from the renormalisation and factorisation scales.

the NLO NNPDF 2.3 PDF set [78] with $\alpha_s(M_Z) = 0.118$ provided by LHAPDF [79].

As already stated above, we have taken $m_c = 1.27 \text{ GeV}$, while choosing $m_c = 1.5 \text{ GeV}$ would generate negligible changes to our results provided that the non-perturbative CEM parameter is chosen coherently. We note that the heavy-quark-mass dependence is de facto absorbed in the CEM parameter, hence the main theoretical uncertainties result from the renormalisation μ_R and factorisation μ_F scale variations which are believed to account for the unknown higher-order α_s corrections. In practice, we have varied them independently within $\frac{1}{2}\mu_0 \leq \mu_R, \mu_F \leq 2\mu_0$ where the central scale μ_0 is the mass of W boson, M_W .

For the current analysis, ATLAS preferred the ratio of the cross section for $W^\pm + J/\psi$ to that for W^\pm in order to discard some efficiency and acceptance corrections related to the W^\pm detection. For our theory evaluation, we use the NNLO W boson production cross section in the ATLAS acceptance corrected by the branching ratio for the decay to muons $\sigma(pp \rightarrow W^\pm) \text{BR}(W \rightarrow \mu\nu) = 5.08 \text{ nb}$ [80–82]. In such a case, the $W^\pm + J/\psi$ cross section in the numerator should also be multiplied by the branching $\text{BR}(W \rightarrow \mu\nu)$ which cancels with that in the denominator. Overall one only needs to consider:

$$\frac{d\sigma(pp \rightarrow W^\pm + J/\psi)}{dy_{J/\psi}} \frac{\text{BR}(J/\psi \rightarrow \mu^+ \mu^-)}{\sigma(pp \rightarrow W^\pm)}, \quad (2)$$

where $y_{J/\psi}$ is the J/ψ rapidity, $\text{BR}(J/\psi \rightarrow \mu^+ \mu^-) = 0.05961 \pm 0.00033$ [83]. We have also assumed $\text{BR}(W \rightarrow \mu\nu) = 0.1063$ and the J/ψ -rapidity range of 4.2 units to evaluate $d\sigma/dy_{J/\psi}$.

Our results (LO, NLO CEM) for the (inclusive) total cross section ratio are gathered in Table 2 along with the experimental results by ATLAS and the LO direct CS yield ratio computed in [58]. One notes that the NLO CEM ratio⁶ is –as expected– nearly three times larger than the direct CSM ratio. In spite of the large experimental uncertainties, the SPS predictions strongly underestimate the measurements, by more than three standard deviations⁷. The same features can be observed in the distribution in $P_T^{J/\psi}$, shown in Fig. 1 with the black (green) hatched histograms, resulting in a 3.1 standard-deviation discrepancy. As such we claim that these are clear indications of the existence of

a DPS yield in this process which is supported by the analysis of the transverse momentum and azimuthal dependences as we will show now.




Figure 1: Comparison of $\frac{d\sigma(pp \rightarrow W^\pm + J/\psi)}{dy_{J/\psi} dP_T^{J/\psi}} \frac{\text{BR}(J/\psi \rightarrow \mu^+ \mu^-)}{\sigma(pp \rightarrow W^\pm)}$ as a function of the transverse momentum of the prompt J/ψ . The division by $dy_{J/\psi}$ here means that we have divided the cross section by the rapidity interval of $|y_{J/\psi}| < 2.1$.

3. Our extraction of the DPS yield

3.1. DPSs and the $P_T^{J/\psi}$ -integrated cross section

The DPS yield results from two uncorrelated scatterings within one proton-proton collision. As such, it is usually parametrised by the rudimentary pocket formula which, for the process under study, reads

$$d\sigma^{\text{DPS}}(J/\psi + W) = \frac{d\sigma(J/\psi) d\sigma(W)}{\sigma_{\text{eff}}}, \quad (3)$$

where σ_{eff} is a supposedly universal parameter with the dimension of a surface. Recent experimental analyses are pointing at a visible impact of the DPSs in many reactions [15, 17, 25–31, 40].

As a reference value, ATLAS used in their data-theory comparison [3] $\sigma_{\text{eff}} = 15 \pm 3(\text{stat.})^{+5}_{-3}(\text{sys.}) \text{ mb}$ (from $W + 2$ -jet data [30]) which results, with $\sigma(J/\psi)$ and $\sigma(W)$ obtained from their data, in a normalised DPS cross section about 3 times smaller than their measurements (compare the first and last column of Table 2).

⁶On the way we note that the K factor for our CEM evaluation is 1.77.

⁷In evaluating the discrepancy, we have not considered the spin uncertainties. Whereas, on general grounds, the ATLAS evaluation is perfectly legitimate, we believe that, under the DPS dominance hypothesis which we suggest, it probably lies outside the range allowed by the single J/ψ polarisation measurement of CMS [84], for instance.

In the present analysis, we will instead extract the DPS yield by assuming that the difference between the ATLAS value and our SPS evaluation is entirely due to DPSs. As such, the σ_{eff} associated to the present observable can also be constrained in a relatively precise manner⁸ and confronted to other extractions. Let us recall that the SPS CEM evaluation should be regarded as an upper limit of the SPS yield.




Figure 2: Comparison of inclusive/fiducial $P_T^{J/\psi}$ -integrated cross section ratios from the CEM NLO SPS, our fit DPS and their sum. The DPS uncertainties reflect that of Eq. (4).

Let us now describe how we evaluate σ_{eff} and its uncertainty. Since the DPS yield is simply obtained by subtracting the SPS yield from the (inclusive) total cross section of ATLAS, the uncertainty on the DPS yield, and consequently⁹ that of σ_{eff} , will then depend on the (statistical and systematic) uncertainties of the data [3] and on the range spanned by the SPS evaluations – the NLO CEM and the direct LO CSM [58]¹⁰. Since the SPS values are in both cases much smaller than the data, the theoretical uncertainties are in fact nearly irrelevant for the determination of σ_{eff} . Our results are reported on Table 3 and on Fig. 2.

In particular, our combined result for σ_{eff} is then

$$\sigma_{\text{eff}} = (6.1^{+3.3}_{-1.9 \text{ exp}} {}^{+0.1}_{-0.3 \text{ theo}}) \text{ mb} \quad (4)$$

which is nearly three times smaller than the ATLAS assumption (15 mb) with somewhat smaller uncertainties than found by other works. It is also consistent with our latest extraction from prompt $J/\psi + Z$ production [55] in Fig. 3, where we also compare our extraction with other measurements [15, 17, 25–31, 40, 48, 49]. Such a low value may hint at the non-universality of σ_{eff} in different processes, with a dependence on the flavour of the initial state, on the kinematics of the final state or on the energy of the proton-proton collision. For example, the 3 values of σ_{eff} from

⁸Assuming the same single particle cross sections, $\sigma(J/\psi)$ and $\sigma(W)$, as ATLAS.

⁹The uncertainties on the measured single-particle cross sections, $\sigma(J/\psi)$ and $\sigma(W)$, are negligible compared to that on $\sigma(J/\psi + W)$.

¹⁰We note that the NLO NRQCD evaluation [57] –despite the aforementioned drawbacks– lies within this range.

the ATLAS J/ψ -associated-production measurements are in general smaller than those from the LHCb measurements at forward rapidities. Such an observation seems to follow –at least qualitatively– the lines of the mean-field approximation [85].

3.2. Consistency with the transverse momentum and azimuthal distributions

Our analysis shows that the DPSs are by far dominant and the SPSs could even be omitted in the analysis of the $P_T^{J/\psi}$ -integrated cross section. Since our procedure simply amounted to assume that any gap between the predicted SPS yield and the data was from DPS, it is important to check its consistency with differential cross sections.

Looking back at Fig. 1, we see that the introduction of a somewhat larger DPS yield perfectly fills the gap where needed and does not create any surplus at large $P_T^{J/\psi}$ where the SPS was closest to the data. We note that the plotted DPS $P_T^{J/\psi}$ spectrum follows from that of ATLAS [3] except for a different normalisation owing to the smaller value of σ_{eff} which we have just discussed. Like for the $P_T^{J/\psi}$ -integrated DPS cross section, ATLAS used the single-particle cross sections, here $d\sigma(J/\psi)/dP_T^{J/\psi}$, which they obtained from their own data.

Indeed, the sum of the SPS and DPS yields, in red, gives a reasonable account of the ATLAS differential yield. This agreement should however not be over-interpreted in view of the large experimental uncertainties. This observation is rather a consistency check than a test. The good agreement at low $P_T^{J/\psi}$ simply follows from the fit of σ_{eff} to the total yield but resolves the 3- σ discrepancy between “theory” and “experiment”.

This also helps illustrate that the low- $P_T^{J/\psi}$ yield is only from DPS contributions (as is the total yield) and that, at high $P_T^{J/\psi}$, DPS and SPS contributions could be of the same size. This is an important point to discuss the azimuthal dependence.

If initial-state-radiation effects are not too important, the SPS yield (dominated by $2 \rightarrow 2$ scatterings) tends to peak at $\Delta\phi_{W\psi} \sim \pi$ –back-to-back scatterings– whereas the DPS one is believed to be evenly spread in $\Delta\phi_{W\psi}$. Fig. 4 shows the event azimuthal distribution for prompt $J/\psi + W^\pm$ production. At first sight, it seems improbable that a yield largely dominated by DPS could be produced in agreement with such a distribution.

Just like for $J/\psi + Z$ [55], we note that the ATLAS acceptance for high- $P_T^{J/\psi}$ events is significantly higher than at low $P_T^{J/\psi}$. This, along with the very different SPS/DPS ratios as a function of $P_T^{J/\psi}$, can lead to a misinterpretation of such an azimuthal dependence which is not corrected in acceptance.

In particular, the peak near $\Delta\phi_{W\psi} \sim \pi$, visible in the raw event distribution, could artificially be accentuated due to a large acceptance of high- $P_T^{J/\psi}$ events.

Since we are not in the position of correcting the data, the simplest way to proceed is to mimic the folding of the theory event distribution with an approximated ATLAS acceptance. To do so, we use the J/ψ -acceptance dependence

CEM, NLO	CSM, direct, LO [58]	Combined
$6.1^{+2.4}_{-1.3} \text{ exp } +1.6 \text{ spin } +0.1 \text{ theo}$	$5.8^{+2.2}_{-1.2} \text{ exp } +1.5 \text{ spin } +0.05 \text{ theo}$	$6.1^{+3.3}_{-1.9} \text{ exp } +0.1 \text{ spin } -0.3 \text{ theo}$

Table 3: Different extractions of σ_{eff} (in units of mb). The experimental uncertainties on our combination conservatively account for the “spin” uncertainty (see the remark above).




Figure 3: Comparison of our range for σ_{eff} ($6.1^{+3.3}_{-1.9}$ mb) extracted from the $J/\psi + W$ data with other extractions [15, 17, 25–31, 40, 48, 49].

inferred in [55] where we assumed the background-over-signal ratio to be like $B/S \propto 1/P_T^{J/\psi}$. Reading out the statistical uncertainties for 4 bins of [3], we can derive the yield in each $P_T^{J/\psi}$ bin. In these, we can then compute the DPS-over-SPS ratio, derive the corresponding azimuthal dependence and finally combine them for the 4 bins using the yield in each. The resulting “theory” distribution is shown on Fig. 4 and agrees within uncertainties with the uncorrected ATLAS distribution. On the way, we note that the number of SPS events following our computation at NLO CEM is 2 ± 1 to be compared to 29^{+8}_{-7} events observed by ATLAS [3]. We expected these 2 ± 1 SPS events to lie at $\Delta\phi_{W\psi} \sim \pi$. Conversely, we expect an updated ATLAS $\Delta\phi_{W\psi}$ analysis in 2 $P_T^{J/\psi}$ bins to show a flat behaviour in the “low” $P_T^{J/\psi}$ bins and a slightly peaked one for the “high” $P_T^{J/\psi}$ bins. It however seems that more statistics is needed to draw final conclusions.

4. Conclusions

We have re-analysed the associated production of a prompt J/ψ with a W boson at the LHC in view of the ATLAS data [3]. To do so, we have performed the first NLO calculation of the SPS cross sections in the CEM which we consider to be a conservative upper limit of the SPS yield. This has allowed us to claim a 3- σ deviation between SPS theory and the ATLAS data, which we interpret as a strong




Figure 4: Comparison between the (uncorrected) ATLAS azimuthal event distribution and our NLO theoretical results for $J/\psi + W$ in the CEM (SPS + DPS) effectively folded with an assumed ATLAS acceptance.

indication for DPSs. Our conclusion does not depend on the quarkonium-production model used to compute the SPS yields.

We have then determined a finite range for σ_{eff} around

6 mb, consistent with extractions from other experiments and other quarkonium-related processes. In particular, we emphasise that our result is consistent with σ_{eff} extracted from our analysis of prompt $J/\psi + Z$ production as measured by ATLAS.

Whereas the azimuthal dependence could be very useful in separating the SPSs from the DPSs, one has to be very cautious with uncorrected raw event distribution. A further analysis with higher statistics allowing one to study the rapidity-separation spectrum and a fully corrected azimuthal distribution will further constrain the range for σ_{eff} since the theoretical uncertainties on the SPS happen to be quasi irrelevant at low $P_T^{J/\psi}$.

Acknowledgements

We thank V. Kartvelishvili and D. Price for useful discussions. The work of J.P.L. is supported in part by the French IN2P3-CNRS via the LIA FCPPL (Quarkonium4AFTER) and the project TMD@NLO. H.S.S. is supported by the ILP Labex (ANR-11-IDEX-0004-02, ANR-10-LABX-63). N.Y. is supported by a JSPS Postdoctoral Fellowships for Research Abroad and by then Riken iTHES Project.

References

- [1] CDF Collaboration, D. Acosta *et al.*, “Search for associated production of Υ and vector boson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV,” *Phys. Rev. Lett.* **90** (2003) 221803.
- [2] CDF Collaboration, T. Aaltonen *et al.*, “Search for production of an $\Upsilon(1S)$ meson in association with a W or Z boson using the full 1.96TeV $p\bar{p}$ collision data set at CDF,” *Phys. Rev. D* **91** no. 5, (2015) 052011, [arXiv:1412.4827 \[hep-ex\]](#).
- [3] ATLAS Collaboration, G. Aad *et al.*, “Measurement of the production cross section of prompt J/ψ mesons in association with a W^\pm boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector,” *JHEP* **04** (2014) 172, [arXiv:1401.2831 \[hep-ex\]](#).
- [4] ATLAS Collaboration, G. Aad *et al.*, “Observation and measurements of the production of prompt and non-prompt J/ψ mesons in association with a Z boson in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector,” *Eur. Phys. J. C* **75** no. 5, (2015) 229, [arXiv:1412.6428 \[hep-ex\]](#).
- [5] D. P. Roy and K. Sridhar, “ $J/\psi + \gamma$ production at the Tevatron energy,” *Phys. Lett. B* **341** (1995) 413–418, [arXiv:hep-ph/9407390 \[hep-ph\]](#).
- [6] P. Mathews, K. Sridhar, and R. Basu, “ $J/\psi + \gamma$ production at the CERN LHC,” *Phys. Rev. D* **60** (1999) 014009, [arXiv:hep-ph/9901276 \[hep-ph\]](#).
- [7] R. Li and J.-X. Wang, “Next-to-Leading-Order QCD corrections to $J/\psi(v) + \gamma$ production at the LHC,” *Phys. Lett. B* **672** (2009) 51–55, [arXiv:0811.0963 \[hep-ph\]](#).
- [8] J. P. Lansberg, “Real next-to-next-to-leading-order QCD corrections to J/ψ and Upsilon hadroproduction in association with a photon,” *Phys. Lett. B* **679** (2009) 340–346, [arXiv:0901.4777 \[hep-ph\]](#).
- [9] R. Li and J.-X. Wang, “Next-to-leading-order study of the associated production of $J/\psi + \gamma$ at the LHC,” *Phys. Rev. D* **89** no. 11, (2014) 114018, [arXiv:1401.6918 \[hep-ph\]](#).
- [10] M. A. Doncheski and C. S. Kim, “Associated J/ψ + gamma production as a probe of the polarized gluon distribution,” *Phys. Rev. D* **49** (1994) 4463–4468, [arXiv:hep-ph/9303248 \[hep-ph\]](#).
- [11] W. J. den Dunnen, J. P. Lansberg, C. Pisano, and M. Schlegel, “Accessing the Transverse Dynamics and Polarization of Gluons inside the Proton at the LHC,” *Phys. Rev. Lett.* **112** (2014) 212001, [arXiv:1401.7611 \[hep-ph\]](#).
- [12] M. N. Doroshenko, V. G. Kartvelishvili, E. G. Chikovani, and S. M. Esakiya, “Vector Quarkonium In Decays Of Heavy Higgs Particles,” *Yad. Fiz.* **46** (1987) 864–868. [*Sov. J. Nucl. Phys.* **46**,493(1987)].
- [13] G. T. Bodwin, F. Petriello, S. Stoynev, and M. Velasco, “Higgs boson decays to quarkonia and the $H\bar{c}c$ coupling,” *Phys. Rev. D* **88** no. 5, (2013) 053003, [arXiv:1306.5770 \[hep-ph\]](#).
- [14] LHCb Collaboration, R. Aaij *et al.*, “Observation of J/ψ pair production in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **707** (2012) 52–59, [arXiv:1109.0963 \[hep-ex\]](#).
- [15] D0 Collaboration, V. M. Abazov *et al.*, “Observation and studies of double J/ψ production at the Tevatron,” *Phys. Rev. D* **90** no. 11, (2014) 111101, [arXiv:1406.2380 \[hep-ex\]](#).
- [16] CMS Collaboration, V. Khachatryan *et al.*, “Measurement of prompt J/ψ pair production in pp collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **09** (2014) 094, [arXiv:1406.0484 \[hep-ex\]](#).
- [17] ATLAS Collaboration, M. Aaboud *et al.*, “Measurement of the prompt J/ψ pair production cross-section in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector,” *Eur. Phys. J. C* **77** no. 2, (2017) 76, [arXiv:1612.02950 \[hep-ex\]](#).
- [18] LHCb Collaboration, R. Aaij *et al.*, “Measurement of the J/ψ pair production cross-section in pp collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **06** (2017) 047, [arXiv:1612.07451 \[hep-ex\]](#).
- [19] NA3 Collaboration, J. Badier *et al.*, “Evidence for $\psi\psi$ Production in π^- Interactions at 150-GeV/c and 280-GeV/c,” *Phys. Lett. B* **114** (1982) 457–460.
- [20] NA3 Collaboration, J. Badier *et al.*, “ $\psi\psi$ Production and Limits on Beauty Meson Production From 400-GeV/c Protons,” *Phys. Lett. B* **158B** (1985) 85. [401(1985)].
- [21] G. T. Bodwin, E. Braaten, and G. P. Lepage, “Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium,” *Phys. Rev. D* **51** (1995) 1125–1171, [arXiv:hep-ph/9407339 \[hep-ph\]](#). [Erratum: *Phys. Rev. D* **55**,5853(1997)].
- [22] J.-P. Lansberg and H.-S. Shao, “Production of $J/\psi + \eta_c$ versus $J/\psi + J/\psi$ at the LHC: Importance of Real α_s^5 Corrections,” *Phys. Rev. Lett.* **111** (2013) 122001, [arXiv:1308.0474 \[hep-ph\]](#).
- [23] L.-P. Sun, H. Han, and K.-T. Chao, “Impact of J/ψ pair production at the LHC and predictions in nonrelativistic QCD,” *Phys. Rev. D* **94** no. 7, (2016) 074033, [arXiv:1404.4042 \[hep-ph\]](#).
- [24] A. K. Likhoded, A. V. Luchinsky, and S. V. Poslavsky, “Production of $J/\psi + \chi_c$ and $J/\psi + J/\psi$ with real gluon emission at LHC,” *Phys. Rev. D* **94** no. 5, (2016) 054017, [arXiv:1606.06767 \[hep-ph\]](#).
- [25] Axial Field Spectrometer Collaboration, T. Akesson *et al.*, “Double Parton Scattering in pp Collisions at $\sqrt{s} = 63$ -GeV,” *Z. Phys. C* **34** (1987) 163.
- [26] UA2 Collaboration, J. Alitti *et al.*, “A Study of multi - jet events at the CERN anti-p p collider and a search for double parton scattering,” *Phys. Lett. B* **268** (1991) 145–154.
- [27] CDF Collaboration, F. Abe *et al.*, “Study of four jet events and evidence for double parton interactions in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV,” *Phys. Rev. D* **47** (1993) 4857–4871.
- [28] CDF Collaboration, F. Abe *et al.*, “Double parton scattering in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV,” *Phys. Rev. D* **56** (1997) 3811–3832.
- [29] D0 Collaboration, V. M. Abazov *et al.*, “Double parton interactions in $\gamma + 3$ jet events in $p\bar{p}$ collisions $\sqrt{s} = 1.96$ TeV,” *Phys. Rev. D* **81** (2010) 052012, [arXiv:0912.5104 \[hep-ex\]](#).
- [30] ATLAS Collaboration, G. Aad *et al.*, “Measurement of hard double-parton interactions in $W(\rightarrow l\nu) + 2$ jet events at $\sqrt{s} = 7$ TeV with the ATLAS detector,” *New J. Phys.* **15** (2013) 033038, [arXiv:1301.6872 \[hep-ex\]](#).
- [31] CMS Collaboration, S. Chatrchyan *et al.*, “Study of double parton scattering using W + 2-jet events in proton-proton collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **03** (2014) 032, [arXiv:1312.5729 \[hep-ex\]](#).
- [32] V. G. Kartvelishvili and S. M. Esakiya, “On Hadron Induced Production OF J / Psi Meson Pairs. (IN RUSSIAN),” *Yad. Fiz.* **38** (1983) 722–726.
- [33] B. Humpert and P. Mery, “psi psi Production At Collider Energies,” *Z. Phys. C* **20** (1983) 83.
- [34] R. Vogt and S. J. Brodsky, “Intrinsic charm contribution to double quarkonium hadroproduction,” *Phys. Lett. B* **349** (1995) 569–575, [arXiv:hep-ph/9503206 \[hep-ph\]](#).
- [35] R. Li, Y.-J. Zhang, and K.-T. Chao, “Pair Production of Heavy

- Quarkonium and $B(c)^*$ Mesons at Hadron Colliders,” *Phys. Rev. D* **80** (2009) 014020, [arXiv:0903.2250 \[hep-ph\]](#).
- [36] C.-F. Qiao, L.-P. Sun, and P. Sun, “Testing Charmonium Production Mechanism via Polarized J/ψ Pair Production at the LHC,” *J. Phys. G* **37** (2010) 075019, [arXiv:0903.0954 \[hep-ph\]](#).
- [37] P. Ko, C. Yu, and J. Lee, “Inclusive double-quarkonium production at the Large Hadron Collider,” *JHEP* **01** (2011) 070, [arXiv:1007.3095 \[hep-ph\]](#).
- [38] A. V. Berezhnoy, A. K. Likhoded, A. V. Luchinsky, and A. A. Novoselov, “Double J/ψ -meson Production at LHC and 4c-tetraquark state,” *Phys. Rev. D* **84** (2011) 094023, [arXiv:1101.5881 \[hep-ph\]](#).
- [39] Y.-J. Li, G.-Z. Xu, K.-Y. Liu, and Y.-J. Zhang, “Relativistic Correction to J/ψ and Upsilon Pair Production,” *JHEP* **07** (2013) 051, [arXiv:1303.1383 \[hep-ph\]](#).
- [40] J.-P. Lansberg and H.-S. Shao, “ J/ψ -pair production at large momenta: Indications for double parton scatterings and large α_s^5 contributions,” *Phys. Lett. B* **751** (2015) 479–486, [arXiv:1410.8822 \[hep-ph\]](#).
- [41] Z.-G. He and B. A. Kniehl, “Complete Nonrelativistic-QCD Prediction for Prompt Double J/ψ Hadroproduction,” *Phys. Rev. Lett.* **115** no. 2, (2015) 022002, [arXiv:1609.02786 \[hep-ph\]](#).
- [42] S. P. Baranov and A. H. Rezaeian, “Prompt double J/ψ production in proton-proton collisions at the LHC,” *Phys. Rev. D* **93** no. 11, (2016) 114011, [arXiv:1511.04089 \[hep-ph\]](#).
- [43] J.-P. Lansberg and H.-S. Shao, “Double-quarkonium production at a fixed-target experiment at the LHC (AFTER@LHC),” *Nucl. Phys. B* **900** (2015) 273–294, [arXiv:1504.06531 \[hep-ph\]](#).
- [44] C. Borschensky and A. Kulesza, “Double parton scattering in pair production of J/ψ mesons at the LHC revisited,” *Phys. Rev. D* **95** no. 3, (2017) 034029, [arXiv:1610.00666 \[hep-ph\]](#).
- [45] J.-P. Lansberg, C. Pisano, F. Scarpa, and M. Schlegel, “Pinning down the linearly-polarized gluons inside unpolarized protons using quarkonium-pair production at the LHC,” [arXiv:1710.01684 \[hep-ph\]](#).
- [46] D0 Collaboration, V. M. Abazov *et al.*, “Evidence for simultaneous production of J/ψ and Υ mesons,” *Phys. Rev. Lett.* **116** no. 8, (2016) 082002, [arXiv:1511.02428 \[hep-ex\]](#).
- [47] H.-S. Shao and Y.-J. Zhang, “Complete study of hadroproduction of a Υ meson associated with a prompt J/ψ ,” *Phys. Rev. Lett.* **117** no. 6, (2016) 062001, [arXiv:1605.03061 \[hep-ph\]](#).
- [48] LHCb Collaboration, R. Aaij *et al.*, “Observation of double charm production involving open charm in pp collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **06** (2012) 141, [arXiv:1205.0975 \[hep-ex\]](#). [Addendum: JHEP03,108(2014)].
- [49] LHCb Collaboration, R. Aaij *et al.*, “Production of associated Υ and open charm hadrons in pp collisions at $\sqrt{s} = 7$ and 8 TeV via double parton scattering,” *JHEP* **07** (2016) 052, [arXiv:1510.05949 \[hep-ex\]](#).
- [50] P. Artoisenet, J. P. Lansberg, and F. Maltoni, “Hadroproduction of J/ψ and Υ in association with a heavy-quark pair,” *Phys. Lett. B* **653** (2007) 60–66, [arXiv:hep-ph/0703129 \[hep-ph\]](#).
- [51] S. P. Baranov, “Topics in associated J/ψ + c + anti-c production at modern colliders,” *Phys. Rev. D* **73** (2006) 074021.
- [52] A. V. Berezhnoy and A. K. Likhoded, “Associated production of and open charm at LHC,” *Int. J. Mod. Phys. A* **30** no. 20, (2015) 1550125, [arXiv:1503.04445 \[hep-ph\]](#).
- [53] S. Mao, M. Wen-Gan, L. Gang, Z. Ren-You, and G. Lei, “QCD corrections to J/ψ plus Z^0 -boson production at the LHC,” *JHEP* **02** (2011) 071, [arXiv:1102.0398 \[hep-ph\]](#). [Erratum: JHEP12,010(2012)].
- [54] B. Gong, J.-P. Lansberg, C. Lorce, and J. Wang, “Next-to-leading-order QCD corrections to the yields and polarisations of J/ψ and Upsilon directly produced in association with a Z boson at the LHC,” *JHEP* **03** (2013) 115, [arXiv:1210.2430 \[hep-ph\]](#).
- [55] J.-P. Lansberg and H.-S. Shao, “Associated production of a quarkonium and a Z boson at one loop in a quark-hadron-duality approach,” *JHEP* **10** (2016) 153, [arXiv:1608.03198 \[hep-ph\]](#).
- [56] J.-P. Lansberg and H.-S. Shao, “Phenomenological analysis of associated production of $Z^0 + b$ in the $b \rightarrow J/\psi X$ decay channel at the LHC,” *Nucl. Phys. B* **916** (2017) 132–142, [arXiv:1611.09303 \[hep-ph\]](#).
- [57] G. Li, M. Song, R.-Y. Zhang, and W.-G. Ma, “QCD corrections to J/ψ production in association with a W-boson at the LHC,” *Phys. Rev. D* **83** (2011) 014001, [arXiv:1012.3798 \[hep-ph\]](#).
- [58] J. P. Lansberg and C. Lorce, “Reassessing the importance of the colour-singlet contributions to direct $J/\psi + W$ production at the LHC and the Tevatron,” *Phys. Lett. B* **726** (2013) 218–222, [arXiv:1303.5327 \[hep-ph\]](#). [Erratum: Phys. Lett. B738,529(2014)].
- [59] V. D. Barger, S. Fleming, and R. J. N. Phillips, “Double gluon fragmentation to J/ψ pairs at the Tevatron,” *Phys. Lett. B* **371** (1996) 111–116, [arXiv:hep-ph/9510457 \[hep-ph\]](#).
- [60] B. A. Kniehl, C. P. Palisoc, and L. Zwierner, “Associated production of heavy quarkonia and electroweak bosons at present and future colliders,” *Phys. Rev. D* **66** (2002) 114002, [arXiv:hep-ph/0208104 \[hep-ph\]](#).
- [61] H. Fritzsch, “Producing Heavy Quark Flavours in Hadronic Collisions: A Test of Quantum Chromodynamics,” *Phys. Lett. B* **67** (1977) 217–221.
- [62] F. Halzen, “Cvc for Gluons and Hadroproduction of Quark Flavours,” *Phys. Lett. B* **69** (1977) 105–108.
- [63] H. S. Shao, H. Han, Y. Q. Ma, C. Meng, Y. J. Zhang, and K. T. Chao, “Yields and polarizations of prompt J/ψ and $\psi(2S)$ production in hadronic collisions,” *JHEP* **05** (2015) 103, [arXiv:1411.3300 \[hep-ph\]](#).
- [64] J. P. Lansberg, “ J/ψ , ψ' and ν production at hadron colliders: A Review,” *Int. J. Mod. Phys. A* **21** (2006) 3857–3916, [arXiv:hep-ph/0602091 \[hep-ph\]](#).
- [65] N. Brambilla *et al.*, “Heavy quarkonium: progress, puzzles, and opportunities,” *Eur. Phys. J. C* **71** (2011) 1534, [arXiv:1010.5827 \[hep-ph\]](#).
- [66] A. Andronic *et al.*, “Heavy-flavour and quarkonium production in the LHC era: from proton-proton to heavy-ion collisions,” *Eur. Phys. J. C* **76** no. 3, (2016) 107, [arXiv:1506.03981 \[nucl-ex\]](#).
- [67] A. Edin, G. Ingelman, and J. Rathsman, “Quarkonium production at the Tevatron through soft color interactions,” *Phys. Rev. D* **56** (1997) 7317–7320, [arXiv:hep-ph/9705311 \[hep-ph\]](#).
- [68] J. Damet, G. Ingelman, and C. Brenner Mariotto, “Prompt J/ψ production at the LHC,” *JHEP* **09** (2002) 014, [arXiv:hep-ph/0111463 \[hep-ph\]](#).
- [69] C. Brenner Mariotto, M. B. Gay Ducati, and G. Ingelman, “Soft and hard QCD dynamics in hadroproduction of charmonium,” *Eur. Phys. J. C* **23** (2002) 527–538, [arXiv:hep-ph/0111379 \[hep-ph\]](#).
- [70] Y.-Q. Ma and R. Vogt, “Quarkonium Production in an Improved Color Evaporation Model,” *Phys. Rev. D* **94** no. 11, (2016) 114029, [arXiv:1609.06042 \[hep-ph\]](#).
- [71] ATLAS Collaboration, G. Aad *et al.*, “Measurement of the differential cross-sections of prompt and non-prompt production of J/ψ and $\psi(2S)$ in pp collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS detector,” *Eur. Phys. J. C* **76** no. 5, (2016) 283, [arXiv:1512.03657 \[hep-ex\]](#).
- [72] R. E. Nelson, R. Vogt, and A. D. Frawley, “Narrowing the uncertainty on the total charm cross section and its effect on the J/ψ cross section,” *Phys. Rev. C* **87** no. 1, (2013) 014908, [arXiv:1210.4610 \[hep-ph\]](#).
- [73] M. Bedjidian *et al.*, “Hard probes in heavy ion collisions at the LHC: Heavy flavor physics,” 2004. [arXiv:hep-ph/0311048 \[hep-ph\]](#). <http://doc.cern.ch/cernrep/2004/2004-009/2004-009.html>.
- [74] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations,” *JHEP* **07** (2014) 079, [arXiv:1405.0301 \[hep-ph\]](#).
- [75] P. Artoisenet, F. Maltoni, and T. Stelzer, “Automatic generation of quarkonium amplitudes in NRQCD,” *JHEP* **02** (2008) 102, [arXiv:0712.2770 \[hep-ph\]](#).
- [76] H.-S. Shao, “HELAC-Onia: An automatic matrix element generator for heavy quarkonium physics,” *Comput. Phys. Commun.* **184** (2013) 2562–2570, [arXiv:1212.5293 \[hep-ph\]](#).

- [77] H.-S. Shao, “HELAC-Onia 2.0: an upgraded matrix-element and event generator for heavy quarkonium physics,” *Comput. Phys. Commun.* **198** (2016) 238–259, [arXiv:1507.03435 \[hep-ph\]](#).
- [78] R. D. Ball *et al.*, “Parton distributions with LHC data,” *Nucl. Phys.* **B867** (2013) 244–289, [arXiv:1207.1303 \[hep-ph\]](#).
- [79] A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rfenacht, M. Schnherr, and G. Watt, “LHAPDF6: parton density access in the LHC precision era,” *Eur. Phys. J.* **C75** (2015) 132, [arXiv:1412.7420 \[hep-ph\]](#).
- [80] ATLAS Collaboration, G. Aad *et al.*, “Measurement of the inclusive W^\pm and Z/γ cross sections in the electron and muon decay channels in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector,” *Phys. Rev.* **D85** (2012) 072004, [arXiv:1109.5141 \[hep-ex\]](#).
- [81] R. Gavin, Y. Li, F. Petriello, and S. Quackenbush, “FEWZ 2.0: A code for hadronic Z production at next-to-next-to-leading order,” *Comput. Phys. Commun.* **182** (2011) 2388–2403, [arXiv:1011.3540 \[hep-ph\]](#).
- [82] R. Gavin, Y. Li, F. Petriello, and S. Quackenbush, “W Physics at the LHC with FEWZ 2.1,” *Comput. Phys. Commun.* **184** (2013) 208–214, [arXiv:1201.5896 \[hep-ph\]](#).
- [83] Particle Data Group Collaboration, C. Patrignani *et al.*, “Review of Particle Physics,” *Chin. Phys.* **C40** no. 10, (2016) 100001.
- [84] CMS Collaboration, S. Chatrchyan *et al.*, “Measurement of the prompt J/ψ and $\psi(2S)$ polarizations in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett.* **B727** (2013) 381–402, [arXiv:1307.6070 \[hep-ex\]](#).
- [85] B. Blok and M. Strikman, “Open charm production in Double Parton Scattering processes in the forward kinematics,” *Eur. Phys. J.* **C76** no. 12, (2016) 694, [arXiv:1608.00014 \[hep-ph\]](#).