

The Covering Principle: A New Approach to Address Multiplicity in Hypotheses Testing

Huajiang Li

Email: HLi@avanir.com

Avanir Pharmaceuticals, Inc.

30 Enterprise, Aliso Viejo, California, 92656, U.S.A.

and

Hong Zhou

Email: hzhou@astate.edu

Department of Mathematics and Statistics, Arkansas State University

P.O.Box 70, State University, Arkansas, 72467, U.S.A.

February 27, 2022

Abstract

The closure and the partitioning principles have been used to build various multiple testing procedures in the past three decades. The essence of these two principles is based on parameter space partitioning. In this article, we propose a novel approach coined the covering principle from the perspective of rejection region coverage in the sample space. The covering principle divides the whole family of null hypotheses into a few overlapped sub-families when there is a priority of making decisions for hypothesis testing. We have proven that the multiple testing procedure constructed by the covering principle strongly controls the familywise error rate as long as the multiple tests for each sub-familiy strongly control the type I error. We have illustrated the covering principle can be applied to solve the general gate-keeping problems.

Keywords: Familywise error rate; Multiple testing; Closure principle; Partitioning principle; Covering principle; Gate Keeping.

1 Introduction

The key issue in the multiple hypotheses testing is to control the familywise error rate. Commonly, there are a few ways to deal with the multiplicity issue. First is to cut the spending of the significance level α as it does in Bonferroni procedure and its modifications (Holm, 1979; Hochberg, 1988; Hommel, 1988; Li *et al.*, 2017). Second is to plan the order of testing the null hypotheses as it does in the gate-keeping procedures (Dmitrienko and Tamhane, 2007; Dmitrienko *et al.*, 2008a; Dmitrienko *et al.*, 2008b; Dmitrienko *et al.*, 2010; Dmitrienko and Tamhane, 2013). Third is to make additional assumptions such as the independence among the null hypotheses or the test statistics following the multivariate normal distribution (Tamhane *et al.*, 1998).

Two important principles: the closure principle (Marcus *et al.*, 1976) and the partitioning principle (Finner and Strassburger, 2002; Sonnemann, 2008), are widely used to construct various multiple test procedures that can strongly control the familywise error rate. In the clinical trials, however, the multiple study objectives usually exhibit a hierarchical structure. That is, the objectives can be divided into different tiers according to their importance, namely, primary, secondary, tertiary, and so on. In this article, we introduce a novel principle termed the covering principle for the construction of the multiple testing procedures. The covering principle analyzes the rejection regions in the sample space based on the priorities of the decisions for testing the null hypotheses and divides the whole family of null hypotheses into a few overlapped sub-families, for which either the closure or the partitioning principle can be used. Section 2 introduces the theorem of the covering principle mathematically. Then we will apply the covering principle to a real clinical trial as well as a general gate-keeping problem in Section 3. The significance and importance using the covering principle are discussed in Section 4. Finally, we will prove Theorem 1 that the familywise error rate is strongly controlled for the whole family as long as it is controlled in each sub-family in Appendix.

2 The Covering Principle

Denote $N = \{1, 2, \dots, n\}$ as the index set of a family of n null hypotheses $\{H_1, H_2, \dots, H_n\}$ with the corresponding test functions $\phi = \{\phi_1, \phi_2, \dots, \phi_n\}$ and rejection regions R_1, R_2, \dots, R_n . Each ϕ_i ($i = 1, 2, \dots, n$) is an elementary test function, where

$$\phi_i = \begin{cases} 1, & \text{if } H_i \text{ is rejected} \\ 0, & \text{if } H_i \text{ is accepted.} \end{cases}$$

For $\emptyset \neq S \subseteq N$, let $\Phi_\alpha(S)$ denote the set of all α -level multiple tests for the family of null hypotheses with an index set S , where $0 < \alpha < 1$. If $\phi = \{\phi_i : i \in S\} \in \Phi_\alpha(S)$, then it indicates the multiple test ϕ strongly controls the familywise error rate on S at the significance level α . For a group of elementary test functions $\phi_i, i \in S$, define

$$\min_{i \in S} \phi_i = \begin{cases} 1, & \text{if } \forall i \in S, \phi_i = 1 \\ 0, & \text{otherwise.} \end{cases}$$

and

$$\max_{i \in S} \phi_i = \begin{cases} 1, & \text{if } \exists i \in S, \phi_i = 1 \\ 0, & \text{otherwise.} \end{cases}$$

For any two elementary test functions ϕ_1 and ϕ_2 , denote $\phi_1 \leq \phi_2$ if $\{\phi_1 = 1\}$ implies $\{\phi_2 = 1\}$; in terms of the rejection regions, $R_1 \subseteq R_2$.

Theorem 1 (Covering principle) Suppose $\emptyset \neq I \subset N$, $\emptyset \neq J \subset N$, $I \cap J = \emptyset$, and $\bigcup_{i \in I} R_i \subseteq \bigcup_{j \in J} R_j$. Denote

$$\phi^j = \{\phi_i^j : i \in N \setminus j\}, \forall j \in J. \quad (1)$$

$$\phi^I = \{\phi_i^I : i \in N \setminus I\}. \quad (2)$$

where ϕ^j denote a group of $|J|$ multiple tests which does not include the j th hypothesis, $j \in J$. Each ϕ^j consists of $|N \setminus j|$ elementary test functions $\phi_i^j, i \in N \setminus j$. Similarly, ϕ^I denote a multiple test which consists of $|N \setminus I|$ elementary test functions $\phi_i^I, i \in N \setminus I$, whose indices are not in I . Define

$$\psi_i = \begin{cases} \min(\min_{j \in J} \phi_i^j, \phi_i^I), & \text{if } i \in N \setminus I \\ \min(\min_{j \in J} \phi_i^j, \max_{j \in J} \psi_j), & \text{if } i \in I. \end{cases} \quad (3)$$

If $\phi^j \in \Phi_\alpha(N \setminus j)$, $\forall j \in J$ and $\phi^I \in \Phi_\alpha(N \setminus I)$, then $\{\psi_i : i \in N\} \in \Phi_\alpha(N)$.

The following is an explanation of Theorem 1. Suppose that there exist two nonempty index sets $I \subset N$ and $J \subset N$, $I \cap J = \emptyset$, i.e. these two sets of hypotheses $H_j, j \in J$, and $H_i, i \in I$, are not overlapped. Furthermore, there are orders when these hypotheses are tested. In order for $H_i, i \in I$, to be tested, at least one of the hypothesis $H_j, j \in J$, must be tested and rejected first. For example, the hypotheses $H_j, j \in J$, could be related to the primary endpoints and $H_i, i \in I$, could be related to the secondary endpoints in clinical trials. For parallel gate-keeping, if a null hypothesis on a secondary endpoint is rejected, then at least one hypothesis on one of primary endpoints already has been rejected. If this is the case, we say the set of hypotheses $H_j, j \in J$, dominates the set of hypotheses $H_i, i \in I$. Simply say, there is a dominance relationship between hypotheses $H_i, i \in I$, and $H_j, j \in J$. From the perspective of hypotheses testing, the orders of testing multiple hypotheses can be illustrated in terms of the logical relationship among their rejection regions: $\bigcup_{i \in I} R_i \subseteq \bigcup_{j \in J} R_j$.

It may seem that the definition of ψ_i in Equation 3 is circular. In fact, domains for the index i are mutually exclusive. For the first part of the definition of the test function ψ_i , hypotheses $H_i, i \in N \setminus I$, could be either hypotheses $H_j, j \in J$, which dominate $H_i, i \in I$, or independent hypotheses. An independent hypothesis is the one that has no dominance relationship with other hypotheses. The first part of ψ_i defines a test function to reject any hypothesis H_i whose index is not in I . That is, $H_i, i \in N \setminus I$, will be rejected if it is rejected in all subsets which contain it.

The second part of ψ_i in Equation 3 defines test functions for those hypotheses whose indices are within I . These hypotheses $H_i, i \in I$, are dominated by $H_j, j \in J$. In order for any hypothesis $H_i, i \in I$, to be rejected, not only at least one $H_j, j \in J$, must be rejected first, but also $H_i, i \in I$, must be rejected in all subsets which contain it.

Then, the covering principle in Theorem 1 states that the original whole family of n null hypotheses can be decomposed into $|J| + 1$ sub-families with index sets $N \setminus I$ and $N \setminus j, \forall j \in J$. In other words, the original multiple testing problem on the family of n null hypotheses with the index set N can be divided into $|J| + 1$ multiple testing problems. The corresponding multiple tests are ϕ^I with the index set $N \setminus I$ and ϕ^j with index sets $N \setminus j, \forall j \in J$. Each subset has fewer null hypotheses than the original family and can be tested using any multiple testing procedure. The multiple testing procedure built on this divide-and-conquer strategy strongly controls the familywise error rate for the whole family at the significance level α if the multiple tests ϕ^I and ϕ^j can control their

familywise error rate at the significance level α for their corresponding subsets. Finally, the decision rule for each individual hypothesis can be reached by summarizing the results as follows:

Step 1. $H_i, i \in N \setminus I$, will be rejected if H_i is rejected in all decomposed subsets in which H_i is contained;

Step 2. $H_i, i \in I$, will be rejected if at least one of the null hypotheses $H_j, j \in J$, is rejected first. In addition, H_i must be also rejected in all subsets in which H_i is contained.

Simply speaking, in order for any individual hypothesis $H_i, i \in N$, to be rejected, not only one of its precedent and dominant hypotheses in the hierarchy of the hypotheses must be rejected first, but it must also be rejected in all subsets which contain H_i . We will illustrate how to use the covering principle to build a multiple testing procedure in Section 3.

The covering principle extends the closure principle to a family of hypotheses with the priority of importance when making decisions. It performs a sample space analysis using the union of rejection regions in contrast to the closure principle using the intersection of hypotheses in the parameter space. The covering principle divides the original family of hypotheses into several sub-families based on the constraints among the relationship of the rejection regions, which is constructed by the usage of the hierarchical structure of the decisions on testing the null hypotheses. The multiple testing problem on the whole family then can be carried out on each sub-family with fewer hypotheses. Furthermore, it can strongly control the familywise error rate for the whole family at the significance level α if the multiple tests for every sub-family control type I error at the significance level α .

3 Applications of covering principle

Dmitrienko et al. (2007, 2008a, 2008b, 2010, 2013) recently proposed an array of the gate-keeping procedures including tree-structured, multistage and mixture procedures for the general gate-keeping problems.

The gate-keeping procedures use the hierarchical structure among the the multiple study objectives. It divides the whole family of the null hypotheses into several ordered sub-families F_1, \dots, F_n . F_i serves as the gatekeeper of F_{i+1} , that means in order for the null hypotheses in F_{i+1} to be tested F_i must be rejected, i.e. either all the null hypotheses in F_i must be rejected (serial gate-keeping) or at least one null hypothesis in F_i must be rejected (parallel gate-keeping). In other words, if F_i is not rejected, then F_{i+1} will be accepted automatically.

The covering principle provides a very general approach to the gate-keeping problems. The following examples will illustrate how to apply the covering principle to a real clinical trial study using parallel gate-keeping and a more general gate-keeping problem.

Example 1: Cummings et al. (1999) and Ettinger et al. (1999) studied the breast cancer with two primary endpoints: the incidence of vertebral fractures H_1 and the incidence of breast cancer H_2 , and one secondary endpoint: the incidence of non-vertebral fractures H_3 . The primary endpoint will result in an independent regulatory claim if one of two primary endpoints is effective. That is, the two primary endpoints are parallel, and anyone can serve as the gatekeeper. The test in the secondary endpoint can be carried out as long as at least one of the null hypotheses in the primary family $\{H_1, H_2\}$ is rejected.

Let R_1, R_2 and R_3 denote three rejection regions according to null hypotheses H_1, H_2 and H_3 , respectively. Because of the logic constraints among the decisions on testing three null hypotheses, the corresponding rejection regions exhibit the coverage relationship: $R_3 \subseteq R_1 \cup R_2$ as illustrated in Figure 1(a). According to the covering principle, $N = \{1, 2, 3\}$, $I = \{3\}$ and $J = \{1, 2\}$, the three

null hypotheses H_1, H_2 and H_3 can be divided into $|J| + 1 = 3$ sub-families as follows: $N \setminus I = \{1, 2\}$, $N \setminus 1 = \{2, 3\}$, and $N \setminus 2 = \{1, 3\}$. The decision rule for each individual hypothesis is as follows:

Step 1. H_1 will be rejected if it is rejected in sub-families $\{H_1, H_2\}$ and $\{H_1, H_3\}$. Similarly, H_2 will be rejected if it is rejected in sub-families $\{H_1, H_2\}$ and $\{H_2, H_3\}$.

Step 2. In order to reject H_3 , either H_1 or H_2 must be rejected at *Step 1*, and H_3 must also be rejected in sub-families $\{H_2, H_3\}$ and $\{H_1, H_3\}$.




Figure 1: (a) Parallel Gate-Keeping (b) General Gate-Keeping

Example 2: Figure 1(b) displays a more complicated scenario for which a general gate-keeping procedure can be used. Suppose that three pairs $\{H_1, H_2\}$, $\{H_3, H_4\}$, and $\{H_5, H_6\}$ are parallel to each other and form three tiers. In addition, $\{H_1, H_3, H_5\}$ forms a serial relation as well as $\{H_2, H_4, H_6\}$. In a clinical trial, H_1 and H_2 may represent hypotheses on the primary endpoints, H_3 and H_4 on the secondary endpoints, H_5 and H_6 on the tertiary endpoints. H_1, H_3, H_5 may represent hypotheses on a treatment, H_2, H_4, H_6 on another treatment. Based on the relationship among the decisions on testing the six null hypotheses, the rejection regions exhibit the following coverage relationship: $R_5 \subseteq R_3 \subseteq R_1$ and $R_6 \subseteq R_4 \subseteq R_2$.

The covering principle will be used recursively in this example. First, the whole family with six null hypotheses $\{H_1, H_2, H_3, H_4, H_5, H_6\}$ will be decomposed into $\{H_1, H_2\}$, $\{H_2, H_3, H_4, H_5, H_6\}$, and $\{H_1, H_3, H_4, H_5, H_6\}$ by using the relationship among rejection regions $R_3 \cup R_4 \cup R_5 \cup R_6 \subseteq R_1 \cup R_2$, that is, $I = \{3, 4, 5, 6\}$ and $J = \{1, 2\}$. Second, the sub-family $\{H_2, H_3, H_4, H_5, H_6\}$ will be decomposed into $\{H_2, H_3, H_5\}$ and $\{H_3, H_4, H_5, H_6\}$ by using the relationship $R_6 \cup R_4 \subseteq R_2$, i.e. $I = \{4, 6\}$ and $J = \{2\}$. Then, $\{H_2, H_3, H_5\}$ is decomposed into $\{H_2, H_3\}$ and $\{H_2, H_5\}$ by using $R_5 \subseteq R_3$. Third, $\{H_3, H_4, H_5, H_6\}$ is decomposed into $\{H_3, H_4, H_6\}$ and $\{H_4, H_5, H_6\}$ by using $R_5 \subseteq R_3$; Fourth, sub-family $\{H_3, H_4, H_6\}$ is further separated into $\{H_3, H_4\}$ and $\{H_3, H_6\}$ by $R_6 \subseteq R_4$. $\{H_4, H_5, H_6\}$ is also separated into $\{H_4, H_5\}$ and $\{H_5, H_6\}$.

Similarly, the sub-family $\{H_1, H_3, H_4, H_5, H_6\}$ can be decomposed into six sub-families: $\{H_1, H_4\}$, $\{H_1, H_6\}$, $\{H_3, H_4\}$, $\{H_4, H_5\}$, $\{H_3, H_6\}$, $\{H_5, H_6\}$.

Combining all sub-families together, the family of six null hypotheses $\{H_1, H_2, H_3, H_4, H_5, H_6\}$ has been divided into nine sub-families with only two hypotheses each: $\{H_1, H_2\}$, $\{H_1, H_4\}$, $\{H_1, H_6\}$, $\{H_2, H_3\}$, $\{H_2, H_5\}$, $\{H_3, H_4\}$, $\{H_3, H_6\}$, $\{H_4, H_5\}$, and $\{H_5, H_6\}$. The original multiple testing problem on the 6-dimension is now reduced into several 2-dimension problems. The decision rule for each hypothesis is as follows:

Step 1. Reject H_1 if it is rejected in all sub-families: $\{H_1, H_2\}$, $\{H_1, H_4\}$, and $\{H_1, H_6\}$. The rejection of H_2 is similar to H_1 due to the symmetry.

Step 2. Reject H_3 if H_1 on its the upper level has been rejected, and H_3 is also rejected in all sub-families: $\{H_3, H_4\}$, $\{H_3, H_6\}$, and $\{H_2, H_3\}$. The rejection of H_4 is similar to H_3 due to the symmetry.

Step 3. Reject H_5 if both H_1 and H_3 on its upper levels have been rejected, and H_5 is also

rejected in all sub-families: $\{H_5, H_6\}$, $\{H_2, H_5\}$, and $\{H_4, H_5\}$. The rejection of H_6 is similar to H_5 due to the symmetry.

4 Closing Remarks

The covering principle can play a key role for the multiple testing problems when there are priorities among the decisions on testing the null hypotheses. It can also be viewed as an extension of two famous closure principle and partitioning principle. Based on the analysis of the rejection regions it decomposed the original family of null hypotheses into a group of sub-families. The merit of this decomposition is threefold. First, it reduces the dimension of the multiple testing problems since each sub-family has fewer hypotheses, which makes the multiple testing problems much easier at the lower dimensions. Second, the hierarchical structure among the decisions on testing the null hypotheses as described in the gate-keeping problems are removed by the decomposition. It makes the consolidation of testing results easy and straightforward. Third, any available multiple testing procedure can be used for each sub-family and will be more powerful with fewer hypotheses. After all, the covering principle is very intuitive and easy to understand and use in practice with the help of the flow chart of the decisions and the diagram of the rejection regions.

Acknowledgement

The research of Hong Zhou is partially supported by the Arkansas Science and Technology Authority Fund, NO. 15-B-09.

5 Appendix

Proof of Theorem 1

By the results (Finner and Strassburger (2002), Sonnemann (2008)), we have $\{\psi_i : i \in N\} \in \Phi_\alpha(N)$ if and only if $\forall \emptyset \neq S \subseteq N$, $\forall \theta \in \cap_{i \in S} H_i$, $P_\theta(\max_{i \in S} \psi_i = 1) \leq \alpha$. For $\forall \emptyset \neq S \subseteq N$, consider two cases of the relationship between S and J .

Case I: $J \not\subseteq S$. There exists a $j_0 \in J$ such that $j_0 \notin S$, then $S \subseteq N \setminus j_0$. By the definition of ψ_i in equation (3), $\psi_i \leq \min_{j \in J} \phi_i^j$, $\forall i \in N$, we have $\psi_i \leq \phi_i^{j_0}$, $j_0 \neq i \in N$. Therefore, $\max_{i \in S} \psi_i \leq \max_{i \in S} \phi_i^{j_0}$. By the assumption of Theorem 1, $\{\phi_i^{j_0} : i \in N \setminus j_0\} \in \Phi_\alpha(N \setminus j_0)$, and $S \subseteq N \setminus j_0$, hence $\forall \theta \in \cap_{i \in S} H_i$, $P_\theta(\max_{i \in S} \phi_i^{j_0} = 1) \leq \alpha$. Therefore, $P_\theta(\max_{i \in S} \psi_i = 1) \leq P_\theta(\max_{i \in S} \phi_i^{j_0} = 1) \leq \alpha$.

Case II: $J \subseteq S$.

If $S \cap I = \emptyset$, then $S = S \setminus I$, hence $\max_{i \in S} \psi_i = \max_{i \in S \setminus I} \psi_i$.

If $S \cap I \neq \emptyset$, by the definition of ψ_i in equation (3), $\psi_i \leq \max_{j \in J} \psi_j$, $i \in I$, and since $\forall i \in S \cap I \subseteq I$, hence $\max_{i \in S \cap I} \psi_i \leq \max_{j \in J} \psi_j$. Because $J \subseteq S$ and $I \cap J = \emptyset$, then $J \subseteq S \setminus I$. Since $S = (S \cap I) \cup (S \setminus I)$, we have $\max_{i \in S} \psi_i = \max(\max_{i \in S \cap I} \psi_i, \max_{i \in S \setminus I} \psi_i) \leq \max(\max_{j \in J} \psi_j, \max_{i \in S \setminus I} \psi_i) = \max_{i \in S \setminus I} \psi_i$. But $S \setminus I \subseteq S$, hence $\max_{i \in S} \psi_i \geq \max_{i \in S \setminus I} \psi_i$. Therefore $\max_{i \in S} \psi_i = \max_{i \in S \setminus I} \psi_i$.

Similarly, by the definition of ψ_i in equation (3), $\psi_i \leq \phi_i^I, i \in N \setminus I$, and since $\forall i \in S \setminus I \subseteq N \setminus I$, hence $\max_{i \in S \setminus I} \psi_i \leq \max_{i \in S \setminus I} \phi_i^I$. By the assumption of Theorem 1, $\{\phi_i^I : i \in N \setminus I\} \in \Phi_\alpha(N \setminus I)$, and $S \setminus I \subseteq N \setminus I$, then $\forall \theta \in \bigcap_{i \in S \setminus I} H_i, P_\theta(\max_{i \in S \setminus I} \phi_i^I = 1) \leq \alpha$. Therefore $\forall \theta \in \bigcap_{i \in S} H_i \subseteq \bigcap_{i \in S \setminus I} H_i, P_\theta(\max_{i \in S} \psi_i = 1) = P_\theta(\max_{i \in S \setminus I} \psi_i = 1) \leq P_\theta(\max_{i \in S \setminus I} \phi_i^I = 1) \leq \alpha$.

Combining Case I and II above, we have $\forall \emptyset \neq S \subseteq N, \forall \theta \in \bigcap_{i \in S} H_i, P_\theta(\max_{i \in S} \psi_i = 1) \leq \alpha$, therefore $\{\psi_i : i \in N\} \in \Phi_\alpha(N)$.

References

- Cummings, S. R. and Eckert, S. and Krueger, K. (1999). The Effect of Raloxifene on Risk of Breast Cancer in Postmenopausal Women. *The Journal of the American Medical Association* **281**, 2189-2197.
- Dmitrienko, A. and Tamhane, A. C. (2007). Gatekeeping procedures with clinical trial applications. *Pharmaceutical Statistics* **6**, 171-180.
- Dmitrienko, A. and Tamhane, A. C. (2013). General theory of mixture procedures for gatekeeping. *Biometrical Journal* **55**, 402-419.
- Dmitrienko A, Tamhane A. C, Bretz F. (2010). *Multiple Testing Problems in Pharmaceutical Statistics*. Chapman and Hall/CRC Press: New York.
- Dmitrienko, A. and Tamhane, A. C. and Liu, W. and Wiens, B. L. (2008a). A note on tree gatekeeping procedures in clinical trials. *Statistics in Medicine*, **27**, 3446-3451.
- Dmitrienko, A. and Tamhane, A. C. and Wiens, B. L. (2008b). General multistage gatekeeping procedures. *Biometrical Journal*, **50**, 667-677.
- Ettinger, B. and Black, D. M. and Mitlak, B. H. and Knickerbocker, R. K. and Nickelsen, T. and Genant, H. K. and Christiansen, C. and Delmas, P. D. and Zanchetta, J. R. and Stakkestad, J. and Gler, C. C. and Krueger, K. and Cohen, F. J. and Eckert, S. and Ensrud, K. E. and Avioli, L. V. and Lips, P. and Cummings, S. R. (1999). Reduction of vertebral fracture risk in postmenopausal women with osteoporosis treated with raloxifene. *The Journal of the American Medical Association* **282**, 637-645.
- Finner, H. and Strassburger, K. (2002). The partitioning principle: a powerful tool in multiple decision theory. *The Annals of Statistics* **30**, 1194-1213.
- Hochberg, Y. (1988). A sharper bonferroni procedure for multiple tests of significance. *Biometrika* **75**, 800-802.
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics* **6**, 65-70.
- Hommel, G. (1988). A stagewise rejective multiple test procedure based on a modified bonferroni test. *Biometrika*, **75**, 383-386.

- Li, H. J. and Yi, M. and Zhou, H. (2017). Generalized Holm's procedure for multiple hypotheses testing problems. *Communications in Statistics - Theory and Methods*, **46**, 7503-7510.
- Marcus, R., Peritzl, E. and Gabriel, K. (1976). On closed testing procedures with special reference to ordered analysis of variance. *Biometrika*, **63**, 655-660.
- Sonnemann, E. (2008). General solution to multiple testing problems. *Biometrical Journal*, **50**, 641-656.
- Tamhane, A. C., Liu, W. and Dunnett, C. W. (1998). A generalized step-up-down multiple test procedure. *The Canadian Journal of Statistics*, **26**, 353-363.