

INSCRIBED AND CIRCUMSCRIBED POLYGONS THAT CHARACTERIZE INNER PRODUCT SPACES

CARLOS BENÍTEZ, PEDRO MARTÍN, AND DIEGO YÁÑEZ

ABSTRACT. Let X be a real normed space with unit sphere S . We prove that X is an inner product space if and only if there exists a real number $\rho = \sqrt{(1 + \cos \frac{2k\pi}{2m+1})}/2$, ($k = 1, 2, \dots, m$; $m = 1, 2, \dots$), such that every chord of S that supports ρS touches ρS at its middle point. If this condition holds, then every point $u \in S$ is a vertex of a regular polygon that is inscribed in S and circumscribed about ρS .

1. INTRODUCTION AND NOTATION

Let X be a real normed space with unit ball B and unit sphere S . X is an inner product space (i.p.s.) if and only if every chord of S supports a sphere homothetic to S at its middle point, namely, if it fulfils the “non-bias” condition

$$u, v \in S \Rightarrow \inf_{t \in [0,1]} \|(1-t)u + tv\| = \|\frac{1}{2}u + \frac{1}{2}v\|.$$

([5]; see [1], p. 29, where this result is used to establish many characterizations of i.p.s.). But in order to characterize an i.p.s. we can only consider the chords of S that supports ρS at its middle point for some $\rho \in (0, 1)$. Namely, given the following property (P- ρS from now on)

$$u, v \in S, \inf_{t \in [0,1]} \|(1-t)u + tv\| = \rho \Rightarrow \frac{1}{2}u + \frac{1}{2}v \in \rho S, \quad (\text{P-}\rho S)$$

X is an i.p.s. if and only if (P- ρS) holds for $\rho = \frac{1}{2}$ (see [2]) or for any real number ρ such that (see [3])

$$0 < \rho < 1, \quad \rho \neq \sqrt{(1 + \cos \frac{2k\pi}{n})}/2, \quad (2k < n; n = 3, 4, \dots).$$

The aim of this paper is to prove (Theorem 11) that X is an i.p.s. if and only if (P- ρS) holds for a real number on the set

$$M = \left\{ \rho \in (0, 1) / \rho = \sqrt{(1 + \cos \frac{2k\pi}{2m+1})}/2 : k = 1, 2, \dots, m; m = 1, 2, \dots \right\}.$$

2010 *Mathematics Subject Classification.* 46B20, 46C15, 52A10, 52A21.

Key words and phrases. characterization, normed spaces, inner product.

Partially supported by Junta de Extremadura grant GR15055 (partially financed with FEDER).

It is known that X is an i.p.s. if and only if so are its 2-dimensional subspaces. This fact and the nature of the property $(P-\rho S)$ allow us to consider that X is a real 2-dimensional space from now on.

Given $u = (u_1, u_2)$ and $v = (v_1, v_2)$ in X , $[u, v]$ denotes the segment meeting u and v , and $u \prec v$ means that u precedes v in the positive orientation (counterclockwise) of X , i.e., the following expression is positive:

$$u \wedge v := u_1 v_2 - u_2 v_1.$$

We say that u is orthogonal to v in the sense of Birkhoff ([4], [6]), denoted by $u \perp v$, if

$$\|u\| \leq \|u + \lambda v\| \quad \forall \lambda \in \mathbb{R}.$$

In other words, $u \perp v$ if and only if the homothetic copy of S with scale factor $\|u\|$ is supported by the line $\{u + \lambda v : \lambda \in \mathbb{R}\}$ at u . If $(P-\rho S)$ holds, and $[u, v]$ supports ρS for some $u, v \in S$, then $u + v \perp v - u$.

If $u, v \in S$, $u \prec v$, B_u^v denotes the sector of B bounded by u, v , and the arc of S from u to v (counterclockwise); and T_u^v denotes the subset of B_u^v bounded by $[u, v]$ and the arc. If $\mathcal{A}(C)$ denotes the area of a set C , since $u \wedge v$ is the area of the parallelogram defined by u and v , then obviously $\mathcal{A}(B_u^v) = \frac{1}{2}(u \wedge v) + \mathcal{A}(T_u^v)$.

This paper is organized as follows. Some preliminary lemmas related to useful maps from $[0, 2\pi]$ to S are studied in Section 2. The concepts of ρ -ellipses and ρ -polygons, already used in [3], are reintroduced in Section 3, and some special results for $\rho \in M$ are proved in Section 4. The main Theorem 11 is obtained in Section 5, and an open problem is proposed in Section 6.

2. PRELIMINARY LEMMAS

Along this section, some essential maps from $[0, 2\pi]$ to S and their properties are introduced. The first lemma and its proof appear in [3].

Lemma 1. *Let $0 < \rho < 1$. For any $u \in S$, there is a unique $u^* \in S$, $u \prec u^*$, such that $[u, u^*]$ supports ρS . The map $u \in S \rightarrow u^* \in S$ is a homeomorphism, and $u \prec v$ implies that $u^* \prec v^*$.*

The next lemma summarizes a set of properties that are consequences of $(P-\rho S)$.

Lemma 2. *Let $0 < \rho < 1$. If X fulfils $(P-\rho S)$, then:*

- (1) X is regular (strictly convex and smooth).
- (2) For any $u \in S$, there is an unique $u^\perp \in S$, $u \prec u^\perp$, such that $u \perp u^\perp$ and the map $u \in S \rightarrow u^\perp \in S$ is a homeomorphism. If $v \in S$ and $u \prec v$, then $u^\perp \prec v^\perp$.
- (3) For any $u \in S$, there exists an unique $\mu > 0$, such that $\rho(u - \mu u^\perp)$ and $\rho(u + \mu u^\perp)$ belong to S .

Proof. Property (P- ρS) implies that $[u, u^*]$ supports ρS at $\frac{1}{2}u + \frac{1}{2}u^*$. Due to this fact and using Lemma 1, the proof of (1) and (2) presented in [2] for the case $\rho = \frac{1}{2}$ can be applied for every $0 < \rho < 1$.

Having in mind (2) and fixed $u \in S$, the convexity of the function $F : \lambda \in \mathbb{R} \rightarrow F(\lambda) = \|\rho u + \lambda \rho u^\perp\|$ implies that there exist only two real numbers $\mu, \nu \in \mathbb{R}_+$ such that $\|\rho u - \mu \rho u^\perp\| = \|\rho u + \nu \rho u^\perp\| = 1$. Applying Lemma 1 to $\rho u - \mu \rho u^\perp$ and (P- ρS), it is concluded that $\mu = \nu$. $\square$

Since S is a convex curve, the following *natural parametrization* s is continuous and of bounded variation

$$\begin{aligned} s : [0, 2\pi] &\longrightarrow S \\ \theta &\longmapsto s(\theta) = (s_1(\theta), s_2(\theta)), \end{aligned}$$

where $(s_1(\theta), s_2(\theta)) = \|(\cos \theta, \sin \theta)\|^{-1}(\cos \theta, \sin \theta)$.

And as a consequence of Lemma 1, the following parametrization (non-natural, in general) is also continuous and of bounded variation

$$s^* : \theta \in [0, 2\pi] \rightarrow s^*(\theta) := s(\theta)^* \in S.$$

Moreover, if X fulfils (P- ρS), the continuity and bounded variation hold for the parametrizations $s^\perp$, $\rho(s + \mu s^\perp)$, and $\rho(s - \mu s^\perp)$, and for the application $\mu s^\perp$ (by (2) and (3) of Lemma 2) defined as follows:

$$\begin{aligned} s^\perp : \theta \in [0, 2\pi] &\rightarrow s^\perp(\theta) := s(\theta)^\perp \in S, \\ \rho(s - \mu s^\perp) : \theta \in [0, 2\pi] &\rightarrow \rho(s(\theta) - \mu(\theta)s^\perp(\theta)) \in S, \\ \rho(s + \mu s^\perp) : \theta \in [0, 2\pi] &\rightarrow \rho(s(\theta) + \mu(\theta)s^\perp(\theta)) \in S, \\ \mu s^\perp : \theta \in [0, 2\pi] &\rightarrow \mu(\theta)s^\perp(\theta) \in X, \end{aligned}$$

where $\mu(\theta)$ is the real number considered for $u = s(\theta)$ in (3) of Lemma 2. Therefore all the Riemann-Stieltjes integrals that we shall write from now on make sense. For example, if t is any of the parametrizations of S introduced above, and $u = t(\alpha)$ and $v = t(\beta)$ ($0 \leq \alpha < \beta \leq 2\pi$), then

$$\mathcal{A}(B_u^v) = \frac{1}{2} \int_\alpha^\beta t(\theta) \wedge dt(\theta) = \frac{1}{2} \int_\alpha^\beta [t_1(\theta)dt_2(\theta) - t_2(\theta)dt_1(\theta)]. \quad (\text{A})$$

Lemma 3. Let $0 < \rho < 1$ and X fulfil (P- ρS). Let $s : \theta \in [0, 2\pi] \rightarrow s(\theta) \in S$ be a natural parametrization for S , and $s^\perp(\theta)$ and $\mu s^\perp(\theta)$ as they are defined above. Then, for any $0 \leq \alpha < \beta \leq 2\pi$:

- (1) $\int_\alpha^\beta \mu(\theta)s^\perp(\theta) \wedge ds(\theta) = 0$.
- (2) $\int_\alpha^\beta s(\theta) \wedge ds^\perp(\theta) = s(\beta) \wedge s^\perp(\beta) - s(\alpha) \wedge s^\perp(\alpha)$.
- (3) $\int_\alpha^\beta s(\theta) \wedge d[\mu(\theta)s^\perp(\theta)] = s(\beta) \wedge \mu(\beta)s^\perp(\beta) - s(\alpha) \wedge \mu(\alpha)s^\perp(\alpha)$.

Proof. Let $\alpha = \theta_0 < \theta_1 < \dots < \theta_n = \beta$ be a partition of $[\alpha, \beta]$. By the Mean Value Theorem, there exist $\eta_1, \eta_2, \dots, \eta_n \in \mathbb{R}$ such that $\theta_0 \leq \eta_1 \leq \theta_1 \leq \dots \leq \theta_{n-1} \leq \eta_n \leq \theta_n$ and $s^\perp(\eta_k) \wedge [s(\theta_k) - s(\theta_{k-1})] = 0$. Thus, the Riemann-Stieltjes sum related to this partition is equal to 0, and (1) holds.

(2) and (3) result from the integration by parts of $\int_\alpha^\beta d[s(\theta) \wedge s^\perp(\theta)]$ and of $\int_\alpha^\beta d[s(\theta) \wedge \mu s^\perp(\theta)]$, respectively, and (1). $\square$

Lemma 4. *Let $0 < \rho < 1$. If X fulfils (P- ρS), the function $u \in S \rightarrow \mathcal{A}(T_u^{u*})$ is constant.*

Proof. Let $u, v \in S$, $u \prec v$, and $s : \theta \in [0, 2\pi] \rightarrow s(\theta) \in S$ be a parametrization of S . By (3) of Lemma 2, there exist $0 \leq \alpha < \beta \leq 2\pi$ such that

$$\begin{aligned} u &= \rho[s(\alpha) - \mu(\alpha)s^\perp(\alpha)], & u^* &= \rho[s(\alpha) + \mu(\alpha)s^\perp(\alpha)] \\ v &= \rho[s(\beta) - \mu(\beta)s^\perp(\beta)], & v^* &= \rho[s(\beta) + \mu(\beta)s^\perp(\beta)]. \end{aligned}$$

By (A),

$$\begin{aligned} \mathcal{A}(B_u^v) &= \frac{\rho^2}{2} \int_\alpha^\beta [s(\theta) - \mu(\theta)s^\perp(\theta)] \wedge d[s(\theta) - \mu(\theta)s^\perp(\theta)], \\ \mathcal{A}(B_{u^*}^{v^*}) &= \frac{\rho^2}{2} \int_\alpha^\beta [s(\theta) + \mu(\theta)s^\perp(\theta)] \wedge d[s(\theta) + \mu(\theta)s^\perp(\theta)]. \end{aligned}$$

Therefore, (1) and (3) of Lemma 3 imply that

$$\begin{aligned} \mathcal{A}(B_{u^*}^{v^*}) - \mathcal{A}(B_u^v) &= \\ &= \rho^2 \int_\alpha^\beta s(\theta) \wedge d[\mu(\theta)s^\perp(\theta)] + \rho^2 \int_\alpha^\beta \mu(\theta)s^\perp(\theta) \wedge ds(\theta) = \\ &= \rho^2[s(\beta) \wedge \mu(\beta)s^\perp(\beta) - s(\alpha) \wedge \mu(\alpha)s^\perp(\alpha)] = \\ &= \frac{1}{2}(v \wedge v^*) - \frac{1}{2}(u \wedge u^*). \end{aligned}$$

If $v \prec u^*$, it is easy to check that $\mathcal{A}(B_{u^*}^{v^*}) = \mathcal{A}(B_u^v) + \mathcal{A}(B_v^{u^*})$ and $\mathcal{A}(B_v^{u^*}) = \mathcal{A}(B_v^{u^*}) + \mathcal{A}(B_{u^*}^{v^*})$. Similarly, $\mathcal{A}(B_u^v) = \mathcal{A}(B_u^{u^*}) + \mathcal{A}(B_{u^*}^{v^*})$ and $\mathcal{A}(B_{u^*}^{v^*}) = \mathcal{A}(B_{u^*}^{v^*}) + \mathcal{A}(B_v^{u^*})$ if $u^* \prec v$. In both cases, it is verified that

$$\mathcal{A}(B_{u^*}^{v^*}) - \mathcal{A}(B_u^v) = \frac{1}{2}(v \wedge v^*) - \frac{1}{2}(u \wedge u^*) + \mathcal{A}(T_v^{v^*}) - \mathcal{A}(T_u^{u^*}),$$

and it is concluded that $\mathcal{A}(T_v^{v^*}) - \mathcal{A}(T_u^{u^*}) = 0$.

□

3. ρ -ELLIPSES AND ρ -POLYGONS

Fixed $0 < \rho < 1$ and $u \in S$, the ρ -**polygon** associated to u is the set of ordered points $P_u = \{u_1, u_2, u_3, \dots\}$ of S and the segments $[u_i, u_{i+1}]$, such that $u = u_1 \prec u_2 \prec u_3, \dots$, and every segment $[u_i, u_{i+1}]$ supports ρS , i.e., $u_{i+1} = u_i^*$. Each $u_i \in P_u$ is a *vertex* of P_u and each $[u_i, u_{i+1}]$ is a *side* of P_u . The ρ -**ellipse** C_u associated to $u \in S$ is the unique ellipse centered at $(0, 0)$ that contains the points u , $\frac{1}{2\rho}(u + u^*)$, and u^* (see Figure 1).

Some properties and examples of ρ -polygons and ρ -ellipses are presented in [3]. For instance, if S is an ellipse and $\rho = \sqrt{(1 + \cos \frac{2k\pi}{n})/2}$ with $\frac{k}{n}$ irreducible, it is proved that P_u is convex with n vertices for $k = 1$ and $n = 3, 4, \dots$; P_u is star-shaped with n vertices for either $k = 2, \dots, \frac{n}{2} - 1$ when n is even, or for $k = 2, \dots, \frac{n-1}{2}$ when n is odd; P_u is dense if ρ is not in any of the previous cases. On the other hand, if S is the unit sphere of ℓ_∞^2

(with vertices $\{(\pm 1, \pm 1)\}$) and $\rho = \frac{1}{2}$, then $P_{(1,0)} = \{(0, \pm 1), (\pm 1, 0)\}$; and if $u \notin \{(0, \pm 1), (\pm 1, 0)\}$, then P_u has infinite vertices, but it is not dense in S (the points $\{(0, \pm 1), (\pm 1, 0)\}$ are the unique points of accumulation of P_u). And the same happens for l_p^2 ($p > 2$) and $\rho = (\frac{1}{2})^{\frac{p-1}{p}}$.

The next lemma presents some properties of ρ -polygons.

Lemma 5. *Let $0 < \rho < 1$. Then:*

- (1) *If there exists $u \in S$ such that P_u is dense in S , then P_v is dense in S for any $v \in S$.*
- (2) *If there exist $u, v \in S$ such that P_u and P_v have a finite number of points, then both have the same number of vertices, and either both are convex or both are star-shaped. Besides, in this last case the number of vertices that are (geometrically situated on S) between u_i and u_{i+1} is equal to the number of vertices between v_i and v_{i+1} .*
- (3) *If $P_u = \{u_1, u_2, u_3, \dots\}$, then $-P_u = \{-u_1, -u_2, -u_3, \dots\} = P_{-u}$.*
- (4) *If P_u has an odd number of vertices, then $P_u \cap P_{-u} = \emptyset$.*

Proof. The statement (1) is proved in Lemma 2.2 of [3].

In order to prove (2), let us consider $P_u = \{u_1, u_2, \dots, u_n\}$ and $P_v = \{v_1, v_2, \dots, v_m\}$ with $m > n$. Let us assume without loss of generality that $u_1 \prec v_1 \prec u_p$, and there is not any other vertex of P_u between u_1 and u_p ($p < n$). By Lemma 1,

$$u_2 \prec v_2 \prec u_{p+1}, \quad \dots, \quad u_n \prec v_n \prec u_{p+n-1}, \quad u_1 \prec v_{n+1} \prec u_p, \quad \dots,$$

and it is concluded that $m = kn$, $k \in \mathbb{N}$. Thus, $u_1 \prec v_1 \prec u_p$ and $u_1 \prec v_{n+1} \prec u_p$. But if $v_1 \prec v_{n+1}$, then $v_1 \prec v_{kn+1}$ ($\forall k \in \mathbb{N}$) by Lemma 1, and this is a contradiction because $v_1 = v_{m+1} = v_{kn+1}$. The analysis is similar if $v_{n+1} \prec v_1$.

Let us assume now that P_u is convex, P_v is star-shaped, and each set has n vertices. Then, $u_1 \prec v_1 \prec u_2 \prec v_2 \prec u_3 \prec \dots \prec u_n \prec v_n \prec u_1$, leads to a contradiction.

For the last assertion of (2), it is enough to consider that $u_i \prec u_m \prec u_{i+1}$ implies (Lemma 1) $u_{i+1} \prec u_{m+1} \prec u_{i+2}$.

The equality $-P_u = P_{-u}$ of (3) is a consequence of the symmetry of S .

Let us see (4). Let us assume that P_u has an odd number of vertices and that there exist $-u_p \in -P_u$ and $u_q \in P_u$ such that $-u_p = u_q$. Then, $(-u_p)^* = -u_{p+1}$ (by symmetry of S) and $(-u_p)^* = u_{q+1}$ (by the construction of P_u). I.e., $-u_{p+1} = u_{q+1}$, and, in general, $-u_{p+k} = u_{q+k}$, ($k \in \mathbb{N}$). Consequently, P_u is symmetric, and this is not possible because it has an odd number of vertices. $\square$

Lemma 6. *Let $u, v \in S$ such that $P_u = \{u_1, u_2, \dots, u_n\}$ and $P_v = \{v_1, v_2, \dots, v_n\}$ for some $0 < \rho < 1$. If X fulfils $(P-\rho S)$, then*

$$u_1 \wedge u_2 + \dots + u_{n-1} \wedge u_n + u_n \wedge u_1 = v_1 \wedge v_2 + \dots + v_{n-1} \wedge v_n + v_n \wedge v_1.$$

Proof. Let r be the constant number of vertices of P_u that are (geometrically situated) between u_i and u_{i+1} (or, equivalently, between v_i and v_{i+1} by (2) of

Lemma 5). The statement is a consequence of Lemma 4 and these equalities

$$\begin{aligned} (r+1)\mathcal{A}(B) &= \mathcal{A}(B_{u_1}^{u_2}) + \dots + \mathcal{A}(B_{u_n}^{u_1}) = \\ &= \frac{1}{2}(u_1 \wedge u_2) + \mathcal{A}(T_{u_1}^{u_2}) + \dots + \frac{1}{2}(u_n \wedge u_1) + \mathcal{A}(T_{u_n}^{u_1}), \\ (r+1)\mathcal{A}(B) &= \mathcal{A}(B_{v_1}^{v_2}) + \dots + \mathcal{A}(B_{v_n}^{v_1}) = \\ &= \frac{1}{2}(v_1 \wedge v_2) + \mathcal{A}(T_{v_1}^{v_2}) + \dots + \frac{1}{2}(v_n \wedge v_1) + \mathcal{A}(T_{v_n}^{v_1}). \end{aligned}$$

□

The following result presents some properties about ρ -ellipses and spheres that are *tangent*. It is said that C_u and S are *tangent* at $v \in S \cap C_u$ if both curves have the same supporting line at v . If X fulfils (P- ρS), then C_u and S are tangent at u if and only if the common supporting line at u is

$$\{u + \lambda[(1 - 2\rho^2)u + u^*] : \lambda \in \mathbb{R}\},$$

that is, if and only if u has the following property (see Figure 1)

$$u \perp (1 - 2\rho^2)u + u^*. \quad (*)$$

Likewise, C_u and S are tangent at $u^* \in S \cap C_u$ if and only if u^* verifies

$$u^* \perp -u - (1 - 2\rho^2)u^*. \quad (**)$$




FIGURE 1. ρ -ellipses C_u associated to $u \in S$. On the right, S and C_u are tangent at u , $\frac{u+u^*}{2\rho}$, and u^* .

Lemma 7. Let $0 < \rho < 1$. If X fulfils (P- ρS), then:

- (1) C_u and S are tangent at $\frac{1}{2\rho}(u + u^*)$ for every $u \in S$.
- (2) If C_u and S are tangent at $u \in S$ (equivalently, if u verifies $(*)$), then C_u and S are tangent at every point of $P_u \cup P_w$, where $w = \frac{1}{2\rho}(u + u^*)$.
- (3) There exists $v \in S$ such that C_v and S are tangent at v .

Proof. Lemma 3.2 in [3] proves (1), (3), and that if C_u and S are tangent at u , then C_u and S are tangent at u^* . The proof of Lemma 3.3 in [3] can be applied for every $u \in S$ such that C_u and S are tangent at u , and (2) holds. $\square$

4. ρ -ELLIPSES AND ρ -POLYGONS: THE SPECIAL CASE $\rho \in M$

We remind the reader the definition of M

$$M = \left\{ \rho \in (0, 1) / \rho = \sqrt{(1 + \cos \frac{2k\pi}{2m+1})/2} : k = 1, 2, \dots, m; m = 1, 2, \dots \right\}.$$

Lemma 8. *Let $\rho \in M$. If X fulfils (P- ρS), the following properties hold for every $v \in S$ such that C_v and S are tangent at v (equivalently, for every v that verifies (*)):*

- (1) $P_v = \{v_1, v_2, \dots, v_n\}$ has $n = 2m + 1$ vertices, and C_v and S are tangent at every $v_i \in P_v$.
- (2) If $w = \frac{1}{2\rho}(v + v^*)$, P_w has $n = 2m + 1$ vertices. Such as vertices are the points $w_i = \frac{1}{2\rho}(v_i + v_{i+1})$, and C_v and S are tangent at every $w_i \in P_w$.
- (3) If k is an odd number, $P_w = P_{-v}$. If k is an even number, $P_w = P_v$.
- (4) $v_1 \wedge v_2 = \dots = v_{n-1} \wedge v_n = v_n \wedge v_1$.
- (5) $\mathcal{A}(B_{v_1}^{v_2}) = \dots = \mathcal{A}(B_{v_{n-1}}^{v_n}) = \mathcal{A}(B_{v_n}^{v_1})$.
- (6) The vertices of P_v and P_{-v} split B into $2n$ disjoint sectors of equal area.

Proof. Lema 7 ensures the existence of $v \in S$ such that C_v and S are tangent at every vertex of $P_v \cup P_w$. Since C_v is an ellipse, P_v has $n = 2m + 1$ vertices for $\rho \in M$ (see Example 1 in [3] or the comments at the beginning of Section 3) and (1) holds, as well as (2).

It is easy to see that (3) and (4) are true when S is an ellipse (see Figure 2). But in the general case, the vertices of P_v and P_w are always the vertices of ρ -polygons inscribed in the ρ -ellipse C_v and circumscribed about its homothetic ellipse of ratio ρ (Lemma 7). Therefore, (3) and (4) hold for every S .

From (4) and Lemma 4, (5) is obtained.

Let us see (6). Since $n = 2m + 1$ is an odd number, then (see (4) of Lemma 5), the vertices of $P_v \cup P_{-v}$ determine $2n$ different vectors in $S \cap C_v$: $\{v_1, \dots, v_n, -v_1, \dots, -v_n\}$. For every $v_i \in P_v$, let $-v_{\sigma(i)} \in P_{-v}$ be such that $v_i \prec -v_{\sigma(i)}$ and there is not any other vertex of $P_v \cup P_{-v}$ between (counterclockwise) v_i and $-v_{\sigma(i)}$ (see Figure 2).

Due to the symmetry of S and (5), $\mathcal{A}(B_{v_i}^{v_{i+1}}) = \mathcal{A}(B_{-v_{\sigma(i)}}^{-v_{\sigma(i+1)}})$. Using arguments similar to those in Lemma 4, it holds that:

$$\begin{aligned} \mathcal{A}(B_{v_i}^{v_{i+1}}) &= \mathcal{A}(B_{v_i}^{-v_{\sigma(i)}}) + \mathcal{A}(B_{-v_{\sigma(i)}}^{v_{i+1}}), \\ \mathcal{A}(B_{-v_{\sigma(i)}}^{-v_{\sigma(i+1)}}) &= \mathcal{A}(B_{-v_{\sigma(i)}}^{v_{i+1}}) + \mathcal{A}(B_{v_{i+1}}^{-v_{\sigma(i+1)}}). \end{aligned}$$

Hence, the n disjoint sectors $B_{v_i}^{-v_{\sigma(i)}}$ have the same area. Since n is an odd number (and again the symmetry of S), (6) holds. $\square$




FIGURE 2. ρ -polygon P_v with $n = 7$ and $k = 1$ (left), $k = 2$ (center), and $k = 3$ (right).

Lemma 9. *Let $\rho \in M$. If X fulfils (P- ρS), then for every $u \in S$ it is verified that $P_u = \{u_1, u_2, u_3, \dots\}$ has $n = 2m + 1$ vertices and C_u supports S at every vertex of P_u . Besides, the function $u \in S \rightarrow u \wedge u^*$ is constant, and $\mathcal{A}(B_u^v) = \mathcal{A}(B_u^{v^*})$ for any $v \in S$ such that $u \prec v$.*

Proof. We remind that if X fulfils (P- ρS) and $u \in S$, then C_u and S are tangent at u if and only if u has property (*). The proof is organized in four steps.

Step 1: there exists $v \in S$ such that the condition () is verified for every $z \in P_v \cup P_{-v}$.*

Since S and the ρ -ellipses are symmetric, it is deduced (by Lemma 7, Lemma 8, and (3) and (4) of Lemma 5) that there exists $v \in S$ such that the condition (*) is verified for every $z \in P_v \cup P_{-v}$.

Let $P_v = \{v_1, v_2, \dots, v_n\}$ be the polygon generated by v . As in Lemma 8, let us denote $-v_{\sigma(i)}$ to the unique point in P_{-v} such that $v_i \prec -v_{\sigma(i)}$ and there is not any other vertex of $P_v \cup P_{-v}$ between (counterclockwise) v_i and $-v_{\sigma(i)}$.

Step 2: for every $i \in \{1, 2, \dots, n\}$, there exists $\bar{v} \in S$ such that $\bar{v}$ verifies () and $v_i \prec \bar{v} \prec -v_{\sigma(i)}$.*

Without loss of generality, let us assume that $i = 1$. Let us consider a parametrization $s : \theta \in [0, 2\pi] \rightarrow s(\theta) \in S$ of S . By (3) of Lemma 2, there exist $0 \leq \theta_1 < \theta'_1 \leq 2\pi$ such that

$$\begin{aligned} v_1 &= \rho[s(\theta_1) - \mu(\theta_1)s^\perp(\theta_1)], & -v_{\sigma(1)} &= \rho[s(\theta'_1) - \mu(\theta'_1)s^\perp(\theta'_1)], \\ v_2 &= \rho[s(\theta_1) + \mu(\theta_1)s^\perp(\theta_1)], & -v_{\sigma(2)} &= \rho[s(\theta'_1) + \mu(\theta'_1)s^\perp(\theta'_1)]. \end{aligned}$$

It is proved just some lines below that

$$\int_{\theta_1}^{\theta'_1} [(1 - \rho^2)s(\theta) + \rho^2\mu(\theta)s^\perp(\theta)] \wedge d[s(\theta) - \mu(\theta)s^\perp(\theta)] = 0, \quad (1)$$

and as a consequence, there exists $\theta_1 < \bar{\theta} < \theta'_1$ such that

$$s(\bar{\theta}) - \mu(\bar{\theta})s^\perp(\bar{\theta}) \perp (1 - \rho^2)s(\bar{\theta}) + \rho^2\mu(\bar{\theta})s^\perp(\bar{\theta}).$$

Thus the points

$$\bar{v} := \rho[s(\bar{\theta}) - \mu(\bar{\theta})s^\perp(\bar{\theta})], \quad \bar{v}^* := \rho[s(\bar{\theta}) + \mu(\bar{\theta})s^\perp(\bar{\theta})],$$

verify $\bar{v} \perp (1 - 2\rho^2)\bar{v} + \bar{v}^*$, with $v_1 \prec \bar{v} \prec -v_{\sigma(1)}$, as Step 2 claims.

In order to see (1), the integral is separated into four parts as follows:

$$\begin{aligned} \int_{\theta_1}^{\theta'_1} [(1 - \rho^2)s(\theta) + \rho^2\mu(\theta)s^\perp(\theta)] \wedge d[s(\theta) - \mu(\theta)s^\perp(\theta)] = \\ (1 - \rho^2) \int_{\theta_1}^{\theta'_1} s(\theta) \wedge ds(\theta) - (1 - \rho^2) \int_{\theta_1}^{\theta'_1} s(\theta) \wedge d[\mu(\theta)s^\perp(\theta)] \\ + \rho^2 \int_{\theta_1}^{\theta'_1} \mu(\theta)s^\perp(\theta) \wedge ds(\theta) - \rho^2 \int_{\theta_1}^{\theta'_1} \mu(\theta)s^\perp(\theta) \wedge d[\mu(\theta)s^\perp(\theta)]. \end{aligned}$$

Let us denote w_1 and $-w_{\sigma(1)}$, respectively, to $\frac{1}{2\rho}(v_1 + v_2)$ and $-\frac{1}{2\rho}(v_{\sigma(1)} + v_{\sigma(2)})$. Then $w_1 = s(\theta_1)$ and $-w_{\sigma(1)} = s(\theta'_1)$. By the calculus of area (see (A)), the first part is

$$(1 - \rho^2) \int_{\theta_1}^{\theta'_1} s(\theta) \wedge ds(\theta) = 2(1 - \rho^2) \mathcal{A}(B_{w_1}^{-w_{\sigma(1)}}).$$

By (3) of Lemma 3 and (4) of Lemma 8, the second part is

$$\begin{aligned} \int_{\theta_1}^{\theta'_1} s(\theta) \wedge d[\mu(\theta)s^\perp(\theta)] = \\ s(\theta'_1) \wedge \mu(\theta'_1)s^\perp(\theta'_1) - s(\theta_1) \wedge \mu(\theta_1)s^\perp(\theta_1) = \\ \frac{1}{2\rho^2} [(-v_{\sigma(1)} \wedge -v_{\sigma(2)}) - (v_1 \wedge v_2)] = 0. \end{aligned}$$

And by (1) of Lemma 3, the third part is

$$\rho^2 \int_{\theta_1}^{\theta'_1} \mu(\theta)s^\perp(\theta) \wedge ds(\theta) = 0.$$

Regarding the last part of the decomposition, using again the calculus of the area (A), and the same statements of Lemma 3 and Lemma 8, it is

deduced that

$$\begin{aligned}
\mathcal{A}(B_{v_1}^{-v_{\sigma(1)}}) &= \\
&\frac{\rho^2}{2} \int_{\theta_1}^{\theta'_1} [s(\theta) - \mu(\theta)s^\perp(\theta)] \wedge d[s(\theta) - \mu(\theta)s^\perp(\theta)] = \\
&\frac{\rho^2}{2} \int_{\theta_1}^{\theta'_1} s(\theta) \wedge ds(\theta) - \frac{\rho^2}{2} \int_{\theta_1}^{\theta'_1} s(\theta) \wedge d[\mu(\theta)s^\perp(\theta)] \\
&- \frac{\rho^2}{2} \int_{\theta_1}^{\theta'_1} \mu(\theta)s^\perp(\theta) \wedge ds(\theta) + \frac{\rho^2}{2} \int_{\theta_1}^{\theta'_1} \mu(\theta)s^\perp(\theta) \wedge d[\mu(\theta)s^\perp(\theta)] = \\
&\rho^2 \mathcal{A}(B_{w_1}^{-w_{\sigma(1)}}) + \frac{\rho^2}{2} \int_{\theta_1}^{\theta'_1} \mu(\theta)s^\perp(\theta) \wedge d[\mu(\theta)s^\perp(\theta)],
\end{aligned}$$

and concluded that

$$\rho^2 \int_{\theta_i}^{\theta'_i} \mu(\theta)s^\perp(\theta) \wedge d[\mu(\theta)s^\perp(\theta)] = 2\mathcal{A}(B_{v_1}^{-v_{\sigma(1)}}) - 2\rho^2 \mathcal{A}(B_{w_1}^{-w_{\sigma(1)}}).$$

Since $\mathcal{A}(B_{v_1}^{-v_{\sigma(i)}}) = \mathcal{A}(B_{w_1}^{-w_{\sigma(1)}})$ by (3) and (6) of Lemma 8, then

$$\rho^2 \int_{\theta_i}^{\theta'_i} \mu(\theta)s^\perp(\theta) \wedge d[\mu(\theta)s^\perp(\theta)] = 2(1 - \rho^2)\mathcal{A}(B_{w_1}^{-w_{\sigma(1)}}),$$

and the equality (1) holds.

Step 3: If $\bar{v}$ verifies $()$ and $v_i \prec \bar{v} \prec -v_{\sigma(i)}$, then there exist $\bar{v}', \bar{v}'' \in S$ such that both $\bar{v}'$ and $\bar{v}''$ verify $(*)$, and $v_i \prec \bar{v}' \prec \bar{v} \prec \bar{v}'' \prec -v_{\sigma(i)}$.*

Let us prove the existence of $\bar{v}'$ (the existence of $\bar{v}''$ can be proved similarly). Only for simplicity, let us assume that $i = 1$. Let us consider

$$\begin{aligned}
\bar{v}_1 &= \bar{v}, & \bar{v}_{j+1} &= \bar{v}_j^* & P_{\bar{v}} &= \{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n\} \\
\bar{w}_j &= \frac{1}{2\rho}(\bar{v}_j + \bar{v}_{j+1}) & \bar{w}_{j+1} &= \bar{w}_j^* & P_{\bar{w}} &= \{\bar{w}_1, \bar{w}_2, \dots, \bar{w}_n\}
\end{aligned}$$

Since $\bar{v}$ verifies $(*)$, then ((2) of Lemma 7) $C_{\bar{v}}$ and S are tangent at every point of $P_{\bar{v}} \cup P_{\bar{w}}$. Moreover:

- (a) $\bar{v}_j \wedge \bar{v}_{j+1} = v_i \wedge v_{i+1}$ (by (4) of Lemma 8 and Lemma 6).
- (b) $\mathcal{A}(B_{v_i \bar{v}_i})$ is constant (by (a) and Lemma 4).
- (c) $\mathcal{A}(B_{v_i \bar{v}_i}) = \mathcal{A}(B_{w_i \bar{w}_i})$ (by (3) of Lemma 8).

Using the above properties, the existence of $\bar{v}'$ of this Step 3 (such that $v_i \prec \bar{v}' \prec \bar{v}$) can be proved applying the arguments of Step 2 to v_i and $\bar{v}$ (instead to v_i and $-v_{\sigma(i)}$).

Step 4: The set of points of S that verify $()$ is dense on S .*

Let x, y be a pair of points of S that verify $(*)$. Step 3 can be applied to x and y (instead of v_i and $\bar{v}$) because x and y have the required properties: both verify property $(*)$ and the conditions for x and y equivalent to (a), (b), and (c) remind true (by the same reasons). Therefore, there exists $z \in S$ such that z verifies $(*)$ and $x \prec z \prec y$.

As a consequence of Step 4, the statements of Lemma 9 are true for every $u \in S$. Particularly, because conditions similar to (a) and to (b) are verified for any pair of points of S (instead of v_i and $\bar{v}_i$), then $u \wedge u^* = v \wedge v^*$ and $\mathcal{A}(B_u^v) = \mathcal{A}(B_{u^*}^{v^*})$ for any $u, v \in S$ such that $u \prec v$. $\square$

5. MAIN RESULT

The last lemma describes some properties of a natural parametrization s and the related parametrization s^* .

Lemma 10. *Let $\rho \in M$ and $s : [0, 2\pi] \rightarrow S$ be a natural parametrization for S . If X fulfils (P- ρS), then:*

- (i) s is continuously differentiable and there is a continuous function $p : [0, 2\pi] \rightarrow \mathbb{R}_+$ such that $s'(\theta) = p(\theta)s^\perp(\theta)$.
- (ii) s^* is continuously differentiable and there is a continuous function $q : [0, 2\pi] \rightarrow \mathbb{R}_+$ such that $s^{*\prime}(\theta) = q(\theta)s^{*\perp}(\theta)$.

Proof. The following conditions holds: X is smooth (by Lemma 2); the function $u \in S \rightarrow u \wedge u^*$ is constant (by Lemma 9); and s, s^* , and $s^\perp$ are continuous functions (see Section 2). Besides, $s'(\theta) \wedge s^*(\theta) \neq 0$ as a consequence of $s(\theta) \perp (1 - 2\rho^2)s(\theta) + s^*(\theta)$ for every $\theta \in [0, 2\pi]$ (by Lemma 9 and because $1 - 2\rho^2 \neq 0$). Therefore, the proof of the statement for the case $\rho = \frac{1}{2}$ (Lemma 2.8 in [2]) can be rewritten for $\rho \in M$. $\square$

And finally, the main result is presented.

Theorem 11. *Given the set*

$$M = \left\{ \rho \in (0, 1) / \rho = \sqrt{(1 + \cos \frac{2k\pi}{2m+1})}/2 : k = 1, 2, \dots, m; m = 1, 2, \dots \right\},$$

a real normed space X is an i.p.s. if and only if there exists $\rho \in M$ such that X fulfils

$$u, v \in S, \inf_{t \in [0, 1]} \|tu + (1-t)v\| = \rho \Rightarrow \frac{1}{2}u + \frac{1}{2}v \in \rho S. \quad (\text{P-}\rho S)$$

Proof. Let X be an i.p.s. such that the scalar product of $u, v \in X$ is $(u|v)$. It is easy to see that for any $u, v \in S$, $u \prec v$, the convex function

$$F(t) = \|(1-t)u + tv\|^2 = 1 - 2t + 2t^2 + 2t(1-t)(u|v)$$

attains its minimum at $t = \frac{1}{2}$ when $(u|v) < 1$. Thus, X fulfils (P- ρS) for every $\rho \in (0, 1)$.

In order to prove the converse, let us fixed a natural parametrization $s : [0, 2\pi] \rightarrow S$ for S . The following conditions holds:

- (1) X is smooth (Lemma 2).
- (2) If $u, v \in S$ such that $u \prec v$, then $u \wedge u^* = v \wedge v^*$; $\mathcal{A}(B_u^v) = \mathcal{A}(B_{u^*}^{v^*})$; $u \perp (1 - 2\rho^2)u + u^*$; and $u^* \perp -u - (1 - 2\rho^2)u^*$ (Lemma 9, property (*) for u , and property (**) for u^*).

- (3) There exist some continuous functions $p, q : [0, 2\pi] \rightarrow \mathbb{R}_+$ such that $s'(\theta) = p(\theta)s^\perp(\theta)$ and $s^{*'}(\theta) = q(\theta)s^{*\perp}(\theta)$ (Lemma 10).

Using (1), (2), and (3), the proof of the statement for the case $\rho = \frac{1}{2}$ (Theorem 3.1 in [2]) can be rewritten for $\rho \in M$ with only very slight and not significant changes. For example, the (non restrictive) initial data $s(0) = (1, 0)$ and $s^*(0) = (-\frac{1}{2}, \frac{\sqrt{3}}{2})$ considered for $\rho = \frac{1}{2}$ would be replaced by $s(0) = (1, 0)$ and $s^*(0) = (\cos \frac{2k\pi}{2m+1}, \sin \frac{2k\pi}{2m+1})$. $\square$

6. CONCLUSION AND OPEN PROBLEM

We conjecture that a real normed space X is an i.p.s. if and only if there exists $0 < \rho < 1$ such that X fulfils property (P- ρS). If $2k < n$ and $n = 3, 4, \dots$, the case $\rho \neq \sqrt{(1 + \cos \frac{2k\pi}{n})}/2$ is proved in [3]. The case $\rho = \sqrt{(1 + \cos \frac{2k\pi}{n})}/2$ is proved in this paper when n is odd (for $n = 3$, $\rho = \frac{1}{2}$, it was solved previously in [2]), but it is left open when n is even. For this unsolved situation, the results of Section 2 and Section 3, as well as some assertions of Lemma 8 ((4), (5); also (1) and (2) considering $n = 2m$) remain true. Besides, regarding (3) of Lemma 8, it is easy to see that $P_v = P_{-v}$ and $P_v \cap P_w = \emptyset$ when n is even. Nevertheless, the authors are not able to prove that the vertices of P_v and P_w split B into $2n$ disjoint sectors of equal area, which would be the property equivalent to (6) of Lemma 8.

ACKNOWLEDGEMENT

The second and third authors are very grateful to the first author for his work along more than 30 years in the University of Extremadura and, in general, for the example of his life. This paper is one of the last mathematical contributions of professor Carlos Benítez, who died on March 7th, 2014.

REFERENCES

- [1] D. AMIR, *Characterization of Inner Product Spaces* (Birkhauser Verlag, Basel, 1986).
- [2] C. BENÍTEZ AND D. YÁÑEZ, ‘Middle points, medians, and inner products’, *Proc. Amer. Math. Soc.* **135** (2007), 1725–1734.
- [3] C. BENÍTEZ AND D. YÁÑEZ, ‘Middle points and inner products’, *Bull. London Math. Soc.* **39** (2007), 811–817.
- [4] G. BIRKHOFF, ‘Orthogonality in linear metric spaces’, *Duke Math. J.* **1** (1935) 169–172.
- [5] N.I. GURARI AND Y.I. SOZONOV, ‘On normed spaces which have no bias of the unit sphere’, *Math. Notes* **7** (1970) 187–189.
- [6] R.C. JAMES, ‘Orthogonality and linear functionals in normed linear spaces’, *Trans. Amer. Math. Soc.* **1** (1947), 265–292.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA. 06071 BADAJOZ, SPAIN

E-mail address: † on March 7th 2014, pjimene@unex.es, dyanez@unex.es