

Some variations of EM algorithms for Marshall-Olkin bivariate Pareto distribution with location and scale

Arabin Kumar Dey · Biplab Paul

the date of receipt and acceptance should be inserted later

Abstract Recently Asimit et. al ([Asimit et al. \(2016\)](#)) used an EM algorithm to estimate Marshall-Olkin bivariate Pareto distribution. The distribution has seven parameters. We describe few alternative approaches of EM algorithm. A numerical simulation is performed to verify the performance of different proposed algorithms. A real-life data analysis is also shown for illustrative purposes.

Keywords Joint probability density function, Bivariate Pareto distribution, EM algorithm, Pseudo likelihood function

1 Introduction

Bivariate Pareto has several forms. In this paper we study Marshal-Olkin formulation ([Marshall and Olkin \(1967\)](#)) of this distribution which includes both location and scale parameters. In a very recent paper, Asimit et al. ([Asimit et al. \(2010\)](#)) used EM algorithm to estimate the parameters of this distribution. We adapt few more variations of the same. We observe that some of our proposed algorithms either works equally well or outperforms their algorithm. We also present some real life data analysis which is absent in their paper.

A. K. Dey
Department of Mathematics,
IIT Guwahati,
Guwahati, India
Assam
Tel.: +91361-258-4620
E-mail: arabin@iitg.ac.in

B. Paul
Department of Mathematics,
IIT Guwahati,
Guwahati, India
E-mail: biplab.paul@iitg.ac.in

In this paper our focus is on **singular** bivariate Pareto distribution whose both the marginals have Pareto type-II distribution. We can obtain bivariate Pareto distribution considering peak over threshold method in a bivariate data. The distribution has wide application in modeling data related to finance, insurance, environmental sciences and internet network. Any analysis based on this distribution requires efficient techniques for estimating parameters of the distribution. We consider a more generalized set up including location and scale parameters in formation of Marshal-Olkin bivariate Pareto distribution. This dependence structure can be described by well-known Marshall-Olkin copula too [Nelsen (2007), Marshall and Olkin (1967), Yeh (2000), Yeh (2004)]. We propose few innovative ways to implement EM algorithm.

A random variable X is said to have Pareto of second kind, i.e. $X \sim Pa(II)(\mu, \sigma, \alpha)$ if it has the survival function

$$\bar{F}_X(x; \mu, \sigma, \alpha) = P(X > x) = (1 + \frac{x - \mu}{\sigma})^{-\alpha}$$

and the probability density function (pdf)

$$f(x; \mu, \sigma, \alpha) = \frac{\alpha}{\sigma} (1 + \frac{x - \mu}{\sigma})^{-\alpha-1}$$

with $x > \mu \in \mathcal{R}$, $\sigma > 0$ and $\alpha > 0$.

Kundu and Gupta (2009), Kundu and Gupta (2010), Kundu and Dey (2009) used EM algorithm for estimating parameters of different bivariate distributions, e.g. bivariate generalized exponential, bivariate Weibull etc. Dey and Kundu (2012) performed discrimination between bivariate Weibull and bivariate generalized exponential distributions where they used parameter estimation through EM algorithm. Sarhan and Balakrishnan (2007) considered estimation issues in Sarhan and Balakrishnan bivariate distribution with extra scale parameter in their model. Many works have been done on multivariate Pareto distribution too [Hanagal (1996), Yeh (2000), Yeh (2004), Asimit et al. (2010)]. Statistical Inference of multivariate Pareto distribution through EM algorithm is attempted by Asimit et al. (2016). This paper also deals with seven parameters including location and scale as its parameters. We handle the same problem in a slightly different manner.

We arrange the paper in the following way. In section 2 we keep the Marshall-Olkin bivariate Pareto formulation and some of its properties. In section 3, we describe our proposed EM algorithms. Some simulation results show the performance of the algorithm in section 4. In section 5 we show the data analysis. Finally we conclude the paper in section 6.

2 Formulation of Marshal-Olkin bivariate Pareto

Let U_0 , U_1 and U_2 be three independent univariate type-II Pareto distributions $Pa(II)(0, 1, \alpha_0)$, $Pa(II)(\mu_1, \sigma_1, \alpha_1)$ and $Pa(II)(\mu_2, \sigma_2, \alpha_2)$.

We define $X_1 = \min\{\sigma_1 U_0 + \mu_1, U_1\}$ and $X_2 = \min\{\sigma_2 U_0 + \mu_2, U_2\}$. We can show that (X_1, X_2) jointly follow bivariate Pareto distribution of second kind, we call it as $BVPA(\mu_1, \mu_2, \sigma_1, \sigma_2, \alpha_0, \alpha_1, \alpha_2)$

The joint distribution can be given by

$$f(x_1, x_2) = \begin{cases} f_1(x_1, x_2) & \text{if } \frac{x_1 - \mu_1}{\sigma_1} < \frac{x_2 - \mu_2}{\sigma_2} \\ f_2(x_1, x_2) & \text{if } \frac{x_1 - \mu_1}{\sigma_1} > \frac{x_2 - \mu_2}{\sigma_2} \\ f_0(x) & \text{if } \frac{x_1 - \mu_1}{\sigma_1} = \frac{x_2 - \mu_2}{\sigma_2} = x \end{cases}$$

where

$$\begin{aligned} f_1(x_1, x_2) &= \frac{\alpha_1}{\sigma_1 \sigma_2} (\alpha_0 + \alpha_2) \left(1 + \frac{x_2 - \mu_2}{\sigma_2}\right)^{-(\alpha_0 + \alpha_2 + 1)} \left(1 + \frac{x_1 - \mu_1}{\sigma_1}\right)^{-(\alpha_1 + 1)} \\ f_2(x_1, x_2) &= \frac{\alpha_2}{\sigma_1 \sigma_2} (\alpha_0 + \alpha_1) \left(1 + \frac{x_1 - \mu_1}{\sigma_1}\right)^{-(\alpha_0 + \alpha_1 + 1)} \left(1 + \frac{x_2 - \mu_2}{\sigma_2}\right)^{-(\alpha_2 + 1)} \\ f_0(x) &= \alpha_0 (1 + x)^{-(\alpha_1 + \alpha_2 + \alpha_0 + 1)} \end{aligned}$$

It can be shown that

1. The distribution of X_j is $Pa(II)(\mu_j, \sigma_j, \alpha_{0j})$, $\alpha_{0j} = \alpha_0 + \alpha_j$.
2. Distribution of the minimum is also Pareto distribution when U_0 , U_1 and U_2 has same location and scale parameter.
3. Maximum Likelihood estimate of μ , σ , α , the parameters of univariate Pareto can be given based on the data set Y_1, Y_2, \dots, Y_n as : $\mu = \min\{Y_1, Y_2, \dots, Y_n\}$, whereas estimates of α and σ can be obtained by solving the fixed point iterations :

$$\sigma = \frac{\alpha + 1}{n} \sum_{i=1}^n \frac{(x_i - \mu)}{(1 + \frac{x_i - \mu}{\sigma})}$$

where,

$$\alpha = \frac{n}{\sum_{i=1}^n \ln(1 + \frac{x_i - \mu}{\sigma})}$$

Surface and contour plots of the absolutely continuous part of the pdf are shown in Figure-1 and Figure-2 respectively. The following four different sets of parameters provide four subfigures in each figure. $\xi_1 : \mu_1 = 0, \mu_2 = 0, \sigma_1 = 1, \sigma_2 = 0.5, \alpha_0 = 1, \alpha_1 = 0.3, \alpha_2 = 1.4$; $\xi_2 : \mu_1 = 1, \mu_2 = 2, \sigma_1 = 0.4, \sigma_2 = 0.5, \alpha_0 = 2, \alpha_1 = 1.2, \alpha_2 = 1.4$; $\xi_3 : \mu_1 = 0, \mu_2 = 0, \sigma_1 = 1.4, \sigma_2 = 0.5, \alpha_0 = 1, \alpha_1 = 1, \alpha_2 = 1.4$; $\xi_4 : \mu_1 = 0, \mu_2 = 0, \sigma_1 = 1.4, \sigma_2 = 0.5, \alpha_0 = 2, \alpha_1 = 0.4, \alpha_2 = 0.5$.




Fig. 1 Surface plots for pdf of BVPA




Fig. 2 Contour plots for pdf of BVPA

3 EM-algorithm

Let us assume μ_1 , μ_2 , σ_1 and σ_2 are known. Now we divide our data into three parts :-

$$I_0 = \left\{ (x_{1i}, x_{2i}) : \frac{x_{1i} - \mu_1}{\sigma_1} = \frac{x_{2i} - \mu_2}{\sigma_2} \right\}, I_1 = \left\{ (x_{1i}, x_{2i}) : \frac{x_{1i} - \mu_1}{\sigma_1} < \frac{x_{2i} - \mu_2}{\sigma_2} \right\}, I_2 = \left\{ (x_{1i}, x_{2i}) : \frac{x_{1i} - \mu_1}{\sigma_1} > \frac{x_{2i} - \mu_2}{\sigma_2} \right\}$$

Usual Likelihood function for the parameters of the BVPA can be written as

$$\begin{aligned} & L(\mu_1, \mu_2, \sigma_1, \sigma_2, \alpha_0, \alpha_1, \alpha_2) \\ &= n_1 \ln \alpha_1 + n_1 \ln(\alpha_0 + \alpha_2) - n_1 \ln \sigma_1 - n_1 \ln \sigma_2 \\ & - (\alpha_0 + \alpha_2 + 1) \sum_{i \in I_1} \ln\left(1 + \frac{x_{2i} - \mu_2}{\sigma_2}\right) - (\alpha_1 + 1) \sum_{i \in I_1} \ln\left(1 + \frac{x_{1i} - \mu_1}{\sigma_1}\right) \\ & - n_2 \ln \sigma_1 - n_2 \ln \sigma_2 + n_2 \ln \alpha_2 + n_2 \ln(\alpha_0 + \alpha_1) \\ & - (\alpha_0 + \alpha_1 + 1) \sum_{i \in I_2} \ln\left(1 + \frac{x_{1i} - \mu_1}{\sigma_1}\right) - (\alpha_2 + 1) \sum_{i \in I_2} \ln\left(1 + \frac{x_{2i} - \mu_2}{\sigma_2}\right) \\ & + n_0 \ln \alpha_0 - n_0 \ln \sigma_1 - (\alpha_0 + \alpha_1 + \alpha_2 + 1) \sum_{i \in I_0} \ln\left(1 + \frac{x_{1i} - \mu_1}{\sigma_1}\right) \end{aligned}$$

Direct maximum likelihood estimates of the parameters based on $(X_{1i}, X_{2i}) : i = 1(1)n$ may not be simple. We implement EM algorithm first given some μ_1, μ_2, σ_1 and σ_2 . It requires identification of some missing structure within the problem.

We do not know X_1 is $\sigma_1 U_0 + \mu_1$ or U_1 and We do not know X_2 is $\sigma_2 U_0 + \mu_2$ or U_2 . So we introduce two new random variables $(\Delta_1; \Delta_2)$ as

$$\Delta_1 = \begin{cases} 0 & \text{if } X_1 = \sigma_1 U_0 + \mu_1 \\ 1 & \text{if } X_1 = U_1 \end{cases}$$

and

$$\Delta_2 = \begin{cases} 0 & \text{if } X_2 = \sigma_2 U_0 + \mu_2 \\ 2 & \text{if } X_2 = U_2 \end{cases}$$

The Δ_1 and Δ_2 are the missing values of the E-M algorithm. To calculate the E-step we need the conditional distribution of Δ_1 and Δ_2 .

Using the definition of X_1, X_2, Δ_1 and Δ_2 we have :

▷ For group I_0 , both Δ_1 and Δ_2 are known,

$$\Delta_1 = \Delta_2 = 0$$

▷ For group I_1 , Δ_1 is known, Δ_2 is unknown,

$$\Delta_1 = 1, \Delta_2 = 0 \text{ or } 2$$

Therefore we need to find out $u_1 = P(\Delta_2 = 0|I_1)$ and $u_2 = P(\Delta_2 = 2|I_1)$

▷ For group I_2 , Δ_2 is known, Δ_1 is unknown,

$$\Delta_1 = 0 \text{ or } 1, \Delta_2 = 2$$

Moreover we need $w_1 = P(\Delta_1 = 0|I_2)$ and $w_2 = P(\Delta_1 = 1|I_2)$

Since, each posterior probability corresponds to one of the ordering from Table-1, we calculate u_1, u_2, w_1, w_2 using the probability of appropriate ordering, where $U_1^* = \frac{(U_1 - \mu_1)}{\sigma_1}, U_2^* = \frac{(U_2 - \mu_2)}{\sigma_2}$.

Ordering	(X_1, X_2)	Group
$U_0 < U_1^* < U_2^*$	(U_0, U_0)	I_0
$U_0 < U_2^* < U_1^*$	(U_0, U_0)	I_0
$U_1^* < U_0 < U_2^*$	(U_1^*, U_0)	I_1
$U_1^* < U_2^* < U_0$	(U_1^*, U_2^*)	I_1
$U_2^* < U_0 < U_1^*$	(U_0, U_2^*)	I_2
$U_2^* < U_1^* < U_0$	(U_1^*, U_2^*)	I_2

Table 1 Groups and corresponding orderings of hidden random variables U_0, U_1^* and U_2^*

We have the following expressions for u_1, u_2, w_1, w_2

$$u_1 = \frac{P(U_1^* < U_0 < U_2^*)}{P(U_1^* < U_0 < U_2^*) + P(U_1^* < U_2^* < U_0)}$$

$$\begin{aligned}
u_2 &= \frac{P(U_1^* < U_2^* < U_0)}{P(U_1^* < U_0 < U_2^*) + P(U_1^* < U_2^* < U_0)} \\
w_1 &= \frac{P(U_2^* < U_0 < U_1^*)}{P(U_2^* < U_0 < U_1^*) + P(U_2^* < U_1^* < U_0)} \\
w_2 &= \frac{P(U_2^* < U_1^* < U_0)}{P(U_2^* < U_0 < U_1^*) + P(U_2^* < U_1^* < U_0)}
\end{aligned}$$

Now we have

$$\begin{aligned}
P(U_1^* < U_0 < U_2^*) &= \int_0^\infty [1 - (1+x)^{-\alpha_1}] \alpha_0 (1+x)^{-(\alpha_0+1)} (1+x)^{-(\alpha_2)} \\
&= \alpha_0 \int_0^\infty (1+x)^{-(\alpha_0+\alpha_2+1)} - (1+x)^{-(\alpha_0+\alpha_1+\alpha_2+1)} \\
&= \frac{\alpha_0 \alpha_1}{(\alpha_0 + \alpha_2 + 1)(\alpha_0 + \alpha_1 + \alpha_2 + 1)}
\end{aligned}$$

Using this above result we evaluate the other probabilities to get values of u_1, u_2, w_1, w_2 as : $u_1 = \frac{\alpha_0}{\alpha_0 + \alpha_2}$ and $u_2 = \frac{\alpha_2}{\alpha_0 + \alpha_2}$ $w_1 = \frac{\alpha_0}{\alpha_0 + \alpha_1}$ and $w_2 = \frac{\alpha_1}{\alpha_0 + \alpha_1}$

3.1 Pseudo-likelihood expression

We define n_0, n_1, n_2 as : $n_0 = |I_0|, n_1 = |I_1|, n_2 = |I_2|$ where $|I_j|$ for $j = 0, 1, 2$ denotes the number of elements in the set I_j . Now the pseudo log-likelihood can be written down as

$$\begin{aligned}
Q &= L(\alpha_0, \alpha_1, \alpha_2) \\
&= -\alpha_0 \left(\sum_{i \in I_0} \ln(1+x_i) + \sum_{i \in I_2} \ln(1+x_{1i}) + \sum_{i \in I_1} \ln(1+x_{2i}) \right) \\
&\quad + (n_0 + u_1 n_1 + w_1 n_2) \ln \alpha_0 - \alpha_1 \left(\sum_{i \in I_0} \ln(1+x_i) + \sum_{i \in I_1 \cup I_2} \ln(1+x_{1i}) \right) \\
&\quad + (n_1 + w_2 n_2) \ln \alpha_1 - \alpha_2 \left(\sum_{i \in I_0} \ln(1+x_i) + \sum_{i \in I_1 \cup I_2} \ln(1+x_{2i}) \right) \\
&\quad + (n_2 + u_2 n_1) \ln \alpha_2
\end{aligned} \tag{1}$$

Therefore M-step involves maximizing (1) with respect to $\alpha_0, \alpha_1, \alpha_2$ at

$$\hat{\alpha}_0^{(t+1)} = \frac{n_0 + u_1^{(t)} n_1 + w_1^{(t)} n_2}{\sum_{i \in I_0} \ln(1+x_i) + \sum_{i \in I_2} \ln(1+x_{1i}) + \sum_{i \in I_1} \ln(1+x_{2i})}. \tag{2}$$

$$\hat{\alpha}_1^{(t+1)} = \frac{(n_1 + w_2^{(t)} n_2)}{\sum_{i \in I_0} \ln(1+x_i) + \sum_{i \in (I_1 \cup I_2)} \ln(1+x_{1i})} \tag{3}$$

$$\hat{\alpha}_2^{(t+1)} = \frac{(n_2 + u_2^{(t)} n_1)}{(\sum_{i \in I_0} \ln(1+x_i) + \sum_{i \in (I_1 \cup I_2)} \ln(1+x_{2i}))} \tag{4}$$

Therefore the algorithm can be given as

Algorithm 1 EM procedure for bivariate Pareto distribution

-
- 1: Calculate the estimates of μ_1, μ_2, σ_1 and σ_2 from marginals of bivariate distribution.
 - 2: Fix I_0, I_1 and I_2 .
 - 3: **while** $\Delta Q/Q < tol$ **do**
 - 4: Compute $u_1^{(i)}, u_2^{(i)}, w_1^{(i)}, w_2^{(i)}$ from $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 5: Update $\alpha_0^{(i+1)}, \alpha_1^{(i+1)}, \alpha_2^{(i+1)}$ using Equation (2), (3) and (4).
 - 6: Calculate Q for the new iterate.
 - 7: **end while**
-

3.2 Algorithm-2 : Modified Algorithm 1

The above algorithm does not work even if all the I_0, I_1 and I_2 are non-empty. In our previous approach we calculate the estimates of location and scale parameters and construct the above three sets based on the transformed variables using those estimated location and scale parameters. Therefore it is high likely that there will be almost no element in I_0 for most of the generated sample. Consequently it should provide the estimate of α_0 as zero most of the time.

We propose to make few modifications in the above EM algorithm. Since we observe the normalized data with respect to the estimated location and scale parameter, the transformation is not going to provide the distribution of normalized data exactly as Pareto with location zero and scale one. The transformation rather form some distribution close to Pareto with location zero and scale one. It is very difficult to know the exact distribution. Approximately even if we assume some Pareto with location parameter near zero and scale parameter near one, calculation of probability that an observation will come from any one of I_0, I_1 or I_2 , becomes another problem.

To avoid such instances, we assume that information of all n_0, n_1 and n_2 are missing because of small perturbation inherited by estimated location and scale parameters. In ideal situation we should get EM algorithm steps as described in 3. Since we have now three more latent information n_0, n_1 and n_2 , we try to approximate them and incorporate in the original EM algorithm.

It is clear that (n_0, n_1, n_2) will jointly follow multinomial distribution with parameters $n = n_0 + n_1 + n_2$ and $(\frac{\alpha_0}{\alpha_0 + \alpha_1 + \alpha_2}, \frac{\alpha_1}{\alpha_0 + \alpha_1 + \alpha_2}, \frac{\alpha_2}{\alpha_0 + \alpha_1 + \alpha_2})$.

We approximate $\tilde{n}_i = n \cdot \frac{\alpha_i}{\alpha_0 + \alpha_1 + \alpha_2}$, $i = 0, 1, 2$.

Therefore M-step involves maximizing (1) with respect to $\alpha_0, \alpha_1, \alpha_2$ at

$$\hat{\alpha}_0^{(t+1)} = \frac{\tilde{n}_0 + u_1^{(t)} \tilde{n}_1 + w_1^{(t)} \tilde{n}_2}{\sum_{i \in I_0} \ln(1 + x_i) + \sum_{i \in I_2} \ln(1 + x_{1i}) + \sum_{i \in I_1} \ln(1 + x_{2i})}. \quad (5)$$

$$\hat{\alpha}_1^{(t+1)} = \frac{(\tilde{n}_1 + w_2^{(t)} \tilde{n}_2)}{\sum_{i \in I_0} \ln(1 + x_i) + \sum_{i \in (I_1 \cup I_2)} \ln(1 + x_{1i})} \quad (6)$$

$$\hat{\alpha}_2^{(t+1)} = \frac{\tilde{n}_2 + u_2^{(t)} \tilde{n}_1}{(\sum_{i \in I_0} \ln(1 + x_i) + \sum_{i \in (I_1 \cup I_2)} \ln(1 + x_{2i}))} \quad (7)$$

Finally modified algorithm would be

Algorithm 2 EM procedure for bivariate Pareto distribution

-
- 1: Calculate the estimates of μ_1, μ_2, σ_1 and σ_2 from marginals of bivariate distribution.
 - 2: Fix I_0, I_1 and I_2 .
 - 3: **while** $\Delta Q/Q < tol$ **do**
 - 4: Compute $u_1^{(i)}, u_2^{(i)}, w_1^{(i)}, w_2^{(i)}, \tilde{n}_0, \tilde{n}_1$ and \tilde{n}_2 from $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 5: Update $\alpha_0^{(i+1)}, \alpha_1^{(i+1)}, \alpha_2^{(i+1)}$ using Equation (5), (6) and (7).
 - 6: Calculate Q for the new iterate.
 - 7: **end while**
-

3.3 Algorithm-3 : Modified Algorithm 2

In this case we update σ_1 and σ_2 along with EM iterations similar to what Asimit et al. has done in their paper. But the main difference of our update is that it is based on one step ahead gradient descent instead of fixed point iteration on scale parameters. Surely gradient descent with respect to bivariate likelihood won't work as bivariate likelihood is a discontinuous function with respect to location and scale parameters. Therefore we estimate scale parameters using density of marginals. At every iteration we use one step ahead gradient descend of σ_1 and σ_2 based on likelihood of its marginal density combined with usual EM steps for other parameters to solve the problem. The idea is similar to stochastic gradient descend. We begin the algorithm estimating the sets I_0, I_1 and I_2 which involves location and scale parameters. Given location and scale parameters, we can update previous EM steps of α_0, α_1 and α_2 . EM steps will ensure to take the correct direction of α_0, α_1 and α_2 starting from any values of this three parameters and also it will enable gradient descend steps of σ_1 and σ_2 to converge faster starting from any values.

This algorithm works even for moderate sample sizes. However it takes lot of time to converge or roam around the actual value for some really bad sample. We calculate the estimates both with full iterations until convergence and with at most 2000 iterations. We observe MSEs and the number of iterations in both the cases. We expect that within 2000 iterations estimated points should be sufficiently closer to the actual value or it should take some value which can be good estimate for starting point of some other optimization algorithms.

Algorithm 3 EM procedure for bivariate Pareto distribution

-
- Take the estimates of μ_1, μ_2 by $\mu_1 = \min\{X_{1i}; i = 1, \dots, n\}$ and $\mu_2 = \min\{X_{2i}; i = 1, \dots, n\}$.
- 2: Start with some initial choice of $\sigma_1, \sigma_2, \alpha_0, \alpha_1, \alpha_2$.
 - while** $|\Delta Q/Q| > tol$ **do**
 - 4: Fix I_0, I_1 and I_2 with estimated $\mu_1, \mu_2, \sigma_1^{(i)}$ and $\sigma_2^{(i)}$.
 - Compute updates of $\sigma_1^{(i)}, \sigma_2^{(i)}$ through one step ahead gradient descend, using $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 6: Compute $u_1^{(i)}, u_2^{(i)}, w_1^{(i)}, w_2^{(i)}, \tilde{n}_0, \tilde{n}_1$ and \tilde{n}_2 from $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - Update $\alpha_0^{(i+1)}, \alpha_1^{(i+1)}, \alpha_2^{(i+1)}$ using Equation (5), (6) and (7).
 - 8: Calculate Q for the new iterate.
 - end while**
-

3.4 Algorithm-4 : Modified Algorithm 3

We can make more variations on it. This is similar to what Asimit et al. has done in their paper. At each iteration we update σ_1 and σ_2 from the marginal density using one step ahead fixed point iteration instead of Gradient decent algorithm and carry on the same EM algorithm. However approach of Asimit et al is different, therefore EM steps are also differing.

Algorithm 4 EM procedure for bivariate Pareto distribution

- 1: Take the estimates of μ_1, μ_2 by $\mu_1 = \min\{X_{1i}; i = 1, \dots, n\}$ and $\mu_2 = \min\{X_{2i}; i = 1, \dots, n\}$.
 - 2: Start with some initial choice of $\sigma_1, \sigma_2, \alpha_0, \alpha_1, \alpha_2$.
 - 3: **while** $|\Delta Q/Q| > tol$ **do**
 - 4: Fix I_0, I_1 and I_2 with estimated $\mu_1, \mu_2, \sigma_1^{(i)}$ and $\sigma_2^{(i)}$.
 - 5: Compute one step ahead update of $\sigma_1^{(i)}, \sigma_2^{(i)}$ through fixed point iteration, using $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 6: Compute $u_1^{(i)}, u_2^{(i)}, w_1^{(i)}, w_2^{(i)}, \tilde{n}_0, \tilde{n}_1$ and \tilde{n}_2 from $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 7: Update $\alpha_0^{(i+1)}, \alpha_1^{(i+1)}, \alpha_2^{(i+1)}$ using Equation (5), (6) and (7).
 - 8: Calculate Q for the new iterate.
 - 9: **end while**
-

3.5 Algorithm-5 : Modified Algorithm 4

This variation is of two fold. We update σ_1 and σ_2 from the marginal density until convergence using fixed point iteration instead of Gradient decent algorithm and carry forward the same EM algorithm. However at the time of using EM iterations, we update the σ_1 and σ_2 using one step ahead fixed point iteration along with other parameters. The detailed algorithmic steps are shown below.

Algorithm 5 EM procedure for bivariate Pareto distribution

- 1: Start with some initial choice of $\sigma_1, \sigma_2, \alpha_0, \alpha_1, \alpha_2$.
 - 2: **while** $|\Delta \sigma| > tol$ **do**
 - 3: Compute one step ahead fixed point iteration of $\sigma_1^{(i+1)}, \sigma_2^{(i+1)}$, using $\sigma_1^{(i)}, \sigma_2^{(i)}, \alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 4: Compute $u_1^{(i)}, u_2^{(i)}, w_1^{(i)}, w_2^{(i)}, \tilde{n}_0, \tilde{n}_1$ and \tilde{n}_2 from $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 5: Update $\alpha_0^{(i+1)}, \alpha_1^{(i+1)}, \alpha_2^{(i+1)}$ using Equation (5), (6) and (7).
 - 6: **end while**
 - 7: **while** $|\Delta Q/Q| > tol$ **do**
 - 8: Fix I_0, I_1 and I_2 with last updated estimates of $\mu_1, \mu_2, \sigma_1^{(i)}, \sigma_2^{(i)}, \alpha_0^{(i)}, \alpha_1^{(i)}$ and $\alpha_2^{(i)}$.
 - 9: Compute updates of $\sigma_1^{(i+1)}, \sigma_2^{(i+1)}$, using $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 10: Compute $u_1^{(i)}, u_2^{(i)}, w_1^{(i)}, w_2^{(i)}, \tilde{n}_0, \tilde{n}_1$ and \tilde{n}_2 from $\alpha_0^{(i)}, \alpha_1^{(i)}, \alpha_2^{(i)}$.
 - 11: Update $\alpha_0^{(i+1)}, \alpha_1^{(i+1)}, \alpha_2^{(i+1)}$ using Equation (5), (6) and (7).
 - 12: Calculate Q for the new iterate.
 - 13: **end while**
-

4 Numerical Results

We use package R 3.2.3 to perform the estimation procedure. All the programs will be available from author on request. First we take four different sets of parameters and observe the average iteration calculated over different sample sizes. We take our sample size as $n = 150, 250, 350, 450$. We can find the results in Table-2 based on 1000 replications. Tolerance limit (denoted as "tol") of stopping criteria is taken as 0.00001. We have used stopping criteria as absolute value of likelihood changes with respect to previous Qseudo-likelihood at each iteration. Results shown in Table-2 provide average iteration by six different approaches. They are (a) Modified Approach 1 (b) Modified Approach 2 which we truncate after 2000 iterations and denote this as Modified Approach 2 (T) (c) Modified Approach 2 without truncating the iteration and denote this as Modified Approach 2 (WT) (d) Modified Approach 3 (e) Modified Approach 4 (f) Asimit's approach. We choose four different sets of parameters to compare the results. These are $\mu_1 = 0, \mu_2 = 0, \sigma_1 = 1, \sigma_2 = 0.5, \alpha_0 = 1, \alpha_1 = 0.3, \alpha_2 = 1.4$; $\mu_1 = 1, \mu_2 = 2, \sigma_1 = 0.4, \sigma_2 = 0.5, \alpha_0 = 2, \alpha_1 = 1.2, \alpha_2 = 1.4$; $\mu_1 = 0, \mu_2 = 0, \sigma_1 = 1.4, \sigma_2 = 0.5, \alpha_0 = 1, \alpha_1 = 1, \alpha_2 = 1.4$; $\mu_1 = 0, \mu_2 = 0, \sigma_1 = 1.4, \sigma_2 = 0.5, \alpha_0 = 2, \alpha_1 = 0.4, \alpha_2 = 0.5$. The proposed EM algorithms work for any initial value. However estimates of μ_1 and μ_2 are always $\hat{\mu}_1 = \min\{X_{1i}; 1, \dots, n\}$ and $\hat{\mu}_2 = \min\{X_{2i}; 1, \dots, n\}$ respectively. Original paper of Asimit et al. uses different stopping criteria and results are provided for large sample sizes. To compare our results, we keep stopping criteria and sample size same across all algorithms.

Average estimates are provided in Table-3 when samples are simulated from bivariate pareto with parameters $\mu_1 = 0, \mu_2 = 0, \sigma_1 = 1, \sigma_2 = 0.5, \alpha_0 = 1, \alpha_1 = 0.3, \alpha_2 = 1.4$. Results of average estimates are shown only by two best approaches (best in the sense of minimum average iteration) i.e. Modified Approach 1 and Modified Approach 4. Table-4 and Table-5 show the MSEs for all procedures where samples are generated from the following parameter sets : $\mu_1 = 0, \mu_2 = 0, \sigma_1 = 1, \sigma_2 = 0.5, \alpha_0 = 1, \alpha_1 = 0.3, \alpha_2 = 1.4$. MSE is an important criteria for selecting the best algorithm. Therefore MSEs are calculated for all the procedures. We also calculate parametric bootstrap confidence interval. We simulate 1000 samples and estimate of the parameters based on the simulated samples. We use them to get the 95% confidence interval by calculating 0.025 and 0.975 sample quantile points of estimated parameters. Table-6 to Table-10 carry the information related to parametric bootstrap confidence intervals.

Important Comments :

1. Mean square error for α_2 is little higher for all methods.
2. MSEs are more or less same for all methods. In the above result we see Modified Approach 1 provides minimum MSE among all procedures. For large sample size Modified Approach 1 comes out as winner for any chosen parameter sets. Performance of Modified Approach 2 with truncation is also worth mentioning, as it provides second best performance with respect to minimum MSE.
3. Modified Approach 4 is the best among other approaches in terms of average iteration. Modified Approach 1 appears to be the second best performer. How-

ever average iteration for Modified Approach 2 with truncation is much higher as compared to Modified Approach 1 and 4.

4. As expected, average iteration is very high in case of Modified Approach 2 (WT).
5. If we compare the algorithms based on minimum MSE and average iteration together, Modified approach 1 stands either winner or closer to the winner.

parameter set		n	150	250	350	450
$\mu_1 = 0, \mu_2 = 0$ $\sigma_1 = 1, \sigma_2 = 0.5$ $\alpha_0 = 1, \alpha_1 = 0.3,$ $\alpha_2 = 1.4$	The average number of Iterations (AI)	Modified Approach 1	244	231	223	220
		Modified Approach 2 (WT)	3402	2163	1542	1220
		Modified Approach 2 (T)	1771	1631	1414	1194
		Modified Approach 3	266	249	238	234
		Modified Approach 4	191	162	151	147
		Asimit's Approach	1002	945	909	902
$\mu_1 = 1, \mu_2 = 2$ $\sigma_1 = 0.4, \sigma_2 = 0.5$ $\alpha_0 = 2, \alpha_1 = 1.2,$ $\alpha_2 = 1.4$	The average number of Iterations (AI)	Modified Approach 1	174	165	159	157
		Modified Approach 2 (WT)	7734	4137	2428	1633
		Modified Approach 2 (T)	1704	1553	1378	1184
		Modified Approach 3	764	551	458	424
		Modified Approach 4	243	185	153	128
		Asimit's Approach	2345	1747	1464	1309
$\mu_1 = 0, \mu_2 = 0$ $\sigma_1 = 1.4, \sigma_2 = 0.5$ $\alpha_0 = 1, \alpha_1 = 1,$ $\alpha_2 = 1.4$	The average number of Iterations (AI)	Modified Approach 1	168	155	150	147
		Modified Approach 2 (WT)	7230	4372	3077	2423
		Modified Approach 2 (T)	1617	1738	1732	1697
		Modified Approach 3	265	237	220	209
		Modified Approach 4	123	101	91	83
		Asimit's Approach	897	547	484	453
$\mu_1 = 0, \mu_2 = 0$ $\sigma_1 = 1.4, \sigma_2 = 0.5$ $\alpha_0 = 2, \alpha_1 = 0.4,$ $\alpha_2 = 0.5$	The average number of Iterations (AI)	Modified Approach 1	138	129	124	122
		Modified Approach 2 (WT)	2172	2707	2237	1821
		Modified Approach 2 (T)	397	591	773	881
		Modified Approach 3	330	261	238	219
		Modified Approach 4	117	100	96	95
		Asimit's Approach	1145	977	919	875

Table 2 Average number of iterations (AI) by six approaches: (a) Modified Approach 1 (b) Modified Approach 2 (WT) (c) Modified Approach 2 (T) (d) Modified Approach 3 (e) Modified Approach 4 (f) Asimit's Approach

n	parameters	μ_1	μ_2	σ_1	σ_2	α_0	α_1	α_2
150	Average (Mod. 1) estimates (AE)	0.0056	0.0014	0.9031	0.5154	0.8993	0.3379	1.5660
	Average (Mod. 4) estimates (AE)	0.0056	0.0014	1.0521	0.5755	0.9372	0.4122	1.7416
250	Average (Mod. 1) Estimates (AE)	0.0031	0.0008	0.8943	0.4989	0.9013	0.3226	1.5046
	Average (Mod. 4) estimates (AE)	0.0031	0.0008	1.0386	0.5457	0.9408	0.3909	1.6292
350	Average (Mod. 1) Estimates (AE)	0.0022	0.0006	0.8854	0.4878	0.9086	0.3107	1.4550
	Average (Mod. 4) estimates (AE)	0.0022	0.0006	1.0252	0.5283	0.9497	0.3745	1.5543
450	Average (Mod. 1) Estimates (AE)	0.0017	0.0005	0.8719	0.4799	0.9099	0.2955	1.4270
	Average (Mod. 4) estimates (AE)	0.0017	0.0005	1.0077	0.5187	0.9492	0.3586	1.5221

Table 3 The average estimates (AE) for $\mu_1 = 0$, $\mu_2 = 0$, $\sigma_1 = 1$, $\sigma_2 = 0.5$, $\alpha_0 = 1$, $\alpha_1 = 0.3$ and $\alpha_2 = 1.4$ through two best approaches i.e. Modified Approach 1 and Modified Approach 4

	parameters	μ_1	μ_2	σ_1
n=150	MSE (Mod. 1)	6.14e-05	3.69e-06	0.0832
	MSE (Mod. 2 (T))	0.0000614	0.0000037	0.0543
	MSE (Mod. 2 (WT))	0.0000614	0.0000370	0.1043
	MSE (Mod. 3)	0.0000614	0.0000037	0.1216
	MSE (Mod. 4)	0.0000614	0.0000037	0.1179
	MSE (Asimit)	0.0000614	0.0000037	0.1218
n = 150	σ_2	α_0	α_1	α_2
MSE (Mod. 1)	0.0477	0.0402	0.0375	0.5178
MSE (Mod. 2 (T))	0.0368	0.0386	0.0342	0.4154
MSE (Mod. 2 (WT))	0.0682	0.0433	0.0609	0.7969
MSE (Mod. 3)	0.1028	0.0439	0.0737	1.2035
MSE (Mod. 4)	0.1090	0.0429	0.0691	1.2536
MSE (Asimit)	0.1011	0.0433	0.0726	1.1860
	parameters	μ_1	μ_2	σ_1
n = 250	MSE (Mod. 1)	0.000020	1.4757e-06	0.0581
	MSE (Mod. 2 (T))	0.0000614	0.00000370	0.0543
	MSE (Mod. 2 (WT))	0.0000198	0.0000014	0.0682
	MSE (Mod. 3)	0.00001978	0.00000147	0.0745
	MSE (Mod. 4)	0.00001978	0.00000147	0.0724
	MSE (Asimit)	0.00001978	0.00000148	0.0746
n = 250	σ_2	α_0	α_1	α_2
MSE (Mod. 1)	0.0278	0.0296	0.0205	0.2817
MSE (Mod. 2 (T))	0.0368	0.0386	0.0342	0.4153
MSE (Mod. 2 (WT))	0.0410	0.0294	0.0386	0.4441
MSE (Mod. 3)	0.0461	0.0296	0.0431	0.5033
MSE (Mod. 4)	0.0460	0.0289	0.0401	0.4978
MSE (Asimit)	0.0462	0.0292	0.0427	0.5638

Table 4 Mean Square Error (MSE) through all approaches when samples are generated from $\mu_1 = 0$, $\mu_2 = 0$, $\sigma_1 = 1$, $\sigma_2 = 0.5$, $\alpha_0 = 1$, $\alpha_1 = 0.3$ and $\alpha_2 = 1.4$

	parameters	μ_1	μ_2	σ_1
n = 350	MSE (Mod. 1)	9.236e-06	6.992e-07	0.0427
	MSE (Mod. 2 (T))	0.00000197	0.000000699	0.0389
	MSE (Mod. 2 (WT))	0.00000924	0.000000699	0.0428
	MSE (Mod. 3)	0.00000923	0.000000699	0.0456
	MSE (Mod. 4)	0.00000924	0.000000699	0.0440
	MSE (Asimit)	0.00000923	0.000000699	0.0454
n=350	σ_2	α_0	α_1	α_2
	MSE (Mod. 1)	0.0155	0.0211	0.0143
	MSE (Mod. 2 (T))	0.02001	0.01907	0.02573
	MSE (Mod. 2 (WT))	0.0213	0.0192	0.0280
	MSE (Mod. 3)	0.0221	0.0194	0.0302
	MSE (Mod. 4)	0.0220	0.0188	0.0275
	MSE (Asimit)	0.0222	0.0194	0.0301
	parameters	μ_1	μ_2	σ_1
n = 450	MSE (Mod. 1)	5.589e-06	4.398e-07	0.0366
	MSE (Mod. 2 (T))	0.0000055	0.000000440	0.0283
	MSE (Mod. 2 (WT))	0.00000559	0.00000044	0.0288
	MSE (Mod. 3)	0.00000559	0.00000047	0.03008
	MSE (Mod. 4)	0.00000558	0.000000439	0.0292
	MSE (Asimit)	0.00000559	0.000000440	0.0300
n=450	σ_2	α_0	α_1	α_2
	MSE (Mod. 1)	0.0107	0.0176	0.0097
	MSE (Mod. 2 (T))	0.0143	0.015	0.0181
	MSE (Mod. 2 (WT))	0.0147	0.0151	0.0184
	MSE (Mod. 3)	0.0151	0.0152	0.0196
	MSE (Mod. 4)	0.0150	0.0148	0.0179
	MSE (Asimit)	0.0151	0.0151	0.0195

Table 5 Mean Square Error (MSE) through all approaches when samples are generated from $\mu_1 = 0$, $\mu_2 = 0$, $\sigma_1 = 1$, $\sigma_2 = 0.5$, $\alpha_0 = 1$, $\alpha_1 = 0.3$ and $\alpha_2 = 1.4$

n	150	250	350	450
parameters				
μ_1	[0.0001, 0.0193]	[0.000077, 0.0125]	[0.00007, 0.0079]	[0.00004, 0.0062]
μ_2	[0.000034, 0.0050]	[0.000027, 0.0034]	[0.00002, 0.0022]	[0.00001, 0.0017]
σ_1	[0.5112, 1.4976]	[0.5513, 1.3754]	[0.6016, 1.2959]	[0.6257, 1.1722]
σ_2	[0.2361, 1.0501]	[0.2796, 0.9011]	[0.3067, 0.7946]	[0.3183, 0.7207]
α_0	[0.6103, 1.2854]	[0.6682, 1.1940]	[0.7004, 1.1586]	[0.7373, 1.1197]
α_1	[0.0974, 0.8052]	[0.1172, 0.6801]	[0.1206, 0.5963]	[0.1241, 0.5377]
α_2	[0.7692, 3.3941]	[0.8701, 2.8723]	[0.9447, 2.3398]	[0.9772, 2.2408]

Table 6 Parametric Bootstrap confidence interval for $\mu_1 = 0$, $\mu_2 = 0$, $\sigma_1 = 1$, $\sigma_2 = 0.5$, $\alpha_0 = 1$, $\alpha_1 = 0.3$ and $\alpha_2 = 1.4$ in Modified Approach 1

5 Data Analysis

We analyze a data set on the indemnity payments (Loss) and allocated loss adjustment expense (ALAE) relating to 1500 general liability claims from insurance companies are available in the R package *evd* ([fbvplot by Chris Ferro \(2015\)](#)). From [Falk and Guillou \(2008\)](#), we know that peak over threshold method on random variable U provides polynomial generalized Pareto distribution for any x_0 with $1 + \log(G(x_0)) \in (0, 1)$ i.e. $P(U > tx_0 | U > x_0) = t^{-\alpha}$, $t \geq 1$ where $G(\cdot)$ is the distribution function of U . We choose appropriate t and x_0 so that data should behave like near Pareto distribution.

n parameters	150	250	350	450
μ_1	[0.0001, 0.01929]	[0.000077, 0.01248]	[0.000067, 0.00792]	[0.0000449, 0.00619]
μ_2	[0.000034, 0.004952]	[0.0000265, 0.00337]	[0.0000203, 0.002219]	[0.0000155, 0.00175]
σ_1	[0.5872, 1.4421]	[0.6316, 1.4810]	[0.6921, 1.4760]	[0.71373, 1.3677]
σ_2	[0.2497, 0.9518]	[0.2876, 0.9674]	[0.3277, 0.8631]	[0.33937, 0.79859]
α_0	[0.6146, 1.3454]	[0.6684, 1.2655]	[0.7089, 1.2288]	[0.75384, 1.17617]
α_1	[0.1340, 0.8017]	[0.1608, 0.7578]	[0.1772, 0.7227]	[0.19451, 0.66522]
α_2	[0.7949, 3.0734]	[0.9092, 3.0089]	[0.9932, 2.5686]	[1.03731, 2.455004]

Table 7 Parametric Bootstrap confidence interval for $\mu_1 = 0$, $\mu_2 = 0$, $\sigma_1 = 1$, $\sigma_2 = 0.5$, $\alpha_0 = 1$, $\alpha_1 = 0.3$ and $\alpha_2 = 1.4$ in Modified Approach 2 (T)

n parameters	150	250	350	450
μ_1	[0.000108, 0.01929]	[0.000077, 0.01248]	[0.000067, 0.00791]	[0.000044, 0.0061]
μ_2	[0.0000343, 0.00495]	[0.0000265, 0.00337]	[0.0000203, 0.0022]	[0.000015, 0.0017]
σ_1	[0.5872, 1.7322]	[0.6316, 0.1.6122]	[0.69207, 1.5152]	[0.7137, 1.3751]
σ_2	[0.2496, 1.1774]	[0.2877, 1.0454]	[0.3277, 0.8711]	[0.3394, 0.7986]
α_0	[0.5967, 1.3852]	[0.6684, 1.2719]	[0.7089, 1.2281]	[0.7538, 1.1762]
α_1	[0.1346, 0.9461]	[0.1585, 0.8536]	[0.1773, 0.7503]	[0.1945, 0.6796]
α_2	[0.7951, 3.7727]	[0.9093, 3.2908]	[0.9932, 2.5707]	[1.0373, 2.4596]

Table 8 Parametric Bootstrap confidence interval for $\mu_1 = 0$, $\mu_2 = 0$, $\sigma_1 = 1$, $\sigma_2 = 0.5$, $\alpha_0 = 1$, $\alpha_1 = 0.3$ and $\alpha_2 = 1.4$ in Modified Approach 2 (WT)

n parameters	150	250	350	450
μ_1	[0.000108, 0.019]	[0.000077, 0.0125]	[0.000067, 0.0079]	[0.000045, 0.0062]
μ_2	[0.000034, 0.0049]	[0.000026, 0.00337]	[0.0000203, 0.0022]	[0.000016, 0.0017]
σ_1	[0.5846, 1.1874]	[0.6355, 1.6567]	[0.6956, 1.5428]	[0.7161, 1.3938]
σ_2	[0.2406, 1.2967]	[0.2875, 1.0817]	[0.3277, 0.8795]	[0.3393, 0.8067]
α_0	[0.5921, 1.4047]	[0.6656, 1.2716]	[0.7075, 1.2319]	[0.7520, 1.1790]
α_1	[0.1378, 0.9898]	[0.1621, 0.8843]	[0.1798, 0.7653]	[0.1946, 0.6894]
α_2	[0.8007, 4.127]	[0.9103, 3.4069]	[0.9941, 2.6199]	[1.0376, 2.4805]

Table 9 Parametric Bootstrap confidence interval for $\mu_1 = 0$, $\mu_2 = 0$, $\sigma_1 = 1$, $\sigma_2 = 0.5$, $\alpha_0 = 1$, $\alpha_1 = 0.3$ and $\alpha_2 = 1.4$ in Modified Approach 3

n parameters	150	250	350	450
μ_1	[0.0001, 0.0193]	[0.000077, 0.0125]	[0.000067, 0.00791]	[0.000044, 0.0062]
μ_2	[0.000034, 0.0050]	[0.000026, 0.00337]	[0.0000203, 0.0022]	[0.000015, 0.0017]
σ_1	[0.5821, 1.7914]	[0.6332, 1.6449]	[0.6936, 1.5209]	[0.7136, 1.3763]
σ_2	[0.2469, 1.2961]	[0.2874, 1.0815]	[0.3246, 0.8798]	[0.3373, 0.8037]
α_0	[0.6102, 1.4071]	[0.6792, 1.2729]	[0.7124, 1.2345]	[0.7562, 1.1791]
α_1	[0.1261, 0.9899]	[0.1518, 0.8474]	[0.1737, 0.7367]	[0.1892, 0.6740]
α_2	[0.7934, 4.1231]	[0.9044, 3.4036]	[0.9831, 2.6059]	[1.0284, 2.4709]

Table 10 Parametric Bootstrap confidence interval for $\mu_1 = 0$, $\mu_2 = 0$, $\sigma_1 = 1$, $\sigma_2 = 0.5$, $\alpha_0 = 1$, $\alpha_1 = 0.3$ and $\alpha_2 = 1.4$ in Modified Approach 4

We assume the data to follow near equal to singular Marshall Olkin bivariate Pareto and try to verify our assumption. We fit the empirical survival functions with the marginals of this bivariate Pareto whose parameters can be obtained from the EM algorithm that we have developed. Figure-3 shows a good fit for both the marginals. The data analysis is performed based on sample size 1500.

Empirical two dimensional density plot in Figure-4 verifies that Marshall-Olkin Pareto can be an alternative model for the transformed dataset. Parameter estimates of this bivariate distribution based on sample size $n = 468$ are provided in Table-11 and Table-12 whereas parametric bootstrap confidence intervals are provided in Table-13 and Table-14 respectively. All algorithms are used separately to calculate the estimates and confidence intervals.




Fig. 3 Survival plots for two marginals of the transformed dataset




Fig. 4 Two dimensional density plots of the transformed dataset

Procedures	parameters	μ_1	μ_2	σ_1
Mod. 1		1	0.4	2.5594
Mod. 2 (T)		1	0.4	2.4936
Mod. 2 (WT)		1	0.4	2.9459
Mod. 3		1	0.4	3.2459
Mod. 4		1	0.4	3.1802

Table 11 EM estimates of the parameters - Data Analysis

Procedures	σ_2	α_0	α_1	α_2
Mod. 1	0.7176	0.7557	0.8162	0.9792
Mod. 2 (T)	0.8164	0.759	0.7878	1.1267
Mod. 2 (WT)	0.8162	0.7960	0.9209	1.0895
Mod. 3	0.8163	0.8171	1.0100	1.0684
Mod. 4	0.7947	0.8330	0.9667	1.0158

Table 12 EM estimates of the parameters - Data Analysis

Procedures	parameters	μ_1	μ_2	σ_1
Mod. 1		[1.00013, 1.0127]	[0.40002, 0.4031]	[1.5793, 2.9953]
Mod. 2 (T)		[1.00013, 1.0127]	[0.40002, 0.4031]	[1.4952, 2.7412]
Mod. 2 (WT)		[1.0001, 1.0135]	[0.40002, 0.4032]	[1.5033, 3.9036]
Mod. 3		[1.00013, 1.0141]	[0.40002, 0.4032]	[2.3578, 4.9274]
Mod. 4		[1.00013, 1.01397]	[0.40002, 0.4032]	[2.3343, 4.7869]

Table 13 Confidence interval - Data Analysis

Procedures	σ_2	α_0	α_1	α_2
Mod. 1	[0.4554, 0.9018]	[0.5435, 0.8524]	[0.5378, 1.0407]	[0.6726, 1.3616]
Mod. 2 (T)	[0.4337, 1.1601]	[0.5234, 0.8712]	[0.4883, 0.9623]	[0.6829, 1.7491]
Mod. 2 (WT)	[0.4448, 1.1833]	[0.5445, 0.9308]	[0.5090, 1.3183]	[0.6909, 1.7151]
Mod. 3	[0.5731, 1.2017]	[0.5869, 0.9988]	[0.7788, 1.6505]	[0.7996, 1.7046]
Mod. 4	[0.5658, 1.1673]	[0.6089, 1.0025]	[0.7526, 1.5866]	[0.7770, 1.6220]

Table 14 Confidence interval - Data Analysis

6 Conclusion

We use different variations of EM algorithms for estimating the parameters of singular bivariate Pareto distribution. All variations work quite well even for moderate sample size (say, 150, 250 etc). Our algorithms outperforms the current state-of-art algorithm by Asimit et al. The data analysis makes the paper more interesting. There was no discussion of any real-life data analysis in the previous paper. We have also calculated parametric bootstrap confidence interval which is absent in Asimit et al. The algorithm shown here can be applicable to other higher dimensional distributions with location and scale parameters. Given a real-life data set model selection can be

an interesting issue among different other form of bivariate pareto distributions. More work is needed in this direction too.

A

A.1

$$S(x_1, x_2) = P(X_1 \geq x_1, X_2 \geq x_2) = \begin{cases} S_1(x_1, x_2) & \text{if } \frac{x_1 - \mu_1}{\sigma_1} < \frac{x_2 - \mu_2}{\sigma_2} \\ S_2(x_1, x_2) & \text{if } \frac{x_1 - \mu_1}{\sigma_1} > \frac{x_2 - \mu_2}{\sigma_2} \\ S_0(x) & \text{if } \frac{x_1 - \mu_1}{\sigma_1} = \frac{x_2 - \mu_2}{\sigma_2} = x \end{cases}$$

$$S_1(x_1, x_2) = \left(1 + \frac{(x_1 - \mu_1)}{\sigma_1}\right)^{-\alpha_1} \left(1 + \frac{x_2 - \mu_2}{\sigma_2}\right)^{-(\alpha_0 + \alpha_2)}$$

$$S_2(x_1, x_2) = \left(1 + \frac{(x_1 - \mu_1)}{\sigma_1}\right)^{-(\alpha_0 + \alpha_1)} \left(1 + \frac{x_2 - \mu_2}{\sigma_2}\right)^{\alpha_2}$$

$$S_0(x) = (1 + x)^{-(\alpha_0 + \alpha_1 + \alpha_2)}$$

From the above expressions we can get the marginal of X_1 and X_2 taking $x_1 \rightarrow \mu_1$ and $x_2 \rightarrow \mu_2$,

$$S_{X_1}(x_1) = \begin{cases} \left(1 + \frac{(x_1 - \mu_1)}{\sigma_1}\right)^{-(\alpha_0 + \alpha_1)} & \text{if } x_1 > \mu_1 \\ 1 & \text{otherwise} \end{cases}$$

$$S_{X_2}(x_2) = \begin{cases} \left(1 + \frac{x_2 - \mu_2}{\sigma_2}\right)^{-(\alpha_0 + \alpha_2)} & \text{if } x_2 > \mu_2 \\ 1 & \text{otherwise} \end{cases}$$

A.2

$$P(\min\{X_1, X_2\} \geq x) = P(X_1 \geq x, X_2 \geq x) \quad (8)$$

This implies

$$P(X_1 > x, X_2 > x) = P(U_0 > x, U_1 > x, U_2 > x) \quad (9)$$

Since,

$$X_k = \min\{U_0, U_k\}$$

Therefore,

$$\begin{aligned} P(X_1 > x, X_2 > x) &= P(U_0 > x, U_1 > x, U_2 > x) \\ &= \left(1 + \frac{x - \mu}{\sigma}\right)^{-\alpha_0} \left(1 + \frac{x - \mu}{\sigma}\right)^{-\alpha_1} \left(1 + \frac{x - \mu}{\sigma}\right)^{-\alpha_2} \\ &= \left(1 + \frac{x - \mu}{\sigma}\right)^{-(\alpha_0 + \alpha_1 + \alpha_2)} \end{aligned}$$

A.3

It is easy to show that maximum likelihood estimate of location parameter is $\mu = \min\{X_1, X_2, \dots, X_n\}$. Rest two equations can be obtained by plug-in the estimates of μ and taking derivative of Log-likelihood with respect to σ and α . Therefore the details of the proof is omitted.

.....

References

- Asimit, A. V., E. Furman, and R. Vernic (2010). On a multivariate pareto distribution. *Insurance: Mathematics and Economics* 46(2), 308–316. 1, 2
- Asimit, A. V., E. Furman, and R. Vernic (2016). Statistical inference for a new class of multivariate pareto distributions. *Communications in Statistics-Simulation and Computation* 45(2), 456–471. 1, 2
- Asimit, A. V., R. Vernic, and R. Zitikis (2013). Evaluating risk measures and capital allocations based on multi-losses driven by a heavy-tailed background risk: The multivariate pareto-ii model. *Risks* 1(1), 14–33.
- Balakrishnan, N., R. C. Gupta, D. Kundu, V. Leiva, and A. Sanhueza (2011). On some mixture models based on the birnbaum–saunders distribution and associated inference. *Journal of Statistical Planning and Inference* 141(7), 2175–2190.
- Block, H. W. and A. Basu (1974). A continuous bivariate exponential extension. *Journal of the American Statistical Association* 69(348), 1031–1037.
- Campbell, J. I. and S. Austin (2002). Effects of response time deadlines on adults strategy choices for simple addition. *Memory & Cognition* 30(6), 988–994.
- D, K. (2012). On sarhan-balakrishnan bivariate distribution. *Journal of Statistics Applications & Probability* 1(3), 163.
- Dey, A. K. and D. Kundu (2012). Discriminating between bivariate generalized exponential and bivariate weibull distributions. *Chilean Journal of Statistics* 3(1), 93–110. 2
- Falk, M. and A. Guillo (2008). Peaks-over-threshold stability of multivariate generalized pareto distributions. *Journal of Multivariate Analysis* 99(4), 715–734. 14
- fbvpot by Chris Ferro, A. S. F. (2015). *evd : Functions for extreme value distributions*. 14
- Gupta, R. D. and D. Kundu (1999). Theory & methods: Generalized exponential distributions. *Australian & New Zealand Journal of Statistics* 41(2), 173–188.
- Hanagal, D. D. (1996). A multivariate pareto distribution. *Communications in Statistics-Theory and Methods* 25(7), 1471–1488. 2
- Khosravi, M., D. Kundu, and A. Jamalizadeh (2015). On bivariate and a mixture of bivariate birnbaum–saunders distributions. *Statistical Methodology* 23, 1–17.
- Kundu, D. and A. K. Dey (2009). Estimating the parameters of the marshall–olkin bivariate weibull distribution by em algorithm. *Computational Statistics & Data Analysis* 53(4), 956–965. 2
- Kundu, D. and R. D. Gupta (2009). Bivariate generalized exponential distribution. *Journal of Multivariate Analysis* 100(4), 581–593. 2
- Kundu, D. and R. D. Gupta (2010). A class of absolutely continuous bivariate distributions. *Statistical Methodology* 7(4), 464–477. 2
- Kundu, D., A. Kumar, and A. K. Gupta (2015). Absolute continuous multivariate generalized exponential distribution. *Sankhya B* 77(2), 175–206.
- Marshall, A. W. (1996). Copulas, marginals, and joint distributions. *Lecture Notes-Monograph Series*, 213–222.
- Marshall, A. W. and I. Olkin (1967). A multivariate exponential distribution. *Journal of the American Statistical Association* 62(317), 30–44. 1, 2
- McLachlan, G. and D. Peel (2004). *Finite mixture models*. John Wiley & Sons.
- Mirhosseini, S. M., M. Amini, D. Kundu, and A. Dolati (2015). On a new absolutely continuous bivariate generalized exponential distribution. *Statistical Methods & Applications* 24(1), 61–83.
- Nelsen, R. B. (2007). *An introduction to copulas*. Springer Science & Business Media. 2
- Rakonczai, P. and A. Zempléni (2012). Bivariate generalized pareto distribution in practice: models and estimation. *Environmetrics* 23(3), 219–227.
- Ristić, M. M. and D. Kundu (2015). Marshall-olkin generalized exponential distribution. *Metron* 73(3), 317–333.
- Sarhan, A. M. and N. Balakrishnan (2007). A new class of bivariate distributions and its mixture. *Journal of Multivariate Analysis* 98(7), 1508–1527. 2
- Smith, R. L. (1994). Multivariate threshold methods. In *Extreme Value Theory and Applications*, pp. 225–248. Springer.
- Smith, R. L., J. A. Tawn, and S. G. Coles (1997). *evd : Functions for extreme value distributions*. <https://cran.r-project.org> 84(2), 249–268.
- Tsay, R. S. (2014). *An introduction to analysis of financial data with R*. John Wiley & Sons.

-
- Yeh, H. C. (2000). Two multivariate pareto distributions and their related inferences. *Bulletin of the Institute of Mathematics, Academia Sinica*. 28(2), 71–86. [2](#)
- Yeh, H.-C. (2004). Some properties and characterizations for generalized multivariate pareto distributions. *Journal of Multivariate Analysis* 88(1), 47–60. [2](#)