

REGULAR VERSUS SINGULAR ORDER OF CONTACT ON PSEUDOCONVEX HYPERSURFACES

J.D. MCNEAL & L. MERNIK

ABSTRACT. The singular and regular type of a point on a real hypersurface \mathcal{H} in \mathbb{C}^n are shown to agree when the regular type is strictly less than 4. If \mathcal{H} is pseudoconvex, we show they agree when the regular type is 4. A non-pseudoconvex example is given where the regular type is 4 and the singular type is infinite.

1. INTRODUCTION

Let $\mathcal{H} \subset \mathbb{C}^n$ be a smooth, real hypersurface. D'Angelo introduced, [4], a measurement of the holomorphic flatness of \mathcal{H} at a point $p \in \mathcal{H}$. Let \mathcal{S} denote the set of parameterized non-constant holomorphic curves $\gamma : V \rightarrow \mathbb{C}^n$ with $\gamma(0) = p$, where $V \subset \mathbb{C}$ is an unspecified neighborhood of 0. Consider a local defining function r for \mathcal{H} : for some neighborhood U of p , $\mathcal{H} \cap U = \{z \in U : r(z) = 0\}$ and $dr \neq 0$ in U . Let $\nu(r \circ \gamma) = \nu(r \circ \gamma)(0)$ denote the order of vanishing of the real-valued function $r \circ \gamma$ at 0. Also let $\nu(\gamma)$ denote the multiplicity of γ at 0, i.e. the unique $M \in \mathbb{Z}^+$ such that

$$\lim_{t \rightarrow 0} \frac{\gamma(t) - \gamma(0)}{t^M} \text{ exists and is } \neq (0, \dots, 0).$$

Let $\mathcal{R} \subset \mathcal{S}$ denote the set of curves with $\nu(\gamma) = 1$.

Definition 1.1. For a given $\gamma \in \mathcal{S}$, the order of contact of γ with \mathcal{H} at p is the (possibly infinite) quantity $O(\gamma; p) = \frac{\nu(r \circ \gamma)}{\nu(\gamma)}$. The singular type of $p \in \mathcal{H}$ is defined

$$(1.2) \quad \Delta_1(p) = \sup_{\gamma \in \mathcal{S}} O(\gamma; p).$$

If $\Delta_1(p) < \infty$, p is of finite singular type.

The regular type of $p \in \mathcal{H}$ is defined

$$(1.3) \quad \Delta_1^{reg}(p) = \sup_{\gamma \in \mathcal{R}} \nu(r \circ \gamma).$$

If $\Delta_1^{reg}(p) < \infty$, p is of finite regular type.

The purpose of this paper is to prove

Theorem 1.4. Let $\mathcal{H} \subset \mathbb{C}^n$ be a smooth real hypersurface and $p \in \mathcal{H}$.

- (i) If $\Delta_1^{reg}(p) \leq 3$, then $\Delta_1(p) \leq 3$ and $\Delta_1(p) = \Delta_1^{reg}(p)$.
- (ii) If \mathcal{H} is pseudoconvex near p and $\Delta_1^{reg}(p) = 4$, then $\Delta_1(p) = 4$.

2010 *Mathematics Subject Classification.* 32W05.

Research of the first author was partially supported by a National Science Foundation grant.

The ratio defining $O(\gamma; p)$ implies this quantity is unchanged if the parameter variable t is changed, e.g., to $t^k, k \in \mathbb{Z}^+$. The subscript in $\Delta_1(p)$ indicates that orders of contact with 1-dimensional curves are considered. There are analogous higher-type conditions, denoted $\Delta_q(p)$ for $q = 2, 3, \dots, n$, discussed in [5], but these conditions are more complicated to define. Finally, $\Delta_1(p)$ does not depend on the choice of defining function r ; see [5], Proposition 5 on page 114 for a proof.

The hypersurface \mathcal{H} can be viewed as the boundary $b\Omega$ of a smoothly bounded domain $\Omega \subset \mathbb{C}^n$. For function theory, $\Delta_1(p)$ is important because it controls quantitative behavior of holomorphic functions on Ω near p ; see [5] and its bibliography for results of this kind. Many analytic estimates in terms of $\Delta_1(p)$ have been obtained, but the story is far from complete. A remarkable result is obtained in [2]: on a smoothly bounded pseudoconvex domain, $\Delta_1(p) < \infty$ for $p \in b\Omega$ is a necessary and sufficient condition for a subelliptic estimate on the $\bar{\partial}$ -Neumann problem to hold near p .

These connections with function theory motivate studying type but are unrelated to computing either $\Delta_1(p)$ or $\Delta_1^{reg}(p)$. Both quantities depend solely on the germ at p of the hypersurface $\{r = 0\}$.

Since $\mathcal{R} \subset \mathcal{S}$, $\Delta_1^{reg}(p) \leq \Delta_1(p)$ for any hypersurface. In general no other relationship between $\Delta_1^{reg}(p)$ and $\Delta_1(p)$ holds. For instance, consider the surface in \mathbb{C}^3 defined by

$$r(z) = \operatorname{Re} z_1 + |z_2^2 - z_3^3|^2$$

near the origin. This example was considered in [3]. It is straightforward to check that $\Delta_1^{reg}(0) = 6$. But the curve $t \rightarrow (0, t^3, t^2)$ is contained in $\{r = 0\}$, so $\Delta_1(0) = \infty$.

However there are hypotheses on \mathcal{H} that imply $\Delta_1^{reg}(p) = \Delta_1(p)$. When $\mathcal{H} \subset \mathbb{C}^2$ this identity holds; see Theorem 9 on page 142 of [5]. In Proposition 4.17 of Section 4, we prove the measurements agree if $\Delta_1^{reg}(p)$ equals 2 or 3. When \mathcal{H} is pseudoconvex, the case $\Delta_1^{reg}(p) = 2$ was known previously; see [6]. Results for more degenerate situations, i.e. when $\Delta_1^{reg}(p) > 3$, are unknown except for one class of hypersurfaces: it was shown in [8] that $\Delta_1^{reg}(p) = \Delta_1(p)$, regardless of the size of $\Delta_1^{reg}(p)$, if \mathcal{H} locally bounds a *convex* domain. A geometric proof and generalization of this fact is given in [1].

In [5], page 148 in Proposition 3, it is stated that $\Delta_1^{reg}(p) = \Delta_1(p)$ if $\Delta_1^{reg}(p) = 4$ on an arbitrary smooth real hypersurface. This turns out to be incorrect – see Section 3 for a counterexample. The main point of Theorem 1.4 is that with the additional hypothesis of pseudoconvexity the conclusion is correct.

In view of [8] and Theorem 1.4 (ii), there may be conditions – intermediate between convexity and pseudoconvexity – that imply $\Delta_1^{reg}(p) = \Delta_1(p)$ for points of type higher than 4. See Remark 4.23 for the obstructions to be controlled.

2. NOTATION

For curves $\gamma \in \mathcal{S}$, let $t \in \mathbb{C}$ denote the parameter variable and write $\gamma(t) = (\gamma^1(t), \dots, \gamma^n(t))$ to indicate components.

Derivatives will be denoted in several ways. For functions defined on \mathbb{C}^n or \mathbb{C} , subscripts will denote derivatives for small number of derivatives, e.g. r_{z_j} or γ_t^k . For higher derivatives, the notation will depend on which function is differentiated.

For the function $r \circ \gamma$,

$$(2.1) \quad D^{a,b}[r \circ \gamma](t) = \frac{\partial^a}{\partial t^a} \frac{\partial^b}{\partial \bar{t}^b} (r \circ \gamma)(t)$$

will distinguish t and \bar{t} derivatives. For the vector-valued function γ , the notation

$$\left(\partial^k \gamma\right)(t) = \left(\frac{\partial^k}{\partial t^k} \gamma^1(t), \dots, \frac{\partial^k}{\partial t^k} \gamma^n(t)\right) \quad \text{and} \quad \left(\bar{\partial}^k \bar{\gamma}\right)(t) = \left(\frac{\partial^k}{\partial \bar{t}^k} \bar{\gamma}^1(t), \dots, \frac{\partial^k}{\partial \bar{t}^k} \bar{\gamma}^n(t)\right)$$

is used.

When differentiating the defining function r , the following notation will be used. Let $a, b \in \mathbb{Z}^+$. Let $X^k(t) = (X_1^k(t), \dots, X_n^k(t))$, $1 \leq k \leq a$, and $Y^\ell(t) = (Y_1^\ell(t), \dots, Y_n^\ell(t))$, $1 \leq \ell \leq b$, be given vector-valued functions. Set

$$(2.2) \quad \nabla^{a,b}[r] \left(X^1, \dots, X^a, Y^1, \dots, Y^b \right) (t) = \sum_{j_1=1}^n \cdots \sum_{j_a=1}^n \sum_{k_1=1}^n \cdots \sum_{k_b=1}^n r_{z_{j_1} \dots z_{j_a} \bar{z}_{k_1} \dots \bar{z}_{k_b}}(\gamma(t)) X_{j_1}^1(t) \dots X_{j_a}^a(t) Y_{k_1}^1(t) \dots Y_{k_b}^b(t)$$

as a definition of the left-hand side. The symbol $\nabla^{a,b}[r]$ is assigned no independent meaning; it will appear only via the action described by (2.2).

Standard multi-index notation is also used. For $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbb{Z}^+)^n$ and h a function of $z \in \mathbb{C}^n$,

$$\left(\frac{\partial}{\partial z}\right)^\alpha h(z) = \left(\frac{\partial}{\partial z_1}\right)^{\alpha_1} \cdots \left(\frac{\partial}{\partial z_n}\right)^{\alpha_n} h(z);$$

$\left(\frac{\partial}{\partial \bar{z}}\right)^\alpha$ is defined analogously.

When displaying derivatives, the underlying variables – always $z \in \mathbb{C}^n$, $t \in \mathbb{C}$, or $\gamma(t) \in \mathbb{C}^n$ – will be suppressed for notational economy. If variables do not appear in an equation, our implied meaning is that the equation holds functionally. To avoid confusion with this, evaluation of derivative expressions at 0 will be explicitly indicated.

3. AN EXAMPLE

Consider the domain $\Omega = \{z \in \mathbb{C}^3 : r(z) < 0\}$ defined by

$$(3.1) \quad r(z_1, z_2, z_3) = \operatorname{Re}(z_1) + |z_2|^2 \operatorname{Re}(z_2^2 - z_3^3) + |z_3|^2 \operatorname{Re}(z_3^2) - \operatorname{Re}(z_2^2 \bar{z}_3).$$

Note that $p = (0, 0, 0) \in b\Omega$.

Claim 1. $\Delta_1(p) = \infty$.

Proof. Consider the curve $\gamma(t) = (0, t^3, t^2)$, $t \in \mathbb{C}$. Then

$$\begin{aligned} (r \circ \gamma)(t) &= 0 + |t^3|^2 \operatorname{Re}(t^6 - t^6) + |t^2|^2 \frac{t^4 + \bar{t}^4}{2} - \frac{t^6 \bar{t}^2 + t^2 \bar{t}^6}{2} \\ &= \frac{1}{2} |t|^4 (t^4 + \bar{t}^4) - \frac{1}{2} |t|^4 (t^4 + \bar{t}^4) = 0. \end{aligned}$$

Thus (the image of) γ lies in $b\Omega$, so $\Delta^1(p) = \infty$. □

Claim 2. $\Delta_1^{reg}(p) = 4$.

Proof. First consider the non-singular curve $\tilde{\gamma}(t) = (0, t, 0)$. Since $(r \circ \tilde{\gamma})(t) = |t|^2 \operatorname{Re}(t^2)$, $\nu(r \circ \tilde{\gamma}) = 4$ and so $\Delta_1^{reg}(p) \geq 4$. We now show $\nu(r \circ \gamma)(0) \leq 4$ for any nonsingular curve $\gamma(t) = (\gamma^1(t), \gamma^2(t), \gamma^3(t))$. Without loss of generality, assume that γ^1 is identically 0 (see Subsection 5.2).

Case 1: $\nu(\gamma^2) = \nu(\gamma^3) = 1$.

In this case, there are two degree 3 terms in $r \circ \gamma$, $t^2 \bar{t}$ and $t \bar{t}^2$, coming from the last term on the right-hand side of (3.1). All other terms in $r \circ \gamma$ are of higher order. Thus $\nu(r \circ \gamma) = 3$, so these curves are irrelevant for $\Delta_1^{reg}(0)$.

For the other cases, a slight rewrite of r is useful:

$$(3.2) \quad r(z) = \operatorname{Re}(z_1) + \frac{1}{2} z_2^3 \bar{z}_2 + \frac{1}{2} z_2 \bar{z}_2^3 - \frac{1}{2} |z_2|^2 (z_3^3 + \bar{z}_3^3) + |z_3|^2 \operatorname{Re}(z_3^2) - \operatorname{Re}(z_2^2 \bar{z}_3).$$

Case 2: $\nu(\gamma^2) = 1$ and $\nu(\gamma^3) > 1$.

In this case, terms $t^3 \bar{t}$ and $t \bar{t}^3$ appear in $r \circ \gamma$, because of the 2nd and 3rd terms on the right-hand side of (3.2). Terms coming from the 4th and 5th term in (3.2) will either be identically 0 or have degree ≥ 8 , by assumption on $\nu(\gamma)$. The last term on the RHS of (3.2) can produce degree 4 terms, but such terms would have *even* holomorphic degree in t or *even* anti-holomorphic degree in \bar{t} , so cannot cancel the terms $t^3 \bar{t}$ and $t \bar{t}^3$.

Thus, $D^{3,1}[r \circ \gamma](0) \neq 0$ for these γ , so $\nu(r \circ \gamma) \leq 4$.

Case 3: $\nu(\gamma^2) > 1$ and $\nu(\gamma^3) = 1$.

Similar reasoning applies to this case. Again there is a $t^3 \bar{t}$ term in $r \circ \gamma$ (and also a $t \bar{t}^3$ term), this time arising from the 5th term on the right-hand side of (3.2). No other degree 4 terms are possible in this case, since we are assuming $\nu(\gamma_2) > 1$.

Thus, $D^{3,1}[r \circ \gamma](0) \neq 0$ for these curves as well, so $\nu(r \circ \gamma) \leq 4$.

□

4. HERE COME THE WARM JETS

If $\gamma \in \mathcal{S}$, the statement $\nu(r \circ \gamma)(0) = T$ means

(i) for all $a, b \in \mathbb{Z}^+$ such that $0 \leq a + b < T$,

$$D^{a,b}[r \circ \gamma](0) = 0, \quad \text{and}$$

(ii) \exists an a_0, b_0 with $a_0 + b_0 = T$ such that

$$D^{a_0, b_0}[r \circ \gamma](0) \neq 0.$$

To prove Theorem 1.4, we examine how derivatives $D^{a,b}[r \circ \gamma]$ for γ with high multiplicity are related to lower order derivatives of $r \circ \tilde{\gamma}$, for $\tilde{\gamma} \in \mathcal{R}$ built from the jets of γ . The computations are local, near some $p \in b\Omega$; henceforth assume $p = 0$.

4.1. Automatic vanishing. Suppose $\gamma \in \mathcal{S}$ and $\nu(\gamma) = M$; the case $M = 1$ is included. Because γ is holomorphic, the chain and product rules simplify when computing $D^{a,b}[r \circ \gamma]$. The early derivatives are

$$(4.1) \quad D^{1,0}[r \circ \gamma] = \nabla^{1,0}[r](\partial^1 \gamma),$$

$$(4.2) \quad D^{1,1}[r \circ \gamma] = \nabla^{1,1}[r](\partial^1 \gamma, \bar{\partial}^1 \bar{\gamma}),$$

$$(4.3) \quad D^{2,1}[r \circ \gamma] = \nabla^{1,1}[r](\partial^2 \gamma, \bar{\partial}^1 \bar{\gamma}) + \nabla^{2,1}[r](\partial^1 \gamma, \partial^1 \gamma, \bar{\partial}^1 \bar{\gamma})$$

$$(4.4) \quad \begin{aligned} D^{2,2}[r \circ \gamma] &= \nabla^{1,1}[r](\partial^2 \gamma, \bar{\partial}^2 \bar{\gamma}) + \nabla^{1,2}[r](\partial^2 \gamma, \bar{\partial}^1 \bar{\gamma}, \bar{\partial}^1 \bar{\gamma}) \\ &\quad + \nabla^{2,1}[r](\partial^1 \gamma, \partial^1 \gamma, \bar{\partial}^2 \bar{\gamma}) \\ &\quad + \nabla^{2,2}[r](\partial^1 \gamma, \partial^1 \gamma, \bar{\partial}^1 \bar{\gamma}, \bar{\partial}^1 \bar{\gamma}) \end{aligned}$$

and

$$(4.5) \quad \begin{aligned} D^{3,1}[r \circ \gamma] &= \nabla^{1,1}[r] (\partial^3 \gamma, \bar{\partial}^1 \bar{\gamma}) + \frac{3}{2} \nabla^{2,1}[r] (\partial^2 \gamma, \partial^1 \gamma, \bar{\partial}^1 \bar{\gamma}) \\ &\quad + \nabla^{3,1}[r] (\partial^1 \gamma, \partial^1 \gamma, \partial^1 \gamma, \bar{\partial}^1 \bar{\gamma}). \end{aligned}$$

Since r is real-valued, $D^{b,a}[r \circ \gamma] = \overline{D^{a,b}[r \circ \gamma]}$. Thus (4.1)–(4.5) contain all derivative information about $r \circ \gamma$ of total order ≤ 4 except for the pure derivatives $D^{a,0}[r \circ \gamma]$ with $a = 2, 3, 4$. These are easily handled. At the same time, we pick coordinates that identify the $T^{1,0}$ part of the tangent space to \mathcal{H} at 0. Choose holomorphic coordinates in a neighborhood of 0 such that

$$(4.6) \quad \begin{aligned} r_{z_1}(0) &= 1, & r_{z_j}(0) &= 0 \quad \text{for } j = 2, \dots, n \\ \left(\frac{\partial}{\partial z}\right)^\alpha r(0) &= \left(\frac{\partial}{\partial \bar{z}}\right)^\alpha r(0) = 0 & \text{for all } 2 \leq |\alpha| \leq 4M. \end{aligned}$$

Such coordinates exist by elementary analysis of the Taylor expansion of r . For example, see the proof of Lemma 3.2.2 in [7]. From (4.1) and (4.6) it follows that

$$(4.7) \quad D^{a,0}[r \circ \gamma](0) = \nabla^{1,0}[r](\partial^a \gamma)(0)$$

for $2 \leq a \leq 4M$.

Return to (4.2)–(4.5), but ignore the first term on the right-hand side of (4.4). At least one factor of $\partial^1 \gamma$ or $\bar{\partial}^1 \bar{\gamma}$ appears in each of the other terms. If $M \geq 2$, these vanish at $t = 0$. Expressions of this kind proliferate as the number of derivatives increases.

To organize these, we make the following

Definition 4.8. Let $\gamma \in \mathcal{S}$. We say a term $A_m(t) = \nabla^{a,b}[r] (\partial^{j_1} \gamma, \dots, \partial^{j_a} \gamma, \bar{\partial}^{k_1} \bar{\gamma}, \dots, \bar{\partial}^{k_b} \bar{\gamma}) (t) \in TZ_\gamma$ if $\min\{j_1, \dots, j_a, k_1, \dots, k_b\} < M$. For a general smooth function $G(t) = G(\gamma(t))$, write $G \in \mathcal{MV}$ if $G(t) = \sum (A_m(t))$ with each $A_m \in TZ_\gamma$. If G, H are two such functions, write

$$G = H \quad \text{mod } (\mathcal{MV})$$

if $G - H \in \mathcal{MV}$.

The symbol \mathcal{MV} is short for multiplicity-vanishing. If $M \geq 2$, the equations (4.2)–(4.5) can be written

$$\begin{aligned} D^{1,1}[r \circ \gamma] &= 0 \quad \text{mod } (\mathcal{MV}), \\ D^{2,1}[r \circ \gamma] &= 0 \quad \text{mod } (\mathcal{MV}), \\ D^{2,2}[r \circ \gamma] &= \nabla^{1,1}[r] (\partial^2 \gamma, \bar{\partial}^2 \bar{\gamma}) \quad \text{mod } (\mathcal{MV}), \\ D^{3,1}[r \circ \gamma] &= 0 \quad \text{mod } (\mathcal{MV}). \end{aligned}$$

4.1.1. Higher derivatives. We now compute $D^{aM,bM}[r \circ \gamma]$ for $(a,b) = (1,1), (2,1), (2,2)$ and $(3,1)$. Computation and display of the results can be done mod \mathcal{MV} , as only values at $t = 0$ are needed.

For $D^{M,M}$, (4.2) implies

$$(4.9) \quad D^{M,M}[r \circ \gamma] = D^{M-1,M-1} D^{1,1}[r \circ \gamma] = \nabla^{1,1}[r] (\partial^M \gamma, \bar{\partial}^M \bar{\gamma}) \quad \text{mod } (\mathcal{MV}).$$

The additional derivatives must all fall on γ or $\bar{\gamma}$ mod (\mathcal{MV}) . There is no mixing of $\frac{\partial}{\partial t}$ and $\frac{\partial}{\partial \bar{t}}$ derivatives since γ is holomorphic.

For $D^{2M,M}[r \circ \gamma]$, (4.3) implies

$$\begin{aligned}
D^{2M,M} [r \circ \gamma] &= D^{2M-2,M-1} \{ D^{2,1} [r \circ \gamma] \} \\
&= D^{2M-2,M-1} \{ \nabla^{1,1} [r] (\partial^2 \gamma, \bar{\partial}^1 \bar{\gamma}) + \nabla^{2,1} [r] (\partial^1 \gamma, \partial^1 \gamma, \bar{\partial}^1 \bar{\gamma}) \} \\
(4.10) \quad &= D^{2M-2,M-1} \{ A + B \},
\end{aligned}$$

where the last equality defines A and B . Many terms arise when $D^{2M-2,M-1} \{ \}$ is expanded. Focus on the “factors” of $\partial^k \gamma$ and $\bar{\partial}^\ell \bar{\gamma}$ appearing in these terms. For example, A contains two factors, one of $\partial^2 \gamma$ and one of $\bar{\partial}^1 \bar{\gamma}$, while B contains three factors, two each of $\partial^1 \gamma$ and one of $\bar{\partial}^1 \bar{\gamma}$. Call these “factors of γ or $\bar{\gamma}$ ” for short. In order for a term *not* to belong to \mathcal{MV} , each factor of γ and $\bar{\gamma}$ that appears must be differentiated at least M times.

For the $\partial/\partial \bar{t}$ derivatives, (4.10) shows all $M-1$ derivatives must fall on the single factor $\bar{\partial}^1 \bar{\gamma} \bmod (\mathcal{MV})$ in both A and B . However the $\frac{\partial}{\partial t}$ derivatives fall on multiple factors of $\partial^k \gamma$ appearing in B and (eventually) in A . For B , no $\frac{\partial}{\partial t}$ derivative may fall on $r \bmod (\mathcal{MV})$: each of the two factors $\partial^1 \gamma$ must be differentiated $M-1$ more times and there are only $2M-2$ total derivatives. For A , one $\frac{\partial}{\partial t}$ derivative may fall on r , but then the remaining derivatives must all be distributed between the newly created factor of $\partial^1 \gamma$ and the originally present $\partial^k \gamma$ factor, resulting in dual factors of $\partial^M \gamma$ at the end. Thus

$$\begin{aligned}
D^{2M,M} [r \circ \gamma] &= \nabla^{1,1} [r] (\partial^{2M} \gamma, \bar{\partial}^M \bar{\gamma}) \\
(4.11) \quad &+ E_0 \nabla^{2,1} [r] (\partial^M \gamma, \partial^M \gamma, \bar{\partial}^M \bar{\gamma}) \bmod (\mathcal{MV}).
\end{aligned}$$

E_0 is a combinatorial constant arising from the product rule; it is computed in the next section. Notice the higher-than-multiplicity derivatives $\partial^{2M} \gamma$ in (4.11). This factor may or may not be 0 for a given $\gamma \in \mathcal{S}$.

Computing the $D^{2M,2M}$ derivative is very similar, as the $\frac{\partial}{\partial t}$ and $\frac{\partial}{\partial \bar{t}}$ derivatives do not intermix. Calculation directly from (4.4) yields

$$\begin{aligned}
(4.12) \quad D^{2M,2M} [r \circ \gamma] &= \nabla^{1,1} [r] (\partial^{2M} \gamma, \bar{\partial}^{2M} \bar{\gamma}) \\
&+ F_0 \nabla^{2,1} [r] (\partial^M \gamma, \partial^M \gamma, \bar{\partial}^{2M} \bar{\gamma}) \\
&+ F_1 \nabla^{1,2} [r] (\partial^{2M} \gamma, \bar{\partial}^M \bar{\gamma}, \bar{\partial}^M \bar{\gamma}) \\
&+ F_2 \nabla^{2,2} [r] (\partial^M \gamma, \partial^M \gamma, \bar{\partial}^M \bar{\gamma}, \bar{\partial}^M \bar{\gamma}) \bmod (\mathcal{MV}).
\end{aligned}$$

The constants $F_0, F_1, F_2 \in \mathbb{Z}^+$ are also computed in subsection 4.1.2.

For the final relevant derivative, $D^{3M,M}$, a variation occurs. The surplus $\frac{\partial}{\partial t}$ derivatives allows several combinations of higher-than-multiplicity derivatives of γ to occur inside $\nabla^{2,1} [r]$. The appearance of these (potentially) non-vanishing terms is the reason $\Delta_1(p) = \Delta_1^{reg}(p)$ does not hold without additional hypothesis.

Starting from (4.5) it follows

$$\begin{aligned}
(4.13) \quad D^{3M,M} [r \circ \gamma] &= \nabla^{1,1} [r] (\partial^{3M} \gamma, \bar{\partial}^M \bar{\gamma}) \\
&+ G_0 \nabla^{2,1} [r] (\partial^{2M} \gamma, \partial^M \gamma, \bar{\partial}^M \bar{\gamma}) \\
&+ \sum_{i=1}^{M-1} G_i \nabla^{2,1} [r] (\partial^{M+i} \gamma, \partial^{2M-i} \gamma, \bar{\partial}^M \bar{\gamma}) \\
&+ G_M \nabla^{3,1} [r] (\partial^M \gamma, \partial^M \gamma, \partial^M \gamma, \bar{\partial}^M \bar{\gamma}) \bmod (\mathcal{MV}),
\end{aligned}$$

for constants $G_0, \dots, G_M \in \mathbb{Z}^+$.

4.1.2. *Combinatorial constants.* The general forms of $D^{aM,bM}$ were derived starting from the corresponding lower-order derivatives (4.3)-(4.5). However the constants in these equations are most easily computed starting from $r \circ \gamma$ itself.

For a term in a derivative expression to not belong to \mathcal{MV} , each factor of γ and $\bar{\gamma}$ present must be differentiated at least M times. Consider (4.11):

$$\begin{aligned} D^{2M,M} [r \circ \gamma] &= \nabla^{1,1}[r] (\partial^{2M} \gamma, \bar{\partial}^M \bar{\gamma}) + E_0 \nabla^{2,1}[r] (\partial^M \gamma, \partial^M \gamma, \bar{\partial}^M \bar{\gamma}) \quad \text{mod } (\mathcal{MV}) \\ &= \tilde{A} + E_0 \tilde{B}. \end{aligned}$$

The factor $\partial^{2M} \gamma$ occurs when all the $\frac{\partial}{\partial t}$ derivatives of $r \circ \gamma$ fall on a single γ . This happens in only one way, so the constant in front of \tilde{A} is 1. The two factors $\partial^M \gamma$ in \tilde{B} occur when the $\frac{\partial}{\partial t}$ derivatives of $r \circ \gamma$ are distributed between two factors of γ . Distinguish these two factor of γ with different symbols: γ_1 and γ_2 . There is more than one way to reach a term of the form $\partial^M \gamma_1 \cdot \partial^M \gamma_2$, as the order of differentiation does not matter. For instance, first γ_1 can be differentiated M times, followed by differentiating γ_2 M times, or γ_2 could first be differentiated M times, followed by M derivatives of γ_1 , as well as all intermediate options. The term \tilde{B} contains the sum of such terms, hence $E_0 \neq 1$.

Counting how many ways we can differentiate both γ_1 and γ_2 exactly M times, using $2M$ total derivatives, is equivalent to counting all sequences of length $2M$ with M 1's and M 2's. For instance, the sequence $\underbrace{1 \dots 1}_M \underbrace{2 \dots 2}_M$ corresponds to first differentiating γ_1 M times followed by differentiating γ_2 M times. The number of all such sequences is $\frac{(2M)!}{M!M!}$. This over-counts E_0 by $2!$, as all sums in $\nabla^{2,1}$ runs from 1 to n so the roles of 1 and 2 can be exchange. Thus

$$(4.14) \quad E_0 = \frac{1}{2} \frac{(2M)!}{M!M!}.$$

The same approach computes the constants in (4.12)-(4.13), yielding

$$(4.15) \quad \begin{aligned} F_0 &= \frac{1}{2} \frac{(2M)!}{M!M!}, & F_1 &= \frac{1}{2} \frac{(2M)!}{M!M!}, & F_2 &= \frac{1}{2} \frac{(2M)!}{M!M!} \frac{1}{2} \frac{(2M)!}{M!M!}, \\ G_0 &= \frac{1}{2} \frac{(3M)!}{(2M)!M!}, & G_i &= \frac{1}{2} \frac{(3M)!}{(M+i)!(2M-i)!}, & \text{and } G_M &= \frac{1}{6} \frac{(3M)!}{M!M!M!}. \end{aligned}$$

4.2. **Connected derivatives.** We now parameterize γ . Let

$$(4.16) \quad \gamma(t) = \left(t^M \sum_{i=0}^{\infty} c_i^1 t^i, \dots, t^M \sum_{i=0}^{\infty} c_i^n t^i \right)$$

for constants $c_i^q \in \mathbb{C}$, $q = 1, \dots, n$ and $i \in \{0, 1, \dots\}$. Since $\nu(\gamma) = M$, the vector $\langle c_0^1, \dots, c_0^n \rangle \neq \langle 0, \dots, 0 \rangle$.

The subscript notation for derivatives is used in the computations below. This notation makes the interaction of the c_j^k in (4.16) – which correspond to components of $\partial^j \gamma$ – with derivatives of r easier to observe. The formulas (4.9)-(4.13) applied to γ of the form (4.16) result in cancellation of various factorials, as seen below.

When $\Delta_1^{reg}(p) < 4$, the computations in 4.1.1 show $\Delta_1(p) = \Delta_1^{reg}(p)$ for a general hypersurface.

Proposition 4.17. *Let $\mathcal{H} \subset \mathbb{C}^n$ be a smooth real hypersurface. Suppose $p \in \mathcal{H}$ and $\Delta_1^{reg}(p) \leq 3$. Then $\Delta_1(p) \leq 3$ and $\Delta_1(p) = \Delta_1^{reg}(p)$.*

Proof. The cases $\Delta_1^{reg}(p) = 2$ and $\Delta_1^{reg}(p) = 3$ will be handled separately.

First suppose that $\Delta_1(0) > 2$. Then there is a curve γ of the form (4.16) satisfying $\nu(r \circ \gamma)(0) > 2M$:

$$(4.18) \quad D^{a,b}[r \circ \gamma](0) = 0 \quad \text{for all } 0 \leq a + b \leq 2M.$$

Consider the non-singular curve $\hat{\gamma}(t) = (c_0^1 t, \dots, c_0^n t)$. It follows from (4.7) that

$$(4.19) \quad D^{M,0}[r \circ \gamma](0) = M! D^{1,0}[r \circ \hat{\gamma}](0).$$

The left-hand side vanishes by (4.18), so (4.19) says $D^{1,0}[r \circ \hat{\gamma}] = 0$. $D^{0,1}[r \circ \hat{\gamma}] = 0$ follows by conjugation. Geometrically this says $\langle c_0^1, \dots, c_0^n \rangle$ is a complex tangent vector to \mathcal{H} at 0. In the same way (4.7) shows that $D^{2,0}[r \circ \hat{\gamma}] = 0 = D^{0,2}[r \circ \hat{\gamma}]$.

It follows from (4.9) that

$$D^{M,M}[r \circ \gamma](0) = M! M! \sum_{k,j=1}^n r_{\bar{z}_k z_j} \bar{c}_0^k c_0^j.$$

(4.18) says the left-hand side vanishes. On the other hand, (4.2) implies

$$D^{1,1}[r \circ \hat{\gamma}(0)] = \sum_{k,j=1}^n r_{\bar{z}_k z_j} \bar{c}_0^k c_0^j.$$

Equating right-hand sides yields $D^{1,1}[r \circ \hat{\gamma}(0)] = 0$. Thus $\hat{\gamma} \in \mathcal{R}$ satisfies $D^{a,b}[r \circ \hat{\gamma}(0)] = 0$ for all $0 \leq a + b \leq 2$, i.e., $\Delta_1^{reg}(0) > 2$.

Turn to type 3 points. Assume γ , of the form (4.16), satisfies $\nu(r \circ \gamma)(0) > 3M$. In this case the complex line $\hat{\gamma}$ above is not the right element of \mathcal{R} . The first term in (4.11) dictates a modification. We claim that

$$\tilde{\gamma}(t) = (c_0^1 t + c_M^1 t^2, \dots, c_0^n t + c_M^n t^2)$$

satisfies $\nu(r \circ \tilde{\gamma})(0) > 3$. Note $\tilde{\gamma} \in \mathcal{R}$.

The computations that gave (4.19) show

$$D^{1,0}[r \circ \tilde{\gamma}] = D^{2,0}[r \circ \tilde{\gamma}] = D^{3,0}[r \circ \tilde{\gamma}] = 0.$$

As before, $D^{0,b}[r \circ \tilde{\gamma}] = 0$ for $b = 1, 2, 3$ because r is real-valued. Additionally, the computation giving $D^{1,1}[r \circ \hat{\gamma}] = 0$ shows $D^{1,1}[r \circ \tilde{\gamma}] = 0$.

It remains to consider $D^{2,1}[r \circ \tilde{\gamma}]$. Applying (4.11) to γ of the form (4.16) gives

$$D^{2M,M}[r \circ \gamma] = (2M)! M! \sum_{k,j=1}^n r_{\bar{z}_k z_j} \bar{c}_0^k c_M^j + \frac{1}{2} (2M)! M! \sum_{\ell,k,j=1}^n r_{z_\ell \bar{z}_k z_j} c_0^\ell \bar{c}_0^k c_0^j \quad \text{mod } (\mathcal{M}\mathcal{V}).$$

However (4.3), applied to $\tilde{\gamma}$, yields

$$D^{2,1}[r \circ \tilde{\gamma}] = \sum_{k,j=1}^n r_{\bar{z}_k z_j} \bar{c}_0^k (2c_M^j) + \sum_{\ell,k,j=1}^n r_{z_\ell \bar{z}_k z_j} c_0^\ell \bar{c}_0^k c_0^j.$$

Simple comparison shows $D^{2,1}[r \circ \tilde{\gamma}](0) = \frac{2}{(2M)! M!} D^{2M,M}[r \circ \gamma](0)$. Since $\nu(r \circ \gamma)(0) > 3M$ forces $D^{2M,M}[r \circ \gamma](0) = 0$, it follows that $D^{2,1}[r \circ \tilde{\gamma}](0) = 0$ as well.

Thus $D^{a,b}[r \circ \tilde{\gamma}](0) = 0$ for $0 \leq a + b \leq 3$, i.e., $\Delta_1^{reg}(0) > 3$.

□

Remark 4.20. If \mathcal{H} is pseudoconvex (see (5.1)), Proposition 5.16 of [6] establishes the following equivalence: $\Delta_1(p) = 2 \Leftrightarrow \Delta_1^{reg}(p) = 2 \Leftrightarrow$ the Levi form of r at p is definite. Our proof of the first equivalence does not require pseudoconvexity and is elementary. The proof in [6] is more complicated, but also connects the equivalence to subelliptic multipliers at p .

The argument in Proposition 4.17 does not extend to type 4 points, as the example in Section 3 shows. We identify where the breakdown occurs. Rewriting (4.13), for γ of the form (4.16), yields

(4.21)

$$\begin{aligned} D^{3M,M}[r \circ \gamma](0) &= (3M)!M! \sum_{j,k=1}^n r_{z_j \bar{z}_k}(0) c_{2M}^j \bar{c}_0^k + \frac{1}{2}(3M)!M! \sum_{j,k,\ell=1}^n r_{z_j z_\ell \bar{z}_k}(0) c_M^j c_0^\ell \bar{c}_0^k + \\ &\underbrace{\frac{1}{2}(3M)!M! \sum_{j,k,\ell=1}^n r_{z_j z_\ell \bar{z}_k}(0) \bar{c}_0^k \left(\sum_{i=1}^{M-1} c_i^j c_{M-i}^\ell \right)}_{B(r;\gamma)} + \frac{1}{6}(3M)!M! \sum_{j,k,\ell,m=1}^n r_{z_j z_\ell z_m \bar{z}_k}(0) c_0^j c_0^\ell c_0^m \bar{c}_0^k \end{aligned}$$

Assume that $\nu(r \circ \gamma) > 4$. The terms labeled $B(r;\gamma)$ obstruct construction of a non-singular curve with order of contact > 4 by the matching-of-derivatives argument used in Proposition 4.17. Note the presence of mixed coefficients of the components of γ^j , γ^k , and γ^ℓ (for various j, k, ℓ) appearing in $B(r;\gamma)$.

This turns out to be the only obstruction:

Lemma 4.22. *Let $\gamma \in \mathcal{S}$, $\nu(\gamma) = M$, and suppose $\nu(r \circ \gamma) > 4M$. If $B(r;\gamma) = 0$ in (4.21), then there exists $\zeta \in \mathcal{R}$ satisfying $\nu(r \circ \zeta) > 4$.*

Proof. For γ of the form (4.16), define

$$\zeta(t) = (c_0^1 t + c_M^1 t^2 + c_{2M}^1 t^3, \dots, c_0^n t + c_M^n t^2 + c_{2M}^n t^3).$$

Since $\langle c_0^1, \dots, c_0^n \rangle \neq \langle 0, \dots, 0 \rangle$, $\zeta \in \mathcal{R}$.

The same computations used in Proposition 4.17 show

$$\frac{(aM)!(bM)!}{a!b!} D^{a,b}[r \circ \zeta](0) = D^{aM,bM}[r \circ \gamma](0) = 0$$

for $0 \leq a + b \leq 3$. A computation using (4.4) and (4.12) shows that

$$\frac{(2M)!(2M)!}{2!2!} D^{2,2}[r \circ \zeta](0) = D^{2M,2M}[r \circ \gamma](0) = 0.$$

This computation is essentially the same as the one showing $M!M!D^{1,1}[r \circ \hat{\gamma}](0) = D^{M,M}[r \circ \gamma](0)$ in Proposition 4.17. Note no extra assumptions were needed to match $D^{aM,bM}$ derivatives of $r \circ \gamma$ with $D^{a,b}$ derivatives of $r \circ \zeta$ in these cases.

The remaining derivatives to consider are $D^{3,1}[r \circ \zeta](0)$ and $D^{3M,M}[r \circ \gamma](0)$. Focusing on the terms $B(r;\gamma)$ in (4.13), note the appearance of derivatives of order not a multiple of the multiplicity M . These are the problematic terms.

Use (4.5) to compute $D^{3,1}[r \circ \zeta](0)$:

$$\begin{aligned}
D^{3,1}[r \circ \zeta](0) &= \sum_{k,j} r_{\bar{z}_k z_j}(0) (3! c_{2M}^j) \bar{c}_0^k + \frac{3}{2} \sum_{\ell,k,j=1}^n r_{z_\ell \bar{z}_k z_j}(0) c_0^\ell \bar{c}_0^k (2c_M^j) \\
&\quad + \sum_{m,\ell,k,j=1}^n r_{z_j z_\ell z_m \bar{z}_k}(0) c_0^m c_0^\ell \bar{c}_0^k c_0^j.
\end{aligned}$$

Comparing this to (4.21), it follows that

$$\frac{(3M)!M!}{3!} D^{3,1}[r \circ \zeta](0) = D^{3M,M}[r \circ \gamma](0) - B(r; \gamma).$$

Therefore if $B(r; \gamma) = 0$, $D^{3,1}[r \circ \zeta](0) = 0$, since $\nu(r \circ \gamma) > 4M$. Thus $\nu(r \circ \zeta) > 4$, completing the proof. \square

Remark 4.23. When seeking an extension of Theorem 1.4 for $\Delta_1^{reg}(p) > 4$, higher order analogs of $B(r; \gamma)$ arise. For instance in the type 6 case, analogs of $B(r; \gamma)$ corresponding to derivatives $D^{4,1}, D^{5,1}, D^{4,2}, D^{3,2}$, and $D^{3,3}$ occur. Label these $B_{4,1}, B_{5,1}, B_{4,2}, B_{3,2}$, and $B_{3,3}$. In each case, $B_{a,b}$ represents the difference between $D^{aM,bM}$ derivatives of $r \circ \gamma$, where $\nu(\gamma) = M$, and $D^{a,b}[r \circ \zeta]$ for a natural non-singular curve ζ constructed from γ . If all $B_{a,b} = 0$, a straightforward modification of the proof of Lemma 4.22 yields the correspondingly modified conclusion.

5. PROOF OF THEOREM 1.4

5.1. Pseudoconvexity. Let $\mathcal{H} \subset \mathbb{C}^n$ be a smooth real hypersurface, $q \in \mathcal{H}$, and let ρ be a defining function for \mathcal{H} near q . The complex tangent space, $T^{1,0}(\mathcal{H}; q)$, to \mathcal{H} at q is defined to be all $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{C}^n$ satisfying $\sum_{j=1}^n \frac{\partial \rho}{\partial z_j}(q) \xi_j = 0$. The vector space $T^{1,0}(\mathcal{H}; q)$ is independent of the choice of defining function.

The Levi form associated to \mathcal{H} is a certain quadratic form on $T^{1,0}(\mathcal{H}; q)$; pseudoconvexity is the condition that the Levi form is semi-definite. There is an arbitrary choice of sign for this semi-definiteness, which we now fix. Say \mathcal{H} is *pseudoconvex* if there exists a defining function r of \mathcal{H} such that

$$(5.1) \quad \sum_{j,k=1}^n \frac{\partial^2 r}{\partial z_j \partial \bar{z}_k}(q) \xi_j \bar{\xi}_k \geq 0 \quad \text{for all } \xi \in T^{1,0}(\mathcal{H}; q),$$

for all $q \in \mathcal{H}$. \mathcal{H} is pseudoconvex near $p \in \mathcal{H}$ if (5.1) holds for all $q \in U \cap \mathcal{H}$, U a neighborhood of p .

Pseudoconvexity implies the existence of good holomorphic coordinates near p :

Proposition 5.2. *Let $\mathcal{H} \subset \mathbb{C}^n$ be a pseudoconvex smooth real hypersurface and $p \in \mathcal{H}$.*

There exists a defining function r for \mathcal{H} , coordinates $(z_1, z_2, \dots, z_n) = (z_1, z')$, and constants $\kappa_2, \dots, \kappa_n \geq 0$, $\lambda_{jkl} \in \mathbb{C}$, such that $p = (0, \dots, 0)$ and

$$(5.3) \quad r(z) = 2\operatorname{Re} z_1 + \sum_{j=2}^n \kappa_j |z_j|^2 + \operatorname{Re} \left(\sum_{j,k,\ell=1}^n \lambda_{jkl} z_j z_k \bar{z}_\ell \right) + \mathcal{O}(|\operatorname{Im} z_1|, |z'|^4)$$

Proof. This is generally well-known. Start with the coordinates (4.6) and an arbitrary defining function ρ for \mathcal{H} . Apply the implicit function theorem to express $\{\rho = 0\}$ as the set $\{2\operatorname{Re} z_1 = F(z', \operatorname{Im} z_1)\}$ locally near 0. A \mathbb{C} -linear rotation of coordinates allows us

to assume that $F(z', \text{Im } z_1)$ vanishes to at least order 2 at 0. Define $r(z) = 2\text{Re } z_1 - F(z', \text{Im } z_1)$.

Taylor's theorem then gives

$$\begin{aligned} r(z) &= 2\text{Re } z_1 + \sum_{j,k=2}^n \kappa_{jk} z_j \bar{z}_k + \sum_{j,k,\ell=1}^n \lambda_{jkl} z_j z_k \bar{z}_\ell + \sum_{j,k,\ell=1}^n \mu_{jkl} z_j \bar{z}_k \bar{z}_\ell + \mathcal{O}(|\text{Im } z_1|, |z'|^4) \\ &= 2\text{Re } z_1 + P_2(z', \bar{z}') + P_3(z, \bar{z}) + \mathcal{O}(|\text{Im } z_1|, |z'|^4). \end{aligned}$$

The fact that r is real-valued implies that $\overline{\mu_{jkl}} = \lambda_{jkl}$; thus $P_3(z, \bar{z})$ is of the form claimed by (5.3). The matrix (κ_{jk}) coming from $P_2(z', \bar{z}')$ is Hermitian, so can be diagonalized; let A be a matrix such that $A^* \cdot (\kappa_{jk}) \cdot A$ is diagonal. Making the linear change of coordinates $z' = Az'$ then puts $P_2(z', \bar{z}')$ in the form claimed by (5.3) – note that pseudoconvexity implies the eigenvalues of the matrix (κ_{jk}) are non-negative – and does not disturb the form of the rest of the right-hand side of (5.3). \square

Our secondary use of (5.1) is in conjunction with curves tangent to \mathcal{H} . The fact we need is proved as Proposition 2 on page 138 of [5]:

Proposition 5.4. *Let \mathcal{H} be a pseudoconvex smooth real hypersurface, $0 \in \mathcal{H}$, and r a smooth defining function for \mathcal{H} . Suppose that $\gamma \in \mathcal{S}$ satisfies*

$$\begin{aligned} 1 &< \nu(r \circ \gamma)(0) = T \\ D^{a,0}[r \circ \gamma](0) &= 0 \quad \forall a \leq T. \end{aligned}$$

Then

- (i) $T = 2K$, for some $K \in \mathbb{Z}^+$, and
- (ii) in the Taylor expansion of $r \circ \gamma(t)$, the coefficient of $|t|^{2K}$ is positive.

Proposition 5.4 is an extension of the following fact. If $u(t)$ is a smooth, subharmonic function with no pure terms in its Taylor expansion and u vanishes to finite order at 0, then the order of vanishing is even ($2K$ above) and the coefficient of $|t|^{2K}$ in the Taylor expansion of u is positive.

5.2. Type 4: pseudoconvexity implies equality. We now prove Theorem 1.4. The coordinates given by Proposition 5.2 are used throughout, in particular p is the origin.

Because of Proposition 4.17, it suffices to consider the case $\Delta_1^{reg}(0) = 4$. Suppose that $\Delta_1(0) > 4$. Then there is a curve γ of the form (4.16),

$$(5.5) \quad \gamma(t) = \left(t^M \sum_{i=0}^{\infty} c_i^1 t^i, \dots, t^M \sum_{i=0}^{\infty} c_i^n t^i \right),$$

with $\langle c_0^1, \dots, c_0^n \rangle \neq \langle 0, \dots, 0 \rangle$ and $M \geq 2$, satisfying $\nu(r \circ \gamma) > 4M$. We will show there is a curve $\zeta \in \mathcal{R}$ with $\nu(r \circ \zeta) > 4$, contradicting the assumption that $\Delta_1^{reg}(0) = 4$.

We first reduce the complexity of γ . Instead of a general curve satisfying $\nu(r \circ \gamma) > 4M$, we may consider one with *maximal* vanishing order. It follows from the proof of Theorem 1 on page 127-128 of [5] that there is such a curve, $\tilde{\gamma} = (\tilde{\gamma}^1, \dots, \tilde{\gamma}^n)$, lying in the complex tangent space $T^{1,0}(\mathcal{H}; 0)$. In the coordinates (4.6), $T^{1,0}(\mathcal{H}; 0) = \{z_1 = 0\}$, so this implies $\tilde{\gamma}^1 \equiv 0$. Thus, without loss of generality, we assume that γ is of the form

$$(5.6) \quad \gamma(t) = \left(0, t^M \sum_{i=0}^{\infty} c_i^2 t^i, \dots, t^M \sum_{i=0}^{\infty} c_i^n t^i \right),$$

at the onset.

Next we infer information about the form of r in (5.3):

Lemma 5.7. *If $\lambda_{j_0 k_0 \ell_0} \neq 0$ in (5.3), for some j_0, k_0, ℓ_0 – not necessarily distinct – then at least one of the constants $\kappa_{j_0}, \kappa_{k_0}$, or κ_{ℓ_0} must be strictly positive. (They may all be strictly positive.)*

Proof. Consider the curve $\eta : t \longrightarrow (0, \eta^2(t), \dots, \eta^n(t))$ with

$$\eta^i(t) = \begin{cases} \delta_i t, & i \in \{j_0, k_0, \ell_0\} \\ 0, & i \notin \{j_0, k_0, \ell_0\} \end{cases},$$

for constants $\delta_i \neq 0$ to be chosen. Choose δ_i such that $r \circ \eta(t)$ has a non-zero $t^2 \bar{t}$ term. Such δ_i can always be chosen as terms in $P_3(z', \bar{z}')$ involving only z_{j_0}, z_{k_0} , or z_{ℓ_0} form a non-zero polynomial. If $P_2(z', \bar{z}')$ contains no terms involving only z_{j_0}, z_{k_0} , and z_{ℓ_0} , $\nu(r \circ \eta)(0) = 3$, contradicting Proposition 5.4 (i). \square

However, a non-vanishing quadratic term in (5.3) forces higher order vanishing of that component of γ :

Lemma 5.8. *If $\kappa_s > 0$ in (5.3) for some s , and γ of the form (5.6) satisfies $\nu(r \circ \gamma) > 4M$, then*

$$c_i^s = 0 \quad \text{for } i = 0, \dots, M-1.$$

Proof. Suppose not. Let $m_0, 0 \leq m_0 \leq M-1$, be the smallest such integer with $c_{m_0}^s \neq 0$. Elementary expansion shows

$$|\gamma^s(t)|^2 = |c_{m_0}^s|^2 |t|^{2(M+m_0)} + \mathcal{O}(|t|^{2(M+m_0)+1}).$$

Thus, in the coordinate direction z_s , i.e. for the curve $\sigma(t) = (0, \dots, \gamma^s(t), \dots, 0)$, it holds that

$$\begin{aligned} P_2(\sigma(t), \bar{\sigma}(t)) &= \kappa_s |c_{m_0}^s|^2 |t|^{2(M+m_0)} + \mathcal{O}(|t|^{2(M+m_0)+1}) \\ &\geq c |t|^{2(M+m_0)} \end{aligned}$$

for $c > 0$ and $t \in \mathbb{C}$ near the origin.

However the other κ_j in (5.3) are ≥ 0 , because of pseudoconvexity. Thus the terms $\kappa_j |\gamma^j(t)|^2$ that arise cannot cancel the $\kappa_s |c_{m_0}^s|^2 |t|^{2(M+m_0)}$ term above. It follows that

$$P_2(\gamma(t), \bar{\gamma}(t)) \geq c |t|^{2(M+m_0)}.$$

The higher-order terms in (5.3) cannot interfere. Any terms coming from $P_3(z', \bar{z}')$ will have holomorphic degree $\geq 2M$ in t or anti-holomorphic degree $\geq 2M$ in \bar{t} . Terms appearing because of degree ≥ 4 terms in r will have degree $\geq 4M$. Lastly, terms involving the tangential z_1 direction need not be considered as $\gamma^1 \equiv 0$. Consequently,

$$|r \circ \gamma(t)| \geq c |t|^{2(M+m_0)}$$

as well. Since $m_0 \leq M - 1$, this says $\nu(r \circ \gamma) \leq 4M - 2$, contradicting the assumption that $\nu(r \circ \gamma) > 4M$. \square

When \mathcal{H} is pseudoconvex, it follows that $B(r; \gamma)$ in (4.21) vanishes:

Proposition 5.9. *Let $\mathcal{H} = \{r = 0\} \subset \mathbb{C}^n$ be pseudoconvex smooth real hypersurface near $0 \in \mathcal{H}$, r of the form (5.3), γ of the form (5.6) satisfying $\nu(r \circ \gamma)(0) > 4M$, and $B(r; \gamma)$ defined in (4.21).*

Then $B(r; \gamma) = 0$.

Proof. Suppose that $B(r; \gamma) \neq 0$. This implies, in particular, that there exist indices j_0, k_0, ℓ_0 – not necessarily distinct – and an integer $0 < u < M$ such that

$$r_{z_{j_0} z_{k_0} \bar{z}_{\ell_0}}(0) c_u^{j_0} c_{M-u}^{k_0} \bar{c}_0^{\ell_0} \neq 0.$$

This means that

$$(5.10) \quad \lambda_{j_0 k_0 \ell_0} \neq 0$$

in $P_3(z, \bar{z})$, as well as

$$(5.11) \quad c_u^{j_0} \neq 0, \quad c_{M-u}^{k_0} \neq 0, \quad \bar{c}_0^{\ell_0} \neq 0.$$

However, (5.10) and Lemma 5.7 force at least one of κ_{j_0} , κ_{k_0} , or κ_{ℓ_0} to be positive. For specificity, say $\kappa_{\ell_0} > 0$. Lemma 5.8 then forces $\bar{c}_0^{\ell_0} = 0$, contradicting (5.11). Thus $B(r; \gamma)$ must be 0, completing the proof. \square

Remark 5.12. The proof of Proposition 5.9 actually shows that each term in $B(r; \gamma)$ vanishes separately.

Proposition 5.9 and Lemma 4.22 finish the proof for the case $\Delta_1^{reg}(0) = 4$. Combining this case with Proposition 4.17, the proof of Theorem 1.4 is complete.

REFERENCES

- [1] BOAS, H.P. & STRAUBE, E. On equality of line type and variety type of real hypersurfaces in \mathbb{C}^n . *J. Geom. Anal.* 2, 2 (1992), 95–98.
- [2] CATLIN, D. W. Subelliptic estimates for the $\bar{\partial}$ -Neumann problem on pseudoconvex domains. *Ann. of Math. (2)* 126, 1 (1987), 131–191.
- [3] D’ANGELO, J. P. Finite type conditions for real hypersurfaces. *J. Diff. Geo.* 14 (1979), 59–66.
- [4] D’ANGELO, J. P. Real hypersurfaces, orders of contact, and applications. *Ann. of Math. (2)* 115, no. 3 (1982), 615–637.
- [5] D’ANGELO, J. P. *Several complex variables and the geometry of real hypersurfaces*. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1993.
- [6] KOHN, J. J. Subellipticity of the $\bar{\partial}$ -Neumann problem on pseudoconvex domains: Sufficient conditions. *Acta Math.* 142 (1979), 79–122.
- [7] KRANTZ, S. G. *Function Theory of Several Complex Variables, 2nd Ed.* Wadsworth & Brooks/Cole Mathematics Series. Wadsworth & Brooks/Cole, 1992.
- [8] MCNEAL, J. D. Convex domains of finite type. *J. Funct. Anal.* 108, 2 (1992), 361–373.

DEPARTMENT OF MATHEMATICS,
THE OHIO STATE UNIVERSITY, COLUMBUS, OHIO, USA
E-mail address: mcneal@math.ohio-state.edu

DEPARTMENT OF MATHEMATICS,
THE OHIO STATE UNIVERSITY, COLUMBUS, OHIO, USA
E-mail address: mernik.1@osu.edu