

On finite MTL-algebras that are representable as poset products of archimedean chains

José Luis Castiglioni and William Javier Zuluaga Botero

January 8, 2022

Abstract

We obtain a duality between certain category of finite MTL-algebras and the category of finite labeled trees. In addition we prove that certain poset products of MTL-algebras are essentially sheaves of MTL-chains over Alexandrov spaces. Finally we give a concrete description for the studied poset products in terms of direct products and ordinal sums of finite MTL-algebras.

Introduction

In [8] Esteva and Godo introduced MTL-logic as the basic fuzzy logic of left-continuous t-norms. Furthermore, a new class of algebras was defined, the variety of MTL-algebras. This variety constitutes an equivalent algebraic semantics for MTL-logic. MTL-algebras are essentially integral commutative residuated lattices with bottom satisfying the pre-linearity equation:

$$(x \rightarrow y) \vee (y \rightarrow x) \approx 1$$

This paper is divided as follows. Section 1 is devoted to present the basic contents that are necessary to understand this work. In Section 2, we characterize the finite archimedean MTL-chains in terms of their nontrivial idempotent elements. In Section 3, we show that there exist a functor from the category of finite MTL-algebras to the category of finite labeled forests. We take advantage of the intimate relation between idempotent elements and filters, that is given for the case of finite MTL-algebras. In Section 4, we study the forest products of MTL-chains. We prove that such construction is, in fact, a sheaf over an Alexandrov space whose fibers are MTL-chains. In Section 5 we use the results obtained in Section 4 in order to establish a functor from the category of finite labeled forest to the category of finite MTL-algebras. We also bring a duality theorem between the category of representable finite MTL-algebras and finite labeled forest. Finally, we present a description of the forest product of finite MTL-algebras in terms of ordinal sums and direct products of finite MTL-algebras.

1 Preliminaries

The aim of the following section is to give a brief survey about the background on MTL-algebras required to read this work. We present some known definitions and some par-

ticular constructions for semihoops that naturally can be extended to MTL-algebras.

We write **Set** to denote the category whose objects are sets and their morphisms are set functions.

A *semihoop*¹ is an algebra $\mathbf{A} = (A, \cdot, \rightarrow, \wedge, \vee, 1)$ of type $(2, 2, 2, 2, 0)$ such that (A, \wedge, \vee) is lattice with 1 as greatest element, $(A, \cdot, 1)$ is a commutative monoid and for every $x, y, z \in A$ the following conditions holds:

$$\begin{array}{ll} (\text{residuation}) & xy \leq z \text{ if and only if } x \leq y \rightarrow z \\ (\text{prelinearity}) & (x \rightarrow y) \vee (y \rightarrow x) = 1 \end{array}$$

Equivalently, a semihoop is an integral, commutative and prelinear residuated lattice.

Remark 1. *It is customary in the literature on semihoops, [15], to present them in the signature $(\cdot, \rightarrow, \wedge, 1)$, beeing \vee a defined operation. However, although less common, it is also possible to present them in the signature $(\cdot, \rightarrow, \vee, 1)$, now beeing the \wedge defined as*

$$x \wedge y := (x \cdot (x \rightarrow y)) \vee (y \cdot (y \rightarrow x)). \quad (1)$$

Let us check that the operation defined in (1) is the infimum in H .

On one hand, since $x \cdot (x \rightarrow y) \leq x \cdot 1 = x$ and $y \cdot (y \rightarrow x) \leq x$, we get $x \wedge y \leq x$. Similarly, we deduce that $x \wedge y \leq y$.

On the other hand, if we assume that $c \leq x, y$, by monotonicity, we get that $x \cdot (x \rightarrow y) \geq c \cdot (x \rightarrow y)$ and $y \cdot (y \rightarrow x) \geq c \cdot (y \rightarrow x)$. Hence, $x \wedge y \geq (c \cdot (x \rightarrow y)) \vee (c \cdot (y \rightarrow x)) = c \cdot ((x \rightarrow y) \vee (y \rightarrow x))$. Since, by prelinearity, the rightmost term of this inequality is c , we get that $x \wedge y \geq c$.

This makes $(H, \cdot, \rightarrow, \wedge, \vee, 1)$ an *integral commutative residuated lattice*. A semihoop \mathbf{A} is *bounded* if $(A, \wedge, \vee, 1)$ has a least element 0. An *MTL-algebra* is a bounded semihoop, hence, MTL-algebras are prelinear integral bounded commutative residuated lattices, as usually defined [8, 11, 15]. An MTL-algebra \mathbf{A} is an *MTL chain* if its semihoop reduct is totally ordered. Let **1** and **2** be the only MTL-chains of one and two elements, respectively. For the rest of this paper we will refer to **1** as the *trivial MTL-chain*.

It is known that the theory of MTL-algebras is a variety so we can realize the presentation of an algebraic theory. We write \mathcal{MTL} for the algebraic category of MTL-algebras.

Let $\mathbf{I} = (I, \leq)$ be a totally ordered set and $\mathcal{F} = \{\mathbf{A}_i\}_{i \in I}$ a family of semihoops. Let us assume that the members of \mathcal{F} share (up to isomorphism) the same neutral element; i.e, for every $i \neq j$, $A_i \cap A_j = \{1\}$. The *ordinal sum* of the family \mathcal{F} , is the structure $\bigoplus_{i \in I} A_i$ whose universe is $\bigcup_{i \in I} A_i$ and whose operations are defined as:

$$x \cdot y = \begin{cases} x \cdot_i y, & \text{if } x, y \in A_i \\ y, & \text{if } x \in A_i, \text{ and } y \in A_j - \{1\}, \text{ with } i > j, \\ x, & \text{if } x \in A_i - \{1\}, \text{ and } y \in A_j, \text{ with } i < j. \end{cases}$$

¹Some authors (for example [15]) name prelinear semihoop what we call here simply semihoop.

$$x \rightarrow y = \begin{cases} x \rightarrow_i y, & \text{if } x, y \in A_i \\ y, & \text{if } x \in A_i, \text{ and } y \in A_j, \text{ with } i > j, \\ 1, & \text{if } x \in A_i - \{1\}, \text{ and } y \in A_j, \text{ with } i < j. \end{cases}$$

where the subindex i denotes the application of operations in A_i .

Moreover, if \mathbf{I} has a minimum \perp , A_i is a totally ordered semihoop for every $i \in \mathbf{I}$ and A_\perp is bounded then $\bigoplus_{i \in \mathbf{I}} A_i$ becomes a MTL-chain.

Let M be an MTL-algebra. A submultiplicative monoid F of M is called a filter if it is an up set respect to the order of M . In particular, for every $x \in F$, we write $\langle x \rangle$ for the filter generated by x ; i.e.,

$$\langle x \rangle = \{a \in F \mid x^n \leq a \text{ for some } n \in \mathbb{N}\}.$$

For any filter F of M , we can define a binary relation \sim , on M by $a \sim b$ if and only if $a \rightarrow b \in F$ and $b \rightarrow a \in F$. A straightforward verification shows that \sim is a congruence on M . For every $a \in M$, we write $[a]$ for the equivalence class of a in M/F . Recall that (Section 3 of [5]) the canonical homomorphism $h : A \rightarrow A/M$ has the *universal property* of forcing all the elements of M to be 1; i.e, for every MTL-algebra B and every MTL-morphism $f : A \rightarrow B$ such that $f(a) = 1$ for every $a \in M$, there exists a unique MTL-morphism $g : A/M \rightarrow B$ making the diagram below

$$\begin{array}{ccc} A & \xrightarrow{h} & A/M \\ & \searrow f & \downarrow g \\ & & B \end{array}$$

commute. A filter F of M is *prime* if $0 \notin F$ and $x \vee y \in F$ entails $x \in F$ or $y \in F$, for every $x, y \in M$. The set of prime filters of an MTL-algebra M ordered by inclusion will be called *spectrum* and will be noted as $\text{Spec}(M)$.

2 Finite archimedean MTL-chains

In this section we bring a characterization for the archimedean finite MTL-chains in terms of their nontrivial idempotent elements. In addition we prove that every morphism of finite archimedean MTL-chains is injective.

A totally ordered MTL-algebra is said to be *archimedean* if for every $x \leq y < 1$, there exists $n \in \mathbb{N}$ such that $y^n \leq x$.

Lemma 1. *Let M be an MTL-chain. If there is an $a \in M$ such that for every $n \in \mathbb{N}$, $a^{n+1} < a^n$, then M is infinite.*

Proposition 1. *A finite MTL-chain M is archimedean if and only if $M = \mathbf{2}$ or M does not have nontrivial idempotent elements.*

Proof. If $M = \mathbf{2}$ the proof is trivial. If $M \neq \mathbf{2}$ and do not have nontrivial idempotents, there exists $a \neq 0, 1$ in M . Since M is finite, by Lemma 1, there exists $n \in \mathbb{N}$ such that $a^{n+1} = a^n$. If $a^n > 0$, $(a^n)^2 = a^n$, and hence, M has a nontrivial idempotent, in contradiction with the fact that M does not have nontrivial idempotents. Hence, there exists $n \in \mathbb{N}$ such that $a^n = 0$. Now, for $b < a$ in M , we have that $a^n \leq b$, from where we can conclude that M is archimedean. On the other hand, let us assume that M is archimedean but there exists an idempotent element $a \neq 0, 1$. Hence, $a^n = a$ for every $n \in \mathbb{N}$. If $b < a$ (for example, if $b = 0$), we have that for every $n \in \mathbb{N}$, $b < a \leq a^n$, contradicting the archimedeanity of M . In consequence, no such idempotent can exist. \square

Corollary 1. *For any finite nontrivial MTL-chain M , there are equivalent:*

- i. M is archimedean,
- ii. M is simple, and
- iii. M does not have nontrivial idempotent elements.

In [11] Horčík and Montagna gave an equational characterization for the archimedean finite MTL-chains.

Lemma 2 ([11], Lemma 6.6). *Let M be a finite MTL-chain. Then, M is archimedean if and only if for every $a, b \in M$,*

$$((a \rightarrow b) \rightarrow b)^2 \leq a \vee b.$$

The last part of this section is devoted to obtain a description of the morphisms between finite archimedean MTL-chains. Let $f : A \rightarrow B$ be a morphism of finite MTL-chains. As usual, we write K_f for the kernel of f ; i.e.,

$$K_f = \{x \in A \mid f(x) = 1\}$$

Lemma 3. *Let $f : A \rightarrow B$ be a morphism of MTL-algebras. Then f is injective if and only if $f(x) = 1$ implies $x = 1$.*

Proof. Let f be an injective morphism of MTL-algebras, then $f(x) = 1 = f(1)$ implies $x = 1$. On the other hand, let us assume that $f(x) = 1$ implies $x = 1$. If $f(a) = f(b)$ then $f(a) \leq f(b)$ and $f(b) \leq f(a)$, thus by general properties of the residual it follows that $f(a) \rightarrow f(b) = 1$ and $f(b) \rightarrow f(a) = 1$ so $f(a \rightarrow b) = 1$ and $f(b \rightarrow a) = 1$. From the assumption we get that $a \rightarrow b = 1$ and $b \rightarrow a = 1$, thus $a \leq b$ and $b \leq a$. Hence, $a = b$ so f is injective. \square

Lemma 4. *Let $f : A \rightarrow B$ be a morphism of finite MTL-chains. If A is archimedean then $B = \mathbf{1}$ or f is injective.*

Proof. Since A is archimedean, by (ii) of Corollary 1 we get that A is also simple so $K_f = A$ or $K_f = \{1\}$. In the first case, we get that $f(a) = 1$ for every $a \in A$, so in particular $f(0) = 0 = 1$, hence $B = \mathbf{1}$. In the last case, it follows that $f(a) = 1$ implies $a = 1$, so by Lemma 3 f is injective. \square

Corollary 2. *Let $f : A \rightarrow B$ be a morphism of finite MTL-chains. If A is archimedean and $B \neq \mathbf{1}$ then $f(x) = 0$ implies $x = 0$.*

Proof. Suppose that there exists $a \in A$ such that $f(a) = 0$ but $a \neq 0$. Since A is arquimedean, by Lemma 4, we get that f is injective so, from

$$f(a^2) = f(a)^2 = 0 = f(a),$$

we obtain that $a = a^2$. On the other hand, since $f(a) \neq 1$, again from Lemma 4, we get that $a \neq 1$ and consequently $0 < a < 1$. So A possesses a non trivial idempotent which, by (iii) of Corollary 1, is absurd. \square

Remark 2. *Observe that every morphism of MTL-algebras between finite archimedean MTL-chains is injective. Let $f : A \rightarrow B$ be a morphism of finite arquimedean MTL-chains. Since A is archimedean, from Lemma 4, it follows that f is injective or $B = \mathbf{1}$. Since B is archimedean, by assumption, it follows, from Proposition 1, that B cannot be trivial. Hence, f must be injective.*

3 Finite labeled forests

It is a very known fact that if M is an BL-algebra, then, its dual spectrum is a forest (c.f. Proposition 6 of [19]). Such relation has been used to establish functorial correspondences before between BL-algebras and certain kind of labeled forests² (c.f. [2]). Motivated by these ideas, in this section we show that there exist a functor from the category of finite MTL-algebras to the category of finite labeled forests. To do so, we will take advantage of the intimate relation between idempotent elements and filters, that is given for the case of finite MTL-algebras. This particular condition allows us to describe the spectrum of a finite MTL-algebra in terms of its join irreducible idempotent elements, as well as characterize the quotients that result arquimedean MTL-chains.

A *forest* is a poset X such that for every $a \in X$ the set

$$\downarrow a = \{x \in X \mid x \leq a\}$$

is a totally ordered subset of X .

This definition is motivated by the following result, whose proof is similar to the dual of Proposition 6 of [19].

Lemma 5. *Let M be a (finite) MTL-algebra. Then $\text{Spec}(M)^{op}$ is a (finite) forest.*

A *tree* is a forest with a least element. A *p-morphism* is a morphism of posets $f : X \rightarrow Y$ satisfying the following property: given $x \in X$ and $y \in Y$ such that $y \leq f(x)$ there exists $z \in X$ such that $z \leq x$ and $f(z) = y$. Let $f\mathcal{MTL}$ be the algebraic category of finite MTL-algebras. We write $fa\mathcal{MTL}$ for the algebraic category of finite archimedean MTL-algebras and $fa\mathcal{MTL}c$ for the full subcategory of finite archimedean MTL-chains. Let

²Actually in [2] the authors use the name *weighted* instead of labeled.

\mathfrak{S} be the skeleton of $fa\mathcal{MTLc}$. A *labeled forest* is a function $l : F \rightarrow \mathfrak{S}$, such that F is a forest and the collection of archimedean MTL-chains $\{l(i)\}_{i \in F}$ (up to isomorphism) shares the same neutral element 1. Consider two labeled forests $l : F \rightarrow \mathfrak{S}$ and $m : G \rightarrow \mathfrak{S}$. A morphism $l \rightarrow m$ is a pair (φ, \mathcal{F}) such that $\varphi : F \rightarrow G$ is a p-morphism and $\mathcal{F} = \{f_x\}_{x \in F}$ is a family of injective morphisms $f_x : (m \circ \varphi)(x) \rightarrow l(x)$ of MTL-algebras.

Let $(\varphi, \mathcal{F}) : l \rightarrow m$ and $(\psi, \mathcal{G}) : m \rightarrow n$ be two morphism between labeled forests. We define the composition $(\varphi, \mathcal{F})(\psi, \mathcal{G}) : l \rightarrow n$ as the pair $(\psi\varphi, \mathcal{M})$, where \mathcal{M} is the family whose elements are the MTL-morphisms $f_x g_{\varphi(x)} : n(\psi\varphi)(x) \rightarrow l(x)$ for every $x \in F$. We will call $f\mathcal{LF}$ the category of labeled forests and its morphisms. The details of checking that $f\mathcal{LF}$ is a category are left to the reader.

Let M be an MTL-algebra. We write $\mathcal{I}(M)$ for the poset of idempotent elements of M ; i.e.,

$$\mathcal{I}(M) := \{x \in M \mid x^2 = x\}.$$

Lemma 6. *In any MTL-algebra M , there are equivalent,*

i. $a \in \mathcal{I}(M)$, and

ii. $\langle a \rangle = \uparrow a$

Proof. Let us assume that $\langle a \rangle = \uparrow a$. Since $a^2 \in \langle a \rangle$ then $a^2 \in \uparrow a$, so $a \leq a^2$. Finally, since M is negatively ordered, $a^2 \leq a$. Therefore $a^2 = a$. The last part of the proof follows directly from the definition. \square

Corollary 3. *Let M be a finite MTL-algebra and $F \subseteq M$ a filter in M . There exists a unique $a \in \mathcal{I}(M)$ such that $F = \uparrow a$.*

Proof. Since M is finite, every filter $F \subseteq M$ is principal, so by Lemma 6, $F = \uparrow a$ for some $a \in \mathcal{I}(M)$. If there exists $a' \in \mathcal{I}(M)$ such that $\uparrow a = \uparrow a'$, then $a \leq a'$ and $a' \leq a$. \square

Let M be a finite MTL-algebra. From the Corollary 3, it follows that there is a bijection between $\mathcal{I}(M)$ and the filters of M . Let $\mathcal{J}(\mathcal{I}(M))$ the subposet of join irreducible elements of $\mathcal{I}(M)$. A direct application of Birkhoff's duality brings the following result.

Corollary 4. *Let M be a finite MTL-algebra and $P \in \text{Spec}(M)$. Then, there exists a unique $e \in \mathcal{J}(\mathcal{I}(M))$ such that $P = \uparrow e$.*

Corollary 5. *Let M be a finite MTL-algebra. The posets $\text{Spec}(M)^{op}$ and $\mathcal{J}(\mathcal{I}(M))$ are isomorphic.*

Proof. Let $\varphi : \mathcal{J}(\mathcal{I}(M)) \rightarrow \text{Spec}(M)$ be the mapping defined as $\varphi(e) = \uparrow e$. From Corollary 4, it follows that φ is bijective. The proofs of the antimonotonicity of φ and φ^{-1} are straightforward. \square

Lemma 7. *Let M be a finite MTL-algebra and $x \in \mathcal{I}(M)$ such that $x \neq 0$. If there exists some $k \in \mathcal{J}(\mathcal{I}(M))$ such that $x \leq k$ then x is join irreducible.*

Proof. Suppose that $0 < x \leq k$ for some $k \in \mathcal{J}(\mathcal{I}(M))$, with $x \in \mathcal{I}(M)$, then $\uparrow k \subseteq \uparrow x$. Suppose that $x \leq a \vee b$ but $x \not\leq a, b$ for some $a, b \in M$, then $a, b \notin \uparrow x$ implies that $a, b \notin \uparrow k$ so $k \not\leq a \vee b$, since $k \in \mathcal{J}(\mathcal{I}(M))$. Thereby $x \not\leq a \vee b$, which is absurd, by assumption. \square

Remark 3. Let M be a finite MTL-algebra. Observe that, from Lemmas 1.2.8 and 6.1.2 of [21] it follows that $M/\uparrow e$ is a MTL-chain if and only if $e \in \mathcal{J}(\mathcal{I}(M))$.

We write $m(M)$ for the minimal elements of $\mathcal{J}(\mathcal{I}(M))$.

Lemma 8. Let M be an MTL-algebra and $e \in \mathcal{J}(\mathcal{I}(M))$. Then, there exists a unique $k \in \mathcal{J}(\mathcal{I}(M)) \cup \{0\}$ such that $k \prec e$, where \prec denotes the covering relation in posets.

Proof. Let $e \in \mathcal{J}(\mathcal{I}(M))$, then either $e \in m(M)$ or $e \notin m(M)$. In the first case, the result follows, since $0 \prec e$. In the second case, by Lemma 5 and Corollary 4 we get that $\downarrow e \cap \mathcal{J}(\mathcal{I}(M))$ is a finite chain. If we consider k as the coatom of the latter chain, the result holds. \square

Let $e \in \mathcal{J}(\mathcal{I}(M))$. In the following, we will write a_e to denote the join irreducible element associated to e in Lemma 8. Note that $a_e = 0$ if and only if $e \in m(M)$.

Lemma 9. Let M be a finite MTL-algebra and $e \in \mathcal{J}(\mathcal{I}(M))$. Then $M/\uparrow e$ is archimedean if and only if $e \in m(M)$.

Proof. Suppose $e \in m(M)$. From Remark 3, we get that $M/\uparrow e$ is an MTL-chain. Now, if $M/\uparrow e$ is not archimedean, by Proposition 1, there exists some $[k] \in \mathcal{I}(M/\uparrow e)$ such that $[k]^2 = [k^2] = [k]$ with $[k]$ different than $[0]$ and $[1]$. So, since $ek^2 \leq k$, $ek \leq ek^2$ and $e \in \mathcal{I}(M)$ we get that $(ek)^2 = ek^2 = ek$. Thus $ek \in \mathcal{I}(M)$. Observe that due to $[k] \neq 0$ we obtain that $ek \neq 0$. Since $ek \leq e$, from Lemma 7 we get that $ek \in \mathcal{J}(\mathcal{I}(M))$. In consequence, $ek \leq e$, which is absurd because $e \in m(M)$.

On the other hand, let assume that $M/\uparrow e$ is archimedean. If there is some $k \in \mathcal{J}(\mathcal{I}(M))$ such that $k \leq e$. Thus $[k] \in \mathcal{I}(M/\uparrow e)$. So by Lemma 1, $[k] = [0]$ or $[k] = [1] = [e]$. If $[k] = [0]$ we get that $ek \leq 0$ then $k \leq e \rightarrow 0$. Since $k \in \mathcal{I}(M)$ and the product is monotone, $k \leq e(e \rightarrow 0) \leq 0$. Thereby $k = 0$, which is absurd because $k \in \mathcal{J}(\mathcal{I}(M))$. In the case of $[k] = [e]$ we obtain that $e \leq k$. Since $k \leq e$ by assumption then we conclude that $e = k$. Therefore, $e \in m(M)$. \square

Remark 4. Let M be a finite MTL-algebra and $F \subseteq M$ a filter. Let us check that $(F, \vee, \wedge, 1, x)$ is an MTL-algebra such that $0_F = x$. By definition of F , $(F, \cdot, 1)$ is a commutative monoid so, if $a, b \in F$ then $ab \in F$. The integrality of M implies that $a \leq b \rightarrow a$ and $b \leq a \rightarrow b$, so since F is an up set of M then for every $a, b \in F$, we get that $a \rightarrow b, b \rightarrow a \in F$. Similarly, since $ab \leq a \wedge b$, by applying the last argument we get that $a \wedge b \in F$. The proof for $a \vee b \in F$ is the same. Finally, due to Corollary 3 there exists a unique $x \in \mathcal{I}(M)$ such that $F = \uparrow x$, which is equivalent to say that $x = 0_F$.

Lemma 10. Let M be a finite MTL-algebra, then $(\uparrow a_e)/(\uparrow e)$ is an archimedean MTL chain. for every $e \in \mathcal{J}(\mathcal{I}(M))$.

Proof. Recall that by Remark 4 and Lemma 8, we get that $\uparrow a_e$ is a finite MTL-algebra whose least element is a_e . Since $a_e \prec e$, it follows that $\uparrow e$ is a proper filter of $\uparrow a_e$ with $e \in m(\uparrow a_e)$. Therefore, from Lemma 9 we get that $\uparrow a_e / \uparrow e$ is an archimedean MTL chain. \square

Let M and N be finite MTL-algebras and $f : M \rightarrow N$ a morphism of MTL-algebras. It is a known fact (c.f. [14]) that the assignments $M \mapsto \text{Spec}(M)$ and $f \mapsto \text{Spec}(f) = f^{-1}$, determines a contravariant functor $\mathbf{Spec} : f\mathcal{MTL} \rightarrow f\mathbf{Coh}$ from the category of finite MTL-algebras into the category of finite Coherent (or Spectral) spaces.

Let φ_M be the isomorphism between $\mathcal{J}(\mathcal{I}(M))$ and $\text{Spec}(M)^{op}$ of Proposition 5.

Lemma 11. *Let M and N be finite MTL-algebras and $f : M \rightarrow N$ an MTL-algebra morphism. There exists a unique p -morphism $f^* : \mathcal{J}(\mathcal{I}(N)) \rightarrow \mathcal{J}(\mathcal{I}(M))$ making the following diagram*

$$\begin{array}{ccc} \mathcal{J}(\mathcal{I}(N)) & \xrightarrow{f^*} & \mathcal{J}(\mathcal{I}(M)) \\ \varphi_N \downarrow & & \downarrow \varphi_M \\ \text{Spec}(N) & \xrightarrow{\text{spec}(f)} & \text{Spec}(M) \end{array}$$

commute.

Proof. Since φ_M is an isomorphism, we get that $f^* = \varphi_M^{-1} \text{spec}(f) \varphi_N$. Observe that this map is defined as $f^*(e) = \min S_e$ where $S_e = f^{-1}(\uparrow e) \cap \mathcal{J}(\mathcal{I}(M))$. In order to check the monotonicity, let $e \leq g$ in $\mathcal{J}(\mathcal{I}(N))$, then $\uparrow g \subseteq \uparrow e$, thus $f^{-1}(\uparrow g) \subseteq f^{-1}(\uparrow e)$ so $\uparrow f^*(g) \subseteq \uparrow f^*(e)$. Thereby, $f^*(e) \leq f^*(g)$. It only remains to check that f^* is a p -morphism. To do so, let $g \in \mathcal{J}(\mathcal{I}(N))$ and $e \in \mathcal{J}(\mathcal{I}(M))$ such that $g \leq f^*(e)$. Since $\mathcal{J}(\mathcal{I}(N))$ is finite, we can consider $m = \min S$, with

$$S = \{k \in \mathcal{J}(\mathcal{I}(N)) \mid k \leq e, g \leq f^*(k)\}.$$

We will prove that $f^*(m) = g$. Let $x \in \mathcal{I}(N)$ be such that $g \leq x$. Since $e \leq f^*(g)$, it follows that $f(x) \neq 0$. Consider $\uparrow mf(x)$. Since $mf(x) \leq m$, by Lemma 7 we get that $mf(x) \in \mathcal{J}(\mathcal{I}(M))$. Let us verify that $g \leq f^*(mf(x))$ by checking $f^{-1}(\uparrow mf(x)) \subseteq \uparrow g$. If $b \in f^{-1}(\uparrow mf(x))$, then $mf(x) \leq f(b)$ and hence $m \leq f(x) \rightarrow f(b) = f(x \rightarrow b)$. Therefore $x \rightarrow b \in f^{-1}(\uparrow m)$. By construction of m , we have that $g \leq f^*(m)$, so $f^{-1}(\uparrow m) \subseteq \uparrow g$. Consequently, $g \leq x \rightarrow b$. Since $g \leq x$, we obtain that $g \leq x(x \rightarrow b) \leq b$. Hence, $mf(x) = m$, because $mf(x) \in S$. Finally, since $mf(x) = m \leq f(x)$, it follows that $x \in f^{-1}(\uparrow m)$. Thus $\uparrow x \subseteq f^{-1}(\uparrow m)$ and since $x \leq g$, we obtain that $\uparrow g \subseteq f^{-1}(\uparrow m)$. Then we conclude that $f^*(m) \leq g$. This concludes the proof. \square

Let M be a finite MTL-algebra and consider the function

$$l_M : \mathcal{J}(\mathcal{I}(M)) \rightarrow \mathfrak{S}$$

defined as $l_M(e) = \uparrow a_e / \uparrow e$. Since from Lemma 5 we know that $\mathcal{J}(\mathcal{I}(M))$ is a finite forest, l_M is a finite labeled forest.

Let \mathbf{F} be a finite forest and $X \subseteq F$. We write $\text{Min}(X)$ for the minimal elements of X .

Lemma 12. *Let $f : X \rightarrow Y$ be a p-morphism. If $x \in \text{Min}(X)$ then $f(x) \in \text{Min}(Y)$.*

Proof. Let us assume $x \in \text{Min}(X)$, and suppose that there exists $y \in Y$ such that $y < f(x)$. Since f is a p-morphism, there exists $z \in X$, with $z \leq x$ such that $y = f(z)$. Since $y \neq f(x)$, then $z \neq x$, so $x \notin \text{Min}(X)$. This fact is absurd by assumption. \square

Lemma 13. *Let M and N be finite MTL-algebras and $f : M \rightarrow N$ an MTL-algebra morphism. Then, for every $e \in \mathcal{J}(\mathcal{I}(N))$, f determines a morphism $\bar{f}_e : \uparrow a_{f^*(e)} \rightarrow \uparrow a_e$ such that there exists a unique MTL-algebra morphism $f_e : \uparrow a_{f^*(e)} / \uparrow f^*(e) \rightarrow \uparrow a_e / \uparrow e$ making the diagram*

$$\begin{array}{ccc} \uparrow a_{f^*(e)} & \xrightarrow{\bar{f}_e} & \uparrow a_e \\ \downarrow & & \downarrow \\ \uparrow a_{f^*(e)} / \uparrow f^*(e) & \xrightarrow{f_e} & \uparrow a_e / \uparrow e \end{array}$$

commute.

Proof. Let $e \in \mathcal{J}(\mathcal{I}(N))$. Then $e \notin m(N)$ or $e \in m(N)$. In the first case, it follows that $a_e > 0_N$ and thus, $\uparrow a_e \subset N$. Since $a_e \leq e$ and f^* is monotone then $f^*(a_e) \leq f^*(e)$. Since $a_{f^*(e)} \prec f^*(e)$, $f^*(a_e) \leq a_{f^*(e)}$. Thereby, $\uparrow a_{f^*(e)} \subseteq \uparrow f^*(a_e)$. Let us define $\bar{f}_e : \uparrow a_{f^*(e)} \rightarrow \uparrow a_e$ as

$$\bar{f}_e(x) = \begin{cases} f(x), & x > a_{f^*(e)} \\ a_e, & x = a_{f^*(e)} \end{cases}$$

From Lemma 11, we get that $\uparrow f^*(a_e) = f^{-1}(\uparrow a_e)$ and \bar{f}_e is a well defined MTL-morphism. Let us consider $a_{f^*(e)} < f^*(e) \leq x$, then, $\bar{f}_e(a_{f^*(e)}) < \bar{f}_e(f^*(e)) \leq \bar{f}_e(x)$ since \bar{f}_e is monotone. By definition of \bar{f}_e we obtain that $a_e < f(f^*(e)) \leq f(x)$. Then, applying Lemma 11, we get that $e \leq f(f^*(e))$, so we can conclude that $e \leq f(x)$. This means that $[\bar{f}_e(x)] = [1]$ in $\uparrow a_e / \uparrow e$. Hence, by the universal property of quotients in \mathcal{MTL} , there exists a unique MTL-morphism $f_e : \uparrow a_{f^*(e)} / \uparrow f^*(e) \rightarrow \uparrow a_e / \uparrow e$ making the diagram above commutes.

Finally, if $e \in m(N)$, we get that $a_e = 0_N$ and $\uparrow a_e = N$. Since f^* is a p-morphism, due to Lemma 12, $f^*(e) \in m(M)$, $a_{f^*(e)} = 0_M$ and consequently, $\uparrow a_{f^*(e)} = M$. Let $\bar{f}_e = f$. The proof of $[\bar{f}_e(x)] = [1]$ in $\uparrow a_e / \uparrow e$ is similar to the given for first case. The rest of the proof follows from the universal property of quotients in \mathcal{MTL} . \square

Let $f : M \rightarrow N$ be an MTL-morphism between finite MTL-algebras and $\mathcal{F}_f := \{f_e\}_{e \in \mathcal{J}(\mathcal{I}(N))}$ be the family of MTL-morphisms obtained in Lemma 13.

Corollary 6. *Let M and N be finite MTL-algebras and $f : M \rightarrow N$ an MTL-algebra morphism. Then the pair (f^*, \mathcal{F}_f) is a morphism between the labeled forests l_N and l_M .*

Theorem 1. *The assignments $M \mapsto l_M$ and $f \mapsto (f^*, \mathcal{F}_f)$ define a contravariant functor*

$$\mathcal{G} : f\mathcal{MTL} \rightarrow f\mathcal{LF}.$$

Proof. Let $f : M \rightarrow N$ and $g : N \rightarrow O$ be morphisms in $f\mathcal{MTL}$ and consider the diagram

$$\begin{array}{ccccc} \text{Spec}(O) & \xrightarrow{\text{Spec}(g)} & \text{Spec}(N) & \xrightarrow{\text{Spec}(f)} & \text{Spec}(M) \\ \varphi_O \uparrow & & \varphi_N \uparrow \downarrow \varphi_N^{-1} & & \varphi_M \uparrow \\ \mathcal{J}(\mathcal{I}(O)) & \xrightarrow{g^*} & \mathcal{J}(\mathcal{I}(N)) & \xrightarrow{f^*} & \mathcal{J}(\mathcal{I}(M)) \end{array}$$

associated to the composition $gf : M \rightarrow O$. Since **Spec** is contravariant,

$$(gf)^* = \varphi_M^{-1} \text{Spec}(gf) \varphi_O = (\varphi_M^{-1} \text{Spec}(f) \varphi_N)(\varphi_N^{-1} \text{Spec}(g) \varphi_O) = f^* g^*. \quad (2)$$

If id_M denotes the identity map of the MTL-algebra M , a straightforward calculation proves that $(id_M)^* = id_{\mathcal{J}(\mathcal{I}(M))}$. On the other hand, let $e \in \mathcal{J}(\mathcal{I}(O))$. We will verify that $\overline{(gf)}_e = \overline{g}_e \overline{f}_{g^*(e)}$. From (2), we conclude that $a_{(gf)^*(e)} = a_{f^*(g^*(e))}$. If $x > a_{(gf)^*(e)}$, then $x > a_{f^*(g^*(e))}$. By the monotonicity of f and Lemma 11, we obtain that $a_{g^*(e)} \leq f(a_{f^*(g^*(e))}) \leq f(x)$. Hence $a_{g^*(e)} \leq f(x)$. In a similar way, by the monotonicity of g and using again Lemma 11, we get that $a_e \leq (gf)(x)$. Therefore, for every $x \in \uparrow a_{(gf)^*(e)}$ it follows that $\overline{(gf)}_e(x) = \overline{g}_e \overline{f}_{g^*(e)}(x)$. Finally, from Lemma 13, we obtain that $g_e f_{g^*(e)} = (gf)_e$. So, for every $e \in \mathcal{J}(\mathcal{I}(N))$ the diagram below

$$\begin{array}{ccccc} & & \overline{(gf)}_e & & \\ & \nearrow & & \searrow & \\ \uparrow a_{(gf)^*(e)} & \xrightarrow{(\overline{f})_{g^*(e)}} & \uparrow a_{g^*(e)} & \xrightarrow{\overline{g}_e} & \uparrow a_e \\ \downarrow & & \downarrow & & \downarrow \\ \uparrow a_{(gf)^*(e)}/\uparrow (gf)^*(e) & \xrightarrow{f_{g^*(e)}} & \uparrow a_{g^*(e)}/\uparrow g^*(e) & \xrightarrow{g_e} & \uparrow a_e/\uparrow e \\ & \searrow & & \nearrow & \\ & & (gf)_e & & \end{array}$$

commutes. Therefore $(g^* f^*, \mathcal{F}) = ((gf)^*, \mathcal{F}_{gf})$, where $\mathcal{F}_{gf} = \{(gf)_e\}_{e \in \mathcal{J}(\mathcal{I}(O))}$ and $\mathcal{F} = \{g_e f_{g^*(e)}\}_{e \in \mathcal{J}(\mathcal{I}(O))}$. Hence $\mathcal{G}(gf) = \mathcal{G}(f)\mathcal{G}(g)$. From the definition of \mathcal{G} it easily follows that $\mathcal{G}(id_M) = id_{\mathcal{G}(M)}$. \square

4 Forest Product of MTL-algebras

In this section we introduce the notion of forest product. It is simply a poset product as defined in [4] when restricted to posets which are forests. For the sake of completeness, we give explicitly the necessary definitions.

Definition 1. Let $\mathbf{F} = (F, \leq)$ be a forest and let $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains such that, up to isomorphism, all they share the same neutral element 1. If $(\bigcup_{i \in \mathbf{F}} \mathbf{M}_i)^F$ denotes the set of functions $h : F \rightarrow \bigcup_{i \in \mathbf{F}} \mathbf{M}_i$ such that $h(i) \in \mathbf{M}_i$ for all $i \in \mathbf{F}$, the forest product $\bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$ is the algebra \mathbf{M} defined as follows:

- (1) The elements of \mathbf{M} are the $h \in (\bigcup_{i \in \mathbf{F}} \mathbf{M}_i)^F$ such that, for all $i \in \mathbf{F}$ if $h(i) \neq 0_i$ then for all $j < i$, $h(j) = 1$.

(2) The monoid operation and the lattice operations are defined pointwise.

(3) The residual is defined as follows:

$$(h \rightarrow g)(i) = \begin{cases} h(i) \rightarrow_i g(i), & \text{if for all } j < i, h(j) \leq_j g(j) \\ 0_i & \text{otherwise} \end{cases}$$

where the subindex i denotes the application of operations and of order in \mathbf{M}_i .

The following result is a slight modification of Theorem 3.5.3 in [4].

Lemma 14. *The forest product of MTL-chains is an MTL-algebra.*

In the following if we refer to a collection $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ of MTL-chains indexed by a forest \mathbf{F} we always will assume that it satisfies the conditions of Definition 1.

Lemma 15. *Let \mathbf{F} be a forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. There are equivalent:*

1. $h \in \bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$,
2. For every $i < j$ in \mathbf{F} , $h(j) = 0_j$ or $h(i) = 1$,
3. For all $i \in \mathbf{F}$ if $h(i) \neq 1$ then for all $i < j$, $h(j) = 0_j$,
4. $\bigcup_{i \in \mathbf{F}} h^{-1}(0_j)$ is an upset of \mathbf{F} , $h^{-1}(1)$ is a downset of \mathbf{F} and

$$C_h = \{i \in \mathbf{F} \mid h(i) \notin \{0_i, 1\}\},$$

is a (possibly empty) antichain of \mathbf{F} .

Proof. Since the implications (1) \Rightarrow (2) \Rightarrow (3) follow straight from definition, we only prove the remaining implications. Let us start proving that (3) implies (4): To prove that $h^{-1}(1)$ is a downset of \mathbf{F} we proceed by contradiction. Suppose $i < j$ with $h(j) = 1$ but $h(i) \neq 1$. Thus, $h(j) = 0_j$, by assumption, which is absurd. To prove that $\bigcup_{i \in \mathbf{F}} h^{-1}(0_j)$ is an upset of \mathbf{F} let us suppose that $i < j$ with $h(i) = 0_i$, thus, since $h(i) \neq 1$, and (3), we get that $h(j) = 0_j$. If C_h is not an antichain, there exist $i, j \in C_h$ comparable. Without loss of generality, we can assume $i < j$, $h(i) \neq 1$ and $h(j) \neq 0_j$, then because of (3), we obtain that $h(j) = 0_j$, which is absurd. Finally, to prove that (4) implies (1), let $h \in (\bigcup_{i \in \mathbf{F}} \mathbf{M}_i)^{\mathbf{F}}$ and suppose that $i < j$ with $h(i) \neq 1$. If $h(j) \neq 0_j$, thus $i, j \in C_h$, which is absurd, since C_h is by assumption an antichain. \square

Remark 5. *Let \mathbf{F} be a chain and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. Let us consider $h \in \bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$ and $j \in \mathbf{F}$. Note that there are only two possible cases for $h(j)$, namely $h(j) \neq 0_j$ or $h(j) = 0_j$. If $h(j) \neq 0_j$, from (4) of Lemma 15, it follows that for every $j < i$, $h(i) = 0_i$ and due to (3) of the same Lemma, $h(k) = 1$ for every $k < j$. If $h(j) = 0_j$, from Definition 1, we get that $h(k) = 0_k$ for every $j < k$ and by (4) of Lemma 15, we have that $h(i) = 1$ for every $i < j$.*

A commutative integral residuated lattice A is called *really local* (Definition 1.2.27 of [21]) if it is not trivial ($0 \neq 1$) and for every $x, y \in A$,

$$x \vee y = 1 \Rightarrow x = 1 \text{ or } y = 1.$$

Lemma 16. *Let A be a non-trivial commutative integral residuated lattice. If the canonical order of A is total then A is really local. If A is also prelinear then the converse holds.*

Proof. A direct consequence of Lemma 13.7 of [5]. □

Lemma 17. *Let \mathbf{F} be a forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. Then \mathbf{F} is a totally ordered set if and only if $\bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$ is an MTL-chain.*

Proof. Suppose \mathbf{F} is a totally ordered set and let $g, h \in \bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$ be such that $(g \vee h) = 1$. Since the lattice operations in $\bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$ are calculated pointwise, for every $i \in \mathbf{F}$, $g(i) \vee h(i) = 1$. If $g, h \neq 1$, there exists some $j \in \mathbf{F}$ such that $g(j), h(j) \neq 1$. From Remark 5 it follows that $g(k) = h(k) = 0_k$, for every $j < k$ so we get that $(g \vee h)(k) = g(k) \vee h(k) = 0_k$, which contradicts our assumption. Hence, since every MTL-algebra is prelinear, from Lemma 16 we get that $\bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$ is a MTL-chain. On the other hand, let us assume that $\bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$ is a MTL-chain. If \mathbf{F} is not a totally ordered set, thus there exist two different elements n and m in \mathbf{F} which are not comparable. Let us consider $g, h \in \bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$, defined as

$$g(i) = \begin{cases} 0_n, & \text{if } i \geq n \\ 1, & \text{otherwise} \end{cases} \quad h(i) = \begin{cases} 0_m, & \text{if } i \geq m \\ 1, & \text{otherwise} \end{cases}$$

Observe that $g \vee h = 1$ but $g, h \neq 1$, which is a contradiction since, by Lemma 16, $\bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$ is really local. □

Let \mathbf{F} be a forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. We write $\mathcal{D}(\mathbf{F})$ for the collection of downsets of \mathbf{F} . Let S be a proper downset of \mathbf{F} and consider

$$X_S := \{h \in \bigotimes_{i \in \mathbf{F}} \mathbf{M}_i \mid h|_S = 1\}$$

Observe that X_S is a proper filter of $\bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$. Since S^c is itself a forest, due to Lemma 14, $\bigotimes_{i \in S^c} \mathbf{M}_i$ is an MTL-algebra. Using the fact that every filter of a MTL-algebra is a semihoop, we obtain the following result.

Lemma 18. *Let \mathbf{F} be a forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains and $S \in \mathcal{D}(\mathbf{F})$. Then X_S and $\bigotimes_{i \in S^c} \mathbf{M}_i$ are isomorphic semihoops.*

Proof. Let $g \in \bigotimes_{i \in S^c} \mathbf{M}_i$. Define $\varphi : \bigotimes_{i \in S^c} \mathbf{M}_i \rightarrow X_S$ as

$$\varphi(g)(i) = \begin{cases} g(i), & \text{if } i \notin S \\ 1, & \text{if } i \in S \end{cases}$$

For this part of the proof we will write h to denote $\varphi(g)$. First, we prove that h is well defined. Let us take $i < j$ in F and suppose that $h(i) \neq 1$. By construction of h , we get that $i \notin S$, so $h(i) = g(i)$. If $h(j) \neq 0_j$, then $h(j) = 1$ or $0_j < h(j) < 1$. In the first case we obtain that $j \in S$ and since $i < j$ and $S \in \mathcal{D}(\mathbf{F})$, $i \in S$, which is absurd. In the second case, since S^c is an upset of \mathbf{F} , from $i \notin S$ and $i < j$ it follows that $j \notin S$. Hence, $h(j) = g(j) \neq 1$. Therefore, there are $i, j \in C_h$ comparable, which by (4) of Lemma 15 is absurd. Consequently, $h(j) = 0_j$ and thus, by (3) of Lemma 15, we get that $h \in \bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$. By construction, it is clear that $h \in X_S$. In order to verify that φ is surjective, let $f \in X_S$ and consider $f|_{S^c}$. Since S^c is an upset and $f \in \bigotimes_{i \in \mathbf{F}} \mathbf{M}_i$, it is clear that $\varphi(f|_{S^c}) = f$. The injectivity of φ is immediate.

Since the monoid and lattice operations in X_S and $\bigotimes_{i \in S^c} \mathbf{M}_i$ are defined pointwise it is clear that φ preserve such operations. We prove that φ preserve the residual. To do so, let $s, t \in \bigotimes_{i \in S^c} \mathbf{M}_i$. Then,

$$(\varphi(s) \rightarrow \varphi(t))(i) = \begin{cases} \varphi(s)(i) \rightarrow_i \varphi(t)(i), & \text{if for all } j < i, \varphi(s)(j) \leq_j \varphi(t)(j) \\ 0_i, & \text{otherwise} \end{cases} \quad (3)$$

and

$$\varphi(s \rightarrow t)(i) = \begin{cases} (s \rightarrow t)(i), & \text{if } i \notin S \\ 1, & \text{if } i \in S \end{cases} \quad (4)$$

If $i \in S$ then $\varphi(s) = \varphi(t) = 1$ so $\varphi(s)(i) \rightarrow_i \varphi(t)(i) = 1$. Since $S \in \mathcal{D}(\mathbf{F})$, if $j < i$, $j \in S$. Therefore $\varphi(s)(j) = \varphi(t)(j) = 1$. Hence, for every $i \in S$, $\varphi(s \rightarrow t)(i) = (\varphi(s) \rightarrow \varphi(t))(i)$. On the other hand, if $i \notin S$ and $(\varphi(s) \rightarrow \varphi(t))(i) \neq 1$ then $(\varphi(s) \rightarrow \varphi(t))(i) = 0_i$ or $0_i < (\varphi(s) \rightarrow \varphi(t))(i) < 1$. In the first case, from the equation (3) it follows that there exists $j \in F$ with $j < i$ such that $\varphi(s)(j) \not\leq_j \varphi(t)(j)$. If $j \in S$, $\varphi(s)(j) = \varphi(t)(j) = 1$. Therefore $j \notin S$. Hence, since $\varphi(s)(j) = s(j)$ and $\varphi(t)(j) = t(j)$ we get that there exists $j \notin S$ with $j < i$ such that $s(j) \not\leq_j t(j)$. Then $(s \rightarrow t)(i) = 0_i = (\varphi(s) \rightarrow \varphi(t))(i)$. In the second case, from the equation (4) we get that $(\varphi(s) \rightarrow \varphi(t))(i) = \varphi(s)(i) \rightarrow_i \varphi(t)(i)$ so, by the definition of φ , we obtain that $\varphi(s)(i) \rightarrow \varphi(t)(i) = (\varphi(s) \rightarrow \varphi(t))(i)$. Finally, in the case $i \notin S$ and $(\varphi(s) \rightarrow \varphi(t))(i) = 1$, $\varphi(s)(i) \leq_i \varphi(t)(i)$ and in consequence $\varphi(s)(i) \rightarrow_i \varphi(t)(i) = 1$. Thus, for every $j < i$, $\varphi(s)(j) \leq_j \varphi(t)(j)$. In particular, if $j \notin S$ and $j < i$, $s(j) \leq_j t(j)$. Therefore $\varphi(s)(i) \rightarrow \varphi(t)(i) = (s \rightarrow t)(i)$. Hence, for every $i \notin S$, $(\varphi(s) \rightarrow \varphi(t))(i) = \varphi(s)(i) \rightarrow_i \varphi(t)(i)$. This concludes the proof. \square

Corollary 7. *Let \mathbf{F} be a forest, $S, T \in \mathcal{D}(\mathbf{F})$ such that $S \subseteq T$ and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. Take $X_S^T := \{h \in \bigotimes_{i \in T} \mathbf{M}_i \mid h|_S = 1\}$. Then, X_S^T and $\bigotimes_{i \in S^c \cap T} \mathbf{M}_i$ are isomorphic semihoops.*

Proof. Since $S \subseteq T$ and $S, T \in \mathcal{D}(\mathbf{F})$ we get that $S \in \mathcal{D}(\mathbf{T})$. The result follows from Lemma 18. \square

4.1 Forest products are sheaves

In every poset \mathbf{F} the collection $\mathcal{D}(\mathbf{F})$ of downsets of \mathbf{F} defines a topology over F called the *Alexandrov topology* on \mathbf{F} . Let $S, T \in \mathcal{D}(\mathbf{F})$ be such that $S \subseteq T$ and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ be a collection of MTL-chains. Observe that if $h \in \bigotimes_{i \in \mathbf{T}} \mathbf{M}_i$ then the restriction $h|_S$ is an element of $\bigotimes_{i \in S} \mathbf{M}_i$, so the assignment that sends $T \in \mathcal{D}(\mathbf{F})$ to $\bigotimes_{i \in \mathbf{T}} \mathbf{M}_i$ defines a presheaf $\mathcal{P} : \mathcal{D}(\mathbf{F})^{op} \rightarrow \mathcal{MTL}$.

Lemma 19. *Let \mathbf{F} be a forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. Then, for every $S \in \mathcal{D}(\mathbf{F})$*

$$\mathcal{P}(S) \cong \mathcal{P}(F)/X_S.$$

Proof. Let $r : \mathcal{P}(F) \rightarrow \mathcal{P}(S)$ be the restriction to S . It is clear that r is a surjective morphism of MTL-algebras such that $r(h) = 1$, for every $h \in X_S$. Then, by the universal property of the canonical homomorphism $\beta : \mathcal{P}(F) \rightarrow \mathcal{P}(F)/X_S$, there exists a unique morphism of MTL-algebras $\alpha : \mathcal{P}(F)/X_S \rightarrow \mathcal{P}(S)$ such that the diagram below

$$\begin{array}{ccc} \mathcal{P}(F) & \xrightarrow{\beta} & \mathcal{P}(F)/X_S \\ & \searrow r & \downarrow \alpha \\ & & \mathcal{P}(S) \end{array}$$

commutes. Observe that $\alpha\beta = r$, so since β is surjective, it follows that α is surjective too. The verification of the injectivity of α is straightforward. \square

Corollary 8. *Let \mathbf{F} be a forest, $S, T \in \mathcal{D}(\mathbf{F})$ such that $S \subseteq T$ and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. Then $\mathcal{P}(S) \cong \mathcal{P}(T)/X_S^T$.*

Proof. Due to Lemma 19, $\mathcal{P}(T) \cong \mathcal{P}(F)/X_T$. Observe that $X_S^T \cong X_S/X_T$, thus the result follows as a direct consequence of the second isomorphism theorem (Theorem 6.15 of [17]). \square

Lemma 20. *Let A be a non-trivial MTL-algebra. Then A/P is an MTL-chain if and only if P is a non trivial prime filter.*

Lemma 21. *Let \mathbf{F} be a forest, $S \in \mathcal{D}(\mathbf{F})$ and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. Then X_S is prime if and only if S is totally ordered.*

Proof. Observe that asking S to be totally ordered, is equivalent, by Lemma 17, to asking $\bigotimes_{i \in S} \mathbf{M}_i$ to be an MTL-chain. By Lemma 19, $\bigotimes_{i \in S} \mathbf{M}_i \cong \bigotimes_{i \in \mathbf{F}} \mathbf{M}_i/X_S$. Hence, the result follows from the first remark and Lemma 20. \square

Let $\mathbf{Shv}(\mathbf{P})$ be the category of sheaves over the Alexandrov space $(P, \mathcal{D}(\mathbf{P}))$. Since the theory of MTL-algebras is algebraic, it is well-known that an MTL-algebra in $\mathbf{Shv}(\mathbf{P})$ is a functor $\mathcal{D}(\mathbf{P})^{op} \rightarrow \mathcal{MTL}$ such that the composite presheaf $\mathcal{D}(\mathbf{P})^{op} \rightarrow \mathcal{MTL} \rightarrow \mathbf{Set}$ is a sheaf.

Lemma 22. *Let \mathbf{F} be a forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. The presheaf $\mathcal{P} : \mathcal{D}(\mathbf{P})^{op} \rightarrow \mathcal{MTL}$, with $\mathcal{P}(T) = \bigotimes_{i \in T} \mathbf{M}_i$, is an MTL-algebra in $\mathbf{Shv}(\mathbf{P})$.*

Proof. Suppose that $T = \bigcup_{\alpha \in I} S_\alpha$, with $S_\alpha, T \in \mathcal{D}(\mathbf{F})$, for every $\alpha \in I$, and let $h_\alpha \in \mathcal{P}(S_\alpha)$ be a matching family. Thus, for every $\alpha \neq \beta$ in I :

$$h_\alpha|_{S_\alpha \cap S_\beta} = h_\beta|_{S_\alpha \cap S_\beta} \quad (5)$$

Let us consider the following function:

$$\begin{aligned} h : I &\rightarrow \bigcup_{i \in \mathbf{T}} \mathbf{M}_i \\ i &\mapsto h_\alpha(i), \text{ if } i \in S_\alpha \end{aligned}$$

Observe that (5) guarantees that h is well defined. To check that $h \in \bigotimes_{i \in \mathbf{T}} \mathbf{M}_i$, let us suppose that $i \in T$ and $h(i) \neq 0_i$. If $j < i$ then, since $T = \bigcup_{\alpha \in I} S_\alpha$, there exists some $\beta \in I$ such that $i \in S_\beta$. In such case, $j \in S_\beta$, since $S_\beta \in \mathcal{D}(\mathbf{F})$. Then $h(i) = h_\beta(i)$ and $h(j) = h_\beta(j)$. Since $h_\beta \in \bigotimes_{i \in \mathbf{S}_\beta} \mathbf{M}_i$, we conclude that $h(j) = h_\beta(j) = 1$. Therefore h amalgamates $\{h_\alpha\}_{\alpha \in I}$. To verify the uniqueness of h , let us suppose that there exists $f \in \mathcal{P}(T)$ such that $f|_{S_\alpha} = h_\alpha$, for every $\alpha \in I$. Then,

$$(f|_{S_\alpha})|_{S_\alpha \cap S_\beta} = (h|_{S_\alpha})|_{S_\alpha \cap S_\beta} = (h|_\beta)|_{S_\alpha \cap S_\beta} = (f|_\beta)|_{S_\alpha \cap S_\beta}.$$

For $i \in S_\alpha$ we have that $f(i) = (f|_\alpha)(i) = h_\alpha(i) = h(i)$. Since this happens for every $\alpha \in I$, $f = h$. \square

Let \mathbf{F} be a forest and $i \in \mathbf{F}$. Since \mathcal{P} is a presheaf of MTL-algebras, its fiber over i is the set of *germs over i* and is written as \mathcal{P}_i (c.f. II.5 [13]). Recall that $f, g \in \mathcal{P}(S)$ have the same germ at i if there exists some $R \in \mathcal{D}(\mathbf{F})$ with $i \in R$, such that $R \subseteq S \cap T$ and $f|_R = g|_R$. Hence, \mathcal{P}_i results to be a “suitable quotient” of the MTL-algebra $\mathcal{P}(T)$. By Lemma 22, \mathcal{P}_i can be described as the filtering colimit over those $T \in \mathcal{D}(\mathbf{F})$ such that $i \in T$, i.e.,

$$\mathcal{P}_i = \varinjlim_{i \in T} \mathcal{P}(T).$$

Thereby, for every $T \in \mathcal{D}(\mathbf{F})$ the map $\varphi_T : \mathcal{P}(T) \rightarrow \mathcal{P}_i$ that sends $h \in \mathcal{P}(T)$ in its equivalence class “modulo germ at i ” result to be a surjective morphism of MTL-algebras. We write $[h]_T$ for the equivalence class of h in \mathcal{P}_i .

Lemma 23. *Let \mathbf{F} be a forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. For every $i \in F$, $\mathcal{P}(\downarrow i) \cong \mathcal{P}_i$ in \mathcal{MTL} .*

Proof. Let $i \in F$ and consider $\varphi_{\downarrow i} : \mathcal{P}(\downarrow i) \rightarrow \mathcal{P}_i$. From the above discussion it is clear that $\varphi_{\downarrow i}$ is surjective. To check that it is injective, let $f, g \in \mathcal{P}(\downarrow i)$ be such that $[f]_{\downarrow i} = [g]_{\downarrow i}$. There exists some $R \in \mathcal{D}(\mathbf{F})$ with $i \in R$, such that $R \subseteq \downarrow i$ and $f|_R = g|_R$. Since $\downarrow i$ is the smallest downset to which i belongs, we get that $R = \downarrow i$. Then $f = g$. Hence $\varphi_{\downarrow i}$ is an isomorphism in \mathcal{MTL} . \square

Corollary 9. *Let \mathbf{F} be a forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. Then \mathcal{P} is a sheaf of MTL-chains.*

Proof. Apply Lemmas 23 and 17. \square

Observe that the same argument used in Example 2 of [4] can be applied to prove that when the index set is a finite chain, the forest product and the ordinal sum of MTL-algebras coincide. The following result will be relevant for the last part of this paper.

Corollary 10. *Let \mathbf{F} be a finite forest and $\{\mathbf{M}_i\}_{i \in \mathbf{F}}$ a collection of MTL-chains. Then for every $i \in \mathbf{F}$, $\mathcal{P}_i \cong \bigoplus_{i \leq j} \mathbf{M}_i$.*

Proof. If \mathbf{F} is a finite forest then $\downarrow j$ is a finite chain for every $j \in \mathbf{F}$. From the observed above respect to the forest product of MTL-algebras indexed by a finite chain, we conclude that $\mathcal{P}(\downarrow j) \cong \bigoplus_{i \leq j} \mathbf{M}_i$, which clearly is an MTL-chain. Therefore, from Lemma 23 the result follows. \square

We can now put together the Lemma 22, and Corollary 9 in the following statement:

The forest product of MTL-chains is essentially a sheaf of MTL-algebras over an Alexandrov space whose fibers are MTL-chains.

5 From finite forest products to MTL-algebras

In this section we show that a wide class of finite MTL-algebras can be represented as finite forest products of finite archimedean MTL-chains. To do so, we begin by showing that there exist a functor \mathcal{H} from the category of finite labeled forests to the category of finite MTL-algebras. Moreover, we will prove that the functor \mathcal{H} is left adjoint to the functor \mathcal{G} and the counit of such adjunction is an isomorphism. It is worth to mention that this result is strongly based in the characterization of the join irreducible elements of a finite forest product of finite archimedean MTL-chains.

In general, the unit of the adjoint pair $\mathcal{G} \dashv \mathcal{H}$ is not an isomorphism. In subsection 5.1 we present a class of finite MTL-algebras which solves that problem. Finally, in subsection 5.2 we give a simple description of the forest product of finite MTL-chains in terms of ordinal sums and direct products.

Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ and $m : \mathbf{G} \rightarrow \mathfrak{S}$ be finite labeled forests. If $(\varphi, \mathcal{F}) : l \rightarrow m$ is a morphism of finite labeled forests (see Section 3) then $\varphi : F \rightarrow G$ is a p-morphism and $\mathcal{F} = \{f_i\}_{i \in F}$ is a family of morphisms $f_i : (m \circ \varphi)(i) \rightarrow l(i)$ of MTL-algebras. Recall that a morphism of posets is a p-morphism if and only if it is open respect to the Alexandrov topologies of the involved posets, so since $F \in \mathcal{D}(\mathbf{F})$ it follows that $\varphi(F) \in \mathcal{D}(\mathbf{G})$. From Lemma 14, we get that $\bigotimes_{k \in \varphi(F)} m(k)$ is an MTL-algebra. Notice that $m \circ \varphi : \mathbf{F} \rightarrow \mathfrak{S}$ is a finite labeled forest so we can consider the forest product $\bigotimes_{i \in \mathbf{F}} (m \circ \varphi)(i)$. Since $\bigcup_{k \in \varphi(F)} m(k) = \bigcup_{i \in \mathbf{F}} (m \circ \varphi)(i)$, we define, for every $h \in \bigotimes_{k \in \varphi(F)} m(k)$, the map $\gamma : \bigotimes_{k \in \varphi(F)} m(k) \rightarrow \bigotimes_{i \in \mathbf{F}} (m \circ \varphi)(i)$ as the composite

$$F \xrightarrow{\varphi} \varphi(F) \xrightarrow{h} \bigcup_{i \in \mathbf{F}} (m \circ \varphi)(i) \\ \searrow \gamma(h) \nearrow$$

Lemma 24. *The map γ , defined above, is a morphism of MTL-algebras.*

Proof. In order to check that γ is well defined, take $h \in \bigotimes_{k \in \varphi(F)} m(k)$ and consider $i \in F$ such that $\gamma(h)(i) \neq 0_{\varphi(i)}$. Assume $j < i$ in F . By the definition of γ , we get that $h(\varphi(i)) \neq 0_{\varphi(i)}$. From the monotonicity of φ , it follows that $\varphi(j) < \varphi(i)$. Then by assumption, when we have that $h(\varphi(j)) = 1$, and consequently $\gamma(h)(i) = 1$. By Definition 1, we have that $\gamma(h) \in \bigotimes_{i \in \mathbf{F}} (m \circ \varphi)(i)$. The proof of the fact that γ is an homomorphism is straightforward. \square

Notice, in addition that the family \mathcal{F} induces a map $\alpha : \bigotimes_{i \in \mathbf{F}} (m \circ \varphi)(i) \rightarrow \bigotimes_{i \in \mathbf{F}} l(i)$ defined as $\alpha(g)(i) = f_i(g(i))$, for every $i \in \mathbf{F}$.

Lemma 25. *The map α , defined above, is a morphism of MTL-algebras.*

Proof. Let $g \in \bigotimes_{i \in \mathbf{F}} (m \circ \varphi)(i)$ be such that $\alpha(g)(i) \neq 1$. Let $j \in \mathbf{F}$ be such that $i < j$. Since, $f_i(g(i)) \neq 1$, we get that, by Lemma 3 $g(i) \neq 1$. Hence, by assumption $g(j) = 0_{\varphi(j)}$. Thereby $\alpha(g)(j) = f_j(g(j)) = 0_j$, and by (2) of Lemma 15, we have that $\alpha(g) \in \bigotimes_{i \in \mathbf{F}} l(i)$. The proof of the fact that α is an homomorphism is straightforward. \square

The Lemmas 24 and 25 allows us to consider the following composite of morphisms of MTL-algebras:

$$\mathcal{P}_m(\mathbf{G}) \xrightarrow{\beta} \bigotimes_{k \in \varphi(F)} m(k) \xrightarrow{\gamma} \bigotimes_{i \in \mathbf{F}} (m \circ \varphi)(i) \xrightarrow{\alpha} \mathcal{P}_l(\mathbf{F})$$

where $\mathcal{P}_m(G) = \bigotimes_{k \in \mathbf{G}} m(k)$, $\mathcal{P}_l(\mathbf{F}) = \bigotimes_{i \in \mathbf{F}} l(i)$ and $\beta : \mathcal{P}_m(G) \rightarrow \bigotimes_{k \in \varphi(F)} m(k)$ is the restriction of $\mathcal{P}_m(G)$ to $\varphi(\mathbf{F})$.

Theorem 2. *The assignments $l \mapsto \mathcal{P}_l(F)$ and $(\varphi, \mathcal{F}) \mapsto \alpha\gamma\beta$ define a contravariant functor*

$$\mathcal{H} : f\mathcal{LF} \rightarrow f\mathcal{MTL}.$$

Proof. Let $l : \mathbf{F} \rightarrow \mathfrak{S}$, $m : \mathbf{G} \rightarrow \mathfrak{S}$ and $n : \mathbf{H} \rightarrow \mathfrak{S}$ be finite labeled forests, and $(\varphi, \mathcal{F}) : l \rightarrow m$ and $(\psi, \mathcal{G}) : l \rightarrow m$ be morphism of finite labeled forests. Let

$$\mathcal{M} = \{f_i g_{\varphi(i)} : n(\psi(\varphi(i))) \rightarrow l(i) \mid i \in \mathbf{F}\}.$$

Consider $s \in \mathcal{H}(n)$ and $i \in \mathbf{F}$. Then from

$$\begin{aligned} \mathcal{H}[(\psi, \mathcal{G})(\varphi, \mathcal{F})](s)(i) &= \mathcal{H}(\psi\varphi, \mathcal{M})(s)(i) \\ &= (f_i g_{\varphi(i)})[s(\psi(\varphi(i)))] \\ &= f_i[g_{\varphi(i)}(s(\psi(\varphi(i))))] \\ &= f_i[\mathcal{H}(\psi, \mathcal{G})(s)(\varphi(i))] \\ &= \mathcal{H}(\varphi, \mathcal{F})[\mathcal{H}(\psi, \mathcal{G})(s)](i) \\ &= [\mathcal{H}(\varphi, \mathcal{F})\mathcal{H}(\psi, \mathcal{G})](s)(i) \end{aligned}$$

we conclude that $\mathcal{H}[(\psi, \mathcal{G})(\varphi, \mathcal{F})] = \mathcal{H}(\varphi, \mathcal{F})\mathcal{H}(\psi, \mathcal{G})$. Since $id_l = (id_F, \mathcal{I})$, where \mathcal{I} is the family formed by the identities of $\{l(i)\}_{i \in \mathbf{F}}$, it is clear from the definition of \mathcal{H} that $\mathcal{H}(id_l) = id_{\mathcal{H}(l)}$. \square

Lemma 26. *Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest. Then $h \in \mathcal{I}(\mathcal{P}_l(\mathbf{F}))$ if and only if $C_h = \emptyset$.*

Proof. Observe that $h \in \mathcal{I}(\mathcal{P}_l(\mathbf{F}))$ if and only if $h(i)^2 = h(i)$, for every $i \in F$, which is equivalent to say that $h(i) \in \mathcal{I}(l(i))$. Since $l(i)$ is an arquimedean MTL-chain, by Proposition 1, the only possible case is $h(i) = 0_i$ or $h(i) = 1$. This concludes the proof. \square

Let \mathbf{F} be a finite forest and $S \subseteq F$. We write $\text{Max}(S)$ for the maximal elements of S .

Lemma 27. *Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest and $S \in \mathcal{D}(\mathbf{F})$, then*

$$X_S = \{h \in \mathcal{P}_l(\mathbf{F}) \mid h(i) = 1, \text{ for every } i \in \text{Max}(S)\}$$

Proof. Let $h \in \mathcal{P}_l(\mathbf{F})$ and suppose $h(i) = 1$ for every $i \in \text{Max}(S)$. If $j \in S$, there exists some $i \in \text{Max}(S)$ such that $j \leq i$. Since $h(i) \neq 0_i$, $h(j) = 1$ and $h \in X_S$. The other inclusion is straightforward. \square

Recall that, from Corollary 3 there exist a unique $h_S \in \mathcal{I}(\mathcal{P}_l(\mathbf{F}))$ such that $X_S = \uparrow h_S$. As a direct consequence of Lemmas 26 and 27 we obtain the following result.

Lemma 28. *Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest and $S \in \mathcal{D}(\mathbf{F})$, then*

$$h_S(j) = \begin{cases} 1, & j \leq i, \text{ for some } i \in \text{Max}(S) \\ 0_j, & \text{otherwise} \end{cases}$$

Corollary 11. *Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest. The following holds for every $i \in F$:*

1. $X_{\downarrow i}$ is a prime filter of $\mathcal{P}_l(\mathbf{F})$,
2. $X_{\downarrow i} = \{h \in \mathcal{P}_l(\mathbf{F}) \mid h(i) = 1\}$,
3. The map

$$h_{\downarrow i}(j) = \begin{cases} 1, & j \leq i \\ 0_j, & \text{otherwise} \end{cases}$$

is a non zero join irreducible element of $\mathcal{P}_l(\mathbf{F})$.

Proof. A direct consequence of Lemmas 21, 27, 28 and Corollary 4. \square

Lemma 29. *Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest and $h \in \mathcal{I}(\mathcal{P}_l(\mathbf{F}))$, then $h \in \mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F})))$ if and only if $h^{-1}(1)$ is a chain.*

Proof. Let us assume that $h^{-1}(1)$ is a chain, then, since F is finite, $h^{-1}(1) = \downarrow i$ for some $i \in F$. Suppose that there are $g, f \in \mathcal{I}(\mathcal{P}_l(\mathbf{F}))$ such that $h = g \vee f$, then $h(k) = g(k) \vee f(k)$, for every $k \in F$. If $k \leq i$, we get that $g(k) \vee f(k) = 1$. Since $l(k)$ is really local, by Lemma 16, $g(k) = 1$ or $f(k) = 1$. Consequently, $g(k) = h(k)$ or $f(k) = h(k)$. If $h(k) = 0_k$, the result follows, since 0_k is join irreducible in $l(k)$. Hence h is join irreducible. On the other hand, suppose that $h \in \mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F})))$. If $h^{-1}(1)$ is not a chain, there exist $i, j \in F$ not comparables such that $h(i) = h(j) = 1$. Let us consider the following functions:

$$g(k) = \begin{cases} h(k), & \text{if } k \neq i \\ 0_i, & \text{otherwise} \end{cases} \quad f(k) = \begin{cases} h(k), & \text{if } k \neq j \\ 0_j, & \text{otherwise} \end{cases}$$

From Lemma 26, it follows that $g, f \in \mathcal{I}(\mathcal{P}_l(\mathbf{F}))$. Thereby, $h = g \vee f$, which is in contradiction with the assumption. \square

Lemma 30. *Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest. There is a poset isomorphism between \mathbf{F} and $\mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F})))$.*

Proof. Let us to consider $\mu : \mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F}))) \rightarrow F$, defined as $\mu(h) = \max h^{-1}(1) = i_h$. From Lemma 29, it follows that μ is well defined and is injective. To verify that μ is surjective, take $i \in F$ and define

$$h_i(j) = \begin{cases} 1, & j \leq i \\ 0_j, & \text{otherwise.} \end{cases}$$

From Lemma 29, it follows that $h_i \in \mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F})))$. It is clear that $\mu(h_i) = i$. In order to check the monotonicity of μ , let us suppose that $h \leq g$, for $h, g \in \mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F})))$. From Corollary 11.3, we have that $h^{-1}(1) \subseteq g^{-1}(1)$, so $i_h \leq i_g$ and consequently $\mu(h) \leq \mu(g)$. The monotonicity of μ^{-1} is straightforward. \square

Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest. For every $i \in F$ we write h_i for the map of 3. in Corollary 11.3. Let us to consider the assignment $\varphi_l : F \rightarrow \mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F})))$, defined as $\varphi_l(i) = h_i$.

Lemma 31. *The assignment φ_l is a p-morphism.*

Proof. The monotonicity of φ_l follows from Corollary 11.3. On the other hand, take $i \in F$ and suppose that $g \leq h_i$. Thus, $h(i) = 1$ implies that $g(i) = 1$ and, due to Corollary 11.2 we get that $g \in X_{\downarrow i}$. Therefore, $h_i \leq g$. In consequence, $g = \varphi_l(i)$, which was our aim. \square

Lemma 32. *The p-morphism φ_l is an isomorphism.*

Proof. We first prove the injectivity of φ_l . Let $i, j \in F$ be such that $\varphi_l(i) = \varphi_l(j)$. Then $h_i = h_j$. In particular $h_i(i) = h_j(i)$ and $h_j(j) = h_i(j)$, which, by definition of h_i and h_j means that $i \leq j$ and $j \leq i$. The surjectivity of φ_l follows from Lemma 29. Finally, we verify that φ_l^{-1} is also a p-morphism. To do so, notice that for every $h \in \mathcal{P}_l(\mathbf{F})$, $\varphi_l^{-1}(h)$ is just the $i \in F$ described in Lemma 29. We will denote such element as i_h . Let us suppose that $j \leq \varphi_l^{-1}(h)$, then $h(j) = h(i_h) = 1$ so, for every $k \leq j \leq i_h$ we get that $h(k) = h(j) = 1$. Take

$$g(k) = \begin{cases} 1, & k \leq j \\ 0_k, & \text{otherwise} \end{cases}$$

It is clear that $g \in \mathcal{P}_l(\mathbf{F})$, $g \leq h$ and $\varphi_l^{-1}(j) = g$. \square

Observe that, from Lemmas 8 and 31 we have that for every $i \in F$, there exists a unique $a_{\varphi_l(i)} \in \mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F})))$ such that $a_{\varphi_l(i)} \prec \varphi_l(i)$. Due to Lemma 10, $\uparrow a_{\varphi_l(i)} / \uparrow \varphi_l(i)$ is an archimedian MTL-chain. Let us consider the assignment $\tau_i : \uparrow a_{\varphi_l(i)} \rightarrow l(i)$, defined

as $\tau_i(h) = h(i)$. It is clear, from the definition, that τ_i preserves all the binary monoid operations. Moreover, it preserves the residual. If $a_{\varphi_l(i)} \leq f, g$, then for every $j < i$, $f(j) = g(j) = 1$ so $f(j) \leq g(j)$, which means that $(f \rightarrow g)(i) = f(i) \rightarrow_i g(i)$, and consequently, $\tau_i(f \rightarrow g) = \tau_i(f) \rightarrow_i \tau_i(g)$. We have proved the following result:

Lemma 33. *The function τ_i , defined above, is a morphism of MTL-algebras.*

Notice that, from the universal property of quotients in \mathcal{MTL} , Lemma 33 implies that for every $i \in F$ there exists a unique morphism of MTL-algebras $f_i : (\uparrow a_{\varphi_l(i)})/(\uparrow \varphi_l(i)) \rightarrow l(i)$ such that the diagram

$$\begin{array}{ccc} \uparrow (a_{\varphi_l(i)}) & \xrightarrow{\quad} & \uparrow (a_{\varphi_l(i)}) / \uparrow (\varphi_l(i)) \\ & \searrow \tau_i & \downarrow f_i \\ & & l(i) \end{array}$$

commutes.

Lemma 34. *For every $i \in F$, $f_i : (\uparrow a_{\varphi_l(i)})/(\uparrow \varphi_l(i)) \rightarrow l(i)$ is an isomorphism.*

Proof. To prove the injectivity of f_i , suppose that $f_i([f]) = f_i([g])$. Then $f(i) = g(i)$. Since $a_{\varphi_l(i)} \leq f, g$, for every $j < i$, $f(j) = g(j) = 1$. Hence, $f(i) \leq g(i)$, for every $i \leq j$, which is equivalent to say that $[f] = [g]$ in $(\uparrow a_{\varphi_l(i)})/(\uparrow \varphi_l(i))$. To check the surjectivity, take $x \in l(i)$. Define

$$h(k) = \begin{cases} 1, & k \leq i \\ x, & k = i \\ 0_k, & \text{otherwise} \end{cases}$$

It is clear that $f_i([h]) = x$. □

Let \mathcal{G} and \mathcal{H} be the functors from Theorems 1 and 2, respectively.

Lemma 35. *Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest. Then $\mathcal{G} \circ \mathcal{H}$ is isomorphic to the identity functor in $f\mathcal{LF}$.*

Proof. A direct application of Lemmas 32 and 34. □

Proposition 2. *The functor \mathcal{G} is left adjoint to the functor \mathcal{H} . Moreover, \mathcal{G} is full and faithful.*

Proof. Let M be a finite MTL-algebra, $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest and $f : M \rightarrow \mathcal{P}_l(\mathbf{F})$ be a morphism of MTL-algebras. By Lemma 11, there exists a unique p-morphism $f^* : \mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F}))) \rightarrow \mathcal{J}(\mathcal{I}(M))$. Let $\varphi : F \rightarrow \mathcal{J}(\mathcal{I}(M))$ be defined as $\varphi = f^* \mu^{-1}$, where μ is the isomorphism between F and $\mathcal{J}(\mathcal{I}(\mathcal{P}_l(\mathbf{F})))$ given in Lemma 30. It is clear that φ is a p-morphism. On the other hand, if we write h_i for $\mu^{-1}(i)$, then from Lemma 13, it follows that for every $i \in F$ there exists a morphism of MTL-algebras $\bar{f}_{h_i} : \uparrow a_{f^*(h_i)} \rightarrow \uparrow a_{h_i}$ which determines a unique morphism of MTL-algebras $f_{h_i} : (\uparrow a_{f^*(h_i)})/(\uparrow f^*(h_i)) \rightarrow (\uparrow a_{h_i})/(\uparrow h_i)$. From Lemma 34 we get that $(\uparrow a_{h_i})/(\uparrow h_i) \cong l(i)$. Let us to consider $f_i : (\uparrow a_{f^*(h_i)})/(\uparrow h_i) \rightarrow l(i)$ as the composition of f_{h_i} with the isomorphism of Lemma 34. This concludes the proof. □

5.1 The duality theorem

In this subsection we present a duality theorem between the class of representable finite MTL-algebras and finite labeled forests. To do so, we restrict the results previously obtained in this section to the class of representable finite MTL-algebras.

Definition 2. Let M be a finite MTL-algebra. An element $e \in \mathcal{I}(M)^*$ is said to be a local unit if for every $x \leq e$, $ex = x$.

Lemma 36. Let M be a finite MTL-algebra and $e \in \mathcal{I}(M)^*$. The following are equivalent:

1. e is a local unit.
2. $ey = e \wedge y$, for every $y \in M$.

Proof. Let us assume that e is a local unit and $y \in M$. Since $e \wedge y \leq y$, $e(e \wedge y) \leq ey$. Hence $e \wedge y \leq ey$, because $e \wedge y \leq e$ and $e(e \wedge y) = e \wedge y$. The other case is straightforward. \square

Definition 3. A finite MTL-algebra M is said to be representable if every non zero idempotent satisfies any of the equivalent conditions of Lemma 36.

Let M be a representable MTL-algebra. For the rest of this section we will write F_M to denote $\mathcal{J}(\mathcal{I}(M))$.

Remark 6. Observe that as a direct consequence of Definition 2, it follows that for every $e \in F_M$, $\uparrow a_e / \uparrow e \cong [a_e, e]$.

Let M be an representable MTL-algebra and $m \in \text{Max}(F_M)$. In what follows we will denote the set $(\downarrow m) \cap F_M$ simply by $(\downarrow m)$.

Lemma 37. Let M be a representable MTL-algebra and $m \in \text{Max}(F_M)$. Then, for every $x \in M$ there exists some $e \in \downarrow m$ such that $a_e \leq x$.

Proof. Suppose that there exist $y > 0$ such that for every $e \in \downarrow m$ it holds that $a_e \not\leq y$. In particular, if $n = \min(\downarrow m)$, it follows that $a_n = 0$; so $0 \not\leq x$, which is clearly absurd. \square

Lemma 38. For every representable MTL-algebra M and $m \in \text{Max}(F_M)$, $M/(\uparrow m) \cong \bigoplus_{e \leq m} [a_e, e]$.

Proof. In order to simplify the notation for this proof, we write B for $\bigoplus_{e \leq m} [a_e, e]$. Let us consider the map $f : M \rightarrow B$, defined as $f(x) = x \wedge m$. It is clear that f is a surjective morphism of MTL-algebras such that $f(m) = 1 \in B$. We stress that B has the universal property of $M/ \uparrow m$. To prove it, let $g : M \rightarrow E$ be a morphism of MTL-algebras such that $g(m) = 1_E$. For every $z \in B$ define $\psi(z) = g(z)$. It easily follows that $\psi \circ f = g$, and since f is surjective, then ψ is unique. Hence $M/(\uparrow m) \cong B$. \square

Let M be a representable MTL-algebra. Recall that $\mathcal{G}(M) = l_M : F_M \rightarrow \mathfrak{S}$ is a finite labeled forest, so, from Lemma 30, it follows that $F_M \cong \mathcal{J}(\mathcal{I}(\mathcal{P}_{l_M}(F_M)))$. As a consequence, we get that $\bigoplus_{e \leq m} [a_e, e] \cong \mathcal{P}_{l_M}(\downarrow m)$. Explicitly, such assignment is defined for every $z \in \bigoplus_{e \leq m} [a_e, e]$ by:

$$h_z(c) = \begin{cases} m, & c \leq e_z \\ z, & c = e_z \\ a_c, & c > e_z \end{cases} \quad (6)$$

Here e_z is the unique idempotent join irreducible below m such that $a_{e_z} \leq z \leq e_z$.

Observe that $F_M = \bigcup_{m \in F_M} (\downarrow m)$. Hence the family $\mathcal{R} = \{\downarrow m\}_{m \in M}$ is a covering for F_M . Let $f_m : M \rightarrow \mathcal{P}_{l_M}(\downarrow m)$ be defined as $f_m(x) = h_{x \wedge m}$.

Lemma 39. *Let M be a representable MTL-algebra. For every $x \in M$, the family $\{f_m(x)\}_{m \in \text{Max}(F_M)}$ is a matching family for the covering \mathcal{R} .*

Proof. Let $m, n \in \text{Max}(F_M)$. Since F_M is a forest, $(\downarrow m) \cap (\downarrow n) = \emptyset$ or $(\downarrow m) \cap (\downarrow n) \neq \emptyset$. In the first case, the result holds, because by Lemma 22, \mathcal{P}_{l_M} is a sheaf. In the second case, there exists an $e \in F_M$ with $e \leq m, n$ such that $(\downarrow m) \cap (\downarrow n) = (\downarrow e)$. Observe that

$$(x \wedge m) \wedge e = x \wedge (m \wedge e) = x \wedge e = x \wedge (n \wedge e) = (x \wedge n) \wedge e.$$

Then, from the description of $h_{x \wedge m}$ and $h_{x \wedge n}$ of equation (6), we obtain that $h_{x \wedge m}|_{\downarrow e} = h_{x \wedge n}|_{\downarrow e}$. \square

Recall that Lemma 22 states that \mathcal{P}_{l_M} is a sheaf so, from Lemma 39, we obtain that every $x \in M$ determines an amalgamation h_x for the family $\{f_m(x)\}_{m \in \text{Max}(F_M)}$. This fact allows us to consider the assignment $f_M : M \rightarrow \mathcal{P}(F_M)$, defined as $f(x) = h_x$. Observe that by construction f is a morphism of MTL-algebras.

Lemma 40. *For every representable MTL-algebra M , the assignment f_M is an isomorphism.*

Proof. Only remains to check that f_M is bijective. To prove the injectivity of f_M , suppose $h_x = h_y$, then, since h_x and h_y are the amalgamations of the families $\{f_m(x)\}_{m \in \text{Max}(F_M)}$ and $\{f_m(y)\}_{m \in \text{Max}(F_M)}$, respectively, it follows that $h_{x \wedge m} = h_{y \wedge m}$ for every $m \in \text{Max}(F_M)$. Then, from equation (6), it follows that $x \wedge m = y \wedge m$, for every $m \in \text{Max}(F_M)$. Then, $\bigvee_{m \in \text{Max}(F_M)} (x \wedge m) = \bigvee_{m \in \text{Max}(F_M)} (y \wedge m)$. Thereby, $x \wedge \bigvee_{m \in \text{Max}(F_M)} m = y \wedge \bigvee_{m \in \text{Max}(F_M)} m$. Since $\bigvee_{m \in \text{Max}(F_M)} m = 1$, $x = y$.

Finally, to prove the surjectivity of f_M , let $h \in \mathcal{P}(F_M)$. Then, $h|_{\downarrow m} \in \mathcal{P}(\downarrow m)$ for every $m \in \text{Max}(F_M)$. Since $\mathcal{P}(\downarrow m) \cong \bigoplus_{e \leq m} [a_e, e]$ we will write z_m for the unique element of $\bigoplus_{e \leq m} [a_e, e]$ which corresponds to $h|_{\downarrow m}$. Observe that for every $m, n \in \text{Max}(F_M)$ we have that

$$(h|_{\downarrow m})|_{\downarrow m \cap \downarrow n} = (h|_{\downarrow n})|_{\downarrow m \cap \downarrow n}$$

Thereby $z_m \wedge e_{mn} = z_n \wedge e_{mn}$, where e_{mn} is the greatest $e \in F_M$ below m and n . Hence

$$z_m \wedge n = (z_m \wedge e_{mn}) \wedge (m \wedge n) = (z_n \wedge e_{mn}) \wedge (m \wedge n) = z_n \wedge m, \quad (7)$$

since $z_m \leq m$ and $z_n \leq n$. If we consider $x = \bigvee_{m \in \text{Max}(F_M)} z_m$, then applying equation (7), in the following calculation

$$x \wedge n = \bigvee_{m \in \text{Max}(F_M)} (z_m \wedge n) = \bigvee_{m \in \text{Max}(F_M)} (z_n \wedge m) = z_n \wedge \left(\bigvee_{m \in \text{Max}(F_M)} m \right) = z_n,$$

we obtain that $h_x|_{\downarrow n} = h|_{\downarrow n}$ for every $n \in \text{Max}(F_M)$. Thereby, since h is the amalgamation of the family $\{h|_{\downarrow m}\}_{m \in \text{Max}(F_M)}$ and \mathcal{P}_{l_M} is a sheaf, it follows that $h_x = h$. \square

Write $r\mathcal{MTL}$ for the category of representable finite MTL-algebras. Let $u\mathcal{LF}$ be the subcategory of $f\mathcal{LF}$ whose objects are the finite labeled forest such that their poset product is a representable MTL-algebra. Let us write \mathcal{G}^* for the restriction of the functor \mathcal{G} to the category $r\mathcal{MTL}$ and \mathcal{H}^* for the restriction of the functor \mathcal{H} to $u\mathcal{LF}$. From Proposition 2, it follows that $\mathcal{G}^* \dashv \mathcal{H}^*$ and that the unit is an isomorphism. Notice that the assignment $f_M : M \rightarrow \mathcal{P}(F_M)$ is the counit of the latter adjunction and, by Lemma 40, it is an isomorphism. We have proved the main result of this paper:

Theorem 3. *The categories $r\mathcal{MTL}$ and $u\mathcal{LF}$ are dually equivalent.*

5.2 An explicit description of finite forest products

The aim of this section is to bring a characterization of the forest product of finite MTL-algebras in terms of ordinal sums and direct products. Unlike the rest of this work, the methods used in this part are completely recursive. Finally, we recall that, along this section, the symbol \oplus will be used indistinctly, to denote the ordinal sum of posets and the ordinal sum of MTL-algebras.

In [1], Aguzzoli suggest that every finite forest can be built recursively. We adapt those ideas in the following definition:

Definition 4. *The set of finite forest \mathbf{fFor} is the smallest collection of finite posets satisfying the following conditions:*

- (F1) $1 \in \mathbf{fFor}$,
- (F2) If $F \in \mathbf{fFor}$, then $1 \oplus F \in \mathbf{fFor}$,
- (F3) If $F_1, \dots, F_m \in \mathbf{fFor}$ then $\biguplus_{k=1}^m F_k \in \mathbf{fFor}$.

Recall that every finite forest \mathbf{F} can be expressed as a finite disjoint union of finite trees. Hence each finite forest can be written as $\mathbf{F} = \biguplus_{k=1}^n \mathbf{T}_k$, where \mathbf{T}_k is a finite tree. We call the family $\{\mathbf{T}_k\}$ the *family of component trees* of \mathbf{F} .

Lemma 41. *Let \mathbf{F} be a finite forest and $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest. Then*

$$\mathcal{P}_l\left(\biguplus_{k=1}^n \mathbf{T}_k\right) \cong \mathcal{P}_{l_1}(\mathbf{T}_1) \times \dots \times \mathcal{P}_{l_n}(\mathbf{T}_n)$$

where $l_i = l|_{\mathbf{T}_i}$, for every $i = 1, \dots, n$.

Proof. Consider $\varphi : \mathcal{P}_l(\biguplus_{k=1}^n \mathbf{T}_k) \rightarrow \mathcal{P}_{l_1}(\mathbf{T}_1) \times \dots \times \mathcal{P}_{l_n}(\mathbf{T}_n)$ defined as $\varphi(h) = (h|_{\mathbf{T}_1}, \dots, h|_{\mathbf{T}_n})$. Also, consider $\tau : \mathcal{P}_{l_1}(\mathbf{T}_1) \times \dots \times \mathcal{P}_{l_n}(\mathbf{T}_n) \rightarrow \mathcal{P}_l(\biguplus_{k=1}^n \mathbf{T}_k)$ defined as $\tau(h_1, \dots, h_n) = h$, with $h(i) = h_j(i)$ if $i \in \mathbf{T}_j$. It is clear that φ and τ are well defined morphisms of MTL-algebras and that one is the inverse of the other. \square

Lemma 42. *Let \mathbf{F} be a finite forest and $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest. If $\mathbf{F} = 1 \oplus \mathbf{F}_0$, where \mathbf{F}_0 is a finite forest, then*

$$\mathcal{P}_l(\mathbf{F}) \cong l(\perp) \oplus \mathcal{P}_{l_0}(\mathbf{F}_0)$$

where $\perp = \text{Min}(\mathbf{F})$ and $l_0 = l|_{\mathbf{T}_0}$.

Proof. Let us assume $\mathbf{F} = 1 \oplus \mathbf{F}_0$ and suppose that $h \in \mathcal{P}_l(\mathbf{F})$. If $A_h = \{i \in \mathbf{F} \mid h(i) \neq 1\}$, then either $\perp \in A_h$, $\perp \notin A_h$ or $h(\perp) = 0_\perp$. In the first case, from the assumption it follows that $h(j) = 0_j$ for every $j \in \mathbf{F}_0$. In the second case, again, by assumption, we obtain that $h(\perp) = 1$. The final case implies that $h = 0$. Thereby, h represents either an element of $l(\perp)$ (those with $h(\perp) \neq 1$) or an element of $\mathcal{P}_{l_0}(\mathbf{F}_0)$ considered as $h = h|_{\mathbf{F}_0}$. Based on this fact we consider $p : \mathcal{P}_l(\mathbf{F}) \rightarrow l(\perp) \oplus \mathcal{P}_{l_0}(\mathbf{F}_0)$, defined for every $h \in \mathcal{P}_l(\mathbf{F})$ as

$$p(h) = \begin{cases} h(\perp), & \text{if } h(\perp) \neq 1 \\ h|_{\mathbf{F}_0}, & \text{if } h(\perp) = 1 \end{cases}$$

Let $a \in l(\perp) \oplus \mathcal{P}_{l_0}(\mathbf{F}_0)$. Then, a is either an element of $l(\perp)$ or $a \in \mathcal{P}_{l_0}(\mathbf{F}_0)$. Let us take $q : l(\perp) \oplus \mathcal{P}_{l_0}(\mathbf{F}_0) \rightarrow \mathcal{P}_l(\mathbf{F})$ as $q(a) = h_a$, where $h_a(i) = a(i)$, if $i \in \mathbf{F}_0$ or

$$h_a(\perp) = \begin{cases} a, & \text{if } a \in l(\perp) \\ 1, & \text{otherwise} \end{cases}, \text{ if } i \notin \mathbf{F}_0.$$

It is clear that p and q are well defined morphisms of MTL-algebras such that one is the inverse of the other. This concludes the proof. \square

Let \mathbf{T} be a finite tree, $i \in \mathbf{T}$ and consider the *set of covering elements* of i :

$$\mathcal{C}_{\mathbf{T}}(i) = \{j \in \mathbf{T} \mid i \prec j\}$$

where \prec denotes the covering relation in posets.

Definition 5. *Let \mathbf{T} be a finite tree and $l : \mathbf{T} \rightarrow \mathfrak{S}$ be a finite labeled forest. For every $i \in \mathbf{T}$ let us define recursively the following MTL-algebra:*

$$M_{\mathbf{T}}(i) = \begin{cases} l(i), & i \in \text{Max}(\mathbf{T}) \\ l(i) \oplus \prod_{j \in \mathcal{C}_{\mathbf{T}}(i)} M_{\mathbf{T}}(j), & i \notin \text{Max}(\mathbf{T}). \end{cases}$$

In the following, we will write $\mathcal{K}_l(\mathbf{T})$ for $M_{\mathbf{T}}(m)$, where m is the last element of \mathbf{T} .

Let \mathbf{F} be a finite forest and $\{\mathbf{T}_k\}$ be its collection of component trees. If $l : \mathbf{F} \rightarrow \mathfrak{S}$ is a finite labeled forest let us consider the MTL-algebra:

$$\mathcal{K}_l(\mathbf{F}) = \prod_{k=1}^n \mathcal{K}_l(\mathbf{T}_k) \tag{8}$$

Proposition 3. *Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest. Then $\mathcal{P}_l(\mathbf{F}) \cong \mathcal{K}_l(\mathbf{F})$.*

Proof. We prove this Proposition by induction over \mathbf{fFor} . If $F = \mathbf{1}$, the conclusion is trivial. On the other hand, let us suppose that $\mathbf{F} = \mathbf{1} \oplus \mathbf{F}_0$, with $\mathbf{F}_0 \in \mathbf{fFor}$ be such that $\mathcal{P}_l(\mathbf{F}_0) \cong \mathcal{K}_l(\mathbf{F}_0)$. From Lemma 42 and the inductive hypothesis, it follows that

$$\mathcal{P}_l(\mathbf{F}) \cong l(\perp) \oplus \mathcal{P}_{l_0}(\mathbf{F}_0) \cong l(\perp) \oplus \mathcal{K}_{l_0}(\mathbf{F}_0).$$

Since \mathbf{F}_0 is a finite forest, $\mathbf{F}_0 = \biguplus_{k=1}^r \mathbf{T}_k^0$. Thus by equation (8):

$$\mathcal{K}_{l_0}(\mathbf{F}_0) = \prod_{k=1}^r \mathcal{K}_{l_0}(\mathbf{T}_k^0) = \prod_{k=1}^r M_{\mathbf{T}_k^0}(m_k),$$


with $m_k = \text{Min}(\mathbf{T}_k^0)$. Since $\mathcal{C}_{\mathbf{F}}(\perp) = \{m_1, \dots, m_r\}$, by Definition 5, we have that $\mathcal{P}_l(\mathbf{F}) \cong l(\perp) \oplus \mathcal{K}_{l_0}(\mathbf{F}_0) = \mathcal{K}_l(\mathbf{F})$. Finally, assume that $\mathbf{F} = \biguplus_{k=1}^n \mathbf{F}_k$, with $\mathbf{F}_k \in \mathbf{fFor}$ be such that $\mathcal{P}_l(\mathbf{F}_k) \cong \mathcal{K}_l(\mathbf{F}_k)$, for every $i = 1, \dots, n$. Since $\mathbf{F}_k = \biguplus_{i=1}^{m_k} \mathbf{T}_{ki}$, with $\{\mathbf{T}_{ki}\}$ be the family of component trees of \mathbf{F}_k , then $\mathbf{F} = \biguplus_{k=1}^n \biguplus_{i=1}^{m_k} \mathbf{T}_{ki}$. By Lemma 41, we have that

$$\mathcal{P}_l(\mathbf{F}) \cong \mathcal{P}_l(\mathbf{T}_{11}) \times \dots \times \mathcal{P}_l(\mathbf{T}_{1m_1}) \times \dots \times \mathcal{P}_l(\mathbf{T}_{n1}) \times \dots \times \mathcal{P}_l(\mathbf{T}_{nm_n}). \quad (9)$$

Since the direct product of algebras is associative, by Lemma 41, $\mathcal{P}_l(\mathbf{F}_k) \cong \prod_{i=1}^{m_k} \mathcal{P}_l(\mathbf{T}_{ki})$. Then, from equation (9), we get that $\mathcal{P}_l(\mathbf{F}) \cong \prod_{k=1}^n \mathcal{P}_l(\mathbf{F}_k)$. Hence, by inductive hypothesis, we have that $\mathcal{P}_l(\mathbf{F}) \cong \prod_{k=1}^n \mathcal{K}_l(\mathbf{F}_k)$. From equation (8), we have that $\mathcal{K}_l(\mathbf{F}_k) = \prod_{i=1}^{m_k} \mathcal{K}_l(\mathbf{T}_{ki})$. Since, the family of component trees of \mathbf{F} is $\bigcup_{k,i=1}^{n,m_k} \{\mathbf{T}_{ki}\}$, using again equation (8), we conclude that $\mathcal{P}_l(\mathbf{F}) \cong \mathcal{K}_l(\mathbf{F})$. \square

In the following example we illustrate how to build an MTL-algebra by applying equation (8) and Definition 5.

Example 1. Let $l : \mathbf{F} \rightarrow \mathfrak{S}$ be a finite labeled forest and regard the following finite tree \mathbf{F} :



As we can see, $\mathbf{F} = \mathbf{T}_1 \uplus \mathbf{T}_2$, where $\mathbf{T}_1 = \{a, c, g, h\}$ and $\mathbf{T}_2 = \{b, d, e, f\}$. Since $\text{Min}(\mathbf{T}_1) = \{a\}$, $\text{Min}(\mathbf{T}_2) = \{b\}$, from equation (8), it follows that:

$$\mathcal{K}_l(\mathbf{F}) = \mathcal{K}_l(\mathbf{T}_1) \times \mathcal{K}_l(\mathbf{T}_2) = M_{\mathbf{T}_1}(a) \times M_{\mathbf{T}_2}(b).$$

Observe that $\mathcal{C}_{\mathbf{T}_1}(a) = \{c\}$ and $\mathcal{C}_{\mathbf{T}_2}(b) = \{d, e, f\}$. Since $a \notin \text{Max}(\mathbf{T}_1)$ and $b \notin \text{Max}(\mathbf{T}_2)$, applying the Definition 5 to $M_{\mathbf{T}_1}(a)$ and $M_{\mathbf{T}_2}(b)$ respectively, we obtain:

$$\mathcal{K}_l(S) = [l(a) \oplus M_{\mathbf{T}_1}(c)] \times [l(b) \oplus (M_{\mathbf{T}_2}(d) \times M_{\mathbf{T}_2}(e) \times M_{\mathbf{T}_2}(f))].$$

Since $c \notin \text{Max}(\mathbf{T}_1)$ and $\mathcal{C}_{\mathbf{T}_1}(c) = \{g, h\}$ but $d, e, f \in \text{Max}(\mathbf{T}_2)$; again, applying Definition 5 to $M_{\mathbf{T}_1}(c)$, $M_{\mathbf{T}_2}(d)$, $M_{\mathbf{T}_2}(e)$ and $M_{\mathbf{T}_2}(f)$ respectively, we get:

$$\mathcal{K}_l(S) = [l(a) \oplus (l(c) \oplus (M_{\mathbf{T}_1}(g) \times M_{\mathbf{T}_1}(h)))] \times [l(b) \oplus (l(d) \times l(e) \times l(f))].$$

Finally, since $g, h \in \text{Max}(\mathbf{T}_1)$, applying the Definition 5 to $M_{\mathbf{T}_1}(g)$ and $M_{\mathbf{T}_1}(h)$, we conclude that:

$$\mathcal{K}_l(S) = [l(a) \oplus (l(c) \oplus (l(g) \times l(h)))] \times [l(b) \oplus (l(d) \times l(e) \times l(f))].$$

Acknowledgements

This work was supported by CONICET [PIP 112-201501-00412].

References

- [1] S. Aguzzoli. Finite Forests. Their Algebras and Logics. *Slides for LATD 2012*. Kanazawa, Japan. 2012
- [2] S. Aguzzoli, S. Bova and V. Marra. Applications of Finite Duality to Locally Finite Varieties of BL-Algebras, in S. Artemov and A. Nerode (eds.), Logical Foundations of Computer Science, *Lecture Notes in Computer Science*, 5407 (2009) 1-15.
- [3] M. Busaniche. Decomposition of BL-chains, *Algebra universalis*, 52 (2004) 519–525.
- [4] M. Busaniche, and F. Montagna. Chapter VII: Basic Fuzzy Logic and BL-algebras, *Handbook of Mathematical Fuzzy Logic*. Vol I. Studies in Logic. College Publications. 2011.
- [5] J. L. Castiglioni, M. Menni and W. J. Zuluaga Botero. A representation theorem for integral rigs and its applications to residuated lattices, *Journal of Pure and Applied Algebra*, 220 (10) (2016) 3533–3566.
- [6] A. H. Clifford. Totally ordered commutative semigroups, *Bull. Amer. Math. Soc.*, 64(6) (1958) 305–316.
- [7] A. Di Nola and A. Lettieri. Finite BL-algebras, *Discrete Math.*, 269 (2003) 93–122.
- [8] F. Esteva and L. Godo. Monoidal t-norm based Logic: Towards a logic for left-continuous t-noms, *Fuzzy Sets and Systems*, 124 (2001) 271–288.
- [9] J. B. Hart, L. Rafter and C. Tsinakakis. The Structure of Commutative Residuated Lattices, *International Journal of Algebra and Computation*, 12(4) (2002) 509–524 .
- [10] R. Horčík. On the Structure of Finite Integral Commutative Residuated Chains, *Journal of Logic and Computation*, 21(5) (2011) 717–728.
- [11] R. Horčík and F. Montagna. Archimedean classes in integral commutative residuated chains, *Math. Log. Quart.*, 55(3) (2009) 320–336.
- [12] N. Kehayopulu and M. Tsingelis. Decomposition of Commutative Ordered Semi-groups into Arquimedean Components, *Lobachevskii Journal of Mathematics*, 22 (2006) 27–34.

- [13] S. MacLane and I. Moerdijk. *Sheaves in Geometry and Logic: A First Introduction to Topos Theory*. Universitext, Springer Verlag (1992).
- [14] C. Mureşan. The Reticulation of a Residuated Lattice, *Bulletin mathématique de la Société des Sciences Mathématiques de Roumanie*, 1 (2008) 47–65.
- [15] C. Noguera, F. Esteva and J. Gispert. On some Varieties of MTL-algebras, *Logic Journal of the IGPL*, 13 (2005) 443–466.
- [16] M. Petrík and Th. Vetterlein. Algorithm to generate the Archimedean, finite negative tomonoids, *SCIS&ISIS 2014*, Kitakyushu, Japan, December 3-6, 2014.
- [17] H. P. Sankappanavar and S. Burris. *A course in universal algebra*, Graduate Texts Math, 78 (1981)
- [18] T. Tamura and N. Kimura. On decompositions of a commutative semigroup, *Kodai Math. Sem. Rep.* , 6(4) (1954) 109–112.
- [19] E. Turunen. BL-algebras of basic fuzzy logic, *Mathware & Soft computing*, 6(1) (1999) 49–61.
- [20] I. A. Xarez and J. J. Xarez. Galois Theories of Commutative Semigroups via Semilattices, *Theory and Applications of Categories*, 28(33) (2013) 1153–1169.
- [21] W. Zuluaga Botero. Representation by Sheaves of riRigs (Spanish). PhD Thesis. Universidad Nacional de La Plata. <http://sedici.unlp.edu.ar/handle/10915/54115>. (2016)