

# $q$ -Virasoro algebra and affine Kac-Moody Lie algebras

Hongyan Guo<sup>a,c</sup>, Haisheng Li<sup>a,b1</sup>, Shaobin Tan<sup>a2</sup> and Qing Wang<sup>a3</sup>

<sup>a</sup>School of Mathematical Sciences

Xiamen University, Xiamen 361005, China

<sup>b</sup>Department of Mathematical Sciences

Rutgers University, Camden, NJ 08102, USA

<sup>c</sup>School of Mathematical and Statistical Sciences

University of Alberta, Edmonton T6G 2G1, Canada

## Abstract

We establish a natural connection of the  $q$ -Virasoro algebra  $D_q$  introduced by Belov and Chaltikian with affine Kac-Moody Lie algebras. More specifically, for each abelian group  $S$  together with a one-to-one linear character  $\chi$ , we define an infinite-dimensional Lie algebra  $D_S$  which reduces to  $D_q$  when  $S = \mathbb{Z}$ . Guided by the theory of equivariant quasi modules for vertex algebras, we introduce another Lie algebra  $\mathfrak{g}_S$  with  $S$  as an automorphism group and we prove that  $D_S$  is isomorphic to the  $S$ -covariant algebra of the affine Lie algebra  $\widehat{\mathfrak{g}_S}$ . We then relate restricted  $D_S$ -modules of level  $\ell \in \mathbb{C}$  to equivariant quasi modules for the vertex algebra  $V_{\widehat{\mathfrak{g}_S}}(\ell, 0)$  associated to  $\widehat{\mathfrak{g}_S}$  with level  $\ell$ . Furthermore, we show that if  $S$  is a finite abelian group of order  $2l + 1$ ,  $D_S$  is isomorphic to the affine Kac-Moody algebra of type  $B_l^{(1)}$ .

## 1 Introduction

In an attempt to find a meaningful  $q$ -analog of the Virasoro algebra, a certain infinite-dimensional Lie algebra  $D_q$  over the field  $\mathbb{C}$  of complex numbers was introduced and studied in [BC] (cf. [N]), where  $q$  is a nonzero complex number. In case that  $q$  is not a root of unity, Lie algebra  $D_q$  was studied in [GLTW] by using vertex algebra theory in an essential way. It was proved therein that the category of (suitably defined) restricted  $D_q$ -modules of a fixed level  $\ell \in \mathbb{C}$  is canonically isomorphic to the category of equivariant quasi modules for the vertex algebra  $V_{\widehat{\mathcal{D}}}(\ell, 0)$  associated to the affine Lie algebra  $\widehat{\mathcal{D}}$  with level  $\ell$  of a certain (infinite-dimensional) Lie algebra  $\mathcal{D}$  with a symmetric invariant bilinear form.

In the current paper, we continue this study to deal with the case where  $q$  is a root of unity. Actually, we work with a slightly more general setting. Let  $S$  be an abelian group with a one-to-one linear character  $\chi$ . (This implies that if  $S$  is finite, it must be cyclic.) Then we define an infinite-dimensional Lie algebra  $D_S$  with

<sup>1</sup>Partially supported by China NSF grants (Nos.11471268, 11571391)

<sup>2</sup>Partially supported by China NSF grants (Nos.11471268, 11531004)

<sup>3</sup>Partially supported by China NSF grants (Nos.11531004, 11622107), Natural Science Foundation of Fujian Province (No. 2016J06002)

generators  $D^\alpha(n)$  for  $\alpha \in S$ ,  $n \in \mathbb{Z}$ , subject to certain relations. In case  $S = \mathbb{Z}$  with  $q = \chi(1)$ ,  $D_S$  reduces to  $D_q$ . Guided by the theory of equivariant quasi modules for vertex algebras (see [L2, L3]), we introduce another Lie algebra  $\mathfrak{g}_S$  with  $S$  as an automorphism group and we prove that  $D_S$  is isomorphic to the  $S$ -covariant algebra of the affine Lie algebra  $\widehat{\mathfrak{g}}_S$ . We then relate restricted  $D_S$ -modules of level  $\ell \in \mathbb{C}$  to equivariant quasi modules for the vertex algebra  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$  associated to  $\widehat{\mathfrak{g}}_S$  with level  $\ell$ . Furthermore, we show that if  $S$  is a finite abelian group of order  $2l + 1$ ,  $D_S$  is actually isomorphic to the affine Kac-Moody algebra of type  $B_l^{(1)}$ .

Now, we give a more detailed account of the contents of this paper. Let  $S$  be an abelian group with a one-to-one linear character  $\chi : S \rightarrow \mathbb{C}^\times$ . We first define an infinite-dimensional Lie algebra  $D_S$  with generators  $D^\alpha(n)$  for  $\alpha \in S$ ,  $n \in \mathbb{Z}$ , subject to certain relations (see Definition 2.2). Following the common practice in vertex algebra theory, we form a generating function  $D^\alpha(x) = \sum_{n \in \mathbb{Z}} D^\alpha(n)x^{-n-1}$  for each  $\alpha \in S$  and then calculate the commutator of two general generating functions. The commutator relations for the generating functions and theory of equivariant quasi modules for vertex algebras indicate that  $D_S$  is closely related to the affine Lie algebra of some (possibly infinite-dimensional) Lie algebra with a symmetric invariant bilinear form. Motivated by this, for the abelian group  $S$  we introduce another Lie algebra  $\mathfrak{g}_S$  with generators  $d^{\alpha, \beta}$  for  $\alpha, \beta \in S$ , subject to relations  $d^{-\alpha, \beta} = -d^{\alpha, \beta}$  and

$$\begin{aligned} [d^{\alpha, \beta}, d^{\mu, \nu}] &= \delta_{\alpha+\mu, \nu-\beta} d^{\alpha+\mu, -\alpha+\nu} - \delta_{\alpha+\mu, \beta-\nu} d^{\alpha+\mu, \alpha+\nu} \\ &\quad - \delta_{\alpha-\mu, \nu-\beta} d^{\alpha-\mu, \nu-\alpha} + \delta_{\alpha-\mu, \beta-\nu} d^{\alpha-\mu, \alpha+\nu} \end{aligned}$$

for  $\alpha, \beta, \mu, \nu \in S$ . It is proved that  $S$  acts on  $\mathfrak{g}_S$  as an automorphism group with

$$\gamma \cdot d^{\alpha, \beta} = d^{\alpha, \beta+\gamma} \quad \text{for } \alpha, \beta, \gamma \in S.$$

We also equip  $\mathfrak{g}_S$  with a symmetric invariant bilinear form. Then we prove that  $D_S$  is isomorphic to the so-called  $S$ -covariant algebra of the affine Lie algebra  $\widehat{\mathfrak{g}}_S$  as we explain next.

It is a classical fact that for any Lie algebra  $K$  with an automorphism group  $G$ , all the  $G$ -fixed points in  $K$  form a Lie subalgebra  $K^G$ . An important result in Kac-Moody algebra theory (see [K]) is that twisted affine Kac-Moody algebras are isomorphic to the fixed-point subalgebras of untwisted affine Kac-Moody algebras with respect to Dynkin diagram automorphisms. Now, assume that  $G$  is a (possibly infinite) automorphism group of  $K$  such that for any  $a, b \in K$ ,

$$[ga, b] = 0 \quad \text{for all but finitely many } g \in G.$$

Under this condition, one can associate another Lie algebra to the pair  $(K, G)$  as follows (see [G-K-K], [L3]). Define a new multiplicative operation  $[\cdot, \cdot]_G$  on  $K$  by

$$[a, b]_G = \sum_{g \in G} [ga, b].$$

Set

$$I_G = \text{span}\{a - ga \mid a \in K, g \in G\}.$$

Then  $I_G$  is a two-sided ideal of the non-associative algebra  $(K, [\cdot, \cdot]_G)$  and the quotient algebra by  $I_G$  is a Lie algebra, which is denoted by  $K/G$  and called the  $G$ -covariant algebra of  $K$ . In case that  $G$  is finite, one can show that Lie algebras  $K^G$  and  $K/G$  are isomorphic.

Now that  $D_S$  is isomorphic to the  $S$ -covariant algebra of the affine Lie algebra  $\widehat{\mathfrak{g}}_S$ , by applying a result of [L3] we show that the category of restricted  $D_S$ -modules of a fixed level  $\ell$  is naturally isomorphic to the category of equivariant quasi  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$ -modules. This generalizes the corresponding result of [GLTW]. Then we continue to determine the Lie algebra  $\mathfrak{g}_S$  for  $S$  a finite abelian group. Assuming  $|S| = 2l + 1$ , we prove that  $\mathfrak{g}_S$  is a simple Lie algebra of type  $B_l$ . It follows that  $D_S$  is isomorphic to the  $S$ -covariant algebra of the affine Kac-Moody algebra of type  $B_l^{(1)}$ . Using this and a result of Kac (see [K]) we show that  $D_S$  is actually isomorphic to the affine Kac-Moody algebra of type  $B_l^{(1)}$ . For the case with  $S$  of an even order, set  $S^0 = \{\alpha \in S \mid 2\alpha = 0\}$  and let  $I$  be the subspace of  $\mathfrak{g}_S$ , linearly spanned by the following vectors

$$d^{\alpha+\gamma, \beta+\gamma} - d^{\alpha, \beta} \quad (\alpha, \beta \in S, \gamma \in S^0).$$

We prove that  $I$  is an ideal of  $\mathfrak{g}_S$  and the quotient algebra  $\mathfrak{g}_S/I$  is a direct sum of simple Lie algebras of type  $B$  or a direct sum of simple Lie algebras of type  $D$  (see Proposition 3.10).

This paper is organized as follows: In Section 2, we introduce Lie algebras  $D_S$  and  $\mathfrak{g}_S$ , and prove that  $D_S$  is isomorphic to the  $S$ -covariant algebra of  $\widehat{\mathfrak{g}}_S$ . In Section 3, we identify  $\mathfrak{g}_S$  as a subalgebra of the general linear algebra  $\mathfrak{gl}_S$  and prove that if  $|S| = 2l + 1$ , then  $\mathfrak{g}_S$  is isomorphic to  $B_l$  and  $D_S$  is isomorphic to the affine Kac-Moody algebra of type  $B_l^{(1)}$ .

## 2 Lie algebras $D_S$ , $\mathfrak{g}_S$ , and equivariant quasi modules for $\Gamma$ -vertex algebras

In this section, to any abelian group  $S$  with a linear character  $\chi$  we associate two Lie algebras  $D_S$  and  $\mathfrak{g}_S$ . Furthermore, we equip Lie algebra  $\mathfrak{g}_S$  with a symmetric invariant bilinear form and show that the vertex algebra  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$  associated to the affine Lie algebra  $\widehat{\mathfrak{g}}_S$  of level  $\ell$  is a  $\Gamma$ -vertex algebra with  $\Gamma = S$ . Then we establish an isomorphism between the category of restricted  $D_S$ -modules of level  $\ell$  and that of equivariant quasi  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$ -modules.

First of all, we fix some basic notations. Throughout this paper,  $\mathbb{C}^\times$  denotes the multiplicative group of nonzero complex numbers (with  $\mathbb{C}$  denoting the complex number field), and the symbols  $x, y, x_1, x_2, \dots$  denote mutually commuting independent formal variables. All vector spaces in this paper are considered to be over  $\mathbb{C}$ . For a vector space  $U$ ,  $U((x))$  is the vector space of lower truncated integer power

series in  $x$  with coefficients in  $U$ ,  $U[[x]]$  is the vector space of nonnegative integer power series in  $x$  with coefficients in  $U$ , and  $U[[x, x^{-1}]]$  is the vector space of doubly infinite integer power series in  $x$  with coefficients in  $U$ .

We recall the  $q$ -analog of the Virasoro algebra from [BC] (cf. [N]). Let  $q$  be a nonzero complex number. By definition,  $D_q$  is the Lie algebra with generators  $\mathbf{c}$  and  $D^\alpha(n)$  ( $\alpha, n \in \mathbb{Z}$ ), subjects to relations

$$D^{-\alpha}(n) = -D^\alpha(n),$$

$$\begin{aligned} [D^\alpha(n), D^\beta(m)] &= (q - q^{-1})[\alpha m - \beta n]_q D^{\alpha+\beta}(m+n) \\ &\quad - (q - q^{-1})[\alpha m + \beta n]_q D^{\alpha-\beta}(m+n) \\ &\quad + ([m]_{q^{\alpha+\beta}} - [m]_{q^{\alpha-\beta}}) \delta_{m+n,0} \mathbf{c} \end{aligned} \quad (2.1)$$

for  $m, n, \alpha, \beta \in \mathbb{Z}$ , where  $\mathbf{c}$  is central and  $[n]_q$  is the  $q$ -integer defined by

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

**Remark 2.1.** Here, it is understood that  $[n]_q = n$  for  $q = 1$  and  $[n]_q = n(-1)^{n-1}$  for  $q = -1$ . This is consistent with the fact that

$$\lim_{q \rightarrow 1} [n]_q = n \quad \text{and} \quad \lim_{q \rightarrow -1} [n]_q = n(-1)^{n-1}.$$

The following are some simple properties of the  $q$ -integers:

$$[-n]_q = -[n]_q, \quad [n]_{q^{-1}} = [n]_q, \quad [m]_{q^n} = \frac{[mn]_q}{[n]_q}. \quad (2.2)$$

We generalize the Lie algebra  $D_q$  as follows:

**Definition 2.2.** Let  $S$  be an abelian group with a group homomorphism  $\chi : S \rightarrow \mathbb{C}^\times$ , i.e., a linear character of  $S$ . Define  $D_S$  to be the Lie algebra with generators  $D^\alpha(n)$  for  $\alpha \in S$ ,  $n \in \mathbb{Z}$ , subjects to the following relations

$$D^{-\alpha}(n) = -D^\alpha(n),$$

$$\begin{aligned} [D^\alpha(n), D^\beta(m)] &= (\chi(m\alpha - n\beta) - \chi(n\beta - m\alpha)) D^{\alpha+\beta}(m+n) \\ &\quad - (\chi(m\alpha + n\beta) - \chi(-m\alpha - n\beta)) D^{\alpha-\beta}(m+n) \\ &\quad + ([m]_{\chi(\alpha+\beta)} - [m]_{\chi(\alpha-\beta)}) \delta_{m+n,0} \mathbf{c} \end{aligned} \quad (2.3)$$

for  $m, n \in \mathbb{Z}$ ,  $\alpha, \beta \in S$ , where  $\mathbf{c}$  is central.

**Remark 2.3.** Consider  $S = \mathbb{Z}$ . Let  $q$  be a nonzero complex number and let  $\chi$  be the linear character of  $\mathbb{Z}$  given by  $\chi(n) = q^n$  for  $n \in \mathbb{Z}$ . Then  $D_S$  is reduced to the Lie algebra  $D_q$ . In the case that  $q$  is not a root of unity, Lie algebra  $D_q$  was studied in [GLTW] in the context of vertex algebras and their quasi modules.

From now on, we assume that  $\chi : S \rightarrow \mathbb{C}^\times$  is one-to-one. Then, for  $\alpha \in S$ ,  $\chi(\alpha) = 1$  if and only if  $\alpha = 0$ . For  $\alpha \in S$ , set

$$D^\alpha(x) = \sum_{n \in \mathbb{Z}} D^\alpha(n) x^{-n-1}. \quad (2.4)$$

Furthermore, we set

$$\tilde{D}^\alpha(x) = D^\alpha(x) + (1 - \delta_{2\alpha,0}) \frac{1}{\chi(\alpha) - \chi(-\alpha)} \mathbf{c} x^{-1}, \quad (2.5)$$

where we use the one-to-one assumption on  $\chi$ .

By a straightforward calculation (cf. [GLTW]) we have:

**Lemma 2.4.** *The defining relations of  $D_S$  are equivalent to*

$$[\mathbf{c}, D_S] = 0, \quad \tilde{D}^{-\alpha}(x) = -\tilde{D}^\alpha(x), \quad (2.6)$$

$$\begin{aligned} [\tilde{D}^\alpha(x_1), \tilde{D}^\beta(x_2)] &= \chi(-\alpha) \tilde{D}^{\alpha+\beta}(\chi(-\alpha)x_2)x_1^{-1} \delta\left(\frac{\chi(-\alpha-\beta)x_2}{x_1}\right) \\ &\quad - \chi(\alpha) \tilde{D}^{\alpha+\beta}(\chi(\alpha)x_2)x_1^{-1} \delta\left(\frac{\chi(\alpha+\beta)x_2}{x_1}\right) \\ &\quad - \chi(-\alpha) \tilde{D}^{\alpha-\beta}(\chi(-\alpha)x_2)x_1^{-1} \delta\left(\frac{\chi(\beta-\alpha)x_2}{x_1}\right) \\ &\quad + \chi(\alpha) \tilde{D}^{\alpha-\beta}(\chi(\alpha)x_2)x_1^{-1} \delta\left(\frac{\chi(\alpha-\beta)x_2}{x_1}\right) \\ &\quad - \chi(\alpha-\beta) \delta_{2(\alpha-\beta),0} \frac{\partial}{\partial x_2} x_1^{-1} \delta\left(\frac{\chi(\alpha-\beta)x_2}{x_1}\right) \mathbf{c} \\ &\quad + \chi(\alpha+\beta) \delta_{2(\alpha+\beta),0} \frac{\partial}{\partial x_2} x_1^{-1} \delta\left(\frac{\chi(\alpha+\beta)x_2}{x_1}\right) \mathbf{c} \end{aligned} \quad (2.7)$$

for  $\alpha, \beta \in S$ .

For  $\alpha, \beta \in S$ , set

$$D^{\alpha,\beta}(x) = \chi(\beta) \tilde{D}^\alpha(\chi(\beta)x). \quad (2.8)$$

From (2.7), we have

$$\begin{aligned} &[D^{\alpha,\beta}(x_1), D^{\mu,\nu}(x_2)] \\ &= \chi(\beta+\nu) [\tilde{D}^\alpha(\chi(\beta)x_1), \tilde{D}^\mu(\chi(\nu)x_2)] \\ &= D^{\alpha+\mu,\nu-\alpha}(x_2)x_1^{-1} \delta\left(\frac{\chi(\nu-\alpha)x_2}{\chi(\mu+\beta)x_1}\right) - D^{\alpha+\mu,\alpha+\nu}(x_2)x_1^{-1} \delta\left(\frac{\chi(\alpha+\nu)x_2}{\chi(\beta-\mu)x_1}\right) \\ &\quad - D^{\alpha-\mu,\nu-\alpha}(x_2)x_1^{-1} \delta\left(\frac{\chi(\nu-\alpha)x_2}{\chi(\beta-\mu)x_1}\right) + D^{\alpha-\mu,\alpha+\nu}(x_2)x_1^{-1} \delta\left(\frac{\chi(\alpha+\nu)x_2}{\chi(\mu+\beta)x_1}\right) \\ &\quad - \chi(\alpha-\mu) \delta_{2(\alpha-\mu),0} \frac{\partial}{\partial x_2} x_1^{-1} \delta\left(\frac{\chi(\alpha-\mu+\nu-\beta)x_2}{x_1}\right) \mathbf{c} \\ &\quad + \chi(\alpha+\mu) \delta_{2(\alpha+\mu),0} \frac{\partial}{\partial x_2} x_1^{-1} \delta\left(\frac{\chi(\alpha+\mu+\nu-\beta)x_2}{x_1}\right) \mathbf{c} \end{aligned} \quad (2.9)$$

for  $\alpha, \beta, \mu, \nu \in S$ .

Using this we get another characterization of the Lie algebra  $D_S$ .

**Proposition 2.5.** *Lie algebra  $D_S$  is isomorphic to the Lie algebra  $\mathcal{L}$  with generators  $\mathbf{c}$  and  $D^{\alpha, \beta}(n)$  for  $\alpha, \beta \in S$ ,  $n \in \mathbb{Z}$ , subject to relations*

$$[\mathbf{c}, \mathcal{L}] = 0, \quad D^{-\alpha, \beta}(n) = -D^{\alpha, \beta}(n), \quad D^{\alpha, \beta + \gamma}(n) = \chi(\gamma)^{-n} D^{\alpha, \beta}(n) \quad (2.10)$$

and the relation (2.9) with  $D^{\alpha, \beta}(x) = \sum_{n \in \mathbb{Z}} D^{\alpha, \beta}(n) x^{-n-1}$ .

*Proof.* Given Lie algebra  $D_S$ , with  $D^{\alpha, \beta}(x)$  defined in (2.8), we see that (2.10) and (2.9) hold. It follows that there is a Lie algebra homomorphism  $\theta$  from  $\mathcal{L}$  onto  $D_S$  sending  $D^{\alpha, \beta}(x)$  to  $\chi(\beta) \tilde{D}^\alpha(\chi(\beta)x)$  for  $\alpha, \beta \in S$ . On the other hand, for Lie algebra  $\mathcal{L}$  we have

$$D^{\alpha, \beta}(x) = \chi(\beta) D^{\alpha, 0}(\chi(\beta)x) \quad \text{and} \quad D^{-\alpha, 0}(x) = -D^{\alpha, 0}(x) \quad \text{for } \alpha, \beta \in S.$$

It is clear that relation (2.9) is equivalent to (2.7) with  $\tilde{D}^\alpha(x) = D^{\alpha, 0}(x)$  for  $\alpha \in S$ . Then there exists a natural Lie algebra homomorphism from  $D_S$  onto  $\mathcal{L}$ , sending  $\tilde{D}^\alpha(x)$  to  $D^{\alpha, 0}(x)$  for  $\alpha \in S$ . Consequently,  $\theta$  is a Lie algebra isomorphism.  $\square$

**Remark 2.6.** Set

$$S^0 = \{\alpha \in S \mid 2\alpha = 0\}, \quad (2.11)$$

a subgroup of  $S$ . Note that if  $S^0 = \{0\}$ , relation (2.9) becomes

$$\begin{aligned} & [D^{\alpha, \beta}(x_1), D^{\mu, \nu}(x_2)] \\ &= D^{\alpha + \mu, \nu - \alpha}(x_2) x_1^{-1} \delta \left( \frac{\chi(\nu - \alpha)x_2}{\chi(\mu + \beta)x_1} \right) - D^{\alpha + \mu, \alpha + \nu}(x_2) x_1^{-1} \delta \left( \frac{\chi(\alpha + \nu)x_2}{\chi(\beta - \mu)x_1} \right) \\ & \quad - D^{\alpha - \mu, \nu - \alpha}(x_2) x_1^{-1} \delta \left( \frac{\chi(\nu - \alpha)x_2}{\chi(\beta - \mu)x_1} \right) + D^{\alpha - \mu, \alpha + \nu}(x_2) x_1^{-1} \delta \left( \frac{\chi(\alpha + \nu)x_2}{\chi(\mu + \beta)x_1} \right) \\ & \quad + (\delta_{\alpha, -\mu} - \delta_{\alpha, \mu}) \frac{\partial}{\partial x_2} x_1^{-1} \delta \left( \frac{\chi(\nu - \beta)x_2}{x_1} \right) \mathbf{c} \end{aligned} \quad (2.12)$$

for  $\alpha, \beta, \mu, \nu \in S$ .

In the following, we shall relate  $D_S$  with the affine algebra of a new Lie algebra.

**Definition 2.7.** Let  $S$  be an abelian group as before. Define  $\mathfrak{g}_S$  to be the Lie algebra with generators  $d^{\alpha, \beta}$  for  $\alpha, \beta \in S$ , subject to relations

$$d^{-\alpha, \beta} = -d^{\alpha, \beta}, \quad (2.13)$$

$$\begin{aligned} [d^{\alpha, \beta}, d^{\mu, \nu}] &= \delta_{\alpha + \mu, \nu - \beta} d^{\alpha + \mu, -\alpha + \nu} - \delta_{\alpha + \mu, \beta - \nu} d^{\alpha + \mu, \alpha + \nu} \\ & \quad - \delta_{\alpha - \mu, \nu - \beta} d^{\alpha - \mu, \nu - \alpha} + \delta_{\alpha - \mu, \beta - \nu} d^{\alpha - \mu, \alpha + \nu} \end{aligned} \quad (2.14)$$

for  $\alpha, \beta, \mu, \nu \in S$ .

The following result can be found in [L3] (Lemma 4.1) (cf. [G-K-K]):

**Lemma 2.8.** *Let  $K$  be any Lie algebra with a symmetric invariant bilinear form  $\langle \cdot, \cdot \rangle$  and let  $G$  be an automorphism group of  $K$ , preserving  $\langle \cdot, \cdot \rangle$ , such that for any  $a, b \in K$ ,*

$$[ga, b] = 0 \quad \text{and} \quad \langle ga, b \rangle = 0 \quad \text{for all but finitely many } g \in G.$$

*Define a new multiplicative operation  $[\cdot, \cdot]_G$  and a new bilinear form  $\langle \cdot, \cdot \rangle_G$  on  $K$  by*

$$[a, b]_G = \sum_{g \in G} [ga, b] \quad \text{and} \quad \langle a, b \rangle_G = \sum_{g \in G} \langle ga, b \rangle.$$

*Set*

$$I_G = \text{span}\{a - ga \mid a \in K, g \in G\}.$$

*Then  $I_G$  is a two-sided ideal of the non-associative algebra  $(K, [\cdot, \cdot]_G)$  and the quotient algebra modulo  $I_G$  is a Lie algebra, which is denoted by  $K/G$  and called the  $G$ -covariant algebra of  $K$ . Furthermore,  $\langle \cdot, \cdot \rangle_G$  reduces to a symmetric invariant bilinear form on  $K/G$ .*

The following is a more explicit construction of the Lie algebra  $\mathfrak{g}_S$ :

**Proposition 2.9.** *Let  $K$  be a vector space with a basis  $\{F_{\alpha, \beta} \mid \alpha, \beta \in S\}$ . Define a multiplicative operation  $*$  on  $K$  by*

$$\begin{aligned} F_{\alpha, \beta} * F_{\mu, \nu} = & \delta_{\alpha+\mu, \nu-\beta} F_{\alpha+\mu, -\alpha+\nu} - \delta_{\alpha+\mu, \beta-\nu} F_{\alpha+\mu, \alpha+\nu} \\ & - \delta_{\alpha-\mu, \nu-\beta} F_{\alpha-\mu, \nu-\alpha} + \delta_{\alpha-\mu, \beta-\nu} F_{\alpha-\mu, \alpha+\nu} \end{aligned}$$

*for  $\alpha, \beta, \mu, \nu \in S$ . Set*

$$J = \text{span}\{F_{-\alpha, \beta} + F_{\alpha, \beta} \mid \alpha, \beta \in S\}.$$

*Then  $J$  is a two-sided ideal of the non-associative algebra  $(K, *)$  and the quotient algebra  $K/J$  is a Lie algebra. Furthermore,  $\mathfrak{g}_S$  is isomorphic to the Lie algebra  $K/J$  with  $d^{\alpha, \beta}$  corresponding to  $F_{\alpha, \beta} + J$  for  $\alpha, \beta \in S$ .*

*Proof.* We shall apply Lemma 2.8. First, define an operation “ $\cdot$ ” on  $K$  by

$$F_{\alpha, \beta} \cdot F_{\mu, \nu} = \delta_{\nu, \alpha+\beta+\mu} F_{\alpha+\mu, \beta+\mu} \tag{2.15}$$

for  $\alpha, \beta, \mu, \nu \in S$ . It is straightforward to show that  $(K, \cdot)$  is an associative algebra. Then  $K$  is naturally a Lie algebra, where

$$\begin{aligned} [F_{\alpha, \beta}, F_{\mu, \nu}] &= \delta_{\nu, \alpha+\beta+\mu} F_{\alpha+\mu, \beta+\mu} - \delta_{\beta, \mu+\nu+\alpha} F_{\mu+\alpha, \nu+\alpha} \\ &= \delta_{\alpha+\mu, \nu-\beta} F_{\alpha+\mu, -\alpha+\nu} - \delta_{\alpha+\mu, \beta-\nu} F_{\alpha+\mu, \alpha+\nu} \end{aligned}$$

for  $\alpha, \beta, \mu, \nu \in S$ . Let  $\theta$  be the linear endomorphism of  $K$  defined by

$$\theta(F_{\alpha,\beta}) = F_{-\alpha,\beta} \quad \text{for } \alpha, \beta \in S.$$

It is straightforward to show that  $\theta$  is an order 2 anti-automorphism of the associative algebra  $K$ . Then  $-\theta$  is an order 2 automorphism of the Lie algebra  $K$ . From Lemma 2.8, we have

$$\begin{aligned} [F_{\alpha,\beta}, F_{\mu,\nu}]_{(-\theta)} &= \delta_{\alpha+\mu,\nu-\beta} F_{\alpha+\mu,-\alpha+\nu} - \delta_{\alpha+\mu,\beta-\nu} F_{\alpha+\mu,\alpha+\nu} \\ &\quad - \delta_{-\alpha+\mu,\nu-\beta} F_{-\alpha+\mu,\alpha+\nu} + \delta_{-\alpha+\mu,\beta-\nu} F_{-\alpha+\mu,\nu-\alpha} \end{aligned}$$

for  $\alpha, \beta, \mu, \nu \in S$ . Then the first part of this lemma follows immediately. As for the second part, we first have a Lie algebra homomorphism  $\psi$  from  $\mathfrak{g}_S$  onto  $K/J$  with  $\psi(d^{\alpha,\beta}) = F_{\alpha,\beta} + J$  for  $\alpha, \beta \in S$ . By using a basis of  $K/J$ , we see that  $\psi$  is actually an isomorphism.  $\square$

From Proposition 2.9 we readily have:

**Lemma 2.10.** *The abelian group  $S$  acts on  $\mathfrak{g}_S$  as an automorphism group with*

$$\gamma \cdot d^{\alpha,\beta} = d^{\alpha,\beta+\gamma} \quad (2.16)$$

for  $\gamma, \alpha, \beta \in S$ .

Now, let  $\chi : S \rightarrow \mathbb{C}^\times$  be a linear character as in the definition of  $D_S$ . It is straightforward to see that  $\mathfrak{g}_S$  admits a bilinear form  $\langle \cdot, \cdot \rangle_\chi$  such that

$$\langle d^{\alpha,\beta}, d^{\mu,\nu} \rangle_\chi = \chi(\alpha + \mu) \delta_{2(\alpha+\mu),0} \delta_{\alpha+\mu,\beta-\nu} - \chi(\alpha - \mu) \delta_{2(\alpha-\mu),0} \delta_{\alpha-\mu,\beta-\nu} \quad (2.17)$$

for  $\alpha, \beta, \mu, \nu \in S$ . We have:

**Lemma 2.11.** *The defined bilinear form  $\langle \cdot, \cdot \rangle_\chi$  on Lie algebra  $\mathfrak{g}_S$  is symmetric and invariant, and the group action of  $S$  preserves  $\langle \cdot, \cdot \rangle_\chi$ . Furthermore, for any  $\alpha, \beta, \mu, \nu \in S$ ,*

$$[d^{\alpha,\beta+\gamma}, d^{\mu,\nu}] = 0 \quad \text{and} \quad \langle d^{\alpha,\beta+\gamma}, d^{\mu,\nu} \rangle_\chi = 0 \quad (2.18)$$

for all but finitely many  $\gamma \in S$ .

*Proof.* We shall use Lemma 2.8. First, recall the associative algebra  $K$  from the proof of Proposition 2.9, where  $K$  has a basis  $\{F_{\alpha,\beta} \mid \alpha, \beta \in S\}$  with

$$F_{\alpha,\beta} \cdot F_{\mu,\nu} = \delta_{\nu,\alpha+\beta+\mu} F_{\alpha+\mu,\beta+\mu}$$

for  $\alpha, \beta, \mu, \nu \in S$ . We define a bilinear form  $\langle \cdot, \cdot \rangle$  on  $K$  by

$$\langle F_{\alpha,\beta}, F_{\mu,\nu} \rangle = \chi(\alpha + \mu) \delta_{2(\alpha+\mu),0} \delta_{\alpha+\mu,\beta-\nu}$$



for  $\alpha, \beta, \mu, \nu \in S$ . It can be readily seen that  $\langle \cdot, \cdot \rangle$  is symmetric. On the other hand,  $\langle \cdot, \cdot \rangle$  is also associative because

$$\langle F_{\alpha, \beta}, F_{\mu, \nu} \rangle = f(F_{\alpha, \beta} \cdot F_{\mu, \nu})$$

for  $\alpha, \beta, \mu, \nu \in S$ , where  $f$  is the linear functional on  $K$  given by

$$f(F_{\alpha, \beta}) = \chi(\alpha) \delta_{2\alpha, 0} \quad \text{for } \alpha, \beta \in S.$$

Consequently,  $\langle \cdot, \cdot \rangle$  is a symmetric and invariant bilinear form on  $K$  viewed as a Lie algebra. It is also clear that the Lie algebra automorphism  $-\theta$  of  $K$  preserves  $\langle \cdot, \cdot \rangle$ . For  $\alpha, \beta, \mu, \nu \in S$ , we have

$$\begin{aligned} \langle F_{\alpha, \beta}, F_{\mu, \nu} \rangle_{\langle -\theta \rangle} &= \langle F_{\alpha, \beta}, F_{\mu, \nu} \rangle - \langle F_{-\alpha, \beta}, F_{\mu, \nu} \rangle \\ &= \chi(\alpha + \mu) \delta_{2(\alpha + \mu), 0} \delta_{\alpha + \mu, \beta - \nu} - \chi(\alpha - \mu) \delta_{2(\alpha - \mu), 0} \delta_{\alpha - \mu, \beta - \nu}. \end{aligned}$$

Then the first part of this lemma follows immediately from Lemma 2.8. The second part is clear.  $\square$

**Remark 2.12.** Note that in case that  $S^0 = \{0\}$  (where  $S^0 = \{\alpha \in S \mid 2\alpha = 0\}$ ), we have

$$\langle d^{\alpha, \beta}, d^{\mu, \nu} \rangle_{\chi} = (\delta_{\alpha, -\mu} - \delta_{\alpha, \mu}) \delta_{\beta, \nu} \quad (2.19)$$

for  $\alpha, \beta, \mu, \nu \in S$ , recalling (2.17).

Let  $(S, \chi)$  be given as in the definition of  $D_S$ . We have a Lie algebra  $\mathfrak{g}_S$  with a symmetric invariant bilinear form  $\langle \cdot, \cdot \rangle_{\chi}$ . Associated to the pair  $(\mathfrak{g}_S, \langle \cdot, \cdot \rangle_{\chi})$ , we have an affine Lie algebra

$$\widehat{\mathfrak{g}}_S = \mathfrak{g}_S \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}\mathbf{k}.$$

Naturally,  $S$  acts on  $\widehat{\mathfrak{g}}_S$  as an automorphism group, where

$$\gamma \cdot \mathbf{k} = \mathbf{k}, \quad \gamma \cdot (d^{\alpha, \beta}(n)) = \chi(\gamma)^n d^{\alpha, \beta + \gamma}(n) \quad (2.20)$$

for  $\gamma, \alpha, \beta \in S$ ,  $n \in \mathbb{Z}$ .

Now, we are in a position to present our first main result.

**Theorem 2.13.** *The linear map  $\pi : \widehat{\mathfrak{g}}_S \rightarrow D_S$ , defined by  $\pi(\mathbf{k}) = \mathbf{c}$  and*

$$\pi(d^{\alpha, \beta}(x)) = D^{\alpha, \beta}(x) = \chi(\beta) \tilde{D}^{\alpha}(\chi(\beta)x) \quad \text{for } \alpha, \beta \in S,$$

*gives rise to a Lie algebra isomorphism from the covariant algebra  $\widehat{\mathfrak{g}}_S/S$  to  $D_S$ .*

*Proof.* Recall the defining relations (2.6) and (2.7) of  $D_S$ . As the relations  $d^{-\alpha, \beta} = -d^{\alpha, \beta}$  for  $\alpha, \beta \in S$  hold in  $\mathfrak{g}_S$ , the linear map  $\pi$  is well defined. On the other hand, note that  $\widehat{\mathfrak{g}}_S/S$  as a vector space is the quotient of  $\widehat{\mathfrak{g}}_S$  by the relations

$$d^{\alpha, \beta + \gamma}(x) = \chi(\gamma) d^{\alpha, \beta}(\chi(\gamma)x) \quad \text{for } \alpha, \beta, \gamma \in S.$$

Then  $\pi$  is a linear isomorphism. Let  $\alpha, \beta, \mu, \nu \in S$ ,  $m, n \in \mathbb{Z}$ . We have

$$\begin{aligned}
& [d^{\alpha, \beta}(m), d^{\mu, \nu}(n)]_S \\
&= \sum_{\gamma \in S} \chi(\gamma)^m [d^{\alpha, \beta + \gamma}(m), d^{\mu, \nu}(n)] \\
&= \sum_{\gamma \in S} \chi(\gamma)^m (\delta_{\alpha + \mu, \nu - \beta - \gamma} d^{\alpha + \mu, -\alpha + \nu}(m + n) - \delta_{\alpha + \mu, \beta + \gamma - \nu} d^{\alpha + \mu, \alpha + \nu}(m + n) \\
&\quad - \delta_{\alpha - \mu, \nu - \beta - \gamma} d^{\alpha - \mu, \nu - \alpha}(m + n) + \delta_{\alpha - \mu, \beta + \gamma - \nu} d^{\alpha - \mu, \alpha + \nu}(m + n)) \\
&\quad + m \delta_{m+n, 0} \mathbf{k} \sum_{\gamma \in S} \chi(\gamma)^m (\chi(\alpha + \mu) \delta_{2(\alpha + \mu), 0} \delta_{\alpha + \mu, \beta + \gamma - \nu} - \chi(\alpha - \mu) \delta_{2(\alpha - \mu), 0} \delta_{\alpha - \mu, \beta + \gamma - \nu}) \\
&= \chi(\nu - \alpha - \beta - \mu)^m d^{\alpha + \mu, -\alpha + \nu}(m + n) - \chi(\alpha + \mu + \nu - \beta)^m d^{\alpha + \mu, \alpha + \nu}(m + n) \\
&\quad - \chi(\mu + \nu - \alpha - \beta)^m d^{\alpha - \mu, \nu - \alpha}(m + n) + \chi(\alpha + \nu - \beta - \mu)^m d^{\alpha - \mu, \alpha + \nu}(m + n) \\
&\quad + m \delta_{m+n, 0} \mathbf{k} (\chi(\alpha + \mu) \delta_{2(\alpha + \mu), 0} \chi(\alpha + \mu + \nu - \beta)^m - \chi(\alpha - \mu) \delta_{2(\alpha - \mu), 0} \chi(\alpha + \nu - \beta - \mu)^m).
\end{aligned}$$

Comparing this with (2.9), we see that  $\pi$  is a Lie algebra homomorphism, and hence it is an isomorphism.  $\square$

Recall the following notion of quasi module for a vertex algebra from [L2]:

**Definition 2.14.** Let  $V$  be a vertex algebra. A *quasi  $V$ -module* is a vector space  $W$  equipped with a linear map

$$\begin{aligned}
Y_W(\cdot, x) : V &\rightarrow \text{Hom}(W, W((x))) \subset (\text{End} W)[[x, x^{-1}]], \\
v &\mapsto Y_W(v, x)
\end{aligned}$$

satisfying the conditions that

$$Y_W(\mathbf{1}, x) = \mathbf{1}_W \quad (\text{the identity operator on } W)$$

and that for any  $u, v \in V$ , there exists a nonzero polynomial  $p(x_1, x_2)$  such that

$$\begin{aligned}
& x_0^{-1} \delta \left( \frac{x_1 - x_2}{x_0} \right) p(x_1, x_2) Y_W(u, x_1) Y_W(v, x_2) \\
& \quad - x_0^{-1} \delta \left( \frac{x_2 - x_1}{-x_0} \right) p(x_1, x_2) Y_W(v, x_2) Y_W(u, x_1) \\
&= x_2^{-1} \delta \left( \frac{x_1 - x_0}{x_2} \right) p(x_1, x_2) Y_W(Y(u, x_0)v, x_2).
\end{aligned}$$

We also recall the definitions of a  $\Gamma$ -vertex algebra and an equivariant quasi module from [L3].

**Definition 2.15.** Let  $\Gamma$  be a group. A  $\Gamma$ -vertex algebra is a vertex algebra  $V$  equipped with group homomorphisms

$$R : \Gamma \rightarrow \text{GL}(V); \quad g \mapsto R_g, \quad \text{and } \chi : \Gamma \rightarrow \mathbb{C}^\times$$

such that  $R_g(\mathbf{1}) = \mathbf{1}$  for  $g \in \Gamma$  and

$$R_g Y(v, x) R_g^{-1} = Y(R_g(v), \chi(g)^{-1}x) \quad \text{for } g \in \Gamma, \quad v \in V.$$

Note that this is a reformulation of the notion of  $\Gamma$ -vertex algebra defined in [L2].

**Definition 2.16.** Let  $V$  be a  $\Gamma$ -vertex algebra. An *equivariant quasi  $V$ -module* is a quasi module  $(W, Y_W)$  for  $V$  viewed as a vertex algebra, satisfying the condition that

$$Y_W(R_g(v), x) = Y_W(v, \chi(g)x) \text{ for } g \in \Gamma, v \in V$$

and for  $u, v \in V$ , there exist  $\alpha_1, \dots, \alpha_k \in \chi(\Gamma) \subset \mathbb{C}^\times$  such that

$$(x_1 - \alpha_1 x_2) \cdots (x_1 - \alpha_k x_2) [Y_W(u, x_1), Y_W(v, x_2)] = 0.$$

**Remark 2.17.** Let  $V$  be a  $\mathbb{Z}$ -graded vertex algebra in the sense that  $V$  is a vertex algebra equipped with a  $\mathbb{Z}$ -grading  $V = \bigoplus_{n \in \mathbb{Z}} V_{(n)}$  such that  $\mathbf{1} \in V_{(0)}$  and

$$u_m V_{(n)} \subset V_{(k+n-m-1)} \quad \text{for } u \in V_{(k)}, k, m, n \in \mathbb{Z}.$$

Denote by  $L(0)$  the linear operator on  $V$ , defined by  $L(0)|_{V_{(n)}} = n$  for  $n \in \mathbb{Z}$ . Define an *automorphism* of a  $\mathbb{Z}$ -graded vertex algebra  $V$  to be an automorphism of vertex algebra  $V$ , which preserves the  $\mathbb{Z}$ -grading. Let  $\Gamma$  be an automorphism group of a  $\mathbb{Z}$ -graded vertex algebra  $V$  and let  $\chi : \Gamma \rightarrow \mathbb{C}^\times$  be a group homomorphism. Then  $V$  becomes a  $\Gamma$ -vertex algebra with  $R_g = \chi(g)^{-L(0)}g$  for  $g \in \Gamma$  (see [L2]).

Let  $\ell$  be a complex number, which is fixed for the rest of this section. Given the affine Lie algebra  $\widehat{\mathfrak{g}}_S$  and the complex number  $\ell$ , we have a  $\mathbb{Z}$ -graded vertex algebra  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$  (cf. [FZ], [LL]). As a vector space,

$$V_{\widehat{\mathfrak{g}}_S}(\ell, 0) = U(\widehat{\mathfrak{g}}_S) \otimes_{U(\mathfrak{g}_S \otimes \mathbb{C}[t] + \mathbb{C}\mathbf{k})} \mathbb{C}_\ell,$$

where  $\mathbb{C}_\ell$  denotes the one-dimensional  $(\mathfrak{g}_S \otimes \mathbb{C}[t] + \mathbb{C}\mathbf{k})$ -module  $\mathbb{C}$  with  $\mathfrak{g}_S \otimes \mathbb{C}[t]$  acting trivially and with  $\mathbf{k}$  acting as scalar  $\ell$ , and that  $\mathbf{1} = 1 \otimes 1$ ,  $Y(d^{\alpha, \beta}, x) = d^{\alpha, \beta}(x)$  for  $\alpha, \beta \in S$ . Furthermore,  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)_{(1)} = \mathfrak{g}_S$  is a generating subspace of  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$ , where  $\mathfrak{g}_S$  is identified as a subspace of  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$  through the linear map  $a \mapsto a(-1)\mathbf{1}$ .

Using a straightforward argument we have:

**Lemma 2.18.** *For each  $\gamma \in S$ , there exists an automorphism  $\sigma_\gamma$  of the  $\mathbb{Z}$ -graded vertex algebra  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$ , which is uniquely determined by  $\sigma_\gamma(d^{\alpha, \beta}) = d^{\alpha, \beta + \gamma}$  for  $\alpha, \beta \in S$ .*

Set

$$\Gamma = S \subset \text{Aut}(V_{\widehat{\mathfrak{g}}_S}(\ell, 0)), \quad (2.21)$$

where  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$  is viewed as a  $\mathbb{Z}$ -graded vertex algebra. Define a group homomorphism

$$R : \Gamma \rightarrow \text{GL}(V_{\widehat{\mathfrak{g}}_S}(\ell, 0))$$

by

$$R_\gamma = \chi(\gamma)^{-L(0)}\gamma \text{ for } \gamma \in S,$$

where  $L(0)$  is the linear operator on  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$  defined by  $L(0)v = nv$  for  $v \in V_{\widehat{\mathfrak{g}}_S}(\ell, 0)_{(n)}$  with  $n \in \mathbb{Z}$ . In view of Remark 2.17, equipped with group homomorphisms  $R$  and  $\chi$ , the  $\mathbb{Z}$ -graded vertex algebra  $V_{\widehat{\mathfrak{g}}_S}(\ell, 0)$  becomes a  $\Gamma$ -vertex algebra.

**Definition 2.19.** A  $D_S$ -module  $W$  is said to be *restricted* if for every  $\alpha \in S$  and  $w \in W$ ,  $D^\alpha(n)w = 0$  for  $n$  sufficiently large, or equivalently, if  $D^\alpha(x) \in \text{Hom}(W, W((x)))$ , or equivalently, if  $\tilde{D}^\alpha(x) \in \text{Hom}(W, W((x)))$  for all  $\alpha \in S$ . We say a  $D_S$ -module  $W$  is of *level*  $\ell \in \mathbb{C}$  if the central element  $\mathbf{c}$  acts as scalar  $\ell$ .

As the second main result of this section we have:

**Theorem 2.20.** *Let  $W$  be a restricted  $D_S$ -module of level  $\ell$ . Then there exists an equivariant quasi  $V_{\widehat{\mathfrak{g}_S}}(\ell, 0)$ -module structure  $Y_W(\cdot, x)$  on  $W$ , which is uniquely determined by*

$$Y_W(d^{\alpha, \beta}, x) = D^{\alpha, \beta}(x) = \chi(\beta) \tilde{D}^\alpha(\chi(\beta)x) \quad \text{for } \alpha, \beta \in S.$$

*On the other hand, let  $(W, Y_W)$  be an equivariant quasi  $V_{\widehat{\mathfrak{g}_S}}(\ell, 0)$ -module. Then  $W$  is a restricted  $D_S$ -module of level  $\ell$  with  $\tilde{D}^\alpha(x) = Y_W(d^{\alpha, 0}, x)$  for  $\alpha \in S$ .*

*Proof.* Recall from Theorem 2.13 that  $D_S$  is isomorphic to the covariant Lie algebra  $\widehat{\mathfrak{g}_S}/S$  of the affine Lie algebra  $\widehat{\mathfrak{g}_S}$ . Note that  $\chi : S \rightarrow \mathbb{C}^\times$  is assumed to be injective. Then all the assertions follow immediately from Theorem 4.9 of [L3].  $\square$

### 3 Identifying Lie algebra $D_S$ with affine Kac-Moody algebras

In this section, we relate Lie algebra  $D_S$  to affine Kac-Moody algebras. To do this, we determine the Lie algebra  $\mathfrak{g}_S$  introduced in Section 2. Especially, we show that if  $S$  is a finite abelian group of order  $2l + 1$ , then  $\mathfrak{g}_S$  is a finite-dimensional simple Lie algebra of type  $B_l$  and Lie algebra  $D_S$  is isomorphic to the affine Kac-Moody algebra of type  $B_l^{(1)}$ . We also obtain some results for  $S$  of an even order.

Let  $S$  be an abelian group as before, which is fixed throughout this section. Define  $\mathfrak{gl}_S$  to be the associative algebra with generators  $E_{\alpha, \beta}$  for  $\alpha, \beta \in S$ , subject to relations

$$E_{\alpha, \beta} \cdot E_{\mu, \nu} = \delta_{\beta, \mu} E_{\alpha, \nu} \quad \text{for } \alpha, \beta, \mu, \nu \in S. \quad (3.1)$$

It is straightforward to see that  $E_{\alpha, \beta}$  for  $\alpha, \beta \in S$  form a basis of  $\mathfrak{gl}_S$ . Notice that if  $S$  is of order  $n$ , then  $\mathfrak{gl}_S$  is isomorphic to the algebra  $M(n, \mathbb{C})$  of  $n \times n$  complex matrices. Naturally,  $\mathfrak{gl}_S$  is a Lie algebra with

$$[E_{\alpha, \beta}, E_{\mu, \nu}] = E_{\alpha, \beta} \cdot E_{\mu, \nu} - E_{\mu, \nu} \cdot E_{\alpha, \beta} = \delta_{\beta, \mu} E_{\alpha, \nu} - \delta_{\nu, \alpha} E_{\mu, \beta}.$$

Equip  $\mathfrak{gl}_S$  with the bilinear form  $\langle \cdot, \cdot \rangle$  defined by

$$\langle E_{\alpha, \beta}, E_{\mu, \nu} \rangle = \frac{1}{2} \delta_{\alpha, \nu} \delta_{\beta, \mu} \quad (3.2)$$

for  $\alpha, \beta, \mu, \nu \in S$ . It is straightforward to show that this form is symmetric, associative (invariant), and nondegenerate. Define a linear endomorphism  $\tau$  of  $\mathfrak{gl}_S$  by

$$\tau(E_{\alpha,\beta}) = -E_{\beta,\alpha} \quad \text{for } \alpha, \beta \in S. \quad (3.3)$$

Then  $\tau$  is an order-2 automorphism of  $\mathfrak{gl}_S$  viewed as a Lie algebra and we see that  $\tau$  preserves the bilinear form.

For  $\alpha, \beta \in S$ , set

$$G_{\alpha,\beta} = E_{\alpha+\beta,\beta-\alpha}. \quad (3.4)$$

We have

$$G_{\alpha,\beta} \cdot G_{\mu,\nu} = \delta_{\beta-\alpha,\mu+\nu} G_{\alpha+\mu,\alpha+\nu} \quad \text{for } \alpha, \beta, \mu, \nu \in S.$$

Furthermore, set

$$\mathcal{A}_S = \text{span}\{G_{\alpha,\beta} \mid \alpha, \beta \in S\} \subset \mathfrak{gl}_S. \quad (3.5)$$

By a straightforward argument we get the following results:

**Lemma 3.1.** *The subspace  $\mathcal{A}_S$  of  $\mathfrak{gl}_S$  is an associative subalgebra with a basis*

$$\{E_{\mu,\nu} \mid \mu, \nu \in S \text{ with } \mu + \nu \in 2S\}. \quad (3.6)$$

*Furthermore, the automorphism  $\tau$  of the Lie algebra  $\mathfrak{gl}_S$  preserves  $\mathcal{A}_S$ , where*

$$\tau(G_{\alpha,\beta}) = -G_{-\alpha,\beta} \quad \text{for } \alpha, \beta \in S. \quad (3.7)$$

Recall that  $S^0 = \{\alpha \in S \mid 2\alpha = 0\}$ . We have:

**Lemma 3.2.** *For  $\alpha, \beta, \alpha', \beta' \in S$ ,  $G_{\alpha,\beta} = G_{\alpha',\beta'}$  if and only if  $\alpha = \alpha' + \gamma$ ,  $\beta = \beta' - \gamma = \beta' + \gamma$  for some  $\gamma \in S^0$ .*

*Proof.* It is straightforward as  $E_{\mu,\nu}$  for  $\mu, \nu \in S$  form a basis of  $\mathfrak{gl}_S$ . □

Set

$$\mathcal{A}_S^\tau = \{a \in \mathcal{A}_S \mid \tau(a) = a\}, \quad (3.8)$$

a Lie subalgebra of  $\mathcal{A}_S$ . For  $\alpha, \beta \in S$ , set

$$G_{\alpha,\beta}^\tau = G_{\alpha,\beta} + \tau(G_{\alpha,\beta}) = G_{\alpha,\beta} - G_{-\alpha,\beta} = E_{\alpha+\beta,\beta-\alpha} - E_{\beta-\alpha,\beta+\alpha} \in \mathcal{A}_S^\tau. \quad (3.9)$$

Then

$$\mathcal{A}_S^\tau = \text{span}\{G_{\alpha,\beta}^\tau \mid \alpha, \beta \in S\}. \quad (3.10)$$

We have

$$G_{\alpha+\gamma, \beta+\gamma}^\tau = G_{\alpha, \beta}^\tau \quad \text{for } \alpha, \beta \in S, \gamma \in S^0, \quad (3.11)$$

$$G_{-\alpha, \beta}^\tau = -G_{\alpha, \beta}^\tau \quad \text{for } \alpha, \beta \in S. \quad (3.12)$$

On the other hand, by a straightforward calculation we get

$$\begin{aligned} [G_{\alpha, \beta}^\tau, G_{\mu, \nu}^\tau] &= \delta_{\alpha+\mu, \beta-\nu} G_{\alpha+\mu, \nu+\alpha}^\tau - \delta_{\alpha+\mu, \nu-\beta} G_{\alpha+\mu, \mu+\beta}^\tau \\ &\quad + \delta_{\alpha-\mu, \nu-\beta} G_{\alpha-\mu, \beta-\mu}^\tau - \delta_{\alpha-\mu, \beta-\nu} G_{\alpha-\mu, \alpha+\nu}^\tau \end{aligned} \quad (3.13)$$

for  $\alpha, \beta, \mu, \nu \in S$ .

**Lemma 3.3.** *The group  $S$  acts on the Lie algebra  $\mathcal{A}_S^\tau$  as an automorphism group with  $\gamma \in S$  acting as  $\sigma_\gamma$ , where*

$$\sigma_\gamma(G_{\alpha, \beta}^\tau) = G_{\alpha, \beta+\gamma}^\tau \quad (3.14)$$

for  $\alpha, \beta \in S$ . Furthermore, the action preserves the bilinear form  $\langle \cdot, \cdot \rangle$  defined in (3.2).

*Proof.* First,  $S$  acts on  $\mathfrak{gl}_S$  as an automorphism group by

$$\sigma_\gamma(E_{\alpha, \beta}) = E_{\alpha+\gamma, \beta+\gamma} \quad \text{for } \gamma, \alpha, \beta \in S. \quad (3.15)$$

It is clear that  $S$  preserves the bilinear form  $\langle \cdot, \cdot \rangle$  on  $\mathfrak{gl}_S$ . Second, for  $\alpha, \beta, \gamma \in S$ , we have

$$\sigma_\gamma(G_{\alpha, \beta}) = E_{\alpha+\beta+\gamma, \beta-\alpha+\gamma} = G_{\alpha, \beta+\gamma}$$

and furthermore, we have

$$\sigma_\gamma(G_{\alpha, \beta}^\tau) = G_{\alpha, \beta+\gamma} - G_{-\alpha, \beta+\gamma} = G_{\alpha, \beta+\gamma}^\tau.$$

Consequently,  $S$  acts on  $\mathcal{A}_S^\tau$  as an automorphism group.  $\square$

Next, we relate the Lie algebra  $\mathfrak{g}_S$  introduced in Section 2 with Lie subalgebra  $\mathcal{A}_S^\tau$  of  $(\mathfrak{gl}_S)^\tau$ . Recall that  $S$  also acts on  $\mathfrak{g}_S$  as an automorphism group. From relations (3.12) and (3.13) and the definition of  $\mathfrak{g}_S$  we immediately have:

**Lemma 3.4.** *There exists a Lie algebra homomorphism  $\pi$  from  $\mathfrak{g}_S$  onto  $\mathcal{A}_S^\tau \subset (\mathfrak{gl}_S)^\tau$  such that*

$$\pi(d^{\alpha, \beta}) = G_{-\alpha, \beta}^\tau = -G_{\alpha, \beta}^\tau = E_{\beta-\alpha, \alpha+\beta} - E_{\alpha+\beta, \beta-\alpha} \quad \text{for } \alpha, \beta \in S. \quad (3.16)$$

Furthermore,  $\pi$  preserves the group actions of  $S$ .

Furthermore, we have:

**Proposition 3.5.** *Assume that  $S$  is an abelian group such that  $S^0 = \{0\}$ . Then the Lie algebra homomorphism  $\pi$  defined in Lemma 3.4 from  $\mathfrak{g}_S$  to  $\mathcal{A}_S^\tau$  is an isomorphism, which also preserves the invariant bilinear form.*

*Proof.* For  $\alpha, \beta, \mu, \nu \in S$ , we have

$$\begin{aligned}
\langle G_{\alpha,\beta}^\tau, G_{\mu,\nu}^\tau \rangle &= \langle E_{\alpha+\beta,\beta-\alpha} - E_{\beta-\alpha,\alpha+\beta}, E_{\mu+\nu,\nu-\mu} - E_{\nu-\mu,\mu+\nu} \rangle \\
&= \delta_{\alpha+\beta,\nu-\mu} \delta_{\beta-\alpha,\mu+\nu} - \delta_{\alpha+\beta,\mu+\nu} \delta_{\beta-\alpha,\nu-\mu} \\
&= \delta_{2(\alpha+\mu),0} \delta_{\beta-\alpha,\mu+\nu} - \delta_{2(\alpha-\mu),0} \delta_{\beta-\alpha,\nu-\mu} \\
&= \delta_{\alpha+\mu,0} \delta_{\beta-\alpha,\mu+\nu} - \delta_{\alpha-\mu,0} \delta_{\beta-\alpha,\nu-\mu} \\
&= (\delta_{\alpha,-\mu} - \delta_{\alpha,\mu}) \delta_{\beta,\nu}.
\end{aligned} \tag{3.17}$$

It follows from this and (2.19) that  $\pi$  preserves the invariant bilinear form. For the isomorphism assertion it remains to show that  $\pi$  is injective. As  $S^0 = \{0\}$ , in view of Lemma 3.2, we have that  $G_{\alpha,\beta} = G_{\alpha',\beta'}$  if and only if  $\alpha = \alpha'$  and  $\beta = \beta'$  for  $\alpha, \beta, \alpha', \beta' \in S$ . Consequently,  $G_{\alpha,\beta}$  for  $\alpha, \beta \in S$  form a basis of  $\mathcal{A}_S$ . Let  $S_-$  be a subset of  $S$  such that  $S = \{0\} \cup S_- \cup (-S_-)$  is a disjoint decomposition. As  $G_{-\alpha,\beta}^\tau = -G_{\alpha,\beta}^\tau$  for  $\alpha, \beta \in S$ ,  $\mathcal{A}_S^\tau$  is linearly spanned by  $G_{\alpha,\beta}^\tau$  for  $\alpha \in S_-$ ,  $\beta \in S$ , which are linearly independent since  $G_{\alpha,\beta}$  for  $\alpha, \beta \in S$  are linearly independent. Then  $G_{\alpha,\beta}^\tau$  for  $\alpha \in S_-$ ,  $\beta \in S$  form a basis of  $\mathcal{A}_S^\tau$ . On the other hand, with (2.13), we see that  $d^{\alpha,\beta}$  with  $\alpha \in S_-$ ,  $\beta \in S$  linearly span  $\mathfrak{g}_S$ . As  $\pi$  is a linear map from  $\mathfrak{g}_S$  onto  $\mathcal{A}_S^\tau$  with  $\pi(d^{\alpha,\beta}) = -G_{\alpha,\beta}^\tau$  for  $\alpha, \beta \in S$ , elements  $d^{\alpha,\beta}$  for  $\alpha \in S_-$ ,  $\beta \in S$  must be linearly independent, so that  $\{d^{\alpha,\beta} \mid \alpha \in S_-, \beta \in S\}$  is a basis of  $\mathfrak{g}_S$ . Consequently,  $\pi$  is injective. This completes the proof.  $\square$

As a consequence of Proposition 3.5, we have:

**Corollary 3.6.** *Assume that  $S$  is a finite abelian group of order  $2l + 1$  with  $l$  a positive integer. Then  $\mathcal{A}_S = \mathfrak{gl}_S$  and  $\mathfrak{g}_S$  is isomorphic to  $(\mathfrak{gl}_S)^\tau$  which is isomorphic to the simple Lie algebra of type  $B_l$ .*

*Proof.* As  $|S|$  is odd, there exist integers  $m$  and  $n$  such that  $1 = 2m + n|S|$ . Then  $\alpha = 2m\alpha$  for  $\alpha \in S$ . It follows from (3.4) that for any  $\alpha, \beta \in S$ ,

$$E_{\alpha,\beta} = G_{m(\alpha-\beta), m(\alpha+\beta)} \in \mathcal{A}_S. \tag{3.18}$$

Thus  $\mathcal{A}_S = \mathfrak{gl}_S$  and hence  $\mathcal{A}_S^\tau = (\mathfrak{gl}_S)^\tau$ . Again, since  $|S|$  is odd, we have  $S^0 = \{0\}$ . By Proposition 3.5,  $\mathfrak{g}_S \simeq \mathcal{A}_S^\tau = (\mathfrak{gl}_S)^\tau$ . It is well known that  $(\mathfrak{gl}_S)^\tau$  is isomorphic to the finite-dimensional simple Lie algebra of type  $B_l$ . This completes the proof.  $\square$

Furthermore, we have:

**Theorem 3.7.** *Assume that  $S$  is a finite abelian group of order  $2l + 1$  with  $l$  a positive integer. Then Lie algebra  $D_S$  is isomorphic to the affine Kac-Moody algebra of type  $B_l^{(1)}$ .*

*Proof.* Note that as  $\chi : S \rightarrow \mathbb{C}^\times$  is assumed to be one-to-one,  $S$  must be cyclic. Assume  $S = \mathbb{Z}/(2l+1)\mathbb{Z}$  and let  $\chi : S \rightarrow \mathbb{C}^\times$  be the group homomorphism defined by  $\chi(\alpha) = q^\alpha$  for  $\alpha \in S$ , where  $q$  is a primitive  $(2l+1)$ -th root of unity. From Theorem 2.13 and Corollary 3.6, we have

$$D_S \simeq \widehat{\mathfrak{g}_S}/S, \quad \mathfrak{g}_S \simeq (\mathfrak{gl}_S)^\tau \simeq B_l. \quad (3.19)$$

Let  $\sigma_1$  be the automorphism of the Lie algebra  $(\mathfrak{gl}_S)^\tau$  ( $\simeq B_l$ ), given by

$$\sigma_1(E_{\alpha,\beta} - E_{\beta,\alpha}) = E_{\alpha+[1],\beta+[1]} - E_{\beta+[1],\alpha+[1]} \quad (3.20)$$

for  $\alpha, \beta \in \mathbb{Z}/(2l+1)\mathbb{Z} = S$ , where  $[1]$  denotes the image of 1 in  $\mathbb{Z}/(2l+1)\mathbb{Z}$ . With  $S$  viewed as an automorphism group of  $(\mathfrak{gl}_S)^\tau$  (recalling (3.15)), we have  $S = \langle \sigma_1 \rangle$ . Then, using (3.19) we get

$$D_S \simeq \widehat{B_l}/S \simeq \widehat{B_l}^S = \widehat{B_l}^{\sigma_1},$$

where  $\widehat{B_l}^S$  (resp.  $\widehat{B_l}^{\sigma_1}$ ) denotes the  $S$ -fixed (resp.  $\sigma_1$ -fixed) points subalgebra. Note that for the isomorphism relation  $\widehat{B_l}/S \simeq \widehat{B_l}^S$ ,  $\mathbf{k}$  corresponds to  $(2l+1)\mathbf{k}$ .

Note that there is no nontrivial Dynkin diagram automorphism for a type  $B_l$  simple Lie algebra. From [K] (Proposition 8.1),  $\sigma_1$  is conjugate to an inner automorphism  $\exp\left(\frac{2\pi i}{2l+1}\text{adh}\right)$ , where  $h$  is an element such that  $\text{adh}$  is semi-simple with only integer eigenvalues. It follows that  $\widehat{B_l}^{\sigma_1}$  is isomorphic to  $\widehat{B_l}$  (see Appendix; cf. Section 8.5, [K]). Consequently,  $D_S$  is isomorphic to the affine Kac-Moody algebra of type  $B_l^{(1)}$ .  $\square$

Next, we consider the general case. Recall  $\mathcal{A}_S = \text{span}\{E_{\alpha+\beta,\alpha-\beta} \mid \alpha, \beta \in S\}$ , which is an associative subalgebra of  $\mathfrak{gl}_S$ , and

$$\mathcal{A}_S^\tau = \text{span}\{G_{\alpha,\beta}^\tau = E_{\beta+\alpha,\beta-\alpha} - E_{\beta-\alpha,\beta+\alpha} \mid \alpha, \beta \in S\},$$

a Lie subalgebra of  $\mathcal{A}_S$ . We also recall  $S^0 = \{\alpha \in S \mid 2\alpha = 0\}$ . The following is a generalization of Proposition 3.5:

**Proposition 3.8.** *Set*

$$I = \text{span}\{d^{\alpha+\gamma,\beta+\gamma} - d^{\alpha,\beta} \mid \alpha, \beta \in S, \gamma \in S^0\} \subset \mathfrak{g}_S. \quad (3.21)$$

*Then  $I$  is an ideal of  $\mathfrak{g}_S$ . Furthermore, the quotient Lie algebra  $\mathfrak{g}_S/I$  is isomorphic to  $\mathcal{A}_S^\tau$ .*

*Proof.* From (2.14), it is straightforward to show that  $I$  is an ideal. On the other hand, by Lemma 3.2, for  $\alpha, \beta \in S$ ,  $\gamma \in S^0$ , we have

$$G_{\alpha+\gamma,\beta+\gamma}^\tau = G_{\alpha+\gamma,\beta+\gamma} - G_{-\alpha-\gamma,\beta+\gamma} = G_{\alpha,\beta} - G_{-\alpha,\beta} = G_{\alpha,\beta}^\tau$$



in  $\mathcal{A}_S^\tau$ . Then the Lie algebra homomorphism  $\pi$  from  $\mathfrak{g}_S$  onto  $\mathcal{A}_S^\tau$  reduces to a Lie algebra homomorphism  $\bar{\pi}$  from  $\mathfrak{g}_S/I$  onto  $\mathcal{A}_S^\tau$  with  $\bar{\pi}(d^{\alpha,\beta} + I) = -G_{\alpha,\beta}^\tau$  for  $\alpha, \beta \in S$ . It suffices to prove that  $\bar{\pi}$  is injective. Let  $T$  be a complete set of equivalence class representatives of  $S$  modulo  $S^0$  with  $0 \in T$ . It follows from Lemma 3.2 that  $\{G_{\alpha,\beta}^\tau \mid (\alpha, \beta) \in T \times S\}$  is a base for  $\mathcal{A}_S$ . Furthermore, consider the equivalence relation  $\sim$  on  $S$  defined by  $\alpha \sim \beta$  if and only if  $\beta = \pm\alpha$ . Let  $T_-$  be a complete set of equivalence class representatives in  $T \setminus \{0\}$  with respect to this particular equivalence relation. Noticing that if  $\alpha \in T_-$ , then  $-\alpha \notin T_-$ , we see that the linear independence of  $\{G_{\alpha,\beta}^\tau \mid (\alpha, \beta) \in T \times S\}$  implies that  $G_{\alpha,\beta}^\tau$  with  $\alpha \in T_-$ ,  $\beta \in S$  are linearly independent. Then  $\{G_{\alpha,\beta}^\tau \mid (\alpha, \beta) \in T_- \times S\}$  is a base for  $\mathcal{A}_S^\tau$ . On the other hand, from the defining relations of  $\mathfrak{g}_S$  and the definition of  $I$ , we see that  $\mathfrak{g}_S/I$  is linearly spanned by  $d^{\alpha,\beta} + I$  with  $(\alpha, \beta) \in T_- \times S$ . It follows that  $\{d^{\alpha,\beta} + I \mid (\alpha, \beta) \in T_- \times S\}$  is a base for  $\mathfrak{g}_S/I$ . Consequently,  $\bar{\pi}$  is a Lie algebra isomorphism.  $\square$

Recall the subgroup  $2S$  of  $S$ . Let

$$S = S_0 \cup S_1 \cup \cdots \cup S_{r-1} \quad (3.22)$$

be the decomposition of  $S$  into distinct cosets of  $2S$  with  $S_0 = 2S$ . We have  $2S \simeq S/S^0$ , which implies  $|2S| \cdot |S^0| = |S|$ . Thus  $r = |S^0|$ . Notice that for  $\alpha, \beta \in S$ , if  $\alpha + \beta \in S_j$  for  $0 \leq j < r$ , we have  $\beta - \alpha = (\alpha + \beta) - 2\alpha \in S_j$ , so that

$$G_{\alpha,\beta}^\tau = E_{\beta+\alpha,\beta-\alpha} - E_{\beta-\alpha,\beta+\alpha} \in \mathfrak{gl}_{S_j}.$$

Thus

$$\mathcal{A}_S^\tau \subset \mathfrak{gl}_{S_0} \oplus \mathfrak{gl}_{S_1} \oplus \cdots \oplus \mathfrak{gl}_{S_{r-1}} \subset \mathfrak{gl}_S. \quad (3.23)$$

For  $0 \leq j \leq r-1$ , set

$$\mathfrak{g}_j = \mathcal{A}_S^\tau \cap \mathfrak{gl}_{S_j},$$

a Lie subalgebra of  $\mathcal{A}_S^\tau$ . Then

$$\mathcal{A}_S^\tau = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_{r-1}, \quad (3.24)$$

a direct sum of Lie algebras.

Furthermore, we have:

**Lemma 3.9.** *For each  $0 \leq j < r$ , we have*

$$\mathfrak{g}_j = \text{span}\{E_{\mu,\nu} - E_{\nu,\mu} \mid \mu, \nu \in S_j\} \quad (3.25)$$

and  $\dim \mathfrak{g}_j = \frac{1}{2}k(k-1)$  where  $k = |2S|$ .

*Proof.* From definition we have

$$\mathfrak{g}_j = \text{span}\{E_{\beta+\alpha, \beta-\alpha} - E_{\beta-\alpha, \beta+\alpha} \mid \alpha, \beta \in S \text{ with } \alpha + \beta \in S_j\},$$

noticing that  $\alpha + \beta \in S_j$  implies  $\beta - \alpha = (\alpha + \beta) - 2\alpha \in S_j$ . It follows that  $\mathfrak{g}_j \subset \text{span}\{E_{\mu, \nu} - E_{\nu, \mu} \mid \mu, \nu \in S_j\}$ . On the other hand, let  $\mu, \nu \in S_j$ . Fix an element  $\alpha_j \in S_j$ . Then  $\mu = \alpha_j + 2\alpha$  and  $\nu = \alpha_j + 2\beta$  for some  $\alpha, \beta \in S$ . Thus

$$E_{\mu, \nu} - E_{\nu, \mu} = G_{\alpha-\beta, \alpha+\beta+\alpha_j}^\tau \in \mathfrak{g}_j.$$

This proves that  $\mathfrak{g}_j \supset \text{span}\{E_{\mu, \nu} - E_{\nu, \mu} \mid \mu, \nu \in S_j\}$ . Hence (3.25) holds. From this the second assertion of the lemma follows immediately.  $\square$

Note that  $\mathfrak{gl}_S$  naturally acts on the group algebra  $\mathbb{C}[S]$  of  $S$ . Equip  $\mathbb{C}[S]$  with a nondegenerate symmetric bilinear form  $\langle \cdot, \cdot \rangle$  defined by  $\langle e^\alpha, e^\beta \rangle = \delta_{\alpha, \beta}$  for  $\alpha, \beta \in S$ . As subalgebras of  $\mathfrak{gl}_S$ ,  $\mathcal{A}_S$  and  $\mathcal{A}_S^\tau$  also act on  $\mathbb{C}[S]$ . We have the following  $\mathcal{A}_S^\tau$ -module decomposition

$$\mathbb{C}[S] = \mathbb{C}[S_0] \oplus \mathbb{C}[S_1] \oplus \cdots \oplus \mathbb{C}[S_{r-1}],$$

where  $\mathfrak{g}_j$  acts on  $\mathbb{C}[S_j]$  for  $0 \leq j < r$ . Denoting by  $\rho$  the algebra homomorphism of  $\mathfrak{gl}_S$  to  $\mathfrak{gl}(\mathbb{C}[S])$ , we have

$$\rho(\mathfrak{g}_j) \subset \{A \in \mathfrak{gl}(\mathbb{C}[S_j]) \mid (Au, v) = -(u, Av) \text{ for } u, v \in \mathbb{C}[S_j]\}.$$

Since  $|S_j| = k$  and  $\dim \mathfrak{g}_j = \frac{1}{2}k(k-1)$  by Lemma 3.9, we see that actually the equality holds. Therefore,  $\mathfrak{g}_j$  is a simple Lie algebra of type either  $B$  or  $D$ , depending on that  $k$  ( $= |2S|$ ) is odd or even. Consequently,  $\mathcal{A}_S^\tau$  is semi-simple.

To summarize, we have:

**Proposition 3.10.** *Let  $S$  be a finite abelian group. Set  $k = |2S|$  and  $r = |S/2S|$ . Then  $\mathfrak{g}_S/I$  is isomorphic to the direct sum of  $r$  copies of simple Lie algebras of type  $B_{\frac{1}{2}(k-1)}$  if  $k$  is odd and of type  $D_{\frac{1}{2}k}$  if  $k$  is even, where  $I$  is the ideal of  $\mathfrak{g}_S$ , which was obtained in Proposition 3.8.*

## Appendix

Here, we provide a proof of a statement at the end of the proof of Theorem 3.7.

**Lemma 3.11.** *Suppose that  $\mathfrak{g}$  is a Lie algebra with a symmetric invariant bilinear form  $\langle \cdot, \cdot \rangle$  and  $h$  is an element of  $\mathfrak{g}$  such that  $\text{ad} h$  is semi-simple with only integer eigenvalues. Set  $\sigma = \exp(\frac{2\pi i}{T} \text{ad} h)$ , where  $T$  is a fixed positive integer. Then  $\hat{\mathfrak{g}}^\sigma \simeq \hat{\mathfrak{g}}$  with  $\mathbf{k}$  corresponding to  $\frac{1}{T}\mathbf{k}$ .*

*Proof.* Recall the affine Lie algebra  $\hat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}\mathbf{k}$ , where

$$[a \otimes t^m, b \otimes t^n] = [a, b] \otimes t^{m+n} + m\delta_{m+n, 0} \langle a, b \rangle \mathbf{k}$$

for  $a, b \in \mathfrak{g}$ ,  $m, n \in \mathbb{Z}$ . On the other hand, we have Lie algebra

$$\widehat{L}(\mathfrak{g}, T) = \mathfrak{g} \otimes \mathbb{C}[t^{\frac{1}{T}}, t^{-\frac{1}{T}}] \oplus \mathbb{C}\mathbf{k},$$

where

$$[a \otimes t^{\frac{m}{T}}, b \otimes t^{\frac{n}{T}}] = [a, b] \otimes t^{\frac{m+n}{T}} + \frac{m}{T} \delta_{m+n,0} \langle a, b \rangle \mathbf{k}$$

for  $a, b \in \mathfrak{g}$ ,  $m, n \in \mathbb{Z}$ . It is straightforward to see that  $\hat{\mathfrak{g}}$  is canonically isomorphic to  $\widehat{L}(\mathfrak{g}, T)$  with  $\mathbf{k}$  corresponding to  $\frac{1}{T}\mathbf{k}$ . View  $\sigma$  as an automorphism of  $\widehat{L}(\mathfrak{g}, T)$  by

$$\sigma(a \otimes t^{\frac{n}{T}}) = \exp\left(\frac{2n\pi i}{T}\right) \sigma(a) \otimes t^{\frac{n}{T}} \quad \text{for } a \in \mathfrak{g}, n \in \mathbb{Z},$$

and then set

$$\hat{\mathfrak{g}}[\sigma] = \widehat{L}(\mathfrak{g}, T)^\sigma.$$

Then  $\hat{\mathfrak{g}}^\sigma$  is canonically isomorphic to  $\hat{\mathfrak{g}}[\sigma]$  with  $\mathbf{k}$  corresponding to  $\frac{1}{T}\mathbf{k}$ . Now, it suffices to prove that  $\hat{\mathfrak{g}}[\sigma]$  is isomorphic to  $\hat{\mathfrak{g}}$  with  $\mathbf{k}$  corresponding to  $\mathbf{k}$ . For  $r \in \mathbb{Z}$ , set  $\mathfrak{g}_r = \{u \in \mathfrak{g} \mid [h, u] = ru\}$ . Then  $\mathfrak{g} = \bigoplus_{r \in \mathbb{Z}} \mathfrak{g}_r$  and  $\langle \mathfrak{g}_r, \mathfrak{g}_s \rangle = 0$  whenever  $r + s \neq 0$ . For  $a \in \mathfrak{g}_r$  with  $r \in \mathbb{Z}$ , set

$$a^\sigma(x) = \sum_{n \in \mathbb{Z}} (a \otimes t^{n-\frac{r}{T}}) x^{-n-1+\frac{r}{T}} \in x^{\frac{r}{T}} \hat{\mathfrak{g}}[\sigma][[x, x^{-1}]].$$

It was essentially proved in [L1] (Remark 5.5) that the linear map  $\psi : \hat{\mathfrak{g}}[\sigma] \rightarrow \hat{\mathfrak{g}}$  defined by  $\psi(\mathbf{k}) = \mathbf{k}$  and

$$\psi(a^\sigma(x)) = x^{\frac{r}{T}} \left( a(x) + \frac{1}{T} \langle h, a \rangle \mathbf{k} x^{-1} \right)$$

for  $a \in \mathfrak{g}_r$  with  $r \in \mathbb{Z}$ , is a Lie algebra isomorphism. In terms of components, we have

$$\psi(a \otimes t^{n-\frac{r}{T}}) = a \otimes t^n + \frac{1}{T} \langle h, a \rangle \delta_{n,0} \mathbf{k},$$

noticing that  $\langle h, a \rangle = \delta_{r,0} \langle h, a \rangle$  as  $h \in \mathfrak{g}_0$ . On the other hand, it is straightforward to show that  $\psi$  is indeed a Lie algebra isomorphism.  $\square$

## References

- [BC] A. Belov, K. Chaltikian,  $q$ -deformation of Virasoro algebra and lattice conformal theories, *Modern Phys. Lett. A* **8**(1993), 1233-1242.
- [FZ] I. B. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* **66** (1992), 123-168.
- [G-K-K] M. Golenishcheva-kutuzova and V. Kac,  $\Gamma$ -conformal algebras, *J. Math. Phys.* **39** (1998), 2290-2305.

- [GLTW] H. Guo, H.-S. Li, S. Tan and Q. Wang,  $q$ -Virasoro algebra and vertex algebras, *Journal of Pure and Applied Algebra* **219** (2015) 1258-1277.
- [K] V. G. Kac, *Infinite-dimensional Lie Algebras*, 3rd ed., Cambridge University Press, Cambridge, 1990.
- [LL] J. Lepowsky and H.-S. Li, *Introduction to Vertex Operator Algebras and Their Representations*, Progress in Math., Vol. 227, Birkhäuser, Boston, 2004.
- [L1] H.-S. Li, Local systems of twisted vertex operators, vertex operator superalgebras and twisted modules, *Contemporary Math.* **193** (1996), 203-236.
- [L2] H.-S. Li, A new construction of vertex algebras and quasi modules for vertex algebras, *Advances in Math.* **202** (2006), 232-286.
- [L3] H.-S. Li, On certain generalizations of twisted affine Lie algebras and quasi-modules for  $\Gamma$ -vertex algebras, *J. Pure and Appl. Algebra* **209** (2007), 853-871.
- [N] A. Nigro, A  $q$ -Virasoro algebra at roots of unity, Free Fermions and Temperley Lieb Hamiltonians, *J. Math. Phys.* **57** (2016), no.4, 041702.