

# SIMPLE CONFORMALLY RECURRENT SPACE-TIMES ARE CONFORMALLY RECURRENT PP-WAVES

CARLO ALBERTO MANTICA AND LUCA GUIDO MOLINARI

ABSTRACT. We show that in dimension  $n \geq 4$  the class of simple conformally recurrent space-times coincides with the class of conformally recurrent pp-waves.

## 1. INTRODUCTION

Riemannian manifolds with recurrent Weyl tensor (conformally recurrent manifolds) were introduced by Adati and Miyazawa [1] and subsequently extended to pseudo-Riemannian manifolds by Derdziński [7], Roter [31, 32, 33, 34], Suh and Kwon [38], and to Lorentzian manifolds (space-times) by Mc Lenaghan et al. [24, 25], Hall [15, 16], De and Mantica [5]. In a recent study on conformally recurrent pseudo-Riemannian and Lorentzian manifolds in dimension  $n \geq 5$ , the authors showed that the Ricci tensor has at most two different eigenvalues, with restrictions on the metric structure, and gave an explicit representation of the Weyl tensor as a Kulkarni-Nomizu product [21, 22].

A pseudo-Riemannian manifold of dimension  $n \geq 4$  is “conformally recurrent” if the Weyl curvature tensor<sup>1</sup> is everywhere non-zero and satisfies the condition

$$(1.1) \quad \nabla_i C_{jklm} = \alpha_i C_{jklm}$$

where  $\alpha_i$  is a non-zero 1-form named recurrence vector.

The natural question arises whether the recurrent vector may be locally cancelled by a conformal transformation. This “simple” case was investigated by Roter [31, 34].

---

*Date:* Nov 2016.

*2010 Mathematics Subject Classification.* Primary 53B30; Secondary 53C80.

*Key words and phrases.* conformally recurrent pseudo-Riemannian manifold, simple conformally recurrent pseudo Riemannian manifold, pp-wave space-time.

<sup>1</sup>The local components of the Weyl tensor are [29]

$$C_{jkl}{}^m = R_{jkl}{}^m + \frac{1}{n-2}(\delta_j^m R_{kl} - \delta_k^m R_{jl} + g_{kl}R_j^m - g_{jl}R_k^m) - R \frac{\delta_j^m g_{kl} - \delta_k^m g_{jl}}{(n-1)(n-2)}$$

with Ricci tensor  $R_{ij} = -R_{kij}{}^k$  and curvature scalar  $R = R^k{}_k$ .

**Definition 1.1** (Roter). A conformally recurrent manifold  $(\mathcal{M}, g)$  is “simple” if the metric is locally conformal to a conformally symmetric metric, i.e. at each point  $x \in \mathcal{M}$  there exist a neighbourhood  $U_x$  and a scalar function  $\sigma$  on  $U_x$  such that  $g'_{ij} = g_{ij}e^{2\sigma}$  and  $\nabla'_i C'_{jklm} = 0$  on  $U_x$ .

The main result of this work is the statement that such spaces, with a Lorentzian metric, coincide with (conformally recurrent) pp-wave Lorentzian space-times.

pp-waves are important in general relativity (see for example [19] and references therein). They were studied in  $n > 4$  by Schimming [36], and appear in Kaluza-Klein theories and in string theory. In the literature (see for example [27]), pp-wave space-times are identified with Brinkmann-waves, i.e. space-times equipped with a null covariantly constant vector field,  $X^i X_i = 0$ ,  $\nabla_j X_k = 0$  [4]. Here we conform to the more restrictive definition used in [36, 19, 17, 18]:

**Definition 1.2.** A pp-wave is a Brinkmann-wave whose Riemann tensor satisfies the condition  $R_{jk}{}^{pq}R_{pqlm} = 0$ .

The layout of the paper is the following. In section 2 we present preliminary material about simple conformally recurrent pseudo-Riemannian manifolds and pp-wave space-times. In section 3 we prove that in dimension  $n \geq 4$  a conformally recurrent pp-wave space-time is a simple conformally recurrent Lorentzian manifold, and establish the converse implication. We obtain new curvature properties of conformally recurrent pp-waves, such as  $\nabla^m \nabla_m R_{ij} = 0$ . In Section 4 we show that conformally recurrent pp-wave space-times are solutions to the extended theory of gravity with pure radiation source.

Throughout the paper, the manifolds are assumed to be connected, Hausdorff, with non-degenerate metric of arbitrary signature, i.e.  $n$ -dimensional pseudo-Riemannian manifolds. Some results are restricted to Lorentzian manifolds (space-times) i.e. to metrics of signature  $n - 2$ .

## 2. PRELIMINARY RESULTS

In this section we present preliminary material about conformally recurrent pseudo-Riemannian manifolds, simple conformally recurrent manifolds, and pp-wave space-times.

A vector field belongs to the Olszak distribution [26] if

$$(2.1) \quad X_i C_{jklm} + X_j C_{kilm} + X_k C_{ijlm} = 0.$$

The condition was extensively studied in the geometric literature on pseudo-Riemannian manifolds (see for example [6, 9, 10, 11]).

A contraction of (2.1) with  $g^{im}$  gives  $X^m C_{jklm} = 0$ . Then, a contraction with  $X^i$  gives  $X^i X_i C_{jklm} = 0$ . If the Weyl tensor is non-zero, the Olszak distribution consists of null vectors,  $X^i X_i = 0$ .

**Lemma 2.1.** *On a  $n$ -dimensional Lorentzian manifold with everywhere non-zero Weyl curvature, the Olszak distribution is one-dimensional.*

*Proof.* Suppose that, besides  $X_i$ , there exists another covector  $Y_i$  satisfying (2.1). Then also  $X_i + Y_i$  belongs to the Olszak distribution,  $X^i X_i = 0$ ,  $Y^i Y_i = 0$  and  $X_i Y^i = 0$ . On a Lorentzian manifold, two null orthogonal vectors must be collinear,  $Y_i = \mu X_i$  [35].  $\square$

The following general identity holds for the Weyl tensor on a pseudo-Riemannian manifold (eq. 3.8 in [1]):

$$(2.2) \quad \nabla_i C_{jkl}^m + \nabla_j C_{kil}^m + \nabla_k C_{ijl}^m = \frac{1}{n-3} \nabla_p [\delta_j^m C_{kil}^p + \delta_k^m C_{ijl}^p + \delta_i^m C_{jkl}^p + g_{kl} C_{ji}^{mp} + g_{il} C_{kj}^{mp} + g_{jl} C_{ik}^{mp}]$$

**Proposition 2.2** (Adati and Myazawa [1]). *Let  $(\mathcal{M}, g)$  be a conformally recurrent pseudo-Riemannian manifold of dimension  $n \geq 4$ . Then:*

- 1)  $\nabla_m C_{jkl}^m = 0$  if and only if  $\alpha_i$  belongs to the Olszak distribution;
- 2) If  $\nabla_m C_{jkl}^m = 0$ , then the recurrent covector is null,  $\alpha^i \alpha_i = 0$ .

*Proof.* Obviously  $\nabla_m C_{jkl}^m = 0$  if and only if  $\alpha_m C_{jkl}^m = 0$ . The vanishing of the right hand side of (2.2) and recurrence imply that  $\alpha_i$  belongs to the Olszak distribution. Conversely, we have  $\alpha^i \alpha_i C_{jklm} = 0$ , i.e. either  $\alpha^i \alpha_i = 0$  or  $C_{jklm} = 0$ .  $\square$

**Theorem 2.3** (Roter [31, 34]). *A pseudo-Riemannian manifold  $\mathcal{M}$  of dimension  $n \geq 4$  is simple conformally recurrent if and only if:*

- 1)  $C_{jklm} \neq 0$  everywhere on  $\mathcal{M}$ ,
- 2)  $\nabla_i C_{jklm} = \alpha_i C_{jklm}$ , with recurrence 1-form  $\alpha_i$  that is locally a gradient,
- 3) the Ricci tensor is a Codazzi tensor,  $\nabla_k R_{jl} = \nabla_j R_{kl}$ .

The contraction of the Codazzi property with the metric tensor gives  $\nabla_k R = 0$ , where  $R$  is the scalar curvature. Moreover, it was shown:

**Proposition 2.4** (Roter [31]). *On a non-locally symmetric simple conformally recurrent manifold,  $R = 0$ .*

**Proposition 2.5** (Roter [34], lemma 12). *If the Ricci tensor of a non-locally symmetric simple conformally recurrent manifold has the form  $R_{ij} = \pm d_i d_j$ ,*

then the following equations hold:

$$(2.3) \quad d_i C_{jklm} + d_j C_{kilm} + d_k C_{ijlm} = 0,$$

$$(2.4) \quad d_i R_{jklm} + d_j R_{kilm} + d_k R_{ijlm} = 0.$$

**Proposition 2.6** (Roter [34], theorem 1). *Every simple conformally recurrent Lorentzian manifold is Ricci-recurrent, i.e. there exists a non-zero covector field  $\omega_i$  such that  $\nabla_k R_{jl} = \omega_k R_{jl}$ .*

Brinkmann and pp-wave space-times arise in the presence of a null covariantly constant vector field. They are special cases of Lorentzian manifolds with a recurrent null vector field, which have been studied for a long time (see for example [39, 4, 36, 17, 18]). A characterization of the metric through a set of canonical coordinates was obtained by Walker [39]. We recall Proposition 1 of [18]:

**Proposition 2.7.** *Let  $(\mathcal{M}, g)$  be a Lorentzian manifold of dimension  $n = d + 2 > 2$  with a recurrent null vector field,  $X^k X_k = 0$ ,  $\nabla_k X_j = p_k X_j$ .*

1) *This is equivalent to the existence of coordinates  $(u, x^1, \dots, x^d, v)$ , in which the metric has the local expression*

$$ds^2 = 2dudv + H(u, \vec{x}, v)du^2 + \sum_{\rho=1}^d a_\rho(u, \vec{x})dudx^\rho + \sum_{\mu, \nu=1}^d g_{\mu\nu}^*(u, \vec{x})dx^\mu dx^\nu,$$

where  $g_{\mu\nu}^*$  and  $a_\rho$  are independent of the coordinate  $v$ , and  $H$  is a smooth function. We refer to these coordinates as Walker coordinates.

2)  $\nabla_k X_j = 0$  if and only if  $H$  does not depend on  $v$ . We refer to these coordinates as Brinkmann coordinates, and to the manifold as a Brinkmann-wave (space-time) [4].

A pp-wave is a Brinkmann-wave with some restrictions. Schimming [36] gave the following coordinate characterization (see also [19, 17, 18]):

**Proposition 2.8** (Schimming [36]). *A Lorentzian manifold of dimension  $d + 2 > 2$  is a pp-wave if and only if there exist coordinates  $(u, \vec{x}, v)$  in which the metric has the following local expression:*

$$(2.5) \quad ds^2 = 2du dv + H(\vec{x}, u)du^2 + \sum_{\rho=1}^d dx^{\rho 2}$$

where  $H(\vec{x}, u)$  is a smooth function independent of  $v$ , usually called the “potential function of the pp-wave”.

The expression of the metric yields the Ricci tensor of a pp-wave:

$$(2.6) \quad R_{kl} = \psi(\vec{x}, u) X_k X_l, \quad \psi(\vec{x}, u) = -\frac{1}{2} \sum_{\rho=1}^d \frac{\partial^2 H}{\partial x^{\rho 2}}$$

where  $X_k = \nabla_k u$  is a covariantly constant null vector ( $X^k X_k = 0$ ,  $\nabla_j X_k = 0$ ) [28]. It follows that the scalar curvature is zero,  $R = 0$ .

**Remark 2.9.** *A metric such that  $R_{kl} = \psi X_k X_l$  with a null recurrent covector is called a pure radiation metric with parallel rays, or aligned pure radiation metric [20].*

**Proposition 2.10** (Schimming [36], see also [19, 17, 18]). *A Lorentzian manifold of dimension  $d + 2 > 2$  with covariantly constant null vector field (i.e. a Brinkmann wave) is a pp-wave if and only if one of the following conditions is satisfied:*

$$(2.7) \quad X_i R_{jklm} + X_j R_{kilm} + X_k R_{ijlm} = 0;$$

$$(2.8) \quad R_{jklm} = X_j X_m D_{kl} - X_j X_l D_{mk} - X_k X_m D_{jl} + X_k X_l D_{jm};$$

$$(2.9) \quad R^p{}_{jk}{}^q R_{plmq} = \chi X_j X_k X_l X_m;$$

$D_{ij}$  is a symmetric tensor and  $\chi$  is a suitable scalar function.

A contraction recovers the form of the Ricci tensor of a pp-wave,  $R_{kl} = \psi X_k X_l$ . Leistner and Nurowski showed that in dimension  $d + 2 = 4$  the conditions are equivalent to  $R_{jk}{}^{pq} R_{pqlm} = 0$  [19].

**Remark 2.11.** *It is worth noting that for a Ricci tensor of the form (2.6) we have*

$$(2.10) \quad X_i C_{jklm} + X_j C_{kilm} + X_k C_{ijlm} = X_i R_{jklm} + X_j R_{kilm} + X_k R_{ijlm}.$$

*A contraction with  $g^{im}$  and the equality  $R_{ij} = \psi X_i X_j$  imply  $X^m C_{jklm} = X^m R_{jklm}$ .*

*For a pp-wave, by eq.(2.7), we have  $X_i C_{jklm} + X_j C_{kilm} + X_k C_{ijlm} = 0$ , and so  $X^m C_{jklm} = X^m R_{jklm} = 0$ .*

### 3. SIMPLE CONFORMALLY RECURRENT LORENTZIAN MANIFOLDS AND PP-WAVES.

In this section we obtain two important results. First we show, in several steps, that a conformally recurrent pp-wave space-time is a simple conformally recurrent space-time. Next, we show that a simple conformally recurrent space-time is a pp-wave space-time. Finally, some curvature properties of conformally recurrent pp-wave space-times are presented, as well as some

technical results.

**Lemma 3.1.** *On a pp-wave space-time of dimension  $n \geq 4$ , the Ricci tensor: 1) is recurrent with a closed recurrence 1-form, 2) it satisfies  $[\nabla_i, \nabla_j]R_{kl} = 0$ , 3) it satisfies*

$$(3.1) \quad \nabla^j \nabla^m C_{jklm} = -\frac{n-3}{n-2} \nabla^2 R_{kl}$$

where  $\nabla^2 = \nabla_k \nabla^k$ . In particular,  $\nabla^2 R_{kl} = 0$  if and only if  $\nabla^j \nabla^m C_{jklm} = 0$ .

*Proof.* In a coordinate system with (2.5) one has  $\nabla_j R_{kl} = (\nabla_j \psi) X_k X_l = (\nabla_j \log |\psi|) R_{kl}$ . From this we infer  $[\nabla_i, \nabla_j]R_{kl} = 0$ , i.e. a pp-wave space-time is Ricci semi-symmetric [30].

The divergence of the Weyl tensor has the general expression

$$(3.2) \quad \nabla_m C_{jkl}{}^m = \frac{n-3}{n-2} (\nabla_k R_{jl} - \nabla_j R_{kl}) + \frac{n-3}{2(n-1)(n-2)} (g_{kl} \nabla_j R - g_{jl} \nabla_k R)$$

A further covariant divergence gives

$$\nabla^j \nabla^m C_{jklm} = -\frac{n-3}{n-2} \left[ \nabla^2 R_{kl} - \frac{g_{kl} \nabla^2 R}{2(n-1)} - [\nabla_j, \nabla_k] R_l^j \right] + \frac{n-3}{2(n-1)} \nabla_k \nabla_l R.$$

Ricci semi-symmetry and the property  $R = 0$  give (3.1).  $\square$

**Lemma 3.2** (see [21] and [23] theorem 6.1.). *A pp-wave space-time of dimension  $d+2 \geq 4$  with the metric (2.5) satisfies  $\nabla_m C_{jkl}{}^m = 0$  if and only if  $\nabla_k \psi = \lambda X_k$ .*

*Proof.* For a pp-wave space-time we have  $\nabla_j R_{kl} = (\nabla_j \psi) X_k X_l$  and  $R = 0$ . Eq.(3.2) becomes:

$$(3.3) \quad \nabla_m C_{jkl}{}^m = \frac{n-3}{n-2} [(\nabla_k \psi) X_j - (\nabla_j \psi) X_k] X_l.$$

It follows that  $\nabla_m C_{jkl}{}^m = 0$  if and only if  $(\nabla_k \psi) X_j = (\nabla_j \psi) X_k$ , which is equivalent to the condition  $\nabla_k \psi = \lambda X_k$  for some scalar function.  $\square$

**Remark 3.3.** *For a pp-wave the condition  $\nabla_m C_{jkl}{}^m = 0$  is equivalent to  $\nabla_k R_{jl} = \nabla_j R_{kl}$ , i.e. to the Ricci tensor being a Codazzi tensor. This follows from (3.2) and  $R = 0$ .*

**Proposition 3.4.** *Let  $(\mathcal{M}, g)$  be a  $d+2 \geq 4$  dimensional pp-wave space-time with the metric (2.5). If  $\psi = -\frac{1}{2} \sum_{\rho=1}^d \partial^2 H / \partial x_\rho^2$  only depends on  $u$ , then  $\nabla^2 R_{kl} = 0$ .*

*Proof.* If  $\psi = \psi(u)$  we have  $\nabla_k \psi = \partial_k \psi(u) = \psi'(u) X_k$  and  $\nabla_m C_{jkl}{}^m = 0$  by the previous lemma.

Therefore  $\nabla_k R_{ij} = \psi' X_k X_i X_j$  and  $\nabla^2 R_{ij} = \psi''(u) X^k X_k X_i X_j = 0$ .  $\square$

Galaev classified the indecomposable conformally recurrent Lorentzian manifolds [12]. He showed that either the manifold is conformally flat, or  $\nabla_i R_{jklm} = 0$ , or it is a pp-wave. Then he found all pp-wave potential functions  $H(\vec{x}, u)$  such that the Weyl tensor in the metric (2.5) is recurrent,  $\nabla_i C_{jklm} = \alpha_i C_{jklm}$ . The potential solves the following system of  $d+1$  equations:

$$(3.4) \quad (\alpha_\rho - \partial_\rho) \Omega_{\mu\nu} = 0 \quad (\rho = 1 \dots d)$$

$$(3.5) \quad (\alpha_u - \partial_u) \Omega_{\mu\nu} = 0$$

where  $\Omega_{\mu\nu} = \partial_\mu \partial_\nu H - \frac{1}{n} \delta_{\mu\nu} \sum_\rho \partial_\rho^2 H$  ( $\mu, \nu = 1 \dots d$ ).

Let  $\Omega^2 = \sum_{\mu\nu} (\Omega_{\mu\nu})^2$ . In an open set  $\mathcal{O} \subset \mathcal{M}$  where  $\Omega^2 \neq 0$  the equations give  $\alpha_\mu = \frac{1}{2} \partial_\mu \log \Omega^2$  and  $\alpha_u = \frac{1}{2} \partial_u \log \Omega^2$ . The potential for which the metric (2.5) is conformally recurrent turns out to be

$$(3.6) \quad H(\vec{x}, u) = \sum_{\rho=1}^d x_\rho^2 [a(u) + F(u) \lambda_\rho^2]$$

with functions  $a(u)$ ,  $F(u)$  and real numbers  $\lambda_1 + \dots + \lambda_d = 0$ . Correspondingly,  $\psi(\vec{x}, u) = -na(u)$  and, since it only depends on  $u$ , we conclude that for conformally recurrent pp-waves the divergence of the conformal tensor vanishes:

**Proposition 3.5.** *On a conformally recurrent pp-wave space-time with the metric (2.5), we have  $\nabla_m C_{jkl}{}^m = 0$ ; moreover there exists a domain  $\mathcal{O}$  where the recurrence covector is a gradient.*

The recurrence 1-form  $\alpha_i$  being closed, it follows that  $[\nabla_p, \nabla_q] C_{jklm} = 0$ , i.e. the manifold is Weyl semi-symmetric (see [8]). Moreover, as  $[\nabla_i, \nabla_j] R_{kl} = 0$ , the manifold is also semi-symmetric,  $[\nabla_p, \nabla_q] R_{jklm} = 0$ .

Now, from Proposition 3.5, Remark 3.3 and Theorem 2.3, we have:

**Theorem 3.6.** *In  $n \geq 4$  any conformally recurrent pp-wave space-time is a simple conformally recurrent Lorentzian manifold.*

Some further properties of conformally recurrent pp-wave space-times are collected here:

**Proposition 3.7.** *For a conformally recurrent pp-wave space-time:*

- 1) *the recurrence covector  $\alpha_i$  is null and collinear with the covariantly constant covector  $X_i = \nabla_i u$*
- 2) *the recurrence covector is recurrent with closed recurrence 1-form.*

*Proof.* From Propositions 3.5 and 2.2 we get  $\alpha_i C_{jklm} + \alpha_j C_{kilm} + \alpha_k C_{ijlm} = 0$  i.e. the recurrence covector of a conformally recurrent pp-wave space-time

belongs to the Olszak distribution. In view of Remark 2.11, for a pp-wave space-time also  $X_i$  belongs to the Olszak distribution; then, by Lemma 2.1,  $\alpha_i = \mu X_i$ . Taking the covariant derivative, we see that  $\nabla_j \alpha_i = (\nabla_j \mu) X_i = (\nabla_j \log |\mu|) \alpha_i \equiv q_j \alpha_i$  (i.e. the recurrence vector is itself recurrent, with closed recurrence 1-form).  $\square$

**Proposition 3.8.** *Let  $(\mathcal{M}, g)$  be a  $d + 2 \geq 4$  dimensional pp-wave space-time with the metric (2.5). Then there exists a coordinate domain  $\mathcal{O}$  in  $\mathcal{M}$  where  $\nabla^2 C_{jklm} = 0$  and  $\nabla^2 R_{jklm} = 0$ .*

*Proof.* From the closedness condition  $\nabla_j \alpha_i = \nabla_i \alpha_j$  we infer  $q_i \alpha_j = q_j \alpha_i$  so that  $q_j = \rho \alpha_i$  and the recurrence 1-form  $\alpha_i$  satisfies  $\nabla_j \alpha_i = \rho \alpha_i \alpha_j$  and  $\nabla^i \alpha_i = 0$ . Thus, since  $\alpha^i \alpha_i = 0$ , we have  $\alpha^i \nabla_i C_{jklm} = 0$  and consequently  $\nabla^i \nabla_i C_{jklm} = \nabla^i (\alpha_i C_{jklm}) = (\nabla^i \alpha_i) C_{jklm} + \alpha^i (\nabla_i C_{jklm}) = 0$ . Moreover we have  $\nabla^i \nabla_i R_{kl} = 0$  so that  $\nabla^i \nabla_i R_{jklm} = 0$ .  $\square$

Now we consider a  $n$ -dimensional Lorentzian simple conformally recurrent manifold and show that it is a conformally recurrent pp-wave space-time.

**Theorem 3.9.** *Let  $(\mathcal{M}, g)$  be a Lorentzian simple conformally recurrent manifold of dimension  $n \geq 4$ : then there exists a coordinate domain  $\mathcal{O}$  in  $\mathcal{M}$  where the manifold is a pp-wave space-time.*

*Proof.* From Theorem 2.3 and Proposition 2.4 we have  $\nabla_m C_{jkl}^m = 0$  so that  $\alpha^m C_{jklm} = 0$  and  $\alpha_i C_{jklm} + \alpha_j C_{kilm} + \alpha_k C_{ijlm} = 0$ . Taking the covariant derivative, we get  $(\nabla_p \alpha_i) C_{jklm} + (\nabla_p \alpha_j) C_{kilm} + (\nabla_p \alpha_k) C_{ijlm} = 0$ . Since in a Lorentzian manifold the Olszak distribution is one-dimensional (Lemma 2.1), this implies proportionality,  $\nabla_i \alpha_j = p_i \alpha_j$  for some 1-form  $p_j$ .

Proposition 2.6 states that a simple conformally recurrent Lorentzian manifold is Ricci recurrent, i.e.  $\nabla_k R_{jl} = \omega_k R_{jl}$  for some 1-form  $\omega_k$ . Furthermore, by Theorem 2.3, the Ricci tensor is Codazzi,  $\nabla_k R_{jl} = \nabla_j R_{kl}$ . Then we have

$$(3.7) \quad \omega_j R_{kl} = \omega_k R_{jl}.$$

Contraction with  $g^{jl}$  gives  $\omega^l R_{kl} = 0$  because  $R = 0$  (theorem 2.3). Thus on multiplying (3.7) by  $\omega_k$  it follows that  $(\omega^k \omega_k) R_{jl} = 0$ , i.e.  $\omega_k$  is a null vector.

Let  $\theta^k$  be a vector such that  $\theta^k \omega_k = 1$ . Eq.(3.7) gives  $R_{jl} = \omega_j \theta^k R_{kl}$  and, by symmetry,  $\omega_j \theta^k R_{kl} = \omega_l \theta^k R_{kj}$ . Thus  $\theta^j R_{jl} = \omega_l (\theta^k \theta^j R_{kj})$  from which we finally get:

$$(3.8) \quad R_{ij} = \psi \omega_i \omega_j$$



where  $\psi = \theta^m \theta^j R_{mj}$  is a scalar function.

Let  $d = \psi/|\psi|$  and  $d_j = \omega_j \sqrt{|\psi|}$ . Then  $R_{ij} = dd_i d_j$  with  $d^k d_k = 0$ . By Proposition 2.5, equations (2.3) and (2.4) are recovered. In this way the vector  $d_j$  belongs to the one-dimensional Olszak distribution: thus  $d_i = \epsilon \alpha_i$  and the Ricci tensor takes the form  $R_{jl} = \phi \alpha_j \alpha_l$ . Furthermore, from

$$(3.9) \quad \alpha_i R_{jklm} + \alpha_j R_{kilm} + \alpha_k R_{ijlm} = 0$$

Remark 2.11 gives:  $\alpha^m R_{jklm} = 0$ . Taking the covariant derivative of  $\nabla_k \alpha_l = p_k \alpha_l$  and using skew symmetrisation, we obtain  $\alpha^m R_{jklm} = (\nabla_j p_k - \nabla_k p_j) \alpha_l = 0$ . Thus the covector  $p_j$  is locally a gradient, i.e. there exists a coordinate domain  $\mathcal{O}$  of  $(\mathcal{M}, g)$  where  $p_j = \nabla_j \eta$ . The covector  $\bar{\alpha}_j = \alpha_j e^{-\eta}$  is covariantly constant ( $\nabla_k \bar{\alpha}_j = 0$ ). Eq.(3.9) with  $\bar{\alpha}_j$  is Schimming's condition (2.7) for the manifold to be locally a pp-wave space-time.  $\square$

#### 4. SOME CONSEQUENCES FOR EXTENDED THEORIES OF GRAVITATION

We derive some consequences for extended theories of gravitation in dimensions  $n \geq 4$ . Since for a pp-wave the scalar curvature is zero, a pp-wave solves the Einstein's field equations (in natural units)

$$R_{ij} - \frac{1}{2} R g_{ij} = 8\pi T_{ij}$$

with a pure radiation source,  $T_{ij} = \Phi^2 k_i k_j$ , where  $k_i k^i = 0$  (see [37], eq. 5.8).

Generic theories of gravitation modify Einstein's theory at short distances by expressing the action integral with zero source as a power series

$$(4.1) \quad I = \int d^n x \sqrt{-g} \left[ R - 2\Lambda_0 + \alpha R^2 + \beta R_{ij} R^{ij} + \gamma (R_{jklm} R^{jklm} - 4R_{jk} R^{jk} + R^2) + \sum_{p>2} C_p (\text{Riem}, \text{Ric}, \nabla \text{Riem}, \dots) \right]$$

where  $\alpha, \beta, \gamma, C_p$  are parameters provided by some microscopic theory such as string theory. The quadratic part represents “quadratic gravity” (Weyl-Eddington terms), and the terms of higher order are contracted products of the Riemann tensor and its derivatives. Such corrections scale as powers of  $\ell_P/L$ , where  $\ell_P$  is Planck's length and  $L$  is a typical length for the variation of the metric [40], and could be significant in the early evolution of the universe.

The field equations of the full theory are complicated; nevertheless Gürses et al. [14] proved that for pp-wave metrics (2.5) the field equations with

cosmological constant  $\Lambda_0 = 0$  may be written as:

$$(4.2) \quad [a_0 + a_1 \nabla^2 + a_2 (\nabla^2)^2 + \dots] R_{kl} = 0$$

where  $a_i$  are constants depending on  $\alpha, \beta, \gamma, C_p$ . If  $\nabla^2 R_{kl} = 0$  the equation reduces to  $a_0 R_{kl} = 0$ , with the vacuum solution. In the same paper the authors noted that for pp-waves with  $\nabla^2 R_{kl} = 0$  the field equations may include a pure radiation source:

$$(4.3) \quad a_0 R_{kl} = T_{kl}$$

It is thus interesting to find non-vacuum pp-wave solutions. According to Proposition 3.4, if  $\psi = -\frac{1}{2} \sum_{\rho=1}^d \partial_\rho^2 H$  depends only on  $u$ , then  $\nabla^2 R_{kl} = 0$ .

**Proposition 4.1.** *Let  $(\mathcal{M}, g)$  be a  $d+2 \geq 4$  dimensional pp-wave space-time with the metric (2.5): if  $\sum_{\rho=1}^d \partial_\rho^2 H$  depends only on  $u$ , then the metric solves the field equations of the generic theory of gravitation with pure radiation source.*

We give two examples where  $\nabla^2 R_{kl} = 0$ :

- 1) the pp-wave metric is conformally recurrent, with potential  $H$  given by (3.6).
- 2) the pp-wave metric is two-symmetric i.e.  $\nabla_p \nabla_i R_{jklm} = 0$ . This occurs if and only if the potential function of the pp-wave has the form

$$(4.4) \quad H(\vec{x}, u) = \sum_{\mu, \nu=1}^d (ua_\mu \delta_{\mu\nu} + b_{\mu\nu}) x^\mu x^\nu$$

where  $0 \leq a_1 \leq \dots \leq a_d$  and  $b_{\mu\nu} = b_{\nu\mu}$  are real numbers (see [2, 3, 13]). One evaluates  $\psi(u) = -\sum_{\mu=1}^d (ua_\mu + b_{\mu\mu})$ . Thus for two-symmetric pp-waves space-times the divergence of the conformal tensor vanishes and, as one may directly calculate,  $\nabla^2 R_{kl} = 0$ .

## REFERENCES

- [1] T. Adati and T. Miyazawa, *On a Riemannian space with recurrent conformal curvature*, Tensor ( N.S.) 18 (1967), 348-354.
- [2] D. V. Alekseevsky and A. S. Galaev, *Two-symmetric Lorentzian manifolds*, J. Geom. Phys. 61 (2011), 2331-2340.
- [3] O. F. Blanco, M. Sánchez and J. M. Senovilla, *Complete classification of second order symmetric spacetimes*, J. Phys.: Conf. ser. 229 (2010), 012021 (5pp).
- [4] H. W. Brinkmann, *Einstein spaces which are mapped conformally on each other*, Math. Ann. 94 (1925), 119-145.

- [5] U.C. De and C. A. Mantica, *Conformally recurrent space-times admitting a proper conformal vector field*, Commun. Korean Math. Soc. 29 (2) (2014) 319–329.
- [6] F. Defever and R. Deszcz, *On semi Riemannian manifolds satisfying the condition  $R \cdot R = Q(S, R)$* , in Geometry and Topology of Submanifolds, III, World Scientific Publ. Singapore. (1991), 108–130.
- [7] A. Derdziński, *Conformally recurrent indefinite metrics on tori*, Tensor (N.S.) 34 (1980), 260–262.
- [8] R. Deszcz, *Examples of four dimensional Riemannian manifolds satisfying some pseudosymmetry curvature conditions*, in Geometry and Topology of Submanifolds II (World Scientific Teaneck, NJ, 1990), pp.134–143.
- [9] R. Deszcz and W. Grycak, *On certain curvature conditions on Riemannian manifolds*, Colloq. Math. 58 (1) (1990), 259–268.
- [10] R. Deszcz and M. Hotłoś, *On a certain extension of the class of semisymmetric manifolds*, Publ. de l’Institut Math. (Beograd) (N.S.), tome 63 (77) (1998), 115–130.
- [11] R. Deszcz, M. Hotłoś, and Z. Sentürk, *On the equivalence of the Ricci-pseudosymmetry and pseudosymmetry*, Colloq. Math. 79 (1999), 211–221.
- [12] A. S. Galaev, *Some applications of the Lorentzian holonomy algebras*, J. Geom. Symmetry Phys. 26 (2012), 13–31.
- [13] S. Gavino-Fernández, *The geometry of Lorentzian Ricci solitons*, Ph.D. Thesis, Publicaciones del Departamento de Geometria y Topologia, Universidade de Santiago de Compostela, (2012) 105pp.
- [14] M. Gürses, T. Ç. Şişman and B. Tekin, *AdS-plane wave and pp-wave solutions of generic gravity theories*, Phys. Rev. D 90 (2014), 124005.
- [15] G. S. Hall, *Complex recurrent space-times*, J. Phys. A: Math. Nucl. Gen. 7 (1974), L 42–4.
- [16] G. S. Hall, *Recurrence condition in space-time*, J. Phys. A: Math. Gen. 10 (1) (1977), 29–42.
- [17] T. Leistner, *Conformal holonomy of C-spaces, Ricci-flat, and Lorentzian manifolds*, Diff. Geom. Appl. 24 (2006), 458–478.
- [18] T. Leistner, *Screen bundles of Lorentzian manifolds and some generalizations of pp-waves*, J. Geom. Phys. 56 (10) (2006), 2117–2134.
- [19] T. Leistner, P. Nurowski, *Ambient metrics for n-dimensional pp-waves*, Commun. Math. Phys. 296 (3) (2010), 881–898.

- [20] T. Leistner and P. Nurowski, *Conformal pure radiation with parallel rays*, Class. Quantum Grav. 29 (2012), 055007, (15pp).
- [21] C. A. Mantica and L. G. Molinari, *On conformally recurrent manifolds of dimension greater than 4*, Int. J. Geom. Meth. Mod. Phys. 13 (2016), 1650053 (17 pages).
- [22] C. A. Mantica and Y. J. Suh, *Recurrent conformal 2-forms on pseudo-Riemannian manifolds*, Int. J. Geom. Meth. Mod. Phys. 11 (6) (2014), 1450056, (29 pp).
- [23] C. A. Mantica and Y. J. Suh, *Pseudo-Z symmetric space-times with divergence-free Weyl tensor and pp-waves*, Int. J. Geom. Meth. Mod. Phys. 13 (2) (2016), 1650015.
- [24] R. G. Mc. Lenaghan and J. Leroy, *Complex recurrent space-times*, Proc. Roy. Soc. London A 327 (1569) (1972), 229–249.
- [25] R. G. Mc Lenaghan and H. A. Thompson, *Second order recurrent space-times in General Relativity*, Lett. Nuovo Cimento 5 (7) (1972) 563–564.
- [26] Z. Olszak, *On conformally recurrent manifolds, I: Special distributions*, Zesz. Nauk. Politech. Śl., Mat.-Fiz. 68 (1993), 213–225.
- [27] M. Ortaggio, V. Pravda and A. Pravdova, *Algebraic classification of higher dimensional spacetimes based on null alignment*, Class. Quantum Grav. 30 (2013), 013001 (57pp).
- [28] J. Podolský, and M. Žofka, *General Kundt spacetimes in higher dimensions*, Class. Quantum Grav. 26 (10) (2009) 105008.
- [29] M. M. Postnikov, *Geometry VI, Riemannian geometry*, Encyclopaedia of Mathematical Sciences, Vol. 91, 2001, Springer-Verlag, Berlin.
- [30] W. Roter, *Some remarks on second order recurrent spaces*, Bull. Acad. Polon. Sci. 12 (1964) 207–211.
- [31] W. Roter, *On a class of conformally recurrent manifolds*, Tensor (N.S.) 39 (1982), 207–217.
- [32] W. Roter, *On conformally recurrent Ricci-recurrent manifolds*, Colloq. Math. 46 (1982), 45–57.
- [33] W. Roter, *On conformally related conformally recurrent metrics. I Some general results*, Colloq. Math. 47 (1982), 315–327.
- [34] W. Roter, *On the existence of certain conformally recurrent metrics*, Colloq. Math. 51 (1987), 39–46.
- [35] J. Santos, M. J. Rebouças and A. F. F. Teixeira, *Segre types of symmetric two-tensors in  $n$ -dimensional space times*, Gen. Rel. Grav. 27 (9) (1995), 989–999.

- [36] R. Schimming, *Riemannsche Räume mit ebenfrontiger und mit ebener Symmetrie*, Mathematische Nachrichten 59 (1974), 128–162.
- [37] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers and E. Hertl, *Exact Solutions of Einstein's Field Equations*, Cambridge University Press, 2nd ed. (2003).
- [38] Y. J. Suh and J-H. Kwon, *Conformally recurrent semi-Riemannian manifolds*, Rocky Mountain J. of Math. 35 (1) (2005), 285–307.
- [39] A. G. Walker, *On parallel fields of partially null vector spaces*, Quart. J. of Math. 20 (1949), 135–145.
- [40] A. Zee, *Einstein gravity in a nutshell*, Princeton University Press (2013).

I.I.S. LAGRANGE, VIA L. MODIGNANI 65, 20161, MILANO, ITALY  
*E-mail address:* carloalberto.mantica@libero.it

PHYSICS DEPARTMENT, UNIVERSITÀ DEGLI STUDI DI MILANO, AND I.N.F.N. SEZ.  
 MILANO, VIA CELORIA 16, 20133 MILANO, ITALY  
*E-mail address:* luca.molinari@unimi.it