

On a topological version of Pach's overlap theorem

Boris Bukh*

Alfredo Hubard[†]

Abstract

Pach showed that every $d + 1$ sets of points $Q_1, \dots, Q_{d+1} \subset \mathbb{R}^d$ contain linearly-sized subsets $P_i \subset Q_i$ such that all the transversal simplices that they span intersect. We show, by means of an example, that a topological extension of Pach's theorem does not hold with subsets of size $C(\log n)^{1/(d-1)}$. We show that this is tight in dimension 2, for all surfaces other than \mathbb{S}^2 . Surprisingly, the optimal bound for \mathbb{S}^2 in the topological version of Pach's theorem is of the order $(\log n)^{1/2}$. We conjecture that, among higher-dimensional manifolds, spheres are similarly distinguished. This improves upon the results of Bárány, Meshulam, Nevo and Tancer.

Introduction

Pach's overlap theorem. Let Q_1, \dots, Q_{d+1} be $d + 1$ sets of points in \mathbb{R}^d . A family of subsets P_1, \dots, P_{d+1} with $P_i \subset Q_i$ is a *Pach family* if all the simplices of the form $p_1 \dots p_{d+1}$ with $p_i \in P_i$ have a common intersection point. Pach's overlap theorem [15] asserts that if each set Q_i is of size n then there is a Pach family with $|P_i| \geq c_d n$. The constant c_d depends only on the dimension and its optimal value is unknown. The same-type lemma of Bárány and Valtr [3] is closely related to Pach's result.

Topological version. Many results in combinatorial geometry admit a topological generalization by replacing linear functions with arbitrary continuous functions. A celebrated example concerns planar graphs: a graph can be drawn in the plane with straight edges if and only if it can be drawn with 'curvy' edges. Closer to the subject of the present paper are the topological Tverberg theorem of Bárány, Shlosman and Szűcs [2] and Gromov's topological overlap theorem [10], which generalize Tverberg's theorem [20] and Bárány's overlap theorem [4, Theorem 5.1] respectively.

It is natural to ask if Pach's overlap theorem admits such a generalization. In order to state it, we need some standard notation. For a simplicial complex Y , let $Y^{\leq d}$ be the d -skeleton of Y , i.e., the complex consisting of all the simplices of dimension $\leq d$. If Y_1, Y_2 are simplicial complexes, $Y_1 * Y_2 \stackrel{\text{def}}{=} \{A \sqcup B : A \in Y_1, B \in Y_2\}$ denotes their join (where \sqcup denotes the disjoint union). In the

*Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh, PA 15213, USA. Supported in part by Sloan Research Fellowship and by U.S. taxpayers through NSF CAREER grant DMS-1555149. Part of the work was done during the visit to Université Paris-Est Marne-la-Vallée supported by LabEx Bézout (ANR-10-LABX-58).

[†]Laboratoire d'Informatique Gaspard-Monge, Université Paris-Est Marne-la-Vallée, France. alfredo.hubard@u-pem.fr.

following, we let V_1, V_2, \dots, V_{d+1} be $d+1$ point sets of size n each. By a slight abuse of notation, we treat each V_i also as a 0-dimensional simplicial complex. Put

$$X \stackrel{\text{def}}{=} V_1 * \dots * V_{d+1}.$$

The complex X consists of d -simplices, which contain one vertex from each V_i , and their faces.

Definition 1. Consider a continuous map $\phi: X \rightarrow M$ into a d -dimensional manifold. A topological Pach family for ϕ consists of sets $P_i \subset V_i$ and a point $p \in M$ such that the ϕ -image of every d -simplex in $P_1 * \dots * P_{d+1}$ contains p . Let $f_M(n)$ be the largest integer such that, for every continuous map $\phi: X \rightarrow M$, there is a topological Pach family all of whose sets are of size $f_M(n)$.

When $M = \mathbb{R}^d$ and ϕ is an affine map, this notion reduces to the usual (non-topological) Pach family. In this paper, the map ϕ is an arbitrary continuous map into a topological manifold M which is assumed to have no boundary, but is not necessarily compact. In [5], it was shown that, unlike in the affine case, a topological Pach family of linear size need not exist. Namely, [5] gave a construction demonstrating that $f_{\mathbb{R}^d}(n) \leq C_d n^{1/d}$; the same paper also showed that $f_{\mathbb{R}^d}(n) \geq c_d (\log n)^{1/d}$. We present a better construction and improve the lower bound in dimension 2, showing that our construction is tight in this case.

Below and everywhere else in the paper the logarithms are to base $e = 2.71\dots$

Theorem 2. For every manifold M of dimension $d \geq 2$,

$$f_M(n) \leq 30(\log n)^{1/(d-1)},$$

for n large enough. This is sharp for $d = 2$: for every surface $M \neq \mathbb{S}^2$,

$$f_M(n) \geq 10^{-14} \log n.$$

Much to our surprise the asymptotics of $f_M(n)$ is sensitive to the target space. The asymptotics of $f_{\mathbb{S}^2}(n)$ is different from the asymptotics of $f_M(n)$ for any other surface $M \neq \mathbb{S}^2$!

Theorem 3. For each $d \geq 1$,


$$\alpha_d \leq \frac{f_{\mathbb{S}^d}(n)}{(\log n)^{1/d}} \leq 2,$$

for n large enough. Here, $\alpha_d = \frac{2^{-d^2-1}}{(d+1)!}$.

We begin with the proof of the easiest result, which is Theorem 3. We then explain the similar, but more complicated Theorem 2. We finish this paper with some remarks and open problems.

The case $M = \mathbb{S}^d$: proof of Theorem 3

Construction. We identify \mathbb{S}^d with the one-point compactification of \mathbb{R}^d . Map the vertices of X to a set in general position, and then extend to the $(d-1)$ -skeleton by linearity. For each d -face of X , there are two natural ways of extending ϕ to that face. We may stay inside \mathbb{R}^d and fill in the simplex by extending ϕ linearly, or we may use the point at infinity to obtain the ‘inverted’ filling. We choose one of these two ways uniformly at random, for each d -face.



The usual (left) and the ‘inverted’ (right) fillings of a triangle

The $(d+1)n$ vertices of X span $\binom{(d+1)n}{d}$ hyperplanes, which in turn partition \mathbb{S}^d into $O(n^{d^2})$ regions. The points in each such region are covered by the same set of ϕ -images of d -faces of X . Hence, the common intersection point of every Pach family can be taken to be a center of one of the regions. Let $m \stackrel{\text{def}}{=} 2(\log n)^{1/d}$. Given a point $p \in \mathbb{S}^d$ and a $(d+1)$ -tuple of m -element sets (P_1, \dots, P_{d+1}) , the probability that p is a common intersection point of all the simplices spanned by P_1, \dots, P_{d+1} is $2^{-m^{d+1}}$ since the fillings are chosen independently. By the union bound over all tuples (P_1, \dots, P_{d+1}) and over all region centers p , it follows that a Pach family of size m exists with probability at most $O\left(\binom{n}{m}^{d+1} n^{d^2} 2^{-m^{d+1}}\right) = O\left((n^{d+1} 2^{-m^d})^m n^{d^2}\right) = O(n^{-m} n^{d^2}) = o(1)$. In particular, for some choice of fillings, a Pach family of this size does not exist.

Lower bound. This argument is from [5], so we only sketch it. Denote by $X^{=d}$ the set of d -simplices in X . By Gromov’s topological overlap theorem [10], for any continuous map $\phi: X \rightarrow \mathbb{S}^d$, there is a point p that is in the ϕ -images of $\alpha_d |X^{=d}|$ many d -faces of X , where $\alpha_d > 0$ is an absolute constant. Let $H = \{\sigma \in X^{=d} : p \in \phi(\sigma)\}$. Since each d -simplex contains exactly one vertex from each V_i , we can regard H as $(d+1)$ -partite $(d+1)$ -uniform hypergraph with parts V_1, \dots, V_{d+1} . Every $(d+1)$ -partite $(d+1)$ -uniform hypergraph of density $\alpha > 0$ contains a complete $(d+1)$ -subhypergraph all of whose parts are of size $\lfloor \alpha(\log n)^{1/d} \rfloor$. This is Nikiforov’s explicit estimate [14] for Erdős’s generalization [9] of the Kővari–Sos–Turán theorem.

The explicit value for α_d of $1/(d+1)!2^{d^2+1}$ will be obtained in the next section, after a more careful discussion of Gromov’s theorem. Plugging this value into Nikiforov’s estimate, we obtain the desired lower bound.

The case $M \neq \mathbb{S}^d$: proof of Theorem 2

Construction. This is similar to the construction for \mathbb{S}^d , except that we cannot randomize the d -faces of X . We randomize the $(d-1)$ -faces instead.

We say that H is a (d, n) -graph if H is a complete $(d+1)$ -partite d -uniform hypergraph in which each part is of size n . In our probabilistic construction we will require an upper bound on the probability that a random subgraph of a (d, n) -graph contains no $(d+1)$ -clique. It is possible to prove a very precise bound using hypergraph containers (see [18, Corollary 2.4] or [1, Theorem 8.1] for similar non-partite results). Instead of doing that, we opt for a simple proof of a cruder upper bound using a restatement of the Loomis–Whitney inequality [12] due to Bollobás–Thomason [6].

For a set family $\mathcal{F} \subset 2^X$ and a set $Y \subset X$, we define the *trace on Y* by $\mathcal{F}|_Y \stackrel{\text{def}}{=} \{S \cap Y : S \in \mathcal{F}\}$.


Lemma 4 (Theorem 7 from [6]). *Let E be a finite set, and $\mathcal{C} \subset 2^E$ be a family of subsets of E such that each $x \in E$ is in equally many sets of \mathcal{C} . Let F be a uniformly random subset of E . Suppose that a family $\mathcal{F} \subset 2^E$ and a constant $c < 1$ satisfy $\Pr[F \cap Y \in \mathcal{F} | Y] \leq c^{|Y|}$ for every $Y \in \mathcal{C}$. Then $\Pr[F \in \mathcal{F}] \leq c^{|E|}$.*

Lemma 5. *Each (d, n) -graph admits a coloring of its edges into two colors so that each (d, m) -subgraph contains monochromatic $(d+1)$ -cliques in each of the two colors, for all $m \geq 25(\log n)^{1/(d-1)}$.*

Proof. Let H be any (d, m) -subgraph of the (d, n) -graph. Let E be the edge set of H , let \mathcal{F} be the set of all subgraphs of H that contain no $(d+1)$ -clique, and let \mathcal{C} be the set of all $(d+1)$ -cliques in H . Then $\Pr[F \cap Y \in \mathcal{F} | Y] = 1 - 2^{-d-1}$ for a random $F \subset E$ and any $(d+1)$ -clique Y . Because each edge is covered by equally many $(d+1)$ -cliques and $|E| = (d+1)m^d$, Lemma 4 implies that $\Pr[F \in \mathcal{F}] \leq (1 - 2^{-d-1})^{m^d}$. Hence the probability that H does not satisfy the conclusion in a random coloring of the (d, n) -graph is at most $2(1 - 2^{-d-1})^{m^d}$. Since $\binom{n}{m}^{d+1} \cdot 2(1 - 2^{-d-1})^{m^d} < n^{m(d+1)} \cdot 2 \exp(-2^{d+1}m^d) < 1$ for $m = \lceil 25(\log n)^{1/(d-1)} \rceil$, by the union bound over all (d, m) -subgraphs H it follows that the probability of some H failing the conclusion is < 1 , and so the desired coloring exists. \square

Recall that $X = V_1 * \dots * V_{d+1}$. The $(d-1)$ -simplices of X form a (d, n) -graph. Choose a 2-edge-coloring $\chi: X^{\leq(d-1)} \rightarrow \{1, -1\}$ of this hypergraph as in the preceding lemma.

We shall use the coloring χ to construct a map $\phi: X \rightarrow M$. We will confine the image of ϕ to a single chart of M ; this way we may assume that $M = \mathbb{R}^d$. We start by mapping the vertices of X to points in general position inside the coordinate hyperplane $\{x_d = 0\}$. Extending this embedding linearly, we obtain a map $X^{\leq(d-1)} \rightarrow \mathbb{R}^d$. Then, for each $(d-1)$ -face σ of X we push the interior of σ up or down in the last coordinate direction according to the color $\chi(\sigma)$. The result of this deformation defines the map ϕ on $X^{\leq(d-1)}$.



Finally, we extend the map to the d -skeleton of X in such a way that if $\sigma \in X^{\leq d}$ and if all the $(d-1)$ -simplices in $\partial\sigma$ are above (below) the $\{x_d = 0\}$ hyperplane, then σ is also above (resp. below) that hyperplane.

Let $p \in \mathbb{R}^d$, and suppose that $\phi^{-1}(p)$ intersects all d -faces of $P_1 * \dots * P_{d+1}$ for some $P_i \subset V_i$. Because ϕ maps the vertices of X to points in general position and because ϕ is linear on $X^{\leq(d-2)}$, the intersection of ϕ -images of any $d/2 + 1$ vertex-disjoint simplices in $X^{\leq(d-2)}$ is empty. So, $\phi^{-1}(p)$ is contained in the interior of at most $d/2$ many faces of $X^{\leq(d-2)}$. Let Q be the union of vertex sets of these faces; evidently, $|Q| \leq d^2/2$.

Let $P'_i = P_i \setminus Q$. If $|P'_i| \geq 30(\log n)^{1/(d-1)}$ and n is sufficiently large, then $|P'_i| \geq 25(\log n)^{1/(d-1)}$. If $|P'_i| \geq 25(\log n)^{1/(d-1)}$ for all i , then $P_1 * \dots * P_{d+1}$ contains monochromatic d -faces of each of the two colors. Let $\sigma \in (P_1 * \dots * P_{d+1})^{\leq d}$ be one of these two monochromatic d -faces. Since σ is monochromatic, the only points of $\phi(\sigma)$ that lie on the hyperplane $\{x_d = 0\}$ are those of $\phi(\sigma^{\leq(d-2)})$. By the definition of Q this implies that $p_d \neq 0$, and hence $\chi(\sigma) = \text{sign } p_d$. Since this holds for monochromatic d -faces of both colors, we reach a contradiction. So, $|P'_i| < 25(\log n)^{1/(d-1)}$ for some i , after all.

Topological intermezzo. Before tackling the proof of the lower bound from Theorem 2, we perform a couple of topological simplifications, and review some definitions and results.

Since the lower bound in Theorem 2 is about surfaces, we may (and do) assume that M carries a triangulation. By slight abuse of notation, we use the same symbol M to denote the triangulation. Consider the Boolean algebra generated by all the ϕ -images of simplices in X . Let $\rho > 0$ be the minimal distance between two disjoint sets in this Boolean algebra. We may assume that the triangulation of M is sufficiently fine so that the diameter of each cell is less than ρ . By the simplicial approximation theorem [16] there is a subdivision X' of X and simplicial map $\phi': X' \rightarrow M$ such that the image $\phi'(x)$ belongs to the (uniquely defined) simplex of M whose relative interior contains $\phi(x)$. If we put $A = \{\sigma \in X : p \in \phi'(\sigma)\}$, then $\bigcap_{\sigma \in A} \phi(\sigma)$ is also non-empty. Therefore, it suffices to find a topological Pach family for ϕ' . To avoid extra notation, we continue writing ϕ for ϕ' , with the added benefit of ϕ now being a piecewise linear map.

We say that the map $\phi: X \rightarrow M$ is in *general position with respect to M* if for every simplex $\sigma \in X$ and every simplex $\tau \in M$, whose dimensions satisfy $\dim \sigma + \dim \tau = d$, $\phi(\sigma)$ and τ intersect transversally. By [21, Ch. VI], we may assume (up to small perturbation) that ϕ is in general position.

A *k-chain in a complex Y* (over \mathbb{F}_2) is a formal \mathbb{F}_2 -linear combination of k -faces of Y . The k -chains of Y form an \mathbb{F}_2 -vector space, which we denote $C_k(Y)$. The *boundary* of a k -face $\sigma \in Y$ is the $(k-1)$ -chain $\partial\sigma$ that is the sum of all $(k-1)$ -faces of σ . The boundary of a general k -chain is defined by linearity, $\partial(\sigma_1 + \dots + \sigma_m) \stackrel{\text{def}}{=} \partial\sigma_1 + \dots + \partial\sigma_m$.

Given a map ϕ and a pair of chains $a \in C_{d-k}(X)$ and $b \in C_k(M)$, we can define the intersection number $\phi(a) \cdot b$. If a and b are simplices, let $\phi(a) \cdot b$ be the parity of the number of the intersection points $\phi(a) \cap b$, which is finite since ϕ is in general position. We extend by linearity to the case when a and b are arbitrary chains. The intersection numbers satisfy

$$\phi(\partial a) \cdot b = \phi(a) \cdot \partial b; \quad (1)$$

see [19, Eq. (5) on p. 256]; for a modern proof of a similar formula for the cap product see [11, p. 240 in Section 3.3].

Finally, we need to state Gromov's overlap theorem [10] slightly more precisely. We follow the exposition in [8]. A *k-cochain in X* is a \mathbb{F}_2 -linear function on $C_k(X)$. The vector space of all k -cochains is denoted $C^k(X)$. A *coboundary* of a cochain $a \in C^k(X)$ is the $(k+1)$ -cochain δa , which is given by $\delta a(\sigma) \stackrel{\text{def}}{=} a(\partial\sigma)$. Define the norm on $C^k(X)$ by

$$\|a\| \stackrel{\text{def}}{=} |\text{supp } a|/|X^{=k}|, \quad (2)$$

where $\text{supp } a$ is the support of a . Up to normalization, this is the Hamming norm. Since we work with \mathbb{F}_2 coefficient, we will occasionally abuse the notation and identify cochains with their supports.

A *k-coboundary* is a cochain of the form δa for some $(k-1)$ -cochain a . We say that a *cofills* δa . In general, there are many $(k-1)$ -cochains a that cofill a given coboundary. Cofillings of small norm are of special interest though. We say that X satisfies an *L-cofilling inequality* in dimension k if for every k -coboundary b there is some $a \in C^{k-1}(X)$ such that $\delta a = b$ and $\|a\| \leq L\|b\|$.

We say that X is ε -*sparse* if, for every face $\tau \in X$ and every $0 \leq k \leq d$, less than ε -fraction of all k -simplices intersect τ .

Theorem 6 (Gromov's overlap theorem, following [8]). *Suppose X is a finite simplicial complex of dimension d , and let $\|\cdot\|$ be defined by (2). Suppose $\phi: X \rightarrow M$ is a continuous map into a piecewise linear manifold M . Suppose that*

- *X satisfies an L -cofilling inequality in dimensions $1, \dots, d$; and*
- *X is ε -sparse; and*
- *the cohomology groups of X vanish in dimensions $0, \dots, d-1$.*

Then there is a point $p \in M$ such that

$$\|\{\sigma \in X^{=d} : \phi(\sigma) \cdot p \neq 0\}\| \geq \frac{1}{2(d+1)!L^d} - O(d\varepsilon).$$

(The statement appearing as Theorem 8 of [8] is slightly different: It is limited to compact connected manifolds, the condition on the cohomology groups is replaced by a condition on cosystoles, and the conclusion asserts only that ϕ -images of a positive fraction of the d -simplices contain p . The compactness condition can be removed as explained in Remark 9 of the same paper, the connectedness condition can be removed by restricting to the component containing the image of X , the cosystole condition is implied by our cohomology condition, and the stronger conclusion follows from the actual proof of the overlap theorem in [8].)

For the complex $X = V_1 * \dots * V_{d+1}$ of interest in this paper, an L -cofilling inequality in dimension k with $L = \frac{|X^{=k}|}{|X^{=(k-1)}|} \cdot \frac{2^k-1}{n} \leq 2^k \leq 2^d$ was proved in [7, Proposition 5.8]. An n -point discrete space can be thought of as a wedge sum of $n-1$ copies of \mathbb{S}^0 , and so X is homotopy equivalent to the wedge sum of $(n-1)^{d+1}$ copies of \mathbb{S}^d , which implies that the cohomology groups vanish in all dimensions less than d . Finally, X is clearly $O(1/n)$ -sparse. Therefore, by Theorem 3, for this X and every continuous map $\phi: X \rightarrow M$ we obtain a point $p \in M$ satisfying

$$\|\{\sigma \in X^{=d} : \phi(\sigma) \cdot p \neq 0\}\| \geq \frac{1}{(d+1)!2^{d^2+1}} - O(1/n). \quad (3)$$

Lower bound in the case $M \neq \mathbb{RP}^2$. Let \widetilde{M} be the universal cover of M . Since the complex X is simply-connected, we may lift the map $\phi: X \rightarrow M$ to a map $\tilde{\phi}: X \rightarrow \widetilde{M}$ satisfying $\phi = \text{prj} \circ \tilde{\phi}$, where $\text{prj}: \widetilde{M} \rightarrow M$ is the projection map. Since M is a manifold, so is \widetilde{M} . Observe that a Pach family for the map $\tilde{\phi}$ is also a Pach family for the map ϕ . So, we may replace M by its universal cover \widetilde{M} if we wish.

If M is compact, and is neither \mathbb{S}^2 nor \mathbb{RP}^2 , then it follows from the classification of compact surfaces that its fundamental group is infinite. When $\pi_1(M)$ is infinite, \widetilde{M} is non-compact because it contains a discrete subset of the form $\text{prj}^{-1}(p)$. In this case, we replace M by \widetilde{M} as discussed above. So, if $M \neq \mathbb{RP}^2$, then we may (and do) assume that M is non-compact.

Since X is compact but M is not, the set $M \setminus \phi(X)$ is non-empty. By Theorem 6 and the explicit bound (3), there is a point $p \in M$ such that $\phi(\sigma) \cdot p \neq 0$ for $\frac{1}{192}n^3$ many triangles σ of X . Let \mathcal{F} be the set of these triangles. Let $R \in C_1(M)$ be a path in the 1-skeleton of M from p to a point in

$M \setminus \phi(X)$. For a triangle $\sigma = x_1x_2x_3 \in X$ with $x_i \in V_i$, let σ_i be its edge obtained by removing vertex x_i . Associate to each $\sigma \in \mathcal{F}$ a vector

$$\pi(\sigma) \stackrel{\text{def}}{=} (\phi(\sigma_1) \cdot R, \phi(\sigma_2) \cdot R, \phi(\sigma_3) \cdot R) \in \mathbb{F}_2^3. \quad (4)$$

By the pigeonhole principle, there is $\mathcal{F}' \subset \mathcal{F}$ of size $|\mathcal{F}'| \geq 2^{-3}|\mathcal{F}|$ on which $\pi(\sigma)$ is constant, say $\pi(\sigma) = \tilde{\pi}$. Because of (1), $\sum_i \tilde{\pi}_i = \phi(\sigma) \cdot \partial R = \phi(\sigma) \cdot p + 0 \neq 0$.

Let $H \subset X^{=2}$ be the 3-partite graph whose edge set consists of all the edges of the triangles $\sigma \in \mathcal{F}'$. We say that $\tau \in E(H)$ is of *type* i if τ contains no vertex from V_i in X (and hence contains a vertex from every other V_j in X). By the definition of \mathcal{F}' we have $\phi(\tau) \cdot R = \tilde{\pi}_i$ whenever τ is of type i . Since $\sum \tilde{\pi}_i \neq 0$, it follows that every triangle in H corresponds to a 2-simplex of X whose ϕ -image contains p .

The lower bound in Theorem 2 now follows from the following result of Nikiforov [13] (stated here for the special case that we need). An alternative proof is in [17, Section 2].

Theorem 7 (Main result of [13]). *Let $0 < c < \frac{1}{2}$. Every n -vertex graph with at least cn^3 triangles contains a complete tripartite subgraph with all parts of size $\lfloor c^3 \log n \rfloor$.*

In our case, the graph has $3n$ vertices and at least $2^{-3} \cdot \frac{1}{192}n^3$ triangles, and so we may take $c = 1/(192 \cdot 2^3 \cdot 3^3)$.

Lower bound in the case $M = \mathbb{RP}^2$. In contrast to the case $M \neq \mathbb{RP}^2$, here we reverse the order of the steps: we start by applying Gromov's theorem, and then pass to the universal cover. So, pick $p \in \mathbb{RP}^2$ such that $\phi(\sigma) \cdot p \neq 0$ for $\frac{1}{192}n^3$ many triangles $\sigma \in X^{=2}$. Let $\tilde{\phi}: X \rightarrow \mathbb{S}^2$ be the lift of ϕ , and let $\text{prj}: \mathbb{S}^2 \rightarrow \mathbb{RP}^2$ be the projection map. Denote by q_1, q_2 the preimages of p under prj . Let $R \in C_1(\mathbb{S}^2)$ be a path from q_1 to q_2 .

Because $\tilde{\phi}$ maps the points of $\phi^{-1}(p)$ to $\{q_1, q_2\}$, it is clear that $\phi(\sigma) \cdot p = \tilde{\phi}(\sigma) \cdot (q_1 + q_2)$. Since $\partial R = q_1 + q_2$, it follows that $\phi(\sigma) \cdot p \neq 0$ if and only if $\tilde{\phi}(\partial\sigma) \cdot R \neq 0$. We can then introduce a vector $\pi(\sigma)$, defined similarly to (4) but with ϕ replaced by $\tilde{\phi}$. The rest of the argument proceeds unchanged.

Remarks and open problems

1. The condition that the manifold M has no boundary is not essential. If $\partial M = B$, we can turn M into a manifold without boundary by attaching $B \times [0, 1)$ to M along B .
2. The lower bound in Theorem 3 carries over without any changes to any piecewise linear manifold. However, we believe that the upper bound in Theorem 2 is sharp for every manifold M that is not a sphere. We are unable to prove this even when M (or its universal cover) is non-compact. Following the argument in Theorem 2, this case would follow from the obvious generalization of Nikiforov's result from [13] to hypergraphs. Alas, this is still an open problem, see [17] for partial results.
3. Assuming a suitable hypergraph generalization of Nikiforov's theorem, the lower bound argument in Theorem 2 generalizes to any non-simply-connected piecewise linear manifold M ; one obtains

$f_M(M) \geq c_M(\log n)^{1/(d-1)}$ for such manifolds. Indeed, we may assume that $\pi_1(M)$ is finite, as otherwise we may reduce to the non-compact case. Let ℓ be the smallest prime divisor of $|\pi_1(M)|$. If $\ell = 2$, we consider the universal cover of M , and proceed as in the case $M = \mathbb{RP}^2$ using 1-chain R which is a sum of $|\pi_1(M)|/2$ paths that form a perfect matching on the preimages of a heavily-covered point. If $\ell > 2$, we need to make minor alterations to the proof. First, we use the Gromov’s overlap theorem with \mathbb{F}_ℓ coefficients in place of \mathbb{F}_2 coefficients; as observed in [8, Remark 18.2] Gromov’s theorem holds under the condition that the manifold is \mathbb{F}_ℓ -orientable. Since $\ell > 2$, the fundamental group of the manifold contains no subgroup of index 2, and so M is orientable [11, Proposition 3.25]. To apply the overlap theorem we must verify the cofilling inequality of X with \mathbb{F}_ℓ -coefficients. Fortunately the proof in [7, Proposition 5.8] remains valid for coefficients in any abelian group without any modification (notice that there is a small typo on page 13 of [7], in the formula for ζ_v , the first row on the left hand side should read $\eta_{v_0,v} + d\alpha_{v_0,v}$, instead of $\eta_{v_0,v} + \alpha_{v_0,v}$). So we let p be a heavily-covered (mod ℓ) point, and we let q_1, \dots, q_N , where $N = |\pi_1(M)|$, be its preimages in the universal cover. Note that $\ell \mid N$ implies that whenever $\phi(\sigma) \cdot p \neq 0 \pmod{\ell}$ then there is a pair of preimages q_i, q_j such that $\tilde{\phi}(\sigma) \cdot (q_i + q_j) \neq 0 \pmod{\ell}$. Therefore, we may take R to be a path connecting the most popular pair of preimages, and finish as in the proof of Theorem 2.

4. The case of a general compact $M \neq \mathbb{S}^d$ will require a new argument. For example, we do not know how to prove the lower bound of $c(\log n)^{1/3}$ for $M = \mathbb{S}^2 \times \mathbb{S}^2$ even assuming the generalization of Nikiforov’s result to hypergraphs.
5. We thank Joseph Briggs for valuable feedback on a previous version of this paper.

References

- [1] József Balogh, Robert Morris, and Wojciech Samotij. Independent sets in hypergraphs. *J. Amer. Math. Soc.*, 28(3):669–709, 2015. [arXiv:1204.6530](#).
- [2] I. Bárány, S. B. Shlosman, and A. Szűcs. On a topological generalization of a theorem of Tverberg. *J. London Math. Soc. (2)*, 23(1):158–164, 1981.
- [3] I. Bárány and P. Valtr. A positive fraction Erdős-Szekeres theorem. *Discrete Comput. Geom.*, 19(3, Special Issue):335–342, 1998. Dedicated to the memory of Paul Erdős.
- [4] Imre Bárány. A generalization of Carathéodory’s theorem. *Discrete Math.*, 40(2-3):141–152, 1982.
- [5] Imre Bárány, Roy Meshulam, Eran Nevo, and Martin Tancer. Pach’s selection theorem does not admit a topological extension. *Discrete Comput. Geom.*, 60(2):420–429, 2018. [arXiv:1610.05053](#).
- [6] Béla Bollobás and Andrew Thomason. Projections of bodies and hereditary properties of hypergraphs. *Bull. London Math. Soc.*, 27(5):417–424, 1995.
- [7] Dominic Dotterer and Matthew Kahle. Coboundary expanders. *J. Topol. Anal.*, 4(4):499–514, 2012. [arXiv:1012.5316](#).

- [8] Dominic Dotterrer, Tali Kaufman, and Uli Wagner. On expansion and topological overlap. In Sándor Fekete and Anna Lubiw, editors, *32nd International Symposium on Computational Geometry (SoCG 2016)*, volume 51 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 35:1–35:10, Dagstuhl, Germany, 2016. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik. [arXiv:1506.04558](https://arxiv.org/abs/1506.04558).
- [9] P. Erdős. On extremal problems of graphs and generalized graphs. *Israel J. Math.*, 2:183–190, 1964. https://www.renyi.hu/~p_erdos/1964-13.pdf.
- [10] Mikhail Gromov. Singularities, expanders and topology of maps. Part 2: From combinatorics to topology via algebraic isoperimetry. *Geom. Funct. Anal.*, 20(2):416–526, 2010.
- [11] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002. Available from author’s website at <https://www.math.cornell.edu/~hatcher/AT/AT.pdf>.
- [12] L. H. Loomis and H. Whitney. An inequality related to the isoperimetric inequality. *Bull. Amer. Math. Soc.*, 55:961–962, 1949.
- [13] Vladimir Nikiforov. Graphs with many r -cliques have large complete r -partite subgraphs. *Bull. Lond. Math. Soc.*, 40(1):23–25, 2008. [arXiv:math/0703554](https://arxiv.org/abs/math/0703554).
- [14] Vladimir Nikiforov. Complete r -partite subgraphs of dense r -graphs. *Discrete Math.*, 309(13):4326–4331, 2009.
- [15] János Pach. A Tverberg-type result on multicolored simplices. *Comput. Geom.*, 10(2):71–76, 1998. [arXiv:math/9603211](https://arxiv.org/abs/math/9603211).
- [16] V. V. Prasolov. *Elements of combinatorial and differential topology*, volume 74 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2006. Translated from the 2004 Russian original by Olga Sipacheva. Russian original is at <https://www.mccme.ru/free-books/prasolov/topol.pdf>.
- [17] Vojtěch Rödl and Mathias Schacht. Complete partite subgraphs in dense hypergraphs. *Random Structures Algorithms*, 41(4):557–573, 2012.
- [18] David Saxton and Andrew Thomason. Hypergraph containers. *Invent. Math.*, 201(3):925–992, 2015. [arXiv:1204.6595](https://arxiv.org/abs/1204.6595).
- [19] Herbert Seifert and William Threlfall. *Seifert and Threlfall: a textbook of topology*, volume 89 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980. Translated from the German edition of 1934 by Michael A. Goldman, With a preface by Joan S. Birman, With “Topology of 3-dimensional fibered spaces” by Seifert, Translated from the German by Wolfgang Heil.
- [20] H. Tverberg. A generalization of Radon’s theorem. *J. London Math. Soc.*, 41:123–128, 1966.
- [21] Eric-Christopher Zeeman. *Seminar on combinatorial topology*. Institut des Hautes Études Scientifiques, 1963. Lecture notes. Available at <http://www.maths.ed.ac.uk/~aar/papers/zeemanpl.pdf>.