arXiv:1708.04422v3 [math.PR] 18 Oct 2018

# Supercritical Superprocesses: Proper Normalization and Non-degenerate Strong Limit

**Yan-Xia Ren*** **Renming Song†** **and** **Rui Zhang‡**

**Abstract**

Suppose that $X = \{X_t, t \geq 0; \mathbb{P}_\mu\}$ is a supercritical superprocess in a locally compact separable metric space $E$. Let $\phi_0$ be a positive eigenfunction corresponding to the first eigenvalue $\lambda_0$ of the generator of the mean semigroup of $X$. Then $M_t := e^{-\lambda_0 t}\langle \phi_0, X_t\rangle$ is a positive martingale. Let $M_\infty$ be the limit of $M_t$. It is known (see, J. Appl. Probab. **46** (2009), 479–496) that $M_\infty$ is non-degenerate iff the $L\log L$ condition is satisfied. In this paper we are mainly interested in the case when the $L\log L$ condition is not satisfied. We prove that, under some conditions, there exist a positive function $\gamma_t$ on $[0,\infty)$ and a non-degenerate random variable $W$ such that for any finite nonzero Borel measure $\mu$ on $E$,

$$\lim_{t\to\infty} \gamma_t\langle \phi_0, X_t\rangle = W, \qquad \text{a.s.-}\mathbb{P}_\mu.$$

We also give the almost sure limit of $\gamma_t\langle f, X_t\rangle$ for a class of general test functions $f$.

**AMS 2010 Mathematics Subject Classification:** 60J68, 60F15.

**Keywords and Phrases**:superprocesses, Seneta-Heyde norming, non-degenerate strong limit, martingales

# 1 Introduction

Suppose $\{Z_n, n \geq 0\}$ is a supercritical Galton-Watson process with offspring number $L$. Let $m := EL \in (1,\infty)$ be the mean of $L$. Then $M_n := \frac{Z_n}{m^n}$ is a non-negative martingale and thus has a finite limit $M_\infty$. The well-known Kesten-Stigum theorem says that the following three statements are equivalent: (i) $P(M_\infty = 0) = P(Z_n = 0 \text{ for large } n)$, i.e., the events $\{M_\infty = 0\}$ and $\{Z_n = 0 \text{ for large } n\}$ are almost the same; (ii) $EM_\infty = 1$; (iii) $E(L\log L) < \infty$. For a classical proof of this result, the reader is referred to the book [1]

*The research of this author is supported by NSFC (Grant No. 11671017 and 11731009).

†Research supported in part by a grant from the Simons Foundation (#429343, Renming Song).

‡Research supported in part by NSFC (Grant No. 11601354) and Youth Innovative Research Team of Capital Normal University. Corresponding author.

of Athreya and Ney. In 1995, Lyons, Pemantle and Peres [18] gave a probabilistic proof of the above $L\log L$ criterion of Kesten and Stigum. If $E(L\log L)=\infty$, then $\lim_{n\to\infty}\frac{Z_n}{m^n}=0$ almost surely, which says that $m^n$ does not give the right growth rate of $Z_n$ conditional on non-extinction. It is natural to ask what is the right growth rate of $Z_n$. In 1968, Seneta [26] proved that there is a sequence of positive numbers $c_n$ such that $c_nZ_n$ converges to a non-degenerate random variable $W$ in distribution. Heyde [10] strengthened the convergence in distribution to almost sure convergence. Later the problem of finding $c_n$ such that $c_nZ_n$ converges to a non-degenerate limit is called the Seneta-Heyde norming problem.

Hoppe [11] generalized the result of Heyde [10] to supercritical multitype branching processes, Grey [8] proved a similar result for continuous state branching processes and Hering [9] obtained a similar result for supercritical branching diffusions. In this paper we are going to consider the Seneta-Heyde norming problem for general superprocesses under some conditions which are easy to check and satisfied by many superprocesses, including superdiffusions in a bounded domain and also superprocesses with discontinuous spatial motions. We emphasize that we are mainly interested in the case when the $L\log L$ condition fails, since the norming problem is already solved in [16] for superdiffusions and [20] for more general superprocesses when the $L\log L$ condition holds.

## 1.1 Superprocesses and assumptions

In this subsection, we describe the setup of this paper and formulate our assumptions.

Suppose that $E$ is a locally compact separable metric space. We will use $\mathcal{B}(E)$ ($\mathcal{B}^+(E)$) to denote the family of (non-negative) Borel functions on $E$, $\mathcal{B}_b(E)$ ($\mathcal{B}_b^+(E)$) to denote the family of (non-negative) bounded Borel functions on $E$, and $C(E)(C_0(E)$, respectively) to denote the family of continuous functions (vanishing at infinity, respectively) on $E$.

Suppose that $\partial$ is a separate point not contained in $E$. We will use $E_\partial$ to denote $E\cup\{\partial\}$. Every function $f$ on $E$ is automatically extended to $E_\partial$ by setting $f(\partial)=0$. We will assume that $\xi=\{\xi_t,\Pi_x\}$ is a Hunt process on $E$ and $\zeta:=\inf\{t>0:\xi_t=\partial\}$ is the lifetime of $\xi$. We will use $\{P_t:t\geq 0\}$ to denote the semigroup of $\xi$. Suppose that $m$ is a $\sigma$-finite Borel measure on $E$ with full support. We will assume below that $\{P_t:t\geq 0\}$ has a dual with respect to the measure $m$ and the dual semigroup is sub-Markovian.

The superprocess $X=\{X_t:t\geq 0\}$ we are going to work with is determined by two parameters: a spatial motion $\xi=\{\xi_t,\Pi_x\}$ on $E$ which is a Hunt process, and a branching mechanism $\varphi$ of the form

$$\varphi(x,s)=-\alpha(x)s+\beta(x)s^2+\int_{(0,+\infty)}(e^{-s\theta}-1+s\theta)n(x,\mathrm{d}\theta),\quad x\in E,\ s\geq 0, \tag{1.1}$$

where $\alpha \in \mathcal{B}_b(E)$, $\beta \in \mathcal{B}_b^+(E)$ and $n$ is a kernel from $E$ to $(0,\infty)$ satisfying

$$\sup_{x\in E}\int_{(0,+\infty)}(\theta\wedge\theta^2)n(x,\mathrm{d}\theta)<\infty. \tag{1.2}$$

Then there exists $M>0$ such that

$$|\alpha(x)|+\beta(x)+\int_{(0,+\infty)}(\theta\wedge\theta^2)n(x,\mathrm{d}\theta)\le M.$$

In this paper, we will exclude the case when $\beta(\cdot)+n(\cdot,(0,\infty))=0$ $m$-almost everywhere.

Let $\mathcal{M}_F(E)$ be the space of finite measures on $E$, equipped with the topology of weak convergence. The superprocess $X$ with spatial motion $\xi$ and branching mechanism $\varphi$ is a Markov process taking values in $\mathcal{M}_F(E)$. The existence of such superprocesses is well-known, see, for instance, [5], [6] or [15]. For any $\mu\in\mathcal{M}_F(E)$, we denote the law of $X$ with initial configuration $\mu$ by $\mathbb{P}_\mu$. As usual, $\langle f,\mu\rangle := \int_E f(x)\mu(dx)$ and $\|\mu\| := \langle 1,\mu\rangle$. Throughout this paper, a real-valued function $u(t,x)$ on $[0,\infty)\times E_\partial$ is said to be locally bounded if, for any $t>0$, $\sup_{s\in[0,t],x\in E_\partial}|u(s,x)|<\infty$. According to [15, Theorem 5.12], there is a Hunt process $X=\{\Omega,\mathcal{G},\mathcal{G}_t,X_t,\mathbb{P}_\mu\}$ taking values in $\mathcal{M}_F(E)$ such that for every $f\in\mathcal{B}_b^+(E)$ and $\mu\in\mathcal{M}_F(E)$,

$$-\log\mathbb{P}_\mu\left(e^{-\langle f,X_t\rangle}\right)=\langle V_tf,\mu\rangle, \tag{1.3}$$

where $V_tf(x)$ is the unique locally bounded non-negative solution to the equation

$$V_tf(x)+\Pi_x\int_0^t\varphi(\xi_s,V_{t-s}f(\xi_s))\mathrm{d}s=\Pi_xf(\xi_t),\quad x\in E_\partial, \tag{1.4}$$

where we use the convention that $\varphi(\partial,s)=0$ for all $s\ge0$. Since $f(\partial)=0$, we have $V_tf(\partial)=0$ for any $t\ge0$. In this paper, the superprocess we deal with is always this Hunt realization.

For any $f\in\mathcal{B}_b(E)$ and $(t,x)\in(0,\infty)\times E$, we define

$$T_tf(x):=\Pi_x\left[e^{\int_0^t\alpha(\xi_s)\mathrm{d}s}f(\xi_t)\right]. \tag{1.5}$$

It is well known that $T_tf(x)=\mathbb{P}_{\delta_x}\langle f,X_t\rangle$ for every $x\in E$.

We will always assume that there exists a family of continuous and strictly positive functions $\{p(t,x,y):t>0\}$ on $E\times E$ such that for any $t>0$ and nonnegative function $f$ on $E$,

$$P_tf(x)=\int_E p(t,x,y)f(y)m(\mathrm{d}y).$$

Define

$$a_t(x):=\int_E p(t,x,y)^2\,m(\mathrm{d}y),\qquad \hat{a}_t(x):=\int_E p(t,y,x)^2\,m(\mathrm{d}y).$$

Our first main assumption is

**Assumption 1** **(i)** *For any $t > 0$, $\int_E p(t,x,y)\, m(\mathrm{d}x) \le 1$.*

**(ii)** *For any $t > 0$, we have*

$$\int_E a_t(x)\, m(\mathrm{d}x) = \int_E \hat{a}_t(x)\, m(\mathrm{d}x) = \int_E \int_E p(t,x,y)^2\, m(\mathrm{d}y)\, m(\mathrm{d}x) < \infty. \tag{1.6}$$

*Moreover, the functions $x \to a_t(x)$ and $x \to \hat{a}_t(x)$ are continuous on $E$.*

Note that, in Assumption 1(i), the integration is with respect to the first space variable. It implies that the dual semigroup $\{\widehat{P}_t : t \ge 0\}$ of $\{P_t : t \ge 0\}$ with respect to $m$ defined by

$$\widehat{P}_t f(x) = \int_E p(t,y,x) f(y) m(\mathrm{d}y)$$

is sub-Markovian. Assumption 1(ii) is a pretty weak $L^2$ condition and it allows us to apply results on operator semigroups in Hilbert spaces. By Hölder's inequality, we have

$$p(t+s,x,y) = \int_E p(t,x,z) p(s,z,y)\, m(\mathrm{d}z) \le (a_t(x))^{1/2} (\hat{a}_s(y))^{1/2}. \tag{1.7}$$

It is well known and easy to check that $\{P_t : t \ge 0\}$ and $\{\widehat{P}_t : t \ge 0\}$ are strongly continuous contraction semigroups on $L^2(E,m)$, see [22] for a proof. We will use $\langle \cdot, \cdot \rangle_m$ to denote the inner product in $L^2(E,m)$. Since $p(t,x,y)$ is continuous in $(x,y)$, by (1.7), Assumption 1(ii) and the dominated convergence theorem, we have that, for any $f \in L^2(E,m)$, $P_t f$ and $\widehat{P}_t f$ are continuous.

It follows from Assumption 1(ii) that, for each $t > 0$, $P_t$ and $\widehat{P}_t$ are compact operators on $L^2(E,m)$. Let $\widetilde{L}$ and $\widehat{\widetilde{L}}$ be the infinitesimal generators of the semigroups $\{P_t\}$ and $\{\widehat{P}_t\}$ in $L^2(E,m)$ respectively. Define $\widetilde{\lambda}_0 := \sup \Re(\sigma(\widetilde{L})) = \sup \Re(\sigma(\widehat{\widetilde{L}}))$. By Jentzsch's theorem (Theorem V.6.6 on page 337 of [25]), $\widetilde{\lambda}_0$ is an eigenvalue of multiplicity 1 for both $\widetilde{L}$ and $\widehat{\widetilde{L}}$, and that an eigenfunction $\widetilde{\phi}_0$ of $\widetilde{L}$ corresponding to $\widetilde{\lambda}_0$ can be chosen to be strictly positive $m$-almost everywhere with $\|\widetilde{\phi}_0\|_2 = 1$ and an eigenfunction $\widetilde{\psi}_0$ of $\widehat{\widetilde{L}}$ corresponding to $\widetilde{\lambda}_0$ can be chosen to be strictly positive $m$-almost everywhere with $\langle \widetilde{\phi}_0, \widetilde{\psi}_0 \rangle_m = 1$. Thus for $m$-almost every $x \in E$,

$$e^{\widetilde{\lambda}_0} \widetilde{\phi}_0(x) = P_1 \widetilde{\phi}_0(x), \qquad e^{\widetilde{\lambda}_0} \widetilde{\psi}_0(x) = \widehat{P}_1 \widetilde{\psi}_0(x).$$

Hence $\widetilde{\phi}_0$ and $\widetilde{\psi}_0$ can be chosen to be continuous and strictly positive everywhere on $E$.

Our second assumption is

**Assumption 2** **(i)** *$\widetilde{\phi}_0$ is bounded.*

**(ii)** *The semigroup $\{P_t, t \ge 0\}$ is intrinsically ultracontractive, that is, there exists $c_t > 0$ such that*

$$p(t,x,y) \le c_t \widetilde{\phi}_0(x) \widetilde{\psi}_0(y). \tag{1.8}$$

Assumption 2 is a pretty strong assumption on the semigroup $\{P_t : t \geq 0\}$. However, this assumption is satisfied in a lot of cases. See [21, Section 1.4] for examples of Markov processes satisfying the assumption above. The concept of intrinsic ultracontractivity was introduced by Davies and Simon in [4] in the setting of symmetric semigroups. This concept was extended to the non-symmetric setting in [12, 13, 14]. Intrinsic ultracontractivity has been studied intensively in the last 30 years and there are many results on the intrinsic ultracontractivity of semigroups, see [12, 13, 14] and the references therein.

We have proved in [21, Lemma 2.1] that there exists a function $q(t, x, y)$ on $(0, \infty) \times E \times E$ which is continuous in $(x, y)$ for each $t > 0$ such that

$$e^{-Mt} p(t, x, y) \leq q(t, x, y) \leq e^{Mt} p(t, x, y), \quad (t, x, y) \in (0, \infty) \times E \times E, \tag{1.9}$$

and that for any bounded Borel function $f$ and any $(t, x) \in (0, \infty) \times E$,

$$T_t f(x) = \int_E q(t, x, y) f(y)\, m(\mathrm{d}y).$$

It follows immediately that

$$\|T_t f\|_2 \leq e^{Mt} \|P_t f\|_2 \leq e^{Mt} \|f\|_2. \tag{1.10}$$

In [22], we have proved that $\{T_t : t \geq 0\}$ is a strongly continuous semigroup on $L^2(E, m)$. Let $\{\widehat{T}_t, t > 0\}$ be the adjoint semigroup on $L^2(E, m)$ of $\{T_t, t > 0\}$, that is, for $f \in L^2(E, m)$,

$$\widehat{T}_t f(x) = \int_E q(t, y, x) f(y)\, m(\mathrm{d}y).$$

We have proved in [22] that $\{\widehat{T}_t : t \geq 0\}$ is also a strongly continuous semigroup on $L^2(E, m)$. We claim that, for all $t > 0$ and $f \in L^2(E, m)$, $T_t f$ and $\widehat{T}_t f$ are continuous. In fact, since $q(t, x, y)$ is continuous in $(x, y)$, by (1.7), (1.9), Assumption 1(ii) and the dominated convergence theorem, we have that, for any $f \in L^2(E, m)$, $T_t f$ and $\widehat{T}_t f$ are continuous.

By Assumption 1(ii) and (1.9), we get that

$$\int_E \int_E q^2(t, x, y)\, m(\mathrm{d}x)\, m(\mathrm{d}y) \leq e^{2Mt} \int_E \int_E p^2(t, x, y)\, m(\mathrm{d}x)\, m(\mathrm{d}y) < \infty.$$

Thus, for each $t > 0$, $T_t$ and $\widehat{T}_t$ are compact operators on $L^2(E, m)$. Let $L$ and $\widehat{L}$ be the infinitesimal generators of the semigroups $\{T_t\}$ and $\{\widehat{T}_t\}$ in $L^2(E, m)$ respectively. Define $\lambda_0 := \sup \Re(\sigma(L)) = \sup \Re(\sigma(\widehat{L}))$. By Jentzsch's theorem, $\lambda_0$ is an eigenvalue of multiplicity 1 for both $L$ and $\widehat{L}$, and that an eigenfunction $\phi_0$ of $L$ corresponding to $\lambda_0$ can be chosen to be strictly positive $m$-almost everywhere with $\|\phi_0\|_2 = 1$ and an eigenfunction $\psi_0$ of $\widehat{L}$ corresponding to $\lambda_0$ can be chosen to be strictly positive $m$-almost everywhere with $\langle \phi_0, \psi_0 \rangle_m = 1$. Thus for $m$-almost every $x \in E$,

$$e^{\lambda_0} \phi_0(x) = T_1 \phi_0(x), \qquad e^{\lambda_0} \psi_0(x) = \widehat{T}_1 \psi_0(x).$$

Hence $\psi_0$ and $\phi_0$ can be chosen to be continuous and strictly positive everywhere on $E$.

Using Assumption 2, the boundedness of $\alpha$ and an argument similar to that used in the proof of [4, Theorem 3.4], one can show the following:

**(i)** $\phi_0$ is bounded.

**(ii)** The semigroup $\{T_t, t \geq 0\}$ is intrinsically ultracontractive, that is, there exists $c_t > 0$ such that

$$q(t, x, y) \leq c_t \phi_0(x) \psi_0(y). \tag{1.11}$$

The main interest of this paper is on supercritical superprocesses, so we assume that

**Assumption 3** $\lambda_0 > 0$.

Define $q_t(x) := \mathbb{P}_{\delta_x}(\|X_t\| = 0)$. Since $\mathbb{P}_{\delta_x}\|X_t\| = T_t 1(x) > 0$, we have $q_t(x) < 1$. Note that $q_t(x)$ is non-decreasing in $t$. Hence the limit

$$q(x) := \lim_{t \to \infty} q_t(x)$$

exists. Then $q(x) = \mathbb{P}_{\delta_x}\{\|X_t\| = 0 \text{ for some } t > 0\}$ is the extinction probability. In this paper, we also assume that

**Assumption 4** *There exists* $t_0 > 0$ *such that*

$$\inf_{x \in E} q_{t_0}(x) > 0. \tag{1.12}$$

In [21, Section 2.2], we gave a sufficient condition (in term of the branching mechnism $\varphi$) for Assumption 4. In particularly, if $\inf_{x \in E} \beta(x) > 0$, then Assumption 4 holds. In Lemma 3.1, we will show that, under our assumptions, $q(x) < 1$, for all $x \in E$.

## 1.2 Main results

Define

$$M_t := e^{-\lambda_0 t} \langle \phi_0, X_t \rangle, \quad t \geq 0.$$

It is easy to prove that (see [20, Theorem 3.2], for example), for every $\mu \in \mathcal{M}_F(E)$, $\{M_t, t \geq 0\}$ is a non-negative $\mathbb{P}_\mu$-martingale with respect to the filtration $\{\mathcal{G}_t : t \geq 0\}$. Thus $\{M_t, t \geq 0\}$ has a $\mathbb{P}_\mu$-a.s. finite limit denoted as $M_\infty$.

Let $n^{\phi_0}(x, \mathrm{d}\theta)$ be the kernel from $E$ to $(0, \infty)$ defined by

$$\int_0^\infty f(\theta) n^{\phi_0}(x, \mathrm{d}\theta) = \int_0^\infty f(\theta \phi_0(x)) n(x, \mathrm{d}\theta).$$

By the boundedness of $\phi_0$ and the assumption (1.2), we get that there exists $\widetilde{M} > 0$ such that

$$\sup_{x\in E}\int_0^\infty (\theta\wedge\theta^2)n^{\phi_0}(x,\mathrm{d}\theta)\le \widetilde{M}. \tag{1.13}$$

Let $l(x) := \int_1^\infty \theta\log\theta\, n^{\phi_0}(x,\mathrm{d}\theta)$. The following $L\log L$ criterion was proved for superdiffusions in [16].

*$L\log L$* ***criterion:*** *$M_\infty$ is non-degenerate under $\mathbb{P}_\mu$ for all nonzero finite measures $\mu$ on $E$ if and only if*

$$\int_E \psi_0(x)l(x)m(\mathrm{d}x) < \infty. \tag{1.14}$$

At first glance, the roles of $\phi_0$ and $\psi_0$ are not symmetric in (1.14). This is not the case. In fact, (1.14) is equivalent to

$$\int_E \phi_0(x)\psi_0(x)m(\mathrm{d}x)\int_1^\infty (\theta\log\theta)n(x,\mathrm{d}\theta) < \infty, \tag{1.15}$$

which says that the spatial average of the "$\theta\log\theta$" moment $\int_1^\infty(\theta\log\theta)n(x,\mathrm{d}\theta)$ with respect to the probability measure $\phi_0(x)\psi_0(x)m(\mathrm{d}x)$ is finite. Note that

$$\begin{aligned} l(x) &= \int_1^\infty \theta\log\theta\, n^{\phi_0}(x,\mathrm{d}\theta) = \phi_0(x)\int_{\phi_0(x)^{-1}}^\infty \theta(\log\theta+\log\phi_0(x))\, n(x,\mathrm{d}\theta)\\ &= \phi_0(x)\int_1^\infty \theta\log\theta\, n(x,\mathrm{d}\theta) + \phi_0(x)\int_{\phi_0(x)^{-1}}^1 \theta\log\theta\, n(x,\mathrm{d}\theta)\\ &\quad + \phi_0(x)\log\phi_0(x)\int_{\phi_0(x)^{-1}}^\infty \theta\, n(x,\mathrm{d}\theta). \end{aligned}$$

Since $\phi_0$ is bounded, $\phi_0(x)|\log\phi_0(x)|$ is bounded above, say by $C$. Thus,

$$\sup_{x\in E}\phi_0(x)|\log\phi_0(x)|\int_{\phi_0(x)^{-1}}^\infty \theta\, n(x,\mathrm{d}\theta) \le C\sup_{x\in E}\int_{\|\phi_0\|_\infty^{-1}}^\infty \theta\, n(x,\mathrm{d}\theta) < \infty.$$

Note also that

$$\begin{aligned} &\sup_{x\in E}\phi_0(x)\left|\int_{\phi_0(x)^{-1}}^1 \theta\log\theta\, n(x,\mathrm{d}\theta)\right| \le \sup_{x\in E}\phi_0(x)|\log\phi_0(x)|\left|\int_{\phi_0(x)^{-1}}^1 \theta\, n(x,\mathrm{d}\theta)\right|\\ &\le C\sup_{x\in E}\left(\mathbf{1}_{\phi_0(x)>1}\int_{\|\phi_0\|_\infty^{-1}}^1 \theta\, n(x,\mathrm{d}\theta) + \mathbf{1}_{\phi_0(x)\le 1}\int_1^\infty \theta\, n(x,\mathrm{d}\theta)\right) < \infty. \end{aligned}$$

In Section 2, we will show that $\psi_0 \in L^1(E,m)$ (see the first paragraph of Section 2). Thus (1.14) is equivalent to (1.15).

Recently, the $L\log L$ criterion above was extended to more general superprocesses with possible non-local branching mechanisms in [20].

The $L\log L$ criterion above says that, under condition (1.14), $e^{\lambda_0 t}$ gives the growth rate of $\langle\phi_0, X_t\rangle$ as $t\to\infty$ conditioned on non-extinction. However, when condition (1.14) is not satisfied, the theorem above does not provide much information about the growth rate of $\langle\phi_0, X_t\rangle$.

The first objective of this paper is to solve the Seneta-Heyde norming problem for the martingale $M_t$, that is, to find a positive function $\gamma_t$ on $[0,\infty)$ such that $\gamma_t\langle\phi_0, X_t\rangle$ has a non-degenerate limit as $t\to\infty$. Although our results (Theorems 1.1, 6.8 and 6.10) also cover the case when (1.14) holds, only the results in the case when (1.14) fails are new, see Theorem 1.2 below. It is easy to find examples such that (1.14) fails. For example, if $n(x,\mathrm{d}\theta) = c(x)[I_{(0,2)}(\theta) + I_{[2,\infty)}(\theta)\theta^{-2}(\log\theta)^{-\beta}]\mathrm{d}\theta$ with $\beta\in(1,2]$ and $c(x)$ being strictly positive bounded measurable function on $E$, then (1.2) is satisfied, but (1.15), which is equivalent to (1.14), fails.

Let $v(x) := -\log q(x)$. By the branching property of $X$, we have

$$\mathbb{P}_\mu(\|X_t\| = 0 \text{ for some } t>0) = e^{-\langle v,\mu\rangle}.$$

**Theorem 1.1** *There exist a positive function $\gamma_t$ on $[0,\infty)$ and a non-degenerate random variable $W$ such that*

$$\lim_{t\to\infty}\frac{\gamma_t}{\gamma_{t+s}} = e^{\lambda_0 s},\quad \forall s\ge 0,$$

*and that for any nonzero $\mu\in\mathcal{M}_F(E)$,*

$$\lim_{t\to\infty}\gamma_t\langle\phi_0, X_t\rangle = W,\qquad a.s.\text{-}\mathbb{P}_\mu$$

*and*

$$\mathbb{P}_\mu(W=0) = e^{-\langle v,\mu\rangle},\qquad \mathbb{P}_\mu(W<\infty) = 1.$$

Moreover, we have the following $L\log L$ criterion.

**Theorem 1.2** *The following conditions are equivalent:*

**(1)** *$M_\infty$ is non-degenerate for some nonzero $\mu\in\mathcal{M}_F(E)$;*

**(2)** *$M_\infty$ is non-degenerate for all nonzero $\mu\in\mathcal{M}_F(E)$;*

**(3)** *$l_0 := \lim_{t\to\infty} e^{\lambda_0 t}\gamma_t \in (0,\infty)$;*

**(4)** *$\int_E \psi_0(x)l(x)m(\mathrm{d}x) < \infty$;*

**(5)** *$\mathbb{P}_\mu W<\infty$ for some nonzero $\mu\in\mathcal{M}_F(E)$.*

**(6)** *$\mathbb{P}_\mu W<\infty$ for all $\mu\in\mathcal{M}_F(E)$.*

Further properties of the limit random variable $W$, such as absolute continuity and tail probabilities, are studied in [23], a sequel to the present paper.

The second objective of this paper is to study the almost sure limit behaviour of $\gamma_t\langle f, X_t\rangle$ as $t \to \infty$ for a class of bounded continuous functions $f$. It turns out that, for $f$ belonging to this class, $\lim_{t\to\infty} \gamma_t\langle f, X_t\rangle = \langle f, \psi_0\rangle_m W$, $\mathbb{P}_\mu$-a.s. for any nonzero $\mu \in \mathcal{M}_F(E)$, see Theorems 6.8 and 6.10.

The rest of the paper is organized as follows. Section 2 contains our basic estimates and Section 3 deals with some properties of the extinction probability. In Section 4, we will define and investigate backward iterates, which is needed in the proof of Theorem 1.1. The proofs of Theorem 1.1 and Theorem 1.2 are given in Section 5. We remark that we will prove Theorem 1.2 without using the $L \log L$ criterion in [16]. The strong limit behaviour of $\gamma_t\langle f, X_t\rangle$ as $t \to \infty$ for a class of general bounded continuous functions $f$ is given in Section 6. In the last section, we give some concluding remarks.

In the remainder of this paper, $C$ will stand for a constant whose value might change from one appearance to the next.

# 2 Some estimates

According to [12, Thorem 2.7], under Assumptions 1–2, for any $\delta > 0$, there exist constants $\gamma = \gamma(\delta) > 0$ and $c = c(\delta) > 0$ such that, for any $(t, x, y) \in [\delta, \infty) \times E \times E$, we have

$$\left|e^{-\lambda_0 t} q(t, x, y) - \phi_0(x)\psi_0(y)\right| \le ce^{-\gamma t}\phi_0(x)\psi_0(y). \tag{2.1}$$

Take $t$ large enough so that $ce^{-\gamma t} < \frac{1}{2}$, then we have

$$e^{-\lambda_0 t} q(t, x, y) \ge \frac{1}{2}\phi_0(x)\psi_0(y).$$

Since $q(t, x, \cdot) \in L^1(E, m)$, we have $\psi_0 \in L^1(E, m)$.

It follows from (2.1) that, if $f \in \mathcal{B}_b^+(E)$ then $\langle f, \psi_0\rangle_m < \infty$ and for any $(t, x) \in [\delta, \infty) \times E$,

$$\left|e^{-\lambda_0 t} T_t f(x) - \langle f, \psi_0\rangle_m \phi_0(x)\right| \le ce^{-\gamma t}\langle |f|, \psi_0\rangle_m \phi_0(x) \tag{2.2}$$

and

$$(1 - ce^{-\gamma t})\langle |f|, \psi_0\rangle_m \phi_0(x) \le e^{-\lambda_0 t} T_t|f|(x) \le (1 + c)\langle |f|, \psi_0\rangle_m \phi_0(x). \tag{2.3}$$

For $x \in E$ and $s > 0$, we define

$$r(x, s) = \varphi(x, s) + \alpha(x)s.$$

**Lemma 2.1** *For any $H \ge 1$,*

$$0 \le r(x, s) \le (3/2 + H/2)Ms^2 + \int_H^\infty \theta\, n(x, \mathrm{d}\theta)s. \tag{2.4}$$

**Proof:** By definition,

$$r(x,s) = \beta(x)s^2 + \int_0^\infty (e^{-\theta s} - 1 + \theta s)\, n(x, \mathrm{d}\theta). \tag{2.5}$$

Note that for any $\theta > 0$,

$$0 < e^{-\theta} - 1 + \theta \le \left(\frac{\theta}{2} \wedge 1\right) \theta. \tag{2.6}$$

Thus for any $H \ge 1$,

$$\begin{aligned} r(x,s) &\le Ms^2 + \frac{1}{2}s^2 \int_0^H \theta^2\, n(x, \mathrm{d}\theta) + s \int_H^\infty \theta\, n(x, \mathrm{d}\theta) \\ &\le Ms^2 + \frac{1}{2}s^2 \left[\int_0^1 \theta^2\, n(x, \mathrm{d}\theta) + H \int_1^H \theta\, n(x, \mathrm{d}\theta)\right] + s \int_H^\infty \theta\, n(x, \mathrm{d}\theta) \\ &\le (3/2 + H/2)Ms^2 + \int_H^\infty \theta\, n(x, \mathrm{d}\theta)s. \end{aligned}$$

□

For any $f \in \mathcal{B}_b^+(E)$ satisfying $m(f > 0) > 0$, define, for any $(t,x) \in [0,\infty) \times E$,

$$R_f(t,x) := T_t f(x) - V_t f(x), \tag{2.7}$$

and

$$g_f(t,x) := \frac{R_f(t,x)}{e^{\lambda_0 t}\langle f, \psi_0\rangle_m \phi_0(x)}. \tag{2.8}$$

**Lemma 2.2** *Assume that $f \in \mathcal{B}_b^+(E)$ and $m(f > 0) > 0$. Then*

$$R_f(t,x) \ge 0, \quad (t,x) \in [0,\infty) \times E, \tag{2.9}$$

*and*

$$\lim_{\|f\|_\infty \to 0} \|g_f(t)\|_\infty = 0, \quad t \ge 0. \tag{2.10}$$

**Proof:** It follows from [15, Theorem 2.23] and (1.4) that $V_t f(x)$ also satisfies

$$V_t f(x) = -\int_0^t T_s\left[r(\cdot, V_{t-s}f(\cdot))\right](x)\mathrm{d}s + T_t f(x), \quad t \ge 0, x \in E, \tag{2.11}$$

which implies that

$$R_f(t,x) = \int_0^t T_s\left[r(\cdot, V_{t-s}f(\cdot)\right](x)\mathrm{d}s, \quad t \ge 0, x \in E. \tag{2.12}$$

Since $r(x,s) \ge 0$, we have $R_f(t,x) \ge 0$, i.e., (2.9) holds.

Since $T_0 f(x) = V_0 f(x) = f(x)$, then $R_f(0,x) = 0$, which implies that $g_f(0,x) = 0$. Thus, it suffices to prove that, for any $\delta > 0$, (2.10) is true for $t > \delta$. It follows from Lemma 2.1 that for any $H \geq 1$,

$$\begin{aligned} r(x, V_{t-s}f(x)) &\leq (2+H)M(V_{t-s}f(x))^2 + V_{t-s}f(x)\int_H^\infty \theta\, n(x, \mathrm{d}\theta) \\ &\leq (2+H)M(T_{t-s}f(x))^2 + T_{t-s}f(x)\int_H^\infty \theta\, n(x, \mathrm{d}\theta). \end{aligned}$$

By (2.12), we have

$$\begin{aligned} R_f(t,x) &\leq (2+H)M\int_0^t T_s[(T_{t-s}f)^2](x)\mathrm{d}s + \int_0^t T_s\left[T_{t-s}f\int_H^\infty \theta\, n(\cdot, \mathrm{d}\theta)\right](x)\mathrm{d}s \\ &=: (I) + (II). \end{aligned}$$

For part (I), since $T_{t-s}f(x) \leq \|f\|_\infty e^{Mt}$, we have $T_s[(T_{t-s}f)^2](x) \leq e^{Mt}\|f\|_\infty T_t f(x)$. Thus, by (2.3), we have that, for any $t > \delta$,

$$\begin{aligned} (I) &\leq (2+H)Me^{Mt}t\|f\|_\infty T_t f(x) \\ &\leq (2+H)Me^{Mt}t(1+c(\delta))e^{\lambda_0 t}\langle f, \psi_0\rangle_m \phi_0(x)\|f\|_\infty. \end{aligned}$$

Hence we have that, for any $t > \delta$,

$$\lim_{\|f\|_\infty \to 0}\sup_{x\in E}\frac{(I)}{e^{\lambda_0 t}\langle f, \psi_0\rangle_m \phi_0(x)} = 0. \tag{2.13}$$

Now we deal with part (II). For any $H > 1$, $t > \delta$ and $0 < \epsilon < t$,

$$\left(\int_0^\epsilon + \int_{t-\epsilon}^t\right) T_s\Big[T_{t-s}f\int_H^\infty \theta\, n(\cdot, \mathrm{d}\theta)\Big](x)\mathrm{d}s \leq 2\epsilon M T_t f(x) \leq 2\epsilon M(1+c(\delta))e^{\lambda_0 t}\langle f, \psi_0\rangle_m \phi_0(x). \tag{2.14}$$

For any $\epsilon < s < t - \epsilon$, by (2.3),

$$T_{t-s}f(x) \leq (1+c(\epsilon))e^{\lambda_0 t}\langle f, \psi_0\rangle_m \phi_0(x).$$

Hence, we have

$$\begin{aligned} &\int_\epsilon^{t-\epsilon} T_s\Big[T_{t-s}f\int_H^\infty \theta\, n(\cdot, \mathrm{d}\theta)\Big](x)\mathrm{d}s \\ &\leq (1+c(\epsilon))e^{\lambda_0 t}\langle f, \psi_0\rangle_m \int_\epsilon^{t-\epsilon} T_s\Big[\int_H^\infty \theta\, n(\cdot, \mathrm{d}\theta)\phi_0\Big](x)\mathrm{d}s \\ &\leq (1+c(\epsilon))^2 e^{2\lambda_0 t}t\Big\langle \int_H^\infty \theta\, n(\cdot, \mathrm{d}\theta)\phi_0, \psi_0\Big\rangle_m \langle f, \psi_0\rangle_m \phi_0(x). \end{aligned} \tag{2.15}$$

Thus, combining (2.14) and (2.15), we get that

$$\limsup_{\|f\|_\infty \to 0}\sup_{x\in E}\frac{(II)}{e^{\lambda_0 t}\langle f, \psi_0\rangle_m \phi_0(x)} \leq 2\epsilon M(1+c(\delta)) + (1+c(\epsilon))^2 e^{\lambda_0 t}t\Big\langle \int_H^\infty \theta\, n(\cdot, \mathrm{d}\theta)\phi_0, \psi_0\Big\rangle_m.$$

Now, first letting $H \to \infty$ and then $\epsilon \to 0$, applying the monotone convergence theorem, we get that

$$\lim_{H\to\infty} \limsup_{\|f\|_\infty \to 0} \sup_{x\in E} \frac{(II)}{e^{\lambda_0 t}\langle f, \psi_0\rangle_m \phi_0(x)} = 0. \tag{2.16}$$

Combining (2.13) and (2.16), we get that

$$\lim_{\|f\|_\infty \to 0} \sup_{x\in E} \frac{R_f(t,x)}{e^{\lambda_0 t}\langle f, \psi_0\rangle_m \phi_0(x)} = 0.$$

□

# 3 Extinction probability

Recall that, for any $t > 0$ and $x \in E$,

$$q_t(x) = \mathbb{P}_{\delta_x}(\|X_t\| = 0) \qquad \text{and} \qquad q(x) = \lim_{t\to\infty} q_t(x).$$

**Lemma 3.1** *For any $x \in E$,*

$$q(x) < 1.$$

**Proof:** Let $v_t(x) := -\log q_t(x)$. Recall that $v(x) = -\log q(x)$. Since $q_t(x) < 1$, we have $v_t(x) > 0$. By Assumption 4, we have for $s > t_0$,

$$\|v\|_\infty \le \|v_s\|_\infty \le \|v_{t_0}\|_\infty = -\log\left(\inf_{x\in E} q_{t_0}(x)\right) < \infty.$$

Recall that, for $\theta > 0$,

$$V_t\theta(x) = -\log \mathbb{P}_{\delta_x} e^{-\langle \theta, X_t\rangle}.$$

By the Markov property of $X$,

$$q_{t+s}(x) = \lim_{\theta\to\infty} \mathbb{P}_{\delta_x}\left(e^{-\theta\|X_{t+s}\|}\right) = \lim_{\theta\to\infty} \mathbb{P}_{\delta_x}\left(e^{-\langle V_s\theta, X_t\rangle}\right) = \mathbb{P}_{\delta_x}\left(e^{-\langle v_s, X_t\rangle}\right). \tag{3.1}$$

It follows from (2.8) that, for any $s > t_0$,

$$v_{t+s}(x) = V_t(v_s)(x) = T_t(v_s)(x) - R_{v_s}(t,x) = T_t(v_s)(x) - g_{v_s}(t,x)e^{\lambda_0 t}\langle v_s, \psi_0\rangle_m \phi_0(x). \tag{3.2}$$

Thus, for $s > t_0$, we have

$$\langle v_{t+s}, \psi_0\rangle_m \ge (1 - \|g_{v_s}(t)\|_\infty)e^{\lambda_0 t}\langle v_s, \psi_0\rangle_m.$$

Since $v_t(x)$ is positive and non-increasing in $t$, we have that for all $t > 0$ and $s > t_0$,

$$(1 - \|g_{v_s}(t)\|_\infty)e^{\lambda_0 t} \le 1. \tag{3.3}$$

We claim that $\langle v, \psi_0\rangle_m > 0$. Otherwise, $\langle v, \psi_0\rangle_m = 0$. By (3.2), we have

$$\|v_{1+s}\|_\infty \le \|T_1(v_s)\|_\infty \le (1+c)e^{\lambda_0}\langle v_s, \psi_0\rangle_m \|\phi_0\|_\infty \to 0$$

as $s \to \infty$. Thus $\lim_{s\to\infty} \|v_s\|_\infty = 0$. Hence by (2.10),

$$\lim_{s\to\infty} \|g_{v_s}(t)\|_\infty = 0.$$

It follows from (3.3) that, for all $t > 0$, $e^{\lambda_0 t} \le 1$, which is a contradiction to the assumption that $\lambda_0 > 0$. Hence the claim above is valid.

By letting $s \to \infty$ in (3.1), we get that

$$\exp\{-v(x)\} = q(x) = \mathbb{P}_{\delta_x} \exp\left\{-\langle v, X_t\rangle\right\}. \tag{3.4}$$

Let $c$ and $\gamma$ be the constants in (2.2) with $\delta = 1$. For $t$ large enough, we have $1 - ce^{-\gamma t} > 0$. Thus for $t$ large enough, we have

$$T_t v(x) \ge (1 - ce^{-\gamma t})e^{\lambda_0 t}\langle v, \psi_0\rangle_m \phi_0(x) > 0.$$

Hence for all $x \in E$ and $t$ large enough,

$$\mathbb{P}_{\delta_x}(\langle v, X_t\rangle > 0) > 0,$$

which implies that

$$q(x) = \mathbb{P}_{\delta_x}(e^{-\langle v, X_t\rangle}) < 1.$$

The proof is now complete. □

**Lemma 3.2** *$V := \lim_{t\to\infty}\langle v, X_t\rangle \in [0, \infty]$ exists, and satisfies that, for all $x \in E$,*

$$\mathbb{P}_{\delta_x}(V = 0) = \exp\{-v(x)\} = q(x)$$

*and*

$$\mathbb{P}_{\delta_x}(V = \infty) = 1 - \exp\{-v(x)\} = 1 - q(x).$$

*Moreover, for any $\theta > 0$, we have*

$$\lim_{t\to\infty} V_t(\theta v)(x) = v(x), \quad x \in E. \tag{3.5}$$

**Proof:** By (3.4) and the Markov property of $X$, $\{e^{-\langle v, X_t\rangle}, t \ge 0\}$ is a bounded martingale. Thus $\lim_{t\to\infty} \exp\left\{-\langle v, X_t\rangle\right\}$ exists and is in $[0, 1]$, which implies that $V := \lim_{t\to\infty}\langle v, X_t\rangle \in [0, \infty]$ exists.

Since $\exp\left\{-\langle v, X_t\rangle\right\} \le 1$, we have

$$\mathbb{P}_{\delta_x} \exp\{-V\} = \exp\{-v(x)\}. \tag{3.6}$$

On the other hand,

$$\mathbb{P}_{\delta_x}(V=0) \geq \lim_{t\to\infty} \mathbb{P}_{\delta_x}(\|X_t\|=0) = \exp\{-v(x)\},$$

which implies that

$$\mathbb{P}_{\delta_x} \exp\{-V\} \geq \mathbb{P}_{\delta_x}(V=0) \geq \exp\{-v(x)\}. \tag{3.7}$$

It follows from (3.6) and (3.7) that

$$\mathbb{P}_{\delta_x}(V=0) = \exp\{-v(x)\} \qquad \text{and} \qquad \mathbb{P}_{\delta_x}(V=\infty) = 1-\exp\{-v(x)\}.$$

By the dominated convergence theorem, we get that, for any $\theta > 0$,

$$\lim_{t\to\infty} V_t(\theta v)(x) = -\log \mathbb{P}_{\delta_x}(e^{-\theta V}) = -\log \mathbb{P}_{\delta_x}(V=0) = v(x).$$

□

# 4 Backward iterates

It is clear that $V_t 0 = 0$. It follows from (3.4) that $V_t v = v$. Thus $v$ and 0 are two fixed points of $V_t$.

**Definition 4.1** *A family $(\eta_t, t \geq 0) \subset \mathcal{B}^+(E)$ is called a family of backward iterates if $\eta_t \leq v$ for all $t \geq 0$ and*

$$\eta_t(x) = V_s(\eta_{t+s})(x), \qquad t, s \geq 0,\ x \in E.$$

*A family $(\eta_t, t \geq 0)$ of backward iterates is said to be non-trivial if, for some $t \geq 0$, neither $\eta_t = 0, m$-a.e. nor $\eta_t = v, m$-a.e.*

It is well known that (see, for instance, [24, 1.44]), there exists a metric $d$ on $C(E)$ such that $(C(E), d)$ is a complete metric space, and convergence in $(C(E), d)$ is equivalent to uniform convergence on each compact subset $K$ of $E$.

For any $a > 0$, let $\mathcal{D}_a(E) := \{f \in \mathcal{B}^+(E) : \|f\|_\infty \leq a\}$.

**Lemma 4.2** *For any $t > 0$, $V_t(\mathcal{D}_a(E))$ is a relatively compact subset of $C(E)$.*

**Proof:** Without loss of generality, we only prove the lemma for $t = 1$ and $a \geq 1$. We first show that, for any compact subset $K \subset E$, $\{V_1 f : f \in \mathcal{D}_a(E)\}$ are equicontinuous on $K$.

Recall that

$$V_1 f(x) = T_1 f(x) - \int_0^1 T_s(r(\cdot, V_{1-s} f))(x) \mathrm{d}s.$$

It is clear that for $f \in \mathcal{D}_a(E)$,

$$|T_1 f(x) - T_1 f(y)| \leq a \int_E |q(1,x,z) - q(1,y,z)|\, m(\mathrm{d}z).$$

Note that by (1.11), we have

$$q(1,x,z) \le c_1\|\phi_0\|_\infty \psi_0(z). \tag{4.1}$$

Since $\psi_0 \in L^1(E,m)$, for any $\epsilon > 0$, we can choose a compact set $\widetilde{K} \subset E$ such that

$$2c_1\|\phi_0\|_\infty \int_{\widetilde{K}^c} \psi_0(z) m(\mathrm{d}z) < \frac{\epsilon}{2}.$$

Using the continuity of $q(1,\cdot,\cdot)$ on $K \times \widetilde{K}$, (4.1) and the fact that $\psi_0 \in L^1(E,m)$, we can find a $\delta > 0$ such that for any $x, y \in K$ with $|x-y| < \delta$,

$$\int_{\widetilde{K}} |q(1,x,z) - q(1,y,z)|\, m(\mathrm{d}z) < \frac{\epsilon}{2}.$$

Thus $\{T_1 f : f \in \mathcal{D}_a(E)\}$ are equicontinuous on $K$. By (2.4) and the fact that $V_t f(x) \le T_t f(x) \le e^{Mt}\|f\|_\infty$, we have that for any $f \in \mathcal{D}_a(E)$, $r(x, V_{1-s}f(x)) \le 2M(e^{2M}a^2 + e^M a)$. Thus

$$\begin{aligned}
&\left| \int_0^1 T_s(r(\cdot V_{1-s}f)(x)\mathrm{d}s - \int_0^1 T_s(r(\cdot, V_{1-s}f)(y)\mathrm{d}s \right| \\
&\le [2M(e^{2M}a^2 + e^M a)] \int_0^1 \int_E |q(s,x,z) - q(s,y,z)|\, m(\mathrm{d}z)\mathrm{d}s.
\end{aligned}$$

Note that, for $\eta \in (0,1)$,

$$\begin{aligned}
&\int_0^1 \int_E |q(s,x,z) - q(s,y,z)|\, m(\mathrm{d}z)\mathrm{d}s \\
&\le \int_0^\eta \int_E q(s,x,z) + q(s,y,z)\, m(\mathrm{d}z)\mathrm{d}s + \int_\eta^1 \int_E |q(s,x,z) - q(s,y,z)|\, m(\mathrm{d}z)\mathrm{d}s \\
&\le 2e^M\eta + \int_\eta^1 \int_E |q(s,x,z) - q(s,y,z)|\, m(\mathrm{d}z)\mathrm{d}s \\
&\le 2e^M\eta + \int_\eta^1 \int_E \int_E |q(\frac{\eta}{2},x,w) - q(\frac{\eta}{2},y,w)| q(s-\frac{\eta}{2},w,z)\, m(\mathrm{d}w)\, m(\mathrm{d}z)\mathrm{d}s.
\end{aligned}$$

For any $\epsilon > 0$, we choose $\eta \in (0,1)$ so that $2e^M\eta < \frac{\epsilon}{3}$. It follows from (2.1) that, for any $s \in [\eta/2, 1)$,

$$q(s,x,z) \le (1 + c(\eta/2))e^{\lambda_0}\|\phi_0\|_\infty \psi_0(z). \tag{4.2}$$

Hence

$$\begin{aligned}
&\int_\eta^1 \int_E \int_E |q(\frac{\eta}{2},x,w) - q(\frac{\eta}{2},y,w)| q(s-\frac{\eta}{2},w,z)\, m(\mathrm{d}w)\, m(\mathrm{d}z)\mathrm{d}s \\
&\le (1 + c(\eta/2))e^{\lambda_0}\|\phi_0\|_\infty \int_\eta^1 \int_E \int_E |q(\frac{\eta}{2},x,w) - q(\frac{\eta}{2},y,w)|\, m(\mathrm{d}w)\psi_0(z)\, m(\mathrm{d}z)\mathrm{d}s.
\end{aligned}$$

Applying (4.2) again, we can find a compact $\widetilde{K} \subset E$ such that

$$(1 + c(\eta/2))e^{\lambda_0}\|\phi_0\|_\infty \int_\eta^1 \int_E \int_{\widetilde{K}^c} |q(\frac{\eta}{2}, x, w) - q(\frac{\eta}{2}, y, w)|\, m(\mathrm{d}w)\psi_0(z)\, m(\mathrm{d}z)\mathrm{d}s < \frac{\epsilon}{3}.$$

Using the continuity of $q(\frac{\eta}{2}, \cdot, \cdot)$ on $K \times \widetilde{K}$, (4.2) and the fact that $\psi_0 \in L^1(E, m)$, we can find a $\delta > 0$ such that for any $x, y \in K$ with $|x - y| < \delta$,

$$(1 + c(\eta/2))e^{\lambda_0}\|\phi_0\|_\infty \int_\eta^1 \int_E \int_{\widetilde{K}} |q(\frac{\eta}{2}, x, w) - q(\frac{\eta}{2}, y, w)|\, m(\mathrm{d}w)\psi_0(z)\, m(\mathrm{d}z)\mathrm{d}s < \frac{\epsilon}{3}.$$

Thus $\{\int_0^1 T_s(r(\cdot, V_{1-s}f))(x)\mathrm{d}s : f \in \mathcal{D}_a(E)\}$ are equicontinuous on $K$. It follows that $\{V_1 f : f \in \mathcal{D}_a(E)\}$ are equicontinuous on $K$. In particular, $V_1(\mathcal{D}_a(E)) \subset C(E)$.

Let $K_n$ be an increasing sequence of compact subsets of $E$. Using the equicontinuity of $\{V_1 f : f \in \mathcal{D}_a(E)\}$ on each $K_n$ and a routine diagonalization argument, we can easily show that any sequence of functions in $V_1(\mathcal{D}_a(E))$ contains a subsequence which converges in $C(E)$. □

**Proposition 4.3** *There exists a non-trivial family of backward iterates.*

**Proof:** Let $\mathcal{D} = \{f \in \mathcal{B}^+(E) : f \le v\}$. For any $f \in \mathcal{D}$, $0 \le V_s f \le V_s v \le v$. Thus $V_{t+s}f = V_t(V_s f) \in V_t(\mathcal{D})$, which implies that $t \to V_t(\mathcal{D})$ is decreasing.

For any $g, h \in \mathcal{D}$ and $t > 0$, it is easy to see that $\lambda \mapsto V_t(\lambda g + (1 - \lambda)h)$, $\lambda \in [0, 1]$, is a continuous curve in $V_t(\mathcal{D})$ connecting $V_t g$ to $V_t h$. Thus $V_t(\mathcal{D})$ is connected. Hence $\mathcal{D}_\infty := \bigcap_{n\in\mathbb{N}} V_n(\mathcal{D})$ is connected. Since $0, v \in \mathcal{D}_\infty$ and $v > 0$, there exists an $\eta \in \mathcal{D}_\infty$, such that $0 < \eta < v$ on a set of positive measure. Thus, for every $n \in \mathbb{N}$, there exists $\eta_{n,n} \in \mathcal{D}$ such that $V_n(\eta_{n,n}) = \eta$. Define $\eta_{n,j} = V_{n-j}(\eta_{n,n})$, $j = 0, 1, 2, \cdots, n - 1$. Note that $\eta_{n,0} = \eta$.

Since $0 \le \eta_{n,j} \le v$ and $\eta_{n,j} = V_1(\eta_{n,j+1})$, it follows from Lemma 4.2 that $\{\eta_{n,j}, n \ge 0, 0 \le j < n\}$ is relatively compact in $C(E)$. Thus there exists a sequence $(\eta_{n_l,j})$ such that $\eta_j := \lim_{l\to\infty} \eta_{n_l,j} \in C(E) \bigcap \mathcal{D}$ exists. Since $\eta_{n_l,j} = V_1(\eta_{n_l,j+1})$, letting $l \to \infty$, we get that $\eta_j = V_1(\eta_{j+1})$.

Define $\eta_t := V_{[t+1]-t}(\eta_{[t+1]})$, for $t \ge 0$. Then

$$V_s(\eta_{t+s}) = V_s V_{[t+s+1]-t-s}(\eta_{[t+s+1]}) = V_{[t+s+1]-t}(\eta_{[t+s+1]}) = V_{[t+1]-t}(\eta_{[t+1]}) = \eta_t.$$

It follows from $\eta_{n,0} = \eta$ that $\eta_0 = \eta$, which implies that the family $\{\eta_t, t \ge 0\}$ of backward iterates is nontrivial. □

**Lemma 4.4** *If $0 \le f \le v$, then*

$$V_t f(x) \ge e^{-at} T_t f(x), \quad t \ge 0, x \in E,$$

*where $a = 2M(1 + \|v\|_\infty)$.*

**Proof:** Using (2.4) with $H = 1$ and the fact that $V_{t-s}f(x) \le V_{t-s}v(x) = v(x) \le \|v\|_\infty$ for any $0 \le s \le t$ and any $x \in E$, we get

$$r(x, V_{t-s}f(x)) \le 2M(1 + \|v\|_\infty)V_{t-s}f(x) = aV_{t-s}f(x), \quad 0 \le s \le t, x \in E.$$

Recall that

$$V_t f(x) = T_t f(x) - \int_0^t T_s(r(\cdot, V_{t-s}f(\cdot)))(x)\mathrm{d}s.$$

Thus we have

$$V_t f(x) = e^{-at}T_t f(x) - \int_0^t e^{-as}T_s(r(\cdot, V_{t-s}f(\cdot)))(x)\mathrm{d}s + a\int_0^t e^{-as}T_s(V_{t-s}f)(x)\mathrm{d}s.$$

Consequently we have

$$V_t f(x) \ge e^{-at}T_t f(x), \quad t \ge 0, x \in E.$$

□

**Lemma 4.5** *If $(\eta_t, t \ge 0)$ is a non-trivial family of backward iterates, then*

$$\lim_{t\to\infty} \|\eta_t\|_\infty = 0.$$

**Proof:** Without loss of generality, we may and will assume that $m(\eta_0 > 0) > 0$ and $m(\eta_0 < v) > 0$. We claim that $\langle \eta_t, \psi_0\rangle_m \to 0$ as $t \to \infty$. Otherwise, there exist a sequence $t_j \uparrow \infty$ and a constant $c_0 > 0$ such that

$$\langle \eta_{t_j}, \psi_0\rangle_m > c_0.$$

Fix $s > 1$ large enough such that $1 - ce^{-\gamma s} > 0$, where $c$ and $\gamma$ are the constants in (2.2) with $\delta = 1$. Then, for $j$ large enough so that $t_j > s$, we have

$$\eta_0(x) = V_{t_j}(\eta_{t_j})(x) = V_{t_j-s}(V_s(\eta_{t_j}))(x), \quad x \in E. \tag{4.3}$$

By Lemma 4.4, we have

$$V_s(\eta_{t_j})(x) \ge e^{-as}T_s(\eta_{t_j})(x) \ge e^{-as}(1-ce^{-\gamma s})e^{\lambda_0 s}\langle \eta_{t_j}, \psi_0\rangle_m \phi_0(x) \ge c_0 e^{-as}(1-ce^{-\gamma s})e^{\lambda_0 s}\phi_0(x).$$

It follows from (2.7) and (2.8) that

$$v(x) = V_1 v(x) \le T_1 v(x) \le (1 + c)e^{\lambda_0}\langle v, \psi_0\rangle_m \phi_0(x). \tag{4.4}$$

Thus

$$V_s(\eta_{t_j})(x) \ge c_0 e^{-as}(1 - ce^{-\gamma s})e^{\lambda_0 s}(1 + c)^{-1}e^{-\lambda_0}\langle v, \psi_0\rangle_m^{-1} v(x) =: C_s v(x). \tag{4.5}$$

Combining (4.5), (4.3) and (3.5), we get

$$\eta_0(x) \ge \lim_{j\to\infty} V_{t_j-s}(C_s v)(x) = v(x),$$

which contradicts the definition of $\eta_0$. Thus the claim above is true.

Note that, for any $s \geq 1$ and $t > 0$,

$$\eta_t(x) = V_s(\eta_{t+s})(x) \leq T_s(\eta_{t+s})(x) \leq (1 + ce^{-\gamma s})e^{\lambda_0 s}\langle \eta_{t+s}, \psi_0\rangle_m \phi_0(x). \tag{4.6}$$

Thus

$$\|\eta_t\|_\infty \leq (1 + c)e^{\lambda_0 s}\langle \eta_{t+s}, \psi_0\rangle_m \|\phi_0\|_\infty \to 0,$$

as $t \to \infty$. The proof is now complete. □

**Lemma 4.6** *If $(\eta_t, t \geq 0)$ is a non-trivial family of backward iterates, then there exist $\{h_t : t \geq 0\} \subset \mathcal{B}_b(E)$ such that*

$$\eta_t(x) = (1 + h_t(x))\langle \eta_t, \psi_0\rangle_m \phi_0(x), \quad t \geq 0, x \in E,$$

*and*

$$\lim_{t\to\infty} \|h_t\|_\infty = 0.$$

*Moreover,*

$$\frac{\langle \eta_t, \psi_0\rangle_m}{\langle \eta_{t+s}, \psi_0\rangle_m} \geq e^{\lambda_0 s}, \forall s, t \geq 0 \quad \textit{and} \quad \lim_{t\to\infty} \frac{\langle \eta_t, \psi_0\rangle_m}{\langle \eta_{t+s}, \psi_0\rangle_m} = e^{\lambda_0 s}, \forall s \geq 0. \tag{4.7}$$

**Proof:** Since $\langle \eta_t, \psi_0\rangle_m \phi_0(x) > 0$, we can define

$$h_t(x) := \frac{\eta_t(x)}{\langle \eta_t, \psi_0\rangle_m \phi_0(x)} - 1, \quad t \geq 0, x \in E.$$

It follows from (2.2) and Lemma 2.2 that for $s > 1$,

$$\eta_t(x) = V_s(\eta_{t+s})(x) = T_s(\eta_{t+s})(x) - R_{\eta_{t+s}}(s, x) \tag{4.8}$$

$$\geq e^{\lambda_0 s}(1 - ce^{-\gamma s} - \|g_{\eta_{t+s}}(s)\|_\infty)\langle \eta_{t+s}, \psi_0\rangle_m \phi_0(x), \tag{4.9}$$

where $c$ and $\gamma$ are the constants in (2.2) with $\delta = 1$. In the remainder of this proof, we assume that $s$ is large enough such that $1 - ce^{-\gamma s} > 0$. By Lemma 4.5 and (2.10), $\lim_{t\to\infty} \|g_{\eta_{t+s}}(s)\|_\infty = 0$. Thus, for large enough $t$, $1 - ce^{-\gamma s} - \|g_{\eta_{t+s}}(s)\|_\infty > 0$. It follows from (4.6) and (4.9) that

$$-\frac{2ce^{-\gamma s} + \|g_{\eta_{t+s}}(s)\|_\infty}{1 + ce^{-\gamma s}} \leq h_t(x) = \frac{\eta_t(x)}{\langle \eta_t, \psi_0\rangle_m \phi_0(x)} - 1 \leq \frac{2ce^{-\gamma s} + \|g_{\eta_{t+s}}(s)\|_\infty}{1 - ce^{-\gamma s} - \|g_{\eta_{t+s}}(s)\|_\infty}.$$

Letting $t \to \infty$ and then $s \to \infty$, we get that

$$\lim_{t\to\infty} \|h_t\|_\infty = 0.$$

It follows from (4.8) that

$$\begin{aligned}\langle \eta_t, \psi_0\rangle_m &= \langle T_s(\eta_{t+s}), \psi_0\rangle_m - \langle R_{\eta_{t+s}}(s,x), \psi_0\rangle_m \\ &= e^{\lambda_0 s}\langle \eta_{t+s}, \psi_0\rangle_m - e^{\lambda_0 s}\langle \eta_{t+s}, \psi_0\rangle_m \langle g_{\eta_{t+s}}(s)\phi_0, \psi_0\rangle_m,\end{aligned}$$

which implies that

$$0 \le e^{\lambda_0 s} - \frac{\langle \eta_t, \psi_0\rangle_m}{\langle \eta_{t+s}, \psi_0\rangle_m} = e^{\lambda_0 s}\langle g_{\eta_{t+s}}(s)\phi_0, \psi_0\rangle_m \le e^{\lambda_0 s}\|g_{\eta_{t+s}}(s)\|_\infty \to 0,$$

as $t \to \infty$, where the last limit follows from (2.10) and Lemma 4.5. The proof is now complete. □

**Remark 4.7** *Let $L(t) := e^{\lambda_0 t}\langle \eta_t, \psi_0\rangle_m$. It follows from (4.7) that $t \to L(t)$ is nondecreasing and, for any $s \ge 0$,*

$$\lim_{t\to\infty} \frac{L(t+s)}{L(t)} = 1.$$

*Therefore, $l_0 := \lim_{t\to\infty} L(t) \in (0, \infty]$ exists.*

# 5 Seneta-Heyde norming for $M$

**Lemma 5.1** *If $f \in \mathcal{B}^+(E)$ satisfies $f(x) \le v(x)$ and $f(x) = V_t f(x)$ for all $t > 0$ and $x \in E$, then either $f(x) = v(x)$ for all $x \in E$ or $f(x) = 0$ for all $x \in E$.*

**Proof:** Let $s_0 > 1$ be such that $1 - ce^{-\gamma s_0} > 0$, where $c$ and $\gamma$ are the constants from (2.2) with $\delta = 1$. By Lemma 4.4 and (4.4), we have

$$f(x) = V_{s_0} f(x) \ge e^{-as_0} T_{s_0} f(x) \ge e^{-as_0}(1 - ce^{-\gamma s_0})e^{\lambda_0 s_0}\langle f, \psi_0\rangle_m \phi_0(x) \ge c_0\langle f, \psi_0\rangle_m v(x)$$

for some constant $c_0 > 0$. If $\langle f, \psi_0\rangle_m > 0$, then by (3.5) we have

$$f(x) = \lim_{t\to\infty} V_t f(x) \ge \lim_{t\to\infty} V_t(c_0\langle f, \psi_0\rangle_m v)(x) = v(x),$$

which implies that $f(x) = v(x)$ for all $x \in E$. If $\langle f, \psi_0\rangle_m = 0$, then

$$f(x) \le T_1 f(x) \le (1 + c)e^{\lambda_0}\langle f, \psi_0\rangle_m \phi_0(x) = 0,$$

which implies that $f(x) = 0$ for all $x \in E$. The proof is now complete. □

**Theorem 5.2** *Let $(\eta_t, t \ge 0)$ be a non-trivial family of backward iterates and define $\gamma_t := \langle \eta_t, \psi_0\rangle_m$. Then there is a non-degenerate random variable $W$ such that for any nonzero $\mu \in \mathcal{M}_F(E)$,*

$$\lim_{t\to\infty} \gamma_t\langle \phi_0, X_t\rangle = W, \qquad a.s.\text{-}\mathbb{P}_\mu$$

*and*

$$\mathbb{P}_\mu(W = 0) = \exp\{-\langle v, \mu\rangle\}, \qquad \mathbb{P}_\mu(W < \infty) = 1. \tag{5.1}$$

**Proof:** By the definition of $\eta_t$, for any nonzero $\mu \in \mathcal{M}_F(E)$,

$$\mathbb{P}_\mu(\exp\{-\eta_{t+s}, X_{t+s}\}|\mathcal{G}_t) = \mathbb{P}_{X_t}(\exp\{-\eta_{t+s}, X_s\}) = \exp\{-\langle \eta_t, X_t\rangle\},$$

which implies that $\{\exp\{-\langle \eta_t, X_t\rangle\}, t \geq 0\}$ is a nonnegative martingale. Thus, by the martingale convergence theorem, $\lim_{t\to\infty} \exp\{-\langle \eta_t, X_t\rangle\}$ exists $\mathbb{P}_\mu$ almost surely and hence $W := \lim_{t\to\infty}\langle \eta_t, X_t\rangle \in [0, \infty]$ exists $\mathbb{P}_\mu$ almost surely.

It follows from Lemma 4.6 that

$$(1 - \|h_t\|_\infty)\gamma_t\langle \phi_0, X_t\rangle \leq \langle \eta_t, X_t\rangle \leq (1 + \|h_t\|_\infty)\gamma_t\langle \phi_0, X_t\rangle.$$

Since $\lim_{t\to\infty} \|h_t\|_\infty = 0$, we have $1 - \|h_t\|_\infty > 0$ for $t$ large enough. Thus for large $t$,

$$(1 + \|h_t\|_\infty)^{-1}\langle \eta_t, X_t\rangle \leq \gamma_t\langle \phi_0, X_t\rangle \leq (1 - \|h_t\|_\infty)^{-1}\langle \eta_t, X_t\rangle.$$

Letting $t \to \infty$, we get that

$$\lim_{t\to\infty} \gamma_t\langle \phi_0, X_t\rangle = W, \qquad \text{a.s.-}\mathbb{P}_\mu.$$

Define

$$\Phi(s, x) := -\log \mathbb{P}_{\delta_x} \exp\{-sW\}. \tag{5.2}$$

Then

$$\begin{aligned} -\log \mathbb{P}_\mu \exp\{-sW\} &= \lim_{t\to\infty} -\log \mathbb{P}_\mu \exp\{-s\gamma_t\langle \phi_0, X_t\rangle\} \\ &= \lim_{t\to\infty} \langle -\log \mathbb{P}_{\delta_x} \exp\{-s\gamma_t\langle \phi_0, X_t\rangle\}, \mu\rangle = \langle \Phi(s, \cdot), \mu\rangle. \end{aligned}$$

Put $\Phi_\infty(x) := \lim_{s\to\infty} \Phi(s, x)$ and $\Phi_0(x) = \lim_{s\to 0} \Phi(s, x)$. Then

$$\mathbb{P}_\mu(W = 0) = \lim_{s\to\infty} \mathbb{P}_\mu \exp\{-sW\} = \exp\{-\langle \Phi_\infty, \mu\rangle\}$$

and

$$\mathbb{P}_\mu(W < \infty) = \lim_{s\to 0} \mathbb{P}_\mu \exp\{-sW\} = \exp\{-\langle \Phi_0, \mu\rangle\}.$$

For any $s, t > 0$, we have

$$\begin{aligned} \exp\{-\Phi(s, x)\} &= \lim_{u\to\infty} \mathbb{P}_{\delta_x}(\exp\{-s\gamma_{t+u}\langle \phi_0, X_{t+u}\rangle\}) \\ &= \lim_{u\to\infty} \mathbb{P}_{\delta_x}\mathbb{P}_{X_t}(\exp\{-s\gamma_{t+u}\gamma_u^{-1}\gamma_u\langle \phi_0, X_u\rangle\}) \\ &= \mathbb{P}_{\delta_x}\mathbb{P}_{X_t} \exp\{-se^{-\lambda_0 t}W\} = \mathbb{P}_{\delta_x} \exp\{-\langle \Phi(se^{-\lambda_0 t}, \cdot), X_t\rangle\}, \end{aligned}$$

which implies that

$$\Phi(s, x) = V_t(\Phi(se^{-\lambda_0 t}, \cdot))(x). \tag{5.3}$$

Thus, letting $s \to \infty$ and $s \to 0$, we get respectively

$$\Phi_\infty(x) = V_t(\Phi_\infty)(x), \qquad \Phi_0(x) = V_t(\Phi_0)(x).$$

Since $s \to \Phi(s,x)$ is increasing, we have $\Phi_0(x) \le \Phi(1,x) \le \Phi_\infty(x)$. Note that

$$\Phi(1,x) = -\log \mathbb{P}_{\delta_x} \exp\{-W\} = \lim_{t\to\infty} -\log \mathbb{P}_{\delta_x} \exp\{-\langle \eta_t, X_t\rangle\} = \eta_0(x).$$

On the other hand,

$$\mathbb{P}_{\delta_x}(W = 0) \ge \mathbb{P}_{\delta_x}(\exists t > 0, \|X_t\| = 0) = \exp\{-v(x)\},$$

which implies that, for all $x \in E$, $\Phi_\infty(x) \le v(x)$. Thus, $\Phi_0(x) \le \eta_0(x) \le \Phi_\infty(x) \le v(x)$. It follows from Lemma 5.1 that

$$\Phi_0 = 0, \quad \Phi_\infty = v.$$

The proof is now complete. □

Now Theorem 1.1 follows immediately from Lemma 4.6 and Theorem 5.2.

**Proposition 5.3** *Let $(\eta_t, t \ge 0)$ be a non-trivial family of backward iterates and $W$ be the limit in Theorem 5.2 corresponding to $(\eta_t, t \ge 0)$. Then a family $(\eta^*_t, t \ge 0)$ is a non-trivial family of backward iterates if and only if there exists a positive number $a$ such that*

$$\eta^*_t(x) = \Phi(ae^{-\lambda_0 t}, x), \quad t \ge 0, x \in E,$$

*where $\Phi$ is defined in* (5.2)*. In particular,*

$$\eta_t(x) = \Phi(e^{-\lambda_0 t}, x), \quad t \ge 0, x \in E. \tag{5.4}$$

**Proof:** For any $a > 0$, by (5.3), we have

$$V_s(\Phi(ae^{-\lambda_0(t+s)}, \cdot))(x) = \Phi(ae^{-\lambda_0 t}, x).$$

Thus $(\Phi(ae^{-\lambda_0 t}, x), t \ge 0)$ is a non-trivial family of backward iterates.

If $(\eta^*_t, t \ge 0)$ is a non-trivial family of backward iterates, then it follows from Lemma 4.6 that, for any $s \ge 0$, $(\eta^*_{t+s}, t \ge 0)$ is also a non-trivial family of backward iterates. Let $W^{(s)}$ be the limit in Theorem 5.2 corresponding to $(\eta^*_{t+s}, t \ge 0)$. By (5.1), we get that, for any $s \ge 0$,

$$\{W > 0\} = \{W^{(s)} > 0\} = \{\forall\, t > 0,\ \|X_t\| > 0\}, \qquad \text{a.s.-}\mathbb{P}_\mu.$$

Thus, we have that, for any $\omega \in \{W > 0\} \cap \{W^{(s)} > 0\}$,

$$\frac{W^{(s)}(\omega)}{W(\omega)} = \lim_{t\to\infty} \frac{\langle \eta^*_{t+s}, \psi_0\rangle_m}{\langle \eta_t, \psi_0\rangle_m} := e(s) > 0,$$

where $e(s)$ is deterministic. Therefore,

$$W^{(s)} = e(s)W, \qquad \text{a.s.-}\mathbb{P}_\mu. \tag{5.5}$$

By (4.7), we have that, for any $s, r \geq 0$,

$$W^{(s)} = \lim_{t\to\infty} \frac{\langle \eta^*_{t+s}, \psi_0\rangle_m}{\langle \eta^*_{t+r}, \psi_0\rangle_m} W^{(r)} = e^{-\lambda_0(s-r)} W^{(r)}, \qquad \text{a.s.-}\mathbb{P}_\mu.$$

It follows that $e(s) = e^{-\lambda_0(s-r)}e(r)$, which implies that there exists a constant $a > 0$ such that

$$e(s) = ae^{-\lambda_0 s}.$$

Note that, for any $s \geq 0$ and $x \in E$,

$$\begin{aligned}\eta^*_s(x) &= \lim_{t\to\infty} -\log \mathbb{P}_{\delta_x} \exp\{-\langle \eta^*_{t+s}, X_t\rangle\} = -\log \mathbb{P}_{\delta_x} \exp\{-W^{(s)}\} \\ &= -\log \mathbb{P}_{\delta_x} \exp\{-ae^{-\lambda_0 s} W\} = \Phi(ae^{-\lambda_0 s}, x).\end{aligned}$$

The proof is now complete. □

Recall from Remark 4.7 that $l_0 = \lim_{t\to\infty} e^{\lambda_0 t}\gamma_t \in (0, \infty]$.

**Proposition 5.4** *Let $(\eta_t, t \geq 0)$ be a non-trivial family of backward iterates and $W$ be the limit in Theorem 5.2 corresponding to $(\eta_t, t \geq 0)$.*

*(1) If $l_0 < \infty$, then $\mathbb{P}_\mu W < \infty$ for any $\mu \in \mathcal{M}_F(E)$.*

*(2) If $\mathbb{P}_\mu W < \infty$ for some nonzero $\mu \in \mathcal{M}_F(E)$, then $l_0 < \infty$.*

*Moreover,*

$$\mathbb{P}_\mu W = \lim_{t\to\infty} \gamma_t \mathbb{P}_\mu \langle \phi_0, X_t\rangle = l_0 \langle \phi_0, \mu\rangle. \tag{5.6}$$

**Proof:** (1) Since $W = \lim_{t\to\infty} \gamma_t \langle \phi_0, X_t\rangle$, we have by Fatou's lemma that

$$\mathbb{P}_\mu W \leq \lim_{t\to\infty} \mathbb{P}_\mu \gamma_t \langle \phi_0, X_t\rangle = \lim_{t\to\infty} \gamma_t e^{\lambda_0 t} \langle \phi_0, \mu\rangle = l_0 \langle \phi_0, \mu\rangle.$$

Thus $l_0 < \infty$ implies $\mathbb{P}_\mu W < \infty$ for any $\mu \in \mathcal{M}_F(E)$.

(2) It follows from (5.4) and Lemma 4.6 that

$$\begin{aligned}\lim_{s\to 0} \frac{\langle \Phi(s,\cdot), \mu\rangle}{s} &= \lim_{t\to\infty} e^{\lambda_0 t} \langle \Phi(e^{-\lambda_0 t}, \cdot), \mu\rangle = \lim_{t\to\infty} e^{\lambda_0 t} \langle \eta_t, \mu\rangle \\ &= \lim_{t\to\infty} e^{\lambda_0 t} \gamma_t \langle (1 + h_t)\phi_0, \mu\rangle = l_0 \langle \phi_0, \mu\rangle. \end{aligned} \tag{5.7}$$

If $\mathbb{P}_\mu W < \infty$ for some nonzero $\mu \in \mathcal{M}_F(E)$, then

$$\mathbb{P}_\mu W = \lim_{s\to 0} \frac{1 - \mathbb{P}_\mu(\exp\{-sW\})}{s} = \lim_{s\to 0} \frac{\langle \Phi(s,\cdot), \mu\rangle}{s} = l_0 \langle \phi_0, \mu\rangle,$$

which implies $l_0 < \infty$. The proof is now complete. □

It was shown in [1, Section 1.10, Lemma 1] that, if $Y$ is a non-negative random variable with $EY < \infty$, then

$$EY \log^+ Y < \infty$$

if and only if, for some $a_0 > 0$

$$\int_0^{a_0} s^{-2} E(e^{-sY} - 1 + sY)\, \mathrm{d}s < \infty.$$

Recall that $l(x) := \int_1^\infty \theta \log \theta\, n^{\phi_0}(x, \mathrm{d}\theta)$.

**Lemma 5.5** *$\int_E \psi_0(x) l(x) m(\mathrm{d}x) < \infty$ if and only if, for any $t > 0$,*

$$\int_E \psi_0(x) \mathbb{P}_{\delta_x}\Big[\langle \phi_0, X_t\rangle \log^+ \langle \phi_0, X_t\rangle\Big] m(\mathrm{d}x) < \infty.$$

**Proof:** Without loss of generality, we only prove the result for $t = 1$. Note that

$$\int_E \psi_0(x) \mathbb{P}_{\delta_x}\Big[\langle \phi_0, X_1\rangle \log^+ \langle \phi_0, X_1\rangle\Big] m(\mathrm{d}x) < \infty$$

if and only if, for some $a_0 > 0$

$$\int_E \psi_0(x) m(\mathrm{d}x) \int_0^{a_0} s^{-2} \mathbb{P}_{\delta_x}(e^{-s\langle \phi_0, X_1\rangle} - 1 + s\langle \phi_0, X_1\rangle)\, \mathrm{d}s < \infty. \tag{5.8}$$

Put $R(f)(x) := R_f(1, x) = T_1 f(x) - V_1 f(x)$. Then, we have

$$\begin{aligned}
&\mathbb{P}_{\delta_x}(e^{-s\langle \phi_0, X_1\rangle} - 1 + s\langle \phi_0, X_1\rangle) \\
&= \exp\{-V_1(s\phi_0)(x)\} - 1 + V_1(s\phi_0)(x) + R(s\phi_0)(x) \\
&\le \frac{1}{2} V_1(s\phi_0)(x)^2 + R(s\phi_0)(x) \\
&\le \frac{1}{2} T_1(s\phi_0)(x)^2 + R(s\phi_0)(x) \\
&= \frac{1}{2} e^{2\lambda_0} s^2 \phi_0(x)^2 + R(s\phi_0)(x).
\end{aligned}$$

On the other hand,

$$\mathbb{P}_{\delta_x}(e^{-s\langle \phi_0, X_1\rangle} - 1 + s\langle \phi_0, X_1\rangle) = \exp\{-V_1(s\phi_0)(x)\} - 1 + T_1(s\phi_0)(x) \ge R(s\phi_0)(x).$$

Thus, (5.8) holds if and only if, for some $a_0 > 0$,

$$\int_0^{a_0} s^{-2} \langle R(s\phi_0), \psi_0\rangle_m \mathrm{d}s < \infty. \tag{5.9}$$

By (2.5), we have

$$\langle r(\cdot, V_{1-t}(s\phi_0)), \psi_0\rangle_m$$

$$= \langle \beta V_{1-t}(s\phi_0)^2, \psi_0\rangle_m + \Big\langle \Big( \int_0^{\phi_0^{-1}} + \int_{\phi_0^{-1}}^{\infty} \Big) \Big( e^{-\theta V_{1-t}(s\phi_0)} - 1 + \theta V_{1-t}(s\phi_0) \Big) n(\cdot, \mathrm{d}\theta), \psi_0 \Big\rangle_m$$
$$=: J_1(s,t) + J_2(s,t) + J_3(s,t).$$

Thus, by (2.12), we have

$$\begin{aligned}
&\int_0^{a_0} s^{-2} \langle R(s\phi_0), \psi_0\rangle_m \mathrm{d}s \\
&= \int_0^{a_0} s^{-2} \mathrm{d}s \int_0^1 e^{\lambda_0 t} \langle r(\cdot, V_{1-t}(s\phi_0)), \psi_0\rangle_m \, \mathrm{d}t \\
&= \int_0^{a_0} s^{-2} \mathrm{d}s \int_0^1 e^{\lambda_0 t} J_1(s,t) \, \mathrm{d}t + \int_0^{a_0} s^{-2} \mathrm{d}s \int_0^1 e^{\lambda_0 t} J_2(s,t) \, \mathrm{d}t \\
&\quad + \int_0^{a_0} s^{-2} \mathrm{d}s \int_0^1 e^{\lambda_0 t} J_3(s,t) \, \mathrm{d}t.
\end{aligned}$$

Since

$$V_{1-t}(s\phi_0)(x) \le T_{1-t}(s\phi_0)(x) \le s e^{\lambda_0} \phi_0(x), \tag{5.10}$$

we have that $J_1(s,t) \le M e^{2\lambda_0} s^2 \|\phi_0\|_\infty$. Thus

$$\int_0^{a_0} s^{-2} \mathrm{d}s \int_0^1 e^{\lambda_0 t} J_1(s,t) \mathrm{d}t < \infty. \tag{5.11}$$

Note that

$$J_2(s,t) \le \Big\langle \int_0^{\phi_0^{-1}} \theta^2 V_{1-t}(s\phi_0)^2 n(\cdot, \mathrm{d}\theta), \psi_0 \Big\rangle_m \le e^{2\lambda_0} s^2 \Big\langle \int_0^1 \theta^2 n^{\phi_0}(\cdot, \mathrm{d}\theta), \psi_0 \Big\rangle_m \le C s^2,$$

which implies that

$$\int_0^{a_0} s^{-2} \mathrm{d}s \int_0^1 e^{\lambda_0 t} J_2(s,t) \mathrm{d}t < \infty. \tag{5.12}$$

Now we deal with $J_3$. By Lemma 4.4 and (2.3) with $s_0 > 1$ large enough such that $c(1) e^{-\gamma(1) s_0} < \frac{1}{2}$,

$$v(x) = V_{s_0} v(x) \ge e^{-a s_0} T_{s_0} v(x) \ge \frac{1}{2} e^{(\lambda_0 - a) s_0} \phi_0(x).$$

We put $A_0 = \frac{1}{2} e^{(\lambda_0 - a) s_0}$. Hence by Lemma 4.4, for any $s \le A_0$,

$$V_{1-t}(s\phi_0)(x) \ge e^{-a} T_{1-t}(s\phi_0)(x) \ge e^{-a} s \phi_0(x). \tag{5.13}$$

Thus, combining (5.10) and (5.13), there exist $C_1, C_2 > 0$ such that for any $s \le A_0$,

$$\begin{aligned}
&\Big\langle \int_{\phi_0^{-1}}^{\infty} \Big( e^{-C_1 \theta s \phi_0} - 1 + C_1 \theta s \phi_0 \Big) n(\cdot, \mathrm{d}\theta), \psi_0 \Big\rangle_m \le J_3(s,t) \\
&\le \Big\langle \int_{\phi_0^{-1}}^{\infty} \Big( e^{-C_2 \theta s \phi_0} - 1 + C_2 \theta s \phi_0 \Big) n(\cdot, \mathrm{d}\theta), \psi_0 \Big\rangle_m.
\end{aligned}$$

Note that for any $C>0$, we have

$$\begin{aligned}
&\int_0^{A_0} s^{-2}\mathrm{d}s\int_0^1 e^{\lambda_0 t}\Big\langle\int_{\phi_0^{-1}}^{\infty}\Big(e^{-C\theta s\phi_0}-1+C\theta s\phi_0\Big)n(\cdot,\mathrm{d}\theta),\psi_0\Big\rangle_m\mathrm{d}t\\
&=\int_0^1 e^{\lambda_0 t}\mathrm{d}t\int_E\psi_0(x)m(\mathrm{d}x)\int_1^{\infty}n^{\phi_0}(x,\mathrm{d}\theta)\int_0^{A_0}s^{-2}\Big(e^{-C\theta s}-1+C\theta s\Big)\mathrm{d}s\\
&=\int_0^1 e^{\lambda_0 t}\mathrm{d}t\int_E\psi_0(x)m(\mathrm{d}x)\int_1^{\infty}\theta n^{\phi_0}(x,\mathrm{d}\theta)\int_0^{A_0\theta}s^{-2}\Big(e^{-Cs}-1+Cs\Big)\mathrm{d}s.
\end{aligned}$$

Since

$$\lim_{\theta\to\infty}\frac{\int_0^{A_0\theta}s^{-2}\Big(e^{-Cs}-1+Cs\Big)\mathrm{d}s}{\log\theta}=C,$$

we have

$$\int_0^{A_0}s^{-2}\mathrm{d}s\int_0^1 e^{\lambda_0 t}J_3(s,t)\mathrm{d}t<\infty\Longleftrightarrow\int_E l(x)\psi_0(x)m(\mathrm{d}x)<\infty.$$

Now the conclusion follows immediately. □

Recall that

$$M_\infty=\lim_{t\to\infty}M_t=\lim_{t\to\infty}e^{-\lambda_0 t}\langle\phi_0,X_t\rangle.$$

**Proposition 5.6** *If $\int_E\psi_0(x)l(x)m(\mathrm{d}x)<\infty$, then for any nonzero $\mu\in\mathcal{M}_F(E)$, $M_\infty$ is non-degenerate under $\mathbb{P}_\mu$ and $\mathbb{P}_\mu M_\infty=\langle\phi_0,\mu\rangle$.*

**Proof:** Suppose $\mu\in\mathcal{M}_F(E)$ is nonzero and fixed. For any $\theta>0$, put

$$\Psi_t(\theta,x):=-\log\mathbb{P}_{\delta_x}(\exp\{-\theta M_t\})\qquad\text{and}\quad\Psi(\theta,x):=-\log\mathbb{P}_{\delta_x}(\exp\{-\theta M_\infty\}).$$

Then for any $x\in E$, $\Psi_t(\theta,x)$ is non-increasing in $t$. By the dominated convergence theorem and monotone convergence theorem, we have

$$-\log\mathbb{P}_\mu(\exp\{-\theta M_\infty\})=\lim_{n\to\infty}\langle\Psi_n(\theta,\cdot),\mu\rangle=\langle\Psi(\theta,\cdot),\mu\rangle.\tag{5.14}$$

We claim that there exists some $\theta>0$ such that

$$\langle\Psi(\theta,\cdot),\psi_0\rangle_m>0.\tag{5.15}$$

By the Markov property of $X$, we have

$$\begin{aligned}
&\Psi_{t+s}(\theta,x)=-\log\mathbb{P}_{\delta_x}\mathbb{P}_{X_t}(\exp\{-\theta e^{-\lambda_0 t}M_s\})\\
&=-\log\mathbb{P}_{\delta_x}\exp\{-\langle\Psi_s(\theta e^{-\lambda_0 t}),X_t\rangle\}=V_t(\Psi_s(\theta e^{-\lambda_0 t}))(x).
\end{aligned}\tag{5.16}$$

Letting $s\to\infty$, we get

$$\Psi(\theta e^{\lambda_0 t},x)=V_t(\Psi(\theta))(x).$$

If (5.15) holds, then by (2.3), we get that, for $t > 1$ large enough,

$$\mathbb{P}_{\delta_x}\langle \Psi(\theta), X_t\rangle = T_t\Psi(\theta)(x) > 0,$$

which implies that $\mathbb{P}_{\delta_x}(\langle \Psi(\theta), X_t\rangle > 0) > 0$. Hence we have that, for any $x \in E$, $\Psi(\theta e^{\lambda_0 t}, x) = V_t(\Psi(\theta))(x) > 0$. Thus, by (5.14),

$$\mathbb{P}_\mu \exp\{-\theta e^{\lambda_0 t} M_\infty\} = \exp\{-\langle \Psi(\theta e^{\lambda_0 t}), \mu\rangle\} < 1,$$

which implies that $\mathbb{P}_\mu(M_\infty = 0) < 1$.

Now we prove claim (5.15). Put $R(f)(x) := R_f(1, x)$. It follows from (5.16) and (2.7) that

$$\begin{aligned}
&\langle \Psi_n(\theta, \cdot), \psi_0\rangle_m = \langle V_1(\Psi_{n-1}(\theta e^{-\lambda_0})), \psi_0\rangle_m \\
&= \langle T_1(\Psi_{n-1}(\theta e^{-\lambda_0})), \psi_0\rangle_m - \langle R(\Psi_{n-1}(\theta e^{-\lambda_0})), \psi_0\rangle_m \\
&= e^{\lambda_0}\langle \Psi_{n-1}(\theta e^{-\lambda_0}), \psi_0\rangle_m - \langle R(\Psi_{n-1}(\theta e^{-\lambda_0})), \psi_0\rangle_m \\
&= e^{\lambda_0(n-1)}\langle \Psi_1(\theta e^{-(n-1)\lambda_0}), \psi_0\rangle_m - \sum_{k=1}^{n-1} e^{\lambda_0(k-1)}\langle R(\Psi_{n-k}(\theta e^{-\lambda_0 k})), \psi_0\rangle_m. \qquad (5.17)
\end{aligned}$$

Note that, by Jensen's inequality, we have

$$\Psi_{n-k}(\theta e^{-\lambda_0 k})(x) = -\log \mathbb{P}_{\delta_x} \exp\{-\theta e^{-\lambda_0 k} M_{n-k}\} \le \mathbb{P}_{\delta_x} \theta e^{-\lambda_0 k} M_{n-k} = \theta e^{-\lambda_0 k}\phi_0(x).$$

By the dominated convergence theorem, we get

$$\lim_{n\to\infty} e^{\lambda_0(n-1)}\langle \Psi_1(\theta e^{-(n-1)\lambda_0}), \psi_0\rangle_m = \theta\langle \mathbb{P}_{\delta\cdot} M_1, \psi_0\rangle_m = \theta.$$

Thus we have

$$\langle \Psi(\theta, \cdot), \psi_0\rangle_m = \lim_{n\to\infty}\langle \Psi_n(\theta, \cdot), \psi_0\rangle_m \ge \theta - \lim_{n\to\infty}\sum_{k=1}^{n-1} e^{\lambda_0(k-1)}\langle R(\theta e^{-\lambda_0 k}\phi_0), \psi_0\rangle_m. \qquad (5.18)$$

Since $t \to \langle R(\theta e^{-\lambda_0 t}\phi_0), \psi_0\rangle_m$ is decreasing, we have

$$\begin{aligned}
&\sum_{k=1}^{\infty} e^{\lambda_0(k-1)}\langle R(\theta e^{-\lambda_0 k}\phi_0), \psi_0\rangle_m \\
&\le \int_0^\infty e^{\lambda_0 t}\langle R(\theta e^{-\lambda_0 t}\phi_0), \psi_0\rangle_m \, \mathrm{d}t \\
&= \lambda_0^{-1}\int_0^1 s^{-2}\langle R(s\theta\phi_0), \psi_0\rangle_m \, \mathrm{d}s \\
&\le \lambda_0^{-1}\int_0^1 s^{-2}\langle T_1(s\theta\phi_0) - (1 - \exp\{-V_1(s\theta\phi_0)\}), \psi_0\rangle_m \, \mathrm{d}s
\end{aligned}$$

$$= \lambda_0^{-1} \int_E \psi_0(x) m(\mathrm{d}x) \mathbb{P}_{\delta_x} \int_0^1 s^{-2} \Big( s\theta \langle \phi_0, X_1 \rangle - 1 + \exp\{-s\theta \langle \phi_0, X_1 \rangle\} \Big) \, \mathrm{d}s.$$

Since $e^{-s} - 1 + s \le s \wedge (s^2/2)$, there exists $C > 0$ such that for any $r \ge 0$,

$$\begin{aligned} &\int_0^1 s^{-2} \Big( rs - 1 + \exp\{-rs\} \Big) \, \mathrm{d}s = r \int_0^r s^{-2} \Big( s - 1 + \exp\{-s\} \Big) \, \mathrm{d}s \\ &\le \frac{1}{2} r^2 \mathrm{I}_{r \le 2} + r \left( 1 + \int_2^r s^{-1} \, \mathrm{d}s \right) \mathrm{I}_{r > 2} \le \frac{1}{2} r^2 \mathrm{I}_{r \le 2} + Cr \log r \mathrm{I}_{r > 2}. \end{aligned}$$

Thus,

$$\begin{aligned} &\sum_{k=1}^{\infty} e^{\lambda_0 (k-1)} \langle R(\theta e^{-\lambda_0 k} \phi_0), \psi_0 \rangle_m \\ &\le \theta \lambda_0^{-1} / 2 \Big( \int_E \mathbb{P}_{\delta_x} (\theta \langle \phi_0, X_1 \rangle^2; \theta \langle \phi_0, X_1 \rangle \le 2) \psi_0(x) m(\mathrm{d}x) \\ &\qquad + 2C \int_E \mathbb{P}_{\delta_x} (\langle \phi_0, X_1 \rangle \log(\theta \langle \phi_0, X_1 \rangle); \theta \langle \phi_0, X_1 \rangle > 2) \psi_0(x) m(\mathrm{d}x) \Big). \end{aligned}$$

Using the dominated convergence theorem, we get

$$\lim_{\theta \to 0} \int_E \mathbb{P}_{\delta_x} (\theta \langle \phi_0, X_1 \rangle^2; \theta \langle \phi_0, X_1 \rangle \le 2) \psi_0(x) m(\mathrm{d}x) = 0.$$

By Lemma (5.5), we have

$$\int_E \mathbb{P}_{\delta_x} (\langle \phi_0, X_1 \rangle \log(\langle \phi_0, X_1 \rangle); \langle \phi_0, X_1 \rangle > 2) \psi_0(x) m(\mathrm{d}x) < \infty.$$

Applying the dominated convergence theorem again, we get

$$\lim_{\theta \to 0} \int_E \mathbb{P}_{\delta_x} (\langle \phi_0, X_1 \rangle \log(\theta \langle \phi_0, X_1 \rangle); \theta \langle \phi_0, X_1 \rangle > 2) \psi_0(x) m(\mathrm{d}x) = 0.$$

Therefore, there exists $\theta_0 > 0$ such that for any $\theta \in (0, \theta_0]$,

$$\sum_{k=1}^{\infty} e^{\lambda_0 (k-1)} \langle R(\theta e^{-\lambda_0 k} \phi_0), \psi_0 \rangle_m \le \theta / 2.$$

It follows from (5.18) that, for $0 < \theta \le \theta_0$,

$$\langle \Psi(\theta, \cdot), \psi_0 \rangle_m \ge \theta / 2 > 0.$$

Now the claim (5.15) is proved, and hence $M_\infty$ is non-degenerate.

It is easy to see that

$$M_\infty = \lim_{t \to \infty} e^{-\lambda_0 t} \langle \phi_0, X_t \rangle = \lim_{t \to \infty} e^{-\lambda_0 t} (\gamma_t)^{-1} \gamma_t \langle \phi_0, X_t \rangle = l_0^{-1} W. \tag{5.19}$$

Since $M_\infty$ is non-degenerate, we have $l_0 < \infty$. Thus by (5.6), $\mathbb{P}_\mu M_\infty = \langle \phi_0, \mu \rangle$. The proof is now complete. □

**Proof of Theorem 1.2:** By (5.19), (1) $\Longleftrightarrow$ (2) $\Longleftrightarrow$ (3). By Proposition 5.4 and Proposition 5.6 , (3) $\Longleftrightarrow$ (5) $\Longleftrightarrow$ (6) and (4) $\Longrightarrow$ (2). Thus, we only need to show that (3) $\Longrightarrow$ (4).

By (2.11) and the fact that $\eta_s = V_t(\eta_{t+s})$, we have

$$\eta_0(x) = V_t(\eta_t)(x) = T_t(\eta_t)(x) - \int_0^t T_s\left[r(\cdot, \eta_s(\cdot))\right](x)\mathrm{d}s.$$

Thus,

$$e^{\lambda_0 t}\gamma_t = \langle T_t(\eta_t), \psi_0 \rangle_m = \gamma_0 + \int_E \psi_0(x) m(\mathrm{d}x) \int_0^t e^{\lambda_0 s} r(x, \eta_s(x))\mathrm{d}s,$$

which implies that

$$l_0 = \gamma_0 + \int_E \psi_0(x) m(\mathrm{d}x) \int_0^\infty e^{\lambda_0 s} r(x, \eta_s(x))\mathrm{d}s. \tag{5.20}$$

Recall that,

$$r(x,s) = \beta(x)s^2 + \int_0^\infty (e^{-s\theta} - 1 + s\theta)\, n(x, \mathrm{d}\theta).$$

Hence,

$$\begin{aligned}
&\int_0^\infty e^{\lambda_0 s} r(x, \eta_s(x))\mathrm{d}s \geq \int_0^\infty e^{\lambda_0 s}\mathrm{d}s \int_{e^{\lambda_0 s}\phi_0(x)^{-1}}^\infty (e^{-\theta\eta_s(x)} - 1 + \theta\eta_s(x))\, n(x, \mathrm{d}\theta) \\
&= \int_{\phi_0(x)^{-1}}^\infty n(x, \mathrm{d}\theta) \int_0^{(\log(\theta\phi_0(x))/\lambda_0} e^{\lambda_0 s}(e^{-\theta\eta_s(x)} - 1 + \theta\eta_s(x))\mathrm{d}s.
\end{aligned} \tag{5.21}$$

Choose $s_0 > 1$ large enough such that $(1 - ce^{-\gamma s_0}) > 0$, where $c = c(1)$ and $\gamma = \gamma(1)$ are the constants in (2.3). By (2.3) and Lemma 4.4, we get that,

$$\eta_s(x) = V_{s_0}(\eta_{s+s_0})(x) \geq e^{-as_0} T_{s_0}(\eta_{s+s_0})(x) \geq e^{-as_0}(1 - ce^{-\gamma s_0}) e^{\lambda_0 s_0}\gamma_{s+s_0}\phi_0(x) \geq C\gamma_{s+s_0}\phi_0(x).$$

Thus, by Remark 4.7, for any $s \leq (\log(\theta\phi_0(x))/\lambda_0$, we have

$$\theta\eta_s(x) \geq C\theta\gamma_{s+s_0}\phi_0(x) \geq Ce^{\lambda_0 s}\gamma_{s+s_0} \geq CL(s+s_0) \geq CL(0) > 0,$$

where $L(t) = e^{\lambda_0 t}\gamma_t$. Therefore,

$$\inf_{s \leq (\log(\theta\phi_0(x))/\lambda_0} \frac{e^{-\theta\eta_s(x)} - 1 + \theta\eta_s(x)}{\theta\eta_s(x)} \geq \inf_{r \geq CL(0)} \frac{e^{-r} - 1 + r}{r} \geq c,$$

for some constant $c > 0$. It follows that

$$\int_{\phi_0(x)^{-1}}^\infty n(x, \mathrm{d}\theta) \int_0^{(\log(\theta\phi_0(x)))/\lambda_0} e^{\lambda_0 s}(e^{-\theta\eta_s(x)} - 1 + \theta\eta_s(x))\mathrm{d}s$$

$$
\begin{aligned}
&\geq c\int_{\phi_0(x)^{-1}}^{\infty}\theta n(x,\mathrm{d}\theta)\int_0^{\log(\theta\phi_0(x))/\lambda_0}e^{\lambda_0 s}\eta_s(x)\mathrm{d}s\\
&\geq C\int_{\phi_0(x)^{-1}}^{\infty}\theta n(x,\mathrm{d}\theta)\int_0^{\log(\theta\phi_0(x))/\lambda_0}e^{\lambda_0 s}\gamma_{s+s_0}\mathrm{d}s\phi_0(x)\\
&\geq C\int_1^{\infty}\theta n^{\phi_0}(x,\mathrm{d}\theta)\int_0^{\log\theta/\lambda_0}L(s+s_0)\mathrm{d}s\\
&\geq CL(0)\int_1^{\infty}\theta\log\theta n^{\phi_0}(x,\mathrm{d}\theta)\geq Cl(x). \qquad (5.22)
\end{aligned}
$$

Combining (5.20),(5.21) and (5.22), we get that $l_0 \geq C\langle l,\psi_0\rangle_m$. Thus (3) $\Longrightarrow$ (4).

The proof is now complete. □

# 6 Strong convergence with general test functions

In this section, we fix a non-trivial family $(\eta_t : t \geq 0)$ of backward iterates and let $\gamma_t := \langle \eta_t, \psi_0\rangle_m$. The goal of this section is to determine the almost sure limit of $\gamma_t\langle f, X_t\rangle$ for general test functions $f$.

## 6.1 The martingale problems of superprocesses

In this subsection, we recall the martingale problem for the superprocess $X$. Let $J$ denote the set of jump times of $X$, i.e.,

$$J := \{s \geq 0 : \triangle X_s = X_s - X_{s-} \neq 0\}.$$

Since $X$ is a càdlàg process in $\mathcal{M}_F(E)$, $J$ is a countable set. Let $N(\mathrm{d}s,\mathrm{d}\nu)$ be the optional random measure on $[0,\infty)\times\mathcal{M}_F(E)$ defined by

$$N(\mathrm{d}s,\mathrm{d}\nu) := \sum_{s\in J}\delta_{(s,\triangle X_s)}(\mathrm{d}s,\mathrm{d}\nu),$$

and $\widehat{N}(\mathrm{d}s,\mathrm{d}\nu)$ be the predictable compensator of $N(\mathrm{d}s,\mathrm{d}\nu)$ which satisfies that for any nonnegative predictable function $F$ on $\mathbb{R}_+\times\mathcal{M}_F(E)\times\Omega$,

$$\int_0^t\int_{\mathcal{M}_F(E)}F(s,\nu)\,\widehat{N}(\mathrm{d}s,\mathrm{d}\nu) = \int_0^t\mathrm{d}s\int_E X_s(\mathrm{d}x)\int_0^{\infty}F(s,\theta\delta_x)n(x,\mathrm{d}\theta), \qquad (6.1)$$

where $n$ is the kernel in the branching mechanism $\varphi$. Define

$$\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu) := N(\mathrm{d}s,\mathrm{d}\nu) - \widehat{N}(\mathrm{d}s,\mathrm{d}\nu).$$

Then $\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu)$ is a martingale measure. The "stochastic integral"

$$\int_0^t\int_{\mathcal{M}_F(E)}F(s,\nu)\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu)$$

can be defined for a wide class of Borel functions $F$ on $[0,t]\times\mathcal{M}_F(E)$. In particular, if $f$ is a bounded Borel function on $[0,t]\times E$ and $F_f(s,\nu):=\int_E f(s,x)\nu(\mathrm{d}x)$, then the integral $\int_0^t\int_{\mathcal{M}_F(E)}F_f(s,\nu)\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu)$ is well-defined. Let $\mathcal{L}_N^2$ be the space of predictable processes $(F(s,\nu):s>0,\nu\in\mathcal{M}_F(E))$ satisfying, for all $\mu\in\mathcal{M}_F(E)$,

$$\mathbb{P}_\mu\int_0^t\mathrm{d}s\int_E X_s(\mathrm{d}x)\int_0^\infty F(s,\theta\delta_x)^2n(x,\mathrm{d}\theta)<\infty.$$

For any $F\in\mathcal{L}_N^2$,

$$M_t^d(F):=\int_0^t\int_{\mathcal{M}_F(E)}F(s,\nu)\,\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu),\qquad t\geq 0,$$

is a square integrable martingale such that

$$\mathbb{P}_\mu(M_t^d(F)^2)=\mathbb{P}_\mu\left[\int_0^t\mathrm{d}s\int_E X_s(\mathrm{d}x)\int_0^\infty F(s,\theta\delta_x)^2n(x,\mathrm{d}\theta)\right].\tag{6.2}$$

Note that $C_0(E)$ is a Banach space under the supremum norm. In the remainder of this paper, we assume that

**Assumption 5** **(i)** *$\{P_t,t\geq 0\}$ is a Feller semigroup, that is, $\{P_t,t\geq 0\}$ preserves $C_0(E)$ and $[0,\infty)\ni t\to P_tf\in C_0(E)$ is continuous for every $f\in C_0(E)$.*

**(ii)**

$$\lim_{a\to\infty}\sup_{x\in E}\int_a^\infty\theta n(x,\mathrm{d}\theta)=0.\tag{6.3}$$

In the reminder of this subsection, we will (also) use $\widetilde{L}$ to denote the infinitesimal generator of $\{P_t,t\geq 0\}$ in the space $C_0(E)$ and use $\mathrm{Dom}(\widetilde{L})$ its domain. It is known (see [5, Section 6.1], for instance) that, for any $f\in\mathrm{Dom}(\widetilde{L})$, we have that

$$\langle f,X_t\rangle=\langle f,X_0\rangle+\int_0^t\langle\widetilde{L}f-\alpha f,X_s\rangle\mathrm{d}s+M_t^c(f)+M_t^d(f),$$

where $M_t^c(f)$ is a continuous local martingale with quadratic variation $\int_0^t 2\langle\beta f^2,X_s\rangle\mathrm{d}s$ and

$$M_t^d(f)=\int_0^t\int_{\mathcal{M}_F(E)}\langle f,\nu\rangle\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu)$$

is a purely discontinuous local martingale. Here we remark that if we assume $\alpha,\beta\in C(E)$ and that $x\to(\theta\wedge\theta^2)n(x,\mathrm{d}\theta)$ is continuous in the topology of weak convergence, then the above result follows from [15, Theorem 7.25]. $M_t^c(f)$ induces a worthy $(\mathcal{G}_t)$-martingale measure $S^C(\mathrm{d}s,\mathrm{d}x)$ (see [15, Section 7.3] for the definition of worthy martingale measure) satisfying

$$M_t^c(f)=\int_0^t\int_E f(x)S^C(\mathrm{d}s,\mathrm{d}x).$$

Standard arguments then show that the "stochastic integral"

$$\int_0^t \int_E f(s,x) S^C(\mathrm{d}s, \mathrm{d}x)$$

can be defined for a wide class of integrands $f$ on $[0,t] \times E$ (see, for example, [15, Theorem 7.25] or [7] for more details). Thus, one can show that (see [7, Proposition 2.13] or [19, Exercise II.5.2] for the case when the branching mechanism has finite variance) for any bounded function $g$ on $E$,

$$\langle g, X_t \rangle = \langle T_t g, X_0 \rangle + \int_0^t \int_{\mathcal{M}_F(E)} \langle T_{t-s} g, \nu \rangle \widetilde{N}(\mathrm{d}s, \mathrm{d}\nu) + \int_0^t \int_E T_{t-s} g(x) S^C(\mathrm{d}s, \mathrm{d}x). \tag{6.4}$$

## 6.2 Discrete times

In this subsection, we show that for any $\delta > 0$ and $f \in \mathcal{B}_b^+(E)$, $\gamma_{n\delta} \langle f\phi_0, X_{n\delta} \rangle$ has an almost sure limit as $n \to \infty$. We will extend this result to continuous times in two different scenarios in the next two subsections.

**Theorem 6.1** *For any $\delta > 0$, $\mu \in \mathcal{M}_F(E)$ and $f \in \mathcal{B}_b^+(E)$, we have*

$$\lim_{n\to\infty} \gamma_{n\delta} \langle \phi_0 f, X_{n\delta} \rangle = \langle f\phi_0, \psi_0 \rangle_m W, \qquad a.s.\text{-}\mathbb{P}_\mu.$$

To prove Theorem 6.1, we first make some preparations. For any $s > 0$, we define

$$\mathcal{D}_{<1}(s) := \{\nu \in \mathcal{M}_F(E) : 0 < \gamma_s \langle \phi_0, \nu \rangle < 1\} \tag{6.5}$$

and

$$\mathcal{D}_{\geq 1}(s) := \{\nu \in \mathcal{M}_F(E) : \gamma_s \langle \phi_0, \nu \rangle \geq 1\}. \tag{6.6}$$

For any $m \in \mathbb{N}$, $\delta > 0$, $\mu \in \mathcal{M}_F(E)$ and $f \in \mathcal{B}_b^+(E)$, by (6.4), we have

$$\begin{aligned}
&\gamma_{(n+m)\delta} \langle \phi_0 f, X_{(n+m)\delta} \rangle \\
&= \gamma_{(n+m)\delta} \langle T_{(n+m)\delta}(\phi_0 f), \mu \rangle + \gamma_{(n+m)\delta} \int_0^{(n+m)\delta} \int_E T_{(n+m)\delta - s}(\phi_0 f)(x) S^C(\mathrm{d}s, \mathrm{d}x) \\
&\quad + \gamma_{(n+m)\delta} \int_0^{(n+m)\delta} \int_{\mathcal{D}_{<1}(s)} \langle T_{(n+m)\delta - s}(\phi_0 f), \nu \rangle \widetilde{N}(\mathrm{d}s, \mathrm{d}\nu) \\
&\quad + \gamma_{(n+m)\delta} \int_0^{(n+m)\delta} \int_{\mathcal{D}_{\geq 1}(s)} \langle T_{(n+m)\delta - s}(\phi_0 f), \nu \rangle \widetilde{N}(\mathrm{d}s, \mathrm{d}\nu) \\
&=: \gamma_{(n+m)\delta} \langle T_{(n+m)\delta}(\phi_0 f), \mu \rangle + C_{(n+m)\delta}(f) + H_{(n+m)\delta}(f) + L_{(n+m)\delta}(f).
\end{aligned}$$

Therefore,

$$\gamma_{(n+m)\delta} \langle \phi_0 f, X_{(n+m)\delta} \rangle - \mathbb{P}_\mu [\gamma_{(n+m)\delta} \langle \phi_0 f, X_{(n+m)\delta} \rangle | \mathcal{G}_{n\delta}]$$

$$= \big(H_{(n+m)\delta}(f) - \mathbb{P}_\mu(H_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})\big) + \big(L_{(n+m)\delta}(f) - \mathbb{P}_\mu(L_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})\big)$$
$$+ \big(C_{(n+m)\delta}(f) - \mathbb{P}_\mu(C_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})\big).$$

We now deal with the three parts separately. Before doing this, we prove a lemma first.

**Lemma 6.2** *If $\{a_n : n \geq 1\}$ is a sequence of positive numbers such that $\lim_{n\to\infty} a_{n+1}/a_n = a > 1$, then*

$$\sup_{x\in E} \sum_{n=1}^{\infty} a_n^{-1} \int_0^{a_n} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta) < \infty \tag{6.7}$$

*and*

$$\sup_{x\in E} \sum_{n=1}^{\infty} a_n \int_{a_n}^{\infty} n^{\phi_0}(x, \mathrm{d}\theta) < \infty. \tag{6.8}$$

**Proof:** Since $\lim_{n\to\infty} a_{n+1}/a_n = a > 1$, for any $a^* \in (1, a)$, there exists $K > 0$ such that for any $n \geq K$,

$$\frac{a_{n+1}}{a_n} > a^*.$$

Without loss of generality, we assume that $a_n \uparrow \infty$. For convenience, we put $a_0 = 0$. Then we have

$$\sum_{n=1}^{\infty} a_n^{-1} \int_0^{a_n} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta) = \sum_{n=1}^{\infty} a_n^{-1} \sum_{k=1}^{n} \int_{a_{k-1}}^{a_k} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta)$$
$$= \sum_{k=1}^{\infty} \int_{a_{k-1}}^{a_k} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta) \sum_{n=k}^{\infty} a_n^{-1}$$
$$\leq \sum_{k=1}^{K} \int_{a_{k-1}}^{a_k} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta) \sum_{n=k}^{\infty} a_n^{-1} + \sum_{k=K+1}^{\infty} \int_{a_{k-1}}^{a_k} \theta n^{\phi_0}(x, \mathrm{d}\theta) a_k \sum_{n=k}^{\infty} a_n^{-1}$$
$$\leq \int_0^{a_K} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta) \sum_{n=1}^{\infty} a_n^{-1} + \sum_{k=K+1}^{\infty} \int_{a_{k-1}}^{a_k} \theta n^{\phi_0}(x, \mathrm{d}\theta) a_k \sum_{n=k}^{\infty} a_n^{-1}.$$

For any $k > K$, we have

$$\sum_{n=k}^{\infty} a_k a_n^{-1} \leq \sum_{n=k}^{\infty} (a^*)^{-(n-k)} = \sum_{n=1}^{\infty} (a^*)^{-n} < \infty.$$

Therefore,

$$\sum_{n=1}^{\infty} a_n^{-1} \int_0^{a_n} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta) \leq C \left[ \int_0^{a_K} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta) + \int_{a_K}^{\infty} \theta n^{\phi_0}(x, \mathrm{d}\theta) \right]$$
$$\leq C \sup_{x\in E} \int_0^{\infty} (\theta \wedge \theta^2) n^{\phi_0}(x, \mathrm{d}\theta) < \infty.$$

Note that

$$\sum_{n=1}^{\infty} a_n \int_{a_n}^{\infty} n^{\phi_0}(x, \mathrm{d}\theta) = \sum_{n=1}^{\infty} a_n \sum_{k=n}^{\infty} \int_{a_k}^{a_{k+1}} n^{\phi_0}(x, \mathrm{d}\theta)$$
$$\leq \sum_{k=1}^{\infty} \int_{a_k}^{a_{k+1}} \theta n^{\phi_0}(x, \mathrm{d}\theta) \Big( a_k^{-1} \sum_{n=1}^{k} a_n \Big).$$

Using elementary calculus, one can easily show that

$$\lim_{k\to\infty} a_k^{-1} \sum_{n=1}^{k} a_n = \lim_{k\to\infty} \frac{a_{k+1}}{a_{k+1} - a_k} = \frac{a}{a-1}.$$

Thus $\sup_{k\geq 1} a_k^{-1} \sum_{n=1}^{k} a_n < \infty$. It follows that

$$\sup_{x\in E} \sum_{n=1}^{\infty} a_n \int_{a_n}^{\infty} n^{\phi_0}(x, \mathrm{d}\theta) \leq C \sup_{x\in E} \int_{a_1}^{\infty} \theta n^{\phi_0}(x, \mathrm{d}\theta) < \infty.$$

The proof is now complete.

□

Define

$$I(a, x) := \int_0^a \theta^2 n^{\phi_0}(x, \mathrm{d}\theta).$$

**Lemma 6.3** *For any $m \in \mathbb{N}$, $\delta > 0$, $\mu \in \mathcal{M}_F(E)$ and $f \in \mathcal{B}_b^+(E)$, we have*

$$\lim_{n\to\infty} H_{(n+m)\delta}(f) - \mathbb{P}_\mu(H_{(n+m)\delta}(f)|\mathcal{G}_{n\delta}) = 0, \qquad a.s.\text{-}\mathbb{P}_\mu.$$

**Proof:** By the conditional Borel-Cantelli lemma, it suffices to prove that

$$\sum_{n=1}^{\infty} \mathbb{P}_\mu \left( [H_{(n+m)\delta}(f) - \mathbb{P}_\mu(H_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})]^2 | \mathcal{G}_{(n-1)\delta} \right) < \infty. \tag{6.9}$$

Recall from (6.5) that

$$\mathcal{D}_{<1}(s) := \{\nu \in \mathcal{M}_F(E) : 0 < \gamma_s \langle \phi_0, \nu \rangle < 1\}.$$

Since $\widetilde{N}(\mathrm{d}s, \mathrm{d}\nu)$ is a martingale measure, we have

$$\mathbb{P}_\mu(H_{(n+m)\delta}(f)|\mathcal{G}_{n\delta}) = \gamma_{(n+m)\delta} \int_0^{n\delta} \int_{\mathcal{D}_{<1}(s)} \langle T_{(n+m)\delta-s}(\phi_0 f), \nu \rangle \widetilde{N}(\mathrm{d}s, \mathrm{d}\nu),$$

which implies that

$$H_{(n+m)\delta}(f) - \mathbb{P}_\mu(H_{(n+m)\delta}(f)|\mathcal{G}_{n\delta}) = \gamma_{(n+m)\delta} \int_{n\delta}^{(n+m)\delta} \int_{\mathcal{D}_{<1}(s)} \langle T_{(n+m)\delta-s}(\phi_0 f), \nu \rangle \widetilde{N}(\mathrm{d}s, \mathrm{d}\nu). \tag{6.10}$$

By (5.4), $\gamma_t = \langle \Phi(e^{-\lambda_0 t}, \cdot), \psi_0 \rangle_m$, which implies that $t \to \gamma_t$ is non-increasing. Thus by (6.2) and (2.3), we have

$$\begin{aligned}
&\mathbb{P}_\mu \left( [H_{(n+m)\delta}(f) - \mathbb{P}_\mu(H_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})]^2 | \mathcal{G}_{(n-1)\delta} \right) \\
&= \gamma^2_{(n+m)\delta} \mathbb{P}_{X_{(n-1)\delta}} \left[ \int_\delta^{(1+m)\delta} \mathrm{d}s \int_E X_s(\mathrm{d}x) \int_0^{\gamma^{-1}_{s+(n-1)\delta}\phi_0(x)^{-1}} \theta^2 [T_{(1+m)\delta - s}(\phi_0 f)(x)]^2 n(x, \mathrm{d}\theta) \right] \\
&\le \|f\|^2_\infty \gamma^2_{(n+m)\delta} \mathbb{P}_{X_{(n-1)\delta}} \left[ \int_\delta^{(1+m)\delta} e^{2\lambda_0((1+m)\delta - s)} \mathrm{d}s \int_E X_s(\mathrm{d}x) \int_0^{\gamma^{-1}_{(n+m)\delta}} \theta^2 n^{\phi_0}(x, \mathrm{d}\theta) \right] \\
&= \|f\|^2_\infty \gamma^2_{(n+m)\delta} \int_\delta^{(1+m)\delta} e^{2\lambda_0((1+m)\delta - s)} \langle T_s[I(\gamma^{-1}_{(n+m)\delta})], X_{(n-1)\delta} \rangle \mathrm{d}s \\
&\le (1 + c(\delta)) \|f\|^2_\infty e^{2\lambda_0(1+m)\delta} \int_\delta^{(1+m)\delta} e^{-\lambda_0 s} \mathrm{d}s \gamma^2_{(n+m)\delta} \langle I(\gamma^{-1}_{(n+m)\delta}), \psi_0 \rangle_m \langle \phi_0, X_{(n-1)\delta} \rangle \\
&\le C \left[ \gamma_{(n+m)\delta} \langle \phi_0, X_{(n-1)\delta} \rangle \right] \gamma_{(n+m)\delta} \langle I(\gamma^{-1}_{(m+n)\delta}), \psi_0 \rangle_m. \qquad (6.11)
\end{aligned}$$

It follows from the fact that $\lim_{n\to\infty} \frac{\gamma^{-1}_{(n+m)\delta}}{\gamma^{-1}_{(n+m-1)\delta}} = e^{\lambda_0 \delta}$ and Lemma 6.2 that

$$\sum_{n=1}^\infty \gamma_{(n+m)\delta} \langle I(\gamma^{-1}_{(m+n)\delta}), \psi_0 \rangle_m < \infty. \qquad (6.12)$$

Since $\lim_{n\to\infty} \gamma_{(n+m)\delta} \langle \phi_0, X_{(n-1)\delta} \rangle = e^{-\lambda_0 (m+1)\delta} W$, combining (6.11) and (6.12), (6.9) follows immediately. The proof is now complete. □

**Lemma 6.4** *For any $m \in \mathbb{N}$, $\delta > 0$, $\mu \in \mathcal{M}_F(E)$ and $f \in \mathcal{B}_b^+(E)$, we have*

$$\lim_{n\to\infty} L_{(n+m)\delta}(f) - \mathbb{P}_\mu(L_{(n+m)\delta}(f)|\mathcal{G}_{n\delta}) = 0, \qquad a.s.\text{-}\mathbb{P}_\mu.$$

**Proof:** Recall the definition of $\mathcal{D}_{\ge 1}(s)$ in (6.6):

$$\mathcal{D}_{\ge 1}(s) = \{\nu \in \mathcal{M}_F(E) : \gamma_s \langle \phi_0, \nu \rangle \ge 1\}.$$

Since $\widetilde{N}(\mathrm{d}s, \mathrm{d}\nu)$ is a martingale measure, we have

$$\mathbb{P}_\mu(L_{(n+m)\delta}(f)|\mathcal{G}_{n\delta}) = \gamma_{(n+m)\delta} \int_0^{n\delta} \int_{\mathcal{D}_{\ge 1}(s)} \langle T_{(n+m)\delta - s}(\phi_0 f), \nu \rangle \widetilde{N}(\mathrm{d}s, \mathrm{d}\nu),$$

which implies that

$$\begin{aligned}
&L_{(n+m)\delta}(f) - \mathbb{P}_\mu(L_{(n+m)\delta}(f)|\mathcal{G}_{n\delta}) \\
&= \gamma_{(n+m)\delta} \int_{n\delta}^{(n+m)\delta} \int_{\mathcal{D}_{\ge 1}(s)} \langle T_{(n+m)\delta - s}(\phi_0 f), \nu \rangle \Big( N(\mathrm{d}s, \mathrm{d}\nu) - \widehat{N}(\mathrm{d}s, \mathrm{d}\nu \Big).
\end{aligned}$$

We claim that

$$\mathbb{P}_\mu\Big(\int_{n\delta}^{(n+m)\delta}\int_{\mathcal{D}_{\geq 1}(s)} N(\mathrm{d}s,\mathrm{d}\nu)>0,\ \text{i.o.}\Big)=0. \tag{6.13}$$

In fact, since $\int_{n\delta}^{(n+m)\delta}\int_{\mathcal{D}_{\geq 1}(s)} N(\mathrm{d}s,\mathrm{d}\nu)$ is a non-negative integer, by the Markov property of $X$,

$$\begin{aligned}
&\sum_{n=1}^{\infty}\mathbb{P}_\mu\Big(\int_{n\delta}^{(n+m)\delta}\int_{\mathcal{D}_{\geq 1}(s)} N(\mathrm{d}s,\mathrm{d}\nu)>0\Big|\mathcal{G}_{(n-1)\delta}\Big)\\
&\leq \sum_{n=1}^{\infty}\mathbb{P}_\mu\Big(\int_{n\delta}^{(n+m)\delta}\int_{\mathcal{D}_{\geq 1}(s)} N(\mathrm{d}s,\mathrm{d}\nu)\Big|\mathcal{G}_{(n-1)\delta}\Big)\\
&= \sum_{n=1}^{\infty}\mathbb{P}_{X_{(n-1)\delta}}\Big(\int_{\delta}^{(1+m)\delta}\int_{\mathcal{D}_{\geq 1}(s+(n-1)\delta)} N(\mathrm{d}s,\mathrm{d}\nu)\Big)\\
&= \sum_{n=1}^{\infty}\mathbb{P}_{X_{(n-1)\delta}}\left[\int_{\delta}^{(1+m)\delta} ds\int_E X_s(\mathrm{d}x)\int_{\phi_0(x)^{-1}\gamma^{-1}_{s+(n-1)\delta}}^{\infty} n(x,\mathrm{d}\theta)\right]\\
&\leq \sum_{n=1}^{\infty}\int_{\delta}^{(1+m)\delta}\mathrm{d}s\left\langle T_s\Big[\int_{\gamma^{-1}_{n\delta}}^{\infty} n^{\phi_0}(\cdot,\mathrm{d}\theta)\Big], X_{(n-1)\delta}\right\rangle\\
&\leq (1+c(\delta))m\delta e^{\lambda_0(m+1)\delta}\sum_{n=1}^{\infty}\left\langle\int_{\gamma^{-1}_{n\delta}}^{\infty} n^{\phi_0}(\cdot,\mathrm{d}\theta),\psi_0\right\rangle_m\langle\phi_0,X_{(n-1)\delta}\rangle,
\end{aligned} \tag{6.14}$$

where in the second to the last inequality, we use the fact that $\gamma_{s+(n-1)\delta}\leq\gamma_{n\delta}$, and the last inequality follows from (2.3). It follows from (6.8) that

$$\sum_{n=1}^{\infty}\gamma^{-1}_{n\delta}\langle\int_{\gamma^{-1}_{n\delta}}^{\infty} n^{\phi_0}(\cdot,\mathrm{d}\theta),\psi_0\rangle_m<\infty. \tag{6.15}$$

By Theorem 5.2, $\gamma_{n\delta}\langle\phi_0,X_{(n-1)\delta}\rangle\to e^{-\lambda_0\delta}W$ as $n\to\infty$. Therefore we have

$$\sum_{n=1}^{\infty}\langle\int_{\gamma^{-1}_{n\delta}}^{\infty} n^{\phi_0}(\cdot,\mathrm{d}\theta),\psi_0\rangle_m\langle\phi_0,X_{(n-1)\delta}\rangle<\infty,$$

which implies that

$$\sum_{n=1}^{\infty}\mathbb{P}_\mu\Big(\int_{n\delta}^{(n+m)\delta}\int_{\mathcal{D}_{\geq 1}(s)} N(\mathrm{d}s,\mathrm{d}\nu)>0\Big|\mathcal{G}_{(n-1)\delta}\Big)<\infty.$$

Now using the the conditional Borel-Cantelli lemma, we immediately get the claim (6.13).

By (6.13), we get

$$\lim_{n\to\infty}\gamma_{(n+m)\delta}\int_{n\delta}^{(n+m)\delta}\int_{\mathcal{D}_{\geq 1}(s)}\langle T_{(n+m)\delta-s}(\phi_0 f),\nu\rangle N(\mathrm{d}s,\mathrm{d}\nu)=0,\qquad \mathbb{P}_\mu\text{-a.s.} \tag{6.16}$$

To complete the proof, we only need to show that

$$\lim_{n\to\infty}\gamma_{(n+m)\delta}\int_{n\delta}^{(n+m)\delta}\int_{\mathcal{D}_{\geq 1}(s)}\langle T_{(n+m)\delta-s}(\phi_0 f),\nu\rangle\widehat{N}(\mathrm{d}s,\mathrm{d}\nu)=0,\qquad \mathbb{P}_\mu\text{-a.s.}\tag{6.17}$$

By (6.1), we have

$$\begin{aligned}
&\gamma_{(n+m)\delta}\int_{n\delta}^{(n+m)\delta}\int_{\mathcal{D}_{\geq 1}(s)}\langle T_{(n+m)\delta-s}(\phi_0 f),\nu\rangle\widehat{N}(\mathrm{d}s,\mathrm{d}\nu)\\
&=\gamma_{(n+m)\delta}\int_{n\delta}^{(n+m)\delta}\mathrm{d}s\int_E T_{(n+m)\delta-s}(\phi_0 f)(x)X_s(\mathrm{d}x)\int_{\phi_0(x)^{-1}\gamma_s^{-1}}^{\infty}\theta n(x,\mathrm{d}\theta)\\
&\leq\|f\|_\infty e^{\lambda_0 m\delta}\gamma_{(n+m)\delta}\int_{n\delta}^{(n+m)\delta}\mathrm{d}s\int_E\phi_0(x)X_s(\mathrm{d}x)\int_{\phi_0(x)^{-1}\gamma_s^{-1}}^{\infty}\theta n(x,\mathrm{d}\theta)\\
&\leq\|f\|_\infty e^{\lambda_0 m\delta}\gamma_{(n+m)\delta}\int_{n\delta}^{(n+m)\delta}\langle\phi_0,X_s\rangle\mathrm{d}s\sup_{x\in E}\left(\int_{\|\phi_0\|_\infty^{-1}\gamma_{n\delta}^{-1}}^{\infty}\theta n(x,\mathrm{d}\theta)\right).
\end{aligned}$$

Note that

$$\begin{aligned}
&\lim_{n\to\infty}\gamma_{(n+m)\delta}\int_{n\delta}^{(n+m)\delta}\langle\phi_0,X_s\rangle\mathrm{d}s=\lim_{n\to\infty}\int_0^{m\delta}\gamma_{(n+m)\delta}\langle\phi_0,X_{s+n\delta}\rangle\mathrm{d}s\\
&=\int_0^{m\delta}e^{-\lambda_0(m\delta-s)}\mathrm{d}sW,
\end{aligned}$$

and by (6.3) we have

$$\lim_{n\to\infty}\sup_{x\in E}\left(\int_{\|\phi_0\|_\infty^{-1}\gamma_{n\delta}^{-1}}^{\infty}\theta n(x,\mathrm{d}\theta)\right)=0.$$

Now we easily see that (6.17) holds. The proof is now complete. □

**Lemma 6.5** *For any $m\in\mathbb{N}$, $\delta>0$, $\mu\in\mathcal{M}_F(E)$ and $f\in\mathcal{B}_b^+(E)$, we have*

$$\lim_{n\to\infty}C_{(n+m)\delta}(f)-\mathbb{P}_\mu(C_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})=0,\qquad a.s.\text{-}\mathbb{P}_\mu.\tag{6.18}$$

**Proof:** Let

$$\widetilde{M}_t:=\int_0^t\int_E T_{(n+m)\delta-s}(\phi_0 f)(x)S^C(\mathrm{d}s,\mathrm{d}x).$$

Then $\{\widetilde{M}_t,0\leq t\leq(n+m)\delta\}$ is a martingale with quadratic variation

$$\langle\widetilde{M}\rangle_t=2\int_0^t\langle\beta(T_{(n+m)\delta-s}(\phi_0 f))^2,X_s\rangle\mathrm{d}s.$$

Note that

$$C_{(n+m)\delta}(f)-\mathbb{P}_\mu(C_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})$$

$$= \gamma_{(n+m)\delta} \int_{n\delta}^{(n+m)\delta} \int_E T_{(n+m)\delta-s}(\phi_0 f)(x) S^C(\mathrm{d}s, \mathrm{d}x)$$
$$= \gamma_{(m+n)\delta} \left(\widetilde{M}_{(n+m)\delta} - \widetilde{M}_{n\delta}\right). \tag{6.19}$$

Using this we get

$$\begin{aligned}
&\mathbb{P}_\mu \left( [C_{(n+m)\delta}(f) - \mathbb{P}_\mu(C_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})]^2 | \mathcal{G}_{(n-1)\delta} \right) \\
&= \gamma^2_{(m+n)\delta} \mathbb{P}_\mu \left( \langle \widetilde{M} \rangle_{(n+m)\delta} - \langle \widetilde{M} \rangle_{n\delta} | \mathcal{G}_{(n-1)\delta} \right) \\
&= \gamma^2_{(m+n)\delta} \mathbb{P}_\mu \left( 2 \int_{n\delta}^{(n+m)\delta} \langle \beta (T_{(n+m)\delta-s}(\phi_0 f))^2, X_s \rangle \mathrm{d}s | \mathcal{G}_{(n-1)\delta} \right) \\
&= \gamma^2_{(m+n)\delta} \mathbb{P}_{X_{(n-1)\delta}} \left( 2 \int_{\delta}^{(1+m)\delta} \langle \beta (T_{(1+m)\delta-s}(\phi_0 f))^2, X_s \rangle \mathrm{d}s \right).
\end{aligned}$$

Note that

$$\beta(x)(T_{(1+m)\delta-s}(\phi_0 f))^2(x) \le \|f\|^2_\infty \beta(x) e^{2\lambda_0((1+m)\delta-s)} \phi_0(x)^2 \le \|f\|^2_\infty \|\beta\phi_0\|_\infty e^{2\lambda_0 m\delta} \phi_0(x).$$

Thus we have

$$\begin{aligned}
&\mathbb{P}_\mu \left( [C_{(n+m)\delta}(f) - \mathbb{P}_\mu(C_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})]^2 | \mathcal{G}_{(n-1)\delta} \right) \\
&\le C\gamma^2_{(m+n)\delta} \mathbb{P}_{X_{(n-1)\delta}} \int_{\delta}^{(1+m)\delta} \langle \phi_0, X_s \rangle \mathrm{d}s = C\gamma^2_{(m+n)\delta} \int_{\delta}^{(1+m)\delta} \langle T_s \phi_0, X_{(n-1)\delta} \rangle \mathrm{d}s \\
&= C\gamma^2_{(m+n)\delta} \int_{\delta}^{(1+m)\delta} e^{\lambda_0 s} \mathrm{d}s \, \langle \phi_0, X_{(n-1)\delta} \rangle \le C\gamma_{(m+n)\delta} \Big( \gamma_{(m+n)\delta} \langle \phi_0, X_{(n-1)\delta} \rangle \Big).
\end{aligned}$$

By (4.7), we have that $\lim_{n\to\infty} \frac{\gamma_{(m+n)\delta}}{\gamma_{(m+n-1)\delta}} = e^{-\lambda_0\delta} < 1$, which implies that

$$\sum_{n=1}^{\infty} \gamma_{(m+n)\delta} < \infty.$$

By Theorem 5.2,

$$\lim_{n\to\infty} \gamma_{(m+n)\delta} \langle \phi_0, X_{(n-1)\delta} \rangle = e^{-\lambda_0(m+1)\delta} W.$$

Thus we have

$$\sum_{n=1}^{\infty} \gamma_{(m+n)\delta} \Big( \gamma_{(m+n)\delta} \langle \phi_0, X_{(n-1)\delta} \rangle \Big) < \infty,$$

which implies that

$$\sum_{n=1}^{\infty} \mathbb{P}_\mu \left( [C_{(n+m)\delta}(f) - \mathbb{P}_\mu(C_{(n+m)\delta}(f)|\mathcal{G}_{n\delta})]^2 | \mathcal{G}_{(n-1)\delta} \right) < \infty.$$

Now using the conditional Bore-Cantelli lemma, we immediately get (6.18). □

Combining the three results above, we get

**Lemma 6.6** *For any $m \in \mathbb{N}$, $\delta > 0$, $\mu \in \mathcal{M}_F(E)$ and $f \in \mathcal{B}_b^+(E)$, we have*

$$\lim_{n\to\infty} \gamma_{(n+m)\delta}\langle \phi_0 f, X_{(n+m)\delta}\rangle - \mathbb{P}_\mu[\gamma_{(n+m)\delta}\langle \phi_0 f, X_{(n+m)\delta}\rangle | \mathcal{G}_{n\delta}] = 0, \qquad a.s.\text{-}\mathbb{P}_\mu.$$

**Proof of Theorem 6.1:** By the Markov property of $X$, we have

$$\gamma_{(n+m)\delta}\mathbb{P}_\mu(\langle \phi_0 f, X_{(n+m)\delta}\rangle | \mathcal{G}_{n\delta}) = \gamma_{(n+m)\delta}\langle T_{m\delta}(\phi_0 f), X_{n\delta}\rangle.$$

It follows from (2.2) that there exist constants $c > 0$ and $\gamma > 0$ such that for any $m \geq 1$,

$$(1 - ce^{-\gamma m\delta})e^{\lambda_0 m\delta}\langle f\phi_0, \psi_0\rangle_m \phi_0(x) \leq T_{m\delta}(\phi_0 f)(x) \leq (1 + ce^{-\gamma m\delta})e^{\lambda_0 m\delta}\langle f\phi_0, \psi_0\rangle_m \phi_0(x).$$

Thus, by Lemma 6.6, we have that, for any $m \in \mathbb{N}$,

$$\begin{aligned}
\limsup_{n\to\infty} \gamma_{n\delta}\langle \phi_0 f, X_{n\delta}\rangle &= \limsup_{n\to\infty} \gamma_{(n+m)\delta}\langle \phi_0 f, X_{(n+m)\delta}\rangle \\
&= \limsup_{n\to\infty} \gamma_{(n+m)\delta}\langle T_{m\delta}(\phi_0 f), X_{n\delta}\rangle \\
&\leq \limsup_{n\to\infty} \gamma_{(n+m)\delta}(1 + ce^{-\gamma m\delta})e^{\lambda_0 m\delta}\langle f\phi_0, \psi_0\rangle_m \langle \phi_0, X_{n\delta}\rangle \\
&= (1 + ce^{-\gamma m\delta})\langle f\phi_0, \psi_0\rangle_m W.
\end{aligned}$$

Letting $m \to \infty$, we get

$$\limsup_{n\to\infty} \gamma_{n\delta}\langle \phi_0 f, X_{n\delta}\rangle \leq \langle f\phi_0, \psi_0\rangle_m W. \tag{6.20}$$

Similarly, we have

$$\liminf_{n\to\infty} \gamma_{n\delta}\langle \phi_0 f, X_{n\delta}\rangle \geq (1 - ce^{-\gamma m\delta})\langle f\phi_0, \psi_0\rangle_m W.$$

Letting $m \to \infty$, we get

$$\liminf_{n\to\infty} \gamma_{n\delta}\langle \phi_0 f, X_{n\delta}\rangle \geq \langle f\phi_0, \psi_0\rangle_m W. \tag{6.21}$$

Combining (6.20) and (6.21), the conclusion follows immediately. □

## 6.3 Continuous times: Case I

Define a new semigroup $(T_t^{\phi_0}, t \geq 0)$ by

$$T_t^{\phi_0} f(x) := \frac{e^{-\lambda_0 t} T_t(f\phi_0)(x)}{\phi_0(x)}, \qquad f \in \mathcal{B}_b(E).$$

Then $(T_t^{\phi_0}, t \geq 0)$ is conservative semigroup with transition density

$$q^{\phi_0}(t, x, y) = \frac{e^{-\lambda_0 t} q(t, x, y)\phi_0(y)}{\phi_0(x)}.$$

In this subsection, we also make the following assumption:

**Assumption 6** *For any $f \in C_0(E)$,*

$$\lim_{t\to 0} \|T_t^{\phi_0} f - f\|_\infty = 0. \tag{6.22}$$

See Examples 4.4, 4.5, 4.7 and Remark 4.6 of [2] for examples satisfying the assumption above, and Assumptions 1 and 2.

**Theorem 6.7** *Under Assumptions 1–6, we have that, for any $\mu \in \mathcal{M}_F(E)$ and $f \in C_0(E)$,*

$$\lim_{t\to\infty} \gamma_t\langle \phi_0 f, X_t\rangle = \langle f\phi_0, \psi_0\rangle_m W, \qquad a.s.\text{-}\mathbb{P}_\mu. \tag{6.23}$$

**Proof:** First, we claim that

$$\lim_{\delta\to 0} \limsup_{n\to\infty} \sup_{t\in[n\delta,(n+1)\delta]} \left|\gamma_t\langle \phi_0 T^{\phi_0}_{(n+1)\delta-t} f, X_t\rangle - \gamma_t\langle \phi_0 f, X_t\rangle\right| = 0. \tag{6.24}$$

In fact, we have

$$\begin{aligned}
&\sup_{t\in[n\delta,(n+1)\delta]} \left|\gamma_t\langle \phi_0 T^{\phi_0}_{(n+1)\delta-t} f, X_t\rangle - \gamma_t\langle \phi_0 f, X_t\rangle\right| \\
&\le \sup_{r\in(0,\delta)} \|T_r^{\phi_0} f - f\|_\infty \sup_{t\in[n\delta,(n+1)\delta]} \gamma_t\langle \phi_0, X_t\rangle.
\end{aligned}$$

Letting $n \to \infty$ and then $\delta \to 0$, using Assumption 6 and Theorem 5.2, one immediately arrives at the claim (6.24). Thus, by (6.24), to obtain (6.23), we only need to prove that, for any $f \in C_0(E)$,

$$\lim_{\delta\to 0} \lim_{n\to\infty} \sup_{t\in[n\delta,(n+1)\delta]} \left|\gamma_t\langle \phi_0 T^{\phi_0}_{(n+1)\delta-t} f, X_t\rangle - \langle f\phi_0, \psi_0\rangle_m W\right| = 0. \tag{6.25}$$

Since $\phi_0 T^{\phi_0}_{(n+1)\delta-t} f = e^{-\lambda_0((n+1)\delta-t)} T_{(n+1)\delta-t}(\phi_0 f)$, we only need to show that, for any $f \in C_0(E)$,

$$\lim_{\delta\to 0} \lim_{n\to\infty} \sup_{t\in[n\delta,(n+1)\delta]} \left|\gamma_t\langle T_{(n+1)\delta-t} f, X_t\rangle - \langle f, \psi_0\rangle_m W\right| = 0. \tag{6.26}$$

By (6.4), we have that, for any $t \in [n\delta, (n+1)\delta]$,

$$\begin{aligned}
&\langle T_{(n+1)\delta-t} f, X_t\rangle = \\
&= \langle T_\delta f, X_{n\delta}\rangle + \int_{n\delta}^t \int_{\mathcal{D}_{<1}(s)} \langle T_{(n+1)\delta-s} f, \nu\rangle \widetilde{N}(\mathrm{d}s, \mathrm{d}\nu) + \\
&\quad + \int_{n\delta}^t \int_{\mathcal{D}_{\ge 1}(s)} \langle T_{(n+1)\delta-s} f, \nu\rangle \widetilde{N}(\mathrm{d}s, \mathrm{d}\nu) + \int_{n\delta}^t \int_E T_{(n+1)\delta-s} f(x) S^C(\mathrm{d}s, \mathrm{d}x) \\
&=: \langle T_\delta f, X_{n\delta}\rangle + H_t^{n,\delta}(f) + L_t^{n,\delta}(f) + C_t^{n,\delta}(f).
\end{aligned}$$

It follows from (2.2) that $\phi_0(x)^{-1} T_\delta f(x) \in \mathcal{B}_b^+(E)$. Thus, by Theorem 5.2, we have

$$\lim_{n\to\infty} \gamma_{n\delta}\langle T_\delta f, X_{n\delta}\rangle = \langle T_\delta f, \psi_0\rangle_m W = e^{\lambda_0\delta}\langle f, \psi_0\rangle_m W.$$

Note that $\gamma_{(n+1)\delta} \le \gamma_t \le \gamma_{n\delta}$. Thus,

$$\lim_{\delta\to 0}\lim_{n\to\infty}\sup_{t\in[n\delta,(n+1)\delta]}|\gamma_t\langle T_\delta f, X_{n\delta}\rangle - \langle f,\psi_0\rangle_m W| = 0.$$

To finish the proof, it suffices to show that

$$\lim_{n\to\infty}\sup_{t\in[n\delta,(n+1)\delta]}\gamma_{(n+1)\delta}|H_t^{n,\delta}(f)| = 0; \tag{6.27}$$

$$\lim_{n\to\infty}\sup_{t\in[n\delta,(n+1)\delta]}\gamma_{(n+1)\delta}|L_t^{n,\delta}(f)| = 0; \tag{6.28}$$

and

$$\lim_{n\to\infty}\sup_{t\in[n\delta,(n+1)\delta]}\gamma_{(n+1)\delta}|C_t^{n,\delta}(f)| = 0. \tag{6.29}$$

Using the Markov property of the superprocess $X$, we get that

$$\begin{aligned}
&\gamma^2_{(n+1)\delta}\mathbb{P}_\mu\left(\sup_{t\in[n\delta,(n+1)\delta]}(H_t^{n,\delta}(f))^2|\mathcal{G}_{(n-1)\delta}\right)\\
&= \gamma^2_{(n+1)\delta}\mathbb{P}_{X_{(n-1)\delta}}\left(\sup_{t\in[\delta,2\delta]}\left[\int_\delta^t\int_{\mathcal{D}_{<1}(s+(n-1)\delta)}\langle T_{2\delta-s}f,\nu\rangle\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu)\right]^2\right)\\
&\le 4\gamma^2_{(n+1)\delta}\mathbb{P}_{X_{(n-1)\delta}}\left(\left[\int_\delta^{2\delta}\int_{\mathcal{D}_{<1}(s+(n-1)\delta)}\langle T_{2\delta-s}f,\nu\rangle\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu)\right]^2\right)\\
&= 4\mathbb{P}_\mu\left([H_{(n+1)\delta}(f)-\mathbb{P}_\mu(H_{(n+1)\delta}(f)|\mathcal{G}_{n\delta})]^2|\mathcal{G}_{(n-1)\delta}\right),
\end{aligned}$$

where the second to the last line follows from the fact that

$$\left(\int_\delta^t\int_{\mathcal{D}_{<1}(s+(n-1)\delta)}\langle T_{2\delta-s}f,\nu\rangle\widetilde{N}(\mathrm{d}s,\mathrm{d}\nu), t\in[\delta,2\delta]\right)$$

is a martingale. Therefore, by (6.9), we have

$$\sum_{n=1}^\infty\gamma^2_{(n+1)\delta}\mathbb{P}_\mu\left(\sup_{t\in[n\delta,(n+1)\delta]}(H_t^{n,\delta}(f))^2|\mathcal{G}_{(n-1)\delta}\right) < \infty.$$

Using the conditional Borel-Cantelli lemma, (6.27) follows immediately.

Similarly, we can prove that (6.29) holds. We omit the details here.

Note that

$$|L_t^{n,\delta}(f)| \le \int_{n\delta}^{(n+1)\delta}\int_{\mathcal{D}_{\ge 1}(s)}\langle T_{(n+1)\delta-s}f,\nu\rangle\left(N(\mathrm{d}s,\mathrm{d}\nu)+\widehat{N}(\mathrm{d}s,\mathrm{d}\nu)\right).$$

Now using (6.16) and (6.17) with $m = 1$, we immediately get (6.28). The proof is now complete. □

**Theorem 6.8** *Suppose that Assumptions 1–6 hold. There exists $\Omega_0 \subset \Omega$ of probability one (that is, $\mathbb{P}_\mu(\Omega_0) = 1$ for every $\mu \in \mathcal{M}_F(E)$) such that, for every $\omega \in \Omega_0$ and for every bounded Borel function $f$ on $E$ satisfying (a) $|f| \le c\phi_0$ for some $c > 0$ and (b) the set of discontinuous points of $f$ has zero $m$-measure, we have*

$$\lim_{t\to\infty} \gamma_t\langle f, X_t\rangle(\omega) = \langle f, \psi_0\rangle_m W(\omega).$$

**Proof:** With the preparation above, the proof of this theorem is similar to that of [2, Theorem 1.4]. We omit the details here. □

## 6.4 Continuous times: Case II

In this subsection, we will consider the almost sure limit of $\gamma_t\langle f, X_t\rangle$ with $f$ being a general bounded continuous function for some class of superdiffusions. The underlying process $\xi$ is a diffusion satisfying the following conditions.

Suppose that $E$ is a domain of finite Lebesgue measure in $\mathbb{R}^d$. Denote by $C_b^1(E)$ the family of bounded differentiable functions on $E$ whose first order partial derivatives are all continuous. The underling process $\{\xi, \Pi_x\}$ is a killed diffusion process on $E$ corresponding to the infinitesimal generator

$$L = \frac{1}{2}\nabla \cdot a\nabla + b \cdot \nabla, \tag{6.30}$$

where $a$ and $b$ satisfy the following conditions:

(a) $a_{ij} \in C_b^1(E)$, $i, j = 1, 2\cdots, d$, and that the matrix $a = (a_{ij})$ is symmetric which satisfying, for all $x \in E$ and $v \in \mathbb{R}^d$,

$$c_0|v|^2 \le \sum_{i,j} a_{ij}v_iv_j,$$

for some positive constant $c_0$.

(b) $b_j \in \mathcal{B}_b(E)$, $j = 1, \cdots, d$.

Using an argument similar to that in [3, section 3.2], one can easily show that $P_t$ has a bounded continuous and strictly positive density $p(t, x, y)$. Thus Assumption 1 holds immediately. Since $m(E) < \infty$ and the first eigenfunction $\widetilde{\phi}_0 \in L^2(E, m)$, we have that $\widetilde{\phi}_0 \in L^1(E, m)$. Then using the fact that $p(1, x, y)$ is bounded and $\widetilde{\phi}_0(x) = e^{-\widetilde{\lambda}_0}P_1\widetilde{\phi}_0(x)$, we get that $\widetilde{\phi}_0$ is bounded on $E$. Similarly, $\widetilde{\psi}_0$ is also bounded, which shows that Assumption 2(i) holds. We assume that the semigroup $P_t$ is intrinsically ultracontractive.

Let $f \in \mathcal{B}_b(E)$, and $U^q f$, $q > 0$, be the $q$-potential of $f$, that is

$$U^q f(x) = \int_0^\infty e^{-qs}T_s^{\phi_0} f(x)\mathrm{d}s.$$

For any $t > 0$,

$$e^{-qt}T_t^{\phi_0}(U^q f)(x) = \int_t^\infty e^{-qs}T_s^{\phi_0} f(x)\mathrm{d}s. \tag{6.31}$$

**Theorem 6.9** *For any $q > 0$, $\mu \in \mathcal{M}_F(E)$ and $f \in \mathcal{B}_b^+(E)$, we have*

$$\lim_{t\to\infty} \gamma_t\langle \phi_0 q U^q f, X_t\rangle = \langle f\phi_0, \psi_0\rangle_m W, \qquad a.s.\text{-}\mathbb{P}_\mu. \tag{6.32}$$

**Proof:** First, we claim that

$$\lim_{\delta\to 0}\lim_{n\to\infty}\sup_{t\in[n\delta,(n+1)\delta]}\left|\gamma_t\langle \phi_0 T^{\phi_0}_{(n+1)\delta-t}(U^q f), X_t\rangle - \gamma_t\langle \phi_0 U^q f, X_t\rangle\right| = 0. \tag{6.33}$$

For any $r \in [0,\delta]$, we have

$$\begin{aligned}
&q\left|T_r^{\phi_0}(U^q f)(x) - U^q f(x)\right| \\
&= q\left|(e^{qr}-1)\int_r^\infty e^{-qs}T_s^{\phi_0} f(x)\mathrm{d}s - \int_0^r e^{-qs}T_s^{\phi_0} f(x)\mathrm{d}s\right| \\
&\le \|f\|_\infty\left(q(e^{qr}-1)\int_r^\infty e^{-qs}\mathrm{d}s + q\int_0^r e^{-qs}\mathrm{d}s\right) \\
&= 2\|f\|_\infty(1-e^{-qr}).
\end{aligned}$$

Thus,

$$\begin{aligned}
&\sup_{t\in[n\delta,(n+1)\delta]}\left|\gamma_t\langle \phi_0 T^{\phi_0}_{(n+1)\delta-t}(qU^q f), X_t\rangle - \gamma_t\langle \phi_0 qU^q f, X_t\rangle\right| \\
&\le 2\|f\|_\infty(1-e^{-q\delta})\sup_{t\in[n\delta,(n+1)\delta]}\gamma_t\langle \phi_0, X_t\rangle.
\end{aligned}$$

Letting $n\to\infty$ and then $\delta\to 0$, the claim (6.33) follows immediately. Note that

$$\langle qU^q f\phi_0, \psi_0\rangle_m = \langle f\phi_0, \psi_0\rangle_m.$$

Thus, applying (6.25) with $f$ replaced by $qU^q f$, we get

$$\lim_{\delta\to 0}\lim_{n\to\infty}\sup_{t\in[n\delta,(n+1)\delta]}\left|\gamma_t\langle \phi_0 T^{\phi_0}_{(n+1)\delta-t}(qU^q f), X_t\rangle - \langle \phi_0 f, \psi_0\rangle_m W\right| = 0. \tag{6.34}$$

Now combining (6.33) and (6.34), (6.32) follows immediately. □

**Theorem 6.10** *Suppose $X$ is a superdiffusion on a domain $E \subset \mathbb{R}^d$ of finite Lebesgue measure with spatial motion being a killed diffusion in $E$ with generator $L$ given in (6.30) satisfying conditions (a) and (b). Suppose that Assumptions 1-5 hold. Then there exists $\Omega_0 \subset \Omega$ of probability one (that is, $\mathbb{P}_\mu(\Omega_0) = 1$ for every $\mu \in \mathcal{M}_F(E)$) such that, for every $\omega \in \Omega_0$ and for every bounded Borel function $f$ on $E$ satisfying (a) $|f| \le c\phi_0$ for some $c > 0$ and (b) the set of discontinuous points of $f$ has zero $m$-measure, we have*

$$\lim_{t\to\infty}\gamma_t\langle f, X_t\rangle(\omega) = \langle f, \psi_0\rangle_m W(\omega).$$

**Proof:** With the preparation above, the proof of this theorem is similar to that of [17, Theorem 1.1]. We omit the details here. □

# 7 Concluding remarks

Suppose that $X = \{X_t, t \geq 0; \mathbb{P}_\mu\}$ is a supercritical superprocess in a locally compact separable metric space $E$ such that the generator of the mean semigroup of $X$ has discrete spectrum. Let $\phi_0$ be a positive eigenfunction corresponding to the first eigenvalue $\lambda_0$ of the generator of the mean semigroup of $X$. Then $M_t := e^{-\lambda_0 t}\langle \phi_0, X_t\rangle$ is a positive martingale. Let $M_\infty$ be the limit of $M_t$. It is known (see [16]) that $M_\infty$ is non-degenerate iff the $L \log L$ condition is satisfied. In this paper, we prove that, under some further conditions, there exist a positive function $\gamma_t$ on $[0, \infty)$ and a non-degenerate random variable $W$ such that for any finite nonzero Borel measure $\mu$ on $E$,

$$\lim_{t\to\infty} \gamma_t\langle \phi_0, X_t\rangle = W, \qquad \text{a.s.-}\mathbb{P}_\mu.$$

We also give the almost sure limit of $\gamma_t\langle f, X_t\rangle$ for a class of general test functions $f$.

In [23], a sequel to the present paper, we studied properties of the limit random variable $W$, such as absolute continuity and tail probabilities.

It would be interesting to extend the results of this paper and [23] to supercritical superprocesses with immigration.

The assumptions of this paper, particularly Assumption 2.(ii), are pretty strong. For example, supercrtical super Brownian motion in $\mathbb{R}^d$ does not satisfy Assumption 2.(ii). It would be interesting to consider corresponding results of this paper and [23] for supercritical superprocesses under weaker conditions. It would be very interesting to get rid of the assumption that the generator of the mean semigroup of $X$ has discrete spectrum.

**Acknowledgements:** We thank the referee for very careful reading of the paper and for very helpful comments.

# References

[1] Athreya K, Ney P. Branching Processes. Springer-Verlag, 1972

[2] Chen Z-Q, Ren Y-X, Song R, Zhang R. Strong law of large numbers for supercritical superprocesses under second moment condition. Front Math China, 2015, 10: 807–838

[3] Chung K L, Zhao Z. From Brownian motion to Schrödinger's equation. Springer-Berlin, 1995

[4] Davies E B, Simon B. Ultracontractivity and the heat kernel for Schrödinger operators and Dirichlet Laplacians. J Funct Anal, 1984, 59: 335–395

[5] Dawson D A. Measure-Valued Markov Processes. Springer-Verlag, 1993

[6] Dynkin E B. Superprocesses and partial differential equations. Ann Probab, 1993, 21: 1185–1262

[7] Fitzsimmons P J. On the martingale problem for measure-valued Markov branching processes. In: Cinlar E, Chung K L, Sharpe M. (eds.) Seminar on Stochastic Processes, 1991, 39–51, Progr Probab, 29. Birkhäuser. Boston 1992.

[8] Grey D R. Asymptotic behaviour of continuous time, continuous state-space branching processes. J Appl Probab, 1974, 11: 669–677

[9] Hering H. The non-degenerate limit for supercritical branching diffusions. Duke Math J, 1978, 45: 561–599

[10] Heyde C C. Extension of a result of Seneta for the supercritical branching process. Ann Math Statist, 1970, 41: 739–742

[11] Hoppe F M. Supercritical multitype branching processes. Ann Prob, 1976, 4: 393–401

[12] Kim P, Song R. Intrinsic ultracontractivity of non-symmetric diffusion semigroups in bounded domains. Tohoku Math J, 2008, 60: 527–547

[13] Kim P, Song R. Intrinsic ultracontractivity of non-symmetric diffusions with measure-valued drifts and potentials. Ann Probab, 2008, 36: 1904–1945

[14] Kim P, Song R. Intrinsic ultracontractivity for non-symmetric Lévy processes. Forum Math, 2009, 21: 43–66

[15] Li Z. Measure-Valued Branching Markov Processes. Springer, Heidelberg, 2011

[16] Liu R-L, Ren Y-X, Song R. $L \log L$ criteria for a class of superdiffusons. J Appl Probab, 2009, 46: 479–496

[17] Liu R-L, Ren Y-X, Song R. Strong law of large numbers for a class of superdiffusions. Acta Appl Math, 2013, 123: 73–97

[18] Lyons R, Pemantle R, Peres Y. Conceptual proofs of $L \log L$ criteria for mean behavior of branching processes. Ann Probab, 1995, 23: 1125–1138

[19] Perkins E. Dawson-Watanabe superprocesses and measure-valued diffusions. In: Lect Notes Math, 2002, 1781:135–192. Springer, Heidelberg

[20] Ren Y-X, Song R, Yang T. Spine decomposition and $L \log L$ criterion for superprocesses with non-local branching mechanisms. Preprint 2016. arXiv:1609.02257.

[21] Ren Y-X, Song R, Zhang R. Limit theorems for some critical superprocesses. Illinois J Math, 2015, 59: 235–276

[22] Ren Y-X, Song R, Zhang R. Central limit theorems for supercritical branching non-symmetric Markov processes. Ann Probab, 2017, 45: 564–623

[23] Ren Y-X, Song R, Zhang R. On properties of a class of strong limits for supercritical superprocesses (in Chinese). Sci Sin Math, (to appear), 2019. (See arXiv: 1803.02973 for the English version.)

[24] Rudin W. Functional Analysis. 2nd Ed. McGraw-Hill, 1991.

[25] Schaefer H H. Banach Lattices and Positive Operators. Springer, New York, 1974

[26] Seneta E. On recent theorems concerning the supercritical Galton-Watson process. Ann Math Statist, 1968, 39: 2098–2102

**Yan-Xia Ren:** LMAM School of Mathematical Sciences & Center for Statistical Science, Peking University, Beijing, 100871, P.R. China. Email: yxren@math.pku.edu.cn

**Renming Song:** Department of Mathematics, University of Illinois, Urbana, IL 61801, U.S.A and School of Mathematical Sciences, Nankai University, Tianjin 300071, P. R. China. Email: rsong@illinois.edu

**Rui Zhang:** School of Mathematical Sciences, Capital Normal University, Beijing, 100048, P.R. China. Email: zhangrui27@cnu.edu.cn