

A metric result for special subsequences of the Halton sequences

Roswitha Hofer*

March 7, 2024

Abstract

In this paper we investigate the special subsequence of the Halton sequence indexed by $\lfloor n\beta \rfloor$ with $\beta \in \mathbb{R}$ and prove a metric almost low-discrepancy result.

KEYWORDS: Halton sequences, Subsequences, Discrepancy.

MSC2010: 11K31, 11K38.

1 Introduction

The *discrepancy* of the first N terms of any sequence $(z_n)_{n \geq 0}$ of points in $[0, 1]^s$ is defined by

$$D_N(z_n) = \sup_J \left| \frac{A_N(J)}{N} - \lambda_s(J) \right|,$$

where the supremum is extended over all half-open subintervals J of $[0, 1]^s$, λ_s denotes the s -dimensional Lebesgue measure, and the counting function $A_N(J)$ is given by

$$A_N(J) = \#\{0 \leq n \leq N-1 : z_n \in J\}.$$

For the sake of simplicity, we will sometimes write D_N instead of $D_N(z_n)$. If the supremum is extended over all half-open subintervals J of $[0, 1]^s$ with the lower left point in the origin then we arrive at the notion of the *star-discrepancy* D_N^* . It is not so hard to see that $D_N^* \leq D_N \leq 2^s D_N^*$. A sequence $(z_n)_{n \geq 0}$ of points in $[0, 1]^s$ is called uniformly distributed if $\lim_{N \rightarrow \infty} D_N = 0$.

It is frequently conjectured in the theory of irregularities of distribution, that for every sequence $(z_n)_{n \geq 0}$ in $[0, 1]^s$ we have

$$D_N \geq c_s \frac{\log^s N}{N}$$

*supported by the Austrian Science Fund (FWF): Project F5505-N26, which is a part of the Special Research Program “Quasi-Monte Carlo Methods: Theory and Applications”

for a constant $c_s > 0$ and for infinitely many N . Therefore sequences whose discrepancy satisfies $D_N \leq C \log^s N/N$ for all N with a constant $C > 0$ that is independent of N (or $D_N = O(\log^s N/N)$), are called *low-discrepancy sequences*. Well-known examples of low-discrepancy sequences are, for example, the s -dimensional Halton sequences and one-dimensional Kronecker sequences $(\{n\alpha\})_{n \geq 0}$ with α irrational and having bounded continued fraction coefficients.

For an integer $b \geq 2$, let $\mathbb{Z}_b = \{0, 1, \dots, b-1\}$ denote the least residue system modulo b . Let $n = \sum_{j=1}^{\infty} a_j(n)b^{j-1}$ with all $a_j(n) \in \mathbb{Z}_b$ and $a_j(n) = 0$ for all sufficiently large j be the unique digit expansion of the integer $n \geq 0$ in base b . The *radical-inverse function* ϕ_b in base b is defined by

$$\phi_b(n) = \sum_{j=1}^{\infty} a_j(n)b^{-j}.$$

The *Halton sequence* $(\mathbf{y}_n)_{n \geq 0}$ (in the bases b_1, \dots, b_s) is given by

$$\mathbf{y}_n = (\phi_{b_1}(n), \dots, \phi_{b_s}(n)) \in [0, 1)^s, \quad n = 0, 1, 2, \dots$$

It is well known (see [7, Theorem 3.6]) that the discrepancy of the Halton sequence in pairwise coprime bases $b_1, \dots, b_s \geq 2$ satisfies a low-discrepancy bound, i.e.,

$$D_N(\mathbf{y}_n) = O_{b_1, \dots, b_s}((\log N)^s/N) \quad \text{for all } N \geq 2. \quad (1)$$

The one-dimensional Kronecker sequences $(\{n\alpha\})_{n \geq 0}$ related to the real number α is uniformly distributed if α is irrational. This sequence is a low-discrepancy sequence if the continued fraction coefficients of α are bounded. Furthermore, the discrepancy D_N satisfies $ND_N \leq C(\alpha, \epsilon) \log^{1+\epsilon} N$ for all N and all $\epsilon > 0$ for almost all $\alpha \in \mathbb{R}$ in the sense of Lebesgue measure. Here the positive constant $C(\alpha, \epsilon)$ may depend on α and $\epsilon > 0$ but is independent of N .


Subsequences of the one-dimensional Kronecker sequence and Halton sequences are frequently studied objects (see for instance [1], [2], [3], [4]). Here we mention a result of Hofer and Ramare [3] who studied the sequence $(\lfloor n\alpha \rfloor \beta)_{n \geq 0}$. It is known to be u.d. mod 1 when α is a nonzero rational real if and only if β is irrational. When $\alpha = 0$ the sequence is for no real β u.d. mod 1. And when α is irrational the uniform distribution modulo one of $(\lfloor n\alpha \rfloor \beta)_{n \geq 0}$ is equivalent to the condition, $1, \alpha, \alpha\beta$ being linearly independent over the rationals. (For a proof see [5, Chapter 5, Theorem 1.8].) In [3] many results on the discrepancy of this sequence were proved. Here we mention just one result, i.e., for almost all pairs of real numbers (α, β) in the sense of Lebesgue measure we have for every $\epsilon > 0$ that $D_N(\lfloor n\alpha \rfloor \beta) \ll_{\alpha, \beta, \epsilon} N^{-1+\epsilon}$.

In this paper we investigate the subsequence $(\mathbf{x}_n)_{n \geq 0}$ of the Halton sequence of this particular form

$$\mathbf{x}_n := (\phi_{b_1}(\lfloor n\beta \rfloor), \dots, \phi_{b_s}(\lfloor n\beta \rfloor)) \in [0, 1)^s, \quad n = 0, 1, 2, \dots \quad (2)$$

with nonzero $\beta \in \mathbb{R}$ and integers $b_1, \dots, b_s \geq 2$ pairwise coprime. See Figure 1 for examples.

Figure 1: The first 500 points in bases 2 and 3 for $\beta = 1, \sqrt{2}, \pi$ and e .



In [2, Example 5.3] the uniform distribution of (2) was studied. A sufficient criterion for the one-dimensional sequence is, for example, the linear independence of $1, \beta$ over \mathbb{Q} . This criterion is not a necessary one, as the choice $\beta = 1/d$ with $d \in \mathbb{N}$ also yields a uniformly distributed sequence (2). If $\beta = d \in \mathbb{N}$ such that $(d, b_i) = 1$ for $i = 1, \dots, s$ then the sequence (2) is even a low-discrepancy sequence (see [4]).

In this paper we prove that the sequence (2) is for almost all $\beta \in \mathbb{R}$ an almost low-discrepancy sequence by the following theorem.

Theorem 1. *For almost all $\beta \in \mathbb{R}$ in the sense of Lebesgue measure the dis-*

crepancy of $(\mathbf{x}_n)_{n \geq 0}$ satisfies for all $\epsilon > 0$

$$ND_N = O_{\beta, \epsilon, b_1, \dots, b_s, s}(\log^{s+1+\epsilon} N).$$

A proof of this theorem will be outsourced in the next section.

2 Proof of Theorem 1

We start with the case $\beta \in (0, 1)$. We write $\alpha := 1/\beta$. Note that $\alpha > 1$. Then Theorem 1 immediately follows from the following three auxiliary results.

Proposition 1. *The star-discrepancy of the first N points of the sequence (2) satisfies*

$$ND_N^*(\mathbf{x}_n) \leq b_1 \dots b_s \sum_{j_1=0}^{f_1} \dots \sum_{j_s=0}^{f_s} N\Delta_N^{(j_1, \dots, j_s)}(\{n/(ab_1^{j_1} \dots b_s^{j_s})\}) + s,$$

where

$$f_i := \left\lceil \frac{1}{\log b_i} \log N \right\rceil \quad \text{for } 1 \leq i \leq s \quad (3)$$

and

$$N\Delta_N^{(j_1, \dots, j_s)} \left(\left\{ \frac{n}{ab_1^{j_1} \dots b_s^{j_s}} \right\} \right) := \sup_{0 \leq R < b_1^{j_1} \dots b_s^{j_s}} \left| A_N \left(\left[\frac{R}{b_1^{j_1} \dots b_s^{j_s}}, \frac{R+1}{b_1^{j_1} \dots b_s^{j_s}} \right) \right) - \frac{N}{b_1^{j_1} \dots b_s^{j_s}} \right|.$$

Proof. The first aim in the proof is to compute or estimate the function $A_N(J) - N\lambda_s(J)$ relative to the points \mathbf{x}_n in (2), where $J \subseteq [0, 1)^s$ is an interval of the form

$$J = \prod_{i=1}^s [0, v_i b_i^{-f_i}) \quad (4)$$

with $v_1, \dots, v_s \in \mathbb{Z}$, $1 \leq v_i < b_i^{f_i}$ for $1 \leq i \leq s$. We extend v_i in base b_i , i.e.,

$$v_i = u_{f_i}^{(i)} + u_{f_i-1}^{(i)} b_i + \dots + u_1^{(i)} b_i^{f_i-1}.$$

We can write J as a union of $\prod_{i=1}^s (\sum_{j=1}^{f_i} u_j^{(i)}) \leq (b_1 - 1) \dots (b_s - 1) f_1 \dots f_s$ elementary intervals of the form

$$\prod_{i=1}^s \left(\bigcup_{j=1}^{f_i} \bigcup_{k=1}^{u_j^{(i)}} \left[\sum_{l=1}^{j-1} \frac{u_l^{(i)}}{b_i^l} + \frac{k-1}{b_i^j}, \sum_{l=1}^{j-1} \frac{u_l^{(i)}}{b_i^l} + \frac{k}{b_i^j} \right) \right)$$

The crucial step is to estimate $A(I_e, N) - N\lambda_s(I_e)$ for an elementary interval I_e by exploiting special properties of the Halton sequence. Using the definition of the sequence we obtain for any integer $n \geq 0$ that the following three assertions are equivalent.

$$(\phi_{b_1}(\lfloor n/\alpha \rfloor), \dots, \phi_{b_s}(\lfloor n/\alpha \rfloor)) \in \prod_{i=1}^s \left[\sum_{l=1}^{j_i-1} \frac{u_l^{(i)}}{b_i^l} + \frac{k_i-1}{b_i^{j_i}}, \sum_{l=1}^{j_i-1} \frac{u_l^{(i)}}{b_i^l} + \frac{k_i}{b_i^{j_i}} \right),$$

where $1 \leq j_i \leq f_i$ and $1 \leq k_i \leq u_{j_i}^{(i)}$ for $i = 1, \dots, s$.

$$\lfloor n/\alpha \rfloor \equiv a_1 \pmod{b_1^{j_1}}, \dots, \lfloor n/\alpha \rfloor \equiv a_s \pmod{b_s^{j_s}} \quad (5)$$

with $a_i = (k_i - 1)b_i^{j_i-1} + \sum_{l=1}^{j_i-1} u_l^{(i)} b_i^{l-1}$.

$$\lfloor n/\alpha \rfloor \equiv R \pmod{b_1^{j_1} \dots b_s^{j_s}}$$

where the R is uniquely determined by the a_1, \dots, a_s .

By observing the fact that

$$\lfloor n/\alpha \rfloor \equiv \lfloor n/\alpha - b_1^{j_1} \dots b_s^{j_s} \lfloor n/(\alpha b_1^{j_1} \dots b_s^{j_s}) \rfloor \rfloor \equiv R \pmod{b_1^{j_1} \dots b_s^{j_s}}$$

we easily obtain the equivalence of the above assertion to

$$\left\{ \frac{n}{\alpha b_1^{j_1} \dots b_s^{j_s}} \right\} \in \left[\frac{R}{b_1^{j_1} \dots b_s^{j_s}}, \frac{R+1}{b_1^{j_1} \dots b_s^{j_s}} \right).$$

Hence,

$$|A(J, N) - N\lambda_s(J)| \leq (b_1 - 1) \dots (b_s - 1) \sum_{j_1=1}^{f_1} \dots \sum_{j_s=1}^{f_s} N \Delta_N^{(j_1, \dots, j_s)}(\{n/(\alpha b_1^{j_1} \dots b_s^{j_s})\})$$

An arbitrary interval $I \subseteq [0, 1]^s$ of the form

$$I = \prod_{i=1}^s [0, w_i) \quad (6)$$

with $0 < w_i \leq 1$ for $1 \leq i \leq s$ can be approximated from below by an interval J of the form (4), by taking the nearest fraction to the left of w_i of the form $v_i b_i^{-f_i}$ with $v_i \in \mathbb{Z}$. We easily get

$$|A(I, N) - N\lambda_s(I)| \leq |A(J, N) - N\lambda_s(J)| + \max(A(I \setminus J, N), N\lambda_s(I \setminus J)).$$

The definition of the f_i in (3) yields $f_i \geq \log_{b_i} N$. Hence $N\lambda_s(I \setminus J) \leq N \sum_{i=1}^s b_i^{f_i} \leq s$. It remains to estimate $A(I \setminus J, N)$: We trivially have $A(I \setminus J, N) \leq A(S, N) \leq |A(S, N) - N\lambda_s(S)| + N\lambda_s(S)$ with S a union of elementary intervals I_e of the form

$$S = \bigcup_{k=1}^s \prod_{i=1}^{k-1} [0, 1) \times \left[\sum_{l=1}^{f_k-1} \frac{u_l^{(i)}}{b_i^l} + \frac{u_{f_k}^{(i)}}{b_i^{f_i}}, \sum_{l=1}^{f_k-1} \frac{u_l^{(i)}}{b_i^l} + \frac{u_{f_k}^{(i)} + 1}{b_i^{f_i}} \right) \times \prod_{i=k+1}^s [0, 1).$$

Now $N\lambda_s(I_e) \leq 1$ by the definition of f_k and

$$|A(I_e, N) - N\lambda_s(I_e)| \leq N\Delta_N^{(0, \dots, 0, f_k, 0, \dots, 0)}(\{n/(\alpha b_k^{f_k})\})$$

and the result follows. \square

The discrepancy of a one-dimensional Kronecker sequence $(\{n\beta\})_{n \geq 0}$ can be related to the continued fraction expansion of β

$$\beta = [\lfloor \beta \rfloor; a_1(\beta), a_2(\beta), a_3(\beta), \dots] = \lfloor \beta \rfloor + \frac{1}{a_1(\beta) + \frac{1}{a_2(\beta) + \frac{1}{a_3(\beta) + \frac{1}{\ddots}}}}$$

This expansion is obtained by setting $\{\beta\} = x_1$ and defining inductively

$$a_k(\beta) = \lfloor 1/x_k \rfloor \text{ and } x_{k+1} = \{1/x_k\}.$$

The convergents p_k/q_k to β are defined by $p_k/q_k = [\lfloor \beta \rfloor; a_1(\beta), a_2(\beta), \dots, a_k(\beta)]$. The denominators q_k obey the following recursion

$$q_k = a_k(\beta)q_{k-1} + q_{k-2} \text{ for } k \geq 1 \text{ and with } q_0 = 1, q_{-1} = 0.$$

Lemma 1. *We have*

$$N\Delta_N^{(j_1, \dots, j_s)}(\{n/(\alpha b_1^{j_1} \dots b_s^{j_s})\}) \leq \alpha + 1 + 2 \left(\sum_{k=1}^{\lceil \log_{3/2}(N) \rceil} a_k((\alpha b_1^{j_1} \dots b_s^{j_s})) \right).$$

Proof. We follow the idea of [5, page 125, proof of Theorem 3.4]. Now we use the fact that $\beta \in (0, 1)$ and hence $\alpha > 1$. Then $\alpha b_1^{j_1} \dots b_s^{j_s} > 1$, and the first coefficient $a_1(1/(\alpha b_1^{j_1} \dots b_s^{j_s})) = \lfloor \alpha b_1^{j_1} \dots b_s^{j_s} \rfloor$. We compute the Ostrowski expansion using the denominators $1 = q_0 = q_1 < q_2 < q_3 < \dots$ of the convergents to $1/(\alpha b_1^{j_1} \dots b_s^{j_s})$:

$$N = N_0 + N_1 q_1 + N_2 q_2 + \dots + N_r q_r$$

with $q_r \leq N < q_{r+1}$ and $N_i \leq a_{i+1}(1/(\alpha b_1^{j_1} \dots b_s^{j_s}))$.

We split

$$N\Delta_N^{(j_1, \dots, j_s)}(\{\frac{n}{\alpha b_1^{j_1} \dots b_s^{j_s}}\}) \leq N_0 \Delta_{N_0}^{(j_1, \dots, j_s)}(\{\frac{n}{\alpha b_1^{j_1} \dots b_s^{j_s}}\}) + (N - N_0) \Delta_{N - N_0}^{(j_1, \dots, j_s)}(\{\frac{n + N_0}{\alpha b_1^{j_1} \dots b_s^{j_s}}\})$$

and proceed for the second term as in [5, page 125, proof of Theorem 3.4] and obtain

$$\begin{aligned} (N - N_0) \Delta_{N - N_0}^{(j_1, \dots, j_s)}(\{\frac{n + N_0}{\alpha b_1^{j_1} \dots b_s^{j_s}}\}) &\leq 2 \sum_{k=2}^{\lceil \log_{3/2} N \rceil + 1} a_k(1/(\alpha b_1^{j_1} \dots b_s^{j_s})) \\ &\leq 2 \sum_{k=1}^{\lceil \log_{3/2} N \rceil} a_k(\alpha b_1^{j_1} \dots b_s^{j_s}). \end{aligned}$$

Finally, it is not so hard to see that $N_0 \Delta_{N_0}^{(j_1, \dots, j_s)}(\{n/(\alpha b_1^{j_1} \dots b_s^{j_s})\}) \leq \alpha + 1$ with $N_0 \leq \lfloor \alpha b_1^{j_1} \dots b_s^{j_s} \rfloor$. Note the fact that $[R/b_1^{j_1} \dots b_s^{j_s}), (R+1)/b_1^{j_1} \dots b_s^{j_s})$ either remains empty or contains at most $\lfloor \alpha \rfloor + 1$ points, and $N_0 1/b_1^{j_1} \dots b_s^{j_s} \leq \lfloor \alpha \rfloor + 1$. \square

Lemma 2. *Let $\alpha \in \mathbb{R}$. For any $L \in \mathbb{N}$ let*

$$S_L(\alpha) := \sum_{j_1=0}^L \dots \sum_{j_s=0}^L \sum_{k=1}^L a_k(\alpha b_1^{j_1} \dots b_s^{j_s}).$$

Then for almost all $\alpha \in \mathbb{R}$ we have for every $\epsilon > 0$

$$S_L(\alpha) = O_{\alpha, \epsilon, b_1, \dots, b_s, s}(L^{s+1+\epsilon}).$$

Proof. See e.g. [6, Lemma 2]. \square

Finally we consider $\beta > 1$ and set $\alpha := \beta/(\beta + 1)$. Then $1/2 < \alpha < 1$, $\lfloor n\alpha \rfloor = n - \lceil n \frac{1}{\beta+1} \rceil$ and hence

$$\{\lfloor n\alpha \rfloor; n \in \mathbb{N}\} = \{\lfloor n\beta \rfloor; n \in \mathbb{N}\} \cup \mathbb{N}_0.$$

Let

$$B := \left\{ \alpha \in (1/2, 1) : D_N(\phi_{b_1}(\lfloor n\alpha \rfloor), \dots, \phi_{b_s}(\lfloor n\alpha \rfloor)) = O_{\alpha, \epsilon, b_1, \dots, b_s} \left(\frac{(\log N)^{s+1+\epsilon}}{N} \right) \text{ for all } \epsilon > 0 \right\}.$$

From the above case we know $\lambda(B) = 1/2$. Let $A := \{\beta \in (1, \infty) : \beta/(\beta + 1) \in B\}$. Then $\lambda((1, \infty) \setminus A) = 0$.

Now let $\beta \in A$. Then $\alpha \in B$. Since

$$\underbrace{(\phi_{b_1}(\lfloor n\alpha \rfloor), \dots, \phi_{b_s}(\lfloor n\alpha \rfloor))_{n \geq 1}}_{D_N = O((\log N)^{s+1+\epsilon}/N)} = (\phi_{b_1}(\lfloor n\beta \rfloor), \dots, \phi_{b_s}(\lfloor n\beta \rfloor))_{n \geq 1} \cup \underbrace{(\phi_{b_1}(n), \dots, \phi_{b_s}(n))_{n \geq 0}}_{D_N = O((\log N)^s/N)}$$

we have

$$D_N(\phi_{b_1}(\lfloor n\beta \rfloor), \dots, \phi_{b_s}(\lfloor n\beta \rfloor)) = O_{\beta, \epsilon, b_1, \dots, b_s} \left(\frac{(\log N)^{s+1+\epsilon}}{N} \right) \text{ for all } \epsilon > 0$$

and the proof is complete.

Acknowledgments

The author would like to thank Gerhard Larcher for valuable comments and fruitful discussions.

References

- [1] C. Aistleitner, R. Hofer, and G. Larcher, *On evil Kronecker sequences and Lacunary Trigonometric Products*, to appear in Annales de l'institut Fourier. (arXiv:1502.06738)
- [2] P. Hellekalek and H. Niederreiter *Constructions of uniformly distributed sequences using the \mathbf{b} -adic method*, Unif. Distrib. Theory **6** (2011), 185–200.
- [3] R. Hofer and O. Ramare, *Discrepancy estimates for some linear generalized monomials*, *Acta Arith.* **173**, 183–196, 2016.
- [4] R. Hofer, P. Kritzer, G. Larcher, F. Pillichshammer. *Distribution properties of generalized van der Corput-Halton sequences and their subsequences*, *Int. J. Number Theory* **5**(4), 719–746, 2009.
- [5] L. Kuipers and H. Niederreiter, *Uniform distribution of sequences*, Wiley, New York, 1974; reprint, Dover Publ., Mineola, NY, 2006.
- [6] G. Larcher, *Probabilistic diophantine approximation and the distribution of Halton-Kronecker sequences*, *J. Complexity* **29**, 397–423.
- [7] H. Niederreiter, *Random Number Generation and Quasi-Monte Carlo Methods*, SIAM, Philadelphia, 1992.

Roswitha Hofer
Institute of Financial Mathematics and Applied Number Theory
Johannes Kepler University Linz
Altenbergerstr. 69
A-4040 Linz, AUSTRIA
roswitha.hofer@jku.at