

COINCIDENCE POINT RESULTS INVOLVING A GENERALIZED CLASS OF SIMULATION FUNCTIONS

D.K. PATEL, P.R. PATLE, L. BUDHIA, D. GOPAL

ABSTRACT. The purpose of this work is to introduce a general class of C_G -simulation functions and obtained some new coincidence and common fixed points results in metric spaces. Some useful examples are presented to illustrate our theorems. Results obtained in this paper extend, generalize and unify some well known fixed and common fixed point results.

1. INTRODUCTION

Because of its large number of applications in various branches of mathematical sciences, fixed point theory enticed many mathematicians. The most relevant result of the theory is Banach contraction principle (BCP). Several authors have studied on various generalizations of Banach contraction mapping principle. For important and famous examples of such generalizations, we may cite Edelstein [7], Kannan [12], Reich [23, 24, 25], Ćirić [5] and Suzuki [27]. Recently, BCP has been generalized in a new way by Khojasteh et al. in [15] defining \mathcal{Z} -contraction with the help of a class of control functions called simulation functions. In [19], Olgun et al. obtained the fixed point results for generalized \mathcal{Z} -contractions. Further, L-de-Hierro et al. [8] enlarged this class of simulation functions and extended some coincidence and common fixed point theorems.

All the above results, require the contraction condition or the contraction type condition to hold for every possible pair of points in domain. In order to relax this requirement in accordance with the outcome of the result Suzuki [27] contributed significantly. The results of Suzuki has inspired many researcher working in the area of metric fixed point theory (see [20],[28] and references therein). Recently, Kumam et al. in [17] defined Suzuki type \mathcal{Z} -contraction and unified some known fixed point results. In [21], Radenovic and Chandok coined the notion of (Z_G, g) -contraction and generalized the results of Jungck [10], L-de-Hierro et al. [8] and Olgun et al. [19]. Motivated by the results of [17] and [21], in this work we introduce the intuition of Suzuki type (Z_G, g) -contractions and obtain some coincidence point theorems. An application to fractional order functional differential equation is presented to illustrate the usability of the obtained result.

2. PRELIMINARIES

We now revisit some concepts and results in the literature.

Definition 2.1. [2, 21] A mapping $G : [0, \infty)^2 \rightarrow \mathbb{R}$ is called C -class function if it is continuous and satisfies the following conditions:

2000 *Mathematics Subject Classification.* 47H10, 54H25.

Key words and phrases. Coincidence Point; \mathcal{Z} -contraction; Suzuki type (Z_G, g) -contractions.

- (i) $G(s, t) \leq s$,
- (ii) $G(s, t) = s$ implies that either $s = 0$ or $t = 0$, for all $s, t \in [0, \infty)$.

Definition 2.2. [21] A mapping $G : [0, \infty)^2 \rightarrow \mathbb{R}$ has a property C_G , if there exists an $C_G \geq 0$ such that

- (i) $G(s, t) > C_G$ implies $s > t$,
- (ii) $G(t, t) \leq C_G$ for all $t \in [0, \infty)$.

Definition 2.3. [4] A simulation function is a mapping $\zeta : [0, \infty)^2 \rightarrow \mathbb{R}$ satisfying the following:

- (i) $\zeta(t, s) < s - t$ for all $t, s > 0$,
- (ii) if $\{t_n\}$ and $\{s_n\}$ are sequences in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} t_n = \lim_{n \rightarrow \infty} s_n > 0$ and $t_n < s_n$, then $\lim_{n \rightarrow \infty} \sup \zeta(t_n, s_n) < 0$.

For examples and related results on simulation functions, one may refer to [6, 8, 9, 13, 15, 14, 16, 18, 22, 26, 29].

Definition 2.4. [3, 21] A C_G -simulation function is a mapping $\zeta : [0, \infty)^2 \rightarrow \mathbb{R}$ satisfying the following:

- (a) $\zeta(t, s) < G(s, t)$ for all $t, s > 0$, where $G : [0, \infty)^2 \rightarrow \mathbb{R}$ is a C -class function;
- (b) if $\{t_n\}$ and $\{s_n\}$ are sequences in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} t_n = \lim_{n \rightarrow \infty} s_n > 0$ and $t_n < s_n$, then $\lim_{n \rightarrow \infty} \sup \zeta(t_n, s_n) < C_G$.

Let Z_G be the family of all C_G -simulation functions $\zeta : [0, \infty)^2 \rightarrow \mathbb{R}$.

Definition 2.5. [21] Let (X, d) be a metric space and $f, g : X \rightarrow X$ be self-mappings. The mapping f is called a (Z_G, g) -contraction if there exists $\zeta \in Z_G$ such that

$$\zeta(d(fx, fy), d(gx, gy)) \geq C_G \quad (2.1)$$

for all $x, y \in X$ with $gx \neq gy$.

If $g = i_X$ (identity mapping on X) and $C_G = 0$, we get \mathcal{Z} -contraction of [15].

Definition 2.6. [21] Let (X, d) be a metric space and $f, g : X \rightarrow X$ be self-mappings. The mapping f is called a generalized (Z_G, g) -contraction if there exists $\zeta \in Z_G$ such that

$$\zeta \left(d(fx, fy), \max \left\{ d(gx, gy), d(gx, fx), d(gy, fy), \frac{d(gx, fy) + d(gy, fx)}{2} \right\} \right) \geq C_G \quad (2.2)$$

for all $x, y \in X$ with $gx \neq gy$.

If $g = i_X$ (identity mapping on X) and $C_G = 0$, we get \mathcal{Z} -contraction of [19].

Definition 2.7. [11] Let f and g be self-mappings of a set X . If $u = fv = gv$ for some $v \in X$, then v is called a coincidence point of f and g and u is called a point of coincidence of f and g . The pair (f, g) is weakly compatible if the mappings commute at their coincidence points.

Proposition 2.1. [1] Let f and g be weakly compatible self-mappings of a set X . If f and g have a unique point of coincidence $u = fx = gx$, then u is the unique common fixed point of f and g .

Lemma 2.2. [21] Let (X, d) be a metric space and $\{x_n\}$ be a sequence in X such that $\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0$. If $\{x_n\}$ is not Cauchy then there exists $\varepsilon > 0$ and two subsequences $\{x_{m(k)}\}$ and $\{x_{n(k)}\}$ of $\{x_n\}$ where $n(k) > m(k) > k$ such that

$$\begin{aligned} \lim_{k \rightarrow \infty} d(x_{m(k)}, x_{n(k)}) &= \lim_{k \rightarrow \infty} d(x_{m(k)}, x_{n(k)+1}) = \lim_{k \rightarrow \infty} d(x_{m(k)-1}, x_{n(k)}) = \varepsilon \\ \text{and } \lim_{k \rightarrow \infty} d(x_{m(k)-1}, x_{n(k)+1}) &= \lim_{k \rightarrow \infty} d(x_{m(k)+1}, x_{n(k)+1}) = \varepsilon. \end{aligned}$$

3. MAIN RESULTS

In this section, we introduce the new concept of Suzuki type contractions involving C_G -simulation functions which is used to prove several coincidence point theorems in complete metric spaces.

Definition 3.1. Let (X, d) be a metric space and $f, g : X \rightarrow X$ be self-mappings. A mapping f is called a Suzuki type (Z_G, g) -contraction if there exists $\zeta \in Z_G$ such that

$$\frac{1}{2}d(gx, fx) < d(gx, gy) \implies \zeta(d(fx, fy), d(gx, gy)) \geq C_G \quad (3.1)$$

for all $x, y \in X$ with $gx \neq gy$.

If $g = i_X$ (identity mapping on X) and $C_G = 0$, we get Suzuki type \mathcal{Z} -contraction of [17].

Definition 3.2. Let (X, d) be a metric space and $f, g : X \rightarrow X$ be self-mappings. The mapping f is called a Suzuki type generalized (Z_G, g) -contraction if there exists $\zeta \in Z_G$ such that

$$\frac{1}{2}d(gx, fx) < d(gx, gy) \implies \zeta(d(fx, fy), M(x, y)) \geq C_G \quad (3.2)$$

for all $x, y \in X$ with $gx \neq gy$, where

$$M(x, y) = \max \left\{ d(gx, gy), d(gx, fx), d(gy, fy), \frac{d(gx, fy) + d(gy, fx)}{2} \right\}.$$

If $g = i_X$ (identity mapping on X) and $C_G = 0$, we get Suzuki type generalized \mathcal{Z} -contraction.

Theorem 3.1. Let (X, d) be a metric space, $f, g : X \rightarrow X$ be self-mappings and f be a Suzuki type generalized (Z_G, g) -contraction. Assume that the following conditions hold:

- (i) $f(X) \subseteq g(X)$,
- (ii) $g(X)$ or $f(X)$ is complete.

Then f and g have unique point of coincidence.

Proof. Suppose that the point of coincidence exists, then the uniqueness follows as: let u_1 and u_2 be two distinct point of coincidence of f and g i.e. $u_1 = fv_1 = gv_1$ and $u_2 = fv_2 = gv_2$ for some $v_1, v_2 \in X$. Now, since

$$\frac{1}{2}d(gv_1, fv_1) = 0 < d(gv_1, gv_2),$$

so by using (3.1), we get

$$\begin{aligned} C_G &\leq \zeta(d(fv_1, fv_2), M(v_1, v_2)) \\ &= \zeta(d(u_1, u_2), d(u_1, u_2)) < G(d(u_1, u_2), d(u_1, u_2)) \leq C_G, \end{aligned}$$

which is a contradiction. Thus, the point of coincidence is unique. Consider the sequences $\{x_n\}$ and $\{y_n\}$ defined as

$$y_n = fx_n = gx_{n+1} \text{ for all } n \in \mathbb{N} \cup \{0\},$$

and assume that $y_n \neq y_{n+1}$ for all $n \in \mathbb{N} \cup \{0\}$. Let $x = x_{n+1}, y = x_{n+2}$, then

$$\frac{1}{2}d(gx_{n+1}, fx_{n+1}) = \frac{1}{2}d(gx_{n+1}, gx_{n+2}) < d(gx_{n+1}, gx_{n+2}).$$

So from (3.2), we have

$$\begin{aligned} C_G &\leq \zeta\left(d(fx_{n+1}, fx_{n+2}), M(x_{n+1}, x_{n+2})\right) \\ &= \zeta\left(d(y_{n+1}, y_{n+2}), \max\{d(y_n, y_{n+1}), d(y_{n+1}, y_{n+2})\}\right) \\ &< G\left(\max\{d(y_n, y_{n+1}), d(y_{n+1}, y_{n+2})\}, d(y_{n+1}, y_{n+2})\right). \end{aligned}$$

Further, using (i) of Definition 2.2, we have

$$\max\{d(y_n, y_{n+1}), d(y_{n+1}, y_{n+2})\} > d(y_{n+1}, y_{n+2}).$$

Hence for all $n \in \mathbb{N} \cup \{0\}$, we have $d(y_n, y_{n+1}) > d(y_{n+1}, y_{n+2})$. So $\{d(y_n, y_{n+1})\}$ is a monotonically decreasing sequence of nonnegative real numbers, and hence there exists $l \geq 0$ such that $\lim_{n \rightarrow \infty} d(y_n, y_{n+1}) = l$. Assume that $l > 0$. Since $d(y_{n+1}, y_{n+2}) < M(x_{n+1}, x_{n+2})$, both $d(y_{n+1}, y_{n+2})$ and $M(x_{n+1}, x_{n+2})$ tends to l as $n \rightarrow \infty$ and

$$\frac{1}{2}d(y_n, y_{n+1}) = \frac{1}{2}d(gx_{n+1}, fx_{n+1}) < d(gx_{n+1}, gx_{n+2}) = d(y_n, y_{n+1}),$$

then using (3.2) and (b) of Definition 2.4, we get

$$C_G \leq \lim_{n \rightarrow \infty} \sup \zeta\left(d(y_{n+1}, y_{n+2}), M(x_{n+1}, x_{n+2})\right) < C_G,$$

which is a contradiction and hence $l = 0$. Now if we assume that $y_n = y_m$ for some $n > m$. Then we can choose $x_{n+1} = x_{m+1}$ and hence also $y_{n+1} = y_{m+1}$. Then using the similar arguments as above we get

$$d(y_n, y_{n+1}) < d(y_{n-1}, y_n) < \cdots < d(y_m, y_{m+1}) = d(y_n, y_{n+1}),$$

which is a contradiction. Similarly if $y_n = y_m$ for some $n < m$, we get a contradiction. Hence $y_n \neq y_m$ for all $n \neq m$.

Now we prove that $\{y_n\}$ is a Cauchy sequence. If not, then by Lemma 2.2 we have

$$\lim_{k \rightarrow \infty} d(x_{m(k)}, x_{n(k)}) = \lim_{k \rightarrow \infty} d(x_{m(k)+1}, x_{n(k)+1}) = \varepsilon,$$

and consequently,

$$\lim_{k \rightarrow \infty} M(x_{m(k)+1}, x_{n(k)+1}) = \varepsilon.$$

We claim that there exists $k_0 \in \mathbb{N}$ such that

$$\frac{1}{2}d(y_{m(k)}, y_{m(k)+1}) < d(y_{m(k)}, y_{n(k)}),$$

$$\text{i.e. } \frac{1}{2}d(gx_{m(k)+1}, fx_{m(k)+1}) < d(gx_{m(k)+1}, gx_{n(k)+1})$$

for all $k \geq k_0$. If not, then letting $k \rightarrow \infty$, we get $\varepsilon \leq 0$, which is not true. Also, using (3.2) and (i) of Definition 2.2, we have $d(y_{m(k)+1}, y_{n(k)+1}) < M(x_{m(k)+1}, x_{n(k)+1})$, then by (b) of Definition 2.4, we get

$$C_G \leq \lim_{k \rightarrow \infty} \sup \zeta \left(d(y_{m(k)+1}, y_{n(k)+1}), M(x_{m(k)+1}, x_{n(k)+1}) \right) < C_G$$

a contradiction. Hence $\{y_n\}$ is a Cauchy sequence.

Suppose $g(X)$ is complete subspace of X . Then $\{y_n\}$ being contained in $g(X)$ has a limit in $g(X)$, say $z \in X$ such that $y_n \rightarrow gz$, i.e. $gx_n \rightarrow gz$ as $n \rightarrow \infty$. We will show that z is the coincidence point of f and g . We can suppose $y_n \neq fz, gz$ for $n \in \mathbb{N} \cup \{0\}$, otherwise we are done. We claim that

$$\frac{1}{2}d(gx_n, gx_{n+1}) < d(gx_n, gz) \text{ or } \frac{1}{2}d(gx_{n+1}, gx_{n+2}) < d(gx_{n+1}, gz) \quad (3.3)$$

for every $n \in \mathbb{N} \cup \{0\}$. If not, then there exists $m \in \mathbb{N}$ for which

$$\frac{1}{2}d(gx_m, gx_{m+1}) \geq d(gx_m, gz) \text{ and } \frac{1}{2}d(gx_{m+1}, gx_{m+2}) \geq d(gx_{m+1}, gz)$$

holds. Then

$$2d(gx_m, gz) \leq d(gx_m, gx_{m+1}) \leq d(gx_m, gz) + d(gz, gx_{m+1})$$

which implies $d(gx_m, gz) \leq d(gz, gx_{m+1}) \leq \frac{1}{2}d(gx_{m+1}, gx_{m+2})$. Further since

$$\frac{1}{2}d(gx_m, fx_m) = \frac{1}{2}d(gx_m, gx_{m+1}) < d(gx_m, gx_{m+1}),$$

so from (3.1), we have

$$C_G \leq \zeta(d(fx_m, fx_{m+1}), M(x_m, x_{m+1})) < G(M(x_m, x_{m+1}), d(fx_m, fx_{m+1})),$$

and in view of (i) of Definition 2.2, we get $M(x_m, x_{m+1}) = d(gx_m, gx_{m+1}) > d(fx_m, fx_{m+1})$. Now

$$\begin{aligned} d(fx_m, fx_{m+1}) &< d(gx_m, gx_{m+1}) \\ &\leq d(gx_m, gz) + d(gz, gx_{m+1}) \\ &\leq \frac{1}{2}d(gx_{m+1}, gx_{m+2}) + \frac{1}{2}d(gx_{m+1}, gx_{m+2}) \\ &= d(gx_{m+1}, gx_{m+2}) = d(fx_m, fx_{m+1}), \end{aligned}$$

which is a contradiction. Hence (3.3) holds. Now from (3.3) and (3.1) we have

$$C_G \leq \zeta \left(d(fx_n, fz), M(x_n, z) \right) < G \left(M(x_n, z), d(fx_n, fz) \right)$$

and then by (i) of Definition 2.2, we get $d(fx_n, fz) < M(x_n, z)$. Now, if we assume $fz \neq gz$, then taking limit as $n \rightarrow \infty$, we get $d(gz, fz) < d(gz, fz)$, a contradiction. Hence $gz = fz$. In same manner if $\frac{1}{2}d(gx_{n+1}, gx_{n+2}) < d(gx_{n+1}, gv)$ we can show that $gz = fz$. In case $f(X)$ is a complete subspace of X , the sequence $\{y_n\}$ converges in $g(X)$ since $f(X) \subseteq g(X)$. So the previous argument works. \square

Theorem 3.2. Let (X, d) be a complete metric space, $f, g : X \rightarrow X$ be self-mappings and f be a Suzuki type generalized (Z_G, g) -contraction. Assume that the following conditions hold:

- (i) $f(X) \subseteq g(X)$,
- (ii) g is a continuous,
- (iii) f and g are commuting.

Then f and g have unique point of coincidence.

Proof. If the coincidence point exist then it is unique (as proved in Theorem 3.1). Construct the sequences $\{x_n\}$ and $\{y_n\}$ as

$$y_n = fx_n = gx_{n+1} \text{ for all } n \in \mathbb{N} \cup \{0\}.$$

Then the sequence $\{y_n\}$ is Cauchy (as shown in proof of Theorem 3.1).

Since the space (X, d) is complete, there exists $z \in X$ such that $y_n \rightarrow z$ or $gx_n \rightarrow z$ as $n \rightarrow \infty$. As g is continuous, $g^2x_n \rightarrow gz$ as $n \rightarrow \infty$. We claim that

$$\frac{1}{2}d(ggx_n, ggx_{n+1}) < d(ggx_n, ggz) \text{ or } \frac{1}{2}d(ggx_{n+1}, ggx_{n+2}) < d(ggx_{n+1}, ggz) \quad (3.4)$$

for every $n \in \mathbb{N} \cup \{0\}$. If not, then there exists $m \in \mathbb{N}$ for which

$$\frac{1}{2}d(ggx_m, ggx_{m+1}) \geq d(ggx_m, ggz) \text{ and } \frac{1}{2}d(ggx_{m+1}, ggx_{m+2}) \geq d(ggx_{m+1}, ggz)$$

holds. Then

$$2d(ggx_m, ggz) \leq d(ggx_m, ggx_{m+1}) \leq d(ggx_m, ggz) + d(ggz, ggx_{m+1})$$

which implies

$$d(ggx_m, ggz) \leq d(ggz, ggx_{m+1}) \leq \frac{1}{2}d(ggx_{m+1}, ggx_{m+2}).$$

Further since f and g are commutative, we have

$$\frac{1}{2}d(ggx_m, fgx_m) = \frac{1}{2}d(ggx_m, gfx_m) < d(ggx_m, ggx_{m+1}).$$

So from (3.2), we have

$$C_G \leq \zeta\left(d(fgx_m, fgx_{m+1}), M(gx_m, gx_{m+1})\right) < G\left(M(gx_m, gx_{m+1}), d(fgx_m, fgx_{m+1})\right)$$

and in view of (i) of Definition 2.2, we get $M(gx_m, gx_{m+1}) = d(ggx_m, ggx_{m+1}) > d(fgx_m, fgx_{m+1})$. Now

$$\begin{aligned} d(fgx_m, fgx_{m+1}) &< d(ggx_m, ggx_{m+1}) \\ &\leq d(ggx_m, ggz) + d(ggz, ggx_{m+1}) \\ &\leq \frac{1}{2}d(ggx_{m+1}, ggx_{m+2}) + \frac{1}{2}d(ggx_{m+1}, ggx_{m+2}) \\ &= d(ggx_{m+1}, ggx_{m+2}) = d(fgx_m, fgx_{m+1}), \end{aligned}$$

which is a contradiction. Hence (3.4) holds. Now from (3.4) and (3.2), we have

$$C_G \leq \zeta\left(d(fgx_n, fz), M(gx_n, z)\right) < G\left(M(gx_n, z), d(fgx_n, fz)\right)$$

and then by (i) of Definition 2.2 and f, g are commuting, we get $d(fgx_n, fz) = d(gfx_n, fz) < M(gx_n, z)$. Now, if we assume $fz \neq gz$, then taking limit as $n \rightarrow \infty$ and by continuity of g , we get $d(gz, fz) < d(gz, fz)$, a contradiction. Hence $gz = fz$. In same manner if $\frac{1}{2}d(ggx_{n+1}, ggx_{n+2}) < d(ggx_{n+1}, ggz)$ we can show that $gz = fz$. \square

Theorem 3.3. Let (X, d) be a metric space, $f, g : X \rightarrow X$ be self-mappings and f be a Suzuki type (Z_G, g) -contraction. Assume that the following conditions hold:

- (i) $f(X) \subseteq g(X)$,
- (ii) $g(X)$ or $f(X)$ is complete.

Then f and g have unique point of coincidence.

Proof. If the coincidence point exist then it is unique (as proved in Theorem 3.1). Construct the sequences $\{x_n\}$ and $\{y_n\}$ as

$$y_n = fx_n = gx_{n+1} \text{ for all } n \in \mathbb{N} \cup \{0\}.$$

Then the sequence $\{y_n\}$ is Cauchy (as shown in proof of Theorem 3.1).

Suppose $g(X)$ is complete subspace of X . Then $\{y_n\}$ being contained in $g(X)$ has a limit in $g(X)$, say $z \in X$ such that $y_n \rightarrow gz$, i.e. $gx_n \rightarrow gz$ as $n \rightarrow \infty$. We will show that z is the coincidence point of f and g . We can suppose $y_n \neq fz$ and $y_n \neq gz$ for $n \in \mathbb{N} \cup \{0\}$, otherwise we are done. We also claim that

$$\frac{1}{2}d(gx_n, gx_{n+1}) < d(gx_n, gz) \text{ or } \frac{1}{2}d(gx_{n+1}, gx_{n+2}) < d(gx_{n+1}, gz) \quad (3.5)$$

for every $n \in \mathbb{N} \cup \{0\}$. If not, then there exists $m \in \mathbb{N}$ for which

$$\frac{1}{2}d(gx_m, gx_{m+1}) \geq d(gx_m, gz) \text{ and } \frac{1}{2}d(gx_{m+1}, gx_{m+2}) \geq d(gx_{m+1}, gz)$$

holds. Then

$$2d(gx_m, gz) \leq d(gx_m, gx_{m+1}) \leq d(gx_m, gz) + d(gz, gx_{m+1})$$

which implies $d(gx_m, gz) \leq d(gz, gx_{m+1}) \leq \frac{1}{2}d(gx_{m+1}, gx_{m+2})$. Further since

$$\frac{1}{2}d(gx_m, fx_m) = \frac{1}{2}d(gx_m, gx_{m+1}) < d(gx_m, gx_{m+1}),$$

so from (3.1), we have

$$C_G \leq \zeta(d(fx_m, fx_{m+1}), d(gx_m, gx_{m+1})) < G(d(gx_m, gx_{m+1}), d(fx_m, fx_{m+1})),$$

and in view of (i) of Definition 2.2, we get $d(gx_m, gx_{m+1}) > d(fx_m, fx_{m+1})$.

$$\begin{aligned} \text{Now } d(fx_m, fx_{m+1}) &< d(gx_m, gx_{m+1}) \\ &\leq d(gx_m, gz) + d(gz, gx_{m+1}) \\ &\leq \frac{1}{2}d(gx_{m+1}, gx_{m+2}) + \frac{1}{2}d(gx_{m+1}, gx_{m+2}) \\ &= d(gx_{m+1}, gx_{m+2}) = d(fx_m, fx_{m+1}) \end{aligned}$$

which is a contradiction. Hence (3.5) holds. Now from (3.5) and (3.1), we have

$$C_G \leq \zeta(d(fx_n, fz), d(gx_n, gz)) < G(d(gx_n, gz), d(fx_n, fz))$$

and then by (i) of Definition 2.2, we get $d(fx_n, fz) < d(gx_n, gz)$. Taking limit as $n \rightarrow \infty$ we get $fx_n \rightarrow fz$. Hence $gz = fz$. In same manner if $\frac{1}{2}d(gx_{n+1}, gx_{n+2}) < d(gx_{n+1}, gz)$, we can show that $gz = fz$. In case $f(X)$ is a complete subspace of X , the sequence $\{y_n\}$ converges in $g(X)$ since $f(X) \subseteq g(X)$. So the previous argument works. \square

Theorem 3.4. Let (X, d) be a complete metric space, $f, g : X \rightarrow X$ be self-mappings and f be a Suzuki type (Z_G, g) -contraction. Assume that the following conditions hold:

- (i) $f(X) \subseteq g(X)$,
- (ii) g is a continuous,
- (iii) f and g are commuting.

Then f and g have unique point of coincidence.

Proof. The proof follows in the same manner as Theorem 3.2 and 3.3. \square

Theorem 3.5. In addition to hypotheses of Theorem 3.1-3.4, if f and g are weakly compatible, then they have a unique common fixed point in X .

Proof. The proof follows from Proposition 2.1. \square

By taking $g = i_X$ (identity mapping on X) and $C_G = 0$ in Theorem 3.5, we get the following result of Kumam et al. [17] as a corollary.

Corollary 3.6. Let (X, d) be a complete metric space and $f : X \rightarrow X$ be a Suzuki type \mathcal{Z} -contraction with respect to ζ . Then f has a unique fixed point x^* in X .

Remark 3.1. We observed that the Property (K) assumed in the main result (Theorem 2.1) of Kumam et al. [17] is not required if the mapping f is asymptotically regular at every $x \in X$.

Example 3.1. Let $X = [0, 1]$ and d be the usual metric on X . Define $f, g : X \rightarrow X$ as follows:

$$fx = \begin{cases} 0 & \text{if } x \in [0, \frac{1}{4}], \\ \frac{1}{20} & \text{if } x \in (\frac{1}{4}, \frac{1}{2}), \\ \frac{4}{25} & \text{if } x \in [\frac{1}{2}, 1] \end{cases} \quad \text{and} \quad gx = \begin{cases} \frac{x}{3} & \text{if } x \in [0, \frac{1}{4}], \\ \frac{1}{4} & \text{if } x \in (\frac{1}{4}, \frac{1}{2}), \\ \frac{1}{3} & \text{if } x \in [\frac{1}{2}, 1]. \end{cases}$$

Consider the functions $\zeta(t, s) = \frac{4}{5}s - t$, $G(s, t) = s - t$ and $C_G = 0$, then for $x \in (\frac{1}{4}, \frac{1}{2})$ and $y \in [\frac{1}{2}, 1]$, the mapping f and g do not satisfy the condition (2.1) and hence Theorem 2.1 of Radenovic and Chandok [21] is not applicable. However, f is a Suzuki type (Z_G, g) -contraction as well as Suzuki type generalized (Z_G, g) -contraction and satisfy all the assumption of Theorem 3.1 and 3.3. Here f and g have a unique coincidence point $x = 0$. Also, the mappings are weakly compatible.

Example 3.2. Let $X = \{(0, 0), (3, 3), (4, 0), (0, 4), (4, 5), (5, 4)\}$ be a metric space endowed with the metric d defined by

$$d((x_1, x_2), (y_1, y_2)) = |x_1 - y_1| + |x_2 - y_2|.$$

Let $f, g : X \rightarrow X$ defined by

$$f(x_1, x_2) = \begin{cases} (0, 0) & \text{if } x_1 = x_2, \\ (x_1, 0) & \text{if } x_1 < x_2, \\ (0, x_2) & \text{if } x_1 > x_2, \end{cases}$$

$$g(x_1, x_2) = \begin{cases} (4, 0) & \text{if } x_1 = x_2 = 3, \\ (x_1, x_2) & \text{otherwise.} \end{cases}$$

Consider the functions $\zeta(t, s) = \frac{4}{5}s - t$, $G(s, t) = s - t$ and $C_G = 0$, then we note that

$$0 \leq \zeta(d(fx, fy), M(x, y))$$

holds if $(x, y) \notin \left\{ \left((4, 5), (5, 4) \right), \left((5, 4), (4, 5) \right) \right\}$. Since in this case

$$\frac{1}{2}d(gx, fx) \geq d(gx, gy),$$

so f is a Suzuki type generalized (Z_G, g) -contraction. Also, f and g satisfy all hypothesis of Theorem 3.5 with the unique common fixed point $(x_1, x_2) = (0, 0)$. Notice that

$$\zeta\left(d(fx, fy), M(x, y)\right) < 0$$

whenever $(x, y) \in \left\{\left((4, 5), (5, 4)\right), \left((5, 4), (4, 5)\right)\right\}$. So the mapping f does not satisfy the condition (2.1) and (2.2) and hence Theorem 2.1 and 2.3 of Radenovic and Chandok [21] are not applicable.

4. COINCIDENCE POINTS OF RATIONAL TYPE QUASI-CONTRACTIONS

If f and g be two self-mappings on a metric space (X, d) satisfying $f(X) \subseteq g(X)$ and $x_0 \in X$, let us define $x_1 \in X$ such that $fx_0 = gx_1$. Having defined $x_n \in X$, let $x_{n+1} \in X$ be such that $fx_n = gx_{n+1}$. We say that $\{fx_n\}$ is a f - g -sequence of initial point x_0 . Letting $fx_n = gx_{n+1} = y_n$, where $n \in \mathbb{N} \cup \{0\}$, denote $\mathcal{O}(y_k; n) = \{y_k, y_{k+1}, y_{k+2}, \dots, y_{k+n}\}$. Let $\delta[\mathcal{O}(y_k; n)]$ denotes the diameter of $\mathcal{O}(y_k; n)$. If $\delta[\mathcal{O}(y_k; n)] > 0$ for $k, n \in \mathbb{N}$ then we have $\delta[\mathcal{O}(y_k; n)] = d(y_i, y_j)$, where $k \leq i < j \leq k + n$.

Definition 4.1. A mapping $G : [0, \infty)^2 \rightarrow \mathbb{R}$ has a property C_G^m , if there exists an $C_G \geq 0$ such that

- (i) $G(s, t) > C_G$ implies $\frac{s}{s+1} > t$,
- (ii) $G(t, t) \leq C_G$ for all $t \in [0, \infty)$.

Denote $m(x, y) = \max\{d(gx, gy), d(gx, fx), d(gy, fy), d(gx, fy), d(gy, fx)\}$. Using the above property we prove following result:

Theorem 4.1. Let (X, d) be a metric space and $f, g : X \rightarrow X$ be self-mappings satisfying the condition

$$C_G \leq \frac{m(x, y)}{m(x, y) + 1} - d(fx, fy) < G(m(x, y), d(fx, fy)) \quad (4.1)$$

where $G : [0, \infty)^2 \rightarrow \mathbb{R}$ is a C -class function. Assume that the following conditions hold:

- (i) $f(X) \subseteq g(X)$,
- (ii) $g(X)$ or $f(X)$ is complete.
- (iii) if $\{t_n\}$ and $\{s_n\}$ are sequences in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} t_n = \lim_{n \rightarrow \infty} s_n > 0$

$$\text{and } t_n < \frac{s_n}{s_n + 1}, \text{ then } \lim_{n \rightarrow \infty} \sup \left(\frac{s_n}{s_n + 1} - t_n \right) < C_G.$$

Then f and g have unique point of coincidence.

Proof. Suppose that the point of coincidence exists, then the uniqueness follows as: let u_1 and u_2 be two distinct points of coincidence of f and g , i.e. $u_1 = fv_1 = gv_1$ and $u_2 = fv_2 = gv_2$ for some $v_1, v_2 \in X$. Then using (4.1) and (ii) of Definition 4.1, we get

$$\begin{aligned} C_G &\leq \frac{m(v_1, v_2)}{m(v_1, v_2) + 1} - d(fv_1, fv_2) < G\left(m(v_1, v_2), d(fv_1, fv_2)\right) \\ &\implies C_G < G(d(u_1, u_2), d(u_1, u_2)) \leq C_G, \end{aligned}$$

which is a contradiction. Thus, the point of coincidence is unique.

Now, consider the sequences $\{x_n\}$ and $\{y_n\}$ defined as

$$y_n = fx_n = gx_{n+1} \text{ for all } n \in \mathbb{N} \cup \{0\},$$

and assume that $y_n \neq y_{n+1}$ for all $n \in \mathbb{N} \cup \{0\}$. Taking $x = x_n, y = x_{n+1}$ in (4.1), we have

$$C_G \leq \frac{m(x_n, x_{n+1})}{m(x_n, x_{n+1}) + 1} - d(fx_n, fx_{n+1}) < G\left(m(x_n, x_{n+1}), d(fx_n, fx_{n+1})\right).$$

Using (i) of Definition 4.1, we get

$$d(fx_n, fx_{n+1}) < \frac{m(x_n, x_{n+1})}{m(x_n, x_{n+1}) + 1}$$

which gives

$$\begin{aligned} d(y_n, y_{n+1}) &< \frac{\max\{d(y_{n-1}, y_n), d(y_n, y_{n+1}), d(y_{n-1}, y_{n+1})\}}{\max\{d(y_{n-1}, y_n), d(y_n, y_{n+1}), d(y_{n-1}, y_{n+1})\} + 1} \\ \text{i.e. } \delta[\mathcal{O}(y_n; 1)] &< \frac{\delta[\mathcal{O}(y_{n-1}; 2)]}{\delta[\mathcal{O}(y_{n-1}; 2)] + 1} \\ \text{or } \delta[\mathcal{O}(y_n; 1)] &< \lambda_1 \delta[\mathcal{O}(y_{n-1}; 2)], \end{aligned} \quad (4.2)$$

where $\lambda_1 = \frac{1}{\delta[\mathcal{O}(y_{n-1}; 2)] + 1} < 1$. Again using (4.1) and (i) of Definition 4.1, we get

$$\begin{aligned} \delta[\mathcal{O}(y_{n-1}; 2)] &= d(y_r, y_p), \text{ where } n-1 \leq r < p \leq n+1 \\ &< \frac{\max\{d(y_{r-1}, y_{p-1}), d(y_{r-1}, y_r), d(y_{p-1}, y_p), d(y_{r-1}, y_p), d(y_{p-1}, y_r)\}}{\max\{d(y_{r-1}, y_{p-1}), d(y_{r-1}, y_r), d(y_{p-1}, y_p), d(y_{r-1}, y_p), d(y_{p-1}, y_r)\} + 1} \\ \text{i.e. } \delta[\mathcal{O}(y_{n-1}; 2)] &< \frac{\delta[\mathcal{O}(y_{n-2}; 3)]}{\delta[\mathcal{O}(y_{n-2}; 3)] + 1} = \lambda_2 \delta[\mathcal{O}(y_{n-2}; 3)]. \end{aligned} \quad (4.3)$$

From (4.2) and (4.3), we have

$$\delta[\mathcal{O}(y_n; 1)] < \lambda_1 \lambda_2 \delta[\mathcal{O}(y_{n-2}; 3)]. \quad (4.4)$$

Continuing in same manner, we get

$$\delta[\mathcal{O}(y_n; 1)] < \lambda_1 \lambda_2 \dots \lambda_n \delta[\mathcal{O}(y_0; n+1)], \quad (4.5)$$

where $\delta[\mathcal{O}(y_0; n+1)] = d(y_r, y_q)$ for some positive $0 \leq r < q \leq n+1$ and $\lambda_i = \frac{1}{\delta[\mathcal{O}(y_{n-i}; i+1)] + 1} < 1$. Let $\lambda = \max\{\lambda_1, \lambda_2, \dots, \lambda_n\}$, then we have

$$\delta[\mathcal{O}(y_n; 1)] < \lambda^n \delta[\mathcal{O}(y_0; n+1)]. \quad (4.6)$$

Now

$$\begin{aligned} \delta[\mathcal{O}(y_0; n+1)] &= d(y_r, y_q) \leq d(y_r, y_{r+1}) + d(y_{r+1}, y_q) \\ &= d(y_r, y_{r+1}) + d(fx_{r+1}, fx_q) \\ &\leq d(y_r, y_{r+1}) + \delta[\mathcal{O}(y_1, n)] \\ &\leq d(y_r, y_{r+1}) + \frac{\delta[\mathcal{O}(y_0; n+1)]}{\delta[\mathcal{O}(y_0; n+1)] + 1} \\ &= d(y_r, y_{r+1}) + \beta \delta[\mathcal{O}(y_0; n+1)], \end{aligned}$$

where $\beta = \frac{1}{\delta[\mathcal{O}(y_0; n+1)] + 1} < 1$. Hence

$$\delta[\mathcal{O}(y_0; n+1)] \leq \frac{1}{1-\beta} d(y_r, y_{r+1}) \quad (4.7)$$

Thus, from (4.6) and (4.7), we have

$$\delta[\mathcal{O}(y_n; 1)] < \frac{\lambda^n}{1-\beta} d(y_0, y_1) \quad (4.8)$$

with $r = 0$. Therefore, because of $\lambda < 1$, we get

$$d(y_n, y_{n+1}) \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (4.9)$$

Now, we claim that the sequence $\{y_n\}$ is Cauchy. If not, then by Lemma 2.2 we have

$$\lim_{k \rightarrow \infty} d(x_{m(k)}, x_{n(k)}) = \lim_{k \rightarrow \infty} d(x_{m(k)+1}, x_{n(k)+1}) = \varepsilon,$$

and consequently,

$$\lim_{k \rightarrow \infty} m(x_{m(k)+1}, x_{n(k)+1}) = \varepsilon.$$

Also, from (4.1) and (i) of Definition 4.1, we have

$$d(y_{m(k)+1}, y_{n(k)+1}) < \frac{m(x_{m(k)+1}, x_{n(k)+1})}{m(x_{m(k)+1}, x_{n(k)+1}) + 1}.$$

So by assumption (iii), we get

$$C_G \leq \limsup_{k \rightarrow \infty} \left(\frac{m(x_{m(k)+1}, x_{n(k)+1})}{m(x_{m(k)+1}, x_{n(k)+1}) + 1} - d(y_{m(k)+1}, y_{n(k)+1}) \right) < C_G, \quad (4.10)$$

a contradiction. Hence $\{y_n\}$ is a Cauchy sequence.

Now, suppose $g(X)$ is complete subspace of X . Then there exists a point, say $z \in X$ such that $y_n \rightarrow gz$, i.e. $gx_n \rightarrow gz$ as $n \rightarrow \infty$. We will show that z is the coincidence point of f and g . We can suppose $y_n \neq fz, gz$ for $n \in \mathbb{N} \cup \{0\}$, otherwise we are done.

Now, using (4.1) for $x = x_n$ and $y = z$, we have

$$C_G \leq \frac{m(x_n, z)}{m(x_n, z) + 1} - d(fx_n, fz) < G\left(m(x_n, z), d(fx_n, fz)\right),$$

and then by (i) of Definition 4.1, we get $d(fx_n, fz) < \frac{m(x_n, z)}{m(x_n, z) + 1}$. If we assume $fz \neq gz$, then taking limit as $n \rightarrow \infty$, we get $d(gz, fz) < d(gz, fz)$, a contradiction. Hence $gz = fz$. In case $f(X)$ is a complete subspace of X , the sequence $\{y_n\}$ converges in $g(X)$ since $f(X) \subseteq g(X)$. In same manner we can show that $gz = fz$. \square

Example 4.1. Let $X = (-2, 2)$ be a metric space endowed with metric $d(x, y) = |x - y|$. Let f and g be self-mappings on X defined by

$$f(x) = \begin{cases} 0 & \text{if } -2 < x < -1 \text{ and } 1 < x < 2, \\ \frac{1+x}{3} & \text{if } -1 \leq x < 0, \\ \frac{1-x}{3} & \text{if } 0 \leq x \leq 1, \end{cases}$$

$$g(x) = \begin{cases} 1 & \text{if } -2 < x \leq -1, \\ x & \text{if } -1 < x < 1, \\ -1 & \text{if } 1 \leq x < 2. \end{cases}$$

Here Jungck's theorem [11] is not applicable since the mapping g is discontinuous. However, for $G(s, t) = s - t$ and $C_G = 0$, f and g satisfy all hypothesis of Theorem 4.1 with the unique coincidence point $x = \frac{1}{4}$.

Remark 4.1. Let $m(x, y) = s$ and $d(fx, fy) = t$ for $x, y \in X$ with the metric d . Define a function $\zeta : [0, \infty)^2 \rightarrow \mathbb{R}$ by $\zeta(t, s) = \frac{s}{s+1} - t$, then clearly $\zeta \in Z_G$. We also observed that the Theorem 4.1 is an approach to provide an partial answer to the question posed by Radenovic and Chandok in [21].

Let $m(x, y) = s$ and $d(fx, fy) = t$ for $x, y \in X$ with the metric d . Define a function $\zeta : [0, \infty)^2 \rightarrow \mathbb{R}$ by $\zeta(t, s) = \frac{s}{s+1} - t$, then clearly $\zeta \in Z_G$. We also observed that the Theorem 4.1 is an approach to provide an partial answer to the question posed by Radenovic and Chandok in [21].

We conclude this section with the following open question:

Question:

Whether the Theorem 4.1 holds when the condition (4.1) is replaced with

$$\frac{1}{2}d(gx, fx) < d(gx, gy) \Rightarrow C_G \leq \frac{m(x, y)}{m(x, y) + 1} - d(fx, fy) < G(m(x, y), d(fx, fy))?$$

REFERENCES

- [1] Abbas M., Jungck J., Common fixed point results for non-commuting mappings without continuity in cone metric spaces, J. Math. Anal. Appl., 2008, 341, 416-420.
- [2] Ansari A., Note on φ - ψ -contractive type mappings and related fixed point, The 2nd Regional Conference on Math. Appl. PNU 377-380 (Sept. 2014).
- [3] Ansari A., Isik H., Radenovic S., Coupled fixed point theorems for contractive mappings involving new function classes and applications, Filomat, 2017, 31 (7), 1893-1907.
- [4] Argoubi H., Samet B., Vetro C., Nonlinear contractions involving simulation functions in a metric space with a partial order, J. Nonlinear Sci. Appl., 2015, 8, 1082-1094.
- [5] Ciric L., A generalization of Banachs contraction principle, Proc. Am. Math. Soc., 1974, 45, 267-273.
- [6] Chen J., Tang X., Generalizations of Darbos fixed point theorem via simulation functions with application to functional integral equations, J. Comput. Appl. Math., 2016, 296, 564-575.
- [7] Edelstein M., On fixed and periodic points under contractive mappings, J. of the London Math. Soc. II, 1962, 37, 74-79.
- [8] R.-L.-de-Hierro A., Karapinar E., R.-L.-de-Hierro C., Moreno J., Coincidence point theorems on metric spaces via simulation functions, J. Comput. Appl. Math., 2015, 275, 345-355.
- [9] R.-L.-de-Hierro A., Samet B., φ -admissibility results via extended simulation functions, J. Fixed Point Theory Appl., 2016, DOI 10.1007/s11784-016-0385.
- [10] Jungck G., Commuting maps and fixed points, Amer. Math. Month., 1976, 83, 261-263.
- [11] Jungck G., Common fixed points for noncontinuous nonself maps on nonmetric spaces, Far East J. Math. Sci., 1996, 4, 199-215.
- [12] Kannan R., Some results on fixed points, Bull. of the Calcutta Math. Soc., 1968, 60, 71-76.
- [13] Karapinar E., Fixed points results via simulation functions, Filomat, 2016, 30 (8), 2343-2350.
- [14] Karapinar E., Khojasteh F., An approach to best proximity points results via simulation functions, J. Fixed Point Theory Appl., 2016, DOI 10. 1007/s11784-016-0380-2.
- [15] Khojasteh F., Shukla S., Radenovic S., A new approach to the study of fixed point theorems via simulation functions, Filomat, 2015, 29(6), 1189-1194.

- [16] Komal S., Kumam P., Gopal D., Best proximity point for \mathcal{Z} -contraction and Suzuki type \mathcal{Z} -contraction mappings with an application to fractional calculus, *Applied General Topology*, 2016, 17, 185-198.
- [17] Kumam P., Gopal D., Budhia L., A new fixed point theorem under Suzuki type \mathcal{Z} -contraction mapping, *Journal of Mathematical Analysis*, 2017, 8 (1), 113-119.
- [18] Nastasi A., Vetro P., Fixed point results on metric and partial metric spaces via simulation functions, *J. Nonlinear Sci. Appl.*, 2015, 8, 1059-1069.
- [19] Olgun M., Bicer O., Alyildiz T., A new aspect to Picard operators with simulation functions, *Turk. J. Math.*, 2016, 40, 832-837.
- [20] Pant R., Singh S., Mishra S., A coincidence and fixed point theorems For semi-quasi contractions, *Fixed Point Theory*, 2016, 17 (2), 449-456.
- [21] Radenovic S., Chandok S., Simulation type functions and coincidence point results, *Filomat* (to appear).
- [22] Radenovic S., Vetro F., Vujakovic J., An alternative and easy approach to fixed point results via simulation functions, *Demonstr. Math.*, 2017, 1-9.
- [23] Reich S., Kannans fixed point theorem, *Boll. Un. Mat. Ital.*, 1971, 4, 1-11.
- [24] Reich S., Some remarks concerning contraction mappings, *Canadian Mathematical Bulletin*, 1971, 14, 121-124.
- [25] Reich S., Fixed points of contractive functions, *Boll. Un. Mat. Ital.*, 1972, 5, 26-42
- [26] Samet B., Best proximity point results in partially ordered metric spaces via simulation functions, *Fixed Point Theory Appl.*, 2015, Article ID 232.
- [27] Suzuki T., A generalized Banach contraction principle that characterizes metric completeness, *Proc. Amer. Math. Soc.*, 2008 136 (5), 1861-1869.
- [28] Suzuki T., A new type of fixed point theorem in metric spaces, *Nonlinear Anal.*, 2009, 71, 5313-5317.
- [29] Tchier F., Vetro C., Vetro F., Best approximation and variational inequality problems involving a simulation function, *Fixed Point Theory Appl.* 2016, Article ID 26.

D.K. PATEL

DEPARTMENT OF MATHEMATICS, VISVESVARAYA NATIONAL INSTITUTE OF TECHNOLOGY, NAGPUR-440010, INDIA

E-mail address: `deepesh456@gmail.com`

P.R. PATLE

DEPARTMENT OF MATHEMATICS, VISVESVARAYA NATIONAL INSTITUTE OF TECHNOLOGY, NAGPUR-440010, INDIA

E-mail address: `pradip.patle12@gmail.com`

L. BUDHIA

DEPARTMENT OF APPLIED MATHEMATICS & HUMANITIES, S.V. NATIONAL INSTITUTE OF TECHNOLOGY, SURAT-395007, INDIA

E-mail address: `lokesh86budhia@gmail.com`

D. GOPAL

DEPARTMENT OF APPLIED MATHEMATICS & HUMANITIES, S.V. NATIONAL INSTITUTE OF TECHNOLOGY, SURAT-395007, INDIA

E-mail address: `gopaldhananjay@yahoo.in`