CONNECTING COMMUTING NORMAL MATRICES

FREDY VIDES

ABSTRACT. In this document we study the local path connectivity of sets of m-tuples of commuting normal matrices with some additional geometric constraints in their joint spectra. In particular, given $\varepsilon > 0$ and any fixed but arbitrary m-tuple $\mathbf{X} \in M_n(\mathbb{C})^m$ in the set of m-tuples of pairwise commuting normal matrix contractions, we prove the existence of paths between arbitrary m-tuples in the intersection of the previously mentioned sets of m-tuples in $M_n(\mathbb{C})^m$ and the δ -ball $B_{\overline{0}}(\mathbf{X}, \delta)$ centered at \mathbf{X} for some $\delta > 0$, with respect to some suitable metric $\overline{0}$ in $M_n(\mathbb{C})^m$ induced by the operator norm. Two of the key features of these matrix paths is that δ can be chosen independent of n, and that the paths stay in the intersection of $B_{\overline{0}}(\mathbf{X}, \varepsilon)$, and the set pairwise commuting normal matrix contractions with some special geometric structure on their joint spectra.

We apply these results to study the local connectivity properties of matrix *-representations of some universal commutative C^* -algebras. Some connections with the local connectivity properties of completely positive linear maps on matrix algebras are studied as well.

1. Introduction

In this document we study the local path connectivity of sets of m-tuples of commuting normal matrices, whose joint spectra may consist of points in some arbitrary element of the family of compact differentiable manifolds contained in \mathbb{R}^m . In particular, given $\varepsilon > 0$ and any fixed but arbitrary m-tuple $\mathbf{X} \in M_n(\mathbb{C})^m$ in the set of m-tuples of pairwise commuting normal matrix contractions (where matrix contraction means a matrix $X \in M_n(\mathbb{C})$ such that $||X|| \leq 1$), we prove the existence of paths between arbitrary m-tuples in the intersection of the previously mentioned sets of m-tuples in $M_n(\mathbb{C})^m$ and the δ -ball $B_{\eth}(\mathbf{X}, \delta)$ centered at \mathbf{X} , with respect to some suitable metric \eth in $M_n(\mathbb{C})^m$ induced by the operator norm (that will be defined further below). Two of the key features of these matrix paths is that δ can be chosen independent of n, and that the paths stay in the intersection of $B_{\eth}(\mathbf{X}, \varepsilon)$, and the set pairwise commuting normal matrix contractions with some special geometric structure on their joint spectra.

As mentioned in the previous paragraph, we start by studying local path connectivity properties of some particular varieties, these varieties consist of m-tuples of pairwise commuting normal matrix contractions with some geometric constraints on their joint spectra. Given r > 0 and a m-tuple of $n \times n$ matrices $\mathbf{X} \in M_n(\mathbb{C})^m$, let us write $B_{\eth}(\mathbf{X}, r)$ to denote the set $\{\mathbf{Y} \in (M_n)^m | \eth(\mathbf{X}, \mathbf{Y}) \leq r\}$. We call $B_{\eth}(\mathbf{X}, r)$ a r-ball centered at \mathbf{X} or a r-neighborhood of \mathbf{X} . Given $\varepsilon > 0$, we will

Date: March 1, 2022.

²⁰¹⁰ Mathematics Subject Classification. 47A58 (primary) and 15A60, 15A27 (secondary). Key words and phrases. Matrix path, spectral variation, commutative matrix algebra, joint spectrum, pseudospectrum.

prove that there is $\delta > 0$ such that for any m-tuple $\mathbf{X} := (X_1, \dots, X_m) \in M_n(\mathbb{C})^m$ in some subset Z^m of the set of commuting m-tuples of normal contractions, and any $\tilde{\mathbf{X}} := (\tilde{X}_1, \dots, \tilde{X}_m) \in B_{\overline{o}}(\mathbf{X}, \delta) \cap Z^m$, we can find a m-tuple of matrix paths $\gamma = (\gamma_1, \dots, \gamma_m) \in C([0, 1], B_{\overline{o}}(\mathbf{X}, \varepsilon)) \cap Z^m$, such that $\gamma(0) = \mathbf{X}, \gamma(1) = \tilde{\mathbf{X}}$, and where $\gamma(t) \in Z^m$ for each 0 < t < 1. As mentioned before, δ does not depend on n.

The problems which motivated the research reported in this document have a topological nature. We started this study in [16], where we found some connections between the map $\Psi := \operatorname{Ad}[W]$ described by Lemma 2.2 in §2 and the local homeomorphims that play a key role in the Kirby Torus Trick (introduced by R. Kirby in [12]) by moving sets of points in homeomorphic copies \mathcal{T}^m of \mathbb{T}^m just a little bit, this analogy together with L.3.5 provides a connection with topologically controlled linear algebra (in the sense of [10]), which can be roughly described as the study of the relations between matrix sets and differentiable manifolds, more specifically, the study of the connections between approximation of functions on differential manifolds and approximation techniques for families of matrices which satisfy some special constraints, like commutativity, hermiticity and spectral behavior among others. The search for the previously mentioned analogies and connections was motivated by a question raised by M. H. Freedman regarding to the role played by the Kirby Torus Trick in linear algebra and matrix approximation theory. In section §3 we obtain uniform versions of the previously mentioned analogies based on Lemma 3.6.

The results presented in section §3 can be used to compute, and to study the approximate and exact simultaneous block diagonalization of matrix m-tuples in the sense of [17, 18], problems of this type appear in biomathematics, image processing and applied spectral graph theory. Another important source of motivation and inspiration for this paper came from physics, more specifically from problems related to the classification of quantum phases in the sense of [22].

Building on some of the ideas developed by M. A. Rieffel in [19, 20], by O. Bratteli, G. A. Elliott, D. E. Evans and A. Kishimoto in [5], by H. Lin in [14], by D. Hadwin in [11], by K. Davidson in [8] and by P. Friis and M. Rördam in [9], we proved Lemma 3.5, Lemma 3.1 and Lemma 3.2, these results together with Lemma 2.2 provide us with the matrix approximation technology that we use to prove the main results. The main results, consisting of Theorem 3.2, Corollary 3.1, Theorem 3.3 are presented in §3. Some applications of the results in §3 to the study of the local connectivity properties of matrix representations of universal commutative C^* -algebras are presented in §4 and some future directions are outlined in §5.

2. Preliminaries and Notation

We will write $M_{m,n}$ to denote the set $M_{m,n}(\mathbb{C})$ of $m \times n$ complex matrices, if m=n we will write M_n , we write M_n^m to denote the set of m-tuples of $n \times n$ complex matrices. The symbols $\mathbf{1}_n$ and $\mathbf{0}_{m,n}$ will be used to denote the identity matrix and the zero matrix in M_n and $M_{m,n}$ respectively, if m=n we will write $\mathbf{0}_n$. Given a matrix $A \in M_n$, we write A^* to denote the conjugate transpose A^{\top} of A. A matrix $X \in M_n$ is said to be normal if $XX^* = X^*X$, a matrix $H \in M_n$ is said to be hermitian if $H^* = H$ and a matrix $U \in M_n$ such that $U^*U = UU^* = \mathbf{1}_n$ is called unitary. We will write \mathbf{i} to denote the number $\sqrt{-1}$.

Let (X,d) be a metric space. We say that $\tilde{X}_{\delta} \subset X$ is a δ -dense subset of X if for all $x \in X$ there exists $\tilde{x} \in \tilde{X}_{\delta}$ such that $d(x,\tilde{x}) \leq \delta$. Given two compact

subsets X,Y of the complex plane, we will write $d_H(X,Y)$ to denote the Hausdorff distance between X and Y that is defined by the expression $d_H(X,Y) := \max\{sup_{x\in X}\inf_{y\in Y}|x-y|, sup_{y\in Y}\inf_{x\in X}|y-x|\}$. Given $N\in\mathbb{Z}^+$, we will write \mathbb{X}_N to denote the finite set $\{x_k=-1+2k/(N-1)|0\leq k\leq N-1\}\subseteq [-1,1]$.

Given $d, m \in \mathbb{Z}^+$ with $d \leq m$, a compact manifold $Z^d \subseteq \mathbb{R}^m$, and a coordinate chart $\varphi : \mathbb{R}^d \to Z^d \subseteq \mathbb{R}^m, (x_1, \dots, x_d) \mapsto (\varphi_1(x_1, \dots, x_d), \dots, \varphi_m(x_1, \dots, x_d))$, we will write φ_j^{-1} to denote the jth component the map $\varphi^{-1} : \mathbb{R}^m \supseteq Z^d \to R^d, (z_1, \dots, z_m) \mapsto (\varphi_1^{-1}(z_1, \dots, z_m), \dots, \varphi_d^{-1}(z_1, \dots, z_m)), 1 \leq j \leq d$.

From here on $\|\cdot\|$ denotes the operator norm defined for any $A \in M_n$ by $\|A\| := \sup_{\|x\|_2=1} \|Ax\|_2$, where $\|\cdot\|_2$ denotes the Euclidean norm in \mathbb{C}^n . Let us denote by \eth the metric in M_n^m defined by $\eth: M_n^m \times M_n^m \to \mathbb{R}_0^+, (\mathbf{S}, \mathbf{T}) \mapsto \max_j \|S_j - T_j\|$. We will write \mathbb{D}^m and \mathbb{T}^m to denote the m-dimensional closed unit disk and the m-dimensional torus respectively. We will write $B(x_0, r)$ to denote the closed ball $\{x \in \mathbb{C} \mid |x - x_0| \leq r\}$.

In this document by a matrix contraction we mean a matrix X in M_n such that $\|X\| \leq 1$. A matrix $P \in M_n$ such that $P^* = P = P^2$ is called a projector or projection. Given two projectors P and Q, if $PQ = QP = \mathbf{0}_n$ we say that P and Q are orthogonal. By an orthogonal partition of unity in M_n , we mean a finite set of pairwise orthogonal projectors $\{P_j\}$ in M_n such that $\sum_j P_j = \mathbf{1}_n$. We will omit the explicit reference to M_n when it is clear from the context.

Given any two matrices $X, Y \in M_n$ we will write [X, Y] and Ad[X](Y) to denote the operations [X, Y] := XY - YX and $Ad[X](Y) := XYX^*$.

Let \mathbb{GL}_n denote the set of invertible elements in M_n . Given a matrix $A \in M_n$, we write $\sigma(X)$ to denote the set $\{\lambda \in \mathbb{C} \mid A - \lambda \mathbf{1}_n \notin \mathbb{GL}_n\}$ of eigenvalues of A, the set $\sigma(A)$ is called the spectrum of A. Given a compact set $\mathbb{X} \subset \mathbb{C}$ and a subset $S \subseteq M_n$, let us denote by $S(\mathbb{X})$ the set $S(\mathbb{X}) := \{X \in S | \sigma(S) \subseteq \mathbb{X}\}$, for instance, the expression $\mathcal{N}(n)(\mathbb{D}^2)$ is used to denote the set normal contractions in M_n .

As a consequence of [3, T.VI.3.3] and [3, C.VI.3.4] we have that.

Proposition 2.1. If A and B are $n \times n$ normal matrices, then

$$d_H(\sigma(A), \sigma(B)) \leq ||A - B||.$$

Definition 2.1. Given $\varepsilon \geq 0$ and a matrix $X \in M_n$, we write $\sigma_{\varepsilon}(X)$ to denote the ε -Pseudospectrum of X which is the set defined by the following relations.

$$\sigma_{\varepsilon}(X) := \{\tilde{\lambda} \in \mathbb{C} \mid \tilde{\lambda} \in \sigma(X + E), \text{ for some } E \in M_n \text{ with } ||E|| \leq \varepsilon\}$$
$$= \{\tilde{\lambda} \in \mathbb{C} \mid ||X\mathbf{v} - \tilde{\lambda}\mathbf{v}|| \leq \varepsilon, \text{ for some } \mathbf{v} \in \mathbb{C}^n \text{ with } ||\mathbf{v}|| = 1\}$$

Definition 2.2 (Semialgebraic Matrix Varieties). Given $J \in \mathbb{Z}^+$, a system of J polynomials $p_1, \ldots, p_J \in \Pi_{\langle N \rangle} = \mathbb{C}\langle x_1, \ldots, x_N \rangle$ in N noncommutative variables and J real numbers $\varepsilon_j \geq 0$, $1 \leq j \leq J$, a matrix representation of the noncommutative semialgebraic set $\mathcal{Z}_n(p_1, \ldots, p_J)$ described by

$$\mathcal{Z}_n(p_1,\ldots,p_J) := \{X_1,\ldots,X_N \in M_n | \|p_j(X_1,\ldots,X_N)\| \le \varepsilon_j, 1 \le j \le J\},$$

will be called a n-semialgebraic matrix variety (n-SMV), if each $\varepsilon_j = 0$, we will refer to the set as a matrix variety and we may replace the normed polynomial relations by polynomial relations.

Example 2.1. As a first example, we will have that the matrix set

$$\mathbf{Z}_{n} := \left\{ (X_{1}, \dots, X_{N}) \in M_{n}^{N} \middle| \begin{array}{l} X_{j} X_{k} - X_{k} X_{j} = \mathbf{0}_{n}, \\ X_{j}^{*} X_{j} = X_{j} X_{j}^{*} = \mathbf{1}_{n}, \end{array} \right. 1 \leq j, k \leq N \right\}$$

is a matrix variety. If for some $\delta > 0$, we set now

$$\mathbf{Z}_{n,\delta} := \left\{ (X_1, \dots, X_N) \in M_n^N \middle| \begin{array}{l} \|X_j X_k - X_k X_j\| \le \delta, \\ \|X_j^* X_j - X_j X_j^*\| = 0, \ 1 \le j, k \le N \\ \|X_j\| \le 1 \end{array} \right\}$$

the set $\mathbf{Z}_{n,\delta}$ is a matrix semialgebraic variety.

Example 2.2. Other example of a matrix semialgebraic variety, that has been useful to understand the geometric nature of the problems solved in this document, is described by the matrix set $\mathbf{Iso}_{\delta}(x,y)$, defined for some given $\delta \geq 0$ and any two normal contractions x and y in M_n , by the expression

$$\mathbf{Iso}_{\delta}(x,y) := \left\{ (z,w) \in \mathcal{N}(n)(\mathbb{D}^2) \times \mathbb{U}(n) \middle| \begin{array}{c} \|xw - wz\| = 0 \\ \|[z,y]\| = 0 \\ \|z - y\| \le \delta \end{array} \right\}.$$

Given a m-tuple $\mathbf{X} \in M_n^m$ we will write either $\mathbf{X}[j]$ or X_j to denote its jth component in M_n , with $1 \leq j \leq m$. In a similar way given a path $\gamma \in C([0,1], M_n^m)$ we will write γ_j to denote its jth component in $C([0,1], M_n)$, and we will write $\gamma_j(t)$ to denote the jth component of $\gamma(t)$ for some $0 \leq t \leq 1$, with $1 \leq j \leq m$.

Given a map $\Phi: M_n \to M_n$, we will write $\hat{\Phi}$ to denote its **extension** to M_n^m defined by $\hat{\Phi}: M_n^m \to M_n^m, (X_1, \dots, X_m) \mapsto (\Phi(X_1), \dots, \Phi(X_m))$.

Definition 2.3 (Morphisms of Matrix Varieties). Given a matrix variety $Z^m \subseteq M_n^m$, a morphism of Z^m is any map $\Psi: Z^m \to Z^m$. Given a map $\Psi: M_n \to M_n$, if the extended map $\hat{\Psi}: M_n^m \to M_n^m$ is a morphism of Z^m , we say that $\hat{\Psi}$ is a morphism induced by Ψ . A morphism $\Phi: Z^m \to Z^m$ is said to be an inner morphism of Z^m if it can be represented in the form $\Phi(X_1, \ldots, X_m) = (\operatorname{Ad}[W_1](X_1), \ldots, \operatorname{Ad}[W_m](X_m))$ for some fixed but arbitrary $W_1, \ldots, W_m \in M_n$ and any $(X_1, \ldots, X_m) \in Z^m$.

Remark 2.1. In particular, given a matrix variety $Z^m \subseteq M_n^m$, and given any inner morphism $\Phi: Z^m \to Z^m$ of the form $\Phi = \widehat{\operatorname{Ad}[W]}$ for some $W \in \mathbb{U}(n)$, it can be seen that if the components X_j of $(X_1, \ldots, X_m) \in Z^m$ are normal matrices, unitaries, commuting matrices, hermitian matrices, projectors, or contractions, then the same will be true for the components $\operatorname{Ad}[W](X_j)$ of $\Phi(X_1, \ldots, X_m) \in Z^m$. Moreover, $\sigma(X_j) = \sigma(\operatorname{Ad}[W](X_j))$ for each $1 \leq j \leq m$.

Definition 2.4 (** operation). Given two matrix paths $\alpha, \beta \in C([0,1], M_n^m)$ we write $\alpha \circledast \beta$ to denote the concatenation of α and β , which is the matrix path defined in terms of α and β by the expression,

$$\alpha\circledast\beta(s):=\left\{\begin{array}{ll}\alpha(2s),\ 0\leq s\leq\frac{1}{2},\\\beta(2s-1),\ \frac{1}{2}\leq s\leq 1.\end{array}\right.$$

From here on, for any two m-tuples of matrices $\mathbf{X}, \mathbf{Y} \in M_n^m$, we will write $\mathbf{X} \leadsto \mathbf{Y}$ to indicate that there is a matrix path $\gamma \in C([0,1], M_n^m)$ such that $\gamma(0) = \mathbf{X}$ and $\gamma(1) = \mathbf{Y}$.

Given $\varepsilon, \delta > 0$, and any two m-tuples of pairwise commuting normal matrix contractions $\mathbf{X} := (X_1, \dots, X_m)$ and $\tilde{\mathbf{X}} := (\tilde{X}_1, \dots, \tilde{X}_m)$ in some (semialgebraic) matrix variety $Z^m \subseteq M_n^m$ such that $\eth(\mathbf{X}, \tilde{\mathbf{X}}) \le \delta$, we will write $\mathbf{X} \leadsto_{\varepsilon} \tilde{\mathbf{X}}$ relative $\operatorname{to} B_{\eth}(\mathbf{X}, \varepsilon) \cap Z^m$, to indicate that there is a piecewise differentiable path $\gamma \in C([0, 1], B_{\eth}(\mathbf{X}, \varepsilon) \cap Z^m)$ such that $\gamma(0) = \mathbf{X}$ and $\gamma(1) = \tilde{\mathbf{X}}$.

Given a matrix $A \in M_n$, we will write $\mathcal{D}(A)$ to denote the diagonal matrix defined by the following operation.

$$\mathscr{D}(A) := \operatorname{diag}[a_{11}, a_{22}, \dots, a_{nn}]$$

(2.2)
$$= \begin{pmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \cdots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & a_{nn} \end{pmatrix}$$

It can be seen that $\mathscr{D}(\mathscr{D}(A)) = \mathscr{D}(A)$ for any $A \in M_n$, the map \mathscr{D} will be called the full pinching.

Remark 2.2. By pinching inequalities (in the sense of [4]) we will have that $\|\mathscr{D}(A) - \mathscr{D}(B)\| = \|\mathscr{D}(A - B)\| \le \|A - B\|$ for any two matrices $A, B \in M_n$.

It is often convenient to have N-tuples (or 2N-tuples) of matrices with real spectra. For this purpose we use the following construction. If $X = (X_1, \ldots, X_N)$ is a N-tuple of n by n matrices then we can always decompose X_j in the form $X_j = X_{1j} + iX_{2j}$ where the X_{kj} all have real spectra. We write $\pi(X) := (X_{11}, \ldots, X_{1N}, X_{21}, \ldots, X_{2N})$ and call $\pi(X)$ a **partition** of X. If the X_{kj} all commute we say that $\pi(X)$ is a commuting partition, and if the X_{kj} are simultaneously triangularizable $\pi(X)$ is a triangularizable partition. If the X_{kj} are all semisimple (diagonalizable) then $\pi(X)$ is called a semisimple partition. From here on, for any m-tuple of pairwise commuting normal matrices $X = (X_1, \ldots, X_m)$, we will assume that the components of $\pi(X) = (X_{11}, \ldots, X_{1m}, X_{21}, \ldots, X_{2m})$ are determined by the relations

(2.3)
$$\begin{cases} X_{1j} = (X_j + X_j^*)/2, \\ X_{2j} = (X_j - X_j^*)/(2\mathbf{i}), \end{cases}$$

for each $1 \leq j \leq m$. In addition, given any operator X in a unital separable C^* -algebra A, we will write Re(X) and Im(X) to denote the self-adjoint operators in A defined by the equations $Re(X) = (X + X^*)/2$ and $Im(X) = (X - X^*)/(2\mathbf{i})$.

Given a 2m-tuple $\mathbf{X} = (X_{11}, \dots, X_{1m}, X_{21}, \dots, X_{2m})$ in M_n^{2m} , the m-tuple obtained by the operation $v(\mathbf{X}) := (X_{11} + \mathbf{i}X_{21}, \dots, X_{1m} + \mathbf{i}X_{2m}) \in M_n^m$ will be called **juncture** of \mathbf{X} .

We say that N normal matrices $X_1, \ldots, X_N \in M_n$ are simultaneously diagonal-izable if there is a unitary matrix $Q \in M_n$ such that Q^*X_jQ is diagonal for each $j=1,\ldots,N$. In this case, for $1 \leq k \leq n$, let $\Lambda^{(k)}(X_j) := (Q^*X_jQ)_{kk}$ the (k,k) element of Q^*X_jQ , and set $\Lambda^{(k)}(X_1,\ldots,X_N) := (\Lambda^{(k)}(X_1),\ldots,\Lambda^{(k)}(X_N))$ in \mathbb{C}^N . The set

$$\Lambda(X_1, \dots, X_N) := \{\Lambda^{(k)}(X_1, \dots, X_N)\}_{1 \le k \le N}$$

is called the joint spectrum of X_1, \ldots, X_N . Given a set $S \subseteq M_n$ of m-tuples of pairwise commuting normal matrices, we will write $\Lambda(S)$ to denote the set $\{\Lambda(X) \mid X \in S\}$, the set $\Lambda(S)$ will be called the joint spectra of S. We will write $\Lambda(X_j)$ to denote the diagonal matrix representation of the j-component of $\Lambda(X_1, \ldots, X_N)$, in other words we will have that

$$\Lambda(X_j) = \operatorname{diag}\left[\Lambda^{(1)}(X_j), \dots, \Lambda^{(n)}(X_j)\right].$$

Given a m-tuple $\mathbf{X} = (X_1, \dots, X_m) \in M_n^m$ of commuting normal matrices, any orthogonal projection $P \in M_n$ such that $X_j P = P X_j = \Lambda^{(r)}(X_j) P$ for each $1 \leq j \leq m$ and some $1 \leq r \leq n$, will be called a joint spectral projector of \mathbf{X} .

Definition 2.5 (C^* -algebras generated by m-tuples). Given a m-tuple of normal matrix contractions $\mathbf{X} = (X_1, \dots, X_m) \in M_n^m$ we write $C^*(\mathbf{X})$ to denote the C^* -algebra $C^*(X_1, \dots, X_m) \subseteq M_n$.

The following result ([13, Lemma 1]) was proved in [13].

Lemma 2.1. Let X be a compact sucset of the plane and let $f \in C(X)$. For any $\varepsilon > 0$, there is $\delta > 0$, for any C^* -algebra A and normal elements $x, y \in A$ with $\sigma(x), \sigma(y) \in X$, if $||x - y|| < \delta$, then $||f(x) - f(y)|| < \varepsilon$.

The following result ([16, Lemma 4.1]) was proved in [16].

Lemma 2.2. Given $\varepsilon > 0$ there is $\delta = \frac{1}{K_m} \varepsilon > 0$ such that, for any two N-tuples of pairwise commuting normal matrices $\mathbf{X} = (X_1, \dots, X_N)$ and $\mathbf{Y} := (Y_1, \dots, Y_N)$ such that $\eth(\mathbf{X}, \mathbf{Y}) \leq \delta$, there is a unitary matrix W such that the *-homomorphism $\Psi = \operatorname{Ad}[W]$ satisfies the condition $\Psi : C^*(\mathbf{X}) \to C^*(\mathbf{Y})'$ together with the constraint $\max\{\eth(\hat{\Psi}(\mathbf{X}), \mathbf{Y}), \eth(\hat{\Psi}(\mathbf{X}), \mathbf{X})\} \leq \varepsilon$.

Remark 2.3. The constant K_m in the statement of L.2.2 depends only on m.

- 3. Spectral Approximation and Local Path Connectivity of Sets of Commuting Normal Matrices
- 3.0.1. Matrix Representations of $C(\mathbb{X}_N)$. Let us consider the universal commutative C^* -algebra $C(\mathbb{X}_N)$ for some fixed but arbitrary $N \in \mathbb{Z}^+$. We will have that $C(\mathbb{X}_N)$ has a universal picture that can be expressed in terms of generators and relations as follows.

(3.1)
$$C(\mathbb{X}_N) := C_1^* \left\langle p_1, \dots, p_N \middle| \begin{array}{l} p_j p_k = p_k p_j \\ p_j^2 - p_j = p_j - p_j^* = 0 \\ p_1^2 + \dots + p_N^2 - 1 = 0 \end{array}, 1 \le j, k \le N \right\rangle,$$

From here on, given a universal C^* -algebra A and a C^* -subalgebra $B \subseteq M_n$, we will write $C^*\text{Rep}(A, B)$ to denote the set of *-representations of A in B.

3.0.2. Eventually uniform matrix representations. Let $N \in \mathbb{Z}^+$ be fix but arbitrary. Given $n \in \mathbb{Z}^+$, let us consider the case n < N first. Without loss of generality we can consider the *-homomorphism induced by the embedding $i_{n,N}: M_n \hookrightarrow M_N$ determined by the map

(3.2)
$$i_{r,s} : M_r \to M_s \\
 : X \mapsto \begin{pmatrix} X & \mathbf{0}_{n,N-n} \\ \mathbf{0}_{N-n,n} & \mathbf{0}_{N-n} \end{pmatrix}$$

for any two positive integers $r \leq s$. We can use (3.2) to compute for any orthogonal partition of unity $\mathbf{P}_n = \{P_1, \dots, P_n\} \subseteq M_n$, a set of orthogonal projections $\mathbf{p}_n = \{p_1, \dots, p_n\} = \{i_{n,N}(p_1), \dots, i_{n,N}(p_n)\} = \{P_1 \oplus \mathbf{0}_{N-n}, \dots, P_n \oplus \mathbf{0}_{N-n}\} \subseteq M_N$. We will have that the map $\kappa_{N,n} : M_N \to M_n : X \mapsto \operatorname{Ad}[K](X)$, with $K = (\mathbf{1}_n \ \mathbf{0}_{n,N-n})$, induces a *-homomorphism $\rho_{N,n}^{>} : C^*(\mathbf{p}_n) \to C^*(\mathbf{P}_n)$ such that $\rho_{N,n}^{>}(p_j) = P_j$ for each $1 \leq j \leq n$, by universality of (3.1) the map $\rho_{N,n}^{>}$ induces a *-homomorphism $\rho_n : C(\mathbb{X}_N) \to C^*(\mathbf{P}_n) \subseteq M_n$, for each n < N. If $n \geq N$, by universality of (3.1),

we will have that for any orthogonal partition of unity $\mathbf{P}_n = \{P_1, \dots, P_N\} \subseteq M_n$, there is a *-homomorphism $\rho_{N,n}^{\leq} : C(\mathbb{X}_N) \to C^*(\mathbf{P}_n) \subseteq M_n$ such that $\rho_{N,n}^{\leq}(p_j) = P_j$ for each $1 \leq j \leq N$. Let us set


(3.3)
$$\rho_{N,n} := \left\{ \begin{array}{l} \rho_{N,n}^{>}, & n < N \\ \rho_{N,n}^{\leq}, & n \ge N \end{array} \right.$$

It can be seen that the map $\rho_{N,n}$ is a *-homomorphism for each $n, N \in \mathbb{Z}^+$. We will call each map $\rho_{N,n}$ of the sequence $\{\rho_{N,n}\}_{n\in\mathbb{Z}^+}$ an **eventually uniform** matrix representation (**EUMR**) of $C(\mathbb{X}_N)$.

In order to provide some perspective on how our connectivity and approximation results do not depend on the matrix size n, we will describe what we call the **approximate eventual uniformity** of matrix representations of C(X), where X is a compact Hausdorff topological space.

Definition 3.1. Given a compact Hausdorff space X, we say that the commutative C^* -algebra C(X) is approximately eventually uniform (**AEU**) if for any finite set of normal contractions $\{z_1, \ldots, z_m\} \subset C(X)$, there is $\delta > 0$ and an integer $N_{\delta,m} > 0$ such that for any integer n > 1 and any $\rho_n \in C^* \operatorname{Rep}(C(X), M_n)$, there is a EUMR $\rho_{N_{\delta,m},n} \in C^* \operatorname{Rep}(C(\mathbb{X}_{N_{\delta,m}}), \rho_n(C(X))')$ together with a *-homomorphism $\rho_{\delta,n} : C(X) \to C^*(\rho_{N_{\delta,n}}(\operatorname{id}_{\mathbb{X}_{N_{\delta}}}))$ such that $\|\rho_n(z_j) - \rho_{\delta,n}(z_j)\| \le \delta$ for each $1 \le j \le m$.

By considering natural embeddings of compact Hausdorff spaces into $[-1,1]^m$ for some $m \in \mathbb{Z}^+$, we have that one of the main problems that we need to solve in this document, is an operator approximation problem that can be stated via the following diagram.



To solve (3.4), given a *-representation $\pi: C(\mathbb{I}^m) \to C^*(H_1, \ldots, H_m)$ such that $\pi(h_j) = H_j$ for each $1 \leq j \leq m$, one needs to find $\pi_n \in C^*\operatorname{Rep}(C(\mathbb{X}_{N_\delta}), \pi(C(\mathbb{I}^m))')$ and a *-representation $\pi_\delta: C(\mathbb{I}^m) \to C^*(\hat{\mathbb{X}}_{N_\delta})$ such that:

$$(3.5) \qquad \left\{ \begin{array}{l} (\imath_2 \circ \pi)(h_j)(\imath_1 \circ \pi_\delta)(h_k) = (\imath_1 \circ \pi_\delta)(h_k)(\imath_2 \circ \pi)(h_j) \\ \|(\imath_2 \circ \pi)(h_j) - (\imath_1 \circ \pi_\delta)(h_j)\| \le \delta \end{array} \right., 1 \le j, k \le m.$$

Here $\hat{\mathbb{X}}_{N_{\delta}}, \hat{\mathbb{X}}_{n} \in C^{*}(H_{1}, \ldots, H_{m})'$ are defined by the equations $\hat{\mathbb{X}}_{N_{\delta}} = \pi_{n}(\mathrm{id}_{\mathbb{X}_{N_{\delta}}})$ and $\hat{\mathbb{X}}_{n} = \rho_{n}(\mathrm{id}_{\mathbb{X}_{n}})$ for some *-homomorphism $\rho_{n} \in C^{*}\mathrm{Rep}(C(\mathbb{X}_{n}), C^{*}(H_{1}, \ldots, H_{m})')$.

Let us define two semialgebraic matrix varieties that will play a key role in this study.

Definition 3.2. The semialgebraic matrix variety $\mathbb{I}^m(n)$ defined by

$$\mathbb{I}^{m}(n) := \left\{ (X_{1}, \dots, X_{m}) \in M_{n}^{m} \middle| \begin{array}{l} [X_{j}, X_{k}] = \mathbf{0}_{n} \\ X_{j} - X_{j}^{*} = \mathbf{0}_{n} \\ \|X_{j}\| \leq 1 \end{array} \right., 1 \leq j, k \leq m \right\}$$

will be called the matrix m-cube.

Definition 3.3. The semialgebraic matrix variety $\mathbb{D}^m(n)$ defined by

$$\mathbb{D}^{m}(n) := \left\{ (X_{1}, \dots, X_{m}) \in M_{n}^{m} \middle| \begin{array}{l} [X_{j}, X_{k}] = \mathbf{0}_{n} \\ X_{j} X_{j}^{*} - X_{j}^{*} X_{j} = \mathbf{0}_{n} \\ \|X_{j}\| \le 1 \end{array}, 1 \le j, k \le m \right\}$$

will be called the matrix m-disk.

Let us now extend the concept of semialgebraic matrix variety to what we will call an *induced compact differential matrix manifold* or *compact differential matrix manifold*.

Definition 3.4. Given a compact differentiable d-manifold $Z^d \subseteq \mathbb{R}^m$ with $m \geq d$, the matrix variety $\mathbb{V}^m[Z^d](n)$ defined by

$$\mathbb{V}^{m}[Z^{d}](n) := \left\{ (X_{1}, \dots, X_{m}) \in M_{n}^{m} \middle| \begin{array}{l} [X_{j}, X_{k}] = \mathbf{0}_{n} \\ X_{j} - X_{j}^{*} = \mathbf{0}_{n} \\ \|X_{j}\| \leq 1 \\ \Lambda(X_{1}, \dots, X_{m}) \subseteq Z^{d} \end{array}, 1 \leq j, k \leq m \right\}$$

will be called the differentiable matrix manifold induced by Z^d .

3.0.3. Borel functional calculus and matrix approximation. By applying Borel functional calculus on commutative von Neumann algebras we have that given any real number $0 < \delta \le 1$, any hermitian matrix $X \in M_n$ such that $\sigma(X) \subset [-1,1]$, and any partition $-1+\delta/2=x_1 < x_2 < \cdots < x_m < x_{m+1}=1+\delta/2$ of $[-1-\delta/2,1+\delta/2]$, with $x_{k+1}-x_k=\delta$ for each $1 \le k \le m+1$, the matrix X has a representation $X = \sum_{k=1}^{m+1} \chi_{[x_k,x_{k+1})}(X)X$, where $\chi_{[a,b)}(X)$ denotes the characteristic function of the interval [a,b), for some real numbers a,b such that a < b. We also have that for each $1 \le k \le m$, the matrix $P_k = \chi_{[x_k,x_{k+1})}(X)$ is an orthogonal projection that commutes with X such that $\sigma(P_kX) \subseteq \sigma(X) \cap [x_k,x_{k+1})$. Moreover, the set $\{P_1,\ldots,P_m\}$ is an orthogonal partition of unity.

Definition 3.5 (Pseudospectral matrix approximant). Given $\delta > 0$ and any matrix $X \in M_n$, we say that the matrix $\tilde{X} \in M_n$ is a δ -Pseudospectral matrix approximant $(\delta \text{-PMA})$ of X if $||X - \tilde{X}|| \le \delta$ and $\sigma(\tilde{X})$ is δ -dense in $\sigma(X)$. If in addition each $\tilde{X}X = X\tilde{X}$, we say that \tilde{X} is a commuting δ -PMA (δ -CPMA).

Lemma 3.1. Given $\delta > 0$ and any hermitian matrix $X \in M_n$ such that $||X|| \le 1$, there is a hermitian δ -CPMA \tilde{X}_{δ} of X. Moreover, there are N_{δ} numbers $x_1, \ldots, x_{N_{\delta}}$ in [-1, 1] and an orthogonal partition of unity $\{P_1, \ldots, P_{N_{\delta}}\}$ such that $\tilde{X}_{\delta} = \sum_{j} x_{j} P_{j}$.

Proof. Let us suppose that $n \geq |\sigma(X)| \geq 2$, as the proof is clear for scalar multiples of $\mathbf{1}_n$. Since $\sigma(X) \subseteq [-1,1]$, we can assume for simplicity that $M_\delta := 1 + (\delta)^{-1} \in \mathbb{Z}^+$. Then the finite set $\hat{R}_\delta(X) := \{x_k := -1 + 2(k-1)\delta \mid 1 \leq k \leq M_\delta\}$ is δ -dense in [-1,1] with $|\hat{R}_\delta(X)| = M_\delta$. Let us set $S_\delta(X) := \{\check{x}_k := -1 + (2k-3)\delta \mid 1 \leq k \leq M_\delta + 1\}$ and $P_k := \chi_{(\check{x}_k,\check{x}_{k+1}]}(X)$ for each $1 \leq k \leq M_\delta$. Then there is a set $R_\delta(X) := \{x_j\} \subseteq \hat{R}_\delta(X)$ that is δ -dense in $\sigma(X)$ with $x_j \in (\check{x}_{k(j)},\check{x}_{k(j)+1}]$ for each $x_j \in R_\delta(X)$ and some $\check{x}_{k(j)},\check{x}_{k(j)+1} \in S_\delta(X)$, and the set $P_\delta(X) := \{P_1,\ldots,P_{N_\delta}\} \subseteq M_n \setminus \{\mathbf{0}_n\}$ of $N_\delta = |R_\delta(X)|$ projections is an orthogonal partition of unity such that $[P_j,X] = \mathbf{0}_n$ for each j. We will have that for each $1 \leq k \leq N_\delta \leq M_\delta$ there is $x_{j(k)} \in R_\delta(X)$

such that $\|XP_k - x_{j(k)}P_k\| \leq \delta$. By the previous facts and Proposition 2.1 we will have that if we set $\tilde{X}_{\delta} := \sum_k x_{j(k)}P_k$ with $x_{j(k)} \in R_{\delta}(X)$ for each k, then $[X, \tilde{X}_{\delta}] = \mathbf{0}_n$ and $d_H(\sigma(X), \sigma(\tilde{X}_{\delta})) \leq \|X - \tilde{X}_{\delta}\| \leq \max_k \|XP_k - x_{j(k)}P_k\| \leq \delta$. If necessary we can renumber the elements of $R_{\delta}(X)$ according to the elements of $P_{\delta}(X)$. This completes the proof.




FIGURE 1. Graphical illustration of a representation grid (consisting of \circ points) for [0,1] together with its support grid (consisting of \bullet points).




FIGURE 2. Graphical illustration of a representation grid (consisting of \circ points) for $[0,1]^2$ together with its support grid (consisting of \bullet points).

Definition 3.6 (CPMA for m-tuples). Given $\delta > 0$ and any m-tuple of pairwise commuting matrices $\mathbf{X} = (X_1, \dots, X_m) \in M_n^m$, we say that the matrix $\tilde{\mathbf{X}} = (\tilde{X}_1, \dots, \tilde{X}_m) \in M_n^m$ is a commuting δ -Pseudospectral matrix approximant (δ -**CPMA**) of \mathbf{X} if $||X_j - \tilde{X}_j|| \leq \delta$, $\sigma(\tilde{X}_j)$ is δ -dense in $\sigma(X_j)$ and $\tilde{X}_k X_j = X_j \tilde{X}_k$ for each $1 \leq j, k \leq m$. If in addition each component \tilde{X}_j of $\tilde{\mathbf{X}}$ is normal, hermitian or unitary, we say that $\tilde{\mathbf{X}}$ is a hermitian, normal or unitary δ -CPMA respectively.

Lemma 3.2. Given $\delta > 0$, an integer $m \ge 1$ and any m-tuple of pairwise commuting hermitian matrix contractions $\mathbf{X} = (X_1, \dots, X_m) \in M_n^m$, there is a hermitian δ -CPMA $\tilde{\mathbf{X}}_{\delta} = (\tilde{X}_{\delta,1}, \dots, \tilde{X}_{\delta,m}) \in M_n^m$ of \mathbf{X} . Moreover, for each $1 \le k \le m$, there are N_{δ} (not necessarily distinct) numbers $x_{k,1}, \dots, x_{k,N_{\delta}}$ counted with multiplicity in [-1,1] and an orthogonal partition of unity $\{P_1, \dots, P_{N_{\delta}}\}$ such that $\tilde{X}_{\delta,k} = \sum_j x_{k,j} P_j$.

Proof. Since the joint spectrum $\sigma(\mathbf{X})$ of \mathbf{X} is a subset of $[-1,1]^m$ we can apply a similar procedure to the one implemented in the proof of Lemma 3.1 to find for each X_j a representation grid $R_{\delta}(X_j)$ together with a support grid $S_{\delta}(X_j)$, an associated δ -projective decomposition $P_{\delta}(X_j)$ and an integer $N_{\delta,j} := |P_{\delta}(X_j)| = |R_{\delta}(X_j)|$. We will have that the set $P_{\delta}(\mathbf{X}) := \{P_1, \dots, P_{N_{\delta}}\} = \{P_{1,j_1}P_{2,j_2}\cdots P_{m,j_m} \mid P_{k,j_k} \in P_{\delta}(X_k), \ 1 \le j_k \le N_{\delta,k}, \ 1 \le k \le m\}$ is an orthogonal partition of unity such that PQ = QP for each $P \in P_{\delta}(\mathbf{X})$ and each $Q \in P_{\delta}(X_k)$, for any $1 \le k \le m$. By setting $N_{\delta} := |P_{\delta}(\mathbf{X})|$, we will have that $N_{\delta} \ge N_{\delta,k}$ for each k. We will also have that $P_{\delta}(X_k) \subseteq \text{span } P_{\delta}(\mathbf{X})$ for each k. For each X_k and each $P_j \in P_{\delta}(\mathbf{X})$ there is $x_{k,j} \in R_{\delta}(X_j) \cup \{0\}$ such that $||X_kP_j - x_{k,j}P_j|| \le \delta$. Moreover, $|[X_j, P_k]| = \mathbf{0}_n$ for each $1 \le j \le m$ and each $1 \le k \le N_{\delta}$. By a similar argument to the one implemented in the proof of Lemma 3.1 it can be seen that the matrix $\tilde{X}_{\delta,k} := \sum_j x_{k,j} P_j$ is a commuting hermitian δ -PMA of X_j for each j. This completes the proof.

Remark 3.1. Given a matrix m-tuple $\mathbf{X} \in M_n^m$, we will refer to the finite sets $R_{\delta}(\mathbf{X}) = \bigcup_j R_{\delta}(X_j) \cup \{0\}$ and $S_{\delta}(\mathbf{X}) = \bigcup_j S_{\delta}(X_j)$ described in the proof of Lemmas 3.1 and 3.2 as representation and support grids respectively. The finite set $P_{\delta}(\mathbf{X})$ described in the previous lemma will be called a δ -projective decomposition of \mathbf{X} . Two graphical examples of representation and support grids are presented in Fogires 1 and 2.

We can now obtain the following approximation result for $C([-1,1]^m)$.

Lemma 3.3. Given $m \in \mathbb{Z}^+$, we have that for any $\delta > 0$ the local approximation problem in $C([-1,1]^m)$ described by the diagram (3.4) is solvable.

Proof. Let $m \in \mathbb{Z}^+$. For any $n \in \mathbb{Z}^+$ and any *-homomorphism ρ_n in the set $C^*\operatorname{Rep}(C([-1,1]^m),M_n)$, we can apply Lemma 3.2 to the m-tuple $\mathbf{H}=(\pi_n(h_j))\in \mathbb{I}^m(n)$ to obtain a δ -**CPMA** $\tilde{\mathbf{H}}=(\tilde{H}_j)\in \mathbb{I}^m(n)$ such that $C^*(\tilde{\mathbf{H}})\subseteq C^*(P_\delta(\mathbf{H}))$. Let us set $N_\delta=|P_\delta(\mathbf{H})|$, by universality of $C(\mathbb{X}_{N_\delta})$ we will have that there is a *-representation $\rho_n:C(\mathbb{X}_{N_\delta})\to C^*(P_\delta(\mathbf{H}))\supseteq C^*(\tilde{\mathbf{H}})$, and by universality of $C([-1,1]^m)$ we will have that there is a *-representation $\pi_\delta:C([-1,1]^m)\to C^*(\tilde{\mathbf{H}})\subseteq C^*(P_\delta(\mathbf{H}))$. It can be seen that π_δ satisfies the conditions (3.5). This completes the proof.

Theorem 3.1. Given any compact Hasudorff space X, we have that the universal C^* -algebra C(X) is AEU.

Proof. Given any m contractions $z_1, \ldots, z_m \in C(X)$, let us consider the 2m selfajoint contractions $Re(z_1), Im(z_1), \ldots, Re(z_m), Im(z_m)$. By universality of the C^* -algebra $C([-1,1]^{2m})$ we will have that any *-representation of the form

$$\hat{\pi}_n: C^*(Re(z_1), Im(z_1), \dots, Re(z_m), Im(z_m)) \to M_n,$$

induces a *-representation of the form

(3.6)
$$\pi_n: C([-1,1]^{2m}) \to C^*(Re(z_1), Im(z_1), \dots, Re(z_m), Im(z_m)) \to M_n,$$
 determined by the equations $\pi_n(h_{2j-1}) = \hat{\pi}_n(Re(z_j))$ and $\pi_n(h_{2j}) = \hat{\pi}_n(Im(z_j)),$ $1 \le j \le m$. We can apply Lemma 3.3 to π_n replacing δ by $\delta/2$ if necessary. This completes the proof.

Lemma 3.4. Given $d, m \in \mathbb{Z}^+$ with $d \leq m$, a compact differentiable d-manifold $Z^d \subseteq [-1,1]^m \subseteq \mathbb{R}^m$, a number $\delta > 0$, and any m-tuple of pairwise commuting hermitian matrix contractions $\mathbf{Z} = (Z_1, \ldots, Z_m) \in \mathbb{V}^m[Z^d](n)$, there is a δ -CPMA $\tilde{\mathbf{Z}}_{\delta} = (\tilde{Z}_{\delta,1}, \ldots, \tilde{Z}_{\delta,m}) \in \mathbb{V}^m[Z^d](n)$ of \mathbf{Z} . Moreover, there are $\mathbf{H} = (H_1, \ldots, H_d)$ and $\tilde{\mathbf{H}}_{\delta} = (H_{\delta,1}, \ldots, \tilde{H}_{\delta,d})$ in $\mathbb{I}^m(n)$ such that for each $1 \leq k \leq m$ and each $1 \leq k' \leq d$, there are N_{δ} (not necessarily distinct) numbers $x_{k',1}, \ldots, x_{k',N_{\delta}}$ counted with multiplicity in \mathbb{R} , N_{δ} (not necessarily distinct) coordinate charts $\varphi_{k,1}, \ldots, \varphi_{k,N_{\delta}}$ of Z^d counted with multiplicity, and an orthogonal partition of unity $\{P_1, \ldots, P_{N_{\delta}}\}$ such that

(3.7)
$$\begin{cases} Z_k = \sum_j \varphi_{k,j}(P_j H_1, \dots, P_j H_d) P_j \\ \tilde{Z}_{\delta,k} = \sum_j \varphi_{k,j}(P_j \tilde{H}_{\delta,1}, \dots, P_j \tilde{H}_{\delta,d}) P_j \\ \varphi_{k,j}(P_j \tilde{H}_{\delta,1}, \dots, P_j \tilde{H}_{\delta,d}) = \varphi_{k,j}(x_{1,j}, \dots, x_{d,j}) P_j, \end{cases}$$

for each $1 \le j \le N_{\delta}$ and each $1 \le k \le m$.

Proof. Let Z^d be a compact differentiable d-manifold given as in the statement of Lemma 3.4. Let us consider a m-tuple of pairwise commuting hermitian matrix contractions $\mathbf{Z} = (Z_1, \ldots, Z_m) \in \mathbb{V}^m[Z^d](n)$. Since Z^d is a compact differentiable d-manifold, for any $\delta > 0$, there is a (finite) family of differentiable charts $\mathcal{C}(Z^d) := \{\varphi_{k,l}\}$ of Z^d together with a number $\nu_{\delta} > 0$ such that

$$(3.8) \qquad (\varphi_{1,l}(x), \dots, \varphi_{m,l}(x)) \in B^{\circ}((\varphi_{1,l}(x), \dots, \varphi_{m,l}(x)); \delta) \cap Z^d,$$

for each $x \in B^{\circ}(x, \nu_{\delta}) \cap dom(\varphi_{k,l}) \subset \mathbb{R}^d$ and for each k, l, where $B^{\circ}(x, r)$ denotes the open ball of radius r in the Euclidean metric in \mathbb{R}^m . For each $1 \leq k \leq m$, let us consider the spectral resolutions

(3.9)
$$Z_k = \sum_{i=1}^n z_{k,i} Q_i,$$

where each $Q_i \in M_n$ is a joint rank one spectral projector of **Z**, by Definition 3.4 and by (3.8) we will have that there is a hermitian matrix contraction

(3.10)
$$H_k = \sum_{i=1}^n \varphi_{k,i}^{-1}(z_{1,i}, \dots, z_{m,i})Q_i$$

for each $1 \leq k \leq d$, where each $\varphi_{i,i} \in \mathcal{C}(\mathbb{Z}^d)$ and where

(3.11)
$$\varphi_{j,i}(\varphi_{1,i}^{-1}(z_{1,i},\ldots,z_{m,i}),\ldots,\varphi_{d,i}^{-1}(z_{1,i},\ldots,z_{m,i}))=z_{j,i}$$

for each $1 \leq i \leq n$ and each $1 \leq j \leq m$. By the previously described facts and by (3.10) we will have that $\mathbf{H} := (H_1, \dots, H_d)$ can be chosen from $\mathbb{I}^d(n)$, and that

we can apply Lemma 3.2 to find a ν_{δ} -PMA $\tilde{\mathbf{H}}_{\delta} := (\tilde{H}_{\delta,1}, \dots, \tilde{H}_{\delta,d})$ of \mathbf{H} in $\mathbb{I}^m(n)$, where each $\tilde{H}_{\delta,k}$ can be decomposed in the form $\tilde{H}_{\delta,k} = \sum_j x_{k,j} P_j$ described in Lemma 3.2 for some ν_{δ} -projective decomposition $\{P_j\} = P_{\delta}(\mathbf{H})$. Because of the smooth structure of Z^d and by (3.8), (3.9), (3.10) and (3.11) we will have that for each $1 \leq k \leq m$, there are N_{δ} (not necessarily distinct) differentiable charts $\varphi_{k,1}, \dots, \varphi_{k,N_{\delta}}$ counted with multiplicity in $\mathcal{C}(Z^d)$ such that for each $1 \leq k \leq m$ the following relations hold.

$$(3.12) \quad \begin{cases} Z_k = \sum_{j=1}^{N_{\delta}} \varphi_{k,j}(P_j H_1, \dots, P_j H_d) P_j, \\ \max_{1 \le j \le N_{\delta}} \|\varphi_{k,j}(P_j H_1, \dots, P_j H_d) P_j - \varphi_{k,j}(x_{1,j}, \dots, x_{d,j}) P_j \| \le \delta \end{cases}$$

If for each k we set

$$(3.13) Z_{k,\delta} := \sum_{j=1}^{N_{\delta}} \varphi_{k,j}(x_{1,j}, \dots, x_{d,j}) P_j = \sum_{j=1}^{N_{\delta}} \varphi_{k,j}(P_j \tilde{H}_{\delta,1}, \dots, P_j \tilde{H}_{\delta,d}) P_j,$$

then the *m*-tuple $\tilde{\mathbf{Z}}_{\delta} := (Z_{1,\delta}, \dots, Z_{m,\delta})$ belongs to $\mathbb{V}^m[Z^d](n)$ and the commutation relations

$$[Z_k, Z_{l,\delta}] = \mathbf{0}_n,$$

hold for each $1 \le k, l \le m$. By (3.11), (3.12) and (3.13) we have that the relations (3.7) hold for each $1 \le k \le m$. By (3.8) and (3.13) we will have that the following estimates hold for each $1 \le k \le m$.

$$||Z_{k} - Z_{k,\delta}|| = ||(\sum_{j=1}^{N_{\delta}} P_{j})(Z_{k} - Z_{k,\delta}))||$$

$$= ||\sum_{j=1}^{N_{\delta}} (\varphi_{k,j}(P_{j}H_{1}, \dots, P_{j}H_{d})P_{j} - \varphi_{k,j}(x_{1,j}, \dots, x_{d,j})P_{j})||$$

$$= \max_{1 \leq j \leq N_{\delta}} ||\varphi_{k,j}(P_{j}H_{1}, \dots, P_{j}H_{d})P_{j} - \varphi_{k,j}(x_{1,j}, \dots, x_{d,j})P_{j}||$$

$$(3.15) \leq \delta.$$

By (3.13), (3.14) and (3.15) we will have that $\tilde{\mathbf{Z}}_{\delta}$ is a hermitian δ -CPMA of \mathbf{Z} . This completes the proof.

Definition 3.7 (Uniform piecewise differentiable local connectivity in M_n^m). If for any given $\varepsilon > 0$, there is $\delta > 0$ such that for any two m-tuples \mathbf{X} and \mathbf{Y} in a matrix variety $\mathbb{Z}^m \subset M_n^m$ such that $\eth(\mathbf{X}, \mathbf{Y}) \leq \delta$ (\mathbf{X} and \mathbf{Y} are $(\delta_{\varepsilon}, \eth)$ -close), we have that $\mathbf{X} \leadsto_{\varepsilon(\delta)} \mathbf{Y}$ relative to $N_{\eth}(\mathbf{X}, \varepsilon) \cap \mathbb{Z}^m$. We will say that the matrix variety $\mathbb{Z}^m \subseteq M_n^m$ is uniformly locally piecewise differentiably connected (**ULPDC**).

Remark 3.2. It is important to remark that ε and δ must not depend on n.

Remark 3.3. It is important to notice that the components of two m-tuples $\mathbf{X} = (X_1, \ldots, X_m)$ and $\mathbf{Y} = (Y_1, \ldots, Y_m)$ in $\mathbb{I}^m(n)$ need not to satisfy the commutation relations $[X_j, Y_k] = \mathbf{0}_n$ for each j, k in general.

3.0.4. Approximate Joint Isospectral Interpolating Paths. Let $\varepsilon > 0$ and let $m \ge 1$ be an integer, in this section we will prove that there is $\delta > 0$ such that for any integer $n \ge 1$, any m-tuple $\mathbf{H} = (H_1, \ldots, H_m) \in \mathbb{I}^m(n)$ and any unitary matrix W in M_n that satisfy the relations $\|WH_j - H_jW\| < \delta$, $1 \le j \le m$. There exist a unitary \tilde{W} and a δ -CPMA $\tilde{\mathbf{H}} = (\tilde{H}_1, \ldots, \tilde{H}_m) \in C^*(H_1, \ldots, H_m)' \cap \mathbb{I}^m(n)$ such

that $W^*\tilde{H}_jW=\tilde{W}^*\tilde{H}_j\tilde{W}$ for each $1\leq j\leq m$. Moreover, there is a piecewise differentiable unitary path $\{W(t)\}_{t\in[0,1]}\subset\mathbb{U}(n)$ such that $W(0)=\mathbf{1}_n,\,W(1)=\tilde{W}$ and $\|W(t)\tilde{H}_j-\tilde{H}_jW(t)\|<\varepsilon$ for each $1\leq j\leq n$ and each $0\leq t\leq 1$.

Remark 3.4. It can be seen that if we set $\Psi = \widehat{\mathrm{Ad}[W]}$, $\widetilde{\Psi} = \widehat{\mathrm{Ad}[\widetilde{W}]}$ and $\psi_t = \widehat{\mathrm{Ad}[W(t)]}$, we will have that.

$$(3.16) \begin{cases} \psi_0(\tilde{\mathbf{H}}) = \mathrm{id}_{\mathbb{I}^m(n)}(\tilde{\mathbf{H}}) = \tilde{\mathbf{H}}, \\ \psi_1(\tilde{\mathbf{H}}) = \Psi(\tilde{\mathbf{H}}) = \tilde{\Psi}(\tilde{\mathbf{H}}), \\ \psi_t(\tilde{\mathbf{H}})[j]\psi_t(\hat{\mathbf{H}})[k] = \psi_t(\tilde{\mathbf{H}})[k]\psi_t(\tilde{\mathbf{H}})[j], \ 1 \leq j, k \leq m, 0 \leq t \leq 1, \\ \psi_t(\tilde{\mathbf{H}}) \in B_{\overline{0}}(\tilde{\mathbf{H}}, \varepsilon), \ 0 \leq t \leq 1, \\ \Lambda(\psi_t(\tilde{\mathbf{H}})) = \Lambda(\tilde{\mathbf{H}}), \ 0 \leq t \leq 1, \end{cases}$$

Any path of inner *-automorphisms like $\{\psi_t\}_{t\in[0,1]}$, that satisfies (3.16) will be called a ε -approximate joint isospectral interpolant with respect to $(\tilde{\Psi}, \tilde{\mathbf{H}}, \mathbb{I}^m(n))$.

Lemma 3.5. Given $r \in \mathbb{Z}^+$ and $\nu > 0$, there is $\delta := \hat{\delta}(\nu, r) > 0$ such that given a unitary W and a normal contraction D in M_n with $n \geq 2$, if $D = \sum_{j=1}^r \alpha_j P_j$ for $2 \leq r \in \mathbb{Z}$ with $\alpha_1, \ldots, \alpha_r \in \mathbb{D}^2$, and the set $\{P_j\}$ is an orthogonal partition of unity consisting of diagonal projectors, then there is a unitary matrix $Z \in M_n$ such that $[Z, D] = \mathbf{0}_n$ and $\|\mathbf{1}_n - WZ\| \leq \nu$ whenever $\|WDW^* - D\| \leq \delta$.

Proof. Let $\mu:=\|WDW^*-D\|=\|WD-DW\|$. Then there are r continuous functions $\ell_1,\dots,\ell_r\in C(\mathbb{D}^2)$ such that $P_k:=\ell_k(D)$. By Lemma 2.1 one can find $\delta:=\hat{\delta}(\nu,r)>0$ such that, $\|WP_kW^*-P_k\|=\|W\ell_k(D)W^*-\ell_k(D)\|=\|\ell_k(WDW^*)-\ell_k(D)\|\leq \nu/(\sqrt{2}r)<1/(\sqrt{2}r)$ whenever $\mu\leq\delta$. Since $\nu<1$, by perturbation theory of amenable C^* -algebras (in the sense of [15, Lemma 2.5.1]) we will have that there is a unitary W_j such that $\|\mathbf{1}-W_j\|\leq\sqrt{2}\|WP_jW^*-P_j\|\leq\nu/r$ and $W_j^*P_jW_j=WP_jW^*$, and this implies that $W_jWP_j=P_jW_jW$ for each j. Since $\{P_j\}$ is an orthogonal partition of unity, we will have that $\tilde{W}:=\sum_j W_jWP_j$ is a unitary matrix which satisfies the commutation relation $[\tilde{W},D]=0$ together with the normed inequalities.

$$||W - \tilde{W}|| = ||W \sum_{j} P_{j} - \sum_{j} W_{j} W P_{j}||$$

$$= ||\sum_{j} ((\mathbf{1}_{n} - W_{j}) W P_{j})||$$

$$\leq \sum_{j} ||\mathbf{1}_{n} - W_{j}||$$

$$\leq \sum_{j} \frac{1}{r} \nu = r(\frac{1}{r} \nu) = \nu$$

Let us set $Z:=\tilde{W}^*$ then $\|\mathbf{1}_n-WZ\|=\|\tilde{W}-W\|\leq\nu$. This completes the proof.

Lemma 3.6. Given $\nu > 0$ and any $m \in \mathbb{Z}^+$, there is $\delta > 0$ such that for any two m-tuples $\mathbf{X}, \mathbf{Y} \in \mathbb{I}^m(n)$ that satisfy the constraint $\eth(\mathbf{X}, \mathbf{Y}) < \delta$, there exist two inner morphisms Ψ and $\tilde{\Psi}$ of $\mathbb{I}^m(n)$, together with a hermitian δ -CPMA $\tilde{\mathbf{X}} \in \mathbb{I}^m(n)$ of \mathbf{X} such that $\tilde{\Psi}(\tilde{\mathbf{X}}) = \Psi(\tilde{\mathbf{X}})$, $\tilde{\Psi}(\tilde{\mathbf{X}})$ is a hermitian δ -CPMA of $\Psi(\mathbf{X})$, $\Psi(\mathbf{X})$ is a

hermitian ν -CPMA of \mathbf{Y} and $\eth(\tilde{\mathbf{X}}, \tilde{\Psi}(\tilde{\mathbf{X}})) < \nu$. Moreover, there is a ν -approximate joint isospectral interpolant $\{\psi_t\}_{t\in[0,1]}$ with respect to $(\tilde{\Psi}, \tilde{\mathbf{X}}, \mathbb{I}^m(n))$.

Proof. Let us consider two m-tuples \mathbf{X} and \mathbf{Y} in $\mathbb{I}^m(n)$, and let us set $\mu := \eth(\mathbf{X}, \mathbf{Y})$. We can assume, by changing basis if necessary, that \mathbf{Y} is a m-tuple of diagonal matrices. By Lemma 2.2 we will have that there exists a C^* -homomorphism $\Phi: M_n \to M_n$ defined by the expression $\Phi = \operatorname{Ad}[W]$ for some $W \in \mathbb{U}(n)$, such that $\hat{\Phi}(\mathbf{X})$ is a m-tuple of diagonal contractions and

(3.17)
$$\mathfrak{J}(\hat{\Phi}(\mathbf{X}), \mathbf{Y}) \le K_m \mu.$$

By applying Lemma 3.2 to the hermitian m-tuple

$$\hat{\Phi}(\mathbf{X}) = (\hat{Y}_1, \dots, \hat{Y}_m),$$

we can obtain a hermitian μ -CPMA

$$\tilde{\mathbf{Y}} = (\tilde{Y}_1, \dots, \tilde{Y}_m)$$

of $\hat{\Phi}(\mathbf{X})$. We will have that the *m*-tuple $\tilde{\mathbf{X}} = \hat{\Phi}^{-1}(\tilde{\mathbf{Y}})$ satisfies the constraint $\eth(\mathbf{X}, \tilde{\mathbf{X}}) \leq \mu$.

As a consequence of Lemma 3.2 there exists a projective decomposition $P_{\delta}(\hat{\Phi}(\mathbf{X}))$ of the form

$$(3.20) P_{\delta}(\hat{\Phi}(\mathbf{X})) = \{P_{1}, \dots, P_{N_{\delta}}\}$$

$$= \begin{cases} P_{1,j_{1}}P_{2,j_{2}} \cdots P_{m,j_{m}} \middle| P_{k,j_{k}} \in P_{\delta}(\hat{Y}_{k}), \\ 1 \leq j_{k} \leq N_{\delta,k}, \\ 1 \leq k \leq m, \end{cases} \},$$

where each \hat{Y}_k is the kth component of m-tuple (3.18). As a consequence of the proof of Lemma 3.2, we have that for each $1 \leq k \leq m$ and each $1 \leq j_k \leq N_\delta$, there exists a continuous function $p_{k,j_k} \in C([-1,1])$ such that

$$(3.21) P_{k,j_k} = p_{k,j_k}(\tilde{Y}_k),$$

where each \tilde{Y}_k is determined by (3.19). By Lemma 2.1 we will have that given $0 < \nu_1 < 1$, there is $\delta > 0$ such that

$$\|\Phi^{-1}(P_{k,j_k}) - P_{k,j_k}\| = \|\Phi^{-1}(p_{k,j_k}(\tilde{Y}_k)) - p_{k,j_k}(\tilde{Y}_k)\|$$

$$= \|W^* p_{k,j_k}(\tilde{Y}_k)W - p_{k,j_k}(\tilde{Y}_k)\|$$

$$= \|p_{k,j_k}(W^* \tilde{Y}_k W) - p_{k,j_k}(\tilde{Y}_k)\|$$

$$< \nu_1/(2\sqrt{2}mN_{\delta}) < 1/(2\sqrt{2}mN_{\delta})$$
(3.22)

whenever $\mu < \delta$. As a consequence of (3.22) we will have that for each projection $P_j = P_{1,j_1} \cdots P_{m,j_m}$ in $P_{\delta}(\hat{\Phi}(\mathbf{X}))$ determined by the relations (3.20), the following estimate will hold.

$$||W^*P_jW - P_j|| = ||\Phi^{-1}(P_j) - P_j||$$

$$\leq \sum_{k=1}^m ||\Phi^{-1}(P_{k,j_k}) - P_{k,j_k}||$$

$$< m\nu_1/(2m\sqrt{2}N_\delta) = \nu_1/(2\sqrt{2}N_\delta)$$

Since the inequalities (3.23) hold for each P_j in $P_{\delta}(\hat{\Phi}(\mathbf{X}))$, we can apply Lemma 3.5 with $r = N_{\delta}$ to compute a unitary $\tilde{W} \in M_n$ that satisfies the constraint

$$||\mathbf{1}_n - \tilde{W}^* W|| = ||W - \tilde{W}|| \le \nu_1/2,$$

together with the commutation relations

$$\tilde{W}P_i = P_i\tilde{W}$$

for each P_j in $P_{\delta}(\hat{\Phi}(\mathbf{X}))$. Let us set $\tilde{\Phi} = \operatorname{Ad}[Z]$ with $Z = \tilde{W}^*W$. By Lemma 3.2 we have that each \tilde{Y}_j determined by (3.19) belongs to span $P_{\delta}(\hat{\Phi}(\mathbf{X}))$, as a consequence of (3.25) we will have that

$$\tilde{\Phi}(\Phi^{-1}(\tilde{Y}_i)) = \tilde{W}^* \tilde{Y}_i \tilde{W} = \tilde{Y}_i = \Phi(\Phi^{-1}(\tilde{Y}_i)),$$

by applying Lemma 2.2 again together with the commutation relations (3.25) we will also have that

$$\|\tilde{\Phi}(\Phi^{-1}(\tilde{Y}_{j})) - \Phi^{-1}(\tilde{Y}_{j})\| = \|\tilde{W}^{*}\tilde{Y}_{j}\tilde{W} - W^{*}\tilde{Y}_{j}W\|$$

$$= \|\tilde{Y}_{j} - W^{*}\tilde{Y}_{j}W\|$$

$$\leq \|\tilde{Y}_{j} - \hat{Y}_{j}\| + \|\hat{Y}_{j} - W^{*}\hat{Y}_{j}W\|$$

$$+ \|W^{*}\hat{Y}_{j}W - W^{*}\tilde{Y}_{j}W\|$$

$$\leq 2\|\tilde{Y}_{j} - \hat{Y}_{j}\| + \|\Phi(X_{j}) - X_{j}\|$$

$$< 2\delta + K_{m}\delta = (K_{m} + 2)\delta.$$

Since $\|\mathbf{1}_n - Z\| < \nu_1/2 < 1$ by (3.24), we will have that there is a hermitian matrix contraction $H \in M_n$ such that $Z = e^{\mathbf{i}\pi H/2}$. If for each $t \in [0,1]$ we set $\phi_t := \mathrm{Ad}[e^{\mathbf{i}\pi t H/2}]$, we will have that $\phi_t : M_n \to M_n$ is a *-homomorphism for each $0 \le t \le 1$ such that $\phi_0 = \mathrm{id}_{M_n}$ and $\phi_1 = \tilde{\Phi}$. We will also have that for each $0 \le t \le 1$ and each \tilde{Y}_i in (3.19)

$$(3.28) \|\phi_t(\Phi^{-1}(\tilde{Y}_i)) - \Phi^{-1}(\tilde{Y}_i)\| \le 2\|\mathbf{1}_n - e^{\mathbf{i}\pi tH/2}\| \le 2\|\mathbf{1}_n - e^{\mathbf{i}\pi H/2}\| < \nu_1.$$

Let us set $\nu := \max\{\nu_1, (K_m + 2)\delta\}$, $\Psi = \hat{\Phi}$, $\tilde{\Psi} := \hat{\Phi}$ and $\psi_t := \hat{\phi}_t$ for each $t \in [0, 1]$. By Remark 2.1 we have that the maps Ψ , $\tilde{\Psi}$ and all members of the family $\{\psi_t\}_{t \in [0, 1]}$ preserve commutativity, hermiticity and the spectrum of each component Z_j of any m-tuple $\mathbf{Z} \in \mathbb{I}^m(n)$, which implies that all of them belong to the set of morphisms of $\mathbb{I}^m(n)$. By the relations (3.26) we have that.

$$\tilde{\Psi}(\tilde{\mathbf{X}}) = (\tilde{\Phi}(\Phi^{-1}(\tilde{Y}_1)), \dots, \tilde{\Phi}(\Phi^{-1}(\tilde{Y}_m)))$$

$$= (\Phi(\Phi^{-1}(\tilde{Y}_1)), \dots, \Phi(\Phi^{-1}(\tilde{Y}_m)))$$

$$= \Psi(\tilde{\mathbf{X}}).$$

By the properties of (3.19), by relation (3.29) and by inequalities (3.17) and (3.27) we also have that

(3.30)
$$\begin{cases} \vec{\eth}(\mathbf{X}, \tilde{\mathbf{X}}) = \vec{\eth}(\Psi(\mathbf{X}), \tilde{\Psi}(\tilde{\mathbf{X}})) = \vec{\eth}(\Psi(\mathbf{X}), \Psi(\tilde{\mathbf{X}})) < \delta, \\ \vec{\eth}(\mathbf{Y}, \Psi(\mathbf{X})) = \vec{\eth}(\mathbf{Y}, \hat{\Phi}(\mathbf{X})) < \nu \\ \vec{\eth}(\tilde{\mathbf{X}}, \tilde{\Psi}(\tilde{\mathbf{X}})) \le \max_{j} \|\tilde{\Phi}(\Phi^{-1}(\tilde{Y}_{j})) - \Phi^{-1}(\tilde{Y}_{j})\| < \nu, \end{cases}$$

and by combining (3.29) and (3.30) we will have that $\tilde{\Psi}(\tilde{\mathbf{X}}) = \Psi(\tilde{\mathbf{X}})$, $\tilde{\Psi}(\tilde{\mathbf{X}})$ is a hermitian δ -CPMA of $\Psi(\mathbf{X})$, $\Psi(\mathbf{X})$ is a hermitian ν -CPMA of \mathbf{Y} and $\tilde{\eth}(\tilde{\mathbf{X}}, \tilde{\Psi}(\tilde{\mathbf{X}})) <$

 ν . By definition of the elements in the family $\{\psi_t\}_{t\in[0,1]}$ we will have that $\psi_0 = \widehat{\mathrm{id}_{M_n}} = \mathrm{id}_{\mathbb{I}^m(n)}, \ \psi_1 = \widehat{\mathrm{Ad}[Z]} = \widetilde{\Psi}$ and for each $t\in[0,1]$ by (3.28) we will have that

$$\eth(\psi_t(\tilde{\mathbf{X}}), \tilde{\mathbf{X}}) \le \max_j \|\phi_t(\Phi^{-1}(\tilde{Y}_j)) - \Phi^{-1}(\tilde{Y}_j)\| < \nu.$$

This completes the proof.

Theorem 3.2. The matrix variety $\mathbb{I}^m(n)$ is uniformly locally piecewise differentiably connected.

Proof. As a consequence of Lemma 3.6 we will have that given $\nu > 0$, there is $\delta > 0$, such that for any two m-tuples $\mathbf{X}, \mathbf{Y} \in \mathbb{I}^m(n)$ that satisfy the constraint $\eth(\mathbf{X}, \mathbf{Y}) < \delta$, there exist a m-tuple $\tilde{X} \in \mathbb{I}^m(n)$, two inner morphisms Ψ and $\tilde{\Psi}$ of $\mathbb{I}^m(n)$, and a family $\{\psi_t\}_{t\in[0,1]}$ of morphisms of $\mathbb{I}^m(n)$ such that $\tilde{\Psi}(\tilde{\mathbf{X}}) = \Psi(\tilde{\mathbf{X}})$, $\tilde{\Psi}(\tilde{\mathbf{X}})$ is a hermitian δ -CPMA of $\Psi(\mathbf{X})$, $\Psi(\mathbf{X})$ is a hermitian ν -CPMA of \mathbf{Y} and

(3.31)
$$\eth(\tilde{\mathbf{X}}, \tilde{\Psi}(\tilde{\mathbf{X}})) < \nu.$$

Moreover, the family of morphisms $\{\psi_t\}_{t\in[0,1]}$ of $\mathbb{D}^m(n)$ satisfies the following conditions.

(3.32)
$$\begin{cases} \psi_0 = \mathrm{id}_{\mathbb{I}^m(n)}, \ \psi_1 = \tilde{\Psi}, \\ \tilde{\eth}(\tilde{\mathbf{X}}, \psi_t(\tilde{\mathbf{X}})) < \nu, 0 \le t \le 1 \end{cases}$$

Given any two *m*-tuples in $\mathbf{H} = (H_1, \dots, H_m)$ and $\mathbf{K} = (K_1, \dots, K_m)$ in $\mathbb{I}^m(n)$, let us write $\ell_{\mathbf{H}, \mathbf{K}}^H$ to denote the path in $C([0, 1], M_n^m)$ defined by the expression

(3.33)
$$\ell_{\mathbf{H},\mathbf{K}}^{H}(t) = ((1-t)H_1 + tK_1, \dots, (1-t)H_m + tK_m), \ 0 \le t \le 1.$$

It can be seen that $(\ell_{j,\mathbf{H},\mathbf{K}}^H(t))^* = ((1-t)H_j + tK_j)^* = (1-t)H_j + tK_j = \ell_{j,\mathbf{H},\mathbf{K}}^H(t)$ for each $1 \leq j \leq m$ and each $0 \leq t \leq 1$. It can also be seen that $\ell_{\mathbf{H},\mathbf{K}}^H(0) = \mathbf{H}$, $\ell_{\mathbf{H},\mathbf{K}}^H(1) = \mathbf{K}$, for any $\ell_{\mathbf{H},\mathbf{K}}^H(1) = \mathbf{K}$, and for each $\ell_{\mathbf{H},\mathbf{K}}^H(1) = \ell_{\mathbf{H},\mathbf{K}}^H(1) = \ell$

(3.34)
$$\kappa(t) = \psi_t(\tilde{\mathbf{X}}), \ 0 \le t \le 1.$$

By Lemma 3.6 and by (3.32) we will have that $\kappa(0) = \tilde{\mathbf{X}}$, $\kappa(1) = \tilde{\Psi}(\tilde{\mathbf{X}})$, $\kappa(t) \in \mathbb{I}^m(n)$ and $\eth(\kappa(t), \tilde{\mathbf{X}}) = \eth(\kappa(t), \kappa(0)) < \nu$ for each $t \in [0, 1]$.

Let us set $\varepsilon := 2(\nu + \delta)$. By the arguments presented in the previous paragraphs, we will have that the piecewise differentiable path $\gamma \in C([0,1], M_n^m)$ defined by the expression

$$\gamma = ((\ell_{\mathbf{X},\tilde{\mathbf{X}}}^H \circledast \kappa) \circledast \ell_{\tilde{\Psi}(\tilde{\mathbf{X}}),\Psi(\mathbf{X})}^H) \circledast \ell_{\Psi(\mathbf{X}),\mathbf{Y}}^H,$$

satisfies the conditions

(3.35)
$$\begin{cases} \gamma(0) = \mathbf{X}, \ \gamma(1) = \mathbf{Y}, \\ \gamma(t) \in \mathbb{I}^m(n), \ 0 \le t \le 1, \\ \eth(\gamma(t), \mathbf{X}) < \varepsilon, \ 0 \le t \le 1. \end{cases}$$

By (3.35) we have that $\mathbf{X} \leadsto_{\varepsilon(\delta)} \mathbf{Y}$ relative to $N_{\eth}(\mathbf{X}, \varepsilon) \cap \mathbb{I}^{m}(n)$. This completes the proof.

Corollary 3.1. The matrix variety $\mathbb{D}^m(n)$ is uniformly locally piecewise differentiably connected.

Proof. Given any two m-tuples \mathbf{Z} and \mathbf{S} in $\mathbb{D}^m(n)$, such that $\eth(\mathbf{Z},\mathbf{S}) \leq r$ for some r > 0, there are two semisimple commuting hermitian partitions $\pi(\mathbf{Z})$ and $\pi(\mathbf{S})$ of \mathbf{Z} and \mathbf{S} respectively, such that $\eth(\pi(\mathbf{Z}), \pi(\mathbf{S})) \leq r$. By the previously described fact, we can apply Theorem 3.2 to the corresponding semisimple commuting hermitian partitions of any two $(\delta_{\varepsilon}, \eth)$ -close m-tuples \mathbf{Z}, \mathbf{S} in $\mathbb{D}^m(n)$, replacing δ_{ε} by $\delta_{\varepsilon}/2$ if necessary, to obtain a path $\gamma^H \in C([0,1], \mathbb{I}^m(n))$ that solves the problem $\pi(\mathbf{Z}) \leadsto_{\varepsilon(\delta)} \pi(\mathbf{S})$ relative to $\mathbb{I}^m(n) \cap B_{\eth}(\pi(\mathbf{Z}), \varepsilon)$. The path $\hat{\gamma}^D$ obtained by taking the juncture $\hat{\gamma}^D(t) = v(\gamma^H(t))$ for each $t \in [0,1]$, will solve the problem $\mathbf{Z} \leadsto_{\varepsilon(\delta)} \mathbf{S}$ relative to $\mathbb{D}^m(n) \cap B_{\eth}(\mathbf{Z}, \varepsilon)$. This completes the proof.

Theorem 3.3. Given a compact differentiable d-manifold $Z^d \subseteq [-1,1]^m \subseteq \mathbb{R}^m$, the induced compact differentiable matrix manifold $\mathbb{V}^m[Z^d](n)$ is uniformly locally piecewise differentiably connected.

Proof. As a consequence of Lemma 3.4 and Remarks 2.1 and 3.4, by a similar argument to the one implemented in the proof of Lemma 3.6 we will have that given $\nu_1 > 0$, there is $\delta > 0$, such that for any two m-tuples $\mathbf{U}, \mathbf{V} \in \mathbb{V}^m[Z^d](n)$ that satisfy the constraint $\eth(\mathbf{U}, \mathbf{V}) \leq \delta$, there exist: a δ -CPMA $\tilde{\mathbf{U}} \in \mathbb{V}^m[Z^d](n)$ of \mathbf{U} , two inner morphisms Ψ and $\tilde{\Psi}$ of $\mathbb{V}^m[Z^d](n)$, and a family $\{\psi_t\}_{t\in[0,1]}$ of morphisms of $\mathbb{V}^m[Z^d](n)$ such that $\tilde{\Psi}(\tilde{\mathbf{U}}) = \Psi(\tilde{\mathbf{U}})$, $\tilde{\Psi}(\tilde{\mathbf{U}})$ is a δ -CPMA of $\Psi(\mathbf{U})$, $\Psi(\mathbf{U})$ is a normal ν_1 -CPMA of \mathbf{V} and

(3.36)
$$\tilde{\eth}(\tilde{\mathbf{U}}, \tilde{\Psi}(\tilde{\mathbf{U}})) < \nu_1.$$

Moreover, the family of morphisms $\{\psi_t\}_{t\in[0,1]}$ of $\mathbb{D}^m(n)$ satisfies the following conditions.

$$(3.37) \qquad \begin{cases} \psi_0 = \mathrm{id}_{\mathbb{V}^m[Z^d](n)}, \ \psi_1 = \tilde{\Psi}, \\ \eth(\tilde{\mathbf{U}}, \psi_t(\tilde{\mathbf{U}})) < \nu_1, 0 \le t \le 1 \\ \psi_t(\tilde{\mathbf{U}})[k]\psi_t(\tilde{\mathbf{U}})[j] = \psi_t(\tilde{\mathbf{U}})[j]\psi_t(\tilde{\mathbf{U}})[k], 1 \le j, k \le m \\ \Lambda(\psi_t(\tilde{\mathbf{U}})) \subseteq Z^d, 0 \le t \le 1 \end{cases}$$

Given any two m-tuples in $\mathbf{W} = (W_1, \dots, W_m)$ and $\mathbf{Z} = (Z_1, \dots, Z_m)$ in $\mathbb{V}^m[Z^d](n)$ such that $Z_j W_k = W_k Z_j$ for each $1 \leq j, k \leq m$, let us set $r := \eth(\mathbf{W}, \mathbf{Z})$ and let us write $\ell_{\mathbf{W}, \mathbf{Z}}^V$ to denote the path in $C([0, 1], M_n^m)$ defined coordinate wise by the expressions:

$$(3.38) \quad \ell_{\mathbf{W},\mathbf{Z}}^{V,k,j}(t) = \varphi_{k,j}((1-t)H_{W,1} + tH_{Z,1}, \dots, (1-t)H_{W,d} + tH_{Z,d})P_j, \quad 0 \le t \le 1,$$

with $1 \leq k \leq m$. Here for each $1 \leq j \leq d$, $H_{W,j}$, $H_{Z,j}$ are the hermitian matrices described by Lemma 3.4, which satisfy the relations $W_k = \sum_j \varphi_{k,j}(H_{W,1}, \dots, H_{W,d})P_j$, $Z_k = \sum_j \varphi_{k,j}(H_{Z,1}, \dots, H_{Z,d})P_j$ and $\eth(\mathbf{H}_W, \mathbf{H}_Z) < \nu(r)$, and ν is the function corresponding to the parametrization of the components of (3.38). It can be seen that

$$(3.39) \qquad \left\{ \begin{array}{l} \ell_{\mathbf{W},\mathbf{Z}}^{V}(0) = \mathbf{W}, \ell_{\mathbf{H},\mathbf{K}}^{V}(1) = \mathbf{Z} \\ \eth(\ell_{\mathbf{W},\mathbf{Z}}^{V}(t), \ell_{\mathbf{W},\mathbf{Z}}^{V}(s)) \leq \nu(\eth(\ell_{\mathbf{H}_{\mathbf{W}},\mathbf{H}_{\mathbf{Z}}}^{H}(t), \ell_{\mathbf{H}_{\mathbf{W}},\mathbf{H}_{\mathbf{Z}}}^{H}(s))), 0 \leq s, t \leq 1, \end{array} \right.$$

with $\ell_{\mathbf{H_W},\mathbf{H_Z}}^H$ defined by (3.33), and for each $0 \leq t \leq 1$ we have that the m-tuple $(\ell_{\mathbf{W},\mathbf{Z}}^{V,1}(t),\ldots,\ell_{\mathbf{W},\mathbf{Z}}^{V,m}(t))$, with $\ell_{\mathbf{W},\mathbf{Z}}^{V,k}(t) = \sum_j \varphi_{k,j}((1-t)H_{W,1}+tH_{Z,1},\ldots,(1-t)H_{W,d}+tH_{Z,d})P_j$, $1 \leq k \leq m$, belongs to $\mathbb{V}^m[Z^d](n)$. By the previous facts we will have that $\ell_{\mathbf{W},\mathbf{Z}}^V(t) \in \mathbb{V}^m[Z^d](n)$ for each $0 \leq t \leq 1$. Let us write κ to denote the path defined by the following expression

(3.40)
$$\kappa(t) = \psi_t(\tilde{\mathbf{U}}), \ 0 \le t \le 1.$$

By Lemma 3.6 and (3.37) we will have that $\kappa(0) = \tilde{\mathbf{U}}$, $\kappa(1) = \tilde{\Psi}(\tilde{\mathbf{U}})$, $\kappa(t) \in \mathbb{V}^m[Z^d](n)$ and $\eth(\kappa(t), \tilde{\mathbf{U}}) = \eth(\kappa(t), \kappa(0)) < \nu_1$ for each $t \in [0, 1]$.

Let us set $\varepsilon := 2(\nu_1 + \max\{\max_{x \in [0,\delta]} \nu(x), \delta\})$. By the arguments presented in the previous paragraphs, we will have that the path $\gamma \in C([0,1], M_n^m)$ defined by the expression

$$\gamma = ((\ell_{\mathbf{U},\tilde{\mathbf{U}}} \circledast \kappa) \circledast \ell_{\tilde{\Psi}(\tilde{\mathbf{U}}),\Psi(\mathbf{U})}) \circledast \ell_{\Psi(\mathbf{U}),\mathbf{V}},$$

satisfies the conditions

(3.41)
$$\begin{cases} \gamma(0) = \mathbf{U}, \ \gamma(1) = \mathbf{V}, \\ \gamma(t) \in \mathbb{V}^m[Z^d](n), \ 0 \le t \le 1, \\ \eth(\gamma(t), \mathbf{U}) < \varepsilon, \ 0 \le t \le 1. \end{cases}$$

By (3.35) we have that $\mathbf{U} \leadsto_{\varepsilon(\delta)} \mathbf{V}$ relative to $N_{\eth}(\mathbf{U}, \varepsilon) \cap \mathbb{V}^m[Z^d](n)$. This completes the proof.

3.1. Graphical analogies of continuous matrix deformations. Given a d-mainfold $Z^d \subseteq [-1,1]^m$, the effect of an approximate joint isospectral interpolant $\psi \in C([0,1], M_n^m)$ on a section of an arbitrary compact matrix manifold $\mathbb{V}^m[Z^d](n) \subseteq \mathbb{I}^m(n)$ in M_n can be interpreted as a matrix representation of a continuous deformation of Z^d , like the one described in Figure 3 in the particular case when $Z^d = \mathbb{S}^2$.




FIGURE 3. Graphical illustration of an approximate isospectral deformation in $V^3[\mathbb{S}^2](n)$.

Given a fixed but arbitrary 3-tuple $\mathbf{X} \in V^3[\mathbb{S}^2](n) \subseteq \mathbb{I}^3(n)$, we can think of the first homeomorphic copy of \mathbb{S}^2 on the left in Figure 3 as the section of $V^3[\mathbb{S}^2](n)$ corresponding to the joint spectra $\Lambda(\hat{\mathscr{D}}(V^3[\mathbb{S}^2](n) \cap C^*(\mathbf{X})))$ (the image of the full pinching operator extended to 3-tuples in $V^3[\mathbb{S}^2](n)$) with respect to a basis in which \mathbf{X} is a m-tuple of diagonal matrices, while the second homeomorphic copy of \mathbb{S}^2 on the right in Figure 3 can be interpreted as the section of $V^3[\mathbb{S}^2](n)$ corresponding

to the joint spectra $\Lambda(\hat{\mathscr{D}}(V^3[\mathbb{S}^2](n) \cap C^*(\psi_1(\mathbf{X}))))$ on the same basis in which **X** is a *m*-tuple of diagonal matrices.

Let us consider two m-tuples \mathbf{X} and \mathbf{Y} in a certain matrix variety $Z^m \subseteq M_n^m$. We can think of each path $X_j \leadsto_{\varepsilon(\delta)} \tilde{X}_j$ relative to $N_{\overline{o}}(X_j, \varepsilon)$, as a noncommutative analogy of the family of paths (**pseudospectral-links**) connecting the point sets determined by the spectra $\sigma(\mathcal{D}(X_j))$ and $\sigma(\tilde{X}_j)$ relative to the embedding of $\mathbb{D}^2 \times [0,1]$ in $[-1,1]^2 \times [0,1]$ (where the embedding is induced by the mapping $\mathbb{D}^2 \to B(0,1) \subseteq [-1,1]^2$). In a similar way we can interpret the induced paths $\mathbf{X} \leadsto_{\varepsilon(\delta)} \tilde{\mathbf{X}}$ as noncommutative analogies of the links connecting the joint spectra $\sigma(\mathbf{X})$ and $\sigma(\tilde{\mathbf{X}})$ relative to $[-1,1]^{2n} \times [0,1]$. An illustration of these analogies are presented in F.4, F.5 and F.6.




FIGURE 4. Graphical illustration of a the pseudospectral-links in $\mathbb{D}^2 \times [0,1]$ corresponding to a matrix path that connects the spectra of two normal contractions X and Y (with Y diagonal) in M_{10} with $\sigma(\mathscr{D}(X)) \subseteq \partial B(0,3/5)$ and $\sigma(\mathscr{D}(Y)) = \sigma(Y) \subseteq \mathbb{T}^1$.




FIGURE 5. Graphical illustration of a the pseudospectral-links in \mathbb{D}^2 corresponding to a matrix path that connects the spectra of two normal contractions X and Y (with Y diagonal) in M_{55} with $\sigma(\mathscr{D}(X)) \subseteq \partial B(0, 7/10)$ and $\sigma(\mathscr{D}(Y)) = \sigma(Y) \subseteq \mathbb{T}^1$.

The paths described in F.5 and F.6 provide a graphical illustration of the ideas behind the *pseudospectral clustering* technique implemented in the proofs of Theorems 3.2 and 3.3.




FIGURE 6. Graphical illustration of a the pseudospectral-paths in \mathbb{S}^2 corresponding to a matrix path that connects the spectra of two triples of pairwise commuting hermitian contractions $\mathbf{X} = (X_1, X_2, X_3)$ and $\mathbf{Y} = (Y_2, Y_2, Y_3)$ in $\mathbb{V}^3[\mathbb{S}^2](7)$.

4. Applications

4.1. Local Connectivity of some particular Compate differentiable Matrix Manifolds. Let us consider the semialgebraic matrix varieties $\mathbb{T}^m(n)$ and $\mathbb{S}^{m-1}(n)$ defined as follows.

(4.1)
$$\mathbb{T}^{m}(n) = \left\{ \mathbf{U} \in \mathbb{D}^{2}(n) \mid U_{k}U_{k}^{*} = \mathbf{1}_{n}, 1 \leq j, k \leq m \right\}$$

(4.2)
$$\mathbb{S}^{m-1}(n) = \left\{ \mathbf{H} \in \mathbb{I}^m(n) \mid \sum_{k=1}^m H_k^2 = \mathbf{1}_n \right\}$$

Lemma 4.1. The matrix semialgebraic variety $\mathbb{T}^m(n)$ is **ULPDC**.

Proof. Let $m \in \mathbb{Z}^+$. Since $\mathbb{T}^m(n)$ is defined by (4.1), we will have that if for any $\mathbf{U} \in \mathbb{T}^m(n)$ we set

(4.3)
$$\pi_H(\mathbf{U}) = (Re(U_1), Im(U_1), \dots, Re(U_m), Im(U_m)),$$

then $\Lambda(\pi_H(\mathbf{U})) \subseteq \mathbb{T}^m = \prod_{j=1}^m \mathbb{S}^1 \subseteq \prod_{j=1}^m [-1,1]^2 \subseteq \mathbb{R}^{2m}$. By considering the mapping $v_H: M_n^{2m} \to M_n^m, (X_1, X_2, \dots, X_{2m-1}, X_{2m}) \mapsto (X_1 + \mathbf{i}X_2, \dots, X_{2m-1} + \mathbf{i}X_{2m})$, it can be seen that the mapping $\pi_H: M_n^m \to M_n^{2m}$ determined by equation (4.3) induces a one to one correspondence between $\mathbb{T}^m(n)$ and $V^{2m}(\mathbb{T}^m)(n)$. Because of this correspondence, we have that any $\mathbf{U} \in \mathbb{T}^m(n)$ has a representation of the form $\mathbf{U} = v_H(\pi_H(\mathbf{U}))$.

Given any two (ν, \eth) -close tuples $\mathbf{U}, \mathbf{V} \in \mathbb{T}^m(n)$ for some given $\nu > 0$, we will have that the tuples $\pi_H(\mathbf{U}), \pi_H(\mathbf{V}) \in V^{2m}[\mathbb{T}^m](n)$ will be (ν, \eth) -close. By Theorem 3.3 we have that $V^{2m}(\mathbb{T}^m)(n)$ is \mathbf{ULPDC} , this implies that given $\varepsilon > 0$, there is $\delta > 0$ such that if $\nu < \delta$, there is a piecewise differentiable path $\gamma \in C([0,1],V^{2m}[\mathbb{T}^m](n))$ that solves the connectivity problem $\pi_H(\mathbf{U}) \leadsto_{\varepsilon/2} \pi_H(\mathbf{V})$ relative to $V^{2m}[\mathbb{T}^m](n) \cap B_{\eth}(\pi_H(\mathbf{U}),\varepsilon/2)$. By the previous facts we will have that

the path $\hat{\gamma}$ defined by the equation $\hat{\gamma}(t) = v_H(\gamma(t))$ for each $t \in [0, 1]$, solves the connectivity problem $\mathbf{U} \leadsto_{\varepsilon} \mathbf{V}$ relative to $\mathbb{T}^m(n) \cap B_{\overline{o}}(\mathbf{U}, \varepsilon)$. This completes the proof.

Lemma 4.2. The matrix semialgebraic variety $\mathbb{S}^{m-1}(n)$ is **ULPDC**.

Proof. Let $m \in \mathbb{Z}^+$. Since $\mathbb{S}^{m-1}(n)$ is defined by (4.2), we will have that for any $\mathbf{H} \in \mathbb{S}^{m-1}(n)$, $\Lambda(\mathbf{H}) \subseteq \mathbb{S}^{m-1} \subseteq \Pi_{j=1}^m[-1,1] \subseteq \mathbb{R}^{m+1}$. It can be seen that the identity mapping $\mathrm{id}_{M_n^m}$ induces a one to one correspondence between $\mathbb{S}^{m-1}(n)$ and $V^m(\mathbb{S}^{m-1})(n)$. Because of this correspondence, we have in fact that $\mathbb{S}^{m-1}(n) = V^m(\mathbb{S}^{m-1})(n)$. Since by Theorem 3.3 we have that $V^m(\mathbb{S}^{m-1})(n)$ is \mathbf{ULPDC} , this implies that $\mathbb{S}^{m-1}(n) = V^m(\mathbb{S}^{m-1}(n))$ is \mathbf{ULPDC} . This completes the proof. \square

4.2. Local Connectivity of Commutative C^* -Subalgebras of Matrix Algebras. Let us consider the universal C^* -algebras $C([-1,1]^m)$, $C(\mathbb{T}^m)$ and $C(\mathbb{T}^1 \times [-1,1])$ described in terms of generators and relations by the expressions.

$$C([-1,1]^{m}) := C_{1}^{*} \left\langle h_{1}, \dots, h_{m} \middle| \begin{array}{c} [h_{j}, h_{k}] = 0 \\ -1 \leq h_{j} \leq 1, \end{array}, 1 \leq j, k \leq m \right\rangle,$$

$$(4.4)$$

$$C(\mathbb{T}^{m}) := C_{1}^{*} \left\langle u_{1}, \dots, u_{m} \middle| \begin{array}{c} [u_{j}, u_{k}] = 0, \\ u_{j} u_{j}^{*} = u_{j}^{*} u_{j} = 1, \end{array}, 1 \leq j, k \leq m \right\rangle,$$

$$(4.5)$$

$$C(\mathbb{S}^{m-1}) := C_{1}^{*} \left\langle h_{1}, \dots, h_{m} \middle| \begin{array}{c} [h_{j}, h_{k}] = 0 = h_{j} - h_{j}^{*}, \ 1 \leq j, k \leq m \\ \sum_{j=1}^{m} h_{j}^{2} = 1 \end{array} \right\rangle.$$

$$(4.6)$$

4.2.1. **ULPDC** property for C^* -algebras. Given a universal C^* -algebra A that can be described by the expression $A := C_1^*\langle x_1, \ldots, x_m, \mathcal{R}(x_1, \ldots, x_m) \rangle$, we say that $C^*\operatorname{Rep}(A, \mathcal{M}_{\infty})$ (the set of all C^* -representations of A in $\mathcal{M}_{\infty} := \overline{\bigcup_{n \in \mathbb{Z}^+} M_n}^{\|\cdot\|}$) is uniformly locally piecewise differentiably connected or (**ULPDC**) if for any two m-tuples of normal contractions $\mathbf{X} := (X_1, \ldots, X_m)$ and $\mathbf{Y} := (Y_1, \ldots, Y_m)$ in M_n^m such that there exist *-representations $\pi_X \in C^*\operatorname{Rep}(A, C^*(\mathbf{X}))$ and $\pi_Y \in C^*\operatorname{Rep}(A, C^*(\mathbf{Y}))$ determined by the equations $\pi_X(x_j) = X_j$ and $\pi_Y(x_j) = Y_j$, $1 \le j \le m$, we have that there are piecewise differentiable contractive normal matrix paths \hat{X}^j that solve the problems $X_j \leadsto_{\varepsilon(\delta)} Y_j$, for some function $\varepsilon : \mathbb{R}_0^+ \to \mathbb{R}_0^+$, with $\eth(\mathbf{X}, \mathbf{Y}) \le \delta$, and if in addition there is $\pi_t \in C^*\operatorname{Rep}(A, C^*(\hat{X}^1(t), \ldots, \hat{X}^m(t)))$ for each $0 \le t \le 1$.

Theorem 4.1. For any integer $m \geq 1$ we have that $C^*\operatorname{Rep}(C[-1,1]^m, \mathcal{M}_{\infty})$ is **ULPDC**.

Proof. Since we have that $C([-1,1]^m)$ is described by 4.4. It can be seen that for any $\rho_n \in C^*\operatorname{Rep}(C([-1,1]^m), M_n)$ the m-tuple $(\rho_n(h_1), \ldots, \rho_n(h_m))$ belongs to $I^m(n)$. By Theorem 3.2, for any given $\varepsilon > 0$, there is $\delta > 0$ such that for any two elements $\rho_n, \rho'_n \in C^*\operatorname{Rep}(C([-1,1]^m), M_n)$ that satisfy the constraint

$$\eth((\rho_n(h_1),\ldots,\rho_n(h_m)),(\rho'_n(h_1),\ldots,\rho'_n(h_m))) \leq \delta,$$

there is a piecewise differentiable path $\gamma \in C([0,1],\mathbb{I}^m(n))$ that solves the problem

$$(\rho_n(h_1),\ldots,\rho_n(h_m)) \leadsto_{\varepsilon} (\rho'_n(h_1),\ldots,\rho'_n(h_m))$$

relative to $\mathbb{I}^m(n) \cap B_{\overline{o}}(\rho_n(h_1), \ldots, \rho_n(h_m), \varepsilon)$. By universality of $C([-1, 1]^m)$ we will have that there is a *-representation

$$\pi_t : C([-1,1]^m) \to C^*(\gamma(t)[1], \dots, \gamma(t)[m]) \subseteq M_n$$

for each $0 \le t \le 1$. This completes the proof.

Theorem 4.2. For any integer $m \geq 1$ we have that $C^*Rep(C(\mathbb{T}^m), \mathcal{M}_{\infty})$ is **ULPDC**.

Proof. Since we have that $C(\mathbb{T}^m)$ is described by 4.5. It can be seen that for any $\rho_n \in C^*\operatorname{Rep}(C(\mathbb{T}^m), M_n)$ the m-tuple $(\rho_n(u_1), \dots, \rho_n(u_m))$ belongs to $\mathbb{T}^m(n)$. By Lemma 4.1, for any given $\varepsilon > 0$, there is $\delta > 0$ such that for any two elements $\rho_n, \rho'_n \in C^*\operatorname{Rep}(C(\mathbb{T}^m), M_n)$ that satisfy the constraint

$$\eth((\rho_n(u_1),\ldots,\rho_n(u_m)),(\rho'_n(u_1),\ldots,\rho'_n(u_m))) \leq \delta,$$

there is a piecewise differentiable path $\gamma \in C([0,1], \mathbb{T}^m(n))$ that solves the problem

$$(\rho_n(u_1),\ldots,\rho_n(u_m)) \leadsto_{\varepsilon} (\rho'_n(u_1),\ldots,\rho'_n(u_m)),$$

relative to $\mathbb{T}^m(n) \cap B_{\eth}(\rho_n(u_1), \ldots, \rho_n(u_m), \varepsilon)$. By universality of $C(\mathbb{T}^m)$ we will have that there is a *-representation

$$\pi_t: C(\mathbb{T}^m) \to C^*(\gamma(t)[1], \dots, \gamma(t)[m]) \subseteq M_n$$

for each $0 \le t \le 1$. This completes the proof.

Theorem 4.3. For any integer $m \geq 1$ we have that $C^*\operatorname{Rep}(C(\mathbb{S}^{m-1}), \mathcal{M}_{\infty})$ is **ULPDC**.

Proof. Since we have that $C(\mathbb{S}^{m-1})$ is described by 4.6. It can be seen that for any $\rho_n \in C^*\operatorname{Rep}(C(\mathbb{S}^{m-1}), M_n)$ the m-tuple $(\rho_n(h_1), \ldots, \rho_n(h_m))$ belongs to $\mathbb{S}^{m-1}(n)$. By Lemma 4.2, for any given $\varepsilon > 0$, there is $\delta > 0$ such that for any two elements $\rho_n, \rho'_n \in C^*\operatorname{Rep}(C(\mathbb{S}^{m-1}), M_n)$ that satisfy the constraint

$$\eth((\rho_n(h_1),\ldots,\rho_n(h_m)),(\rho'_n(h_1),\ldots,\rho'_n(h_m))) \leq \delta,$$

there is a piecewise differentiable path $\gamma \in C([0,1],\mathbb{S}^{m-1}(n))$ that solves the problem

$$(\rho_n(h_1),\ldots,\rho_n(h_m)) \leadsto_{\varepsilon} (\rho'_n(h_1),\ldots,\rho'_n(h_m)),$$

relative to $\mathbb{S}^{m-1}(n) \cap B_{\eth}(\rho_n(h_1), \dots, \rho_n(h_m), \varepsilon)$. By universality of $C(\mathbb{S}^{m-1})$ we will have that there is a *-representation

$$\pi_t: C(\mathbb{S}^{m-1}) \to C^*(\gamma(t)[1], \dots, \gamma(t)[m]) \subset M_n$$

for each $0 \le t \le 1$. This completes the proof.

4.2.2. Spherical Unitaries and Completely Positive Maps on Matrix Algebras. Given $m, n \in \mathbb{Z}^+$, let us consider any unital completely positive (**CP**) map (in the sense of [6] and [23]) $\psi_{\mathbf{S}} : \mathcal{M}_{\infty} \to \mathcal{M}_{\infty}$ determined for each $X \in M_n$ by some m-tuple $\mathbf{S} = (S_1, \ldots, S_m) \in M_n^m$ by the expression

(4.7)
$$\psi_{\mathbf{S}}(X) = \sum_{k=1}^{m} S_j X S_j^*,$$

We will call the map $\psi_{\mathbf{S}}$, the CP map induced by **S**. If in addition we have that $\mathbf{S} \in \mathbb{D}^m(n)$, and if the elements of **S** satisfy the contraint $\sum_{j=1}^m S_j S_j^* = \mathbf{1}_n$, we will call a map like $\psi_{\mathbf{S}}$ a m-spherical CP (m-SCP) map, since the m-tuple **S** is

a matrix representation of the universal operators known as spherical unitaries in the sense of [8] and [23]. The set of m-SCP maps acting on M_n will be denoted by $\mathrm{SCP}_m(M_n)$, similarly the set of m-SCP maps acting on \mathcal{M}_{∞} will be denoted by $\mathrm{SCP}_m(\mathcal{M}_{\infty})$. We will call any m-tuple like \mathbf{S} a spherical unitary, and will denote by $\mathbb{SU}^m(n)$ the set of all spherical unitaries in M_n^m described as follows.

(4.8)
$$\mathbb{SU}^m(n) = \left\{ \mathbf{S} \in \mathbb{D}^m(n) \middle| \sum_{k=1}^m S_k S_k^* = \mathbf{1}_n \right\}$$

Lemma 4.3. The matrix semialgebraic variety $\mathbb{SU}^m(n)$ is **ULPDC**.

Proof. Let $m \in \mathbb{Z}^+$ and let us consider the maps π_H and v_H implemented in the proof of Lemma 4.1. Since $\mathbb{SU}^m(n)$ is defined by (4.8), we will have that for any $\mathbf{S} \in \mathbb{SU}^m(n)$ we have that $\Lambda(\pi_H(\mathbf{S})) \subseteq \mathbb{S}^{2m-1} \subseteq \Pi_{j=1}^{2m}[-1,1] \subseteq \mathbb{R}^{2m}$. As in the proof of Lemma 4.1 we will have that there is a one to one correspondence between $\mathbb{SU}^m(n)$ and $V^{2m}(\mathbb{S}^{2m-1})(n)$. Because of this correspondence, we have that any $\mathbf{S} \in \mathbb{SU}^m(n)$ has a representation of the form $\mathbf{S} = v_H(\pi_H(\mathbf{S}))$.

Given any two (ν, \eth) -close tuples $\mathbf{S}, \mathbf{T} \in \mathbb{SU}^m(n)$ for some given $\nu > 0$, we will have that the tuples $\pi_H(\mathbf{S}), \pi_H(\mathbf{T}) \in V^{2m}[\mathbb{S}^{2m-1}](n)$ will be (ν, \eth) -close. By Theorem 3.3 we have that $V^{2m}(\mathbb{S}^{2m-1})(n)$ is \mathbf{ULPDC} , this implies that given $\varepsilon > 0$, there is $\delta > 0$ such that if $\nu < \delta$, there is a piecewise differentiable path $\gamma \in C([0,1],V^{2m}[\mathbb{S}^{2m-1}](n))$ that solves the connectivity problem $\pi_H(\mathbf{S}) \leadsto_{\varepsilon/2} \pi_H(\mathbf{T})$ relative to $V^{2m}[\mathbb{S}^{2m-1}](n) \cap B_{\eth}(\pi_H(\mathbf{U}),\varepsilon/2)$. By the previous facts we will have that the path $\hat{\gamma}$ defined by the equation $\hat{\gamma}(t) = \upsilon_H(\gamma(t))$ for each $t \in [0,1]$, solves the connectivity problem $\mathbf{S} \leadsto_{\varepsilon} \mathbf{T}$ relative to $\mathbb{SU}^m(n) \cap B_{\eth}(\mathbf{U},\varepsilon)$. This completes the proof.

Theorem 4.4. For any integer $m \ge 1$ and any universal spherical unitary $\mathbf{S} = (s_1, \ldots, s_m)$ we have that $C^*\operatorname{Rep}(C^*(\mathbf{S}), \mathcal{M}_{\infty})$ is **ULPDC**.

Proof. Since we have that $C^*(\mathbf{S})$ is described in terms of generators and relations by the expression.

(4.9)
$$C^*(\mathbf{S}) := C_1^* \left\langle s_1, \dots, s_m \middle| \begin{array}{l} [s_j, s_k] = 0 = s_j s_j^* - s_j^* s_j, & 1 \le j, k \le m \\ \sum_{j=1}^m s_j s_j^* = 1 \end{array} \right\rangle.$$

It can be seen that for any $\rho_n \in C^*\operatorname{Rep}(C^*(\mathbf{S}), M_n)$ the *m*-tuple $(\rho_n(s_1), \dots, \rho_n(s_m))$ belongs to $\mathbb{SU}^m(n)$. By Lemma 4.3, for any given $\varepsilon > 0$, there is $\delta > 0$ such that for any two elements $\rho_n, \rho'_n \in C^*\operatorname{Rep}(C^*(\mathbf{S}), M_n)$ that satisfy the constraint

$$\eth((\rho_n(s_1),\ldots,\rho_n(s_m)),(\rho'_n(s_1),\ldots,\rho'_n(s_m))) \leq \delta,$$

there is a piecewise differentiable path $\gamma \in C([0,1], \mathbb{SU}^m(n))$ that solves the problem

$$(\rho_n(s_1),\ldots,\rho_n(s_m)) \leadsto_{\varepsilon} (\rho'_n(s_1),\ldots,\rho'_n(s_m)),$$

relative to $\mathbb{SU}^m(n) \cap B_{\eth}(\rho_n(s_1), \ldots, \rho_n(s_m), \varepsilon)$. By universality of $C^*(\mathbf{S})$ we will have that there is a *-representation

$$\pi_t: C^*(\mathbf{S}) \to C^*(\gamma(t)[1], \dots, \gamma(t)[m]) \subseteq M_n$$

for each 0 < t < 1. This completes the proof.

Let us write $CP(M_n)$ to denote the set of completely positive linear maps on M_n , let us extend the **ULPDC** property to subsets of $CP(M_n)$.

Definition 4.1. We say that s subset $S \subseteq CP(M_n)$ is **ULPDC** if given two maps $\psi_{\mathbf{S}}, \psi_{\mathbf{T}} \in S$ induced by two m-tuples $\mathbf{S}, \mathbf{T} \in M_n^m$ and given $\varepsilon > 0$, there is $\delta > 0$ such that if $\eth(\mathbf{S}, \mathbf{T}) < \delta$, then there is a piecewise differentiable path $\{\Psi_t\}_{t \in [0,1]} \subseteq S$ such that $\Psi_0 = \psi_{\mathbf{S}}, \Psi_1 = \psi_{\mathbf{T}}$, and $\|\Psi_t(X) - \Psi_{\mathbf{S}}(X)\| < \varepsilon$ for each $t \in [0,1]$ and any contraction $X \in M_n$.

The following lemma was motivated by some connections between the connectivity properties of $CP(\mathcal{M}_{\infty})$ and the classification of quantum phases in the sense of [22].

Lemma 4.4. For any integer $m \geq 1$ we have that $SCP_m(\mathcal{M}_{\infty})$ is **ULPDC**.

Proof. Let $m \in \mathbb{Z}^+$. It can be seen that for any $\mathbf{S}, \mathbf{T} \in \mathbb{SU}^m(n)$ and any $X \in M_n$ such that $||X|| \leq 1$, the following estimate holds for the corresponding induced CP maps.

$$\|\psi_{\mathbf{S}}(X) - \psi_{\mathbf{T}}(X)\| = \left\| \sum_{j=1}^{m} S_{j} X S_{j}^{*} - \sum_{j=1}^{m} T_{j} X T_{j}^{*} \right\|$$

$$\leq \sum_{j=1}^{m} \|S_{j} X S_{j}^{*} - T_{j} X T_{j}^{*}\|$$

$$\leq \sum_{j=1}^{m} 2 \|S_{j} - T_{j}\| \|X\|$$

$$\leq 2m \eth(\mathbf{S}, \mathbf{T}).$$
(4.10)

Since $\mathbb{SU}^m(n)$ is **ULPDC** by Lemma 4.3, we have that given $\varepsilon > 0$, there is $\delta > 0$ such that for any two m-tuples $\mathbf{S}, \mathbf{T} \in \mathbb{SU}^m(n)$ that satisfy the constraint $\eth(\mathbf{S}, \mathbf{T}) < \delta$, there is a piecewise differentiable path $\gamma \in C([0, 1], \mathbb{SU}^m(n))$ that solves the problem $\mathbf{S} \leadsto_{\varepsilon/(2m)} \mathbf{T}$ relative to $\mathbb{SU}^m(n) \cap B_{\eth}(\mathbf{S}, \varepsilon/(2m))$. Let us set $\Psi_t = \psi_{\gamma(t)}$ for each $t \in [0, 1]$. It can be seen that $\Psi_0 = \psi_{\mathbf{S}}, \ \Psi_1 = \psi_{\mathbf{T}}, \ \Psi_t \in \mathrm{SCP}_m(M_n)$ for each $t \in [0, 1]$, and by the estimate (4.10) we will have that for any $X \in M_n$ such that $\|X\| \leq 1$,

$$\|\Psi_t(X) - \psi_{\mathbf{S}}(X)\| \le 2m\mathfrak{F}(\gamma(t), \mathbf{S}) < 2m(\varepsilon/(2m)) = \varepsilon.$$

This completes the proof.

5. Hints and Future Directions

The implications of the results in sections §3 and §4 in uniform local path connectivity of other subsets of $CP(\mathcal{M}_{\infty})$ will be further explored, some connections to fixed pont approximations for particular types of elements in $CP(\mathcal{M}_{\infty})$ will be studied as well.

By considering matrix paths as continuous/differentiable analogies of numerical linear algebra algorithms in the sense of [7], we will work on the adaptation and application of the results in sections §3 and §4 to the structure-preserving approximation/perturbation of families of structured matrices with some particular spectral behavior in the sense of [1, 2, 18, 17, 21].

ACKNOWLEDGEMENT

I am very grateful with the Fields Institute for Research in Mathematical Sciences at the University of Toronto, for their support during my research visit to the Institute from July 31 to August 4, 2017. Much of the research reported in this document was carried out while I was visiting the Institute.

I am grateful with Terry Loring, Alexandru Buium, Alexandru Chirvasitu, Moody Chu, Marc Rieffel, Stanly Steinberg, Yossi Avron, Masoud Khalkhali and Concepción Ferrufino, for several interesting questions and comments that have been very helpful for the preparation of this document.

References

- M. AHUES, A. LARGILLIER, F. DIAS D'ALMEIDA AND P. B. VASCONCELOS. Spectral refinement on quasi-diagonal matrices. Linear Algebra Appl. 401 (2005) 109117.
- [2] M. AHUES, F. DIAS D'ALMEIDA, A. LARGILLIER AND P. B. VASCONCELOS. Spectral refinement for clustered eigenvalues of quasi-diagonal matrices. Linear Algebra Appl. 413 (2006) 394402.
- [3] R. Bhatia. Matrix Analysis. Gaduate Texts in Mathematics 169. Springer-Verlag. 1997.
- [4] R. Bhatia Pinching, Trimming, Truncating, and Averaging of Matrices. Am. Math. Monthly Vol. 107, No. 7 (Aug. - Sep., 2000), pp. 602-608
- [5] O. Bratteli, G. A. Elliott, D. E. Evans and A. Kishimoto. Homotopy of a Pair of Approximately Commuting Unitaries in a Simple C*-Algebra, J. Funct. Anal., 160, p. 466-523. 1998.
- [6] M.-D. Choi. Completely Positive Linear Maps on Complex Matrices. Linear Algebra Appl. 10, 285-290 (1975).
- [7] M. T. Chu. Linear Algebra Algorithms as Dynamical Systems. Acta Numer. (2008), pp. 001-086, 2008.
- [8] K. R. DAVIDSON AND LE T. Commutant Lifting for Commuting Row Contractions. Bull. Lond. Math. Soc., Vol. 42, Issue 3, 1 June 2010, Pages 506516.
- [9] P. Friis and M. Rördam. Almost commuting self-adjoint matrices a short proof of Huaxin Lin's theorem. J. Reine Angew. Math., 479:121131, 1996.
- [10] M. H. FREEDMAN AND W. H. PRESS. Truncation of Wavelet Matrices: Edge Effects and the Reduction of Topological Control Linear Algebra Appl. 2:34:1-19 (1996)
- [11] D. Hadwin. Lifting Algebraic Elements in C*-Algebras. J. Funct. Anal. 127, 431-437 (1995).
- [12] R. C. Kirby. Stable homeomorphisms and the annulus conjecture. Ann. Math., Second Series, Vol. 89, No. 3 (May, 1969), pp. 575-582
- [13] H. Lin. Approximation by Normal Elements with Finite Spectra in Simple AF-Algebras. J. Operator Theory 31(1994), 83-98.
- [14] H. Lin. Almost Commuting Selfadjoint Matrices and Applications. In Operator algebras and their applications (Waterloo, ON, 1994/1995), volume 13 of Fields Inst. Commun., pages 193-233. Amer. Math. Soc., Providence, RI, 1997.
- [15] H. Lin. An introduction to the classification of amenable C*-algebras. World Scientific, River Edge, NJ, ISBN: 981-02-4680-3, pp 1320. 2001.
- [16] T. A. LORING AND F. VIDES. Local Matrix Homotopies and Soft Tori. arXiv:1605.06590 [math.OA]. 2016.
- [17] T. MAEHARA AND K. MUROTA. Algorithm for Error-Controlled Simultaneous Block-Diagonalization of Matrices. SIAM J. Matrix Anal. Appl., Vol. 32, No. 2, pp. 605-620, 2011.
- [18] K. C. O'MEARA AND C. VINSONHALER. On approximately simultaneously diagonalizable matrices. Linear Algebra Appl. 412 (2006) 39-74.
- [19] M. A. RIEFFEL. Actions of Finite Groups on C*-Algebras. Math. Scand. 47 (1980), 157-176. 1980.
- [20] M. A. RIEFFEL. Vector Bundles and Gromov-Hausdorff Distance. J. K-theory 5(2010), 39-103.
- [21] B. SAVAS AND I. S. DHILLON. Clustered Matrix Approximation. SIAM J. Matrix Anal. Appl. Vol. 37, No. 4, pp. 1531-1555, 2016.
- [22] O. SZEHR AND M. M. WOLF. Connected components of irreducible maps and 1D quantum phases. J. Math. Phys., 57, 081901 (2016); doi: 10.1063/1.4960557.

[23] H. ZHANG AND M. XUE. Fixed points of trace preserving completely positive maps. Linear and Multilinear Algebra, 2016 Vol. 64, No. 3, 404411.

DEPARTMENT OF APPLIED MATHEMATICS, SCHOOL OF MATHEMATICS, FACULTY OF SCIENCE, UNIVERSIDAD NACIONAL AUTÓNOMA DE HONDURAS, CIUDAD UNIVERSITARIA, TEGUCIGALPA, HONDURAS.

 $E\text{-}mail\ address: \verb|fredy.vides@unah.edu.hn||$