

# Pentavalent symmetric graphs admitting transitive non-abelian characteristically simple groups

Jia-Li Du, Yan-Quan Feng\*

*Department of Mathematics, Beijing Jiaotong University, Beijing 100044, China*

## Abstract

Let  $\Gamma$  be a graph and let  $G$  be a group of automorphisms of  $\Gamma$ . The graph  $\Gamma$  is called *G-normal* if  $G$  is normal in the automorphism group of  $\Gamma$ . Let  $T$  be a finite non-abelian simple group and let  $G = T^l$  with  $l \geq 1$ . In this paper we prove that if every connected pentavalent symmetric  $T$ -vertex-transitive graph is  $T$ -normal, then every connected pentavalent symmetric  $G$ -vertex-transitive graph is  $G$ -normal. This result, among others, implies that every connected pentavalent symmetric  $G$ -vertex-transitive graph is  $G$ -normal except  $T$  is one of 57 simple groups. Furthermore, every connected pentavalent symmetric  $G$ -regular graph is  $G$ -normal except  $T$  is one of 20 simple groups, and every connected pentavalent  $G$ -symmetric graph is  $G$ -normal except  $T$  is one of 17 simple groups.

**Keywords:** Vertex-transitive graph, symmetric graph, Cayley graph, regular permutation group, simple group.

**2010 Mathematics Subject Classification:** 05C25, 20B25.

## 1 Introduction

Throughout this paper, all groups and graphs are finite, and all graphs are simple and undirected. Denote by  $\mathbb{Z}_n$ ,  $D_n$ ,  $A_n$  and  $S_n$  the cyclic group of order  $n$ , the dihedral group of order  $2n$ , the alternating group and the symmetric group of degree  $n$ , respectively. Let  $G$  be a permutation group on a set  $\Omega$  and let  $\alpha \in \Omega$ . Denote by  $G_\alpha$  the stabilizer of  $\alpha$  in  $G$ , that is, the subgroup of  $G$  fixing the point  $\alpha$ . We say that  $G$  is *semiregular* on  $\Omega$  if  $G_\alpha = 1$  for every  $\alpha \in \Omega$ , and *regular* if it is semiregular and transitive. For a graph  $\Gamma$ , we denote its vertex set and automorphism group by  $V(\Gamma)$  and  $\text{Aut}(\Gamma)$ , respectively. The graph  $\Gamma$  is said to be *G-vertex-transitive* or *G-regular* for  $G \leq \text{Aut}(\Gamma)$  if  $G$  acts transitively or regularly on  $V(\Gamma)$  respectively, and *G-symmetric* if  $G$  acts transitively on the arc set of  $\Gamma$  (an arc is an ordered pair of adjacent vertices). In particular,  $\Gamma$  is *vertex-transitive* or *symmetric* if it is  $\text{Aut}(\Gamma)$ -vertex-transitive or  $\text{Aut}(\Gamma)$ -symmetric, respectively. A graph  $\Gamma$  is said to be *G-normal* for  $G \leq \text{Aut}(\Gamma)$  if  $G$  is normal in  $\text{Aut}(\Gamma)$ .

---

\*Corresponding author. E-mails: JiaLiDu@bjtu.edu.cn, yqfeng@bjtu.edu.cn

For a non-abelian simple group  $T$ ,  $T$ -vertex-transitive graphs have received wide attentions, specially for the two extreme cases:  $T$ -symmetric graphs and  $T$ -regular graphs. It was shown in [2] that a connected pentavalent symmetric  $T$ -vertex-transitive graph  $\Gamma$  is either  $T$ -normal or  $\text{Aut}(\Gamma)$  contains a non-abelian simple normal subgroup  $L$  such that  $T \leq L$  and  $(T, L)$  is one of 58 possible pairs of non-abelian simple groups.

A  $T$ -regular graph is also called a *Cayley graph* over  $T$ , and the Cayley graph is called *normal* if it is  $T$ -normal. Investigation of Cayley graphs over a non-abelian simple group is currently a hot topic in algebraic graph theory. One of the most remarkable achievements is the complete classification of connected trivalent symmetric non-normal Cayley graphs over non-abelian simple groups. This work was began in 1996 by Li [12], and he proved that a connected trivalent symmetric Cayley graph  $\Gamma$  over a non-abelian simple group  $T$  is either normal or  $T = A_5, A_7, \text{PSL}(2, 11), M_{11}, A_{11}, A_{15}, M_{23}, A_{23}$  or  $A_{47}$ . In 2005, Xu *et al* [16] proved that either  $\Gamma$  is normal or  $T = A_{47}$ , and two years later, Xu *et al* [17] further showed that if  $T = A_{47}$  and  $\Gamma$  is not normal, then  $\Gamma$  must be 5-arc-transitive and up to isomorphism there are exactly two such graphs. Du *et al* [2] showed that a connected pentavalent symmetric Cayley graph  $\Gamma$  over  $T$  is either normal, or  $\text{Aut}(\Gamma)$  contains a non-abelian simple normal subgroup  $L$  such that  $T \leq L$  and  $(T, L)$  is one of 13 possible pairs of non-abelian simple groups.

For  $T$ -symmetric graphs, Fang and Praeger [4, 5] classified such graphs when  $T$  is a Suzuki or Ree simple group acting transitively on the set of 2-arcs of the graphs. For a connected cubic  $T$ -symmetric graph  $\Gamma$ , it was proved by Li [12] that either  $\Gamma$  is  $T$ -normal or  $(T, \text{Aut}(\Gamma)) = (A_7, A_8), (A_7, S_8), (A_7, 2.A_8), (A_{15}, A_{16})$  or  $(\text{GL}(4, 2), \text{AGL}(4, 2))$ . Fang *et al* [3] proved that none of the above five pairs can happen, that is,  $T$  is always normal in  $\text{Aut}(\Gamma)$ . Du *et al* [2] showed that a connected pentavalent  $T$ -symmetric graph  $\Gamma$  is either  $T$ -normal or  $\text{Aut}(\Gamma)$  contains a non-abelian simple normal subgroup  $L$  such that  $T \leq L$  and  $(T, L)$  is one of 17 possible pairs of non-abelian simple groups.

Let  $G$  be the characteristically simple group  $T^l$  with  $l \geq 1$ . In this paper, we extend the above results on connected pentavalent  $T$ -vertex graphs to  $G$ -vertex graphs.

**Theorem 1.1** *Let  $T$  be a non-abelian simple group and let  $G = T^l$  with  $l \geq 1$ . Assume that every connected pentavalent symmetric  $T$ -vertex-transitive graph is  $T$ -normal. Then every connected pentavalent symmetric  $G$ -vertex-transitive graph is  $G$ -normal.*

In 2011, Hua *et al* [10] proved that if every connected cubic symmetric  $T$ -vertex-transitive graph is  $T$ -normal, then every connected cubic symmetric  $G$ -vertex-transitive graph is  $G$ -normal. By Theorem 1.1 and [2, Theorem 1.1], we have the following corollaries.

**Corollary 1.2** *Let  $T$  be a non-abelian simple group and let  $G = T^l$  with  $l \geq 1$ . Then every connected pentavalent symmetric  $G$ -vertex-transitive graph is  $G$ -normal except for  $T = \text{PSL}(2, 8), \Omega_8^-(2)$  or  $A_{n-1}$  with  $n \geq 6$  and  $n \mid 2^9 \cdot 3^2 \cdot 5$ .*

**Corollary 1.3** *Let  $T$  be a non-abelian simple group and let  $G = T^l$  with  $l \geq 1$ . Then every connected pentavalent  $G$ -symmetric graph is  $G$ -normal except for  $T = A_{n-1}$  with  $n = 2 \cdot 3, 2^2 \cdot 3, 2^4, 2^3 \cdot 3, 2^5, 2^2 \cdot 3^2, 2^4 \cdot 3, 2^3 \cdot 3^2, 2^5 \cdot 3, 2^4 \cdot 3^2, 2^6 \cdot 3, 2^5 \cdot 3^2, 2^7 \cdot 3, 2^6 \cdot 3^2, 2^7 \cdot 3^2, 2^8 \cdot 3^2$  or  $2^9 \cdot 3^2$ .*

**Corollary 1.4** *Let  $T$  be a non-abelian simple group and let  $G = T^l$  with  $l \geq 1$ . Then every connected pentavalent symmetric  $G$ -regular graph is  $G$ -normal except for  $T = \text{PSL}(2, 8)$ ,  $\Omega_8^-(2)$  or  $A_{n-1}$  with  $n = 2 \cdot 3, 2^3, 3^2, 2 \cdot 5, 2^2 \cdot 3, 2^2 \cdot 5, 2^3 \cdot 3, 2^3 \cdot 5, 2 \cdot 3 \cdot 5, 2^4 \cdot 5, 2^3 \cdot 3 \cdot 5, 2^4 \cdot 3^2 \cdot 5, 2^6 \cdot 3 \cdot 5, 2^5 \cdot 3^2 \cdot 5, 2^7 \cdot 3 \cdot 5, 2^6 \cdot 3^2 \cdot 5, 2^7 \cdot 3^2 \cdot 5$  or  $2^9 \cdot 3^2 \cdot 5$ .*

## 2 Preliminaries

In this section, we describe some preliminary results which will be used later. The first one is the vertex stabilizers of connected pentavalent symmetric graphs. By [7, Theorem 1.1], we have the following proposition.

**Proposition 2.1** *Let  $\Gamma$  be a connected pentavalent  $G$ -symmetric graph with  $v \in V(\Gamma)$ . Then  $G_v \cong \mathbb{Z}_5, D_5, D_{10}, F_{20}, F_{20} \times \mathbb{Z}_2, F_{20} \times \mathbb{Z}_4, A_5, S_5, A_4 \times A_5, S_4 \times S_5, (A_4 \times A_5) \rtimes \mathbb{Z}_2, \text{ASL}(2, 4), \text{AGL}(2, 4), \text{A}\Sigma\text{L}(2, 4), \text{A}\Gamma\text{L}(2, 4)$  or  $\mathbb{Z}_2^6 \rtimes \Gamma\text{L}(2, 4)$ , where  $F_{20}$  is the Frobenius group of order 20,  $A_4 \rtimes \mathbb{Z}_2 = S_4$  and  $A_5 \rtimes \mathbb{Z}_2 = S_5$ . In particular,  $|G_v| = 5, 2 \cdot 5, 2^2 \cdot 5, 2^2 \cdot 5, 2^3 \cdot 5, 2^4 \cdot 5, 2^2 \cdot 3 \cdot 5, 2^3 \cdot 3 \cdot 5, 2^4 \cdot 3^2 \cdot 5, 2^6 \cdot 3^2 \cdot 5, 2^5 \cdot 3^2 \cdot 5, 2^6 \cdot 3 \cdot 5, 2^6 \cdot 3^2 \cdot 5, 2^7 \cdot 3 \cdot 5, 2^7 \cdot 3^2 \cdot 5$  or  $2^9 \cdot 3^2 \cdot 5 \cdot 5$ , respectively.*

Connected pentavalent symmetric graphs admitting vertex-transitive non-abelian simple groups were classified in [2].

**Proposition 2.2** [2, Theorem 1.1] *Let  $T$  be a non-abelian simple group and  $\Gamma$  a connected pentavalent symmetric  $T$ -vertex-transitive graph. Then either  $T \leq \text{Aut}(\Gamma)$ , or  $T = \Omega_8^-(2), \text{PSL}(2, 8)$  or  $A_{n-1}$  with  $n \geq 6$  and  $n \mid 2^9 \cdot 3^2 \cdot 5$ .*

The following is straightforward (also see the short proof of [2, Lemma 3.2]).

**Proposition 2.3** *Let  $\Gamma$  be a connected pentavalent symmetric  $G$ -vertex-transitive graph with  $v \in V(\Gamma)$  and let  $A = \text{Aut}(\Gamma)$ . If  $H \leq A$  and  $GH \leq A$ , then  $|H|/|H \cap G| = |(GH)_v|/|G_v| \mid 2^9 \cdot 3^2 \cdot 5$ , and if  $\Gamma$  is further  $G$ -symmetric then  $|H|/|H \cap G| \mid 2^9 \cdot 3^2$ .*

The following proposition follows the classification of three-factor simple groups.

**Proposition 2.4** [11, Theorem I] *Let  $G$  be a non-abelian simple  $\{2, 3, 5\}$ -group. Then  $G = A_5, A_6$  or  $\text{PSU}(4, 2)$ .*

By Guralnick [8, Theorem 1], we have the following proposition.

**Proposition 2.5** *Let  $G$  be a non-abelian simple group with a subgroup  $H$  such that  $|G : H| = p^a$  with  $p$  a prime and  $a \geq 1$ . Then*

- (1)  $G = A_n$  and  $H = A_{n-1}$  with  $n = p^a$ ;
- (2)  $G = \text{PSL}(2, 11)$  and  $H = A_5$  with  $|G : H| = 11$ ;

- (3)  $G = M_{23}$  and  $H = M_{22}$  with  $|G : H| = 23$ , or  $G = M_{11}$  and  $H = M_{10}$  with  $|G : H| = 11$ ;
- (4)  $G = \text{PSU}(4, 2) \cong \text{PSp}(4, 3)$  and  $H$  is the parabolic subgroup of index 27;
- (5)  $G = \text{PSL}(n, q)$  and  $H$  is the stabilizer of a line or hyperplane with  $|G : H| = (q^n - 1)/(q - 1) = p^a$ .

By [13, Theorem 1] and Proposition 2.5, we have the following proposition.

**Proposition 2.6** *Let  $G$  be a non-abelian simple group and  $H$  a maximal subgroup of  $G$  such that  $|G : H| = 2^a \cdot 3^b \geq 6$  with  $0 \leq a \leq 9$  and  $0 \leq b \leq 2$ . Then  $G$ ,  $H$  and  $|G : H|$  are listed in Table 1.*

| $G$                | $H$                                   | $ G : H $       | $G$                | $H$   | $ G : H $               |
|--------------------|---------------------------------------|-----------------|--------------------|---|-------------------------|
| $M_{11}$           | $\text{PSL}(2, 11)$                   | $2^2 \cdot 3$   | $M_{12}$           | $M_{11}$  | $2^2 \cdot 3$           |
| $M_{24}$           | $M_{23}$                              | $2^3 \cdot 3$   | $\text{PSU}(3, 3)$ | $\text{PSL}(2, 7)$                                | $2^2 \cdot 3^2$         |
| $A_9$              | $S_7$                                 | $2^2 \cdot 3^2$ | $\text{PSU}(4, 2)$ | $S_6$   | $2^2 \cdot 3^2$         |
| $\text{PSp}(6, 2)$ | $S_8$                                 | $2^2 \cdot 3^2$ | $M_{12}$           | $\text{PSL}(2, 11)$                               | $2^4 \cdot 3^2$         |
| $\text{PSL}(2, 8)$ | $D_7$                                 | $2^2 \cdot 3^2$ | $\text{PSL}(3, 3)$ | $\mathbb{Z}_{13} \rtimes \mathbb{Z}_3$            | $2^4 \cdot 3^2$         |
| $\text{PSL}(2, 9)$ | $A_5$                                 | $2 \cdot 3$     | $\text{PSL}(2, p)$ | $\mathbb{Z}_p \rtimes \mathbb{Z}_{\frac{p-1}{2}}$ | $p + 1 = 2^a \cdot 3^b$ |
| $\text{PSL}(2, 8)$ | $\mathbb{Z}_2^3 \rtimes \mathbb{Z}_7$ | $3^2$           | $A_n$              | $A_{n-1}$   | $n = 2^a \cdot 3^b$     |

Table 1: Non-abelian simple group pairs of index  $2^a \cdot 3^b$

Let  $G$  be a group. The *inner automorphism group*  $\text{Inn}(G)$  of  $G$  is the group of automorphisms of  $G$  induced by conjugate action of elements in  $G$ , which is a normal subgroup in the automorphism group  $\text{Aut}(G)$  of  $G$ . The quotient group  $\text{Aut}(G)/\text{Inn}(G)$  is called the *outer automorphism group* of  $G$ . By the classification of finite simple groups, we have the following proposition, which is the famous Schreier conjecture.

**Proposition 2.7** [6, Theorem 1.64] *Every finite simple group has a solvable outer automorphism group.*

Baddeley and Praeger [1] considered almost simple groups containing a direct product of at least two isomorphic non-abelian simple groups.

**Proposition 2.8** [1, Theorem 1.4] *Let  $H$  be an almost simple group, that is,  $S \leq H \leq \text{Aut}(S)$  for a non-abelian simple group  $S$ , and suppose that  $H = AB$ , where  $A$  is a proper subgroup of  $H$  not containing  $S$ , and  $B \cong T^r$  for a non-abelian simple group  $T$  and integer  $k \geq 2$ . Then  $S = A_n$  and  $A \cap S = A_{n-1}$ , where  $n = |H : A| = |S : A \cap S| \geq 10$ .*

Let  $\Gamma$  be a graph and  $N \leq \text{Aut}(\Gamma)$ . The *quotient graph*  $\Gamma_N$  of  $\Gamma$  relative to  $N$  is defined as the graph with vertices the orbits of  $N$  on  $V(\Gamma)$  and with two orbits adjacent if there is an edge in  $\Gamma$  between these two orbits.

**Proposition 2.9** [14, Theorem 9] *Let  $\Gamma$  be a connected  $G$ -symmetric graph of prime valency, and let  $N \trianglelefteq G$  have at least three orbits on  $V(\Gamma)$ . Then  $N$  is the kernel of  $G$  on  $V(\Gamma_N)$ , and semiregular on  $V(\Gamma)$ . Furthermore,  $\Gamma_N$  is  $G/N$ -symmetric with  $G/N \leq \text{Aut}(\Gamma_N)$ .*

### 3 Proof of Theorem 1.1

Let  $T$  be a non-abelian simple group and let  $G = T^l$  with  $l \geq 1$ . Let  $\Gamma$  be a connected pentavalent symmetric  $G$ -vertex-transitive graph with  $v \in V(\Gamma)$  and let  $A = \text{Aut}(\Gamma)$ . We make the following assumption throughout this section.

**Assumption:** Every connected pentavalent symmetric  $T$ -vertex-transitive graph is  $T$ -normal.

Since the complete graph  $K_6$  of order 6 has automorphism group  $S_6$  and is  $A_5$ -symmetric, we have  $T \neq A_5$  by Assumption. Since  $G$  is vertex-transitive and has no subgroup of index 2,  $\Gamma$  is not bipartite.

To prove Theorem 1.1, we apply induction on  $l$ . It suffices to show that  $G$  contains a minimal normal subgroup of  $A$ , and this is done in Lemmas 3.1 and 3.2 when  $\text{rad}(A) = 1$ , where  $\text{rad}(A)$  is the largest solvable normal subgroup of  $A$ . For  $\text{rad}(A) \neq 1$ , we need the fact  $\text{rad}(A)G = \text{rad}(A) \times G$ , which is proved in Lemmas 3.3 and 3.4.

**Lemma 3.1** *Let  $\Gamma$  be  $X$ -symmetric with  $G \leq X$ , and let  $X$  have a minimal normal subgroup that is a direct product of  $T$ . Then  $G$  contains a minimal normal subgroup of  $X$ .*

**Proof:** By Assumption, if  $l = 1$  then  $G \trianglelefteq A$  and the lemma is true. Assume  $l \geq 2$ .

Let  $N$  be a minimal normal subgroup of  $X$  such that  $N = T^m$  for a positive integer  $m$ . Since  $N \cap G \trianglelefteq G$ , we have  $N \cap G = T^n$  with  $n \leq m$  and  $n \leq l$ . Set  $D = NG$ . By Proposition 2.3,  $|N|/|N \cap G| = |T|^{m-n} = |D|/|G| = |D_v|/|G_v| \mid 2^9 \cdot 3^2 \cdot 5$ . Since  $T$  is non-abelian simple, we have either  $m - n = 1$  or  $m = n$ .

Suppose  $m - n = 1$ . Then  $|T| \mid 2^9 \cdot 3^2 \cdot 5$ , and by Proposition 2.4,  $T = A_6$  as  $T \neq A_5$ . Thus,  $|G| = |A_6|^l$ ,  $|D| = |A_6|^{l+1}$  and  $|D_v|/|G_v| = |A_6|$ , implying  $5 \mid |D_v|$  and  $5 \nmid |G_v|$ . It follows that  $\Gamma$  is  $D$ -symmetric, but not  $G$ -symmetric.

Let  $\bar{D} = D/G_D$ ,  $\bar{G} = G/G_D$  and  $\bar{N} = NG_D/G_D$ , where  $G_D$  is the largest normal subgroup of  $D$  contained in  $G$ . Then  $\bar{D} = \bar{G}\bar{N}$ . Since  $\Gamma$  is  $D$ -symmetric,  $D_v$  is primitive on the set of 5 neighbours of  $v$  in  $\Gamma$ , and since  $G_D \trianglelefteq D$ , we have either  $(G_D)_v = 1$  or  $5 \mid |(G_D)_v|$ . Thus,  $(G_D)_v = 1$  because  $\Gamma$  is not  $G$ -symmetric. If  $G = G_D$ , then  $G_v = 1$  and  $|D_v| = |T| = 2^3 \cdot 3^2 \cdot 5$ , contrary to Proposition 2.1. Thus,  $G/G_D = T^{l'}$  with  $l' \geq 1$ .

If  $G_D$  is transitive on  $V(\Gamma)$ , then  $|G_v| = |G_v|/|(G_D)_v| = |G|/|G_D| = |T|^{l'}$ . Since  $T = A_6$ , we have  $5 \mid |G_v|$ , a contradiction. If  $G_D$  has two orbits on  $V(\Gamma)$ , then  $\Gamma$  is bipartite, which is impossible. Thus,  $G_D$  has at least three orbits on  $V(\Gamma)$ . By Proposition 2.9,  $\Gamma_{G_D}$  is a connected pentavalent  $G/G_D$ -vertex-transitive and  $D/G_D$ -symmetric graph. If  $l' = 1$ , by Proposition 2.2,  $G/G_D \trianglelefteq D/G_D$ , that is,  $G \trianglelefteq D$ , which is also impossible as otherwise  $G_v = 1$  and  $|D_v| = |T| = 2^3 \cdot 3^2 \cdot 5$ . Thus,  $\bar{G} = G/G_D = T^{l'}$  with  $l' \geq 2$  and  $|\bar{D}| = |T|^{l'+1}$  as  $|\bar{D} : \bar{G}| = |D : G| = |T|$ .

Let  $\bar{M}$  be a maximal subgroup of  $\bar{D}$  containing  $\bar{G}$ . Then  $\bar{G} \leq \bar{M}$  and  $|T|^{l'} \leq |\bar{M}| < |T|^{l'+1}$ . Consider the right multiplication of  $\bar{D}$  on  $[\bar{D} : \bar{M}]$ , the set of right cosets of  $\bar{M}$  in  $\bar{D}$ . Then  $\bar{D}/\bar{M}_{\bar{D}}$  is a primitive permutation group on  $[\bar{D} : \bar{M}]$  and  $\bar{D}/\bar{M}_{\bar{D}} \leq S_{|T|}$ , where  $\bar{M}_{\bar{D}}$  is the largest normal subgroup of  $\bar{D}$  contained in  $\bar{M}$ . Since  $|\bar{D} : \bar{M}| \mid |T| = |A_6|$ , the primitive group  $\bar{D}/\bar{M}_{\bar{D}}$  has degree dividing 360.

Let  $\bar{L}$  be a minimal normal subgroup of  $\bar{D}$ . Then  $\bar{L} \cap \bar{N} = \bar{L}$  or 1. For the former,  $\bar{L} \leq \bar{N}$ , and for the latter,  $\bar{L} = \bar{L}/\bar{L} \cap \bar{N} \cong \bar{N}\bar{L}/\bar{N} \trianglelefteq \bar{D}/\bar{N} \cong \bar{G}/\bar{G} \cap \bar{N}$ . In both cases,  $\bar{L}$  is a direct product of  $T$ , and hence  $\text{soc}(\bar{D}) = T^k$  for a positive integer  $k$ , where  $\text{soc}(\bar{D})$  is the product of all minimal normal subgroups of  $\bar{D}$ , that is, the socle of  $\bar{D}$ .

Assume  $\bar{M}_{\bar{D}} \neq 1$ . Then  $\bar{D}$  has a minimal normal subgroup contained in  $\bar{M}_{\bar{D}}$ , say  $\bar{K}$ . Then  $\bar{K} \trianglelefteq \text{soc}(\bar{D}) = T^k$ . If  $\bar{K} \not\leq \bar{G}$ , then  $|T|^{l'+1} = |\bar{D}| > |\bar{M}| \geq |\bar{G}\bar{K}| = |\bar{G}|(|\bar{K}|/|\bar{K} \cap \bar{G}|) \geq |\bar{G}||T| = |T|^{l'+1}$ , a contradiction. Thus,  $\bar{K} \leq \bar{G}$ , and we may assume  $\bar{K} = K/G_D$ , where  $G_D \leq K \leq G$ . Since  $\bar{K} \trianglelefteq \bar{D}$ , we have  $K \trianglelefteq D$ , and hence  $K \leq G_D$ . It follows that  $K = G_D$  and  $\bar{K} = 1$ , a contradiction.

Thus,  $\bar{M}_{\bar{D}} = 1$  and so  $\bar{D}$  is a primitive group on  $[\bar{D} : \bar{M}]$  of degree dividing  $|T| = 360$ . Since  $\text{soc}(\bar{D}) = T^k = A_6^k$ , we have  $k \leq 2$  by [15, Section 7], and the primitivity implies that  $|\bar{D}| \leq |\text{Aut}(\text{soc}(\bar{D}))| \leq 32|T|^2$ , which is impossible because  $|\bar{D}| = |T|^{l'+1} \geq |T|^3$ .

It follows that  $m - n = 0$  and hence  $N \leq G$ , as required.  $\square$

**Remark:** It is easy to check that Lemma 3.1 is also true if Assumption is replaced by the conditions that  $l \geq 2$  and  $T \neq A_5$ .

**Lemma 3.2** *Let  $G \leq X$  such that  $\text{rad}(X) = 1$  and  $\Gamma$  is  $X$ -symmetric. Then  $X$  has a minimal normal subgroup that is a direct product of  $T$ .*

**Proof:** Suppose to the contrary that every minimal normal subgroup of  $X$  is not a direct product of  $T$ . Then  $G = T^l \not\trianglelefteq X$ , and by Assumption,  $l \geq 2$ .

Let  $L$  be the socle of  $X$ , the product of all minimal normal subgroups of  $X$ . Since  $\text{rad}(X) = 1$ , we have  $L = S_1 \times \cdots \times S_t$  with  $t$  a positive integer, and  $S_i \not\cong T$  is a non-abelian simple group for each  $1 \leq i \leq t$ . Furthermore,  $L \cap G = T^r$  with  $0 \leq r \leq l$  as  $L \cap G \trianglelefteq G$ . By Proposition 2.3,  $|L|/|L \cap G| = |(GL)_v|/|G_v| \mid 2^9 \cdot 3^2 \cdot 5$ .

For convenience, write  $L \cap G = T^r = T_1 \times \cdots \times T_r$ . Let  $P_i$  be the projection from  $L \cap G$  to  $S_i$  for a given  $1 \leq i \leq t$ , that is,  $P_i(x) = x_i$  for any  $x = x_1 \cdots x_t \in L \cap G$  with  $x_i \in S_i$  ( $1 \leq i \leq t$ ). Then  $P_i$  is a homomorphism from  $L \cap G$  to  $S_i$  and hence  $(L \cap G)/K_i \cong I_i$ , where  $K_i$  is the kernel of  $P_i$  and  $I_i \leq S_i$  is the image of  $L \cap G$  under  $P_i$ . Since  $K_i \trianglelefteq L \cap G$ , we have  $(L \cap G)/K_i \cong T^{r_i} = I_i$ . Clearly,  $K_1 \leq S_2 \times \cdots \times S_t$  and  $K_1 \cap \cdots \cap K_t = 1$ . It follows  $T^r = L \cap G = (L \cap G)/(K_1 \cap \cdots \cap K_t) \lesssim (L \cap G)/K_1 \times \cdots \times (L \cap G)/K_t \cong T^{r_1} \times \cdots \times T^{r_t} = I_1 \times \cdots \times I_t$ , and hence  $(|S_1 : I_1| \cdots |S_t : I_t|) \mid |L : (L \cap G)| \mid 2^9 \cdot 3^2 \cdot 5$ .

Assume  $t \geq 5$ . Then at least four  $|S_i : I_i|$  are divisors of  $2^9 \cdot 3^2$ , say  $|S_i : I_i| = 2^{a_i} \cdot 3^{b_i}$  with  $2 \leq i \leq 5$ . Since  $T \neq A_5$ , Proposition 2.6 implies  $2^{a_i} \cdot 3^{b_i} \geq 2^3$  and  $|S_2 : I_2| \cdots |S_t : I_t| \geq 2^{12}$ . If  $5 \nmid |S_1 : I_1|$ , then  $|S_1 : I_1| \geq 2^3$  and  $|S_1 : I_1| \cdots |S_t : I_t| \geq 2^{15}$ . Since  $(|S_1 : I_1| \cdots |S_t : I_t|) \mid 2^9 \cdot 3^2 \cdot 5$ , we have  $2^{15} \leq 2^9 \cdot 3^2 \cdot 5$ , a contradiction. Thus,  $5 \mid |S_1 : I_1|$ . If  $|S_1 : I_1| \neq 5$  then  $|S_1 : I_1| \geq 2 \cdot 5$  and hence  $2^{13} \cdot 5 \leq 2^9 \cdot 3^2 \cdot 5$ , a contradiction. It follows  $|S_1 : I_1| = 5$ , which is also impossible by Proposition 2.5. This yields  $t \leq 4$ .

Let  $C = C_X(L)$  be the centralizer of  $L$  in  $X$ . Then  $C \cap L = 1$  and hence  $C = 1$  because  $L$  contains every minimal normal subgroup of  $X$ . Thus,  $X = X/C \leq \text{Aut}(L)$  and  $G/G \cap L \cong GL/L \leq X/L \leq \text{Out}(L)$ . Since  $t \leq 4$ , Proposition 2.7 implies that  $\text{Out}(L)$  is solvable, and hence  $G/G \cap L = 1$  and  $G \leq L$ . Since  $G \not\trianglelefteq X$ , we have  $G < L$ , that is,  $G$  is a proper subgroup of  $L$ . It follows that  $L_v \neq 1$  and since  $L \trianglelefteq X$ , we have  $5 \mid |L_v|$  and so  $L$  is symmetric.



If  $t = 1$ , the Frattini argument implies that  $L = GL_v$ , and by Proposition 2.8,  $L_v = A_n$  with  $n \geq 9$ , which is impossible. Thus,  $2 \leq t \leq 4$ . Since  $(|S_1 : I_1| \cdots |S_t : I_t|) \mid 2^9 \cdot 3^2 \cdot 5$ , there is at least one  $|S_i : I_i|$  that is a divisor of  $2^9 \cdot 3^2$ , say  $|S_1 : I_1| \mid 2^9 \cdot 3^2$ . Since  $I_1 = T^{r_1}$ , Proposition 2.6 implies  $I_1 = T$ , and since  $I_1 \cong G/K_1$ , we have  $K_1 = T^{l-1}$ . Set  $\bar{S}_1 = S_2 \times \cdots \times S_t$ . Then  $L = S_1 \times \bar{S}_1$  and  $K_1 = G \cap \bar{S}_1$  as  $K_1 \leq \bar{S}_1$  and  $G = T^l \neq K_1$ .

If  $\bar{S}_1$  is transitive on  $V(\Gamma)$ , then  $|L_v|/|(\bar{S}_1)_v| = |L|/|\bar{S}_1| = |S_1| \mid 2^9 \cdot 3^2 \cdot 5$  and so  $S_1 = A_5$  or  $A_6$ , forcing  $I_1 = T = A_5$ , a contradiction. If  $\bar{S}_1$  has two orbits, then  $\Gamma$  is bipartite, which is also impossible. Thus,  $\bar{S}_1$  has at least three orbits on  $V(\Gamma)$  and Proposition 2.9 implies that  $\Gamma_{\bar{S}_1}$  is a connected pentavalent  $G\bar{S}_1/\bar{S}_1$ -vertex-transitive and  $L/\bar{S}_1$ -symmetric graph. Since  $G\bar{S}_1/\bar{S}_1 \cong G/G \cap \bar{S}_1 = T$ , by Assumption,  $T \cong G\bar{S}_1/\bar{S}_1 \trianglelefteq L/\bar{S}_1 \cong S_1$ , which is impossible because  $S_1$  is a non-abelian simple group. This completes the proof.  $\square$

**Remark:** Let

$$\begin{aligned} \Delta &= \{ \text{PSL}(2, 8), \Omega_8^-(2), A_{n-1} \mid n \geq 6, n \mid 2^9 \cdot 3^2 \cdot 5 \} \\ \Delta_1 &= \{ \text{PSL}(2, 8), \Omega_8^-(2), A_{n-1} \mid n = 2 \cdot 3, 2^3 \cdot 3^2, 2^2 \cdot 3 \cdot 2 \cdot 5, 2^2 \cdot 5, 2^3 \cdot 3, \\ &\quad 2^3 \cdot 5, 2 \cdot 3 \cdot 5, 2^5 \cdot 3^2 \cdot 5, 2^7 \cdot 3 \cdot 5, 2^6 \cdot 3^2 \cdot 5, 2^7 \cdot 3^2 \cdot 5, 2^9 \cdot 3^2 \cdot 5 \}. \end{aligned}$$

Then Lemma 3.2 is true if Assumption is replaced by the conditions that  $l \geq 2$  and  $T = A_{n-1} \in \Delta - \Delta_1$ . To prove it, we only need to change the last paragraph in the proof of Lemma 3.2 as following.

Since  $|S_1 : I_1| \mid 2^9 \cdot 3^2$ , Proposition 2.6 implies that  $I_1 = T$ , and since  $T \in \Delta - \Delta_1$ , we have  $(S_1, I_1) = (A_n, A_{n-1})$  with  $n = 2^i, 2^j \cdot 3$  or  $2^k \cdot 3^2$ , where  $4 \leq i, j \leq 9$  and  $1 \leq k \leq 9$ . Recall that  $(|S_1 : I_1| \cdots |S_t : I_t|) \mid 2^9 \cdot 3^2 \cdot 5$  and  $2 \leq t \leq 4$ . Then for each  $2 \leq i \leq t$ , we have  $|S_i : I_i| \mid 2^5 \cdot 3^2 \cdot 5$  or  $2^8 \cdot 5$  and hence  $S_i$  is a primitive permutation group of degree dividing 1440 or 1280. By [15, Section 7],  $(S_i : I_i) = (A_n, A_{n-1})$ , which implies that  $t = 2$  and  $(S_2, S_1, T) = (A_n, A_n, A_{n-1})$  with  $n = 2^4$  or  $2^4 \cdot 3$ . Thus,  $G = T^2 < L = S_1 \times S_2$ ,  $G \cap S_2 = T$  (note that  $\bar{S}_1 = S_2$ ),  $|L_v|/|G_v| = |L|/|G| = n^2$  and  $2^8 \mid |L_v|$ . By Proposition 2.1,  $|L_v| = 2^9 \cdot 3^2 \cdot 5$  and so  $|G_v| = 2 \cdot 3^2 \cdot 5$  or  $2 \cdot 5$ . The former is impossible by Proposition 2.1. For the latter,  $|(GS_2)_v|/|G_v| = |GS_2|/|G| = |S_2|/|S_2 \cap G| = |S_2|/|T| = 2^4 \cdot 3$  and  $|(GS_2)_v| = 2^5 \cdot 3 \cdot 5$ , which is also impossible by Proposition 2.1.

**Lemma 3.3** Assume  $5 \mid |\text{rad}(A)|$ . Then  $\text{rad}(A)G = \text{rad}(A) \times G$ .

**Proof:** Since  $\text{rad}(A)$  is solvable,  $\text{rad}(A) \cap G = 1$ . If  $l = 1$  then by Assumption,  $G \trianglelefteq A$ , and hence  $\text{rad}(A)G = \text{rad}(A) \times G$ . Thus, we may assume  $l \geq 2$ .

Set  $B = \text{rad}(A)G$ . Then  $|B| = |\text{rad}(A)||G|$ , and by Proposition 2.3,  $|\text{rad}(A)| = |B|/|G| = |B_v|/|G_v| \mid 2^9 \cdot 3^2 \cdot 5$ . Since  $5 \mid |\text{rad}(A)|$ , we have  $5 \mid |B_v|$  and  $5 \nmid |G_v|$ , that is,  $\Gamma$  is  $B$ -symmetric, but not  $G$ -symmetric. In particular,  $G_v$  is a  $\{2, 3\}$ -group and  $5^2 \nmid |\text{rad}(A)|$ .

Since  $\text{rad}(A)$  is a solvable  $\{2, 3, 5\}$ -group,  $\text{rad}(A)$  has a Hall  $\{2, 3\}$ -subgroup, say  $H$ . Set  $\Omega = \{H^r \mid r \in \text{rad}(A)\}$ . By [9], all Hall  $\{2, 3\}$ -subgroups of  $\text{rad}(A)$  are conjugate and so the conjugate action of  $B$  on  $\Omega$  is transitive. Let  $K$  be the kernel of the action of  $G$  on  $\Omega$ . Since  $5^2 \nmid |\text{rad}(A)|$ , we have  $|\Omega| = 1$  or  $5$ , and hence  $G/K = T^r \leq S_5$ . Since  $T \neq A_5$ , we have  $G = K$  and so  $G$  fixes  $H$ . It follows  $H \trianglelefteq GH \leq B$  and  $G \cap H = 1$ .

Set  $Y = GH$  and  $\Delta = \{Yb \mid b \in B\}$ . Then  $|\Delta| = |B : Y| = 5$ . Let  $Y_B$  be the kernel of the right multiplication action of  $B$  on  $\Delta$ . Then  $Y_B$  is the largest normal subgroup of  $B$

contained in  $Y$ . It follows  $B/Y_B \leq S_5$ . Suppose  $G \not\leq Y_B$ . Then  $G/G \cap Y_B \cong GY_B/Y_B \leq B/Y_B \leq S_5$ , and so  $G/G \cap Y_B = T = A_5$ , a contradiction. Thus,  $G \leq Y_B$ .

Since  $|Y_v|/|G_v| = |Y|/|G| = |H|$  and  $G_v$  is a  $\{2, 3\}$ -group,  $Y_v$  is a  $\{2, 3\}$ -group. Since  $\Gamma$  is  $B$ -symmetric,  $B_v$  is primitive on the set of the 5 neighbors of  $v$  in  $\Gamma$ , and since  $Y_B \trianglelefteq B$ , we have either  $(Y_B)_v = 1$  or  $5 \mid |(Y_B)_v|$ . The latter cannot happen as  $Y_v$  is a  $\{2, 3\}$ -group. Thus,  $(Y_B)_v = 1$ , and the Frattini argument implies  $Y_B = G(Y_B)_v = G$ . Thus,  $G \trianglelefteq B$  and  $B = \text{rad}(A) \times G$ .  $\square$

**Lemma 3.4** *Let  $G \leq X \leq A$  and let  $\Gamma$  be  $G$ -symmetric. Then  $\text{rad}(X)G = \text{rad}(X) \times G$ .*

**Proof:** The lemma is true for  $|\text{rad}(X)| = 1$ . In what follows we assume that  $|\text{rad}(X)| \neq 1$ . Recall that  $G = T^l$  with  $l \geq 1$ . Since  $\Gamma$  is  $G$ -symmetric,  $5 \mid |G_v|$  and hence  $5 \mid |T|$ . Set  $B = \text{rad}(X)G$ . Then  $G \cap \text{rad}(X) = 1$  as  $\text{rad}(X)$  is solvable, and so  $|B| = |\text{rad}(X)||G|$ . By Proposition 2.3,  $|\text{rad}(X)| = |B_v|/|G_v| \mid 2^9 \cdot 3^2$ .

Let  $N$  be a minimal normal subgroup of  $X$  contained in  $\text{rad}(X)$ . Then  $N \cong \mathbb{Z}_2^s$  or  $N \cong \mathbb{Z}_3^t$  for some  $1 \leq s \leq 9$  or  $1 \leq t \leq 2$ , and hence  $\text{Aut}(N) \leq \text{GL}(s, 2)$  or  $\text{GL}(t, 3)$ .

First we claim  $D := GN = G \times N$ . Consider the conjugate action of  $G$  on  $N$  and let  $K$  be the kernel of  $G$  in this action. Then  $K = T^r \trianglelefteq G$  for some  $r \leq l$ , and  $G/K \leq \text{Aut}(N)$ . It is easy to see that  $D = G \times N$  if and only if  $G = K$ .

Suppose  $G \neq K$ . Then  $r < l$  and  $G/K = T^{l-r}$  is insolvable. In particular,  $\text{Aut}(N)$  is insolvable. Recall that  $\text{Aut}(N) \leq \text{GL}(s, 2)$  or  $\text{GL}(t, 3)$ . It follows that  $N \cong \mathbb{Z}_2^s$  with  $3 \leq s \leq 9$  as both  $\text{GL}(2, 2)$  and  $\text{GL}(2, 3)$  are solvable.

Note that  $G \cap N = 1$ . By Proposition 2.3,  $|N| = |D|/|G| = |D_v|/|G_v|$ , and so  $|D_v|/|G_v|$  is a 2-power. If  $G_v \cong \mathbb{Z}_5$  then Proposition 2.1 implies that  $D_v \cong D_5, D_{10}, F_{20}, F_{20} \times \mathbb{Z}_2$  or  $F_{20} \times \mathbb{Z}_4$ . It follows that  $|N| \mid 2^5$ , and this is always true by checking all other possible cases for  $G_v$  in Proposition 2.1. This means that  $N \cong \mathbb{Z}_2^s$  with  $3 \leq s \leq 5$  and hence  $G/K \leq \text{Aut}(N) \leq \text{PSL}(5, 2)$ . Since  $5^2 \nmid |\text{PSL}(5, 2)|$ , we have  $l - r = 1$  and  $G/K = T$ .

If  $K$  has at least three orbits, Proposition 2.9 implies that  $\Gamma_K$  is a connected pentavalent  $G/K$ -symmetric graph, and by Assumption,  $G/K \trianglelefteq D/K$ . It follows that  $G \trianglelefteq D$  and hence  $G = K$ , a contradiction. Thus,  $K$  has one or two orbits. If  $K$  has two orbits, then  $\Gamma$  is bipartite, a contradiction. This yields that  $K$  is transitive. Then  $|G_v|/|K_v| = |G|/|K| = |T|$ , and since  $K \trianglelefteq G$  and  $\Gamma$  is  $G$ -symmetric, we have either  $K_v = 1$  or  $5 \mid |K_v|$ . If  $5 \mid |K_v|$ , then  $5 \nmid |T|$  because  $|G_v|/|K_v| = |T|$ , a contradiction. It follows  $K_v = 1$ . Let  $L = KN$ . Then  $L \trianglelefteq D$  and so  $L_v = 1$  or  $5 \mid |L_v|$ . On the other hand,  $|L_v| = |L_v|/|K_v| = |L|/|K| = |N| = 2^s$  ( $3 \leq s \leq 5$ ), a contradiction. Therefore,  $D = G \times N$ , as claimed.

Now we finish the proof by induction on  $|\text{rad}(X)|$ . Since  $5 \nmid |N|$ , we have  $N_v = 1$ . Assume that  $N$  has one or two orbits on  $V(\Gamma)$ . Then  $5 \nmid |V(\Gamma)|$ , and so  $5^2 \nmid |G|$ , which implies that  $G$  is a simple group. By Assumption,  $G \trianglelefteq X$  and hence  $\text{rad}(X)G = G \times \text{rad}(X)$ . Assume that  $N$  has at least three orbits. By Proposition 2.9,  $\Gamma_N$  is a connected pentavalent  $GN/N$ -symmetric graph. Note that  $GN/N \cong T^l$  and  $GN/N \leq X/N$ . Since  $\text{rad}(X/N) = \text{rad}(X)/N$ , we have  $|\text{rad}(X/N)| < |\text{rad}(X)|$  and the inductive hypothesis implies that  $\text{rad}(X)G/N = \text{rad}(X/N) \cdot GN/N = \text{rad}(X)/N \times GN/N$ . Thus,  $GN \trianglelefteq \text{rad}(X)G$ . Since  $GN = G \times N$ ,  $G$  is characteristic in  $GN$  and hence  $G$  is normal in  $\text{rad}(X)G$ . It follows  $\text{rad}(X)G = \text{rad}(X) \times G$ .  $\square$



Now, we are ready to prove Theorem 1.1.

**The proof of Theorem 1.1:** Recall that  $G = T^l$  and  $\Gamma$  is a connected pentavalent symmetric  $G$ -vertex-transitive graph with  $A = \text{Aut}(\Gamma)$ . We apply induction on  $l$ . If  $l = 1$ , Theorem 1.1 is true by Assumption. Assume  $l \geq 2$ . Let  $R$  be the radical of  $A$ , and set  $B = RG$ . Then  $G \cap R = 1$  and  $|R| = |B|/|G| = |B_v|/|G_v|$ .

If  $R$  is transitive on  $V(\Gamma)$ , Proposition 2.3 implies  $|G| = |GR|/|R| = |(GR)_v|/|R_v| \mid 2^9 \cdot 3^2 \cdot 5$ , which is impossible because  $l \geq 2$ . Furthermore,  $R$  cannot have two orbits as  $\Gamma$  is not bipartite. Thus,  $R$  has at least three orbits. By Proposition 2.9,  $\Gamma_R$  is a connected pentavalent  $B/R \cong G = T^l$ -vertex-transitive and  $A/R$ -symmetric graph. Since  $R$  is the largest solvable normal subgroup of  $A$ , we have  $\text{rad}(A/R) = 1$ . By Lemmas 3.2 and 3.1,  $A/R$  has a minimal normal subgroup  $M/R = T^r$  contained in  $B/R$  with  $1 \leq r \leq l$ . It follows that  $R \leq M \leq B$  and  $M \trianglelefteq A$ . In particular,  $R \neq M$ .

Assume  $B = R \times G$ . Then  $M = M \cap RG = R(M \cap G) = R \times (M \cap G)$  and  $M \cap G$  is characteristic in  $M$ . It follows  $M \cap G \trianglelefteq A$  and  $|M \cap G| = |T|^r$ . If  $M \cap G$  is transitive, then  $|G_v|/|(M \cap G)_v| = |G|/|M \cap G| = |T|^{l-r} \mid 2^9 \cdot 3^2 \cdot 5$ . It follows  $l - r = 0$  or  $l - r = 1$ . The former implies  $G = M \cap G \trianglelefteq A$  and we are done. For the latter,  $T = A_6$  as  $T \neq A_5$  and hence  $5 \mid |G_v|$  and  $5 \nmid |(M \cap G)_v|$ , that is,  $\Gamma$  is  $G$ -symmetric, but not  $(M \cap G)$ -symmetric. Since  $M \cap G \trianglelefteq G$ , we have  $(M \cap G)_v = 1$  and  $|G_v| = |T| = 2^3 \cdot 3^2 \cdot 5$ , which is impossible by Proposition 2.1. If  $M \cap G$  has two orbits on  $V(\Gamma)$ , then  $\Gamma$  is bipartite, which is also impossible. Thus,  $M \cap G$  has at least three orbits on  $V(\Gamma)$  and Proposition 2.9 implies that  $\Gamma_{M \cap G}$  is a connected pentavalent  $G/(M \cap G)$ -vertex-transitive and  $A/(M \cap G)$ -symmetric graph. By inductive hypothesis,  $G/(M \cap G) \trianglelefteq A/(M \cap G)$  and so  $G \trianglelefteq A$ .

By the above paragraph, to finish the proof we only need to show  $B = R \times G$ , and to do this, we now claim  $B \trianglelefteq A$ . If  $M$  is transitive, then  $|B_v|/|M_v| = |B|/|M| = |B/R|/|M/R| = |T|^{l-r} \mid 2^9 \cdot 3^2 \cdot 5$  and so  $l - r = 0$  or  $l - r = 1$ . For the former,  $B = M \trianglelefteq A$ , as claimed. For the latter,  $T = A_6$  as  $T \neq A_5$ , and so  $|B_v|/|M_v| = |A_6|$ . It follows  $M_v = 1$  and  $|B_v| = |T| = 2^3 \cdot 3^2 \cdot 5$ , which is impossible by Proposition 2.1. If  $M$  has two orbits, then  $\Gamma$  is bipartite, which is also impossible. Thus,  $M$  has at least three orbits, and by Proposition 2.9,  $\Gamma_M$  is a connected pentavalent  $B/M$ -vertex-transitive and  $A/M$ -symmetric graph. Since  $B/M \cong (B/R)/(M/R) = T^{l-r}$  with  $l - r < l$ , the inductive hypothesis implies that  $B/M \trianglelefteq A/M$  and so  $B \trianglelefteq A$ , as claimed.

Since  $G \leq B \trianglelefteq A$  and  $A$  is symmetric, we have  $B_v = 1$  or  $5 \mid |B_v|$ . If  $B_v = 1$ , the Frattini argument implies  $B = GB_v = G$ , and hence  $R = 1$  and  $B = R \times G$ . If  $5 \mid |B_v|$ , then  $5 \mid |R|$  or  $5 \mid |G_v|$  as  $|B_v| = |G_v||R|$ . By Lemmas 3.3 and 3.4,  $B = R \times G$ .  $\square$

**The proof of Corollary 1.2:** This follows from Theorem 1.1 and [2, Theorem 1.1].  $\square$

**The proof of Corollary 1.3:** It is easy to check that Lemmas 3.1-3.4 are true if Assumption is replaced by the following assumption:

**Assumption 1:** Every connected pentavalent  $T$ -symmetric graph is  $T$ -normal.

Then a similar proof to the proof of Theorem 1.1 implies that under Assumption 1, every connected pentavalent  $G$ -symmetric graph is  $G$ -normal, and by [2, Corollary 1.2], we have Corollary 1.3.  $\square$

**The proof of Corollary 1.4:** By assumption of Corollary 1.4,  $\Gamma$  is a connected pentavalent symmetric  $G$ -regular graph with  $G = T^l$ , and hence it is  $G$ -vertex-transitive. Recall that

$$\begin{aligned}\Delta &= \{\text{PSL}(2, 8), \Omega_8^-(2), A_{n-1} \mid n \geq 6, n \mid 2^9 \cdot 3^2 \cdot 5\}, \\ \Delta_1 &= \{\text{PSL}(2, 8), \Omega_8^-(2), A_{n-1} \mid n = 2 \cdot 3, 2^3, 3^2, 2^2 \cdot 3, 2 \cdot 5, 2^2 \cdot 5, 2^3 \cdot 3, \\ &\quad 2^3 \cdot 5, 2 \cdot 3 \cdot 5, 2^5 \cdot 3^2 \cdot 5, 2^7 \cdot 3 \cdot 5, 2^6 \cdot 3^2 \cdot 5, 2^7 \cdot 3^2 \cdot 5, 2^9 \cdot 3^2 \cdot 5\}.\end{aligned}$$

By Corollary 1.2,  $\Gamma$  is  $G$ -normal except for  $T \in \Delta$ , and to prove Corollary 1.4, it suffices to show that  $\Gamma$  is  $G$ -normal for  $T \in \Delta - \Delta_1$ . In what follows we always assume  $T \in \Delta - \Delta_1$ .

Clearly,  $T \neq A_5$ . If  $l = 1$ , Corollary 1.4 is true by [2, Corollary 1.3]. Assume  $l \geq 2$ . Then Lemmas 3.1 and 3.2 are true by the Remarks of these two lemmas. Then a similar argument to the proof of Theorem 1.1 (the first three paragraphs) implies that  $\Gamma$  is  $G$ -normal if  $RG = R \times G$ , where  $R$  is the radical of  $A = \text{Aut}(\Gamma)$ . Set  $B = RG$ . To finish the proof, we only need to show  $B = R \times G$ .

Since  $R$  is solvable,  $G \cap R = 1$  and  $|B| = |G||R|$ , and by Proposition 2.3,  $|R| = |R|/|R \cap G| \mid 2^9 \cdot 3^2 \cdot 5$ . We may write  $|R| = 2^m \cdot 3^h \cdot 5^k$ , where  $0 \leq m \leq 9$ ,  $0 \leq h \leq 2$  and  $0 \leq k \leq 1$ . Since  $R$  is solvable, there exists a series of principle subgroups of  $B$ :

$$B > R = R_s > \cdots R_1 > R_0 = 1$$

such that  $R_i \trianglelefteq B$  and  $R_{i+1}/R_i$  is an elementary abelian  $r$ -group with  $0 \leq i \leq s-1$ , where  $r = 2, 3$  or  $5$ . Clearly,  $G$  has a natural action on  $B_{i+1}/B_i$  by conjugation.

Suppose to the contrary that  $B \neq R \times G$ . Then there exists some  $0 \leq j \leq s-1$  such that  $GR_j = G \times R_j$ , but  $GR_{j+1} \neq G \times R_{j+1}$ . If  $G$  acts trivially on  $R_{j+1}/R_j$  by conjugation, then  $[GR_j/R_j, R_{j+1}/R_j] = 1$ . Furthermore, since  $GR_j/R_j \cong G = T^l$ , we have  $(GR_j/R_j) \cap (R_{j+1}/R_j) = 1$ , and since  $|GR_{j+1}/R_j| = |GR_{j+1}/R_{j+1}||R_{j+1}/R_j| = |G||R_{j+1}/R_j| = |GR_j/R_j||R_{j+1}/R_j|$ , we have  $GR_{j+1}/R_j = GR_j/R_j \times R_{j+1}/R_j$ . In particular,  $GR_j \trianglelefteq GR_{j+1}$  and so  $G \trianglelefteq GR_{j+1}$  because  $GR_j = G \times R_j$  implies that  $G$  is characteristic in  $GR_j$ . It follows that  $GR_{j+1} = G \times R_{j+1}$ , a contradiction. Thus,  $G$  acts non-trivially on  $R_{j+1}/R_j$ . Let  $K$  be the kernel of this action. Then  $G/K = T^{l'}$  with  $l' \geq 1$ .

Since  $|R| = 2^m \cdot 3^h \cdot 5^k$  with  $0 \leq m \leq 9$ ,  $0 \leq h \leq 2$  and  $0 \leq k \leq 1$ ,  $R_{j+1}/R_j$  is an elementary abelian group of order  $r^\ell$ , where  $\ell \leq m \leq 9$  for  $r = 2$ ,  $\ell \leq 2$  for  $r = 3$  and  $\ell \leq 1$  for  $r = 5$ . Then  $G/K \leq \text{GL}(\ell, r)$ . Since  $\text{GL}(2, 3)$  and  $\text{GL}(1, 5)$  are solvable, we have  $G/K \leq \text{GL}(9, 2)$ , and since  $T \in \Delta - \Delta_1$ , we have  $11 \mid |T|$  and hence  $11 \mid |G/K|$ . It follows  $11 \mid |\text{GL}(9, 2)|$ , a contradiction. This implies  $B = R \times G$ , as required.  $\square$

**Acknowledgements:** This work was supported by the National Natural Science Foundation of China (11571035) and by the 111 Project of China (B16002).

## References

- [1] R. W. Baddeley and C. E. Praeger, On primitive overgroups of quasiprimitive permutation groups, J. Alg. 263 (2003) 294-344.

- [2] J.-L. Du, Y.-Q. Feng, J.-X. Zhou, Pentavalent symmetric graphs admitting vertex-transitive non-abelian simple groups, *Europ. J. Combin.* 63 (2017) 132-145.
- [3] X.G. Fang, L.J. Jia, J. Wang, On the automorphism groups of symmetric graphs admitting an almost simple group, *Europ. J. Combin.* 29 (2008) 1467-1472.
- [4] X.G. Fang, C.E. Praeger, Finite two-arc transitive graphs admitting a Suzuki simple group, *J. Algebraic Combin.* 27 (1999) 3727-3754.
- [5] X.G. Fang and C.E. Praeger, Finite two-arc-transitive graphs admitting a Ree simple group, *Comm. Algebra* 27 (1999) 3755-3769.
- [6] D. Gorenstein, *Finite simple groups* (Plenum Press, New York, 1982).
- [7] S.-T. Guo, Y.-Q. Feng, A note on pentavalent  $s$ -transitive graphs, *Discrete Math.* 312 (2012) 2214-2216.
- [8] R.M. Guralnick, Subgroups of prime power index in a simple group, *J Algebra.* 81 (1983) 304-311.
- [9] P. Hall, A note on soluble groups, *J. London Math. Soc.* 3 (1928) 98-105.
- [10] X.-H. Hua, Y.-Q. Feng, Cubic graphs admitting transitive non-abelian characterically simple groups, *Proceedings of the Edinburgh Mathematical Soc.* 54 (2011) 113-123.
- [11] B. Huppert, W. Lempken, Simple groups of order divisible by at most four primes, *Proc. of the F. Scorina Gemel State University* 16 (2000) 64-75.
- [12] C.H. Li, *Isomorphisms of finite Cayley graphs*, Ph.D. Thesis, The University of Western Australia, 1996.
- [13] X.H. Li, M.Y. Xu, The primitive permutation groups of degree  $2^a \cdot 3^b$ , *Arch Math.* 86 (2006) 385-391.
- [14] P. Lorimer, Vertex-transitive graphs: symmetric graphs of prime valency, *J. Graph Theory* 8 (1984) 55-68.
- [15] C. M. Roney-Dougal, The primitive permutation groups of degree less than 2500, *J. Alg.* 292 (2005) 154-183.
- [16] S.J. Xu, X.G. Fang, J. Wang, M.Y. Xu, On cubic  $s$ -arc transitive Cayley graphs of finite simple groups, *Europ. J. Combin.* 26 (2005) 133-143.
- [17] S.J. Xu, X.G. Fang, J. Wang, M.Y. Xu, 5-arc transitive cubic Cayley graphs on finite simple groups, *Europ. J. Combin.* 28 (2007) 1023-1036.