

The distinguishing number and the distinguishing index of line and graphoidal graphs

Saeid Alikhani Samaneh Soltani *

March 7, 2024

Department of Mathematics, Yazd University, 89195-741, Yazd, Iran
alikhani@yazd.ac.ir, s.soltani1979@gmail.com

Abstract

The distinguishing number (index) $D(G)$ ($D'(G)$) of a graph G is the least integer d such that G has an vertex labeling (edge labeling) with d labels that is preserved only by a trivial automorphism. A graphoidal cover of G is a collection ψ of (not necessarily open) paths in G such that every path in ψ has at least two vertices, every vertex of G is an internal vertex of at most one path in ψ and every edge of G is in exactly one path in ψ . Let $\Omega(G, \psi)$ denote the intersection graph of ψ . A graph H is called a graphoidal graph, if there exists a graph G and a graphoidal cover ψ of G such that $H \cong \Omega(G, \psi)$. In this paper, we study the distinguishing number and the distinguishing index of the line graph and the graphoidal graph of a simple connected graph G .

Keywords: distinguishing number; distinguishing index; line and graphoidal graph

AMS Subj. Class.: 05C25

1 Introduction and definitions

Let $G = (V, E)$ be a simple, finite, connected and undirected graph, and let $\text{Aut}(G)$ be its *automorphism group*. A labeling of G , $\phi : V \rightarrow \{1, 2, \dots, r\}$, is said to be *r-distinguishing*, if no non-trivial automorphism of G preserves all of the vertex labels. The point of the labels on the vertices is to destroy the symmetries of the graph, that is, to make the automorphism group of the labeled graph trivial. Formally, ϕ is *r-distinguishing* if for every non-identity $\sigma \in \text{Aut}(G)$, there exists x in V such that $\phi(x) \neq \phi(\sigma(x))$. The *distinguishing number* of a graph G is defined by

$$D(G) = \min\{r \mid G \text{ has a labeling that is } r\text{-distinguishing}\}.$$

This number has defined in [1]. Similar to this definition, in [7] the *distinguishing index* $D'(G)$ of G , which is the least integer d such that G has an edge colouring with

*Corresponding author

d colours that is preserved only by a trivial automorphism, has been defined. $D'(G)$ can be arbitrary smaller than $D(G)$, for example $D'(K_{p,p}) = 2$ and $D(K_{p,p}) = p + 1$, for $p \geq 4$.

The *line graph* $L(G)$ of a graph G is the graph whose vertices are edges of G and two edges $e, e' \in V(L(G)) = E(G)$ are adjacent if they share an endpoint in common. The line graphs can be viewed a special case of graphoidal graphs. The concept of graphoidal cover was introduced by Acharya and Sampathkumar [2].

Definition 1.1 [2] *A graphoidal cover of a graph G is a collection ψ of (not necessarily open) paths in G satisfying the following conditions:*

- (i) *Every path in ψ has at least two vertices.*
- (ii) *Every vertex of G is an internal vertex of at most one path in ψ .*
- (iii) *Every edge of G is in exactly one path in ψ .*

Definition 1.2 [2] *If $F = \{S_1, S_2, \dots, S_n\}$ is a family of distinct nonempty subsets of a set S , whose union is S , then the intersection graph of F , denoted by $\Omega(F)$, is the graph whose vertex set and edge set are given by*

$$V(F) = \{S_1, S_2, \dots, S_n\}, \quad E(F) = \{S_i S_j : i \neq j, S_i \cap S_j \neq \emptyset\}.$$

Let \mathcal{G}_G be the set of all graphoidal covers of G and $\psi \in \mathcal{G}_G$. The intersection graph on ψ is denoted by $\Omega(G, \psi)$. A graph H is called a *graphoidal graph* if there exists a graph G and $\psi \in \mathcal{G}_G$ such that $H \cong \Omega(G, \psi)$. Since $E(G)$ is obviously a graphoidal cover of G , all line graphs are graphoidal graphs. In [2] Acharya and Sampathkumar have proved that all the Beineke's forbidden subgraphs of line graphs (see Figure 1) are graphoidal graphs and hence they conjectured that all graphs are graphoidal. In [4] Arumugam and Pakkiam disproved this conjecture by showing that complete bipartite graphs $K_{m,n}$ with $mn > 2(m + n)$ are not graphoidal graphs. They also obtained a forbidden subgraph characterization of all bipartite graphs which are graphoidal. Note that if $P = (v_0, v_1, \dots, v_k)$ is a path, not necessarily open, in a graph G , then v_0 and v_k are called *terminal vertices* and v_1, \dots, v_{k-1} are called *internal vertices* of P . For cycles (considered as closed paths), there is an inherent ordering of vertices as in paths. So, when we say that a cycle C of a graph G is a member of a graphoidal cover ψ of G , we should mention the vertex at which the cycle C begins, and this particular vertex is considered as the terminal vertex of C and all other vertices on C are called internal vertices of C . Given a graphoidal cover ψ of G , a vertex v is said to be interior to ψ if v is an internal vertex of an element of ψ and is called exterior to ψ otherwise.

In the next section, we study the distinguishing number and the distinguishing index of line graphs. In Section 3, we consider graphoidal graphs and study their distinguishing number and distinguishing index.




Figure 1: The nine forbidden subgraph characterization of line graphs.

2 Study of $D(G)$ and $D'(G)$ for the line graphs

For a simple graph G , the line graph $L(G)$ is the graph whose vertices are edges of G and two edges $e, e' \in V(L(G)) = E(G)$ are adjacent if they share an endpoint in common. The following theorem characterize the line graphs.

Theorem 2.1 [5] *The following statements are equivalent for a graph G .*

- (i) *The graph G is the line graph of some graphs.*
- (ii) *The edges of G can be partitioned into complete subgraphs in such a way that no vertex belongs to more than two of the subgraphs.*
- (iii) *The graph $K_{1,3}$ is not an induced subgraph of G ; and if abc and bcd are distinct odd triangles, then a and d are adjacent.*
- (iv) *None of the nine graphs in Figure 1 is an induced subgraph of G .*

To study the distinguishing number and the distinguishing index of $L(G)$, we need more information about the automorphism group of $L(G)$. Let $\gamma_G : \text{Aut}(G) \rightarrow \text{Aut}(L(G))$ be given by $(\gamma_G \phi)(\{u, v\}) = \{\phi(u), \phi(v)\}$ for every $\{u, v\} \in E(G)$. In [11], Sabidussi proved the following Theorem which we need it later.

Theorem 2.2 [11] *Suppose that G is a connected graph that is not P_2 , Q , or $L(Q)$ (see Figure 2). Then G is a group isomorphism, and so $\text{Aut}(G) \cong \text{Aut}(L(G))$.*

Now we are ready to obtain the distinguishing number of line graph of a graph. We note that if $G = L(Q)$, then it is easy to see that $D'(L(Q)) = 2$, while $D(L(L(Q))) = 3$.

Theorem 2.3 *Suppose that G is a connected graph that is not P_2 and $L(Q)$. Then $D(L(G)) = D'(G)$.*



Figure 2: graphs Q and $L(Q)$ of Theorem 2.2.

Proof. If $G = Q$, then it is easy to see that $D'(Q) = D(L(Q)) = 2$. If $G \neq Q$, First we show that $D(L(G)) \leq D'(G)$. Let $c : E(G) \rightarrow \{1, \dots, D'(G)\}$ be an edge distinguishing labeling of G . We define $c' : V(L(G)) \rightarrow \{1, \dots, D'(G)\}$ such that $c'(e) = c(e)$, where $e \in V(L(G)) = E(G)$. The vertex labeling c' is a distinguishing vertex labeling of $L(G)$, because if f is an automorphism of $L(G)$ preserving the labeling, then $c'(f(e)) = c'(e)$, and hence $c(f(e)) = c(e)$ for any $e \in E(G)$. On the other hand, by Theorem 2.2, $f = \gamma_G \phi$ for some automorphism ϕ of G . Thus from $c(f(e)) = c(e)$ for any $e \in E(G)$, we can conclude that $c(\gamma_G \phi(e)) = c(e)$ and so $c(\{\phi(u), \phi(v)\}) = c(\{u, v\})$ for every $\{u, v\} \in E(G)$. This means that ϕ is an automorphism of G preserving the labeling c , and so ϕ is the identity automorphism of G . Therefore f is the identity automorphism of $L(G)$, and hence $D(L(G)) \leq D'(G)$. For the converse, suppose that $c : V(L(G)) \rightarrow \{1, \dots, D(L(G))\}$ is a vertex distinguishing labeling of $L(G)$. We define $c' : E(G) \rightarrow \{1, \dots, D(L(G))\}$ such that $c'(e) = c(e)$ where $e \in E(G)$. The edge labeling c' is a distinguishing edge labeling of G . Because if f is an automorphism of G preserving the labeling, then $c'(f(e)) = c'(e)$, and hence $c(f(e)) = c(e)$ for any $e \in E(G)$. Then, there exist the automorphism $\gamma_G f$ of $L(G)$ such that $\gamma_G f(\{u, v\}) = \{f(u), f(v)\}$ for every $\{u, v\} \in E(G)$, by Theorem 2.2. Thus from $c(f(e)) = c(e)$ for any $e \in E(G)$, we can conclude that $c(\{u, v\}) = c(\{f(u), f(v)\}) = c(\gamma_G f(\{u, v\}))$ for every $\{u, v\} \in E(G)$, which means that $\gamma_G f$ preserves the distinguishing vertex labeling of $L(G)$, and hence $\gamma_G f$ is the identity automorphism of $L(G)$. Therefore f is the identity automorphism of G , and so $D'(G) \leq D(L(G))$. \square

An upper bound for the distinguishing index of line graphs follows immediately from the following result of Pilśniak in [9]. A $K_{1,3}$ -free graph, called also a *claw-free graph*, is a graph containing no copy of $K_{1,3}$ as an induced subgraph.

Theorem 2.4 [9] *If G is a connected claw-free graph, then $D'(G) \leq 3$.*

Now by Part (iii) of Theorem 2.1 and Theorem 2.4, we have the following theorem.

Theorem 2.5 *If G is a connected graph, then $D'(L(G)) \leq 3$.*

3 Study of $D(G)$ and $D'(G)$ for graphoidal graphs

We recall that a graphoidal cover of a graph G is a collection ψ of (not necessarily open) paths in G satisfying the following conditions: every path in ψ has at least two




Figure 3: Graph G in the proof of Theorem 3.1.

vertices, every vertex of G is an internal vertex of at most one path in ψ , and every edge of G is in exactly one path in ψ . Let ψ be a graphoidal cover of G and $\Omega(G, \psi)$ denote the intersection graph of ψ . Thus the vertices of $\Omega(G, \psi)$ are the paths in ψ and two paths in ψ are adjacent in $\Omega(G, \psi)$ if and only if they have a common vertex. A graph G is said to be graphoidal if there exists a graph H and a graphoidal cover ψ of H such that G is isomorphic to $\Omega(H, \psi)$. First we state and prove the following theorem:

Theorem 3.1 *There exists a graph G such that the values $|D(G) - D(\Omega(G, \psi))|$ and $|D'(G) - D'(\Omega(G, \psi))|$ can be arbitrarily large.*

Proof. Let G be a graph as shown in Figure 3. It is easy to see that $D(G) = D'(G) = 2$. A graphoidal cover of G is $\psi = \{P_1, P_2, \dots, P_{n+1}\}$ where $P_1 = (v_1, v_2, \dots, v_{n+2})$ and $P_i = (v_i, w_i)$ for $2 \leq i \leq n+1$. Thus $\Omega(G, \psi) = K_{1, n}$, and hence $D(\Omega(G, \psi)) = D'(\Omega(G, \psi)) = n$. Therefore we have the result. \square

To study the distinguishing index of graphoidal graphs, we need some theorems.

Theorem 3.2 [7] *If G is a connected graph of order $n \geq 3$, then $D'(G) \leq \Delta(G)$, unless G is C_3, C_4 or C_5 .*

Theorem 3.3 [9] *Let G be a connected graph that is neither a symmetric nor a bisymmetric tree. If the maximum degree of G is at least 3, then $D'(G) \leq \Delta(G) - 1$, unless G is K_4 or $K_{3,3}$.*

Theorem 3.4 [6] *If T is a tree of order $n \geq 3$, then $D(T) \leq \Delta(T)$. Furthermore, equality is achieved if and only if T is a symmetric tree or a path of odd length.*

Before we state the next theorem we need to define some terms: Let v be a vertex of a graph G and let f be a k -distinguishing edge labeling of G . We say that f is v -distinguishing if f is preserved only by the trivial automorphism among the automorphisms $\alpha \in \text{Aut}(G)$ for which $\alpha(v) = v$ holds. If vw is an edge of a tree T , then let T_v and T_w be the components of $T - \{vw\}$ with $v \in T_v$ and $w \in T_w$. A family \mathcal{T} consists of those trees T of order at least 3, for which the following conditions are fulfilled: (1) T is a bicentric tree with the central edge $e = vw$, (2) T_v and T_w are isomorphic trees, (3) T_v admits a unique v -distinguishing edge $D(T)$ -labeling.




Figure 4: The graph G and its graphoidal of Theorem 3.7.

Theorem 3.5 [3] *Let T be a tree of order $n \geq 3$. Then $D'(T) = D(T) + 1$ if T belongs to \mathcal{T} , and $D'(T) = D(T)$ for all other trees.*

Now, we obtain bounds for the distinguishing index of graphoidal graphs.

Theorem 3.6 *Let G be a connected graph and ψ be a graphoidal cover of G such that the order of $\Omega(G, \psi)$ is at least 3. If $\Omega(G, \psi) \neq C_3, C_5$, then $1 \leq D'(\Omega(G, \psi)) \leq |\psi| - 1$. Moreover, $D'(\Omega(G, \psi)) = |\psi| - 1$ if and only if $\Omega(G, \psi)$ is C_4 , K_4 or $K_{1,|\psi|-1}$.*

Proof. By Figure 5, we can conclude the left inequality. For the right inequality, by contradiction, we suppose that $D'(\Omega(G, \psi)) \geq |\psi|$. Since $D'(\Omega(G, \psi)) \leq \Delta(\Omega(G, \psi))$ by Theorem 3.2, we get $|\psi| \leq \Delta(\Omega(G, \psi))$, which is a contradiction, since $\Delta(\Omega(G, \psi)) \leq |V(\Omega(G, \psi))| - 1 = |\psi| - 1$.

If $D'(\Omega(G, \psi)) = |\psi| - 1$, then $D'(\Omega(G, \psi)) = \Delta(\Omega(G, \psi))$, by Theorem 3.2, unless for $\Omega(G, \psi) = C_4$. If $\Delta(\Omega(G, \psi)) \geq 3$, then we have $D'(\Omega(G, \psi)) = \Delta(\Omega(G, \psi))$, if and only if $\Omega(G, \psi)$ is a symmetric, a bisymmetric tree, K_4 or $K_{3,3}$, by Theorem 3.3. Now since $\Delta(\Omega(G, \psi)) = |\psi| - 1$, so $D'(\Omega(G, \psi)) = |\psi| - 1$ if and only if $\Omega(G, \psi)$ is C_4 , K_4 or $K_{1,|\psi|-1}$, by Theorems 3.4 and 3.5. \square

The following theorem states that for every i , $2 \leq i \leq |\psi| - 1$, there exists a connected graphoidal graph with the distinguishing index i .

Theorem 3.7 *Suppose that x and p are positive integers and $(x, p) \neq (1, 1)$. If $k = 1 + xp$ and $d = \lceil \sqrt[2]{x} \rceil$, then there exist a graph G of order $n \geq x(p + 2)$ and graphoidal cover ψ of G with $|\psi| = k$ such that $D'(\Omega(G, \psi)) = d$.*

Proof. Let G be a graph in Figure 4. If $\psi = \{P_0, P_1^{(1)}, \dots, P_p^{(1)}, \dots, P_1^{(x)}, \dots, P_p^{(x)}\}$ where $P_0 = (v_1, \dots, v_{t_1}, w_1, \dots, w_x, z_1, \dots, z_{t_2})$ and for every $1 \leq i \leq x$, $P_1^{(i)} = (w_i, y_1^{(i)})$, $P_j^{(i)} = (y_j^{(i)}, y_{j+1}^{(i)})$ for each $2 \leq j \leq p$, then $\Omega(G, \psi)$ is as shown in Figure 4. It is easy to compute that $D'(\Omega(G, \psi)) = \lceil \sqrt[2]{x} \rceil$. With respect to integers t_1 and t_2 , we get that $n \geq x(p + 2)$. \square

In sequel, we gives bounds for the distinguishing number of the graphoidal graphs.




Figure 5: An example for sharpness of inequality of Corollary 3.9.

Theorem 3.8 *Let G be a connected graph of order $n \geq 3$. Then*

$$D'(G) - 2 \leq D(\Omega(G, \psi)) \leq |\psi|.$$

Proof. Since $\Omega(G, \psi)$ has $|\psi|$ vertices, so it is clear that $D(\Omega(G, \psi)) \leq |\psi|$. To prove the left inequality, let $D(\Omega(G, \psi)) = t$ and $X_i = \{P_{i1}, \dots, P_{is_i}\}$, $1 \leq i \leq t$ be the set of vertices of $\Omega(G, \psi)$ having label i in the distinguishing labeling of $\Omega(G, \psi)$. Now we want to label the edges of G distinguishingly, using $t + 2$ labels. For this purpose, for every i , $1 \leq i \leq t$, we label the edge set of all paths of ψ (not necessarily open) in X_i of length j with j -tuple $(i, \underbrace{t + 1, t + 2, \dots, t + 2}_{(j-2)\text{-times}})$ of labels. We note that since

there exist the closed paths (cycles) in ψ , so we need at most three different labels to distinguishing these closed paths. Then we have an edge labeling of G with $t + 2$ labels. To show that this edge labeling is distinguishing, we prove that if f is an automorphism of G preserving this edge labeling, then f maps ψ to ψ , setwise. In fact, if $P \in \psi$, then $f(P)$ is a path of the same length as P . Since f preserves the edge labeling of G , so the label of edges of $f(P)$ is the same as edges of P . With respect to the method of labeling of edge set of each path in ψ , we conclude that $f(P) \in \psi$. Hence f maps a path in ψ to a path in ψ . Thus f can be considered as an automorphism of $\Omega(G, \psi)$ preserves the distinguishing vertex labeling of $\Omega(G, \psi)$ with t labels. Thus f is the identity automorphism of $\Omega(G, \psi)$, and this means that $f(P) = P$ for all $P \in \psi$. On the other hand, since we labeled the edge set of each path in ψ distinguishingly with at most three labels, so f fixes all vertices of path P , where $P \in \psi$. Thus f is the identity automorphism of G . \square

The bounds of Theorem 3.8 are sharp. For the left inequality, it is sufficient to consider $G = C_i$, $3 \leq i \leq 5$, and $\psi = \{C_i\}$. Thus $\Omega(G, \psi) = K_1$, and hence $D'(G) = 3$ and $D(\Omega(G, \psi)) = 1$. To show that the upper bound of this theorem is sharp, we consider $G = C_3$ and $\psi = \{e_1, e_2, e_3\}$. Thus $\Omega(G, \psi) = C_3$, and hence $|\psi| = 3$ and $D(\Omega(G, \psi)) = 3$.

Theorem 3.9 *Let G be a connected graph of order $n \geq 3$. If graphoidal cover ψ of G contains only open paths, then $D'(G) \leq D(\Omega(G, \psi)) + 1$.*

Proof. The proof is exactly the same as proof of Theorem 3.8, except the method of labeling of edges of paths in ψ . Since all paths in ψ are open, then by notation of

Theorem 3.8, for every i , $1 \leq i \leq t$, we label the edge set of all open paths of ψ in X_i of length j with the j -tuple $(i, \underbrace{t+1, \dots, t+1}_{(j-1)\text{-times}})$ of labels. Then we have an edge labeling

of G with $t+1$ labels. To show this edge labeling is distinguishing, we prove that if f is an automorphism of G preserving this edge labeling, then f maps ψ to ψ setwise. In fact, if $P \in \psi$, then $f(P)$ is a path of the same length as P . Since f preserves the edge labeling of G , so the label of edges of $f(P)$ is the same as edges of P . With respect to the method of labeling of edge set of each path in ψ , we conclude that $f(P) \in \psi$. Hence f maps a path in ψ to a path in ψ . Thus f can be considered as an automorphism of $\Omega(G, \psi)$ preserves the distinguishing vertex labeling of $\Omega(G, \psi)$ with t labels. Thus f is the identity automorphism of $\Omega(G, \psi)$, and this means that $f(P) = P$ for all $P \in \psi$. On the other hand, since we labeled the edge set of each path in ψ distinguishingly with at most two labels, so f fixes all vertices of path P where $P \in \psi$. Thus f is the identity automorphism of G . \square

The bound of Theorem 3.9 is sharp. let G be as shown in Figure 5. It is easy to obtain that $D'(G) = 2$. A graphoidal cover of G is $\psi = \{P_1, P_2, \dots, P_7\}$ where $P_1 = (v_1, v_2)$, $P_2 = (v_2, v_3)$, $P_3 = (v_3, v_4)$, $P_4 = (v_4, v_5)$, $P_5 = (v_5, v_6)$, $P_6 = (v_5, v_7, v_8)$, and $P_7 = (v_8, v_9)$. Thus $\Omega(G, \psi)$ is an asymmetric graph, and hence $D(\Omega(G, \psi)) = 1$.

We conclude the paper with the following theorem, which shows that the value $|D'(G) - D(\Omega(G, \psi))|$ can be arbitrary.

Theorem 3.10 *For every i , $i \geq 0$, there exist a connected graph G and a graphoidal cover ψ of G such that $|D'(G) - D(\Omega(G, \psi))| = i$.*

Proof. Let G and $\Omega(G, \psi)$ be the two graphs of Theorem 3.7. Since the automorphism group of G has at most one nonidentity automorphism, so $D'(G) \leq 2$. On the other hand $\Omega(G, \psi)$ is a tree with a central vertex, and so $D(\Omega(G, \psi)) = D'(\Omega(G, \psi)) = \lceil \sqrt[p]{x} \rceil$, by Theorem 3.5. Now, with respect to the values of x and p , we obtain the result. \square

References

- [1] M.O. Albertson and K.L. Collins, *Symmetry breaking in graphs*, Electron. J. Combin. 3 (1996), #R18.
- [2] B. D. Acharya and E. Sampathkumar, *Graphoidal covers and graphoidal covering number of a graph*, Indian J. Pure Appl. Math., 18 (10) (1987), 882-890.
- [3] S. Alikhani, S. Klavžar, F. Lehner and S. Soltani, *Trees with distinguishing index equal distinguishing number plus one*, arXiv:1608.03501v4 [math.CO].
- [4] S. Arumugam and C. Pakkiam, *Graphoidal bipartite graphs*, Graphs Combin. 10 (1994), 305-310.
- [5] L.W. Beineke, *Characterizations of derived graphs*, J. Combin. Theory 9 (2) (1970), 129-135.

- [6] K.L. Collins, A.N. Trenk, *The distinguishing chromatic number*, Electron. J. Combin. 13 (2006), #R16.
- [7] R. Kalinowski and M. Pilśniak, *Distinguishing graphs by edge colourings*, European J. Combin. 45 (2015), 124-131.
- [8] J. Krausz, *Démonstration nouvelle d'un théorème de Whitney sur les réseaux*, Mat. Fiz. Lapok 50 (1943), 75-85.
- [9] M. Pilśniak, *Improving upper bounds for the distinguishing index*, Ars Math. Contemp. 13 (2017) 259-274.
- [10] A. V. Rooij and H. Wilf, *The interchange graph of a finite graph*, Acta Math. Acad. Sci. Hungar. 16 (1965), 263-269.
- [11] G. Sabidussi, *Graph derivatives*, Math. Z. 76 (1961), 385-401.
- [12] I. Shipman, *The distinguishing number of the iterated line graph*, arXiv preprint math/0509296 (2005).