

SPANNING SIMPLICIAL COMPLEXES OF UNI-CYCLIC MULTIGRAPHS

IMRAN AHMED, SHAHID MUHMOOD

ABSTRACT. A multigraph is a nonsimple graph which is permitted to have multiple edges, that is, edges that have the same end nodes. We introduce the concept of spanning simplicial complexes $\Delta_s(\mathcal{G})$ of multigraphs \mathcal{G} , which provides a generalization of spanning simplicial complexes of associated simple graphs. We give first the characterization of all spanning trees of a uni-cyclic multigraph $\mathcal{U}_{n,m}^r$ with n edges including r multiple edges within and outside the cycle of length m . Then, we determine the facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{U}_{n,m}^r))$ of spanning simplicial complex $\Delta_s(\mathcal{U}_{n,m}^r)$ and its primary decomposition. The Euler characteristic is a well-known topological and homotopic invariant to classify surfaces. Finally, we device a formula for Euler characteristic of spanning simplicial complex $\Delta_s(\mathcal{U}_{n,m}^r)$.

Key words: multigraph, spanning simplicial complex, Euler characteristic.
2010 Mathematics Subject Classification: Primary 05E25, 55U10, 13P10, Secondary 06A11, 13H10.

1. INTRODUCTION

Let $\mathcal{G} = \mathcal{G}(V, E)$ be a multigraph on the vertex set V and edge-set E . A spanning tree of a multigraph \mathcal{G} is a subtree of \mathcal{G} that contains every vertex of \mathcal{G} . We represent the collection of all edge-sets of the spanning trees of a multigraph \mathcal{G} by $s(\mathcal{G})$. The facets of spanning simplicial complex $\Delta_s(\mathcal{G})$ is exactly the edge set $s(\mathcal{G})$ of all possible spanning trees of a multigraph \mathcal{G} . Therefore, the spanning simplicial complex $\Delta_s(\mathcal{G})$ of a multigraph \mathcal{G} is defined by

$$\Delta_s(\mathcal{G}) = \langle F_k \mid F_k \in s(\mathcal{G}) \rangle,$$

which gives a generalization of the spanning simplicial complex $\Delta_s(G)$ of an associated simple graph G . The spanning simplicial complex of a simple connected finite graph was firstly introduced by Anwar, Raza and Kashif in [1]. Many authors discussed algebraic and combinatorial properties of spanning simplicial complexes of various classes of simple connected finite graphs, see for instance [1], [5], [6] and [9].

Let Δ be simplicial complex of dimension d . We denote f_i by the number of i -cells of simplicial complex Δ . Then, the Euler characteristic of Δ is given

by

$$\chi(\Delta) = \sum_{i=0}^{d-1} (-1)^i f_i,$$

which is a well-known topological and homotopic invariant to classify surfaces, see [4] and [7].

The uni-cyclic multigraph $\mathcal{U}_{n,m}^r$ is a connected graph having n edges including r multiple edges within and outside the cycle of length m . Our aim is to give some algebraic and topological characterizations of spanning simplicial complex $\Delta_s(\mathcal{U}_{n,m}^r)$.

In Lemma 3.1, we give characterization of all spanning trees of a uni-cyclic multigraph $\mathcal{U}_{n,m}^r$ having n edges including r multiple edges and a cycle of length m . In Proposition 3.2, we determine the facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{U}_{n,m}^r))$ of spanning simplicial complex $\Delta_s(\mathcal{U}_{n,m}^r)$ and its primary decomposition. In Theorem 3.3, we give a formula for Euler characteristic of spanning simplicial complex $\Delta_s(\mathcal{U}_{n,m}^r)$.

2. BASIC SETUP

A simplicial complex Δ on $[n] = \{1, \dots, n\}$ is a collection of subsets of $[n]$ satisfying the following properties.

- (1) $\{j\} \in \Delta$ for all $j \in [n]$;
- (2) If $F \in \Delta$ then every subset of F will belong to Δ (including empty set).

The elements of Δ are called faces of Δ and the dimension of any face $F \in \Delta$ is defined as $|F| - 1$ and is written as $\dim F$, where $|F|$ is the number of vertices of F . The vertices and edges are 0 and 1 dimensional faces of Δ (respectively), whereas, $\dim \emptyset = -1$. The maximal faces of Δ under inclusion are said to be the facets of Δ . The dimension of Δ is denoted by $\dim \Delta$ and is defined by $\dim \Delta = \max\{\dim F \mid F \in \Delta\}$.

If $\{F_1, \dots, F_q\}$ is the set of all the facets of Δ , then $\Delta = \langle F_1, \dots, F_q \rangle$. A simplicial complex Δ is said to be pure, if all its facets are of the same dimension.

A subset M of $[n]$ is said to be a vertex cover for Δ if M has non-empty intersection with every F_k . M is said to be a minimal vertex cover for Δ if no proper subset of M is a vertex cover for Δ .

Definition 2.1. Let $\mathcal{G} = \mathcal{G}(V, E)$ be a multigraph on the vertex set V and edge-set E . A spanning tree of a multigraph \mathcal{G} is a subtree of \mathcal{G} that contains every vertex of \mathcal{G} .

Definition 2.2. Let $\mathcal{G} = \mathcal{G}(V, E)$ be a multigraph on the vertex set V and edge-set E . Let $s(\mathcal{G})$ be the edge-set of all possible spanning trees of \mathcal{G} . We define a simplicial complex $\Delta_s(\mathcal{G})$ on E such that the facets of $\Delta_s(\mathcal{G})$ are exactly the elements of $s(\mathcal{G})$, we call $\Delta_s(\mathcal{G})$ as the spanning simplicial complex of \mathcal{G} and given by

$$\Delta_s(\mathcal{G}) = \langle F_k \mid F_k \in s(\mathcal{G}) \rangle.$$

Definition 2.3. A uni-cyclic multigraph $\mathcal{U}_{n,m}^r$ is a connected graph having n edges including r multiple edges within and outside the cycle of length m .

Let Δ be simplicial complex of dimension d . Then, the chain complex $C_*(\Delta)$ is given by

$$0 \rightarrow C_d(\Delta) \xrightarrow{\partial_d} C_{d-1}(\Delta) \xrightarrow{\partial_{d-1}} \dots \xrightarrow{\partial_2} C_1(\Delta) \xrightarrow{\partial_1} C_0(\Delta) \xrightarrow{\partial_0} 0.$$

Each $C_i(\Delta)$ is a free abelian group of rank f_i . The boundary homomorphism $\partial_d : C_d(\Delta) \rightarrow C_{d-1}(\Delta)$ is defined by

$$\partial_d(\sigma_\alpha^d) = \sum_{i=0}^d (-1)^i \sigma_\alpha^d|_{[v_0, v_1, \dots, \hat{v}_i, \dots, v_d]}.$$

Of course,

$$H_i(\Delta) = Z_i(\Delta) / B_i(\Delta) = \text{Ker} \partial_i / \text{Im} \partial_{i+1},$$

where $Z_i(\Delta) = \text{Ker} \partial_i$ and $B_i(\Delta) = \text{Im} \partial_{i+1}$ are the groups of simplicial i -cycles and simplicial i -boundaries, respectively. Therefore,

$$\text{rank } H_i(\Delta) = \text{rank } Z_i(\Delta) - \text{rank } B_i(\Delta).$$

One can easily see that $\text{rank } B_d(\Delta) = 0$ due to $B_d(\Delta) = 0$. For each $i \geq 0$ there is an exact sequence

$$0 \rightarrow Z_i(\Delta) \rightarrow C_i(\Delta) \xrightarrow{\partial_i} B_{i-1}(\Delta) \rightarrow 0.$$

Moreover,

$$f_i = \text{rank } C_i(\Delta) = \text{rank } Z_i(\Delta) + \text{rank } B_{i-1}(\Delta).$$

Therefore, the Euler characteristic of Δ can be expressed as

$$\begin{aligned} \chi(\Delta) &= \sum_{i=0}^d (-1)^i f_i = \sum_{i=0}^d (-1)^i (\text{rank } Z_i(\Delta) + \text{rank } B_{i-1}(\Delta)) \\ &= \sum_{i=0}^d (-1)^i \text{rank } Z_i(\Delta) + \sum_{i=0}^d (-1)^i \text{rank } B_{i-1}(\Delta). \end{aligned}$$

Changing index of summation in the last sum and using the fact that $\text{rank } B_{-1}(\Delta) = 0 = \text{rank } B_d(\Delta)$, we get

$$\begin{aligned} \chi(\Delta) &= \sum_{i=0}^d (-1)^i \text{rank } Z_i(\Delta) + \sum_{i=0}^d (-1)^{i+1} \text{rank } B_i(\Delta) \\ &= \sum_{i=0}^d (-1)^i (\text{rank } Z_i(\Delta) - \text{rank } B_i(\Delta)) = \sum_{i=0}^d (-1)^i \text{rank } H_i(\Delta). \end{aligned}$$

Thus, the Euler characteristic of Δ is given by

$$\chi(\Delta) = \sum_{i=0}^d (-1)^i \beta_i(\Delta),$$

where $\beta_i(\Delta) = \text{rank } H_i(\Delta)$ is the i -th Betti number of Δ , see [4] and [7].

3. TOPOLOGICAL CHARACTERIZATIONS OF $\Delta_s(\mathcal{U}_{n,m}^r)$

Let $\mathcal{U}_{n,m}^r$ be a uni-cyclic multigraph having n edges including r multiple edges within and outside the cycle of length m . We fix the labeling of the edge set E of $\mathcal{U}_{n,m}^r$ as follows:

$E = \{e_{11}, \dots, e_{1t_1}, \dots, e_{r'1}, \dots, e_{r't_{r'}}, e_{(r'+1)1}, \dots, e_{m1}, e_{(m+1)1}, \dots, e_{(m+1)t_{m+1}}, \dots, e_{(m+r'')1}, \dots, e_{(m+r'')t_{m+r''}}, e_1, \dots, e_v\}$, where e_{i1}, \dots, e_{it_i} are the multiple edges of i -th edge of cycle with $1 \leq i \leq r'$ while $e_{(r'+1)1}, \dots, e_{m1}$ are the single edges of the cycle and e_{j1}, \dots, e_{jt_j} are the multiple edges of the j -th edge outside the cycle with $m+1 \leq j \leq m+r''$, moreover, e_1, \dots, e_v are single edges appeared outside the cycle.

We give first the characterization of $s(\mathcal{U}_{n,m}^r)$.

Lemma 3.1. *Let $\mathcal{U}_{n,m}^r$ be the uni-cyclic multigraph having n edges including r multiple edges and a cycle of length m with the edge set E , given above. A subset $E(T_{wi_w}) \subset E$ will belong to $s(\mathcal{U}_{n,m}^r)$ if and only if $T_{wi_w} = \{e_{1i_1}, \dots, e_{r'i_{r'}}, e_{(r'+1)1}, \dots, e_{m1}, e_{(m+1)i_{(m+1)}}, \dots, e_{(m+r'')i_{(m+r'')}}, e_1, \dots, e_v\} \setminus \{e_{wi_w}\}$ for some $i_h \in \{1, \dots, t_h\}$ with $1 \leq h \leq r'$, $m+1 \leq h \leq m+r''$ and $i_w \in \{1, \dots, t_w\}$ with $1 \leq w \leq r'$ or $i_w = 1$ with $r'+1 \leq w \leq m$ for some $w = h$ and $i_w = i_h$ appeared in T_{wi_w} .*

Proof. By cutting down method [3], the spanning trees of $\mathcal{U}_{n,m}^r$ can be obtained by removing exactly $t_h - 1$ edges from each multiple edge such that $1 \leq h \leq r'$, $m+1 \leq h \leq m+r''$ and in addition, an edge from the resulting cycle need to be removed. Therefore, the spanning trees will be of the form $T_{wi_w} = \{e_{1i_1}, \dots, e_{r'i_{r'}}, e_{(r'+1)1}, \dots, e_{m1}, e_{(m+1)i_{(m+1)}}, \dots, e_{(m+r'')i_{(m+r'')}}, e_1, \dots, e_v\} \setminus \{e_{wi_w}\}$ for some $i_h \in \{1, \dots, t_h\}$ with $1 \leq h \leq r'$, $m+1 \leq h \leq m+r''$ and $i_w \in \{1, \dots, t_w\}$ with $1 \leq w \leq r'$ or $i_w = 1$ with $r'+1 \leq w \leq m$ for some $w = h$ and $i_w = i_h$ appeared in T_{wi_w} . \square

In the following result, we give the primary decomposition of facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{U}_{n,m}^r))$.

Proposition 3.2. *Let $\Delta_s(\mathcal{U}_{n,m}^r)$ be the spanning simplicial complex of uni-cyclic multigraph $\mathcal{U}_{n,m}^r$ having n edges including r multiple edges within and outside the cycle of length m . Then,*

$$I_{\mathcal{F}}(\Delta_s(\mathcal{U}_{n,m}^r)) =$$

$$\left(\bigcap_{1 \leq a \leq v} (x_a) \right) \cap \left(\bigcap_{1 \leq i \leq r'; r'+1 \leq k \leq m} (x_{i1}, \dots, x_{it_i}, x_{k1}) \right) \cap \left(\bigcap_{r'+1 \leq k < l \leq m} (x_{k1}, x_{l1}) \right) \\ \cap \left(\bigcap_{1 \leq i < b \leq r'} (x_{i1}, \dots, x_{it_i}, x_{b1}, \dots, x_{bt_b}) \right) \cap \left(\bigcap_{m+1 \leq j \leq m+r''} (x_{j1}, \dots, x_{jt_j}) \right),$$

where t_i with $1 \leq i \leq r'$ is the number of multiple edges appeared in the i -th edge of the cycle and t_j with $m+1 \leq j \leq m+r''$ is the number of multiple edges appeared in the j th-edge outside the cycle.

Proof. Let $I_{\mathcal{F}}(\Delta_s(\mathcal{U}_{n,m}^r))$ be the facet ideal of the spanning simplicial complex $\Delta_s(\mathcal{U}_{n,m}^r)$. From ([8], Proposition 1.8), minimal prime ideals of the facet ideal $I_{\mathcal{F}}(\Delta)$ have one-to-one correspondence with the minimal vertex covers of the simplicial complex Δ . Therefore, in order to find the primary decomposition of the facet ideal $I_{\mathcal{F}}(\Delta_s(\mathcal{U}_{n,m}^r))$; it is sufficient to find all the minimal vertex covers of $\Delta_s(\mathcal{U}_{n,m}^r)$.

As e_a , $a \in [v]$ is not an edge of the cycle of uni-cyclic multigraph $\mathcal{U}_{n,m}^r$ and does not belong to any multiple edge of $\mathcal{U}_{n,m}^r$. Therefore, it is clear by definition of minimal vertex cover that $\{e_a\}$, $1 \leq a \leq v$ is a minimal vertex cover of $\Delta_s(\mathcal{U}_{n,m}^r)$. Moreover, a spanning tree is obtained by removing exactly $t_h - 1$ edges from each multiple edge with $h = 1, \dots, r', m+1, \dots, m+r''$ and

in addition, an edge from the resulting cycle of $\mathcal{U}_{n,m}^r$. We illustrate the result into the following cases.

Case 1. If atleast one multiple edge is appeared in the cycle of $\mathcal{U}_{n,m}^r$. Then, we cannot remove one complete multiple edge and one single edge from the cycle of $\mathcal{U}_{n,m}^r$ to get spanning tree. Therefore, $(x_{i1}, \dots, x_{it_i}, x_{k1})$ with $1 \leq i \leq r'$, $r' + 1 \leq k \leq m$ is a minimal vertex cover of the spanning simplicial complex $\Delta_s(\mathcal{U}_{n,m}^r)$ having non-empty intersection with all the spanning trees of $\mathcal{U}_{n,m}^r$. Moreover, two single edges cannot be removed from the cycle of $\mathcal{U}_{n,m}^r$ to get spanning tree. Consequently, (x_{k1}, x_{l1}) for $r' + 1 \leq k < l \leq m$ is a minimal vertex cover of $\Delta_s(\mathcal{U}_{n,m}^r)$ having non-empty intersection with all the spanning trees of $\mathcal{U}_{n,m}^r$.

Case 2. If atleast two multiple edges are appeared in the cycle of $\mathcal{U}_{n,m}^r$. Then, two complete multiple edges cannot be removed from the cycle of $\mathcal{U}_{n,m}^r$ to get spanning tree. Consequently, $(x_{i1}, \dots, x_{it_i}, x_{b1}, \dots, x_{bt_b})$ for $1 \leq i < b \leq r'$ is a minimal vertex cover of $\Delta_s(\mathcal{U}_{n,m}^r)$ having non-empty intersection with all the spanning trees of $\mathcal{U}_{n,m}^r$.

Case 3. If atleast one multiple edge appeared outside the cycle of $\mathcal{U}_{n,m}^r$. Then, one complete multiple edge outside the cycle of $\mathcal{U}_{n,m}^r$ cannot be removed to get spanning tree. So, $(x_{j1}, \dots, x_{jt_j})$ for $m + 1 \leq j \leq m + r''$ is a minimal vertex cover of $\Delta_s(\mathcal{U}_{n,m}^r)$ having non-empty intersection with all the spanning trees of $\mathcal{U}_{n,m}^r$. This completes the proof. \square

We give now formula for Euler characteristic of $\Delta_s(\mathcal{U}_{n,m}^r)$.

Theorem 3.3. Let $\Delta_s(\mathcal{U}_{n,m}^r)$ be spanning simplicial complex of uni-cyclic multigraph $\mathcal{U}_{n,m}^r$ having n edges including r multiple edges and a cycle of length m . Then, $\dim(\Delta_s(\mathcal{U}_{n,m}^r)) = n - \sum_{i=1}^{r'} t_i - \sum_{j=m+1}^{m+r''} t_j + r - 2$ and the Euler characteristic of $\Delta_s(\mathcal{U}_{n,m}^r)$ is given by

$$\begin{aligned} \chi(\Delta_s(\mathcal{U}_{n,m}^r)) &= \sum_{i=0}^{n-1} (-1)^i \left[\binom{n}{i+1} - \prod_{i=1}^{r'} t_i \left[\binom{n-\alpha+r'-m}{i+1-m} \right] \right. \\ &\quad \left. - \sum_{j=2}^{\beta} \left(\binom{\beta}{j} - \left(\sum_{m+1 \leq i_1 < \dots < i_j \leq m+r''} \prod_{k=i_1}^{i_j} \binom{t_k}{1} \right) \right) \sum_{l=j}^{\beta} (-1)^{l-j} \binom{\beta-j}{l-j} \binom{n-\alpha+r'-m-l}{i+1-m-l} \right] \\ &\quad - \sum_{j=2}^{\alpha+\beta} \left(\binom{\alpha+\beta}{j} - \left(\sum_{1 \leq i_1 < \dots < i_j \leq m+r''} \prod_{k=i_1}^{i_j} \binom{t_k}{1} \right) \right) \sum_{l=j}^{\alpha+\beta} (-1)^{l-j} \binom{\alpha+\beta-j}{l-j} \binom{n-l}{i+1-l} \right] \end{aligned}$$

with $\alpha = \sum_{i=1}^{r'} t_i$ and $\beta = \sum_{j=m+1}^{m+r''} t_j$ such that r' and r'' are the number of multiple edges appeared within and outside the cycle, respectively.

Proof. Let $E = \{e_{11}, \dots, e_{1t_1}, \dots, e_{r'1}, \dots, e_{r't_{r'}}, e_{(r'+1)1}, \dots, e_{m1}, e_{(m+1)1}, \dots, e_{(m+1)t_{m+1}}, \dots, e_{(m+r'')1}, \dots, e_{(m+r'')t_{m+r''}}, e_1, \dots, e_v\}$ be the edge set of uni-cyclic multigraph $\mathcal{U}_{n,m}^r$ having n edges including r multiple edges and a cycle of length m such that r' and r'' are the number of multiple edges appeared within and outside the cycle, respectively.

One can easily see that each facet T_{wi_w} of $\Delta_s(\mathcal{U}_{n,m}^r)$ is of the same dimension $n - \sum_{i=1}^{r'} t_i - \sum_{j=m+1}^{m+r''} t_j + r - 2 = n - \alpha - \beta + r - 2$ with $\alpha = \sum_{i=1}^{r'} t_i$ and $\beta = \sum_{j=m+1}^{m+r''} t_j$, see Lemma 3.1.

By definition, f_i is the number of subsets of E with $i + 1$ elements not containing cycle and multiple edges. There are $\prod_{i=1}^{r'} t_i \binom{n-\alpha+r'-m}{i+1-m}$ number of subsets of E containing cycle but not containing any multiple edge within the cycle. There are

$$\binom{n-\alpha+r'-m-\beta}{i+1-m-\beta} = \binom{\beta}{\beta} \sum_{l=\beta}^{\beta} (-1)^{l-\beta} \binom{\beta-\beta}{l-\beta} \binom{n-\alpha+r'-m-l}{i+1-m-l}$$

subsets of E containing cycle and β multiple edges of $\mathcal{U}_{n,m}^r$ outside the cycle but not containing any multiple edge within the cycle. There are

$$\begin{aligned} & \binom{\beta}{\beta-1} \left[\binom{n-\alpha+r'-m-(\beta-1)}{i+1-m-(\beta-1)} - \binom{n-\alpha+r'-m-\beta}{i+1-m-\beta} \right] \\ &= \binom{\beta}{\beta-1} \sum_{l=\beta-1}^{\beta} (-1)^{l-(\beta-1)} \binom{\beta-(\beta-1)}{l-(\beta-1)} \binom{n-\alpha+r'-m-l}{i+1-m-l} \end{aligned}$$

subsets of E containing cycle and $\beta-1$ multiple edges of $\mathcal{U}_{n,m}^r$ outside the cycle but not containing any multiple edge within the cycle. Continuing in similar manner, the number of subsets of E containing cycle and two edges from a multiple edge outside the cycle but not containing any multiple edge within the cycle is given by

$$\begin{aligned} & \binom{\beta-2}{0} \binom{n-\alpha+r'-m-2}{i+1-m-2} - \binom{\beta-2}{1} \binom{n-\alpha+r'-m-3}{i+1-m-3} + \binom{\beta-2}{2} \binom{n-\alpha+r'-m-4}{i+1-m-4} + \dots \\ &+ (-1)^{\beta-2} \binom{\beta-2}{\beta-2} \binom{n-\alpha+r'-m-\beta}{i+1-m-\beta} = \sum_{l=2}^{\beta} (-1)^{l-2} \binom{\beta-2}{l-2} \binom{n-\alpha+r'-m-l}{i+1-m-l}. \end{aligned}$$

There are $\left(\binom{\beta}{2} - \left(\sum_{m+1 \leq i_1 < i_2 \leq m+r''} \prod_{k=i_1}^{i_2} \binom{t_k}{1} \right) \right)$ choices of two edges from a multiple edge outside the cycle. Therefore, we obtain

$$\left(\binom{\beta}{2} - \left(\sum_{m+1 \leq i_1 < i_2 \leq m+r''} \prod_{k=i_1}^{i_2} \binom{t_k}{1} \right) \right) \sum_{l=2}^{\beta} (-1)^{l-2} \binom{\beta-2}{l-2} \binom{n-\alpha+r'-m-l}{i+1-m-l}$$

the number of subsets of E with $i+1$ elements containing cycle and all possible choices of two edges from a multiple edge outside the cycle but not containing multiple edges within the cycle.

Now, we use inclusion exclusion principal to obtain
(number of subsets of E containing cycle but not containing multiple edges) =
 $\prod_{i=1}^{r'} t_i$ [(number of subsets of E with $i+1$ elements containing cycle but not containing multiple edges within the cycle) - (number of subsets of E with $i+1$ elements containing cycle and β multiple edges outside the cycle but


not containing multiple edges within the cycle) – (number of subsets of E with $i + 1$ elements containing cycle and $\beta - 1$ multiple edges outside the cycle but not containing multiple edges within the cycle) – \dots – (number of subsets of E with $i + 1$ elements containing cycle and two edges from a multiple edge outside the cycle but not containing multiple edges within the cycle)]

$$\begin{aligned}
&= \prod_{i=1}^{r'} t_i \left[\binom{n-\alpha+r'-m}{i+1-m} - \binom{\beta}{\beta} \sum_{l=\beta}^{\beta} (-1)^{l-\beta} \binom{\beta-\beta}{l-\beta} \binom{n-\alpha+r'-m-l}{i+1-m-l} \right. \\
&\quad - \binom{\beta}{\beta-1} \sum_{l=\beta-1}^{\beta} (-1)^{l-(\beta-1)} \binom{\beta-(\beta-1)}{l-(\beta-1)} \binom{n-\alpha+r'-m-l}{i+1-m-l} \\
&\quad - \dots - \left(\binom{\beta}{2} - \left(\sum_{m+1 \leq i_1 < i_2 \leq m+r''} \prod_{k=i_1}^{i_2} \binom{t_k}{1} \right) \right) \sum_{l=2}^{\beta} (-1)^{l-2} \binom{\beta-2}{l-2} \binom{n-\alpha+r'-m-l}{i+1-m-l} \Big] \\
&= \prod_{i=1}^{r'} t_i \left[\binom{n-\alpha+r'-m}{i+1-m} \right. \\
&\quad - \sum_{j=2}^{\beta} \left(\binom{\beta}{j} - \left(\sum_{m+1 \leq i_1 < \dots < i_j \leq m+r''} \prod_{k=i_1}^{i_j} \binom{t_k}{1} \right) \right) \sum_{l=j}^{\beta} (-1)^{l-j} \binom{\beta-j}{l-j} \binom{n-\alpha+r'-m-l}{i+1-m-l} \Big].
\end{aligned}$$

Therefore, we compute

$f_i =$ (number of subsets of E with $i+1$ elements) – (number of subsets of E with $i + 1$ elements containing cycle but not containing multiple edges) – (number of subsets of E with $i+1$ elements containing $\alpha + \beta$ multiple edges) – (number of subsets of E with $i+1$ elements containing $\alpha + \beta - 1$ multiple edges) – \dots – (number of subsets of E with $i + 1$ elements containing two edges from a multiple edge of $\mathcal{U}_{n,m}^r$)

$$\begin{aligned}
&= \binom{n}{i+1} - \prod_{i=1}^{r'} t_i \left[\binom{n-\alpha+r'-m}{i+1-m} \right. \\
&\quad - \sum_{j=2}^{\beta} \left(\binom{\beta}{j} - \left(\sum_{m+1 \leq i_1 < \dots < i_j \leq m+r''} \prod_{k=i_1}^{i_j} \binom{t_k}{1} \right) \right) \sum_{l=j}^{\beta} (-1)^{l-j} \binom{\beta-j}{l-j} \binom{n-\alpha+r'-m-l}{i+1-m-l} \Big] \\
&\quad - \binom{\alpha+\beta}{\alpha+\beta} \sum_{l=\alpha+\beta}^{\alpha+\beta} (-1)^{l-(\alpha+\beta)} \binom{(\alpha+\beta)-(\alpha+\beta)}{l-(\alpha+\beta)} \binom{n-l}{i+1-l} \\
&\quad - \binom{\alpha+\beta}{\alpha+\beta-1} \sum_{l=\alpha+\beta-1}^{\alpha+\beta} (-1)^{l-(\alpha+\beta-1)} \binom{(\alpha+\beta)-(\alpha+\beta-1)}{l-(\alpha+\beta-1)} \binom{n-l}{i+1-l} \\
&\quad - \dots - \left(\binom{\alpha+\beta}{2} - \left(\sum_{1 \leq i_1 < i_2 \leq m+r''} \prod_{k=i_1}^{i_2} \binom{t_k}{1} \right) \right) \sum_{l=2}^{\alpha+\beta} (-1)^{l-2} \binom{\alpha+\beta-2}{l-2} \binom{n-l}{i+1-l} \\
&= \binom{n}{i+1} - \prod_{i=1}^{r'} t_i \left[\binom{n-\alpha+r'-m}{i+1-m} \right. \\
&\quad - \sum_{j=2}^{\beta} \left(\binom{\beta}{j} - \left(\sum_{m+1 \leq i_1 < \dots < i_j \leq m+r''} \prod_{k=i_1}^{i_j} \binom{t_k}{1} \right) \right) \sum_{l=j}^{\beta} (-1)^{l-j} \binom{\beta-j}{l-j} \binom{n-\alpha+r'-m-l}{i+1-m-l} \Big] \\
&\quad - \sum_{j=2}^{\alpha+\beta} \left(\binom{\alpha+\beta}{j} - \left(\sum_{1 \leq i_1 < \dots < i_j \leq m+r''} \prod_{k=i_1}^{i_j} \binom{t_k}{1} \right) \right) \sum_{l=j}^{\alpha+\beta} (-1)^{l-j} \binom{\alpha+\beta-j}{l-j} \binom{n-l}{i+1-l}. \quad \square
\end{aligned}$$

FIGURE 1. $\mathcal{U}_{7,3}^2$

Example 3.4. Let $E = \{e_{11}, e_{12}, e_{13}, e_{21}, e_{31}, e_{41}, e_{42}\}$ be the edge set of unicyclic multigraph $\mathcal{U}_{7,3}^2$ having 7 edges including 2 multiple edges and a cycle of length 3, as shown in Figure 1. By cutting-down method, we obtain

$$s(\mathcal{U}_{7,3}^2) = \{\{e_{21}, e_{31}, e_{41}\}, \{e_{11}, e_{31}, e_{41}\}, \{e_{11}, e_{21}, e_{41}\}, \{e_{12}, e_{31}, e_{41}\}, \{e_{12}, e_{21}, e_{41}\}, \{e_{13}, e_{31}, e_{41}\}, \{e_{13}, e_{21}, e_{41}\}, \{e_{21}, e_{31}, e_{42}\}, \{e_{11}, e_{31}, e_{42}\}, \{e_{11}, e_{21}, e_{42}\}, \{e_{12}, e_{31}, e_{42}\}, \{e_{12}, e_{21}, e_{42}\}, \{e_{13}, e_{31}, e_{42}\}, \{e_{13}, e_{21}, e_{42}\}\}.$$

By definition, f_i is the number of subsets of E with $i + 1$ elements not containing cycle and multiple edges. Since, $\{e_{11}\}, \{e_{12}\}, \{e_{13}\}, \{e_{21}\}, \{e_{31}\}, \{e_{41}\}$ and $\{e_{42}\}$ are subsets of E containing one element. It implies that $f_0 = 7$. There are $\{e_{11}, e_{21}\}, \{e_{11}, e_{31}\}, \{e_{11}, e_{41}\}, \{e_{11}, e_{42}\}, \{e_{12}, e_{21}\}, \{e_{12}, e_{31}\}, \{e_{12}, e_{41}\}, \{e_{12}, e_{42}\}, \{e_{13}, e_{21}\}, \{e_{13}, e_{31}\}, \{e_{13}, e_{41}\}, \{e_{13}, e_{42}\}, \{e_{21}, e_{31}\}, \{e_{21}, e_{41}\}, \{e_{21}, e_{42}\}, \{e_{31}, e_{41}\}, \{e_{31}, e_{42}\}$ subsets of E containing two elements but not containing cycle and multiple edges. So, $f_1 = 17$. We know that the spanning trees of $\mathcal{U}_{7,3}^2$ are 2-dimensional facets of the spanning simplicial complex $\Delta_s(\mathcal{U}_{7,3}^2)$. Therefore, $f_2 = 14$. Thus,

$$\chi(\Delta_s(\mathcal{U}_{7,3}^2)) = f_0 - f_1 + f_2 = 7 - 17 + 14 = 4.$$

Now, we compute the Euler characteristic of $\Delta_s(\mathcal{U}_{7,3}^2)$ by using Theorem 3.3. We observe that, $n = 7$, $r' = 1$, $r'' = 1$, $r = 2$, $m = 3$, $\alpha = 3$, $\beta = 2$ and $0 \leq i \leq d$, where $d = n - \alpha - \beta + r - 2 = 2$ is the dimension of $\Delta_s(\mathcal{U}_{7,3}^2)$.

By substituting these values in Theorem 3.3, we get

$$\begin{aligned} f_i &= \binom{7}{i+1} - 3 \left[\binom{2}{i+1-3} - \binom{2}{2} \binom{0}{i+1-5} \right] - [\binom{5}{2} - \binom{3}{1} \binom{2}{1}] \left(\binom{3}{0} \binom{5}{i+1-2} - \binom{3}{1} \binom{4}{i+1-3} + \right. \\ &\quad \left. \binom{3}{2} \binom{3}{i+1-4} - \binom{3}{3} \binom{2}{i+1-4} \right) - \binom{5}{3} \left(\binom{2}{0} \binom{4}{i+1-3} - \binom{2}{1} \binom{3}{i+1-4} + \binom{2}{2} \binom{2}{i+1-5} \right) - \\ &\quad \binom{5}{4} \left(\binom{1}{0} \binom{3}{i+1-4} - \binom{1}{1} \binom{2}{i+1-5} \right) - \binom{5}{5} \binom{2}{i+1-5}. \end{aligned}$$

Alternatively, we compute $(f_0, f_1, f_2) = (7, 17, 14)$ and $\chi(\Delta_s(\mathcal{U}_{7,3}^2)) = 4$.

We compute now the Betti numbers of $\Delta_s(\mathcal{U}_{7,3}^2)$. The facet ideal of $\Delta_s(\mathcal{U}_{7,3}^2)$ is given by

$$I_{\mathcal{F}}(\Delta_s(\mathcal{U}_{7,3}^2)) = \langle x_{F_{21,31,41}}, x_{F_{11,31,41}}, x_{F_{11,21,41}}, x_{F_{12,31,41}}, x_{F_{12,21,41}}, x_{F_{13,31,41}}, x_{F_{13,21,41}}, x_{F_{21,31,42}}, x_{F_{11,31,42}}, x_{F_{11,21,42}}, x_{F_{12,31,42}}, x_{F_{12,21,42}}, x_{F_{13,31,42}}, x_{F_{13,21,42}} \rangle.$$

We consider the chain complex of $\Delta_s(\mathcal{U}_{7,3}^2)$

$$0 \xrightarrow{\partial_3} C_2 \xrightarrow{\partial_2} C_1 \xrightarrow{\partial_1} C_0 \xrightarrow{\partial_0} 0$$

The homology groups are given by $H_i = \frac{Ker(\partial_i)}{Im(\partial_{i+1})}$ with $i = 0, 1, 2$.

Therefore, the i -th Betti number of $\Delta_s(\mathcal{U}_{7,3}^2)$ is given by

$$\beta_i = rank(H_i) = rank(Ker(\partial_i)) - rank(Im(\partial_{i+1})) \text{ with } i = 0, 1, 2.$$

Now, we compute rank and nullity of the matrix ∂_i of order $f_i \times f_{i+1}$ with $i = 0, 1, 2$.

The boundary homomorphism $\partial_2 : C_2(\Delta_s(\mathcal{U}_{7,3}^2)) \rightarrow C_1(\Delta_s(\mathcal{U}_{7,3}^2))$ can be expressed as

$$\begin{aligned} \partial_2(F_{21,31,41}) &= F_{31,41} - F_{21,41} + F_{21,31}; \quad \partial_2(F_{11,31,41}) = F_{31,41} - F_{11,41} + F_{11,31}; \\ \partial_2(F_{11,21,41}) &= F_{21,41} - F_{11,41} + F_{11,21}; \quad \partial_2(F_{12,31,41}) = F_{31,41} - F_{12,41} + F_{12,31}; \\ \partial_2(F_{12,21,41}) &= F_{21,41} - F_{12,41} + F_{12,21}; \quad \partial_2(F_{13,31,41}) = F_{31,41} - F_{13,41} + F_{13,31}; \\ \partial_2(F_{13,21,41}) &= F_{21,41} - F_{13,41} + F_{13,21}; \quad \partial_2(F_{21,31,42}) = F_{31,42} - F_{21,42} + F_{21,31}; \\ \partial_2(F_{11,31,42}) &= F_{31,42} - F_{11,42} + F_{11,31}; \quad \partial_2(F_{11,21,42}) = F_{21,42} - F_{11,42} + F_{11,21}; \\ \partial_2(F_{12,31,42}) &= F_{31,42} - F_{12,42} + F_{12,31}; \quad \partial_2(F_{12,21,42}) = F_{21,42} - F_{12,42} + F_{12,21}; \\ \partial_2(F_{13,31,42}) &= F_{31,42} - F_{13,42} + F_{13,31}; \quad \partial_2(F_{13,21,42}) = F_{21,42} - F_{13,42} + F_{13,21}. \end{aligned}$$

The boundary homomorphism $\partial_1 : C_1(\Delta_s(\mathcal{U}_{7,3}^2)) \rightarrow C_0(\Delta_s(\mathcal{U}_{7,3}^2))$ can be written as

$$\begin{aligned} \partial_1(F_{11,21}) &= e_{21} - e_{11}; \quad \partial_1(F_{11,31}) = e_{31} - e_{11}; \quad \partial_1(F_{11,41}) = e_{41} - e_{11}; \quad \partial_1(F_{11,42}) = e_{42} - e_{11}; \\ \partial_1(F_{12,21}) &= e_{21} - e_{12}; \quad \partial_1(F_{12,31}) = e_{31} - e_{12}; \quad \partial_1(F_{12,41}) = e_{41} - e_{12}; \\ \partial_1(F_{12,42}) &= e_{42} - e_{12}; \quad \partial_1(F_{13,21}) = e_{21} - e_{13}; \quad \partial_1(F_{13,31}) = e_{31} - e_{13}; \quad \partial_1(F_{13,41}) = e_{41} - e_{13}; \\ \partial_1(F_{13,42}) &= e_{42} - e_{13}; \quad \partial_1(F_{21,31}) = e_{31} - e_{21}; \quad \partial_1(F_{21,41}) = e_{41} - e_{21}; \\ \partial_1(F_{21,42}) &= e_{42} - e_{21}; \quad \partial_1(F_{31,41}) = e_{41} - e_{31}; \quad \partial_1(F_{31,42}) = e_{42} - e_{31}. \end{aligned}$$

Then, by using MATLAB, we compute rank of $\partial_2 = 11$; nullity of $\partial_2 = 3$; rank of $\partial_1 = 6$; nullity of $\partial_1 = 11$.

Therefore, the Betti numbers are given by $\beta_0 = rank(Ker(\partial_0)) - rank(Im(\partial_1)) = 7 - 6 = 1$; $\beta_1 = rank(Ker(\partial_1)) - rank(Im(\partial_2)) = 11 - 11 = 0$; $\beta_2 = rank(Ker(\partial_2)) - rank(Im(\partial_3)) = 3 - 0 = 3$.

Alternatively, the Euler characteristic of $\Delta_s(\mathcal{U}_{7,3}^2)$ is given by

$$\chi(\Delta_s(\mathcal{U}_{7,3}^2)) = \beta_0 - \beta_1 + \beta_2 = 1 - 0 + 3 = 4.$$

REFERENCES

- [1] I. Anwar, Z. Raza and A. Kashif, Spanning Simplicial Complexes of Uni-Cyclic Graphs, *Algebra Colloquium*, **22** (2015), no.4, 707-710.
- [2] S. Faridi, The Facet Ideal of a Simplicial Complex, *Manuscripta*, **109** (2002), 159-174.
- [3] F. Harary, *Graph Theory*, Reading, MA: Addison-Wesley, 1994.
- [4] A. Hatcher, *Algebraic Topology*, Cambridge University Press, 2002.
- [5] A. Kashif, I. Anwar and Z. Raza, On the Algebraic Study of Spanning Simplicial Complex of r -cycles Graphs $G_{n,r}$, *ARS Combinatoria*, **115** (2014), 89-99.
- [6] Y. Pan, R. Li and G. Zhu, Spanning Simplicial Complexes of n -Cyclic Graphs with a Common Vertex, *International Electronic Journal of Algebra*, **17** (2015), 180-187.

- [7] J. J. Rotman, *An Introduction to Algebraic Topology*, Springer-Verlag, New York, 1988.
- [8] R. H. Villarreal, *Monomial Algebras*. Dekker, New York, 2001.
- [9] G. Zhu, F. Shi and Y. Geng, Spanning Simplicial Complexes of n -Cyclic Graphs with a Common Edge, *International Electronic Journal of Algebra*, **15** (2014), 132-144.

COMSATS INSTITUTE OF INFORMATION TECHNOLOGY, LAHORE, PAKISTAN
E-mail address: `drimranahmed@ciitlahore.edu.pk`

COMSATS INSTITUTE OF INFORMATION TECHNOLOGY, LAHORE, PAKISTAN
E-mail address: `shahid_nankana@yahoo.com`