

Convergence of the HX Preconditioner for Maxwell's Equations with Jump Coefficients (i): Extensions of the Regular Decomposition

Qiya Hu¹

Abstract

This paper is the first of the two serial articles, aiming to prove the convergence of the HX preconditioner for Maxwell's equations with jump coefficients, originally proposed by Hiptmair and Xu [17]. In this paper, we establish several extensions of the discrete regular decomposition for edge finite element functions defined in three-dimensional domains. The functions defined by the new discrete regular decompositions can inherit zero degrees of freedom of the considered edge finite element function on some faces and edges of polyhedral domains as well as of some non-Lipschitz domains, and possess nearly optimal stability with only a *logarithmic* factor. These regular decompositions will be used to prove the convergence of the HX preconditioner for Maxwell's equations with jump coefficients in a subsequent paper [19].

Key Words. Maxwell's equations, Nedelec elements, regular decomposition, stability

AMS(MOS) subject classification. 65N30, 65N55

Contents

1	Introduction	2
2	Preliminaries	3
2.1	Sobolev spaces and norms	3
2.2	Edge and nodal element spaces	3
2.3	Some special notions and notations	4
3	Main results	5
4	Basic regular decompositions	7
4.1	Curl -bounded regular decompositions on Lipschitz polyhedrons	7
4.2	Curl -bounded and L^2 -stable regular decompositions on non-convex polyhedrons	11
5	Proof of Theorem 3.1	17
6	Proof of Theorem 3.2-Theorem 3.4	27

¹1. LSEC, ICMSEC, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; 2. School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China (hqy@lsec.cc.ac.cn). This author was supported by the Natural Science Foundation of China G11571352.

1 Introduction

It is known that any vector-valued function in $H(\mathbf{curl})$ space can be decomposed into the sum of a H^1 vector-valued function and the gradient of a H^1 scalar-valued function (refer to [12] and [11]), and the decomposition is stable with respect to the standard norms. If the sum is orthogonal with respect to the L^2 inner product, the decomposition is called the Helmholtz decomposition, otherwise, it is called a regular decomposition (which is not unique). The regular decompositions are more popular than the Helmholtz decomposition because the regular decompositions are valid on the general Lipschitz domains, whereas the Helmholtz decomposition holds only on smooth or convex domains (if H^1 -regularity of the vector-valued function is not required, the Helmholtz decomposition can always be found for the general Lipschitz domains). In the study of Maxwell's equations, the Helmholtz decompositions and regular decompositions can be used to transform a problem for a $H(\mathbf{curl})$ function into problems on two H^1 functions.

The Nedelec edge finite element method (refer to [28]) is a popular discretization method for Maxwell's equations. Suitable decompositions for edge finite element functions, known as discrete (Helmholtz or regular) decompositions, play a key role in the convergence analysis of preconditioners for Maxwell's equations (see, for example, [14, 17, 21, 22, 23, 29, 32, 33]). In this analysis, Maxwell's equations in a *homogeneous* medium are often assumed because the stability of the standard Helmholtz decomposition only holds for the usual norms. However, nonhomogeneous media often occur in practical applications; therefore, some weighted norms have to be introduced. A natural question is: whether the discrete decompositions are still stable with respect to the weighted norms? This question was first positively answered in a study [21] by constructing a discrete *weighted orthogonal* Helmholtz decomposition that was proved to be almost stable with respect to weighted norms.

For the numerical analysis of Maxwell's equations in a nonhomogeneous medium, a discrete *regular* decomposition needs to be developed for edge finite element functions defined in three-dimensional domains such that the decomposition is stable with respect to weighted norms. This study is complex; therefore, it will be published in two parts. Herein, some technical tools are developed to derive some extensions of the discrete regular decomposition for edge finite element functions. The standard regular (and Helmholtz) decompositions possess a vital property: when the vector-valued function being considered has zero trace on the boundary of the underlying domain, the functions defined by the decomposition also have zero trace on this boundary. We will construct a discrete *regular* decomposition on a polyhedral domain such that the inheritance of the zero trace on the boundary can be maintained when the boundary is replaced by a union of some local faces and edges of the polyhedral domain. In particular, we also establish the corresponding decompositions for edge finite element function on some non-Lipschitz domains, which are unions of two polyhedral domains intersecting at one edge or vertex. We will show that the resulting regular decompositions possess stability estimates with only one *logarithmic* factor. These interesting results will be used in a subsequent article [19] to develop a discrete regular decomposition that is nearly stable with respect to weighted norms, and the convergence of the HX preconditioner for the case with jump coefficients will be further proved.

The remainder of this paper is organized as follows. Section 2 defines some edge finite element subspaces. Section 3 presents the main results. Section 4 discusses the regular decompositions that preserve zero-tangential traces on faces. Section 5 presents several discrete regular decompositions that preserve local zero-tangential components on edges and faces; moreover, this section proves the result for polyhedral domains as well. In Section 6, discrete regular decompositions on some non-Lipschitz domains are derived.

2 Preliminaries

In this section we introduce some fundamental notions and notations.

2.1 Sobolev spaces and norms

For an open and connected bounded domain G in $\mathbf{R}^3$, let $H^1(G)$ be the standard Sobolev space consisting of the functions whose weak derivatives belong to $L^2(G)$. Define the $H(\mathbf{curl})$ space as follows

$$H(\mathbf{curl}; G) = \{\mathbf{v} \in L^2(G)^3; \mathbf{curl} \mathbf{v} \in L^2(G)^3\}.$$

For the two spaces, we use natural norms

$$\|\mathbf{v}\|_{1,G} = (|\mathbf{v}|_{1,G}^2 + \|\mathbf{v}\|_{0,G}^2)^{\frac{1}{2}}, \quad \mathbf{v} \in (H^1(G))^3$$

and

$$\|\mathbf{v}\|_{\mathbf{curl},G} = (\|\mathbf{curl} \mathbf{v}\|_{0,G}^2 + \|\mathbf{v}\|_{0,G}^2)^{\frac{1}{2}}, \quad \mathbf{v} \in H(\mathbf{curl}; G).$$

For a polyhedron G , let Γ be a (closed) face or a union of several faces of G . Define

$$H_\Gamma(\mathbf{curl}; G) = \{\mathbf{v} \in H(\mathbf{curl}; G) : \mathbf{v} \times \mathbf{n} = 0 \text{ on } \Gamma\}$$

and

$$H_\Gamma^1(G) = \{v \in H^1(G) : v = 0 \text{ on } \Gamma\}.$$

2.2 Edge and nodal element spaces

For a polyhedron G , let G be divided into a union of small tetrahedral elements of size h , and let $\mathcal{T}_h$ denote the resulting triangulation of the domain G . For convenience, we assume that the triangulation $\mathcal{T}_h$ is quasi-uniform. If the assumption is not satisfied, the estimates built in this paper may depend on the shape parameters of elements in $\mathcal{T}_h$. We use $\mathcal{E}_h$ and $\mathcal{N}_h$ to denote the set of edges of $\mathcal{T}_h$ and the set of nodes in $\mathcal{T}_h$ respectively. Then the Nedelec edge element space, of the lowest order, is a subspace of piecewise linear polynomials defined on $\mathcal{T}_h$:

$$V_h(G) = \left\{ \mathbf{v} \in H(\mathbf{curl}; G); \mathbf{v}|_K \in R(K), \forall K \in \mathcal{T}_h \right\},$$

where $R(K)$ is a subset of all linear polynomials on the element K of the form:

$$R(K) = \left\{ \mathbf{a} + \mathbf{b} \times \mathbf{x}; \mathbf{a}, \mathbf{b} \in \mathbf{R}^3, \mathbf{x} \in K \right\}.$$

It is known that, for any $\mathbf{v} \in V_h(G)$, its tangential components are continuous on all edges in $\mathcal{E}_h$, and $\mathbf{v}$ is uniquely determined by its moments on edges e of $\mathcal{T}_h$:

$$M_h(\mathbf{v}) = \left\{ \lambda_e(\mathbf{v}) = \int_e \mathbf{v} \cdot \mathbf{t}_e ds; e \in \mathcal{E}_h \right\},$$

where $\mathbf{t}_e$ denotes the unit vector on an edge e , and this notation will be used to denote the unit vector on any edge or union of edges, either from an element $K \in \mathcal{T}_h$ or from G itself. For example, for a face F of G , the notation $\mathbf{t}_{\partial F}$ denotes the unit vector along ∂F .

For a vector-valued function $\mathbf{v}$ with appropriate smoothness, we introduce its edge element “projection” $\mathbf{r}_h \mathbf{v}$ such that $\mathbf{r}_h \mathbf{v} \in V_h(G)$, and $\mathbf{r}_h \mathbf{v}$ and $\mathbf{v}$ have the same moments as in $M_h(\mathbf{v})$. Such an operator $\mathbf{r}_h$ can be chosen as the standard edge element interpolation

operator or the edge element projector $\mathbf{R}_D^1$ introduced in [16]. The operator $\mathbf{r}_h$ will be used in the construction of a stable decomposition for any function $\mathbf{v}_h \in V_h(G)$.

As we will see, the edge element analysis also involves frequently the nodal element space. For this purpose, we use $Z_h(G)$ to denote the standard continuous piecewise linear finite element space associated with the triangulation $\mathcal{T}_h$.

2.3 Some special notions and notations

Throughout this paper, we shall frequently use the notations $\lesssim$. For any two non-negative quantities x and y , $x \lesssim y$ means that $x \leq Cy$ for some constant C independent of mesh size h . For a non-negative quantity z , when $1 \lesssim z \lesssim 1$, we simply write $z = O(1)$.

From now on, a “usual” polyhedron G means that G is a bounded polyhedron with a diameter $O(1)$ and vanishing first Betti number, and G has a fixed number (independent of h) of flat faces only, but it may be non-convex. A face F of G is always understood as a “complete” face, i.e., F satisfies $F = \pi \cap \partial G$ for the plane π containing F , which means that F is just a closed (flat) face instead of only a part of a face.

We will often use F , E and v to denote a general face, edge and vertex of G , respectively, but use e to denote a general edge in $\mathcal{E}_h$ associated with a triangulation $\mathcal{T}_h$ on G . Throughout this paper a face F or an edge E is understood as a closed subset of ∂G .

A connected union Γ of several faces of G is called a *connected “Lipschitz” union of some faces* if there is no isolated vertex on Γ (see Figure 1), i.e., for any face $F \subset \Gamma$, there exists another face $F' \subset \Gamma$ such that the intersection $F \cap F'$ is an edge; otherwise, it is called a *connected “non-Lipschitz” union of some faces*. In essence, *connected “Lipschitz” union of some faces* can be understood as a Lipschitz subset of ∂G , but a *connected “non-Lipschitz” union of some faces* is not a Lipschitz subset of ∂G .




Figure 1. F_1 and F_2 are two opposite lateral faces of four-sided pyramid G , and F_3 denotes another lateral face that has a common edge with each of F_1 and F_2 . The set $\Gamma = F_1 \cup F_2 \cup F_3$ is a *connected “Lipschitz” union of three faces* (left), but the set $\Gamma = F_1 \cup F_2$ is a *connected “non-Lipschitz” union of two faces* (right), where the point v is an isolated vertex of Γ .

As we will see, a connected union of several edges has no essential difference from an edge. For simplicity of exposition, an “edge” is always understood as an “edge” or a “connected union of edges” later.

3 Main results

In this section we describe the new regular decompositions (and their stabilities), which will be constructed and analyzed in the upcoming sections.

Let G be a usual polyhedron or a connected union of two usual polyhedra, on which a triangulation $\mathcal{T}_h$ is given, and let Γ denote a union of faces and edges of G . The following assumption will be used repeatedly:

Assumption 3.1. The vector-valued function $\mathbf{v}_h \in V_h(G)$ has zero degrees of freedom $\lambda_e(\mathbf{v}_h) = 0$ for all $e \subset \Gamma$, where $e \in \mathcal{E}_h$.

The first result can be regarded as a variant of the existing discrete regular decompositions on polyhedral domains.

Theorem 3.1 *Let G be a usual polyhedron, and let Γ be a (may be non-connected) union of some faces and edges of G . Suppose that $\mathbf{v}_h$ satisfies **Assumption 3.1**. Then the function $\mathbf{v}_h$ admits a decomposition*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h \quad (3.1)$$

for some $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$ such that $p_h = 0$ and $\mathbf{w}_h = \mathbf{0}$ on Γ , and $\lambda_e(\mathbf{R}_h) = 0$ for all $e \subset \Gamma$. Moreover, we have

$$\|\mathbf{w}_h\|_{1,G} + h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\text{curl},G} \quad (3.2)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\text{curl},G}. \quad (3.3)$$

We would like to emphasize that, in Theorem 3.1, the union set Γ may contain some edges of the polyhedron, which is different from existing results.

Remark 3.1 *In particular, when Γ is a connected set of ∂G , the norm on the right side of (3.2) can be replaced by the **curl** semi-norm. If there is not an isolated edge in Γ , i.e., Γ is a union of faces only, then the norm on the right side of (3.3) can be replaced by the L^2 norm. When $\Gamma = \cup_{j=1}^J \Gamma_j$ with Γ_j being a connected “Lipschitz” union of some faces of G , the factor $\log(1/h)$ in the estimates (3.2) and (3.3) can be dropped.*

We would like to extend the above theorem to the case of non-Lipschitz polyhedra. Let G_1 and G_2 be two intersecting (usual) polyhedra, and define G as their union: $\bar{G} = \bar{G}_1 \cup \bar{G}_2$. We consider two particular cases: (1) the intersection $\bar{G}_1 \cap \bar{G}_2$ is an edge E of G_1 and G_2 ; (2) the intersection $\bar{G}_1 \cap \bar{G}_2$ is a vertex v of G_1 and G_2 . For the two cases, G is not a Lipschitz domain.

We assume that the two triangulations on G_1 and G_2 are matching on the intersection $\bar{G}_1 \cap \bar{G}_2$, i.e., their nodes are coincident on $\bar{G}_1 \cap \bar{G}_2$. An edge finite element function $\mathbf{v}_h \in V_h(G)$ means that $\mathbf{v}_h|_{G_i} \in V_h(G_i)$ ($i = 1, 2$) and $\mathbf{v}_h$ has continuous tangential complements on E for Case (1). A nodal finite element function $p_h \in Z_h(G)$ means that $p_h|_{G_i} \in Z_h(G_i)$ ($i = 1, 2$) and p_h is continuous on the intersection $\bar{G}_1 \cap \bar{G}_2$. Norms of finite element functions on G are understood in the natural manner, for instance, the norm of $\mathbf{v}_h \in V_h(G)$ is defined as $\|\mathbf{v}_h\|_{\text{curl},G} = (\|\mathbf{v}_h\|_{\text{curl},G_1}^2 + \|\mathbf{v}_h\|_{\text{curl},G_2}^2)^{\frac{1}{2}}$. For a union Γ of faces and edges of G , define $\Gamma_i = \Gamma \cap \partial G_i$ ($i = 1, 2$).

The following result is a direct extension of Theorem 3.1 to Case (1).

Theorem 3.2 *Let G be a union of two usual polyhedra G_1 and G_2 , with $\bar{G}_1 \cap \bar{G}_2$ being the common edge E of G_1 and G_2 , and let Γ be a union of some faces and edges of G . Suppose that $\mathbf{v}_h$ satisfies **Assumption 3.1**. Then $\mathbf{v}_h$ admits a decomposition*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h \quad (3.4)$$

for some $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$ such that $p_h = 0$ and $\mathbf{w}_h = \mathbf{0}$ on Γ , and $\lambda_e(\mathbf{R}_h) = \mathbf{0}$ for all $e \subset \Gamma$. Moreover, we have

$$\|\mathbf{w}_h\|_{1,G} + h^{-1}\|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h)\|\mathbf{v}_h\|_{\mathbf{curl},G} \quad (3.5)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h)\|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (3.6)$$

In particular, when Γ contains faces only and $E \subset \Gamma_i$ for $i = 1, 2$, the logarithmic factor in the above estimates can be dropped. If Γ is connected, then the complete norm on the right side of (3.5) can be replaced by the **curl** semi-norm.

Now we consider Case (2), i.e., $\bar{G}_1 \cap \bar{G}_2 = \mathbf{v}$ is a vertex of G_1 and G_2 . For this case, the problem is a bit complicated, and a similar result with Theorem 3.2 cannot be built yet except that the considered function $\mathbf{v}_h$ meets an additional condition.

Theorem 3.3 *Let G be a union of two usual polyhedra G_1 and G_2 , with $\bar{G}_1 \cap \bar{G}_2$ being the common vertex $\mathbf{v}$ of G_1 and G_2 . Let Γ denote a union of some faces and edges of G_1 and G_2 . Suppose that $\mathbf{v}_h$ satisfies **Assumption 3.1**. Then there exists a functional $\mathcal{F}$ (whose definition will be given in Section 6) such that, if $\mathbf{v}_h$ satisfies the constraint $\mathcal{F}\mathbf{v}_h = 0$, the function $\mathbf{v}_h$ has a decomposition*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h \quad (3.7)$$

for some $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$ such that p_h , $\mathbf{w}_h$ and $\mathbf{R}_h$ have zero degrees of freedom on Γ . Moreover, we have

$$\|\mathbf{w}_h\|_{1,G} + h^{-1}\|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h)\|\mathbf{v}_h\|_{\mathbf{curl},G} \quad (3.8)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h)\|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (3.9)$$

When both Γ_1 and Γ_2 are connected, the norm on the right side of (3.8) can be replaced by the **curl** semi-norm.

Remark 3.2 *The additional condition $\mathcal{F}\mathbf{v}_h = 0$ in Theorem 3.3 seems absolutely necessary. The value of an edge finite element function $\mathbf{v}_h$ is not uniquely defined at the common vertex $\mathbf{v}$ of G_1 and G_2 , but the two nodal finite element functions defined by the regular decomposition of $\mathbf{v}_h$ are required to be continuous at the vertex $\mathbf{v}$. Intuitively understanding, there exists a gap between $\mathbf{v}_h$ and the gradient of the scalar finite element function p_h . The role of the additional condition on $\mathbf{v}_h$ is to fill this gap.*

We can also consider the case that G is a union of more polyhedrons. Let $G_1, G_2, \dots, G_s$ be usual polyhedrons that may be non-convex, and let G be a union of $G_1, G_2, \dots, G_s$ (then G is a non-Lipchitz domain) such that the intersection of any two polyhedrons in $G_1, G_2, \dots, G_s$ just is the same vertex $\mathbf{v}$, i.e., $\bar{G}_i \cap \bar{G}_j = \mathbf{v}$ for $i \neq j$, which implies that $\bar{G}_1 \cap \bar{G}_2 \cap \dots \cap \bar{G}_s = \mathbf{v}$.

Theorem 3.4 *Let G_i ($i = 1, \dots, s$; $s \geq 3$) and G be defined as above, and let Γ denote a union of some faces and edges of $G_1, \dots, G_s$. Suppose that $\mathbf{v}_h$ satisfies **Assumption 3.1**. Then there exist $s-1$ functionals $\mathcal{F}_i$ (whose definition will be given at the ending of Section 6) such that, if $\mathbf{v}_h$ satisfies the constraints $\mathcal{F}_i \mathbf{v}_h = 0$ for $i = 1, \dots, s-1$, the function $\mathbf{v}_h$ admits a regular decomposition like (3.7) and the resulting functions satisfy the conditions and estimates described in Theorem 3.3.*

4 Basic regular decompositions

In this section we develop regular decompositions for vector-valued functions that have zero tangential trace on a part of the boundary of a Lipschitz polyhedron.

A regular decomposition is called **curl**-bounded if the H^1 norms of the two H^1 functions defined by the decomposition can be controlled by the **curl** semi-norm of the original vector-valued function; a regular decomposition is called L^2 -stable if the L^2 norm of the vector-valued H^1 function and the H^1 norm of the scalar H^1 function can be controlled by the L^2 norm of the original vector-valued function.

In this section we first build **curl**-bounded regular decompositions on a class of Lipschitz polyhedrons and then develop **curl**-bounded and L^2 -stable regular decompositions on a class of non-convex polyhedrons .

4.1 Curl-bounded regular decompositions on Lipschitz polyhedrons

Let G be a bounded Lipschitz polyhedron with a diameter $O(1)$ and vanishing first Betti number. Assume that G is a union of tetrahedral elements, associated with a triangulation $\mathcal{T}_h$ ($h \ll 1$), with h being the mesh width. For each r ($1 \leq r \leq J$), let $\Gamma_r \subset \partial G$ be a connected union of faces of some elements of $\mathcal{T}_h$ (see Figure 2).

Assumption 4.1. For each Γ_r , there exists a bounded Lipschitz domain D_r with a diameter $O(1)$ and vanishing first Betti number, such that (i) $\bar{D}_r \cap \bar{G} = \Gamma_r$; (ii) $\text{dist}(D_i, D_j) \geq \delta$ for any $i \neq j$, where δ is a positive constant; (iii) $B = G \cup (\cup_{r=1}^J \bar{D}_r)$ is a Lipschitz domain with vanishing first Betti number.

The condition (ii) implies that $\text{dist}(\Gamma_i, \Gamma_j) \geq \delta$ for any $i \neq j$. This assumption can be viewed as an extension of the assumption given in [18] and is essentially equivalent to the one introduced in [16], but it seems more direct. Before presenting the main result of this subsection, we would like to give two examples satisfying Assumption 4.1.




Figure 2: The cross-section of the domains described in Assumption 4.1, where the red parts denote the connected components of Γ , the blue parts denote the auxiliary subdomains.

Example 4.1. Let domain G be a usual polyhedron with fixed number of (plane) faces only, and let each Γ_r be a complete face or a connected union of several complete faces. Assume that $\bar{\Gamma}_i \cap \bar{\Gamma}_j = \emptyset$ for $i \neq j$.

We will explain that Assumption 4.1 is met as long as each Γ_r is a *connected “Lipschitz” union of some faces*, which means that $\partial\Gamma_r$ is connected and closed but is not self intersecting. We only need to define a suitable domain D_r for each Γ_r . It is easy to see that $\partial\Gamma_r$ can

be written as $\partial\Gamma_r = \cup_{l=1}^{n_r} E_l^{(r)}$, where $E_l^{(r)}$ is an edge of some face $F_l^{(r)} \subset \Gamma_r$ ($l = 1, \dots, n_r$). For each $E_l^{(r)}$, we choose a half plane $\pi_l^{(r)}$ such that $\pi_l^{(r)} \cap \bar{G} = E_l^{(r)}$ and the interfacial angle between $\pi_l^{(r)}$ and the face $F_l^{(r)}$ is an acute angle (this means that the boundary line of $\pi_l^{(r)}$ contains $E_l^{(r)}$ and $\pi_l^{(r)}$ extends to the exterior of G). For a convex polyhedron G , $\pi_l^{(r)}$ can be simply chosen as the extending half plane of the face that has the common edge $E_l^{(r)}$ with $F_l^{(r)}$ (see the left graph in Figure 3). If Γ_r and all the half planes $\pi_1^{(r)}, \dots, \pi_{n_r}^{(r)}$ just surround a polyhedron on one side of G , then this polyhedron can be chosen as D_r and it satisfies $\bar{D}_r \cap \bar{G} = \Gamma_r$. Otherwise, we can add one (or several) suitable plane π^* such that Γ_r, π^* and all the half planes $\pi_1^{(r)}, \dots, \pi_{n_r}^{(r)}$ surround a polyhedron on one side of G , and we choose this polyhedron as D_r . It is easy to see that Assumption 4.1 is met for this choice of D_r . In particular, when G is convex or Γ contains all the concave parts of ∂G (its precise definition will be given later), the resulting extended domain B is convex.

However, if some Γ_r is a connected “non-Lipschitz” union of some faces of G (for example, G is a four-sided pyramid and Γ_r is a union of two opposite lateral faces of G , see the right graph of Figure 1), then $\partial\Gamma_r$ is self intersecting and so there does not exist a Lipschitz domain D_r satisfying Assumption 4.1.




Figure 3: In the left graph, G denote a union of the cube and the four-sided pyramid, Γ_r denotes a union of two neighboring triangle faces of the four-sided pyramid, the polyhedron D_r is generated by four half planes that extend to the exterior of G from the red edges $E_1^{(r)}, \dots, E_4^{(r)}$ (the blue triangle face of D_r is a part of the half plane $\pi_1^{(r)}$ that contains the edge $E_1^{(r)}$ and extends above). In the right graph, G is generated from a cube domain Ω in the manner described in Example 4.2, D_r is the Lipschitz polyhedron at the upper-left corner and with yellow edges (the tetrahedron with purple edges denotes the cut domain $\hat{D}_r$, π denote the cut plane), the global green polyhedron is a union of G and one D_r .

Example 4.2: For a usual polyhedron Ω , let $\Omega^\partial \subset \Omega$ be the polyhedron generated by cutting its convex angles with some planes, such that every cut domain $\hat{D}_r$ has much larger size than h and has a suitable distance from the other cut domains. Notice that the polyhedron Ω^∂ may be not a union of elements. Let G be a union of all the elements, each of which has at least one vertex in Ω^∂ (see the right graph in Figure 3). Then G is a Lipschitz polyhedron, which can be regarded as a small perturbation of the usual polyhedron Ω^∂ . Let D_r be a union of all the elements contained in $\hat{D}_r$ and set $\Gamma_r = \bar{G} \cap \bar{D}_r$. We define $B = G \cup (\cup_{l=1}^J \bar{D}_r) = \Omega$. It is clear that Assumption 4.1 is met for this example.

Set $\Gamma = \cup_{l=1}^J \Gamma_l$. We use the same notations introduced in Section 2. The following result on regular decompositions can be found in [16].

Lemma 4.1 *Let Assumption 4.1 be satisfied. For any $\mathbf{v} \in H_\Gamma(\mathbf{curl}; G)$, there exists a vector-valued function $\Phi \in (H_\Gamma^1(G))^3$ and a scalar function $p \in H_\Gamma^1(G)$ such that*

$$\mathbf{v} = \Phi + \nabla p. \quad (4.1)$$

Moreover, we have

$$\|\Phi\|_{1,G} \lesssim \|\mathbf{v}\|_{\mathbf{curl},G} \quad \text{and} \quad \|\Phi\|_{0,G} + \|p\|_{1,G} \lesssim \|\mathbf{v}\|_{0,G}. \quad (4.2)$$

In particular, if $\mathbf{v} = \mathbf{v}_h \in V_h(G) \cap H_\Gamma(\mathbf{curl}; G)$, then there exist $p_h \in Z_h(G) \cap H_\Gamma^1(G)$, $\Phi_h \in (Z_h(G) \cap H_\Gamma^1(G))^3$ and $\mathbf{R}_h \in V_h(G) \cap H_\Gamma(\mathbf{curl}; G)$ such that

$$\mathbf{v}_h = \mathbf{r}_h \Phi_h + \nabla p_h + \mathbf{R}_h, \quad (4.3)$$

where $\mathbf{r}_h$ denotes the edge element projector $\mathbf{R}_D^1$ introduced in [16]. Moreover, they satisfy the stabilities

$$\|\Phi_h\|_{1,G} + h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \|\mathbf{v}\|_{\mathbf{curl},G} \quad \text{and} \quad \|\Phi_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \|\mathbf{v}\|_{0,G}. \quad (4.4)$$

‡

In the following lemma, we present **curl**-bounded regular decompositions, which have different stability estimates from that in Lemma 4.1.

Lemma 4.2 *Let $J = 1$, i.e., Γ be a connected union of closed faces of elements, and let Assumption 4.1 be satisfied. For any $\mathbf{v} \in H_\Gamma(\mathbf{curl}; G)$, there exists a vector-valued function $\Phi \in (H_\Gamma^1(G))^3$ and a scalar function $p \in H_\Gamma^1(G)$ satisfying (4.1), such that*

$$\|\Phi\|_{1,G} \lesssim \|\mathbf{curl} \mathbf{v}\|_{0,G} \quad \text{and} \quad \|p\|_{1,G} \lesssim \|\mathbf{v}\|_{\mathbf{curl},G}. \quad (4.5)$$

When $\mathbf{v} = \mathbf{v}_h \in V_h(G) \cap H_\Gamma(\mathbf{curl}; G)$, there exist $p_h \in Z_h(G) \cap H_\Gamma^1(G)$, $\Phi_h \in (Z_h(G) \cap H_\Gamma^1(G))^3$ and $\mathbf{R}_h \in V_h(G) \cap H_\Gamma(\mathbf{curl}; G)$, such that they satisfy the decomposition (4.3) with $\mathbf{r}_h$ being the standard edge element interpolation, and they have the stability estimates

$$\|\Phi_h\|_{1,G} + h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \|\mathbf{curl} \mathbf{v}_h\|_{0,G} \quad \text{and} \quad \|p_h\|_{1,G} \lesssim \|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (4.6)$$

In particular, if the extended domain B is convex, the second inequality in (4.5) (and (4.6)) can be replaced by

$$\|\Phi\|_{0,G} + \|p\|_{1,G} \lesssim \|\mathbf{v}\|_{0,G} \quad \text{and} \quad \|\Phi_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \|\mathbf{v}_h\|_{0,G}. \quad (4.7)$$

Proof. The basic ideas follow the arguments in [11, 29], here the proof seems more direct. When $J = 1$, we have $\Gamma = \Gamma_1$ and $D = D_1$. Let B be the extended domain defined in Assumption 4.1. We extend $\mathbf{v}$ onto the global B by zero, i.e., the extension $\tilde{\mathbf{v}}$ satisfying $\tilde{\mathbf{v}} = \mathbf{v}$ on G and $\tilde{\mathbf{v}} = \mathbf{0}$ on $\bar{D}$. Since $\mathbf{v} \times \mathbf{n} = \mathbf{0}$ on Γ , we have $\tilde{\mathbf{v}} \in H(\mathbf{curl}; B)$. By the regular decomposition (Lemma 2.4 in [15]), we get²

$$\tilde{\mathbf{v}} = \mathbf{w} + \nabla \varphi \quad \text{on } B, \quad (4.8)$$

²Let $\mathbf{N}_\mathcal{H}$ and $\mathbf{R}$ be the mappings defined in Lemma 2.4 of [15]. We define $\mathbf{w} = \mathbf{R}\tilde{\mathbf{v}}$. Since the first Betti number of B vanishes and so we have $\mathbf{N}_\mathcal{H}\tilde{\mathbf{v}} = \mathbf{0}$. The property $\mathbf{R}|_{\mathbf{H}(\mathbf{curl}; B)} = \mathbf{0}$ means that $\|\mathbf{w}\|_{1,B} \lesssim \|\mathbf{curl} \tilde{\mathbf{v}}\|_{0,B}$.

with $\mathbf{w} \in (H^1(B))^3$ and $\varphi \in H^1(B)$. Moreover, $\mathbf{w}$ and φ satisfy

$$\|\mathbf{w}\|_{1,B} \lesssim \|\mathbf{curl} \tilde{\mathbf{v}}\|_{0,B} = \|\mathbf{curl} \mathbf{v}\|_{0,G} \quad \text{and} \quad \|\nabla \varphi\|_{0,B} \lesssim \|\tilde{\mathbf{v}}\|_{\mathbf{curl},B} = \|\mathbf{v}\|_{\mathbf{curl},G}. \quad (4.9)$$

Here we require that φ has zero average value on D (not on B), which is written as $\gamma_D(\varphi) = 0$. When B is convex, by the orthogonal Helmholtz decomposition in [12], we have

$$\|\mathbf{w}\|_{1,B} \lesssim \|\mathbf{curl} \mathbf{v}\|_{0,G} \quad \text{and} \quad \|\mathbf{w}\|_{0,B} + \|\nabla \varphi\|_{0,B} \lesssim \|\mathbf{v}\|_{0,G}. \quad (4.10)$$

Notice that $\tilde{\mathbf{v}} = \mathbf{0}$ on $\bar{D}$, we have $\nabla \varphi = -\mathbf{w}$ on D , and so $\varphi \in H^2(D)$. Let $\tilde{\varphi} \in H^2(B)$ be a stable extension of φ from D onto the global B . It follows by (4.8) that

$$\tilde{\mathbf{v}} = (\mathbf{w} + \nabla \tilde{\varphi}) + \nabla(\varphi - \tilde{\varphi}) \quad \text{on } B. \quad (4.11)$$

Define $\Phi = \mathbf{w} + \nabla \tilde{\varphi}$ and $p = \varphi - \tilde{\varphi}$. Then we have $\Phi \in (H^1(B))^3$ and $p \in H^1(B)$, and they satisfy (4.1). Since $\tilde{\varphi} = \varphi$ on $\bar{D}$, we have $p = 0$ on $\bar{D}$, which means that $p \in H_\Gamma^1(G)$ and $\nabla p = \mathbf{0}$ on $\bar{D}$. Therefore $\Phi = \mathbf{0}$ on $\bar{D}$ and so $\Phi = \mathbf{0}$ on Γ , which implies that $\Phi \in (H_\Gamma^1(G))^3$.

By the definition of Φ and the first inequality in (4.9), together with Friedrichs inequality and the relation $\nabla \varphi = -\mathbf{w}$ on D , we get (notice that D is simply-connected, so Friedrichs inequality can be used on D)

$$\begin{aligned} \|\Phi\|_{1,G} &\leq \|\mathbf{w}\|_{1,G} + \|\tilde{\varphi}\|_{2,G} \lesssim \|\mathbf{curl} \mathbf{v}\|_{0,G} + \|\varphi\|_{2,D} \\ &\lesssim \|\mathbf{curl} \mathbf{v}\|_{0,G} + \|\nabla \varphi\|_{1,D} \lesssim \|\mathbf{curl} \mathbf{v}\|_{0,G} + \|\mathbf{w}\|_{1,D} \\ &\lesssim \|\mathbf{curl} \mathbf{v}\|_{0,G} + \|\mathbf{w}\|_{1,B} \lesssim \|\mathbf{curl} \mathbf{v}\|_{0,G}. \end{aligned} \quad (4.12)$$

We can further obtain the second inequality of (4.5) by (4.1) since $p = 0$ on Γ .

Let $\mathbf{Q}_h : (H_\Gamma^1(G))^3 \rightarrow (Z_h(G) \cap H_\Gamma^1(G))^3$ denote the Scott-Zhang interpolation (see [31]), which can preserve Dirichlet zero boundary condition, and define $\Phi_h = \mathbf{Q}_h \Phi$. In addition, let $\mathbf{r}_h$ be the standard edge element interpolation operator. By Lemma 4.7 in [5] (notice that $\mathbf{curl} \Phi = \mathbf{curl} \mathbf{v}_h$), we know that the interpolation functions $\mathbf{r}_h \Phi$ and $\mathbf{r}_h \Phi_h$ are well defined. Define

$$\mathbf{R}_h = \mathbf{r}_h(\Phi - \Phi_h) = (\mathbf{I} - \mathbf{Q}_h)\Phi + (\mathbf{r}_h - \mathbf{I})(\Phi - \Phi_h).$$

Then

$$\mathbf{v}_h = \mathbf{r}_h \Phi + \mathbf{r}_h(\nabla p) = \mathbf{r}_h \Phi + \nabla p_h = \mathbf{r}_h \Phi_h + \nabla p_h + \mathbf{R}_h \quad (4.13)$$

with $p_h \in (Z_h(G) \cap H_\Gamma^1(G))^3$. By the H^1 stability of the operator $\mathbf{Q}_h$ and the first estimate in (4.5), we directly get

$$\|\Phi_h\|_{1,G} \lesssim \|\Phi\|_{1,G} \lesssim \|\mathbf{curl} \mathbf{v}_h\|_{0,G}. \quad (4.14)$$

From the proof of Lemma 4.3 in [23], we can see that $(\mathbf{curl}(\Phi - \Phi_h) = \mathbf{curl} \mathbf{v}_h - \mathbf{curl} \Phi_h$ is a piecewise constant vector)

$$\|(\mathbf{r}_h - \mathbf{I})(\Phi - \Phi_h)\|_{0,G} \lesssim h(|\Phi - \Phi_h|_{1,G} + \|\mathbf{curl}(\Phi - \Phi_h)\|_{0,G}) \lesssim h(|\Phi|_{1,G} + |\Phi_h|_{1,G}).$$

This, together with the first inequalities in (4.5)-(4.6), yields

$$\|(\mathbf{r}_h - \mathbf{I})(\Phi - \Phi_h)\|_{0,G} \lesssim h\|\mathbf{curl} \mathbf{v}_h\|_{0,G}.$$

Then, by the L^2 approximation estimates for $\mathbf{Q}_h$ (see Theorem 4.1 of [31]), we further obtain

$$\begin{aligned} \|\mathbf{R}_h\|_{0,G} &\leq \|(\mathbf{I} - \mathbf{Q}_h)\Phi\|_{0,G} + \|(\mathbf{r}_h - \mathbf{I})(\Phi - \Phi_h)\|_{0,G} \\ &\lesssim h|\Phi|_{1,G} + h\|\mathbf{curl} \mathbf{v}_h\|_{0,G} \lesssim h\|\mathbf{curl} \mathbf{v}_h\|_{0,G}. \end{aligned}$$

This, together with (4.14), gives the first inequality in (4.6). The second inequality in (4.6) can be derived immediately by (4.13).

If B is convex, then we can use the second inequality in (4.10) to verify the first estimate in (4.7) in an analogous way with (4.12). Using the L^2 approximation of $\mathbf{Q}_h$ and the first estimate in (4.7) yields

$$\|\Phi_h\|_{0,G} \lesssim \|\Phi\|_{0,G} + h\|\Phi\|_{1,G} \lesssim \|\Phi\|_{0,G} + h\|\mathbf{curl} \mathbf{v}_h\|_{1,G} \lesssim \|\Phi\|_{0,G} + \|\mathbf{v}_h\|_{0,G} \lesssim \|\mathbf{v}_h\|_{0,G}. \quad (4.15)$$

Similarly, using the equality (4.13) and the L^2 approximation estimates for $\mathbf{r}_h$, together with the first estimate in (4.7), leads to (notice that $\mathbf{curl} \Phi = \mathbf{curl} \mathbf{v}_h$)

$$\begin{aligned} \|\nabla p_h\|_{0,G} &\leq \|\mathbf{v}_h\|_{0,G} + \|\mathbf{r}_h \Phi\|_{0,G} \lesssim \|\mathbf{v}_h\|_{0,G} + \|\Phi\|_{0,G} + \|(\mathbf{r}_h - \mathbf{I})\Phi\|_{0,G} \\ &\lesssim \|\mathbf{v}_h\|_{0,G} + h(\|\Phi\|_{1,G} + \|\mathbf{curl} \Phi\|_{0,G}) \\ &\lesssim \|\mathbf{v}_h\|_{0,G} + h\|\mathbf{curl} \mathbf{v}_h\|_{0,G} \lesssim \|\mathbf{v}_h\|_{0,G}. \end{aligned}$$

This, together with (4.15), gives the second inequality in (4.7).

We would like to emphasize that all the constants omitted in the proof are independent of h , since the involved domains have Lipschitz constants independent of h . $\sharp$

Remark 4.1 *It can be seen from Lemma 4.1 and Lemma 4.2 that the functions defined by different discrete regular decompositions may have different stabilities (since the functions are constructed in different manners). In general, the $\mathbf{curl}$ -bounded stability*

$$\|\Phi_h\|_{1,G} + h^{-1}\|\mathbf{R}_h\|_{0,G} \lesssim \|\mathbf{curl} \mathbf{v}_h\|_{0,G} \quad (4.16)$$

and the L^2 stability

$$\|\Phi_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \|\mathbf{v}_h\|_{0,G} \quad (4.17)$$

cannot be satisfied simultaneously. The inequality (4.16) implies that a $\mathbf{curl}$ -free vector-valued function must be the gradient of some scalar H^1 function. Lemma 4.2 gives some conditions that can guarantee both the estimates to be held. It is clear that, when $\Gamma = \partial G$ (which is connected), the conditions are met, where the extended domain B can be chosen as a ball (for this case, the result has essentially been built in [29]). Other situations satisfying the conditions was described in Example 4.1. As we will see in [19], when higher order term in the considered Maxwell equations is dominant, the semi-norm controlled estimate (4.16) will play a key role in the analysis. Unfortunately, if Γ is non-connected (i.e., $J \geq 2$), this estimate is false (a counterexample can be constructed). In fact, the domain D is non-connected for this case, so Friedrichs inequality cannot be used to prove (4.12). By the way, if we require that the function φ satisfying (4.8) has zero average value on B , which is a natural idea, the semi-norm controlled estimate (4.12) cannot be obtained.

4.2 Curl-bounded and L^2 -stable regular decompositions on non-convex polyhedrons

In this subsection we establish slightly weaker results than (4.16)-(4.17) for a class of non-convex polyhedrons. To this end, we first give several basic auxiliary results.

Proposition 4.1. Let Ω be a usual polyhedron, and assume that $\mathbf{w}_h \in (Z_h(\Omega))^3$. Then we have $\mathbf{curl}(\mathbf{r}_h \mathbf{w}_h) = \mathbf{curl} \mathbf{w}_h$ and $\|\mathbf{r}_h \mathbf{w}_h\|_{0,\Omega} \lesssim \|\mathbf{w}_h\|_{0,\Omega}$.

Proof. Let $W_h(\Omega)$ denote the Raviart-Thomas finite element space of the lowest order,

and let Π_h be the interpolation operator into $W_h(\Omega)$. Since $\mathbf{w}_h \in (Z_h(\Omega))^3$, we have $\mathbf{curl} \mathbf{w}_h \in W_h(\alpha)$. Then

$$\mathbf{curl} (\mathbf{r}_h \mathbf{w}_h) = \Pi_h \mathbf{curl} \mathbf{w}_h = \mathbf{curl} \mathbf{w}_h.$$

The desired inequality can be derived by the approximation property of $\mathbf{r}_h$ and the inverse estimate of finite element functions. $\sharp$

The following two results come from the so called “edge” lemma and “face” lemma (cf. Lemma 4.16 and Lemma 4.24 in [34], or Lemma 4.9-Lemma 4.10 in [35]). Here the considered domain Ω has a diameter $O(1)$ and the space $H_{00}^{\frac{1}{2}}(\mathbb{F})$ is defined in the standard manner (cf. [35]).

Lemma 4.3 *Let Ω be a usual polyhedron, and $\mathbb{E}$ be an edge of Ω . Then, for $v_h \in Z_h(\Omega)$, we have*

$$\|v_h\|_{0,\mathbb{E}} \lesssim \log^{\frac{1}{2}}(1/h) \|v_h\|_{1,\Omega}. \quad (4.18)$$

Lemma 4.4 *Let Ω be a usual polyhedron, and $\mathbb{F}$ be a face of Ω . For $v_h \in Z_h(\Omega)$, define $I_{\mathbb{F}}^0 v_h \in V_h(\partial\Omega)$ such that $I_{\mathbb{F}}^0 v_h$ equals v_h at the nodes in the interior of $\mathbb{F}$ and vanishes at all the other nodes on $\partial\Omega$. Then*

$$\|I_{\mathbb{F}}^0 v_h\|_{H_{00}^{\frac{1}{2}}(\mathbb{F})} \lesssim \log(1/h) \|v_h\|_{1,\Omega}. \quad (4.19)$$

By Lemma 4.24 in [34], we can easily derive the following result

Lemma 4.5 *Let Ω be a usual polyhedron, and $\mathbb{F}$ be a face of Ω . Assume that $v_h \in Z_h(\Omega)$. Then there exists a function $v_h^{\mathbb{F}} \in Z_h(\Omega)$ such that $v_h^{\mathbb{F}}$ equals v_h at all the nodes in the interior of $\mathbb{F}$ and vanishes on $\partial\Omega \setminus \mathbb{F}$. Moreover, the function $v_h^{\mathbb{F}}$ satisfies the estimates*

$$\|v_h^{\mathbb{F}}\|_{1,\Omega} \lesssim \log(1/h) \|v_h\|_{1,\Omega}. \quad (4.20)$$

and

$$\|v_h^{\mathbb{F}}\|_{0,\Omega} \lesssim \|v_h\|_{0,\Omega}. \quad (4.21)$$

The inequality (4.20) is included in the results of Lemma 4.24 in [34], and the inequality (4.21) can be directly obtained by the definition of the function $v_h^{\mathbb{F}}$, which depends on the weight function $\vartheta_{\mathbb{F}}$ constructed in Lemma 3.3.6 of [8] and Lemma 4.23 of [34].

Let G be a usual non-convex polyhedron, which is a non-overlapping union of several convex polyhedra $G_1, \dots, G_{m_0}$. As usual, we assume that each G_r is a union of elements and the intersection of any two neighboring polyhedra G_i and G_j is either their common face, or their common edge, or their common vertex, i.e., the intersection cannot be only a part of one face or one edge of G_i or G_j .

We need to give precise definitions of convex “complete” face and “concave part” of ∂G . Let $\mathbb{F} \subset \partial G$ be a connected union of some triangles, and let π be the plane containing $\mathbb{F}$. When $\pi \cap \bar{G} = \pi \cap \partial G = \mathbb{F}$, we call $\mathbb{F}$ as a convex “complete” face of G . If $\pi \cap \bar{G} \neq \pi \cap \partial G$, then $\mathbb{F}$ is called a “concave part” of ∂G . In particular, when $\mathbb{F}$ is a convex “complete” face of some G_r , then $\mathbb{F}$ is called a concave “local” face of G (see Figure 4). A point $\mathbf{v} \in \partial G$ is called a concave vertex of G if $\mathbf{v}$ is a common vertex of several concave “local” faces of G ; similarly, an edge $\mathbb{E} \subset \partial G$ is called a concave edge of G if $\mathbb{E}$ is a common edge of several concave “local” faces of ∂G . For a concave vertex $\mathbf{v}$ (resp. concave edge $\mathbb{E}$) of G , the connected union of all concave “local” faces that contain $\mathbf{v}$ (resp. $\mathbb{E}$) as their common vertex (resp. common edge) is called a concave “complete” face of G .




Figure 4: G denotes the union of four cubes G_1, G_2, G_3, G_4 . The red square F is a convex “complete” face of G_1 but it is a concave “local” face of G , the purple point V is the unique concave vertex of G and the green edge E is a concave edge of G .

Let all the convex polyhedra $\{G_r\}_{r=1}^{m_0}$ be divided into two sets Σ_1 and Σ_2 . For convenience, set

$$\Lambda_k = \{r : G_r \in \Sigma_k\} \quad \text{and} \quad \Gamma_k = \bigcup_{r \in \Lambda_k} \partial G_r \quad (k = 1, 2).$$

Let Γ be a connected “Lipschitz” union of some complete faces. We assume that the sets Σ_1 and Σ_2 satisfy the conditions: (i) the set Σ_1 consists of polyhedrons that do not intersect each other, and the intersection of Γ with each polyhedron in Σ_1 is either an empty set or a connected union of faces of the polyhedron; (ii) the intersection of each polyhedron in Σ_2 with $\Gamma \cup \Gamma_1$ is a connected union of faces of the polyhedron; (iii) the intersection of any polyhedron G_l in Σ_2 with another polyhedron in Σ_2 is a subset of $G_l \cap (\Gamma \cup \Gamma_1)$ (which is a connected union of faces of G_l) but cannot be one face.

The following result gives a slightly weak L^2 stability of the regular decomposition on a class of non-convex polyhedra.

Theorem 4.1 *Let G be a usual non-convex polyhedron, and Γ be a connected “Lipschitz” union of some complete faces of G , with G and Γ satisfying the above assumptions. Assume that $\mathbf{v}_h \in V_h(G)$ satisfies $\mathbf{v}_h \times \mathbf{n} = \mathbf{0}$ on Γ . Then there exist $p_h \in Z_h(G) \cap H_\Gamma^1(G)$, $\mathbf{w}_h \in (Z_h(G) \cap H_\Gamma^1(G))^3$ and $\mathbf{R}_h \in V_h(G) \cap H_\Gamma(\mathbf{curl}; G)$ such that*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h, \quad (4.22)$$

with the following estimates

$$\|\mathbf{w}_h\|_{1,G} + h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G} \quad (4.23)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{0,G}. \quad (4.24)$$

In particular, when Γ contains all the “concave parts” of ∂G , the above logarithmic factor can be dropped.

Proof. When Γ contains all the “concave parts” of ∂G , the results directly follow from Lemma 4.2 and Example 4.1. We only need to consider the case that Γ does not contain the concave parts of ∂G . In order to make the ideas be understood more easily, we first assume that G is a union of three cubes: $G = D_1 \cup D_2 \cup D_3$ with $D_1 = [0, \frac{1}{2}]^3$, $D_2 = [\frac{1}{2}, 1] \times [0, \frac{1}{2}]^2$ and $D_3 = [0, \frac{1}{2}] \times [\frac{1}{2}, 1] \times [0, \frac{1}{2}]$ (see Figure 5). For this case, we can define $\Sigma_1 = \{D_1\}$ and $\Sigma_2 = \{D_2, D_3\}$.




Figure 5: A non-convex polyhedron composed of three cubes

We divide the proof into two steps.

Step 1. Build the desired decomposition.

We first build a decomposition of $\mathbf{v}_h$ on D_1 by Lemma 4.2. The function $\mathbf{v}_{h,1} = \mathbf{v}_h|_{D_1}$ admits the decomposition

$$\mathbf{v}_{h,1} = \nabla p_{h,1} + \mathbf{r}_h \mathbf{w}_{h,1} + \mathbf{R}_{h,1} \quad \text{on } D_1, \quad (4.25)$$

with $p_{h,1} \in Z_h(D_1)$, $\mathbf{w}_{h,1} \in (Z_h(D_1))^3$ and $\mathbf{R}_{h,1} \in V_h(D_1)$, which satisfy $p_{h,1} = 0$, $\mathbf{w}_{h,1} = \mathbf{0}$ and $\mathbf{R}_{h,1} \times \mathbf{n} = 0$ on $\partial D_1 \cap \Gamma$ (when $\partial D_1 \cap \Gamma = \emptyset$, we can require that $p_{h,1}$ has the zero average value on D_1). Moreover, since D_1 is convex (which implies the L^2 stability of the decomposition), we have

$$\|\mathbf{w}_{h,1}\|_{1,D_1} \lesssim \|\mathbf{curl} \mathbf{v}_{h,1}\|_{0,D_1}, \quad \|\mathbf{w}_{h,1}\|_{0,D_1} + \|p_{h,1}\|_{1,D_1} \lesssim \|\mathbf{v}_{h,1}\|_{0,D_1} \quad (4.26)$$

and

$$h^{-1} \|\mathbf{R}_{h,1}\|_{0,D_1} \lesssim \|\mathbf{curl} \mathbf{v}_{h,1}\|_{0,D_1}. \quad (4.27)$$

Secondly, we extend $\mathbf{w}_{h,1}$ and $p_{h,1}$ into D_2 and D_3 in a special manner such that some stability can be satisfied.

For $k = 2, 3$, set $F_{1k} = \partial D_1 \cap \partial D_k$. By Lemma 4.5, there exists a function $\mathbf{w}_{h,1}^{F_{1k}} \in (Z_h(D_1))^3$ such that $\mathbf{w}_{h,1}^{F_{1k}}$ equals $\mathbf{w}_{1k}$ at all the nodes in the interior of F_{1k} and $\mathbf{w}_{h,1}^{F_{1k}} = \mathbf{0}$ on $\partial D_1 \setminus F_{1k}$ ($k = 2, 3$). By the extension theorem and the Scott-Zhang interpolation [31], we can show (refer to the proof of Lemma 4.5 in [25]) there exists an extension $\tilde{\mathbf{w}}_{h,1}^{F_{1k}} \in (Z_h(G))^3$ such that $\tilde{\mathbf{w}}_{h,1}^{F_{1k}} = \mathbf{w}_{h,1}^{F_{1k}}$ on D_1 , $\tilde{\mathbf{w}}_{h,1}^{F_{1k}}$ vanishes on $\partial D_k \setminus F_{1k}$ ($k = 2, 3$) and satisfies

$$\|\tilde{\mathbf{w}}_{h,1}^{F_{1k}}\|_{1,D_k} \lesssim \|\mathbf{w}_{h,1}^{F_{1k}}\|_{1,D_1} \quad \text{and} \quad \|\tilde{\mathbf{w}}_{h,1}^{F_{1k}}\|_{0,D_k} \lesssim \|\mathbf{w}_{h,1}^{F_{1k}}\|_{0,D_1} \quad (k = 2, 3). \quad (4.28)$$

Here we have used L^2 approximation of the quasi-interpolation and the inverse inequality to derive the L^2 stability (refer to (4.15)). Set $F^\partial = \partial F_{12} \cup \partial F_{13}$, and let $\tilde{\mathbf{w}}_{h,1}^\partial \in (Z_h(G))^3$ denote the natural zero extension of $\mathbf{w}_{h,1}|_{F^\partial}$. Define $\tilde{\mathbf{w}}_{h,1}$ as follows:

$$\tilde{\mathbf{w}}_{h,1} = \mathbf{w}_{h,1} \quad \text{on } D_1; \quad \tilde{\mathbf{w}}_{h,1} = \tilde{\mathbf{w}}_{h,1}^{F_{1k}} + \tilde{\mathbf{w}}_{h,1}^\partial|_{D_k} \quad \text{on } D_k \quad (k = 2, 3).$$

It is easy to see that $\tilde{\mathbf{w}}_{h,1} \in (Z_h(G))^3$.

We define an extension $\tilde{p}_{h,1} \in Z_h(G)$ as follows: $\tilde{p}_{h,1} = p_{h,1}$ on $\bar{D}_1$; $\tilde{p}_{h,1}$ vanishes at all the nodes in $\partial D_k \setminus \bar{F}_{1k}$ ($k = 2, 3$); $\tilde{p}_{h,1}$ is discrete harmonic in D_k ($k = 2, 3$). Let $\tilde{\mathbf{R}}_{h,1} \in V_h(G)$ be the natural zero extension of $\mathbf{R}_{h,1}$. For $k = 2, 3$, we define

$$\mathbf{v}_{h,k}^* = \mathbf{v}_h|_{D_k} - (\nabla \tilde{p}_{h,1} + \mathbf{r}_h \tilde{\mathbf{w}}_{h,1} + \tilde{\mathbf{R}}_{h,1})|_{D_k} \quad \text{on } D_k. \quad (4.29)$$

It is easy to see that $\mathbf{v}_{h,k}^* \times \mathbf{n} = \mathbf{0}$ on $\bar{F}_{1k} \cup (\partial D_k \cap \Gamma)$ ($k = 2, 3$).

Now we build the desired decomposition based on a regular decomposition of the function $\mathbf{v}_{h,k}^*$ ($k = 2, 3$) defined above.

It follows by Lemma 4.2 that the function $\mathbf{v}_{h,k}^*$ admits a decomposition

$$\mathbf{v}_{h,k}^* = \nabla p_{h,k}^* + \mathbf{r}_h \mathbf{w}_{h,k}^* + \mathbf{R}_{h,k}^* \quad \text{on } D_k \quad (k = 2, 3) \quad (4.30)$$

with $p_{h,k}^* \in Z_h(D_k)$, $\mathbf{w}_{h,k}^* \in (Z_h(D_k))^3$ and $\mathbf{R}_{h,k}^* \in V_h(D_k)$ ($k = 2, 3$), which satisfy $p_{h,k}^* = 0$, $\mathbf{w}_{h,k}^* = \mathbf{0}$ and $\mathbf{R}_{h,k}^* \times \mathbf{n} = \mathbf{0}$ on $\bar{F}_{1k} \cup (\partial D_k \cap \Gamma)$ ($k = 2, 3$). Moreover, notice that D_k is a convex polyhedron (which implies the L^2 stability of the decomposition), for $k = 2, 3$ we have

$$\|\mathbf{w}_{h,k}^*\|_{1,D_k} \lesssim \|\mathbf{curl} \mathbf{v}_{h,k}^*\|_{0,D_k}, \quad \|\mathbf{w}_{h,k}^*\|_{0,D_k} + \|p_{h,k}^*\|_{1,D_k} \lesssim \|\mathbf{v}_{h,k}^*\|_{0,D_k} \quad (4.31)$$

and

$$h^{-1} \|\mathbf{R}_{h,k}^*\|_{0,D_k} \lesssim \|\mathbf{curl} \mathbf{v}_{h,k}^*\|_{0,D_k}. \quad (4.32)$$

Since $p_{h,k}^*$, $\mathbf{w}_{h,k}^*$ and $\mathbf{R}_{h,k}^*$ have the zero degrees of freedom on $\bar{F}_{1k}$, we can naturally extend them onto G by zero. We denote the resulting zero extensions by $\tilde{p}_{h,k}^*$, $\tilde{\mathbf{w}}_{h,k}^*$ and $\tilde{\mathbf{R}}_{h,k}^*$. Define

$$p_h = \tilde{p}_{h,1} + \sum_{k=2}^3 \tilde{p}_{h,k}^*, \quad \mathbf{w}_h = \tilde{\mathbf{w}}_{h,1} + \sum_{k=2}^3 \tilde{\mathbf{w}}_{h,k}^* \quad \text{and} \quad \mathbf{R}_h = \tilde{\mathbf{R}}_{h,1} + \sum_{k=2}^3 \tilde{\mathbf{R}}_{h,k}^*.$$

It is easy to see that p_h , $\mathbf{w}_h$ and $\mathbf{R}_h$ have the zero degrees of freedom on Γ . Using the local decompositions (4.25) and (4.30), together with the relation (4.29), we get the global decomposition of $\mathbf{v}_h$

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h. \quad (4.33)$$

Step 2. Derive the stability estimates.

From the definition of $\mathbf{w}_h$, we have

$$\|\mathbf{w}_h\|_{1,G} \lesssim \|\tilde{\mathbf{w}}_{h,1}\|_{1,G} + \sum_{k=2}^3 \|\tilde{\mathbf{w}}_{h,k}^*\|_{1,D_k}. \quad (4.34)$$

For $k = 2, 3$, by (4.31) and (4.29) we can deduce that

$$\|\tilde{\mathbf{w}}_{h,k}^*\|_{1,D_k} \lesssim \|\mathbf{curl} \mathbf{v}_{h,k}^*\|_{0,D_k} \lesssim \|\mathbf{curl} \mathbf{v}_h\|_{0,D_k} + \|\mathbf{curl}(\mathbf{r}_h \tilde{\mathbf{w}}_{h,1})\|_{0,D_k} + \|\mathbf{curl} \tilde{\mathbf{R}}_{h,1}\|_{0,D_k}.$$

Applying Proposition 4.1 and the inverse inequality, we further get

$$\begin{aligned} \|\tilde{\mathbf{w}}_{h,k}^*\|_{1,D_k} &\lesssim \|\mathbf{curl} \mathbf{v}_h\|_{0,D_k} + \|\mathbf{curl} \tilde{\mathbf{w}}_{h,1}\|_{0,D_k} + h^{-1} \|\tilde{\mathbf{R}}_{h,1}\|_{0,D_k} \\ &\lesssim \|\mathbf{curl} \mathbf{v}_h\|_{0,D_k} + \|\tilde{\mathbf{w}}_{h,1}\|_{1,D_k} + h^{-1} \|\mathbf{R}_{h,1}\|_{0,D_1}. \end{aligned} \quad (4.35)$$

Here we have used the relation $\|\tilde{\mathbf{R}}_{h,1}\|_{0,D_k} \lesssim \|\mathbf{R}_{h,1}\|_{0,D_1}$, which can be verified directly by the definition of $\tilde{\mathbf{R}}_{h,1}$. Substituting (4.35) into (4.34), and using (4.26)-(4.27) yields

$$\|\mathbf{w}_h\|_{1,G} \lesssim \|\mathbf{curl} \mathbf{v}_h\|_{0,G} + \sum_{k=2}^3 \|\tilde{\mathbf{w}}_{h,1}\|_{1,D_k}. \quad (4.36)$$

Similarly, we can show

$$\|\mathbf{w}_h\|_{0,G} \lesssim \|\mathbf{v}_h\|_{0,G} + \sum_{k=2}^3 \|\tilde{\mathbf{w}}_{h,1}\|_{0,D_k}. \quad (4.37)$$

It suffices to estimate $\|\tilde{\mathbf{w}}_{h,1}\|_{1,D_k}$ and $\|\tilde{\mathbf{w}}_{h,1}\|_{0,D_k}$ ($k = 2, 3$). It follows by Lemma 4.5 that

$$\|\mathbf{w}_{h,1}^{F_{1k}}\|_{1,D_1} \lesssim \log(1/h) \|\mathbf{w}_{h,1}\|_{1,D_1} \quad (k = 2, 3).$$

Combining this inequality with (4.28) leads to

$$\|\tilde{\mathbf{w}}_{h,1}^{F_{1k}}\|_{1,D_k} \lesssim \log(1/h) \|\mathbf{w}_{h,1}\|_{1,D_1}. \quad (4.38)$$

On the other hand, from Lemma 4.3, we have

$$\|\tilde{\mathbf{w}}_{h,1}^\partial\|_{1,D_k} \lesssim \|\mathbf{w}_{h,1}\|_{0,\mathbb{F}^\partial} \lesssim \log^{\frac{1}{2}}(1/h) \|\mathbf{w}_{h,1}\|_{1,D_1} \quad (k = 2, 3).$$

By the definition of $\tilde{\mathbf{w}}_{h,1}$, together with (4.38) and the above estimate, we deduce that

$$\|\tilde{\mathbf{w}}_{h,1}\|_{1,D_k} \lesssim \log(1/h) \|\mathbf{w}_{h,1}\|_{1,D_1} \quad (k = 2, 3).$$

Plugging this in (4.36) gives the first estimate of the inequality (4.23).

By Lemma 4.5 and the definition of $\tilde{\mathbf{w}}_{h,1}^\partial$, we get

$$\|\mathbf{w}_{h,1}^{F_{1k}}\|_{0,D_1} \lesssim \|\mathbf{w}_{h,1}\|_{0,D_1} \quad \text{and} \quad \|\tilde{\mathbf{w}}_{h,1}^\partial\|_{0,D_k} \lesssim \|\mathbf{w}_{h,1}\|_{0,D_1} \quad (k = 2, 3). \quad (4.39)$$

This, together with the second inequality in (4.28), leads to

$$\|\tilde{\mathbf{w}}_{h,1}\|_{0,D_k} \lesssim \|\mathbf{w}_{h,1}\|_{0,D_1} \quad (k = 2, 3). \quad (4.40)$$

Substituting this into (4.37) yields

$$\|\mathbf{w}_h\|_{0,G} \lesssim \|\mathbf{v}_h\|_{0,G}. \quad (4.41)$$

In the following we estimate $\|p_h\|_{1,G}$. It suffices to consider $\|\tilde{p}_{h,1}\|_{1,D_k}$ for $k = 2, 3$. Since $\tilde{p}_{h,1}$ is discrete harmonic in D_k , we have

$$\|\tilde{p}_{h,1}\|_{1,D_k} \lesssim \|\tilde{p}_{h,1}\|_{\frac{1}{2},\partial D_k} \quad (k = 2, 3). \quad (4.42)$$

Define the interpolation operators $I_{F_{1k}}^0$ and $I_{\partial F_{1k}}^0$ as follows: for $\psi_h \in Z_h(G)$, the function $I_{F_{1k}}^0 \psi_h$ (resp. $I_{\partial F_{1k}}^0 \psi_h$) equals ψ_h at the nodes in the interior of F_{1k} (resp. on ∂F_{1k}) and vanishes at all the other nodes on ∂D_k (resp. at all the nodes not on ∂F_{1k}). From the definition of $\tilde{p}_{h,1}$, we have $\tilde{p}_{h,1} = I_{F_{1k}}^0 \tilde{p}_{h,1} + I_{\partial F_{1k}}^0 \tilde{p}_{h,1}$ on ∂D_k . Then, it follows by (4.42) that

$$\|\tilde{p}_{h,1}\|_{1,D_k} \lesssim \|I_{F_{1k}}^0 \tilde{p}_{h,1}\|_{\frac{1}{2},\partial D_k} + \|I_{\partial F_{1k}}^0 \tilde{p}_{h,1}\|_{\frac{1}{2},\partial D_k}$$

$$\begin{aligned}
&\lesssim \|I_{F_{1k}}^0 \tilde{p}_{h,1}\|_{H_{00}^{\frac{1}{2}}(F_{1k})} + \|\tilde{p}_{h,1}\|_{0,\partial F_{1k}} \\
&= \|I_{F_{1k}}^0 p_{h,1}\|_{H_{00}^{\frac{1}{2}}(F_{1k})} + \|p_{h,1}\|_{0,\partial F_{1k}}.
\end{aligned}$$

Therefore, by using Lemma 4.4 and Lemma 4.3, we further obtain

$$\|\tilde{p}_{h,1}\|_{1,D_k} \lesssim \log(1/h) \|p_{h,1}\|_{1,D_1} \quad (k = 2, 3).$$

Then, as in the estimate for $\|\mathbf{w}_h\|_{1,G}$ (but (4.40) needs to be used), we get

$$\|p_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{0,G}.$$

Combining this with (4.41) gives the inequality (4.24). Moreover, we can similarly derive the second estimate in the inequality (4.23) by (4.27) and (4.32).

Next we extend the above ideas to the general case. Due to the condition (i), we can first independently define a regular decomposition of $\mathbf{v}_h$ on every convex polyhedron in Σ_1 and extend the functions defined by the regular decompositions to all the polyhedra in Σ_2 in the same manner considered above. Then, by the conditions (ii) and (iii), we separately build regular decompositions of the residual functions on all the polyhedra in Σ_2 . Finally we sum all the functions generated by these regular decompositions to get the desired regular decomposition. The condition (ii) makes Lemma 4.2 can be used for the residual functions on the polyhedra in Σ_2 . Moreover, the condition (iii) makes two possibly neighboring polyhedra (which have a common edge or a common vertex) in Σ_2 can be separately handled since the functions defined by regular decompositions for the residual functions vanish on the intersection of the two polyhedra. $\sharp$

Remark 4.2 *The construction of the decomposition (4.33) is a bit technical. The main difficulty comes from the definition of the vector-valued function $\mathbf{w}_h$, which must satisfy L^2 stability. A natural idea is to extend $\mathbf{w}_{h,1}$ onto G such that the extension is discrete harmonic in D_2 and D_3 , but the resulting extension may not satisfy the L^2 stability (4.40).*

5 Proof of Theorem 3.1

In this section we are devoted to the proof of Theorem 3.1. Throughout this section we always assume that G is a usual (simply-connected) polyhedron, and a face means a complete closed face of G . For convenience, we use the same notation $\mathbf{r}_h$ to denote the standard edge element interpolation operator (if a connected union of some faces and edges of G is involved) or the edge element projector $\mathbf{R}_D^1$ introduced in [16] (if a non-connected union of some faces and edges of G is involved). Of course, we can always choose $\mathbf{r}_h$ as the projector $\mathbf{R}_D^1$.

Let div_τ be the tangential divergence defined in [2], which was called surface *curl* in [33]. For $\mathbf{v}_h \in V_h(G)$, we have $\text{curl}_S \mathbf{v}_h = \text{div}_\tau(\mathbf{n} \times \mathbf{v}_h) = (\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}$; see [2]. For the purpose of clarity, we directly use the notation $(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}$ in the rest of this paper.

Let $\mathbf{E}_h : V_h(\partial G) \rightarrow V_h(G)$ be the discrete **curl**-harmonic extension operator defined in [1]. This following result can be seen from the proof (see (4.2)) of Theorem 2.3 in [1].

Proposition 5.1 (stability of the **curl**-harmonic extension) For any $\mathbf{v}_h \in V_h(G)$, we have

$$\|\mathbf{curl} (\mathbf{E}_h(\mathbf{v}_h \times \mathbf{n})|_{\partial G})\|_{0,G} \lesssim \|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2},\partial G} \quad (5.1)$$

and

$$\|\mathbf{E}_h(\mathbf{v}_h \times \mathbf{n})|_{\partial G}\|_{0,G} \lesssim \|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2},\partial G} + \|\mathbf{v}_h \times \mathbf{n}\|_{-\frac{1}{2},\partial G}. \quad (5.2)$$

In the rest of this paper, we will repeatedly use $H^{-\frac{1}{2}}$ -norm on a part of ∂G . For a connected “Lipschitz” union Γ of some faces of G , let $H^{-\frac{1}{2}}(\Gamma)$ denote the dual space of $H^{\frac{1}{2}}(\Gamma)$ and define the corresponding norm as

$$\|\mu\|_{-\frac{1}{2},\Gamma} = \sup_{v \in H^{\frac{1}{2}}(\Gamma)} \frac{\langle \mu, v \rangle}{\|v\|_{\frac{1}{2},\Gamma}}, \quad \mu \in H^{-\frac{1}{2}}(\Gamma).$$

Notice that $H^{-\frac{1}{2}}(\Gamma)$ is different from the dual space of $H_{00}^{\frac{1}{2}}(\Gamma)$, which is often denoted by $H_{00}^{-\frac{1}{2}}(\Gamma)$ (see, for example, Subsection 7.1 of [33]).

The following result can be directly built by the known $H^{-\frac{1}{2}}$ -stability of the face extension (cf. Lemma 6 in [20], Lemma 6.7 in [22] and Lemma 7.2 in [33])

Proposition 5.2 Let $\mathbf{v}_h \in V_h(G)$ and Γ be a connected “Lipschitz” union of some faces of G . Then

$$\|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2},\Gamma} \lesssim \log(1/h) \|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2},\partial G}. \quad (5.3)$$

Based on the above results we can build a discrete regular decomposition preserving zero tangential trace on the boundary of a face of G .

Lemma 5.1 Let F be a (closed) face of G . Assume that $\mathbf{v}_h \in V_h(G)$ satisfies $\mathbf{v}_h \cdot \mathbf{t}_{\partial F} = 0$ on ∂F . Then there exist $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$, which satisfy $p_h = 0$, $\mathbf{w}_h = \mathbf{0}$ and $\mathbf{R}_h \cdot \mathbf{t}_{\partial F} = 0$ on ∂F , such that

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h \quad (5.4)$$

with the following estimates

$$\|\mathbf{w}_h\|_{1,G} + h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G} \quad (5.5)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}\|_{\mathbf{curl},G}. \quad (5.6)$$

Proof. We separate the proof into two steps.

Step 1: Establish the desired decomposition.

Define $F_c = (\partial G \setminus F) \cup \partial F$, which is a connected “Lipschitz” union of faces. From the assumptions, we have a decomposition

$$\mathbf{v}_h = \mathbf{v}_h^F + \mathbf{v}_h^{F_c}, \quad (5.7)$$

where $\mathbf{v}_h^F \in V_h(G) \cap H_F(\mathbf{curl}; G)$ is discrete $\mathbf{curl}$ -harmonic on G , and $\mathbf{v}_h^{F_c} \in V_h(G) \cap H_{F_c}(\mathbf{curl}; G)$ (it may be not discrete $\mathbf{curl}$ -harmonic on G). For convenience, we use Γ to denote F or F_c . By Lemma 4.2, the function $\mathbf{v}_h^\Gamma$ ($\Gamma = F, F_c$) admits a discrete regular decomposition

$$\mathbf{v}_h^\Gamma = \mathbf{r}_h \Phi_h^\Gamma + \nabla p_h^\Gamma + \mathbf{R}_h^\Gamma \quad (\Gamma = F, F_c), \quad (5.8)$$

where $\Phi_h^\Gamma \in (Z_h(G) \cap H_\Gamma^1(G))^3$, $p_h^\Gamma \in Z_h(G) \cap H_\Gamma^1(G)$ and $\mathbf{R}_h^\Gamma \in V_h(G) \cap H_\Gamma(\mathbf{curl}; G)$ ($\Gamma = F, F_c$). Moreover, they possess the stability estimates

$$\|\Phi_h^\Gamma\|_{1,G} + h^{-1} \|\mathbf{R}_h^\Gamma\|_{0,G} \lesssim \|\mathbf{curl} \mathbf{v}_h^\Gamma\|_{0,G} \quad (\Gamma = F, F_c) \quad (5.9)$$

and

$$\|\Phi_h^\Gamma\|_{0,G} + \|p_h^\Gamma\|_{1,G} \lesssim \|\mathbf{v}_h^\Gamma\|_{\mathbf{curl},G} \quad (\Gamma = F, F_c). \quad (5.10)$$

Define

$$\mathbf{w}_h = \Phi_h^F + \Phi_h^{F^c}, \quad p_h = p_h^F + p_h^{F^c} \quad \text{and} \quad \mathbf{R}_h = \mathbf{R}_h^F + \mathbf{R}_h^{F^c}.$$

Then, from (5.7) and (5.8), these functions satisfy the regular decomposition (5.4). Moreover, the functions $\mathbf{w}_h$, p_h and $\mathbf{R}_h$ have zero degrees of freedom on ∂F since $\partial F = F \cap F_c$.

Step 2: Verify the desired estimate (5.5) for the decomposition (5.4).

From the definition of $\mathbf{w}_h$ and the triangle inequality, we have

$$\|\mathbf{w}_h\|_{1,G} \lesssim \|\Phi_h^F\|_{1,G} + \|\Phi_h^{F^c}\|_{1,G}.$$

This, along with (5.9), leads to

$$\|\mathbf{w}_h\|_{1,G} \lesssim \|\mathbf{curl} \mathbf{v}_h^F\|_{0,G} + \|\mathbf{curl} \mathbf{v}_h^{F^c}\|_{0,G}. \quad (5.11)$$

By the definition of $\mathbf{v}_h^F$ and using the stability (5.1) of $\mathbf{curl}$ -harmonic extension, we have

$$\|\mathbf{curl} \mathbf{v}_h^F\|_{0,G} \lesssim \|(\mathbf{curl} \mathbf{v}_h^F) \cdot \mathbf{n}\|_{-\frac{1}{2}, \partial G} \lesssim \|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2}, F_c}. \quad (5.12)$$

Moreover, using the estimate (5.3) and the trace inequality for $H(\mathbf{curl})$, we deduce

$$\|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2}, F_c} \lesssim \log(1/h) \|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2}, \partial G} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G}.$$

Substituting this into (5.12) yields

$$\|\mathbf{curl} \mathbf{v}_h^F\|_{0,G} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G}.$$

This, together with (5.7), further gives

$$\|\mathbf{curl} \mathbf{v}_h^{F^c}\|_{0,G} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G}.$$

Combining (5.11) with the above two inequalities, we obtain the first estimate of the inequality (5.5). The second estimate in (5.5) and the inequality (5.6) can be derived similarly (the stability (5.2) and the trace inequality need to be used). $\sharp$

Remark 5.1 *The conclusions in Lemma 5.1 are also valid when F is replaced by a connected “Lipschitz” union of some faces of G (the union must have one connected boundary, so F cannot be ∂G itself). But, if F is a non-connected union of some faces, then the semi-norm on the right side of (5.5) must be replaced by the complete norm.*

Remark 5.2 *Can the $\mathbf{curl}$ norm on the right side of (5.6) be replaced by the L^2 norm when the semi-norm on the right side of (5.5) is replaced by the complete norm (comparing Lemma 4.1 and Lemma 4.2)? Unfortunately, we failed to get the positive answer (notice that the $\mathbf{curl}$ norm on the right side of the trace inequality cannot be replaced by the L^2 norm).*

From the above proof, we can obtain the following result

Corollary 5.1. Let Γ be a non-connected union of $\Gamma_1, \dots, \Gamma_J$, where each Γ_j is *connected* “Lipschitz” union of some faces of G , and let F be a (closed) face of G satisfying $F \cap \Gamma = \emptyset$. Assume that $\mathbf{v}_h \in V_h(G)$ satisfies $\mathbf{v}_h \cdot \mathbf{t}_{\partial F} = 0$ on ∂F and $\mathbf{v}_h \times \mathbf{n} = \mathbf{0}$ on Γ . Then there exist $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$, which satisfy $p_h = 0$, $\mathbf{w}_h = \mathbf{0}$ on $\Gamma \cup \partial F$ and $\lambda_e(\mathbf{R}_h) = 0$ for any $e \subset \Gamma \cup \partial F$, such that

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h. \quad (5.13)$$

Moreover, we have the following estimates

$$\|\mathbf{w}_h\|_{1,G} + h^{-1}\|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h)\|\mathbf{v}_h\|_{\mathbf{curl},G} \quad (5.14)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h)\|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (5.15)$$

If $J = 1$ and $F \cap \Gamma$ is an edge of F , then the norm on the right side of (5.14) can be replaced by the **curl** semi-norm.

Proof. As in the proof of Lemma 5.1, we set $F_c = (\partial G \setminus F) \cup \partial F$ and use Lemma 4.1 for F_c and $\Gamma \cup F$ (since Assumption 4.1 is met for them), respectively. If $J = 1$ and $F \cap \Gamma$ is an edge of F , we can use Lemma 4.2 since $F \cup \Gamma$ is connected. $\sharp$

By Lemma 5.1 we can further build a regular decomposition preserving zero tangential component on one edge.

Lemma 5.2 *Let E be a (closed) edge of G , and $\mathbf{v}_h$ be a finite element function in $V_h(G)$ such that $\mathbf{v}_h \cdot \mathbf{t}_E = 0$ on E . Then $\mathbf{v}_h$ admits a decomposition*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h$$

for some $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$, which satisfy $p_h = 0$, $\mathbf{w}_h = \mathbf{0}$, $\mathbf{R}_h \cdot \mathbf{t}_E = 0$ on E . Moreover, the following estimates hold

$$\|\mathbf{w}_h\|_{1,G} + h^{-1}\|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h)\|\mathbf{curl} \mathbf{v}_h\|_{0,G} \quad (5.16)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h)\|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (5.17)$$

The conclusion is also valid for the case when E is replaced by a connected union of several edges of G .

Proof. We separate the proof into three steps.

Step 1: Establish an edge-related decomposition.

Let F be a face containing the edge E . We first consider a decomposition of the tangential component $\mathbf{v}_h \cdot \mathbf{t}_{\partial F}$ of $\mathbf{v}_h$ on ∂F . For convenience, we write $E_c = \partial F \setminus E$. Let s be the arc-length variable along E_c , taking values from 0 to l_0 , where l_0 is the total length of E_c . In terms of s , the function $\mathbf{v}_h \cdot \mathbf{t}_{E_c}$ is piecewise linear on the interval $[0, l_0]$, denoted by $\hat{v}(s)$. Then we define

$$C_E = \frac{1}{l_0} \int_0^{l_0} \hat{v}(s) ds \quad \text{and} \quad \phi_E(t) = \int_0^t (\hat{v}(s) - C_E) ds, \quad \forall t \in [0, l_0].$$

Clearly we see $\phi_E(t)$ vanishes at $t = 0$ and l_0 . We can extend ϕ_E naturally by zero onto E , then into ∂G and G such that its extension $\tilde{\phi}_E \in Z_h(G)$ and vanishes at all of the nodes that do not in the interior of E_c . In the following, we define an extension $\tilde{C}_E$ of C_E such that $\tilde{C}_E$ belongs to $(Z_h(G))^3$ and vanishes on E . Moreover, we require that $\tilde{C}_E$ satisfies $(\mathbf{r}_h \tilde{C}_E) \cdot \mathbf{t}_{\partial F} = C_E$ on $E_c = \partial F \setminus E$ and

$$\|\tilde{C}_E\|_{1,\hat{\Omega}} \lesssim |C_E|. \quad (5.18)$$

Let $\mathcal{N}_h^G$ denote the set of the nodes on G , and let $\mathcal{N}_h^{E_c}$ denote the set of the nodes in E_c . Then the values of the vector-valued function $\tilde{C}_E$ at all of the nodes in $\mathcal{N}_h^G \setminus \mathcal{N}_h^{E_c}$ are defined

to be zero. Moreover, the values of the vector-valued function $\tilde{C}_E$ at the nodes in $\mathcal{N}_h^{E_c}$ are defined such that $\tilde{C}_E$ is linear on each (coarse) edge on E_c , and $\|\tilde{C}_E\|_{0,E_c}^2$ reaches the minimal value under the constraint $(\mathbf{r}_h \tilde{C}_E) \cdot \mathbf{t}_{\partial F} = C_E$ on E_c . Notice that the number of degrees of freedom of the function $\tilde{C}_E$, which equals three times the number of vertices in E_c , is greater than the number of coarse edges contained in E_c . Then the minimization problem (with a quadric objective functional and compatible linear constraints) has a solution. In particular, if $C_E = 0$, the desired vector-valued function $\tilde{C}_E = \mathbf{0}$. Since each edge on E^c is of size $O(1)$, we have

$$\|\tilde{C}_E\|_{0,E_c} \lesssim |C_E|.$$

Moreover, by the definition of $\tilde{C}_E$ and the discrete norms, we get

$$\|\tilde{C}_E\|_{1,\hat{\Omega}} \lesssim \|\tilde{C}_E\|_{0,E_c}.$$

Thus the inequality (5.18) indeed holds. Using the definitions of $\tilde{\phi}_E$ and $\tilde{C}_E$, and noting that $\mathbf{v}_h \cdot \mathbf{t}_E = 0$, one can verify that (cf. [33])

$$\mathbf{v}_h \cdot \mathbf{t}_{\partial F} = (\nabla \tilde{\phi}_E) \cdot \mathbf{t}_{\partial F} + (\mathbf{r}_h \tilde{C}_E) \cdot \mathbf{t}_{\partial F}. \quad (5.19)$$

Moreover, by the discrete norms and Lemma 7.8 of [33], we can deduce that

$$\|\tilde{\phi}_E\|_{1,G} \lesssim \|\phi_E\|_{0,\partial F} \lesssim \log(1/h)(\|\mathbf{v}_h\|_{0,G} + \|\mathbf{curl} \mathbf{v}_h\|_{0,G}). \quad (5.20)$$

For the derivation of the second inequality in (5.20), we have used the following relations (see [33] for the details)

$$\|\phi_E\|_{0,\partial F} \lesssim \|\mathbf{v}_h \cdot \mathbf{t}_{\partial F} - C_E\|_{H^{-1}(\partial F)} \lesssim \|\mathbf{v}_h \times \mathbf{n}\|_{-\frac{1}{2},F} + \|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2},F}.$$

The logarithm in (5.20) essentially comes from the $H^{-\frac{1}{2}}$ -stability of the face extension of finite element functions.

Step 2: Construct the desired decomposition in Lemma 5.2.

For this purpose, we set

$$\hat{\mathbf{v}}_{h,E} = \mathbf{v}_h - (\nabla \tilde{\phi}_E + \mathbf{r}_h \tilde{C}_E). \quad (5.21)$$

From (5.19), we know that $\hat{\mathbf{v}}_{h,E} \cdot \mathbf{t}_{\partial F} = 0$ on ∂F . For the function $\hat{\mathbf{v}}_{h,E}$ defined in (5.21), by Lemma 5.1 one can find functions $\hat{p}_h \in Z_h(G)$, $\hat{\mathbf{w}}_h \in (Z_h(G))^3$ and $\hat{\mathbf{R}}_h \in V_h(G)$, which vanish on ∂F , such that

$$\hat{\mathbf{v}}_{h,E} = \nabla \hat{p}_h + \mathbf{r}_h \hat{\mathbf{w}}_h + \hat{\mathbf{R}}_h$$

with the following estimates

$$\|\hat{\mathbf{w}}_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{curl} \hat{\mathbf{v}}_{h,E}\|_{0,G}, \quad \|\hat{p}_h\|_{1,G} \lesssim \log(1/h) \|\hat{\mathbf{v}}_{h,E}\|_{0,G} \quad (5.22)$$

and

$$h^{-1} \|\hat{\mathbf{R}}_h\|_{0,\hat{\Omega}} \lesssim \log(1/h) \|\mathbf{curl} \hat{\mathbf{v}}_{h,E}\|_{0,\hat{\Omega}}. \quad (5.23)$$

Now by defining

$$p_h = \tilde{\phi}_E + \hat{p}_h, \quad \mathbf{w}_h = \tilde{C}_E + \hat{\mathbf{w}}_h \quad \text{and} \quad \mathbf{R}_h = \hat{\mathbf{R}}_h,$$

we get the final decomposition

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h \quad (5.24)$$

such that $p_h = 0$ and $\mathbf{w}_h = 0$ on E .

Step 3: Derive the desired estimate in Lemma 5.2 for the decomposition (5.24).

By the definition of C_E and noting that $\mathbf{v}_h \cdot \mathbf{t}_E = 0$ on E , which implies that $\mathbf{v}_h \cdot \mathbf{t}_{\partial F} = 0$ on E , we have

$$C_E = \frac{1}{l_0} \int_{\mathbf{t}_{E_c}} \mathbf{v}_h \cdot \mathbf{t}_{E_c} ds = \frac{1}{l_0} \int_{\partial F} \mathbf{v}_h \cdot \mathbf{t}_{\partial F} ds.$$

Thus, by the Stokes' theorem on F , we deduce

$$C_E = \frac{1}{l_0} \int_F (\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n} ds.$$

Then

$$|C_E| \lesssim \left| \int_F (\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n} ds \right| \lesssim \|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2}, F} \cdot \|1\|_{\frac{1}{2}, F}. \quad (5.25)$$

Using the estimate (5.3) again, we have

$$\|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2}, F} \lesssim \log(1/h) \|(\mathbf{curl} \mathbf{v}_h) \cdot \mathbf{n}\|_{-\frac{1}{2}, \partial G},$$

This, along with (5.25), leads to

$$|C_E| \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0, G}.$$

By (5.18), we further obtain

$$\|\tilde{C}_E\|_{1, \hat{\Omega}} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0, G}.$$

By the definition of $\mathbf{w}_h$ and the above inequality, together with (5.22), (5.21) and (5.20), we deduce the first estimate in the inequality (5.16)

In a similar way, we can prove the second estimate in (5.16) and the inequality (5.17) by (5.23).

When E is replaced by a connected union of some edges, the proof is almost the same as that given above, where the face F may be replaced by a connected ‘‘Lipschitz’’ union of several faces (the union has one connected boundary, so it cannot be ∂G itself). $\sharp$

Remark 5.3 *There is a key difference in the proof of the above lemma from that of Lemma 4.3 in [21]: since the extension $\tilde{C}_E$ in the above proof must belong to the space $(Z_h(G))^3$, $\tilde{C}_E$ cannot be defined to be the natural zero extension of C_E as in [21].*

From the explanations in Example 4.1, we know that Lemma 4.1 and Lemma 4.2 do not hold when some Γ_r is a connected ‘‘non-Lipschitz’’ union of some faces. The following result can be regarded as extensions of Lemma 4.1 and Lemma 4.2.

Lemma 5.3 *Let Γ be a connected union of some faces of G . Assume that $\mathbf{v}_h \in V_h(G)$ has zero tangential trace on Γ . Then the function $\mathbf{v}_h$ has a decomposition*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h$$

with some $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$ satisfying $p_h = 0$ and $\mathbf{w}_h = \mathbf{0}$ on Γ . Moreover, we have

$$\|\mathbf{w}_h\|_{1, G} + h^{-1} \|\mathbf{R}_h\|_{0, G} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0, G} \quad (5.26)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (5.27)$$

If Γ is a non-connected union of $\Gamma_1, \Gamma_2, \dots, \Gamma_J$, with each Γ_r being a connected union of some faces of G , then the semi-norm on the right side of (5.26) should be replaced by the complete **curl** norm.

Proof. Here we need not assume that Γ is a connected “Lipschitz” union of some faces, i.e., Γ may contain one isolated vertex: there exists one face on Γ such that the intersection of this face with any neighboring face on Γ is a vertex in Γ . For this case, Assumption 4.1 is not met and so Lemma 4.2 cannot be used directly. For simplicity of exposition, we only focus on the simplest case that $\Gamma \subset \partial G$ is a union of two faces that just have a common vertex. There is no essential difference in the proof for more general cases (see the explanations given later).

Let $\Gamma = F_1 \cup F_2$ with F_1 and F_2 being two closed faces of G , where the intersection of F_1 and F_2 is a vertex v (i.e., $v = F_1 \cap F_2$ is an isolated vertex). We use $E_i \subset F_i$ ($i = 1, 2$) to denote one of the closed edges containing v as their endpoint, such that $E = E_1 \cup E_2$ is on the same side of Γ . Then there exists a connected “Lipschitz” union $\hat{\Gamma}$ of some faces such that $E \subset \partial \hat{\Gamma}$ ($\hat{\Gamma}$ may be just one face), see Figure 6.

Since E is connected and $\mathbf{v}_h$ has zero tangential components on E by the assumption, we can define $\tilde{C}_E \in (Z_h(G))^3$ and $\tilde{\phi}_E \in Z_h(G)$ as in Lemma 5.2 (choosing $F = \hat{\Gamma}$). Define

$$\hat{\mathbf{v}}_{h,E} = \mathbf{v}_h - (\nabla \tilde{\phi}_E + \mathbf{r}_h \tilde{C}_E).$$




Figure 6: The blue faces denote F_1 and F_2 , which are two opposite lateral faces of four pyramid domain G , the red lateral face denotes $\hat{\Gamma}$.

Then $\hat{\mathbf{v}}_{h,E} \cdot \mathbf{t}_{\partial \hat{\Gamma}} = 0$ on $\partial \hat{\Gamma}$. Set $\hat{\Gamma}_c = (\partial G \setminus \hat{\Gamma}) \cup \partial \hat{\Gamma}$ and $\tilde{\Gamma} = \Gamma \cup \hat{\Gamma}$. It is easy to see that both $\hat{\Gamma}_c$ and $\tilde{\Gamma}$ are connected “Lipschitz” unions of some faces (i.e., any vertex of them is not an isolated vertex). By the assumptions, we have a decomposition

$$\hat{\mathbf{v}}_{h,E} = \hat{\mathbf{v}}_{h,E}^{\hat{\Gamma}_c} + \hat{\mathbf{v}}_{h,E}^{\tilde{\Gamma}},$$

where $\hat{\mathbf{v}}_{h,E}^{\hat{\Gamma}_c} \in V_h(G) \cap H_{\hat{\Gamma}_c}(\mathbf{curl}; G)$ is discrete **curl**-harmonic on G , and $\hat{\mathbf{v}}_{h,E}^{\tilde{\Gamma}} \in V_h(G) \cap H_{\tilde{\Gamma}}(\mathbf{curl}; G)$. Then we can build discrete regular decompositions respectively for the two

functions by Lemma 4.2 since Assumption 4.1 is met for $\hat{\Gamma}_c$ and $\tilde{\Gamma}$. The resulting functions have zero traces on $\Gamma = \hat{\Gamma}_c \cap \tilde{\Gamma}$. Finally, we can prove the desired results as in the proof of Lemma 5.2 and Lemma 5.1.

If Γ is a non-connected union of $\Gamma_1, \Gamma_2, \dots, \Gamma_J$, with some Γ_r being a union of two faces that just have a common vertex, we can handle Γ_r in the same manner as the one used above, and transform it into the cases that each considered part is connected “Lipschitz” union of some faces of G . Then we use Lemma 4.1 several times to get the desired results. $\sharp$

By Lemma 5.3 and the explanations in Example 4.1, the discrete results in Lemma 4.2 are essentially valid for any connected union Γ of faces of a polyhedron. With the help of Lemma 5.3, we need not to emphasize the word “Lipschitz” in *connected “Lipschitz” union* and simply say *connected union* in the following discussions.

Lemma 5.4 *Let Γ be a connected union of some faces and edges of G . Suppose that $\mathbf{v}_h$ satisfies Assumption 3.1. Then $\mathbf{v}_h$ can be decomposed as*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h \quad (5.28)$$

for some $p_h \in Z_h(G)$ and $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$ such that p_h and $\mathbf{w}_h$ vanish on Γ . Moreover, we have

$$\|\mathbf{w}_h\|_{1,G} + h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G} \quad (5.29)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (5.30)$$

The results are also valid when E is replaced by a connected union of several edges of G .

Proof. Let $E \subset \Gamma$ be an isolated edge (or a union of several edges) E of Γ , i.e., E is not an edge of any face in Γ . Since Γ is connected, there exists a face $F \subset \Gamma$ (provided that Γ indeed contains faces) such that the intersection $E \cap F$ is an endpoint of E . Let E' be an edge of the face F such that $E' \cap E$ is just this endpoint, which implies that $E' \cup E$ is connected. Let $\hat{\Gamma}$ be a connected “Lipschitz” union of some faces of G such that $E \cup E' \subset \partial \hat{\Gamma}$. By the assumptions, the function $\mathbf{v}_h$ has zero degrees of freedom on $E \cup E'$.

Without loss of generality, we assume that all of the faces in Γ constitute a connected “Lipschitz” union of some faces, otherwise, we need to add a union of some faces as in the proof of Lemma 5.3. Under this assumption, the set $\Gamma \cup \hat{\Gamma}$ is also a connected “Lipschitz” union of some faces. Then the results in this lemma can be built as in the proof of Lemma 5.2 (replacing E by $E \cup E'$), by using Lemma 4.2 on $\hat{\Gamma}_c = (\partial G \setminus \hat{\Gamma}) \cup \partial \hat{\Gamma}$ and $\Gamma \cup \hat{\Gamma}$, respectively. For this case, the norm on the right side of (5.29) can be replaced by the $\mathbf{curl}$ semi-norm. $\sharp$

Lemma 5.5 *Let $E_1, \dots, E_n$ be (closed) edges of G , which satisfy $E_i \cap E_j = \emptyset$ for $i \neq j$. Assume that $\mathbf{v}_h \in V_h(G)$ satisfies $\mathbf{v}_h \cdot \mathbf{t}_{E_l} = 0$ on each E_l . Then $\mathbf{v}_h$ can be decomposed as*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h$$

for some $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$ such that p_h and $\mathbf{w}_h$ vanish on each edge E_l . Moreover, we have

$$\|\mathbf{w}_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G}, \quad \|p_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G} \quad (5.31)$$

and

$$h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (5.32)$$

The results are also valid when E_l is replaced by a connected union of edges.

Proof. We first consider a simple case that, for each E_l , there exists a face $F_l \subset \partial G$ such that F_l contains E_l and the face F_l does not adjoin the other edges. For this case, regarding E_l as the edge E in the proof of Lemma 5.2 and almost repeating the proof process (but using Lemma 4.1 for $\Gamma = \cup_{l=1}^J F_l$), we can obtain the desired results.

If the above condition is not met, the proof of this lemma is a bit technical. Without loss of generality, we assume that this condition is not satisfied for each E_l (An example is that E_1, E_2, E_3, E_4 just are four parallel edges of a cube G , see Figure 7). This means that, for each edge E_l , any face containing E_l also contains another different edge $E_{l'}$ (which does not intersect E_l). In this situation, the above proof is not practical since the functions $\tilde{\phi}_E$ and $\tilde{C}_E$ defined in the proof of Lemma 5.2 may not vanish on $E_{l'}$.




Figure 7: An example with four edges: the left shaded polyhedron denotes the subdomain G'_1 , the middle polyhedron denotes G'_0 .

Since the considered edges $\{E_r\}$ do not intersect each other, we have $\text{dist}(E_i, E_j) = O(1)$ for $i \neq j$ (notice that G is a usual polyhedron with a diameter $O(1)$). Thus, we can decompose G into a union of non-overlapping subdomains $G'_0, G'_1, \dots, G'_m$ such that: (i) each subdomain G'_l is a polyhedron with a diameter $O(1)$; (ii) $\bar{G}'_l \cap \bar{G}'_j = \emptyset$ for $j \neq l$ ($l, j \neq 0$), and G'_0 just has a common face Γ'_{0l} with each G'_l ($l \neq 0$); (iii) for $l = 1, \dots, m$, the subdomain G'_l contains E_l as one of its edges, but the subdomain G'_0 does not intersect any E_l . In general we cannot require each subdomain G'_l to be a union of some elements. Because of this, for each l we choose another subdomain G_l generated from G'_l (with a small perturbation only), where G_l is a union of all the elements K satisfying $\text{meas}(K \cap G_l) \geq \frac{1}{2} \text{meas}(K)$, where G_l can be also generated as in Example 4.2. It is clear that G_l is not a usual polyhedron since $\Gamma_{0l} = \bar{G}_0 \cap \bar{G}_l$ is not a plane face yet. Fortunately, all the subdomains $\{G_l\}$ still constitute a union of G and preserve the other properties of $\{G'_l\}$.

For each E_l , we can use Lemma 5.2 to build a regular decomposition³

$$\mathbf{v}_h = \mathbf{r}_h \mathbf{w}_h^{(l)} + \nabla p_h^{(l)} + \mathbf{R}_h^{(l)} \quad \text{on } G_l \quad (5.33)$$

with $\mathbf{w}_h^{(l)} \in (Z_h(G_l))^3$ and $p_h^{(l)} \in Z_h(G_l)$, which vanish on E_l but may not vanish on the other edges. The decomposition is stable with a logarithmic factor. Let d_0 be a given positive number independent of h . For $l = 1, \dots, m$, we choose a ball D_l containing G_l such that $\text{dist}(\partial D_l, \partial G_l) \geq d_0$ and D_l does not intersect any G_j for $j \neq 0, l$. By the extension

³Regarding E_l as E in Lemma 5.2, we first get three finite element functions by the regular decomposition of $\mathbf{v}_h$ on G , then we define $\mathbf{w}_h^{(l)}$, $p_h^{(l)}$ and $\mathbf{R}_h^{(l)}$ as the restrictions of the three functions on G_l , respectively.

theorem, there exists an extension $\tilde{\mathbf{w}}^{(l)}$ (resp. $\tilde{p}^{(l)}$) of $\mathbf{w}_h^{(l)}|_{G_l}$ (resp. $p_h^{(l)}|_{G_l}$) such that: (a) $\tilde{\mathbf{w}}^{(l)} \in (H^1(\mathbb{R}^3))^3$ (resp. $\tilde{p}^{(l)} \in H^1(\mathbb{R}^3)$); (b) $\tilde{\mathbf{w}}^{(l)}$ and $\tilde{p}^{(l)}$ vanish on the outside of D_l ; (c) $\|\tilde{\mathbf{w}}^{(l)}\|_{1,G_0} \lesssim \|\mathbf{w}_h^{(l)}\|_{1,G_l}$ and $\|\tilde{p}^{(l)}\|_{1,G_0} \lesssim \|p_h^{(l)}\|_{1,G_l}$. For each l , let $\tilde{\mathbf{R}}_h^{(l)} \in V_h(G)$ denote the standard zero extension of $\mathbf{R}_h^{(l)}|_{G_l}$. Define

$$\tilde{\mathbf{v}}_h^{(0)} = \mathbf{v}_h - \sum_{l=1}^m (\mathbf{r}_h \tilde{\mathbf{w}}^{(l)} + \nabla \tilde{p}^{(l)} + \tilde{\mathbf{R}}_h^{(l)}) \quad \text{on } G_0. \quad (5.34)$$

It is clear that $\tilde{\mathbf{v}}_h^{(0)} \in H(\mathbf{curl}; G_0)$. Since $\tilde{\mathbf{w}}^{(l)}$, $\tilde{p}^{(l)}$ and $\tilde{\mathbf{R}}_h^{(l)}$ vanish on $\bar{G}_j$ for $j \neq 0, l$, by (5.33) we have $\tilde{\mathbf{v}}_h^{(0)} \times \mathbf{n} = \mathbf{0}$ on Γ_{0l} for $l = 1, \dots, m$. From the explanations in Example 4.2, we know that Assumption 4.1 is satisfied for G_0 . By Lemma 4.1, the function $\tilde{\mathbf{v}}_h^{(0)}$ admits a stable decomposition

$$\tilde{\mathbf{v}}_h^{(0)} = \tilde{\mathbf{w}}^{(0)} + \nabla \tilde{p}^{(0)} \quad \text{on } G_0 \quad (5.35)$$

for $\tilde{\mathbf{w}}^{(0)} \in (H^1(G_0))^3$ and $\tilde{p}^{(0)} \in H^1(G_0)$, with $\tilde{\mathbf{w}}^{(0)}$ and $\tilde{p}^{(0)}$ vanishing on Γ_{0l} for $l = 1, \dots, m$. Combining (5.34) and (5.35), we have

$$\mathbf{v}_h = \tilde{\mathbf{w}}^{(0)} + \nabla \tilde{p}^{(0)} + \sum_{l=1}^m (\mathbf{r}_h \tilde{\mathbf{w}}^{(l)} + \nabla \tilde{p}^{(l)} + \tilde{\mathbf{R}}_h^{(l)}) \quad \text{on } G_0.$$

Thus

$$\mathbf{v}_h = \mathbf{r}_h (\tilde{\mathbf{w}}^{(0)} + \sum_{l=1}^m \tilde{\mathbf{w}}^{(l)}) + \nabla p_h^{(0)} + \sum_{l=1}^m \tilde{\mathbf{R}}_h^{(l)} \quad \text{on } G_0 \quad (5.36)$$

with $p_h^{(0)}$ satisfying $\nabla p_h^{(0)} = \mathbf{r}_h \nabla (\tilde{p}^{(0)} + \sum_{l=1}^m \tilde{p}^{(l)})$.

Let $\Pi_h : (H^1(G_0))^3 \rightarrow (Z_h(G_0))^3$ denote the Scott-Zhang interpolation operator, which can preserve the values of a linear polynomial on some elements of the boundary ∂G_0 . Define

$$\mathbf{w}_h^{(0)} = \Pi_h(\tilde{\mathbf{w}}^{(0)} + \sum_{l=1}^m \tilde{\mathbf{w}}^{(l)}) \quad \text{and} \quad \mathbf{R}_h^{(0)} = \mathbf{r}_h (I - \Pi_h)(\tilde{\mathbf{w}}^{(0)} + \sum_{l=1}^m \tilde{\mathbf{w}}^{(l)}) + \sum_{l=1}^m \tilde{\mathbf{R}}_h^{(l)}.$$

Then (5.36) can be written as

$$\mathbf{v}_h = \mathbf{r}_h \mathbf{w}_h^{(0)} + \nabla p_h^{(0)} + \mathbf{R}_h^{(0)} \quad \text{on } G_0. \quad (5.37)$$

It is clear that $\mathbf{w}_h^{(0)} = \mathbf{w}_h^{(l)}$ and $p_h^{(0)} = p_h^{(l)}$ on Γ_{0l} for $l = 1, \dots, m$, which implies that $\mathbf{R}_h^{(0)} \times \mathbf{n} = \mathbf{R}_h^{(l)}$ on Γ_{0l} for $l = 1, \dots, m$. Thus we naturally define $\mathbf{w}_h = \mathbf{w}_h^{(l)}$, $p_h = p_h^{(l)}$ and $\mathbf{R}_h = \mathbf{R}_h^{(l)}$ on G_l for $l = 0, 1, \dots, m$, and we have $\mathbf{w}_h \in (Z_h(G))^3$, $p_h \in Z_h(G)$ and $\mathbf{R}_h \in V_h(G)$, which have zero degrees of freedom on all of the edges E_l . It is easy to see from (5.33) and (5.37) that the desired regular decomposition is valid for the defined functions. Besides, we can verify that the resulting regular decomposition is also stable with a logarithmic factor. $\sharp$

Remark 5.4 *In the proofs of Lemma 5.1-Lemma 5.5, our main ideas are to transform the problem vanishing on an edge E into a problem vanishing on a face F of G and then to use Lemma 4.1 or Lemma 4.2 to build a regular decomposition, which can preserve zero trace on this face.*

Notice that, in all the previous Lemmas, a connected union of several edges has no essential difference from an edge. Now we can prove Theorem 3.1 by Lemma 5.4 and Lemma 5.5.

Proof of Theorem 3.1. We decompose Γ into a union of $\Gamma_1, \Gamma_2, \dots, \Gamma_J$, where each Γ_r is a connected union of some faces and edges of G and any two different Γ_i and Γ_j do not intersect (i.e., $\Gamma_i \cap \Gamma_j = \emptyset$).

Since $\Gamma_i \cap \Gamma_j = \emptyset$ and G is a usual polyhedron with a diameter $O(1)$, we have $\text{dist}(\Gamma_i, \Gamma_j) = O(1)$ for $i \neq j$. As in the second part in the proof of Lemma 5.5, we can decompose G into a union of $G_0, G_1, G_2, \dots, G_J$ such that: (i) each subdomain G_l has the size $O(1)$ and is a union of some elements; (ii) $\bar{G}_l \cap \bar{G}_j = \emptyset$ for $j \neq l$ ($l, j \neq 0$), and G_0 just has a common face Γ_{0l} with each G_l ($l \neq 0$); (iii) for $l = 1, \dots, J$, $\Gamma_l \subset \partial G_l$, but the subdomain G_0 does not intersect any Γ_l . Here G_l may not be a usual polyhedron yet. For $l = 1, \dots, J$, we can build a stable discrete regular decomposition of $\mathbf{v}_h|_{G_l}$ as in Lemma 5.4 (detailed explanations can refer to the footnote in the proof of Lemma 5.5)

$$\mathbf{v}_h|_{G_l} = \mathbf{r}_h \mathbf{w}_h^{(l)} + \nabla p_h^{(l)} + \mathbf{R}_h^{(l)}$$

with $p_h^{(l)} \in Z_h(G_l)$ and $\mathbf{w}_h^{(l)} \in (Z_h(G_l))^3$ vanishing on Γ_l . We stably extend $\mathbf{w}_h^{(l)}$, $p_h^{(l)}$ and $\mathbf{R}_h^{(l)}$ into $\mathbb{R}^3$ (see the proof of Lemma 5.5), and define

$$\tilde{\mathbf{v}}_h^{(0)} = \mathbf{v}_h - \sum_{l=1}^J (\mathbf{r}_h \tilde{\mathbf{w}}^{(l)} + \nabla \tilde{p}^{(l)} + \tilde{\mathbf{R}}_h^{(l)}) \quad \text{on } G_0.$$

Then we can build a stable discrete regular decomposition of $\tilde{\mathbf{v}}_h^{(0)}$ on G_0 and furthermore build the desired regular decomposition (3.1) of $\mathbf{v}_h$ on G as in Lemma 5.5. $\sharp$

6 Proof of Theorem 3.2-Theorem 3.4

In this section we are devoted to the proof of Theorem 3.2-Theorem 3.4. We use the same notations as in Section 3 (for example, $\Gamma_i = \Gamma \cap \partial G_i$ for $i = 1, 2$).

Proof of Theorem 3.2. We recall that $\bar{G} = \bar{G}_1 \cup \bar{G}_2$, and $\bar{G}_1 \cap \bar{G}_2$ is the common edge E of G_1 and G_2 . We consider three different relations of position between E and Γ_i ($i = 1, 2$).

(i) $E \subset \Gamma_i$ for $i = 1, 2$. In this case, we have $E = \Gamma_1 \cap \Gamma_2$. We use Theorem 3.1 to build a regular decomposition of $\mathbf{v}_h|_{G_i}$ independently for $i = 1, 2$. Then the resulting functions $p_{h,i}$ and $\mathbf{w}_{h,i}$ vanish on Γ_i and so they also vanish on E . Thus we can directly extend $p_{h,i}$ and $\mathbf{w}_{h,i}$ onto the domain $G \setminus G_i$ by zero to get extension functions $\tilde{p}_{h,i}$ and $\tilde{\mathbf{w}}_{h,i}$ ($i = 1, 2$). Define $p_h = \tilde{p}_{h,1} + \tilde{p}_{h,2}$ and $\mathbf{w}_h = \tilde{\mathbf{w}}_{h,1} + \tilde{\mathbf{w}}_{h,2}$. By the two functions, we can build the regular decomposition (3.4) of $\mathbf{v}_h$ on the global G . In this case, there is no logarithmic factor in the stability estimates.

(ii) E is contained in only one of Γ_1 and Γ_2 , for example, E is contained in Γ_1 but is not contained in Γ_2 . We use Theorem 3.1 to build a regular decomposition of $\mathbf{v}_h|_{G_1}$ associated with Γ_1 , but use Theorem 3.1 to get a regular decomposition of $\mathbf{v}_h|_{G_2}$ associated with $\Gamma_2 \cup E$ (notice that $\mathbf{v}_h|_{G_2}$ vanishes on Γ_2 and E since $E \subset \Gamma_1$). It is clear that the resulting functions $p_{h,i}$ and $\mathbf{w}_{h,i}$ vanish on Γ_i and E . Then the desired regular decomposition can be built as in the above situation.

(iii) $E \cap \Gamma_i = \emptyset$ for $i = 1, 2$. In this case, we have $\Gamma_1 \cap \Gamma_2 = \emptyset$. If both Γ_1 and Γ_2 are non-empty sets, then Γ is non-connected; when Γ is connected, one of Γ_1 and Γ_2 is connected and the other is empty.

Without loss of generality, we assume that both Γ_1 and Γ_2 are non-empty sets. We first use Theorem 3.1 to build a regular decomposition of $\mathbf{v}_h|_{G_1}$

$$\mathbf{v}_h = p_{h,1} + \mathbf{r}_h \mathbf{w}_{h,1} + \mathbf{R}_{h,1} \quad \text{on } G_1 \quad (6.1)$$

with $p_{h,1} \in Z_h(G_1)$ and $\mathbf{w}_{h,1} \in (Z_h(G_1))^3$ vanishing on Γ_1 (and so $\mathbf{R}_{h,1}$ has zero degrees of freedom on Γ_1). Moreover, we have (if Γ_1 is connected and contains edges)

$$\|\mathbf{w}_{h,1}\|_{1,G_1} + h^{-1} \|\mathbf{R}_{h,1}\|_{0,G_1} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G_1} \quad (6.2)$$

and

$$\|\mathbf{w}_{h,1}\|_{0,G_1} + \|p_{h,1}\|_{1,G_1} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G_1}. \quad (6.3)$$

Notice that $p_{h,1}$, $\mathbf{w}_{h,1}$ and $\mathbf{R}_{h,1}$ may do not vanish on E since $E \cap \Gamma_1 = \emptyset$. We extend $p_{h,1}$, $\mathbf{w}_{h,1}$ and $\mathbf{R}_{h,1}$ onto G_2 such that the resulting extensions $\tilde{p}_{h,1} \in Z_h(G)$, $\tilde{\mathbf{w}}_{h,1} \in (Z_h(G))^3$ and $\tilde{\mathbf{R}}_{h,1} \in V_h(G)$ have zero degrees of freedom on the nodes or the edges in $G_2 \setminus E$. Define

$$\mathbf{v}_h^* = \mathbf{v}_h - (\tilde{p}_{h,1} + \mathbf{r}_h \tilde{\mathbf{w}}_{h,1} + \tilde{\mathbf{R}}_{h,1}) \quad \text{on } G. \quad (6.4)$$

Next we build a regular decomposition of $\mathbf{v}_h^*$. Since $(\tilde{p}_{h,1} + \mathbf{r}_h \tilde{\mathbf{w}}_{h,1} + \tilde{\mathbf{R}}_{h,1}) \times \mathbf{n} = \mathbf{v}_h \times \mathbf{n}$ on ∂G_1 , we have $\lambda_e(\mathbf{v}_h^*) = 0$ for $e \subset E \subset \partial G_1$. Moreover, noting that $\Gamma_2 \cap E = \emptyset$, we have $\lambda_e(\mathbf{v}_h^*) = \lambda_e(\mathbf{v}_h)$ for $e \subset \Gamma_2 \subset \partial G_2 \setminus E$, which implies that $\lambda_e(\mathbf{v}_h^*) = 0$ for $e \subset \Gamma_2$. In summary, $\mathbf{v}_h^*$ has zero degrees of freedom on $E \cup \Gamma_2$. Thus, by Theorem 3.1 the function $\mathbf{v}_h^*$ admits a regular decomposition

$$\mathbf{v}_h^* = p_{h,2}^* + \mathbf{r}_h \mathbf{w}_{h,2}^* + \mathbf{R}_{h,2}^* \quad \text{on } G_2, \quad (6.5)$$

with $p_{h,2}^* \in Z_h(G_2)$ and $\mathbf{w}_{h,2}^* \in (Z_h(G_2))^3$ vanishing on $E \cup \Gamma_2$. Moreover, we have (the set $E \cup \Gamma_2$ is non-connected since $E \cap \Gamma_2 = \emptyset$ and $\Gamma_2 \neq \emptyset$)

$$\|\mathbf{w}_{h,2}^*\|_{1,G_2} + h^{-1} \|\mathbf{R}_{h,2}^*\|_{0,G_2} \lesssim \log(1/h) \|\mathbf{v}_h^*\|_{\mathbf{curl},G_2} \quad (6.6)$$

and

$$\|\mathbf{w}_{h,2}^*\|_{0,G_2} + \|p_{h,2}^*\|_{1,G_2} \lesssim \log(1/h) \|\mathbf{v}_h^*\|_{\mathbf{curl},G_2}. \quad (6.7)$$

When Γ is connected, for example, Γ_1 is connected but $\Gamma_2 = \emptyset$, the norm on the right side of (6.6) can be replaced by the $\mathbf{curl}$ semi-norm.

Since $p_{h,2}^*$, $\mathbf{w}_{h,2}^*$ and $\mathbf{R}_{h,2}^*$ have zero degrees of freedom on E , these functions can be extended onto G_1 by zero and the resulting extensions $\tilde{p}_{h,2}^*$, $\tilde{\mathbf{w}}_{h,2}^*$ and $\tilde{\mathbf{R}}_{h,2}^*$ satisfy $\tilde{p}_{h,2}^* \in Z_h(G)$, $\tilde{\mathbf{w}}_{h,2}^* \in (Z_h(G))^3$ and $\tilde{\mathbf{R}}_{h,2}^* \in V_h(G)$, respectively. Define

$$p_h = \tilde{p}_{h,1} + \tilde{p}_{h,2}^*, \quad \mathbf{w}_h = \tilde{\mathbf{w}}_{h,1} + \tilde{\mathbf{w}}_{h,2}^* \quad \text{and} \quad \mathbf{R}_h = \tilde{\mathbf{R}}_{h,1} + \tilde{\mathbf{R}}_{h,2}^*.$$

Then p_h , $\mathbf{w}_h$ and $\mathbf{R}_h$ have zero degrees of freedom on $\Gamma = \Gamma_1 \cup \Gamma_2$ and the regular decomposition (3.4) follows by (6.4) and (6.5).

In an analogous way with Step 2 in the proof of Theorem 4.1, we can verify the estimates (3.5) and (3.6) by using (6.2)-(6.3) and (6.6)-(6.7). $\sharp$

In the following we want to prove Theorem 3.3 (and Theorem 3.4). For this purpose, we first prove two auxiliary results involving one vertex on a usual polyhedron domain.

Lemma 6.1 *Let G be a usual polyhedron and v be a vertex of G . Assume that $\mathbf{v}_h$ is a function in $V_h(G)$. Then we can write $\mathbf{v}_h$ as*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h$$

for some $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$ satisfying $p_h(v) = 0$ and $\mathbf{w}_h(v) = \mathbf{0}$. Moreover, we have

$$\|\mathbf{w}_h\|_{1,G} + h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G} \quad (6.8)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|\nabla p_h\|_{0,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (6.9)$$

Proof. Consider a (closed) face F containing v as its vertex. Like Step 1 in the proof of Lemma 5.2, we can define $\phi_{\partial F}$ to be a function that is piecewise linear and continuous on ∂F such that $\phi_{\partial F}(v) = 0$, and define $C_{\partial F}$ to be a constant such that $\mathbf{v}_h \cdot \mathbf{t}_{\partial F} = \phi'_{\partial F} + C_{\partial F}$ on ∂F . In fact, they can be defined as

$$C_{\partial F} = \frac{1}{l} \int_0^l (\mathbf{v}_h \cdot \mathbf{t}_{\partial F})(s) ds, \quad \phi_{\partial F}(t) = \int_0^t (\mathbf{v}_h \cdot \mathbf{t}_{\partial F} - C_{\partial F})(s) ds, \quad \forall t \in [0, l], \quad (6.10)$$

where l is the length of ∂F and $t = 0$ (and $t = l$) corresponds to the vertex v . Let $c = \gamma_E(\phi_{\partial F})$ denote the average of $\phi_{\partial F}$ on E , where E is an edge or a union of several edges of F . Define an extension $\tilde{\phi}_V \in Z_h(G)$ of $\phi_{\partial F}$, such that $\tilde{\phi}_V$ equals to the average $c = \gamma_E(\phi_{\partial F})$ at all of the nodes on G except those on ∂F . Then (cf. Lemma 7.6 and Lemma 7.1 of [33])

$$\|\nabla \tilde{\phi}_V\|_{0,G} = \|\nabla(\tilde{\phi}_V - c)\|_{0,G} \lesssim \|\phi_{\partial F} - c\|_{0,\partial F} \lesssim \|\phi'_{\partial F}\|_{H^{-1}(\partial F)} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (6.11)$$

We define a similar extension $\tilde{C}_V$ of $C_{\partial F}$ with $\tilde{C}_E$ (defined in the proof of Lemma 5.2), such that $\tilde{C}_V$ belongs to $(Z_h(G))^3$ and vanishes at v , and it satisfies the condition $(\mathbf{r}_h \tilde{C}_V) \cdot \mathbf{t}_{\partial F} = C_{\partial F}$ on ∂F and the stability

$$\|\tilde{C}_V\|_{1,G} \lesssim \|\tilde{C}_V\|_{0,\partial F} \lesssim |C_{\partial F}| \lesssim \log^{\frac{1}{2}}(1/h) \|\mathbf{curl} \mathbf{v}_h\|_{0,G}.$$

Define

$$\hat{\mathbf{v}}_{h,V} = \mathbf{v}_h - (\nabla \tilde{\phi}_V + \mathbf{r}_h \tilde{C}_V).$$

Then we have $\hat{\mathbf{v}}_{h,V} \cdot \mathbf{t}_{\partial F} = 0$. As in the proof of Lemma 5.2, we can use Lemma 5.1 for $\hat{\mathbf{v}}_{h,V}$ to build the desired decomposition of $\mathbf{v}_h$. $\sharp$

Remark 6.1 *Differently from the second inequality in (5.17), the inequality (6.9) holds only for the H^1 semi-norm of p_h . The main reason is that a stable estimate of $\|\tilde{\phi}_V\|_{0,G}$ cannot be built except that the constant c in (6.11) vanishes. In the above proof, the face F can be replaced by a connected “Lipschitz” union of some faces of G , provided that union set has one connected boundary containing v as one of its vertices (the union cannot be ∂G itself).*

Corollary 6.1. Let $\bar{G} = \bar{G}_1 \cup \bar{G}_2$ with G_i be a usual polyhedron domain. Assume that $\bar{G}_1 \cap \bar{G}_2$ is just the common vertex v of G_1 and G_2 . Then any function $\mathbf{v}_h \in V_h(G)$ admits a discrete regular decomposition on G and the resulting functions have the stability estimates given in Lemma 6.1.

Proof. We use Lemma 6.1 to build a regular decomposition of $\mathbf{v}_h|_{G_i}$ independently for $i = 1, 2$. Since the resulting nodal finite element functions vanish at v , the two regular decompositions naturally define a regular decomposition for $\mathbf{v}_h$ on the global domain G . $\sharp$

In the rest of this section, for a face (or a connected “Lipschitz” union of some faces) F , we always use $\phi_{\partial F}$ to denote the function defined in the proof of Lemma 6.1.

Lemma 6.2 *Let G be a usual polyhedron, and let Γ be a (closed) union of some faces and edges of G , and v be a vertex of G ($v \notin \Gamma$). Assume that $\mathbf{v}_h$ satisfies **Assumption 3.1**. Suppose that there exists a face (or a connected “Lipschitz” union of some faces) F containing v such that $\gamma_E(\phi_{\partial F}) = 0$ for an edge E of F , where either $F \cap \Gamma = \emptyset$ or $\partial F \cap \Gamma$ is a union of edges of G (in this case, we require that $E \subset \partial F \cap \Gamma$). Then $\mathbf{v}_h$ can be decomposed as*

$$\mathbf{v}_h = \nabla p_h + \mathbf{r}_h \mathbf{w}_h + \mathbf{R}_h \quad (6.12)$$

for some $p_h \in Z_h(G)$, $\mathbf{w}_h \in (Z_h(G))^3$ and $\mathbf{R}_h \in V_h(G)$ such that p_h and $\mathbf{w}_h$ vanish on Γ and v . Moreover, we have

$$\|\mathbf{w}_h\|_{1,G} + h^{-1} \|\mathbf{R}_h\|_{0,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G} \quad (6.13)$$

and

$$\|\mathbf{w}_h\|_{0,G} + \|p_h\|_{1,G} \lesssim \log(1/h) \|\mathbf{v}_h\|_{\mathbf{curl},G}. \quad (6.14)$$

In particular, when Γ is connected and $E \subset \partial F \cap \Gamma$, the norm on the right side of (6.13) can be replaced by the **curl** semi-norm.

Proof. A key step in the proof is to define two functions $\tilde{\phi}_V \in Z_h(G)$ and $\tilde{C}_V \in (Z_h(G))^3$ such that they have zero degrees of freedom at v and Γ and satisfy $\mathbf{v}_h \cdot \mathbf{t}_{\partial F} = (\nabla \tilde{\phi}_V + \mathbf{r}_h \tilde{C}_V) \cdot \mathbf{t}_{\partial F}$ on ∂F .

We first consider the case with $F \cap \Gamma = \emptyset$. For this case, we can simply define $\tilde{\phi}_V$ and $\tilde{C}_V$ as in the proof of Lemma 6.1. Set

$$\hat{\mathbf{v}}_{h,V} = \mathbf{v}_h - (\nabla \tilde{\phi}_V + \mathbf{r}_h \tilde{C}_V).$$

Then we have $\lambda_e(\hat{\mathbf{v}}_{h,V}) = 0$ for any $e \subset \partial F \cup \Gamma$ by the definitions of $\tilde{\phi}_V$ and $\tilde{C}_V$, together with the assumption on $\mathbf{v}_h$. Thus we can use Theorem 3.1 (with Γ replacing by $\partial F \cup \Gamma$) to build a regular decomposition of $\hat{\mathbf{v}}_{h,V}$ such that the functions defined by the decomposition have zero degrees of freedom on $\partial F \cup \Gamma$, which implies that the two H^1 finite element functions vanish at v (and Γ) since $v \in \partial F$. Based on this decomposition we further get the desired decomposition of $\mathbf{v}_h$ and the estimates (refer to the proof of Lemma 6.1). Here we need to use the assumption $\gamma_E(\phi_{\partial F}) = 0$ to get the L^2 stability of $\tilde{\phi}_V$ (see Remark 6.1).

Now we investigate the case with $\partial F \cap \Gamma$ being a union of edges of G . For this case, the analysis is a bit complicated because the functions $\tilde{\phi}_V$ and $\tilde{C}_V$ defined in the proof of Lemma 6.1 may do not vanish on $\partial F \cap \Gamma$ although they vanish at v . Based on this consideration, we have to define two new functions $\tilde{\phi}_V$ and $\tilde{C}_V$ such that they can satisfy the requirements mentioned above.




Figure 8: An example of cube: the blue face denote Γ , the red face denotes F , $F \cap \Gamma = E$.

Without loss of generality, we assume that F is a face and $\partial F \cap \Gamma = E$ is an edge of G (see Figure 8). For more general situation, there is no essential change in the analysis. Let $\phi_{\partial F}$ and $C_{\partial F}$ be defined by (6.10). The basic ideas are to choose a piecewise constant function $\varepsilon_{\partial F}$ on ∂F and define $\tilde{\phi}_V \in Z_h(G)$ such that $\tilde{\phi}_V$ vanishes at all of the nodes on $\bar{G} \setminus \partial F$ and is given by the following formula on ∂F

$$\tilde{\phi}_V(t) = \phi_{\partial F}(t) - \int_0^t \varepsilon_{\partial F} ds, \quad t \in [0, |\partial F|], \quad (6.15)$$

where $t = 0$ and $t = |\partial F|$ are the arc-length coordinates of the vertex v .

In the following we give a definition of $\varepsilon_{\partial F}$. Notice that $\phi_{\partial F}(|\partial F|) = 0$, the function $\varepsilon_{\partial F}$ should satisfy $\int_0^{|\partial F|} \varepsilon_{\partial F} ds = 0$ so that $\tilde{\phi}_V(|\partial F|) = 0$. Besides, since we hope that $\tilde{\phi}_V$ vanishes on E , we naturally define $\varepsilon_{\partial F} = \phi'_{\partial F}$ on E . Let $E_1, E_2 \subset \partial F$ be the two neighboring edges with E (see Figure 8), and let t_1 and t_2 denote the arc-length coordinates of the two endpoints of E ($t_1 < t_2$). Then the piecewise constant function $\varepsilon_{\partial F}$ can be defined as

$$\varepsilon_{\partial F} = \begin{cases} \phi'_{\partial F} & \text{on } E, \\ \phi_{\partial F}(t_1)/|E_1| & \text{on } E_1, \\ -\phi_{\partial F}(t_2)/|E_2| & \text{on } E_2, \\ 0 & \text{on } \partial F \setminus (E \cup E_1 \cup E_2). \end{cases}$$

It can be verified that, for the above $\varepsilon_{\partial F}$, the function $\tilde{\phi}_V$ defined in (6.15) indeed vanishes on v and E .

For the new function $\tilde{\phi}_V$, we need to define the corresponding function $\tilde{C}_V$, which can be regarded as a variant of $C_{\partial F}$. Since $E \subset \partial F \cap \Gamma$ and $\mathbf{v}_h$ has zero degrees of freedom on Γ , we have $(\mathbf{v}_h \cdot \mathbf{t}_{\partial F})|_E = \mathbf{v}_h \cdot \mathbf{t}_E = 0$. Then, by the direct calculation, we get

$$\phi_{\partial F}(t) = \int_0^t (\mathbf{v}_h \cdot \mathbf{t}_{\partial F} - C_{\partial F}) ds = \int_0^{t_1} \mathbf{v}_h \cdot \mathbf{t}_{\partial F} ds - t C_{\partial F}, \quad t \in [t_1, t_2].$$

Thus we have $\varepsilon_{\partial F} = \phi'_{\partial F} = -C_{\partial F}$ on E . Define $\tilde{C}_V \in (Z_h(G))^3$ such that $\tilde{C}_V \cdot \mathbf{t}_{\partial F} = C_{\partial F} + \varepsilon_{\partial F}$ on ∂F , and $\tilde{C}_V$ vanishes at v and all of the nodes on $\bar{G} \setminus \partial F$. It is clear that $\tilde{C}_V \cdot \mathbf{t}_{\partial F} = 0$ on E by the relation $\varepsilon_{\partial F} = -C_{\partial F}$. Moreover, the following decomposition holds for the modified functions $\tilde{\phi}_V$ and $\tilde{C}_V$

$$\mathbf{v}_h \cdot \mathbf{t}_{\partial F} = C_{\partial F} + \phi'_{\partial F} = \tilde{C}_V \cdot \mathbf{t}_{\partial F} + \tilde{\phi}'_V \quad \text{on } \partial F. \quad (6.16)$$

From the above discussions, we know that the functions $\tilde{C}_V$ and $\tilde{\phi}_V$ have zero degrees of freedom on Γ and v (since the two functions naturally have zero degrees of freedom on $\Gamma \setminus E$ by their definitions). In addition, the functions $\tilde{C}_V$ and $\tilde{\phi}_V$ possess the same stabilities as the corresponding functions defined in the proof of Lemma 6.1. Here we use the condition that $\tilde{\phi}_V$ vanishes on E to get the L^2 stability of $\tilde{\phi}_V$. Define

$$\hat{\mathbf{v}}_{h,V} = \mathbf{v}_h - (\nabla \tilde{\phi}_V + \mathbf{r}_h \tilde{C}_V).$$

By (6.16) and the assumption on $\mathbf{v}_h$, we have $\lambda_e(\hat{\mathbf{v}}_{h,V}) = 0$ for any $e \subset \partial F \cup \Gamma$. Thus we can prove the desired results as in the case with $F \cap \Gamma = \emptyset$ (see the previous part). In particular, if Γ is connected, then $F \cup \Gamma$ is also connected, and so the complete norm on the right side of (6.13) can be replaced by the **curl** semi-norm (see the conclusion given at the bottom of Theorem 3.1). $\sharp$

Remark 6.2 We find, from Lemma 6.2, that a vertex is essentially different from an edge. This phenomenon can be intuitively explained as follows: the value of an edge finite element function $\mathbf{v}_h$ is not uniquely defined at a vertex, but the two nodal finite element functions defined by the regular decomposition of $\mathbf{v}_h$ are required to vanish at the vertex. Thus there is a gap between $\mathbf{v}_h$ and the gradient of the scalar finite element function, which needs to be filled by a constraint of $\mathbf{v}_h$.

Now we prove Theorem 3.3 by Theorem 3.1, Lemma 6.1 and Lemma 6.2.

Proof of Theorem 3.3. We recall that $\bar{G} = \bar{G}_1 \cup \bar{G}_2$, where G_1 and G_2 are two usual polyhedra that intersect at a vertex v , and $\Gamma_i = \Gamma \cap \partial G_i$. We consider three different situations, respectively.

(i) Both Γ_1 and Γ_2 contain the vertex v , which implies that the intersection of Γ_1 and Γ_2 is just v .

We use Theorem 3.1 to build a regular decomposition of $\mathbf{v}_h|_{G_i}$ independently for $i = 1, 2$, such that the resulting nodal finite element functions have zero degrees of freedom on Γ_i ($i = 1, 2$). Since $v = \Gamma_1 \cap \Gamma_2$, the nodal finite element functions also vanish at v . Thus the two regular decompositions can naturally define the desired regular decomposition on the global domain G . In this case, the constraint is unnecessary, i.e., we can choose $\mathcal{F}$ as the zero functional.

(ii) One of Γ_1 and Γ_2 contains the vertex v , for example, $v \in \Gamma_1$ but $v \notin \Gamma_2$.

We choose a face F of G_2 such that $v \in F$ (see the left graph in Figure 9). Assume that either $F \cap \Gamma_2$ is an empty set or $F \cap \Gamma_2$ is a union of edges. Let s be the arc-length variable along ∂F , taking values from 0 to $|\partial F|$, which is the total length of ∂F . Define

$$C_{\partial F} = \frac{1}{|\partial F|} \int_{\partial F} (\mathbf{v}_h \cdot \mathbf{t}_{\partial F})(s) ds, \quad \phi_{\partial F}(t) = \int_0^t (\mathbf{v}_h \cdot \mathbf{t}_{\partial F} - C_{\partial F})(s) ds, \quad \forall t \in [0, |\partial F|].$$

For an edge E of F , let $\gamma_E(\phi_{\partial F})$ denote the integration average of $\phi_{\partial F}$ on E . Define the functional $\mathcal{F}$ as $\mathcal{F}\mathbf{v}_h = \gamma_E(\phi_{\partial F})$, where we require that $E \subset F \cap \Gamma_2$ in the case of $F \cap \Gamma_2 \neq \emptyset$. Then the constraint $\mathcal{F}\mathbf{v}_h = 0$ means that $\gamma_E(\phi_{\partial F}) = 0$.




Figure 9. The blue part of ∂G_i denotes Γ_i . Situation (i) is shown by the left graph: $v \in \Gamma_1$, $v \notin \Gamma_2$, $F \subset \partial G_2$ and $F \cap \Gamma_2$ is an edge, the red part of ∂G_2 denotes F ; Situation (ii) is shown by the right graph: $v \notin \Gamma_1 \cup \Gamma_2$, $F_1 \subset \partial G_1$ and $F_2 \subset \partial G_2$, the red part of ∂G_i denotes F_i .

We use Theorem 3.1 to build a discrete regular decomposition on G_1 such that the resulting functions have zero degrees of freedom on Γ_1 . Since $v \in \Gamma_1$, the H^1 finite element functions vanish at v . Moreover, we use Lemma 6.1 (if $\Gamma_2 = \emptyset$) or Lemma 6.2 (if $\Gamma_2 \neq \emptyset$) to build a discrete regular decomposition on G_2 such that the resulting three functions have zero degrees of freedom on Γ_2 and the two H^1 finite element functions vanish at v . Then the two regular decompositions naturally define the desired regular decomposition on G . The stability of the L^2 norm of p_h can be guaranteed by the condition $\gamma_E(\phi_{\partial F}) = 0$.

(iii) Neither Γ_1 nor Γ_2 contains the vertex v , i.e., $v \notin \Gamma = \Gamma_1 \cup \Gamma_2$.

We choose two faces $F_1 \subset \partial G_1$ and $F_2 \subset \partial G_2$ such that both of them contain v (see the right graph in Figure 9). Suppose that either $F_i \cap \Gamma_i$ is an empty set or $\partial F_i \cap \Gamma_i$ is a union of some edges of G_i ($i = 1, 2$). The face F_i can be replaced by a connected “Lipschitz” union of some faces of G_i , provided that the union has one connected boundary containing v as one of its vertices (the union cannot be ∂G_i itself). Define

$$C_{\partial F_i} = \frac{1}{|\partial F_i|} \int_{\partial F_i} (\mathbf{v}_h \cdot \mathbf{t}_{\partial F_i})(s) ds, \quad \phi_{\partial F_i}(t) = \int_0^t (\mathbf{v}_h \cdot \mathbf{t}_{\partial F_i} - C_{\partial F_i})(s) ds + c_i, \quad \forall t \in [0, |\partial F_i|],$$

where the constant c_i is chosen such that $\gamma_{E_i}(\phi_{\partial F_i}) = 0$ for some edge $E_i \subset \partial F_i$. When $\partial F_i \cap \Gamma_i$ is a union of some edges of G_i , we require that $E_i \subset \partial F_i \cap \Gamma_i$.

Define the functional $\mathcal{F}$ as $\mathcal{F}\mathbf{v}_h = \phi_{\partial F_1}(|\partial F_1|) - \phi_{\partial F_2}(|\partial F_2|)$. The constraint $\mathcal{F}\mathbf{v}_h = 0$ means that $\phi_{\partial F_1}(|\partial F_1|) = \phi_{\partial F_2}(|\partial F_2|)$.

For convenience, we only consider the case with $F_i \cap \Gamma_i = \emptyset$ ($i = 1, 2$), otherwise, we need to repeat the technique in Lemma 6.2. As in the proof of Lemma 6.1, we respectively define the extensions $\tilde{\phi}_V^{(i)}$ and $\tilde{C}_V^{(i)}$ of $\phi_{\partial F_i}$ and $C_{\partial F_i}$, but define the values of $\tilde{\phi}_V^{(i)}$ as zero at all of the nodes in G_i except on ∂F_i . Set

$$\hat{\mathbf{v}}_{h,V}^{(i)} = \mathbf{v}_h|_{G_i} - (\tilde{\phi}_V^{(i)} + \mathbf{r}_h \tilde{C}_V^{(i)}) \quad \text{on } G_i.$$

Then we have $\lambda_e(\hat{\mathbf{v}}_{h,V}^{(i)}) = 0$ for any $e \subset \partial F_i \cup \Gamma_i$ by the definitions of $\tilde{\phi}_V^{(i)}$ and $\tilde{C}_V^{(i)}$. Thus we can use Theorem 3.1 for $\hat{\mathbf{v}}_{h,V}^{(i)}$ to build a regular decomposition of $\mathbf{v}_h|_{G_i}$ and the resulting functions have the corresponding stability estimates by the condition $\gamma_{E_i}(\phi_{\partial F_i}) = 0$ (refer to the proof of Lemma 6.1 and Remark 6.1). Using the definitions of the functions $p_{h,i}$ and $\mathbf{w}_{h,i}$ in the regular decomposition of $\mathbf{v}_h|_{G_i}$, together with the condition $\phi_{\partial F_1}(|\partial F_1|) = \phi_{\partial F_2}(|\partial F_2|)$, yields $p_{h,1} = p_{h,2}$ and $\mathbf{w}_{h,i} = \mathbf{0}$ at the vertex v . Therefore, we can naturally define a regular decomposition of $\mathbf{v}_h$ on the global G and obtain the desired stability estimates. $\sharp$

Proof of Theorem 3.4. Let F_i be a face (or a connected “Lipschitz” union of faces) satisfying $v \in \partial F_i \subset \partial G_i$ and define a function $\phi_{\partial F_i}$ as in the proof of Theorem 3.3 ($i = 1, \dots, s$). We define the functional $\mathcal{F}_i$ by

$$\mathcal{F}_i \mathbf{v}_h = \phi_{\partial F_{i+1}}(v) - \phi_{\partial F_i}(v) \quad (i = 1, \dots, s-1);$$

Then we can prove the desired results in an analogous way with the proof of Theorem 3.3. $\sharp$

Acknowledgments. The author would like to thank the anonymous reviewer, who gives many insightful comments to improve the presentation of this paper. Moreover, the author wishes to thank Professor Jinchao Xu for suggesting him to study the convergence of the HX preconditioner for the case with jump coefficients. In particular, Professor Jinchao Xu suggested the author to consider regular decomposition on non-Lipchitz domains, which makes the results in this paper be richer and more complete.

References

- [1] M. Ainsworth, J. Guzman and F. Sayas, *Discrete extension operators for mixed finite element spaces on locally refined meshes*, Math. Comput., **85**(2016), pp. 2639-2650.
- [2] A. Alonso and A. Valli, *Some remarks on the characterization of the space of tangential traces of $H(\mathbf{curl}; \Omega)$ and the construction of an extension operator*, Manuscr. Math., **89**(1996), 159-178.
- [3] A. Alonso and A. Valli, *An optimal domain decomposition preconditioner for low-frequency time-harmonic Maxwell equations*, Math. Comp., **68**(1999), 607-631.
- [4] R. Albanese and G. Rubinacci, *Magnetostatic field computations in terms of two-component vector potentials*, Int. J. Numer. Meth. Engng., **29**(1990), 515-532.
- [5] C. Amrouche, C. Bernardi, M. Dauge and V. Girault. Vector Potentials in three dimensional nonsmooth domains, Math. Meth. Appl. Sci., 21(1998), 823-864
- [6] A. Bossavit, Computational Electromagnetism. Variational Formulation, Complementarity, Edge Elements, Electromagnetism 2, Academic Press, San Diego, CA, 1998.
- [7] F. Brezzi and M. Fortin, Mixed and Hybrid Finite Element Methods, Springer-Verlag, New York, 1991.
- [8] M. Casarin, Schwarz Preconditioners for Spectral and Mortar Finite Element Methods with Applications to Incompressible Fluids. PhD thesis, Courant Institute of Mathematics Sciences, March 1996.
- [9] P. Ciarlet, Jr. and J. Zou, *Fully discrete finite element approaches for time-dependent Maxwell's equations*, Numer. Math., **82**(1999), 193-219.
- [10] P. Degond and P.-A. Raviart, *An analysis of the Darwin model of approximation to Maxwell's equations*, Forum Math., **4**(1992), 13-44.
- [11] A. S. B.B. Dhia, C. Hazard, and S. Lohrengel. A singular field method for the solution of Maxwells equations in polyhedral domains. SIAM J. Appl. Math., 59(6):2028C2044 (electronic), 1999. ISSN 0036-1399.
- [12] V. Girault and P. A. Raviart, Finite Element Methods for Navier-Stokes Equations, Springer- Verlag, New York, 1986
- [13] P. Grisvard, Elliptic Problems in Nonsmooth Domains, Monographs and Studies in Mathematics, 24, Pitman, London, 1985.
- [14] R. Hiptmair, *Multigrid method for Maxwell's equations*, SIAM J. Numer. Anal., **36**(1998), 204-225.
- [15] R. Hiptmair, Finite elements in computational electromagnetism, Acta Numer., 11 (2002), pp. 237-339.
- [16] R. Hiptmair and C. Pechstein, *Discrete regular decompositions of tetrahedral discrete 1-forms*, Tech. Rep. 2017-47, Seminar for Applied Mathematics, ETH Zurich, Switzerland, 2017

- [17] R. Hiptmair and J. Xu, *Nodal auxiliary space preconditioning in $H(\text{curl})$ and $H(\text{div})$ spaces*, SIAM J. Numer. Anal., **45**(2007), 2483-2509.
- [18] R. Hiptmair and W. Zheng, *Local multigrid in $H(\text{curl})$* , J. Comp. Math. 27 (2009), 573-603
- [19] Q. Hu, *Convergence of HX Preconditioner for Maxwell's Equations with Jump Coefficients (ii): The main results*. (which will be submitted to this journal in near future)
- [20] Q. Hu, G. Liang and J. Lui, *The construction of preconditioner for domain decomposition methods with polynomial Lagrangian multipliers*, J. Comp. Math. 19 (2001) 213-224.
- [21] Q. Hu, Shi Shu and J. Zou, *A discrete weighted Helmholtz decomposition and its applications*, Numer. Math., **125**(2013), 153-189.
- [22] Q. Hu and J. Zou, *A non-overlapping domain decomposition method for Maxwell's equations in three dimensions*, SIAM J. Numer. Anal., **41**(2003), 1682-1708.
- [23] Q. Hu and J. Zou, *Substructuring preconditioners for saddle-point problems arising from Maxwell's equations in three dimensions*, Math. Comput., **73**(2004), 35-61.
- [24] P. Jones, *Quasiconformal mappings and extendability of functions in Sobolev spaces*, Acta Math., 147 (1981), pp. 71-88.
- [25] A. Klawonn, O. Rheinbach, and O. B. Widlund, *An analysis of a FETI-DP algorithm on irregular subdomains in the plane*, SIAM J. Numer. Anal., 46 (2008), pp. 2484-2504.
- [26] P. Monk, *Analysis of a finite element method for Maxwell's equations*, SIAM J. Numer. Anal., **29**(1992), 32-56.
- [27] Peter Monk, *Finite Element Methods for Maxwell's Equations*, Oxford University Press, Oxford, 2003.
- [28] J. Nédélec, *Mixed finite elements in R^3* , Numer. Math., **35**(1980), 315-341.
- [29] J.E. Pasciak and J. Zhao. *Overlapping Schwarz methods in $H(\text{curl})$ on polyhedral domains*. J. Numer. Math., 10(3):221-234, 2002.
- [30] J. Saranen, *On electric and magnetic problems for vector fields in anisotropic nonhomogeneous media*, J. Math. Anal. Appl., **91** (1983), 254-275.
- [31] L.R. Scott and S. Zhang, *Finite element interpolation of nonsmooth functions satisfying boundary conditions*, Math. Comp. 54 (1990), 483-493
- [32] A. Toselli, *Overlapping Schwarz methods for Maxwell's equations in three dimensions*, Numer. Math., **86**(2000), 733-752.
- [33] A. Toselli, *Dual-primal FETI algorithms for edge element approximations: three-dimensional h finite elements on shape-regular meshes*, IMA J. Numer. Anal., **26**(2006), 96-130.
- [34] A. Toselli and Olof Widlund, *Domain Decomposition Methods - Algorithms and Theory*, Volume 34, Springer Series in Computational Mathematics, Springer-Verlag, Berlin, 2005.

- [35] J. Xu and J. Zou, *Some non-overlapping domain decomposition methods*, SIAM Review, **40**(1998), 857-914