

# Martingale representation on enlarged filtrations: the role of the accessible jump times

Antonella Calzolari<sup>\*</sup>      Barbara Torti<sup>\*</sup>

## Abstract

We consider a filtration  $\mathbb{G}$  obtained as enlargement of a filtration  $\mathbb{F}$  by a filtration  $\mathbb{H}$ . We assume that all  $\mathbb{F}$ -local martingales are represented by a martingale  $M$  and all  $\mathbb{H}$ -local martingales are represented by a martingale  $N$ .  $M$  and  $N$  are not necessarily quasi-left continuous processes and their jump times may overlap. We first analyze the contribution of the accessible jump times of  $M$  and  $N$  to the Jacod's dimension of the space of the  $\mathcal{H}^1(\mathbb{G})$ -martingales. Then we prove a new martingale representation theorem on  $\mathbb{G}$ .

**Keywords:** Predictable representations property; enlargement of filtration; marked point processes; compensators.

## 1 Introduction

Martingale representation is a classical topic of stochastic analysis with relevant theoretical and practical applications. As well known, the martingale representation property is crucial to prove existence and uniqueness of BSDE's solutions, to derive the stochastic filtering equation by means of the innovation approach and to compute options hedging strategies in financial markets. It is interesting to study what happens concerning martingale representation when the available information grows and typically the class of local martingales changes. In this paper we investigate this problem assuming that the "representable" martingales may have jump times with non trivial *accessible component*. As far as we know, this assumption is an important novelty of our analysis compared to most of the papers on this subject. We are motivated by the growing interest of the literature in modeling situations of the real world which may present critical "announced" random times, that is, from a mathematical point of view, *predictable random times* (see e.g. Fontana and Smith [25]). A second point of novelty is that we allow the *additional information* to be not completely exogenous w.r.t. to the *basic information*. In fact, rarely in the literature the two kinds of information, basic and additional, are produced by processes which may jump at the same random times (see e.g. Jiao and Li [33], Jeanblanc and Song [31] and Aksamit, Jeanblanc and Rutkowski [3]). We stress also that only a few authors deal, as we do here, with martingale representation when the growth of information is due to a process possibly different from the occurrence process of a random time  $\tau$  (see

---

<sup>\*</sup>Dipartimento di Matematica - Università di Roma "Tor Vergata", via della Ricerca Scientifica 1, I 00133 Roma, Italy

e.g. Kchia and Protter [36], El Karoui, Jeanblanc and Jiao [24], Di Tella and Jeanblanc [22], Calzolari and Torti [14] and Biagini, Mazzon and Oberpriller [8]). Moreover, the additional information is always generated by the observation of a process, while here we model the extra information as the reference filtration of a martingale, not necessarily its natural one<sup>1</sup>. Based on the features highlighted in the above discussion, our work meets the need suggested by the recent literature of new martingale representations on extended filtrations (see e.g. Bandini, Confortola and Di Tella [7] and Bandini, Calvia and Colaneri [6]).

The framework of our study is the following. On a given probability space  $(\Omega, \mathcal{F}, P)$  let  $M = (M_t)_{t \in [0, T]}$  and  $N = (N_t)_{t \in [0, T]}$  be two square-integrable martingales enjoying the *strong predictable representation property (PRP)* w.r.t. the filtrations  $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$  and  $\mathbb{H} = (\mathcal{H}_t)_{t \in [0, T]}$ , respectively. This means that all  $(P, \mathbb{F})$  ( $(P, \mathbb{H})$ )-local martingales can be represented, up to the initial condition, as a stochastic integral w.r.t.  $M$  (w.r.t.  $N$ ). Let  $\mathbb{G}$  be the smallest right-continuous filtration containing both  $\mathbb{F}$  and  $\mathbb{H}$ . If there exists an equivalent probability measure such that  $\mathbb{F}$  and  $\mathbb{H}$  are independent (*equivalent decoupling measure*), then it has been proved that the  $\mathbb{R}^3$ -valued process  $(M, N, [M, N])$  enjoys the  $\mathbb{G}$ -PRP under the *martingale preserving measure*,  $P^*$ , which is the decoupling measure under which  $M$  and  $N$  are  $\mathbb{G}$ -martingales (see [11]). In other words any  $(P^*, \mathbb{G})$ -local martingale can be written as a *vector stochastic integral* driven by  $(M, N, [M, N])$ . The fundamental argument is that the covariation process  $[M, N]$  is a  $(P^*, \mathbb{G})$ -square-integrable martingale strongly orthogonal to  $M$  and  $N$ . So the set  $(M, N, [M, N])$  turns out to be a  $(P^*, \mathbb{G})$ -*basis of martingales* in the sense of Davis and Varaiya (see [19]). The role of  $\mathbb{F}$  and  $\mathbb{H}$  is symmetric and we can look at  $\mathbb{G}$  either as the *enlargement of  $\mathbb{F}$  by  $\mathbb{H}$*  or viceversa.

In this paper we face two different issues linked to the above result.

Our first problem starts from the following remark. The process  $[M, N]$ , when not trivial, is a *pure jump process* whose jump times arise by the overlapping of the accessible components of the jump times of  $M$  and  $N$ . Therefore  $[M, N]$  vanishes almost surely as soon as at least one of the two martingales has *totally inaccessible* jump times only. The non-triviality of  $[M, N]$  reflects on the dimension of the  $(P^*, \mathbb{G})$ -martingale driving the representation. We investigate the consequences of this fact on the Jacod's dimension of the *martingale space*  $\mathcal{H}^1(P, \mathbb{G})$ , that is the minimal possible dimension of a vector-valued local martingale enjoying the  $(P, \mathbb{G})$ -PRP. We show that the maximum value of  $\dim(\mathcal{H}^1(P, \mathbb{G}))$  only depends on the mutual singularity of the kernels associated with the sharp brackets of  $M$  and  $N$  joint with the mutual singularity of the kernels associated with the sharp brackets of their accessible martingale parts. Based on the mutual behavior of the sharp brackets, we provide a condition equivalent to get  $\dim(\mathcal{H}^1(P, \mathbb{G}))$  equal to one and a condition sufficient to get  $\dim(\mathcal{H}^1(P, \mathbb{G}))$  equal to two.

The second problem we deal with is the construction of a local martingale driving the  $(P, \mathbb{G})$ -martingale representation. More precisely, taking the point of view of the enlargement of  $\mathbb{F}$  by  $\mathbb{H}$ , we compute explicitly a multidimensional local martingale enjoying the  $(P, \mathbb{G})$ -PRP. Indeed  $N$  is a special  $(P, \mathbb{G})$ -semi-martingale and we denote by  $N'$  its local martingale part. Our main hypothesis is that  $P$  is the *minimal martingale measure* for

---

<sup>1</sup> for an example of a martingale enjoying the PRP w.r.t. a filtration larger than the natural one refer to the notion of *Weak Brownian Filtration* in [37]

$N'$  w.r.t.  $P^*$  (see [5]). We show that this condition implies that any  $(P, \mathbb{F})$ -martingale is a  $(P, \mathbb{G})$ -martingale (*immersion of  $\mathbb{F}$  in  $\mathbb{G}$  under  $P$* ) and coincides with this property when  $[M, N] \equiv 0$ . Under our assumptions we prove that  $(M, N', [M, N])$  is a  $(P, \mathbb{G})$ -local martingale representing all the others.

The addressed issues lead us to prove two results of stochastic analysis of general interest. More precisely we write the explicit expression of the compensator of a jump process with accessible jump times only and we show the existence of two predictable sets whose  $\omega$ -sections are disjoint supports of the random measures generated by two predictable increasing processes with mutually singular associated kernels.

The paper is organized as follows. Section 2 is devoted to preliminaries. After introducing the notations and recalling the necessary definitions and statements, we present in separated subsections some original results of the theory of semi-martingales interesting by themselves. Section 3 describes the specific setting of our work. In Section 4 we prove the theorem about the dimension of  $\mathcal{H}^1(P, \mathbb{G})$ . The result on martingale representation is the object of Section 5. First we discuss the case of progressive enlargement by  $\tau$  and then the general case. Finally in Section 6 we formulate conclusions and future perspectives.

## 2 Preliminaries

Throughout the paper we consider processes on the time interval  $[0, T]$ . Given a standard filtered probability space  $(\Omega, \mathcal{A}, \mathbb{A} = (\mathcal{A}_t)_{t \leq T}, R)$  we denote by  $\mathcal{M}(R, \mathbb{A})$  the set of  $(R, \mathbb{A})$ -martingales ( $\mathcal{M}_0(R, \mathbb{A})$  when the initial value is null) and by  $\mathcal{M}_{loc}(R, \mathbb{A})$  ( $\mathcal{M}_{loc,0}(R, \mathbb{A})$ ) the set of  $(R, \mathbb{A})$ -local martingales. Analogously  $\mathcal{M}^2(R, \mathbb{A})$  ( $\mathcal{M}_0^2(R, \mathbb{A})$ ) is the set of square integrable  $(R, \mathbb{A})$ -martingales and  $\mathcal{M}_{loc}^2(R, \mathbb{A})$  ( $\mathcal{M}_{loc,0}^2(R, \mathbb{A})$ ) its localization.

If  $Y$  and  $Y'$  are two semi-martingales we denote by  $[Y, Y']$  their *quadratic covariation process* and, when  $Y = Y'$ , we simply write  $[Y]$  in place of  $[Y, Y]$ . Given two local martingales  $Z$  and  $Z'$  we denote by  $\langle Z, Z' \rangle^{R, \mathbb{A}}$  their *sharp bracket process* and, when  $Z = Z'$ , we simply write  $\langle Z \rangle^{R, \mathbb{A}}$  in place of  $\langle Z, Z \rangle^{R, \mathbb{A}}$  (for definitions and existence's conditions we refer to Chapter VII in [21]).

Given a semi-martingale  $Y$  we denote by  $\mathbb{P}(Y, \mathbb{A})$  the set of the local martingale measures for  $Y$  on  $(\Omega, \mathcal{A}_T)$  equivalent to  $R|_{\mathcal{A}_T}$ .

Following Definition 4.22, page 121, in [26], a random time  $\eta$  is a *jump time* of a càdlàg process  $X$  if  $R(X_\eta \neq X_{\eta-}, \eta < +\infty) = R(\eta < +\infty)$ . As usual we denote with  $\Delta X_\eta$  the random variable  $X_\eta - X_{\eta-}$  with the convention that  $\Delta X_\eta = 0$  on  $\{\eta = +\infty\}$ . When, as in this paper,  $X$  lives on the time interval  $[0, T]$ , its jump times are random variables taking values in  $[0, T] \cup \{+\infty\}$ .

A stopping time  $\eta$  satisfies

$$\eta = \eta^{dp} \wedge \eta^{dq} \tag{1}$$

where  $\eta^{dp}$  and  $\eta^{dq}$  are the *accessible component* and the *totally inaccessible component*, respectively (see e.g. Theorem 3, page 104, in [38]). More precisely there exist two disjoint events  $A$  and  $B$  such that  $P$ -a.s.

$$A \cup B = (\eta < \infty)$$

and

$$\eta^{dp} = \eta \mathbb{I}_A + \infty \mathbb{I}_{A^c}, \quad \eta^{dq} = \eta \mathbb{I}_B + \infty \mathbb{I}_{B^c}. \quad (2)$$

Therefore

$$\eta \mathbb{I}_{\eta < +\infty} = \eta^{dp} \mathbb{I}_A + \eta^{dq} \mathbb{I}_B.$$

It is worthwhile to recall that

$$[[\eta^{dp}]] \subset \cup_m [[\eta_m^{dp}]], \quad (3)$$

where  $\{\eta_m^{dp}\}_{m \in \mathbb{N}}$  is a sequence of predictable stopping times. Such a sequence is not unique and it may be chosen in such a way that the corresponding graphs are pairwise disjoint (see Theorem 3.31, page 95, in [26]). As far as  $\eta^{dq}$  is concerned, we recall that by definition

$$R(\eta^{dq} = \sigma < +\infty) = 0,$$

for any predictable stopping time, and therefore for any accessible stopping time,  $\sigma$ .

*From now on we call enveloping sequence of  $\eta^{dp}$  any sequence of predictable stopping times satisfying (3) and with pairwise disjoint graphs.*

**Remark 2.1.** *We note that any finite totally inaccessible random time cannot coincide with positive probability with an independent random time.*

Let us state Yoeurp's result about orthogonal decomposition of a local martingale.

**Theorem 2.2.** *(Theorem 1-4 in [40])*

*If  $Z \in \mathcal{M}_{loc,0}(R, \mathbb{A})$  then  $Z$  can be uniquely decomposed as*

$$Z = Z^c + Z^{dp} + Z^{dq}, \quad (4)$$

*where  $Z^c$ ,  $Z^{dp}$  and  $Z^{dq}$  belong to  $\mathcal{M}_{loc,0}(R, \mathbb{A})$  and  $Z^c$  has continuous trajectories,  $Z^{dp}$  has accessible jump times only and is strongly orthogonal to any local martingale with at most totally inaccessible jump times,  $Z^{dq}$  has totally inaccessible jump times only and is strongly orthogonal to any local martingale with at most accessible jump times.*

As usual we refer to  $Z^c$ ,  $Z^{dp}$  and  $Z^{dq}$  as the *continuous*, the *accessible martingale part* and the *totally inaccessible martingale part* of  $Z$ , respectively. We recall that  $Z^{dp}$  and  $Z^{dq}$  are purely discontinuous local martingales (see Definition 4.11, page 40, in [29]). Observe that to any jump time of  $Z^{dp}$  ( $Z^{dq}$ ) corresponds the accessible (totally inaccessible) component of a jump time of  $Z$  and viceversa.

It is worthwhile to stress that the above decomposition depends on the reference filtration. In fact the nature of a random time is linked to the choice of the filtration and in particular *accessibility is preserved by enlarging the filtration and viceversa total inaccessibility is preserved under restriction of the filtration.*

The crucial notion of this paper is the *strong predictable representation property* of a local martingale. We refer to [15] for the definition of the *vector stochastic integral* and its relation with the *componentwise stochastic integral*. We recall that the two notions coincide when the components of the driving local martingale are pairwise strongly orthogonal (see Theorem 3.1 in [15]).

**Definition 2.3.** (Definition 13.1, page 362, in [26])

$\mathbf{Z} = (Z_1, \dots, Z_m)$  with  $Z_i \in M_{loc}(R, \mathbb{A})$ ,  $i = 1, \dots, m$ , enjoys the  $(R, \mathbb{A})$ -strong predictable representation property  $((R, \mathbb{A})$ -PRP) if each  $V \in M_{loc,0}(R, \mathbb{A})$  can be represented as vector stochastic integral w.r.t.  $\mathbf{Z}$ .

Next theorem is known as *II Fundamental Theorem of Asset Pricing*.

**Theorem 2.4.** (Theorem 13.9, page 366, in [26])

Let  $\mathcal{A}_0$  be trivial. Then  $\mathbf{Z} \in \mathcal{M}_{loc}(R, \mathbb{A})$  enjoys the  $(R, \mathbb{A})$ -PRP if and only if  $\mathbb{P}(\mathbf{Z}, \mathbb{A})$  is a singleton.

By the above theorem and the comment to Corollary 11.4, page 340, in [27] it follows that  $\mathbf{Z} = (Z_1, \dots, Z_m)$ , with  $Z_i \in \mathcal{M}_{loc}^2(R, \mathbb{A})$ ,  $i = 1, \dots, m$ , enjoys the  $(R, \mathbb{A})$ -PRP if and only for each  $V \in \mathcal{M}^2(R, \mathbb{A})$  holds

$$V_t = V_0 + (\xi^V \cdot \mathbf{Z})_t,$$

where  $V_0$  is  $\mathcal{A}_0$ -measurable,  $\xi^V = (\xi_1^V, \dots, \xi_m^V)$  is an  $m$ -dimensional predictable process such that

$$E^R \left[ \sum_{i,j} \int_0^T \xi_i^V(t) \xi_j^V(t) d[Z_i, Z_j]_t \right] < +\infty \quad (5)$$

and  $(\xi^V \cdot \mathbf{Z})_t$  denotes the vector stochastic integral of  $\xi^V$ .

The isometry between  $\mathcal{M}^2(R, \mathbb{A})$  and  $L^2(\Omega, \mathcal{A}_T, R)$  implies that, if  $\mathbf{Z} = (Z_1, \dots, Z_m)$ , with  $Z_i \in \mathcal{M}_{loc}^2(R, \mathbb{A})$ ,  $i = 1, \dots, m$ , enjoys the  $(R, \mathbb{A})$ -PRP, then each  $K \in L^2(\Omega, \mathcal{A}_T, R)$  satisfies  $K = K_0 + (\xi^K \cdot \mathbf{Z})_T$ , where  $K_0$  is  $\mathcal{A}_0$ -measurable and  $\xi^K$  is an  $m$ -dimensional predictable process such that the analogous of (5) holds (see pages 27-28 in [27]).

To conclude the *PRP is invariant by change of probability measures* in the following sense. For a probability measure  $\tilde{R}$  equivalent to  $R|_{\mathcal{A}_T}$ , set

$$L_t := \frac{d\tilde{R}}{dR} \Big|_{\mathcal{A}_t}, \quad t \leq T.$$

$L = (L_t)_{t \in [0, T]}$  is a positive  $(R, \mathbb{A})$ -martingale and the following result holds.

**Proposition 2.5.** (Lemma 2.5 in [31])

Let  $\mathbf{Z} = (Z_1, \dots, Z_m)$ , with  $Z_i \in \mathcal{M}_{loc}(R, \mathbb{A})$ , enjoy the  $(R, \mathbb{A})$ -PRP. Assume that for any component  $Z^i$  of  $\mathbf{Z}$  there exists  $\langle L, Z^i \rangle^{R, \mathbb{A}}$ . Set

$$\tilde{Z}_t^i := Z_t^i - \int_0^t \frac{1}{L_{s-}} d\langle L, Z^i \rangle_s^{R, \mathbb{A}}.$$

Then  $\tilde{\mathbf{Z}}$  enjoys the  $(\tilde{R}, \mathbb{A})$ -PRP.

Now we introduce the notion of *Jacod's dimension of the set  $\mathcal{H}^1(R, \mathbb{A})$* , where

$$\mathcal{H}^1(R, \mathbb{A}) := \{M \in \mathcal{M}(R, \mathbb{A}) : E^R[\sup_{t \in [0, T]} |M_t|] < +\infty\}.$$

Let  $\mathbf{Z}$  be a vector local martingale. We denote by  $\mathcal{Z}^1(\mathbf{Z}, R)$  the *stable subspace in  $\mathcal{H}^1(R, \mathbb{A})$  generated by  $\mathbf{Z}$*  (see Definition (4.4), page 114, in [27]).

**Definition 2.6.** (Definition (4.38), pages 130-131, in [27])

$\dim(\mathcal{H}^1(R, \mathbb{A}))$  is the minimal dimension of a 1-generator of  $\mathcal{H}^1(R, \mathbb{A})$ , where a 1-generator of  $\mathcal{H}^1(R, \mathbb{A})$  is any vector local martingale  $\mathbf{Z}$  such that  $\mathcal{Z}^1(\mathbf{Z}, R) = \mathcal{H}_0^1(R, \mathbb{A})$ .

By Theorem 13.4, page 363, in [26] it follows immediately that  $\dim(\mathcal{H}^1(R, \mathbb{A}))$  is the minimal possible dimension of a vector-valued local martingale enjoying the  $(R, \mathbb{A})$ -PRP.

**Proposition 2.7.** (Proposition 2.10 in [2])

If  $\tilde{R}$  is a probability measure on  $(\Omega, \mathcal{A})$  equivalent to  $R$ , then  $\dim(\mathcal{H}^1(\tilde{R}, \mathbb{A})) = \dim(\mathcal{H}^1(R, \mathbb{A}))$ .

For the definition of *basis* and *multiplicity* of a filtration we follow [19].

**Definition 2.8.** An  $(R, \mathbb{A})$ -basis is a (at most countable) subset  $\{Z_1, Z_2, \dots\}$  of  $\mathcal{M}^2(R, \mathbb{A})$  whose elements are pairwise strongly orthogonal and such that each  $V \in \mathcal{M}^2(R, \mathbb{A})$  satisfies

$$V_t = V_0 + \sum_i \int_0^t \Phi_i^V(s) dZ_i(s)$$

where  $\Phi_i^V$  is a predictable process such that  $E^R \left[ \int_0^T (\Phi_i^V)^2(s) d[Z_i]_s \right] < +\infty$  and  $V_0$  is  $\mathcal{A}_0$ -measurable.

**Remark 2.9.** If  $\mathbf{Z} = (Z_1, \dots, Z_m)$  with  $Z_i \in \mathcal{M}^2(R, \mathbb{A})$  enjoys the  $(R, \mathbb{A})$ -PRP and has pairwise strongly orthogonal components then  $(Z_1, \dots, Z_m)$  is a  $(R, \mathbb{A})$ -basis.

**Definition 2.10.** The multiplicity of the filtration  $\mathbb{A}$  under the measure  $R$  is the smallest integer  $k \in \mathbb{N} \cup \{+\infty\}$  such that there exists an  $(R, \mathbb{A})$ -basis of dimension  $k$ .

For stating our main result we will need the notion of *minimal martingale measure*.

**Definition 2.11.** (Definition 1 in [5])

Let  $Y$  be a special semi-martingale with local martingale part,  $Z$ . A measure  $Q$  in  $\mathbb{P}(Y, \mathbb{A})$  is the minimal martingale measure for  $Y$  under  $R$  if  $Y$  is a  $(Q, \mathbb{A})$ -martingale,  $Q|_{\mathcal{A}_0} = R|_{\mathcal{A}_0}$  and any  $V \in \mathcal{M}_{loc}^2(R, \mathbb{A})$  orthogonal to  $Z$  belongs to  $\mathcal{M}_{loc}(Q, \mathbb{A})$ .

Finally we provide a result linking the regularity of a special semi-martingale to the regularity of the elements of its canonical decomposition.

**Proposition 2.12.** Let  $Y$  be a square-integrable special semi-martingale and let  $Z$  be its local martingale part. Then  $Z \in \mathcal{M}^2(R, \mathbb{A})$ .

*Proof.* The regularity of  $Y$  implies that its quadratic variation process  $[Y]$  is integrable (see Theorem 2, page 246, in [38]) and by Theorem VII-55, page 245, in [21] it follows that  $[Z]$  is integrable. Then the thesis follows applying Theorem 11.4.5, page 243, in [18].  $\square$

## 2.1 The sharp bracket of the accessible part of a martingale

It is well-known that any adapted process of locally integrable variation admits a unique dual predictable projection also called (*predictable*) *compensator* (see Theorem VI-80, page 139, in [21]).

Here we explicitly provide the compensator of a pure jump process with accessible jump times only and we use its expression to compute the sharp bracket of the accessible part of any square-integrable martingale.



**Lemma 2.13.** *Consider an  $\mathbb{A}$ -adapted pure jump process  $U$  of integrable variation with accessible jump times only. Then its  $(R, \mathbb{A})$ -compensator  $B^U$  admits the representation*

$$B_t^U = \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}] \mathbb{I}_{\eta_{n,m} \leq t} \quad (6)$$

where  $\{\eta_n\}_{n \in \mathbb{N}}$  is the sequence of the jump times of  $U$  and, for each  $n \in \mathbb{N}$ ,  $\{\eta_{n,m}\}_{m \in \mathbb{N}}$  is an enveloping sequence of  $\eta_n$ .

*Proof.*  $U$  satisfies

$$U_t = \sum_{s \leq t} \Delta U_s = \sum_{n \in \mathbb{N}} \Delta U_{\eta_n} \mathbb{I}_{\eta_n \leq t} = \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} \Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mathbb{I}_{\eta_{n,m} \leq t}. \quad (7)$$

Moreover for any  $\mathbb{A}$ -predictable stopping time  $\eta$  it has to

$$\Delta B_\eta^U = E^R [\Delta U_\eta \mid \mathcal{A}_{\eta-}]$$

(see Theorem VI-76, page 136, in [21]). Observe that the right hand side of the above equality makes sense by the integrability assumption on  $U$ . We stress that the same assumption provides the integrability conditions necessary for the rest of the proof.

Equality (6) is proved once we show that

$$U_t - \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}] \mathbb{I}_{\eta_{n,m} \leq t}, \quad t \in [0, T]$$

is an  $(R, \mathbb{A})$ -martingale. In fact, since the second addend in the previous expression is an  $\mathbb{A}$ -predictable process (see Lemma 22.3 ii), page 411, in [34]), the statement follows by the uniqueness of the compensator.

Using representation (7), for any  $s \leq t$ , we get

$$\begin{aligned} & E^R \left[ U_t - \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}] \mathbb{I}_{\eta_{n,m} \leq t} \mid \mathcal{A}_s \right] = \\ & U_s - \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}] \mathbb{I}_{\eta_{n,m} \leq s} + \\ & E^R \left[ \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} (\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} - E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}]) \mathbb{I}_{s < \eta_{n,m} \leq t} \mid \mathcal{A}_s \right]. \end{aligned}$$

Last term in previous expression can be written as

$$\sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} E^R \left[ \left( \Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} - E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}] \right) \mathbb{I}_{\eta_{n,m} \leq t} \mid \mathcal{A}_s \right] \mathbb{I}_{s < \eta_{n,m}}$$

Since  $\sigma(\eta_{n,m}) \subset \mathcal{A}_{\eta_{n,m}-}$  (see Chapter III Theorem 3.4 point 1, page 80, in [26]), then the general term of last sum is null if and only if

$$\begin{aligned} & E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mathbb{I}_{\eta_{n,m} \leq t} \mid \mathcal{A}_s] \mathbb{I}_{s < \eta_{n,m}} = \\ & = E^R \left[ E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mathbb{I}_{\eta_{n,m} \leq t} \mid \mathcal{A}_{\eta_{n,m}-}] \mid \mathcal{A}_s \right] \mathbb{I}_{s < \eta_{n,m}}, \end{aligned}$$

that is if and only if for any set  $A \in \mathcal{A}_s$

$$\begin{aligned} & \int_{A \cap (s < \eta_{n,m})} E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mathbb{I}_{\eta_{n,m} \leq t} \mid \mathcal{A}_s] dR = \\ & = \int_{A \cap (s < \eta_{n,m})} E^R \left[ E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mathbb{I}_{\eta_{n,m} \leq t} \mid \mathcal{A}_{\eta_{n,m}-}] \mid \mathcal{A}_s \right] dR. \end{aligned}$$

Last equality is true. In fact,  $A \cap (s < \eta_{n,m})$  belongs either to  $\mathcal{A}_s$  or to  $\mathcal{A}_{\eta_{n,m}-}$ , so that both expressions coincide with

$$\int_{A \cap (s < \eta_{n,m})} \Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mathbb{I}_{\eta_{n,m} \leq t} dR.$$

□

**Remark 2.14.** The process  $B^U$ , although unique, has different representations according to the choice of the enveloping sequences of the jump times of  $U$ . Nevertheless the representation of  $B^U$  as multivariate point process is unique and is obtained from (6), whichever family  $\{\{\eta_{n,m}\}_{m \in \mathbb{N}}, n \in \mathbb{N}\}$  is chosen. More precisely,

$$B_t^U = \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} E^R [\Delta U_{\tilde{\eta}_{n,m}} \mathbb{I}_{\eta_n = \tilde{\eta}_{n,m}} \mid \mathcal{A}_{\tilde{\eta}_{n,m}-}] \mathbb{I}_{\tilde{\eta}_{n,m} \leq t}, \quad (8)$$

where

$$\tilde{\eta}_{n,m} = \begin{cases} \eta_{n,m} & \text{if } E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}] \neq 0; \\ +\infty & \text{if } E^R [\Delta U_{\eta_{n,m}} \mathbb{I}_{\eta_n = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}] = 0, \end{cases}$$

and the sequence of increasing stopping times and marks of  $B^U$  is identified by reordering the set  $\{\tilde{\eta}_{n,m}\}_{n,m \in \mathbb{N}}$ . Note that, fixed  $n$ , the set  $\{\tilde{\eta}_{n,m}\}_{m \in \mathbb{N}}$  is an enveloping sequence of  $\eta_n$ . In particular, for any  $m$ ,  $\tilde{\eta}_{n,m}$  is a predictable random time by Proposition 2.10, page 17, in [29].

**Proposition 2.15.** If  $Z \in \mathcal{M}^2(R, \mathbb{A})$ , then

$$\langle Z^{dp} \rangle_t^{R, \mathbb{A}} = \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} E^R [(\Delta Z_{\eta_{n,m}})^2 \mathbb{I}_{\eta_n^{dp} = \eta_{n,m}} \mid \mathcal{A}_{\eta_{n,m}-}] \mathbb{I}_{\eta_{n,m} \leq t},$$

where  $Z^{dp}$  is the accessible martingale part of  $Z$  in the decomposition (4),  $\{\eta_n^{dp}\}_{n \in \mathbb{N}}$  is the sequence of the jump times of  $Z^{dp}$  and, for each  $n \in \mathbb{N}$ ,  $\{\eta_{n,m}\}_{m \in \mathbb{N}}$  is an enveloping sequence of  $\eta_n^{dp}$ .

*Proof.* We recall that  $\langle Z^{dp} \rangle^{R, \mathbb{A}}$  exists since  $Z^{dp} \in \mathcal{M}^2(R, \mathbb{A})$  and it is the compensator of  $[Z^{dp}]$ . Therefore it is enough to apply previous lemma to  $[Z^{dp}]$ , which is the adapted purely discontinuous integrable increasing process with accessible jump times only which satisfies

$$[Z^{dp}]_t = \sum_{s \leq t} (\Delta Z_s^{dp})^2 = \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} (\Delta Z_{\eta_{n,m}}^{dp})^2 \mathbb{I}_{\eta_{n,m} \leq t} = \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} (\Delta Z_{\eta_{n,m}})^2 \mathbb{I}_{\eta_{n,m} \leq t}.$$

□



## 2.2 The compensated occurrence process of a random time

Let  $\eta$  be a random time in  $[0, +\infty]$  and consider the occurrence process of  $\eta$  restricted to  $[0, T]$ , that is the process

$$\mathbb{I}_{\eta \leq \cdot} := (\mathbb{I}_{\eta \leq t})_{t \in [0, T]}.$$

It is a pure jump bounded process with a unique jump time corresponding to

$$\eta \mathbb{I}_{\eta \leq T} + \infty \mathbb{I}_{\eta > T}.$$

In the rest of the paper, for notational convenience, we will identify it with  $\eta$ .

Define

$$\mathcal{H}_t^\eta := \sigma(\eta \wedge t).$$

Then  $\mathbb{H}^\eta = (\mathcal{H}_t^\eta)_{t \in [0, T]}$  is the natural filtration of  $\mathbb{I}_{\eta \leq \cdot}$ , that is the smallest filtration which makes  $\eta$  a stopping time. Let  $A^{\eta, R, \mathbb{H}^\eta}$  be the  $(R, \mathbb{H}^\eta)$ -compensator of  $\mathbb{I}_{\eta \leq \cdot}$  or simply the (natural) compensator of  $\eta$  and let  $H^\eta = (H_t^\eta)_{t \in [0, T]}$  be the (naturally) compensated occurrence process of  $\eta$ , that is the  $(R, \mathbb{H}^\eta)$ -martingale defined by

$$H_t^\eta := \mathbb{I}_{\eta \leq t} - A_t^{\eta, R, \mathbb{H}^\eta}. \quad (9)$$

**Remark 2.16.** It is to stress that any random time  $\eta$  in  $[0, T] \cup \{+\infty\}$ , according to its law  $\mu^\eta$ , is an  $\mathbb{H}^\eta$ -totally inaccessible stopping time if and only if  $\mu^\eta$  restricted to  $[0, T]$  is a diffusive measure and it is an  $\mathbb{H}^\eta$ -accessible stopping time if and only if  $\mu^\eta$  is atomic. This is a trivial generalization of Theorem IV-107, page 241, in [20], which only covers the case when  $\eta$  is finite. Therefore  $\eta$ , unless trivial, can never be  $\mathbb{H}^\eta$ -predictable, so that the martingale  $H^\eta$  cannot be identically null.

According to (1) we denote by  $\eta^{dp}$  and  $\eta^{dq}$  the  $\mathbb{H}^\eta$ -accessible and the  $\mathbb{H}^\eta$ -totally inaccessible component of  $\eta$ , respectively. Next proposition highlights their different contribution in the expression of the compensator of  $\eta$ .

**Proposition 2.17.** The compensated occurrence process of  $\eta$  admits the representation

$$H_t^\eta = \mathbb{I}_{\{\eta \leq t\}} - A_t^{dq, R, \mathbb{H}^\eta} - \sum_{m \in \mathbb{N}} R(\eta^{dp} = \eta_m^{dp} \mid \mathcal{H}_{\eta_m^{dp}-}^\eta) \mathbb{I}_{\{\eta_m^{dp} \leq t\}}, \quad (10)$$

where  $A^{dq, R, \mathbb{H}^\eta}$  denotes the (continuous)  $(R, \mathbb{H}^\eta)$ -compensator of  $\eta^{dq}$  and  $(\eta_m^{dp})_{m \in \mathbb{N}}$  is an enveloping sequence of  $\eta^{dp}$ .

*Proof.* We start by the simple equality

$$\mathbb{I}_{\eta \leq t} = \mathbb{I}_{\eta = \eta^{dq}} \mathbb{I}_{\eta^{dq} \leq t} + \mathbb{I}_{\eta = \eta^{dp}} \mathbb{I}_{\eta^{dp} \leq t}. \quad (11)$$

Observing that

$$\{\eta^{dq} \leq t\} \subset \{\eta = \eta^{dq}\}, \quad \{\eta^{dp} \leq t\} \subset \{\eta = \eta^{dp}\},$$

we get

$$\mathbb{I}_{\eta \leq t} = \mathbb{I}_{\eta^{dq} \leq t} + \mathbb{I}_{\eta^{dp} \leq t}. \quad (12)$$

In analogy with the notation  $A^{dq, R, \mathbb{H}^\eta}$ , we indicate with  $A^{dp, R, \mathbb{H}^\eta}$  the  $(R, \mathbb{H}^\eta)$ -compensator of  $\eta^{dp}$ . Then by the uniqueness of the compensator of  $\eta$  we derive

$$A^{\eta, R, \mathbb{H}^\eta} = A^{dq, R, \mathbb{H}^\eta} + A^{dp, R, \mathbb{H}^\eta}. \quad (13)$$

Therefore

$$H^\eta = (\mathbb{I}_{\eta^{dq} \leq \cdot} - A^{dq, R, \mathbb{H}^\eta}) + (\mathbb{I}_{\eta^{dp} \leq \cdot} - A^{dp, R, \mathbb{H}^\eta}), \quad (14)$$

that is

$$H^\eta = \mathbb{I}_{\eta \leq \cdot} - A^{dq, R, \mathbb{H}^\eta} - A^{dp, R, \mathbb{H}^\eta}.$$

Finally Lemma 2.13 yields

$$A_t^{dp, R, \mathbb{H}^\eta} = \sum_{m \in \mathbb{N}} R(\eta^{dp} = \eta_m^{dp} \mid \mathcal{H}_{\eta_m^{dp}-}) \mathbb{I}_{\{\eta_m^{dp} \leq t\}}. \quad (15)$$

□

We stress that, since the two martingales at the right-hand side of (14) are strongly orthogonal, by the uniqueness of Yoeurp's decomposition (4), the  $\mathbb{H}$ -totally inaccessible and the  $\mathbb{H}$ -accessible martingale parts of  $H$  satisfy the equalities

$$H_t^{dq} = \mathbb{I}_{\eta^{dq} \leq t} - A_t^{dq, R, \mathbb{H}^\eta}, \quad H_t^{dp} = \mathbb{I}_{\eta^{dp} \leq t} - A_t^{dp, R, \mathbb{H}^\eta}.$$

We conclude recalling a key result to be used in the rest of the paper.

**Theorem 2.18.** (*Proposition 2 in [16]*)

$H$  enjoys the  $(R, \mathbb{H}^\eta)$ -PRP.

### 2.3 Mutual singularity of the kernels associated with two predictable increasing processes

In this subsection we prove a result on the supports of the random measures generated by two increasing processes, which will be the core of the study of the Jacod's dimension of the space of martingales on the enlarged filtration  $\mathbb{G}$ .

Let us start with two definitions given in analogy with those at pages 19 and 374 in [9], respectively.

**Definition 2.19.** A support of a measure  $\mu$  on a measurable space  $(E, \mathcal{E})$  is any set  $C \in \mathcal{E}$  such that  $\mu(E \setminus C) = 0$ .

**Definition 2.20.** Two measures  $\mu$  and  $\nu$  on the measurable space  $(E, \mathcal{E})$  are mutually singular if they admit disjoint supports, that is if there exist two measurable disjoint sets  $C^\mu$  and  $C^\nu$  such that  $\mu(E \setminus C^\mu) = 0$  and  $\nu(E \setminus C^\nu) = 0$ .

**Remark 2.21.** If  $\mu$  and  $\nu$  are mutually singular with disjoint supports  $C^\mu$  and  $C^\nu$ , then  $\mu(C^\nu) = \nu(C^\mu) = 0$ .

We call *random set* any subset of the product space  $\Omega \times [0, T]$ , *measurable random set* any random set belonging to the product sigma-algebra  $\mathcal{A} \otimes \mathcal{B}([0, T])$  and *predictable random set* any random set belonging to the predictable sigma algebra  $\mathcal{P}(\mathbb{A})$  on  $\Omega \times [0, T]$  (see pages 3 and 16 in [29]). Finally given a random set  $C$  we denote by  $\overline{C}$  its complementary set and by  $C^\omega$  its  $\omega$ -section, that is

$$C^\omega := \{t, (t, \omega) \in C\}. \quad (16)$$

Moreover we also introduce next definition (see pages 29 and 30 in [35]).

**Definition 2.22.** A function  $\kappa : \Omega \times \mathcal{B}([0, T]) \rightarrow \mathbb{R}^+$ ,  $(\omega, G) \rightarrow \kappa(\omega, G)$ , is a kernel from  $\Omega$  to  $[0, T]$  if and only if, for any fixed  $G \in \mathcal{B}([0, T])$ ,  $\omega \rightarrow \kappa(\omega, G)$  is  $\mathcal{A}$ -measurable and, for any fixed  $\omega$ ,  $G \rightarrow \kappa(\omega, G)$  is a measure on  $([0, T], \mathcal{B}([0, T]))$ .

Two kernels  $\kappa^i$ ,  $i = 1, 2$ , from  $\Omega$  to  $[0, T]$  are mutually singular if there exist two disjoint measurable random sets  $\Gamma_i$ ,  $i = 1, 2$ , with  $\Gamma_i^\omega$  support of the measure  $\kappa^i(\omega, \cdot)$ , that is  $\kappa^i(\omega, \overline{\Gamma_i^\omega}) = 0$ ,  $i = 1, 2$ .

**Remark 2.23.** Observe that any increasing process  $(V_t)_{t \in [0, T]}$  uniquely defines a kernel  $dV : \Omega \times \mathcal{B}([0, T]) \rightarrow \mathbb{R}^+$ ,  $(\omega, G) \rightarrow \int_G dV_s(\omega)$  (for further details see VI-86, page 145, in [21]).

**Proposition 2.24.** Let  $(A_t)_{t \in [0, T]}$  and  $(B_t)_{t \in [0, T]}$  be two non-negative increasing predictable processes. Assume that the associated kernels  $dA$  and  $dB$  are mutually singular. Then there exists a predictable random set  $C^A$  such that,  $R$ -a.s.,  $C^{A, \omega}$  and  $\overline{C^{A, \omega}}$  are disjoint supports of the measures  $dA(\omega, \cdot)$  and  $dB(\omega, \cdot)$ , respectively.

*Proof.* By assumption there exist two disjoint measurable random sets  $\Gamma^A$  and  $\Gamma^B$  such that, for any  $\omega$ ,  $dA(\omega, \overline{\Gamma^{A, \omega}}) = dB(\omega, \overline{\Gamma^{B, \omega}}) = 0$  and in particular

$$dB(\omega, \Gamma^{A, \omega}) = 0. \quad (17)$$

First, by means of a projection procedure, we prove that it is possible to construct a predictable random set  $C^A$  whose  $\omega$ -sections are  $R$ -a.s. supports of the measures  $dA(\omega, \cdot)$ . Then, we prove that  $R$ -a.s. the  $\omega$ -section of the predictable random set  $\overline{C^A}$  is support of the measure  $dB(\omega, \cdot)$  and, since  $(\overline{C^A})^\omega = \overline{C^{A, \omega}}$ , this ends the proof. Let us be more precise.

We construct  $C^A$  as the *predictable support* of the measurable random set  $\Gamma^A$ , that is

$$C^A := \{(\omega, t), \text{ } {}^p\mathbb{I}_{\Gamma^A}(\omega, t) > 0\}, \quad (18)$$

where  ${}^p\mathbb{I}_{\Gamma^A}$  denotes the predictable projection of the process  $\mathbb{I}_{\Gamma^A}$  (see Chapter 1 Definition 2.32, page 24, in [29]). Now we prove that  $R$ -a.s. the  $\omega$ -section  $C^{A, \omega}$  is a support of the measure  $dA(\omega, \cdot)$ .

To this end it suffices to establish that

$$\int_{\overline{C^{A, \omega}}} dA_s(\omega) = 0, \quad R\text{-a.s.} \quad (19)$$

In fact  $\int_{\overline{C^{A, \omega}}} dA_s(\omega) \geq 0$  and moreover the following equalities hold

$$E^R \left[ \int_{\overline{C^{A, \cdot}}} dA_s(\cdot) \right] = E^R \left[ \int_{\overline{C^{A, \cdot}}} \mathbb{I}_{\Gamma^A}(\cdot, s) dA_s(\cdot) \right] = E^R \left[ \int_{[0, T]} \overline{\mathbb{I}_{\Gamma^A}}(s) \mathbb{I}_{\Gamma^A}(\cdot, s) dA_s(\cdot) \right]. \quad (20)$$

Observe that for any  $(\omega, t) \in \Omega \times [0, T]$  it holds

$$\overline{\mathbb{I}_{\Gamma^A}}(t) = \mathbb{I}_{\overline{\Gamma^A}}(\omega, t).$$

Then the last term in (20) coincides with

$$E^R \left[ \int_{[0, T]} \zeta(\cdot, s) dA_s(\cdot) \right], \quad (21)$$

where  $\zeta(\omega, t) := \mathbb{I}_{\overline{C^A}}(\omega, t) \mathbb{I}_{\Gamma^A}(\omega, t)$ . Again, due to the fact that, by assumption, the process  $A$  is predictable, the expression in (21) turns to be equal to (see Theorem VI.57, page 122, in [21])

$$E^R \left[ \int_{[0,T]} {}^p\zeta(\cdot, s) dA_s(\cdot) \right],$$

where, being  $\overline{C^A}$  a predictable random set,

$${}^p\zeta(\omega, t) = \mathbb{I}_{\overline{C^A}}(\omega, t) {}^p\mathbb{I}_{\Gamma^A}(\omega, t)$$

(see Chapter 1 Theorem 2.28 c) in [29]). Then

$$E^R \left[ \int_{[0,T]} {}^p\zeta(\cdot, s) dA_s(\cdot) \right] = E^R \left[ \int_{(\overline{C^A})} {}^p\mathbb{I}_{\Gamma^A}(\cdot, s) dA_s(\cdot) \right].$$

Since by definition for any  $\omega$  the function the function  ${}^p\mathbb{I}_{\Gamma^A}(\omega, \cdot)$  is null on the set  $\overline{C^{A,\omega}}$ , the right hand side in previous equality is equal to zero, so that (19) follows.

Let us prove that  $\overline{C^{A,\omega}}$  is a support of  $dB(\omega, \cdot)$ . This follows as soon as we show that  $R$ -a.s.

$$\int_{C^{A,\omega}} dB_s(\omega) = 0. \quad (22)$$

In fact

$$0 = E^R \left[ \int_{\Gamma^A} dB_s \right] = E^R \left[ \int_{[0,T]} \mathbb{I}_{\Gamma^A, \cdot}(s) dB_s(\cdot) \right] = E^R \left[ \int_{[0,T]} {}^p\mathbb{I}_{\Gamma^A}(\cdot, s) dB_s(\cdot) \right],$$

where last equality again follows by Theorem VI.57 in [21]. Therefore

$$E^R \left[ \int_{C^{A,\cdot}} {}^p\mathbb{I}_{\Gamma^A}(\cdot, s) dB_s(\cdot) \right] = 0$$

which is equivalent to  $\int_{C^{A,\omega}} {}^p\mathbb{I}_{\Gamma^A}(\omega, s) dB_s(\omega) = 0$ ,  $R$ -a.s. Since by definition, fixed  $\omega$ ,  ${}^p\mathbb{I}_{\Gamma^A}(\omega, s) > 0$  for all  $s \in C^{A,\omega}$ , then (22) follows,  $R$ -a.s.  $\square$

### 3 The setting

In the following we introduce the setting we will work in looking for a new martingale representation theorem.

From now on we will consider a fixed probability space  $(\Omega, \mathcal{F}, P)$  and two standard filtrations on it,  $\mathbb{F} = (\mathcal{F}_t)_{t \in [0,T]}$  and  $\mathbb{H} = (\mathcal{H}_t)_{t \in [0,T]}$ , with trivial initial  $\sigma$ -algebras and such that  $\mathcal{F}_T \subset \mathcal{F}$  and  $\mathcal{H}_T \subset \mathcal{F}$ . We will set

$$\mathbb{G} := \mathbb{F} \vee \mathbb{H}. \quad (23)$$

From now on we will standing assume the following condition (D).

(D) There exists an *equivalent decoupling measure* that is a probability measure  $Q$  on

$(\Omega, \mathcal{G}_T)$  equivalent to  $P|_{\mathcal{G}_T}$  under which  $\mathcal{F}_T$  and  $\mathcal{H}_T$  are independent.

Then also  $\mathbb{G}$  under  $P$  is a standard filtration (see Lemma 2.2 in [4]).

Let  $M = (M_t)_{t \in [0, T]}$  and  $N = (N_t)_{t \in [0, T]}$  be a square-integrable  $(P, \mathbb{F})$ -martingale and a square-integrable  $(P, \mathbb{H})$ -martingale respectively. We define on  $(\Omega, \mathcal{G}_T)$  the *martingale preserving measure*  $P^*$  associated to  $Q$  by

$$dP^* := \frac{dP}{dQ} \Big|_{\mathcal{F}_T} \cdot \frac{dP}{dQ} \Big|_{\mathcal{H}_T} dQ. \quad (24)$$

Then  $M$  and  $N$  are independent  $(P^*, \mathbb{G})$ -martingales and each of them preserves under  $P^*$  its law since

$$P^*|_{\mathcal{F}_T} = P|_{\mathcal{F}_T} \quad P^*|_{\mathcal{H}_T} = P|_{\mathcal{H}_T}. \quad (25)$$

In the light of Theorem 2.2 we consider the  $(P^*, \mathbb{G})$ -decompositions of  $M$  and  $N$

$$M = M^c + M^{dp} + M^{dq}, \quad N = N^c + N^{dp} + N^{dq}. \quad (26)$$

**Proposition 3.1.** *All  $\mathbb{F}$ -totally inaccessible stopping times and all  $\mathbb{H}$ -totally inaccessible stopping times are also  $\mathbb{G}$ -totally inaccessible.*

*Proof.* It is enough to prove the statement for  $\mathbb{F}$  and  $M$ .

Let  $\eta$  be an  $\mathbb{F}$ -totally inaccessible stopping time. Then, since condition (D) is in force, Remark 2.1 ensures  $\eta$  to avoid  $\mathbb{H}$ -stopping times and therefore  $\eta$  turns out to be  $\mathbb{G}$ -totally inaccessible (see Lemma 3.5 in [17]).  $\square$

Recalling that all  $\mathbb{F}$ -accessible random times are also  $\mathbb{G}$ -accessible, by previous proposition we immediately derive next result.

**Corollary 3.2.** *The  $(P, \mathbb{F})$ -decomposition of  $M$  and the  $(P, \mathbb{H})$ -decomposition of  $N$  coincide with the decompositions given in (26).*

**Proposition 3.3.** *For all  $t$  in  $[0, T]$  it holds  $P^*$ -a.s. and  $P$ -a.s.*

$$[M, N]_t = [M^{dp}, N^{dp}]_t. \quad (27)$$

*Proof.* By the representation formula (26), the linearity of the covariation operator and its invariance under equivalent changes of measures it follows  $P^*$ -a.s. and  $P$ -a.s.

$$[M, N] = [M^c, N] + [M^{dp}, N] + [M^{dq}, N].$$

In particular  $M^c$  and  $N$  as well as  $M^{dq}$  and  $N$  are  $(P^*, \mathbb{G})$ -independent martingales. Then, as well known, one has  $[M^c, N] = 0$  (see Theorem 2.2 and Corollary 4.55 points b) and c) in [29]). By Remark 2.1 it also holds  $[M^{dq}, N] = 0$ . Summarizing

$$[M, N] = [M^{dp}, N]$$

and similarly, by using the representation (26) for  $N$ , we get the thesis.  $\square$

It is worthwhile to note that (27) can be explicitly written as

$$[M, N]_t = [M^{dp}, N^{dp}]_t = \sum_{n,l} \Delta M_{\eta_n^{dp}}^{dp} \Delta N_{\tau_l^{dp}}^{dp} \mathbb{I}_{\eta_n^{dp} = \tau_l^{dp}} \mathbb{I}_{\eta_n^{dp} \leq t}, \quad (28)$$

where  $(\eta_n^{dp})_n$  and  $(\tau_l^{dp})_l$  are sequences of  $\mathbb{G}$ -accessible jump times of  $M$  and  $N$ , respectively.

In the previous proposition the covariation process  $[M, N]$  is derived by using the  $(P^*, \mathbb{G})$ -decompositions defined in (26), which coincide with the  $(P, \mathbb{F})$ -decomposition of  $M$  and the  $(P, \mathbb{H})$ -decomposition of  $N$  (see Corollary 3.2). It is to note that  $\mathbb{F}$  ( $\mathbb{H}$ ) could be different from the natural filtration  $\mathbb{F}^M$  of  $M$  ( $\mathbb{F}^N$  of  $N$ ). When using the natural filtrations different Yoeurp's decompositions for  $M$  and  $N$  could arise. As an example, a jump time of  $M$  could be a  $\mathbb{G}$ -accessible stopping time and an  $\mathbb{F}^M$ -totally inaccessible one, and this would imply that  $M^{dp} \neq M^{\mathbb{F}^M, dp}$ . However, as we will show now, the covariation process  $[M, N]$  still coincides with the covariation process of the natural accessible martingale parts.

**Lemma 3.4.** *Let  $\eta$  and  $\tau$  be an  $\mathbb{F}^M$ -stopping time and an  $\mathbb{F}^N$ -stopping time, respectively. If  $P(\eta = \tau) > 0$  then there exist an  $\mathbb{F}^M$ -accessible stopping time  $\eta^{dp}$  and an  $\mathbb{F}^N$ -accessible stopping time  $\tau^{dp}$  such that on the set  $(\eta = \tau)$  it holds  $\eta = \eta^{dp}$  and  $\tau = \tau^{dp}$ ,  $P$ -a.s.*

*Proof.* The representation of  $\eta$  given by (1) and (2) yields

$$\eta \mathbb{I}_{(\eta=\tau)} = \eta \mathbb{I}_{(\eta=\tau=+\infty)} + \eta \mathbb{I}_{(\eta=\tau) \cap A} + \eta \mathbb{I}_{(\eta=\tau) \cap B}.$$

Since  $A = (\eta = \eta^{dp} < +\infty)$  and  $B = (\eta = \eta^{dq} < +\infty)$  and therefore  $(\eta = \eta^{dp} = \eta^{dq} = +\infty)$  on  $(A \cup B)^c$ , one immediately derives that  $P$ -a.s.

$$\eta \mathbb{I}_{(\eta=\tau)} = \eta^{dp} \mathbb{I}_{(\eta=\tau=+\infty)} + \eta^{dp} \mathbb{I}_{(\eta^{dp}=\tau) \cap A} + \eta^{dq} \mathbb{I}_{(\eta^{dq}=\tau) \cap B}.$$

Consider now the equivalent decoupling measure  $Q$  introduced in condition (D). By Remark 2.1  $(\eta^{dq} = \tau) \cap B$  has null measure under  $Q$  so that a.s.

$$\eta \mathbb{I}_{(\eta=\tau)} = \eta^{dp} \mathbb{I}_{(\eta=\tau=+\infty)} + \eta^{dp} \mathbb{I}_{(\eta^{dp}=\tau) \cap A}$$

which implies  $\eta = \eta^{dp}$  on  $(\eta = \tau)$ .

Analogously one shows that  $\tau = \tau^{dp}$  on  $(\eta = \tau)$ . □

**Proposition 3.5.** *For all  $t$  in  $[0, T]$  it holds  $P^*$ -a.s. and  $P$ -a.s.*

$$[M, N]_t = [M^{\mathbb{F}^M, dp}, N^{\mathbb{F}^N, dp}]_t$$

where  $M^{\mathbb{F}^M, dp}$  and  $N^{\mathbb{F}^N, dp}$  are the accessible martingale parts in the  $(P, \mathbb{F}^M)$  and the  $(P, \mathbb{F}^N)$ -decompositions of  $M$  and  $N$ , respectively.

*Proof.* If  $\eta$  and  $\tau$  are jump times of  $M^{dp}$  and  $N^{dp}$ , respectively, then they are  $\mathbb{F}^M$ -stopping time and  $\mathbb{F}^N$ -stopping time, respectively. If  $P(\eta = \tau) > 0$  then, by Lemma 3.4, there exist an  $\mathbb{F}^M$ -accessible stopping time of  $M$ ,  $\hat{\eta}^{dp}$ , and an  $\mathbb{F}^N$ -accessible stopping time of  $N$ ,  $\hat{\tau}^{dp}$ , such that  $P$ -a.s.

$$\eta \mathbb{1}_{\eta=\tau} = \hat{\eta}^{dp} \mathbb{1}_{\eta=\tau} \text{ and } \tau \mathbb{1}_{\eta=\tau} = \hat{\tau}^{dp} \mathbb{1}_{\eta=\tau}.$$



Moreover  $\hat{\eta}^{dp}$  is a jump times of the accessible martingale part  $M^{\mathbb{F}^M, dp}$  in the  $(P, \mathbb{F}^M)$ -decomposition of  $M$  and  $\hat{\tau}^{dp}$  is a jump times of the accessible martingale part  $N^{\mathbb{F}^N, dp}$  in the  $(P, \mathbb{F}^N)$ -decomposition of  $N$ . Then using (28)

$$\begin{aligned} [M, N]_t &= \sum_{n,l} \mathbb{I}_{\hat{\eta}_n^{dp} \leq t} \mathbb{I}_{\hat{\eta}_n^{dp} = \hat{\tau}_l^{dp}} \Delta M_{\hat{\eta}_n^{dp}}^{dp} \Delta N_{\hat{\tau}_l^{dp}}^{dp} = \\ &= \sum_{n,l} \mathbb{I}_{\hat{\eta}_n^{dp} \leq t} \mathbb{I}_{\hat{\eta}_n^{dp} = \hat{\tau}_l^{dp}} \Delta M_{\hat{\eta}_n^{dp}}^{\mathbb{F}^M, dp} \Delta N_{\hat{\tau}_l^{dp}}^{\mathbb{F}^N, dp} = [M^{\mathbb{F}^M, dp}, N^{\mathbb{F}^N, dp}]_t, \end{aligned}$$

where  $(\hat{\eta}_n^{dp})_n$  is the set of all  $\mathbb{F}^M$ -accessible jump times of  $M$  and  $(\hat{\tau}_l^{dp})_l$  is the set of all  $\mathbb{F}^N$ -accessible jump times of  $N$ .  $\square$

**Remark 3.6.** Under condition (D) the covariation process  $[M, N]$  is identically zero whenever  $M$  or  $N$  are càdlàg quasi-left continuous martingales (see pages 121-122 in [26]). Among the others this is the case when at least one of the reference filtrations of  $M$  and  $N$  is quasi-left continuous. In fact any martingale with quasi-left continuous reference filtration is quasi-left continuous. In particular this happens if  $M$  or  $N$  belongs to the class of Lévy processes without deterministic jump times (see page 190 and Exercise 8, page 148, in [38]).

## 4 The Jacod's dimension of $\mathcal{H}^1(P, \mathbb{G})$

In this section we clarify how the Jacod's dimension of  $\mathcal{H}^1(P, \mathbb{G})$  is affected by the mutual behavior either of the sharp brackets of  $M$  and  $N$  or of their accessible parts,  $M^{dp}$  and  $N^{dp}$ .

**Lemma 4.1.** The kernels  $d\langle M^{dp} \rangle_t^{P, \mathbb{F}}$  and  $d\langle N^{dp} \rangle_t^{P, \mathbb{H}}$  are mutually singular if and only if the covariation process  $[M, N]$  is  $P$ -a.s. null.

*Proof.* Let  $\{\eta_n^{dp}\}_{n \in \mathbb{N}}$  and  $\{\tau_l^{dp}\}_{l \in \mathbb{N}}$  be the sequences of accessible jump times of  $M$  and of  $N$ , respectively, and, fixed  $n \in \mathbb{N}$  and  $l \in \mathbb{N}$ , let  $\{\eta_{n,m}\}_{m \in \mathbb{N}}$  and  $\{\tau_{l,h}\}_{h \in \mathbb{N}}$  be enveloping sequences of  $\eta_n^{dp}$  and  $\tau_l^{dp}$ , respectively.

Proposition 2.15 applies to give the following representations

$$\langle M^{dp} \rangle_t^{P, \mathbb{F}} = \sum_{n \in \mathbb{N}} \sum_{m \in \mathbb{N}} U_{n,m} \mathbb{I}_{\eta_{n,m} \leq t}, \quad \langle N^{dp} \rangle_t^{P, \mathbb{H}} = \sum_{l \in \mathbb{N}} \sum_{h \in \mathbb{N}} V_{l,h} \mathbb{I}_{\tau_{l,h} \leq t}, \quad (29)$$

with  $U_{n,m}$  and  $V_{l,h}$  defined by

$$U_{n,m} := E^P \left[ (\Delta M_{\eta_{n,m}}^{dp})^2 \mathbb{I}_{\eta_{n,m} = \eta_{n,m}} \mid \mathcal{F}_{\eta_{n,m}-} \right], \quad V_{l,h} := E^P \left[ (\Delta N_{\tau_{l,h}}^{dp})^2 \mathbb{I}_{\tau_{l,h} = \tau_{l,h}} \mid \mathcal{H}_{\tau_{l,h}-} \right].$$

Assume first the mutual singularity of the kernels  $d\langle M^{dp} \rangle_t^{P, \mathbb{F}}$  and  $d\langle N^{dp} \rangle_t^{P, \mathbb{H}}$ . Fixed  $n, m$  and  $l, h$ , when

$$P(\eta_{n,m} = \tau_{l,h} < +\infty) > 0,$$

representations (29) implies that

$$\mathbb{I}_{\{\eta_{n,m} = \tau_{l,h} < +\infty\}} U_{n,m} V_{l,h} = 0, \quad P\text{-a.s.} \quad (30)$$

In other words, if we introduce the measurable sets

$$A_{n,m} := \{U_{n,m} = 0\}, \quad B_{l,h} := \{V_{l,h} = 0\},$$

then by the mutual singularity the set

$$A_{n,m}^c \cap B_{l,h}^c \cap \{\eta_{n,m} = \tau_{l,h} < +\infty\}$$

has null measure.

Note that equality (28) can be written in terms of enveloping sequences as

$$[M, N]_t = \sum_{n,l} \sum_{m,h} \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \Delta M_{\eta_{n,m}}^{dp} \Delta N_{\tau_{l,h}}^{dp} \mathbb{I}_{\eta_{n,m}=\tau_{l,h}} \mathbb{I}_{\eta_{n,m} \leq t}. \quad (31)$$

Therefore  $[M, N]$  is  $P$ -a.s. null if for any choice of  $n, m$  and  $l, h$  such that  $P(\eta_{n,m} = \tau_{l,h} < +\infty) > 0$  it holds

$$\mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}} \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} \Delta M_{\eta_{n,m}}^{dp} \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \Delta N_{\tau_{l,h}}^{dp} = 0, \quad P\text{-a.s.}$$

Equivalently  $[M, N]$  is  $P$ -a.s. null if

$$\tilde{A}_{n,m}^c \cap \tilde{B}_{l,h}^c \cap \{\eta_{n,m} = \tau_{l,h} < +\infty\}$$

is a set of null measure, where

$$\tilde{A}_{n,m} := \{\Delta M_{\eta_{n,m}}^{dp} \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} = 0\}, \quad \tilde{B}_{l,h} := \{\Delta N_{\tau_{l,h}}^{dp} \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} = 0\}.$$

This easily follows considering that up to a null measure set it holds

$$\tilde{A}_{n,m}^c \cap \tilde{B}_{l,h}^c \subset A_{n,m}^c \cap B_{l,h}^c.$$

In fact by construction  $A_{n,m} \in \mathcal{F}_{\eta_{n,m}-}$  and therefore

$$0 = \int_{A_{n,m}} U_{n,m} dP = \int_{A_{n,m}} (\Delta M_{\eta_{n,m}}^{dp})^2 \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} dP$$

so that  $A_{n,m} \subset \tilde{A}_{n,m}$  up to a null measure set. A similar procedure proves that  $B_{l,h} \subset \tilde{B}_{l,h}$  up to a null measure set and we conclude that  $[M, N] \equiv 0$   $P$ -a.s.

Viceversa, assume  $[M, N] \equiv 0$   $P$ -a.s. This implies that  $P$ -a.s. any addend in the sum of the right-hand side of (31) is null or equivalently  $P$ -a.s. for all  $n, m$  and  $l, h$  for any time  $t$

$$\mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \mathbb{I}_{\eta_{n,m}=\tau_{l,h}} \mathbb{I}_{\eta_{n,m} \leq t} = 0$$

since, when  $\eta_{n,m}$  and  $\tau_{l,h}$  are finite,  $\Delta M_{\eta_{n,m}}^{dp} \Delta N_{\tau_{l,h}}^{dp}$  is  $P$ -a.s. different from zero. More briefly

$$\mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}} = 0 \quad \forall n, m, l, h, \quad P\text{-a.s.} \quad (32)$$

In order to prove the singularity of the kernels we need to show that, if  $P(\eta_{n,m} = \tau_{l,h} < +\infty) > 0$ , then (30) holds. Actually recalling that  $P^* \sim P$  it is enough to show that

$$\mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}} U_{n,m} V_{l,h} = 0, \quad P^*\text{-a.s.}$$

or equivalently

$$E^{P^*}[\mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}}U_{n,m}V_{l,h}] = 0.$$

Since (25) holds then

$$U_{n,m} = E^{P^*} \left[ (\Delta M_{\eta_{n,m}}^{dp})^2 \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} \mid \mathcal{F}_{\eta_{n,m}-} \right], \quad V_{l,h} = E^{P^*} \left[ (\Delta N_{\tau_{l,h}}^{dp})^2 \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \mid \mathcal{H}_{\tau_{l,h}-} \right]$$

and therefore

$$\begin{aligned} E^{P^*}[\mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}}U_{n,m}V_{l,h}] &= \\ E^{P^*} \left[ \mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}} E^{P^*} \left[ (\Delta M_{\eta_{n,m}}^{dp})^2 \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} \mid \mathcal{F}_{\eta_{n,m}-} \right] E^{P^*} \left[ (\Delta N_{\tau_{l,h}}^{dp})^2 \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \mid \mathcal{H}_{\tau_{l,h}-} \right] \right] &= \end{aligned}$$

Now applying Lemma 4.3 in [11] and observing that the random variable  $\mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}}$  is  $\mathcal{F}_{\eta_{n,m}-} \vee \mathcal{H}_{\tau_{l,h}-}$ -measurable, we get

$$\begin{aligned} E^{P^*}[\mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}}U_{n,m}V_{l,h}] &= \\ E^{P^*} \left[ \mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}} E^{P^*} \left[ (\Delta M_{\eta_{n,m}}^{dp})^2 \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} (\Delta N_{\tau_{l,h}}^{dp})^2 \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \mid \mathcal{F}_{\eta_{n,m}-} \vee \mathcal{H}_{\tau_{l,h}-} \right] \right] &= \\ E^{P^*} \left[ E^{P^*} \left[ \mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}} (\Delta M_{\eta_{n,m}}^{dp})^2 \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} (\Delta N_{\tau_{l,h}}^{dp})^2 \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \mid \mathcal{F}_{\eta_{n,m}-} \vee \mathcal{H}_{\tau_{l,h}-} \right] \right] &= \\ E^{P^*} \left[ (\Delta M_{\eta_{n,m}}^{dp})^2 (\Delta N_{\tau_{l,h}}^{dp})^2 \mathbb{I}_{\{\eta_{n,m}=\tau_{l,h}<+\infty\}} \mathbb{I}_{\eta_n^{dp}=\eta_{n,m}} \mathbb{I}_{\tau_l^{dp}=\tau_{l,h}} \right] &= 0, \end{aligned}$$

where last equality follows by (32).  $\square$

Next result states the equivalence between the mutual behavior of the sharp brackets of  $M$  and  $N$  and the existence of a  $\mathbb{G}$ -basis of dimension one. The starting point is Theorem 4.5 in [11], which requires both the  $P$ -triviality of the  $\sigma$ -algebras  $\mathcal{F}_0$  and  $\mathcal{H}_0$  and the  $(P, \mathbb{F})$ -PRP of  $M$  and the  $(P, \mathbb{H})$ -PRP of  $N$ .

From now on, by virtue of Theorem 2.4, we will assume the following condition (A1).

$$(A1) \quad \mathbb{P}(M, \mathbb{F}) = \{P|_{\mathcal{F}_T}\} \text{ and } \mathbb{P}(N, \mathbb{H}) = \{P|_{\mathcal{H}_T}\}.$$

**Proposition 4.2.** *Let  $P^*$  be the probability measure defined by (24).*

- (i) *If the kernels  $d\langle M \rangle^{P, \mathbb{F}}$  and  $d\langle N \rangle^{P, \mathbb{H}}$  are mutually singular, then  $M + N$  enjoys the  $(P^*, \mathbb{G})$ -PRP.*
- (ii) *If there exists  $Z \in \mathcal{M}^2(P^*, \mathbb{G})$  which enjoys the  $(P^*, \mathbb{G})$ -PRP, then the kernels  $d\langle M \rangle^{P, \mathbb{F}}$  and  $d\langle N \rangle^{P, \mathbb{H}}$  are mutually singular.*

*Proof.* (i) Under  $P^*$  the triplet  $(M, N, [M, N])$  is a  $(P^*, \mathbb{G})$ -basis (see Theorem 4.5 in [11]). If  $d\langle M \rangle^{P, \mathbb{F}}$  and  $d\langle N \rangle^{P, \mathbb{H}}$  are mutually singular, then the same holds for  $d\langle M^{dp} \rangle^{P, \mathbb{F}}$  and  $d\langle N^{dp} \rangle^{P, \mathbb{H}}$ . By Lemma 4.1 the covariation process  $[M, N]$  is  $P$ -a.s. null, so that the  $(P^*, \mathbb{G})$ -basis reduces to the pair  $(M, N)$ .

As a consequence, if  $K \in L^2(\Omega, \mathcal{G}_T, P^*)$ , then

$$K = k_0 + \int_0^T \gamma_s dM_s + \int_0^T \eta_s dN_s, \quad (33)$$

where  $k_0$  is a constant,  $(\gamma)_{t \in [0, T]}$  and  $(\eta)_{t \in [0, T]}$  are  $\mathbb{G}$ -predictable processes satisfying  $E^{P^*} \left[ \int_0^T \gamma_t^2 d[M]_t \right] < +\infty$  and  $E^{P^*} \left[ \int_0^T \eta_t^2 d[N]_t \right] < +\infty$ , respectively.

Moreover, since  $P^*|_{\mathcal{F}_T}$  coincides with  $P|_{\mathcal{F}_T}$  then on one side  $\langle M \rangle^{P, \mathbb{F}} = \langle M \rangle^{P^*, \mathbb{F}}$  and, since  $P^*$  decouples  $\mathbb{F}$  and  $\mathbb{H}$ , on the other side  $\langle M \rangle^{P^*, \mathbb{F}} = \langle M \rangle^{P^*, \mathbb{G}}$  and therefore

$$\langle M \rangle^{P, \mathbb{F}} = \langle M \rangle^{P^*, \mathbb{G}}.$$

Analogously  $\langle N \rangle^{P, \mathbb{H}} = \langle N \rangle^{P^*, \mathbb{G}}$ . Then by Proposition 2.24 there exist two  $\mathbb{G}$ -predictable subsets of  $\Omega \times [0, T]$ ,  $C^M$  and  $C^N$ , with  $C^N = \overline{C^M}$ , such that

$$\int_{C^N} d\langle M \rangle_t^{P^*, \mathbb{G}} = 0, \quad \int_{C^M} d\langle N \rangle_t^{P^*, \mathbb{G}} = 0.$$

Let  $(\lambda)_{t \in [0, T]}$  be defined at time  $t$  by

$$\lambda_t = \gamma_t \mathbb{I}_{C^M}(t) + \eta_t \mathbb{I}_{C^N}(t).$$

Since the indicator functions  $(\mathbb{I}_{C^M}(t))_{t \in [0, T]}$  and  $(\mathbb{I}_{C^N}(t))_{t \in [0, T]}$  of the predictable sets  $C^M$  and  $C^N$ , respectively, are predictable processes,  $\lambda$  is a  $\mathbb{G}$ -predictable process.

Moreover  $E^{P^*} \left[ \int_0^T \lambda_t^2 d[M + N]_t \right] < +\infty$  and the following equalities hold

$$E^{P^*} \left[ \int_0^T (\lambda_s - \gamma_s)^2 d\langle M \rangle_s^{P^*, \mathbb{G}} \right] = E^{P^*} \left[ \int_0^T (\lambda_s - \eta_s)^2 d\langle N \rangle_s^{P^*, \mathbb{G}} \right] = 0.$$

As a consequence we get

$$\int_0^T \gamma_s dM_s = \int_0^T \lambda_s dM_s, \quad \int_0^T \eta_s dN_s = \int_0^T \lambda_s dN_s,$$

so that (33) can be rewritten as

$$K = k_0 + \int_0^T \lambda_s d(M_s + N_s)$$

and by the arbitrariness of  $K$  we derive that  $M + N$  enjoys the  $(P^*, \mathbb{G})$ -PRP.

(ii) Let  $Z$  enjoy the  $(P^*, \mathbb{G})$ -PRP. Then

$$M_t = \int_0^t \alpha_s dZ_s, \quad N_t = \int_0^t \beta_s dZ_s,$$

with  $(\alpha)_{t \in [0, T]}$  and  $(\beta)_{t \in [0, T]}$   $\mathbb{G}$ -predictable processes which satisfy  $E^{P^*} \left[ \int_0^T \alpha_t^2 d[Z]_t \right] < +\infty$  and  $E^{P^*} \left[ \int_0^T \beta_t^2 d[Z]_t \right] < +\infty$ , respectively.

By construction  $M$  and  $N$  are independent  $(P^*, \mathbb{G})$ -martingales, so that

$$0 = \langle M, N \rangle^{P^*, \mathbb{G}} = \int_0^T \alpha_s \beta_s d\langle Z \rangle_s^{P^*, \mathbb{G}},$$

that is

$$\alpha.\beta. = 0, \quad d\langle Z \rangle^{P^*, \mathbb{G}}\text{-a.s.} \quad (34)$$

Observe that

$$\langle M \rangle^{P, \mathbb{F}} = \langle M \rangle^{P^*, \mathbb{G}} = \int_0^\cdot \alpha_s^2 d\langle Z \rangle_s^{P^*, \mathbb{G}},$$

and

$$\langle N \rangle^{P, \mathbb{H}} = \langle N \rangle^{P^*, \mathbb{G}} = \int_0^\cdot \beta_s^2 d\langle Z \rangle_s^{P^*, \mathbb{G}}.$$

In order to prove that the kernels  $d\langle M \rangle^{P, \mathbb{F}}$  and  $d\langle N \rangle^{P, \mathbb{H}}$  are mutually singular, let us introduce the predictable set

$$\Gamma := \{(\omega, t), \alpha_t^2(\omega) > 0\}.$$

Then

$$\int_{\Gamma^\omega} \alpha_s^2(\omega) d\langle Z \rangle_s^{P^*, \mathbb{G}}(\omega) = 0,$$

or equivalently

$$d\langle M \rangle^{P, \mathbb{F}}(\omega)(\overline{\Gamma^\omega}) = 0,$$

that is the  $\omega$ -section of  $\Gamma$  is a support of the measure  $d\langle M \rangle^{P, \mathbb{F}}(\omega)$ . Moreover

$$\begin{aligned} & \int_{\Gamma^\omega} \beta_s^2(\omega) d\langle Z \rangle_s^{P^*, \mathbb{G}}(\omega) = \\ & \int_{\{t, \alpha_t(\omega) \neq 0\}} \beta_s^2(\omega) d\langle Z \rangle_s^{P^*, \mathbb{G}}(\omega) = \\ & \int_{[0, T]} \mathbb{1}_{\{t, \alpha_t(\omega) \neq 0\}}(s) \beta_s^2(\omega) d\langle Z \rangle_s^{P^*, \mathbb{G}}(\omega) = \\ & \int_{[0, T]} \mathbb{1}_{\{t, \alpha_t(\omega) \neq 0\}}(s) \mathbb{1}_{\{t, \beta_t(\omega) \neq 0\}}(s) \beta_s^2(\omega) d\langle Z \rangle_s^{P^*, \mathbb{G}}(\omega) = \\ & \int_{[0, T]} \mathbb{1}(s)_{\{t, (\alpha_t \beta_t)(\omega) \neq 0\}} \beta_s^2(\omega) d\langle Z \rangle_s^{P^*, \mathbb{G}}(\omega) = 0, \end{aligned}$$

where the last equality follows by (34). Therefore

$$d\langle N \rangle^{P, \mathbb{H}}(\omega)(\Gamma^\omega) = 0,$$

that is the  $\omega$ -section of  $\overline{\Gamma}$  is a support of the measure  $d\langle N \rangle^{P, \mathbb{H}}(\omega)$  and we get proof.  $\square$

Finally we state the main result of this section. For the sake of clarity we stress that  $\langle M \rangle^{P^*, \mathbb{F}} = \langle M \rangle^{P, \mathbb{F}}$  and  $\langle N \rangle^{P^*, \mathbb{H}} = \langle N \rangle^{P, \mathbb{H}}$ .

**Theorem 4.3.** *The dimension of  $\mathcal{H}^1(P, \mathbb{G})$  is at most three and in particular is*

- (i) *equal to one if and only if the kernels  $d\langle M \rangle^{P, \mathbb{F}}$  and  $d\langle N \rangle^{P, \mathbb{H}}$  are mutually singular;*
- (ii) *equal to two if the kernels  $d\langle M \rangle^{P, \mathbb{F}}$  and  $d\langle N \rangle^{P, \mathbb{H}}$  are not mutually singular but the kernels  $d\langle M^{dp} \rangle^{P, \mathbb{F}}$  and  $d\langle N^{dp} \rangle^{P, \mathbb{H}}$  are mutually singular.*

*Proof.*  $\dim(\mathcal{H}^1(P, \mathbb{G}))$  coincides with  $\dim(\mathcal{H}^1(P^*, \mathbb{G}))$  by Proposition 2.7. By Theorem 4.5 in [11] the set  $\mathbb{P}((M, N, [M, N]), \mathbb{G})$  is a singleton, that is  $(M, N, [M, N])$  enjoys the  $(P^*, \mathbb{G})$ -PRP and therefore, by the remark following Definition 2.6, one gets  $\dim(\mathcal{H}^1(P^*, \mathbb{G}))$  is less or equal to three.

(i) This point immediately follows by Proposition 4.2.

(ii) If  $\dim(\mathcal{H}^1(P^*, \mathbb{G}))$  is greater than one previous point implies that the kernels  $d\langle M \rangle^{P, \mathbb{F}}$  and  $d\langle N \rangle^{P, \mathbb{H}}$  are not mutually singular. Moreover if the kernels  $d\langle M^{dp} \rangle^{P, \mathbb{F}}$  and  $d\langle N^{dp} \rangle^{P, \mathbb{H}}$  are mutually singular, as stated by Lemma 4.1,  $[M, N] = 0$  and  $\mathbb{P}((M, N), \mathbb{G}) = \{P^*|_{\mathbb{G}}\}$ , so that  $\dim(\mathcal{H}^1(P^*, \mathbb{G}))$  is equal to two.  $\square$

**Remark 4.4.** Proposition 4.2 and Theorem 4.3 have been inspired by Theorem 9.5.2.4, page 540, Theorem 9.5.2.5, page 541, and subsequent arguments in [32] and in particular by the following example. A Brownian motion  $W$  and a compensated Poisson martingale  $\Pi$  are considered. If  $f$  and  $g$  are deterministic functions such that  $fg = 0$  then the process  $dX_t = f(t) dW_t + g(t) d\Pi_t$  enjoys the PRP with respect to its natural filtration.

## 5 $(P, \mathbb{G})$ -martingale representation

In this section, in the same setting of last two sections and in particular under the standing assumptions (D) and (A1), we present our martingale representation result on  $\mathbb{G}$  under  $P$ . First we deal with the case when  $\mathbb{G}$  is the progressive enlargement of  $\mathbb{F}$  by a general random time  $\tau$  and then we consider the case when  $\mathbb{G}$  is the enlargement of  $\mathbb{F}$  by the reference filtration  $\mathbb{H}$  of a martingale enjoying the PRP.

In view of possible applications to financial models and although this is not essential for the validity of the result, we assume that  $M$  is the martingale part of a square-integrable special semi-martingale  $X = (X_t)_{t \in [0, T]}$  on  $(\Omega, \mathcal{F}, P)$  with canonical decomposition

$$X_t = X_0 + M_t + A_t, \quad (35)$$

such that

$$E^P [X_0^2 + [M]_T + |A|_T^2] < +\infty.$$

As usual  $A$  is a predictable process of finite variation,  $M_0 = A_0 = 0$  and  $|A|$  denotes the total variation process of  $A$  (see VII-98, page 294, in [21]). In particular we assume that there exists a predictable process  $\alpha = (\alpha_t)_{t \in [0, T]}$   $P$ -a.s. in the space  $L_{loc}^2([0, T], \mathcal{B}([0, T]), d\langle M \rangle^{P, \mathbb{F}})$  such that

$$A_t = \int_0^t \alpha_s d\langle M \rangle_s^{P, \mathbb{F}}. \quad (36)$$

**Remark 5.1.** We note that if  $\mathbb{P}(X, \mathbb{F})$  is a singleton then condition (A1) for  $M$  holds, that is  $\mathbb{P}(M, \mathbb{F}) = \{P|_{\mathcal{F}_T}\}$  (see Proposition 2.2 in [13]). Moreover if  $\mathbb{P}(X, \mathbb{F}) = \{P^X\}$  and the derivative  $dP^X/dP|_{\mathcal{F}_T}$  is locally square-integrable, then (36) holds (see Proposition 4 in [39]).

**Proposition 5.2.**  $X$  and  $M$  share the same jump times.

*Proof.* By Yoeurp's decomposition  $\langle M \rangle^{P, \mathbb{F}}$  coincides with the sum of  $\langle M^c \rangle^{P, \mathbb{F}}$ ,  $\langle M^{dq} \rangle^{P, \mathbb{F}}$  and  $\langle M^{dp} \rangle^{P, \mathbb{F}}$ . The first two processes are continuous and, by Proposition 2.15,  $\langle M^{dp} \rangle^{P, \mathbb{F}}$  is a pure jump process which shares its jump times with  $M^{dp}$ . It follows that the jump times of  $A$  are a subset of those of  $M$  and the thesis follows by equality (35).  $\square$



## 5.1 Progressive enlargement by a random time

Let  $\tau$  be any random time in  $[0, +\infty]$ . For the sake of notational convenience we will rename by  $H$  the compensated occurrence process of  $\tau$ ,  $H^\tau$ , defined as in Subsection 2.2 and by  $\mathbb{H}$  its natural filtration. By Proposition 2.12  $H$  is square-integrable and by Theorem 2.18 it enjoys the  $(P, \mathbb{H})$ -PRP. Here  $H$  plays the role of  $N$  and the filtration  $\mathbb{G}$  defined in (23) is the progressive enlargement of  $\mathbb{F}$  by  $\tau$ . Condition (D) is assumed.

According to (1) we denote by  $\tau^{dp}$  and  $\tau^{dq}$  the  $\mathbb{H}$ -accessible and the  $\mathbb{H}$ -totally inaccessible component of  $\tau$ , respectively. We stress that Proposition 3.1 applies to prove that  $\tau^{dp}$  and  $\tau^{dq}$  are also the  $\mathbb{G}$ -accessible and the  $\mathbb{G}$ -totally inaccessible component of  $\tau$ , respectively. So, we will simply refer to  $\tau^{dp}$  ( $\tau^{dq}$ ) as to the accessible (totally inaccessible) component of  $\tau$  and, as usual,  $(\tau_h^{dp})_{h \in \mathbb{N}}$  will denote an enveloping sequence of  $\tau^{dp}$ .

**Proposition 5.3.** *(i) The triplet  $(M, H, [M, H])$  enjoys the  $(P^*, \mathbb{G})$ -PRP and is a  $(P^*, \mathbb{G})$ -basis of martingales.*

*(ii)  $[M, H]$  admits the representation*

$$[M, H]_t = \sum_{h \in \mathbb{N}} \Delta M_{\tau_h^{dp}} \left( \mathbb{I}_{\{\tau^{dp} = \tau_h^{dp}\}} - P(\tau^{dp} = \tau_h^{dp} \mid \mathcal{H}_{\tau_h^{dp}-}) \right) \mathbb{I}_{\{\tau_h^{dp} \leq t\}}. \quad (37)$$

*Proof.* The  $(P^*, \mathbb{G})$ -martingales  $M$  and  $H$  are independent and therefore strongly orthogonal so that point (i) follows by Theorem 4.5 in [11].

As far as point (ii) is concerned, by Proposition 3.3 it is enough to compute  $[M^{dp}, H^{dp}]$ , where  $H^{dp}$  denotes the accessible part of the Yoeurp decomposition of  $H$  (see (26)). Then representation (37) follows immediately recalling representation (10) and equality (28).  $\square$

Point (i) above joint with Proposition 2.5 assures that a triplet of local martingales driving the  $(P, \mathbb{G})$ -representation exists. Nevertheless, in order to give an explicit description of its components, we need some preliminaries results.

Let  $A^{\tau, P, \mathbb{G}}$  be the  $(P, \mathbb{G})$ -compensator of  $\tau$ . Then  $H' = (H'_t)_{t \in [0, T]}$  defined by

$$H'_t := \mathbb{I}_{\{\tau \leq t\}} - A_t^{\tau, P, \mathbb{G}}$$

is a  $(P, \mathbb{G})$ -martingale. We may refer to  $H'$  as to the  $(P, \mathbb{G})$ -compensated occurrence process of  $\tau$ .

**Proposition 5.4.**

*(i)  $H' \in \mathcal{M}^2(P, \mathbb{G})$  and satisfies*

$$H'_t = \mathbb{I}_{\{\tau \leq t\}} - A_t^{dq, P, \mathbb{G}} - \sum_{h \in \mathbb{N}} P(\tau^{dp} = \tau_h^{dp} \mid \mathcal{G}_{\tau_h^{dp}-}) \mathbb{I}_{\{\tau_h^{dp} \leq t\}},$$

*where  $A^{dq, P, \mathbb{G}}$  denotes the  $(P, \mathbb{G})$ -compensator of  $\tau^{dq}$ .*

*(ii)  $H'$  is a special  $(P^*, \mathbb{G})$ -semi-martingale and  $H$  is its martingale part.*

*Proof.* Let  $A^{dp,P,\mathbb{G}}$  denote the  $(P, \mathbb{G})$ -compensator of  $\tau^{dp}$ . In order to show point (i) observe that Proposition 2.12 applies to prove that  $H'$  is square-integrable and the same procedure used to get (13) yields

$$A^{\tau,P,\mathbb{G}} = A^{dp,P,\mathbb{G}} + A^{dq,P,\mathbb{G}}. \quad (38)$$

The process  $A^{dq,P,\mathbb{G}}$  is a continuous process, while by Lemma 2.13  $A^{dp,P,\mathbb{G}}$  obeys the equality

$$A_t^{dp,P,\mathbb{G}} = \sum_{h \in \mathbb{N}} P(\tau^{dp} = \tau_h^{dp} \mid \mathcal{G}_{\tau_h^{dp}-}) \mathbb{I}_{\{\tau_h^{dp} \leq t\}}. \quad (39)$$

Point (ii) follows immediately by the equality

$$H'_t = H_t + A_t^{\tau,P,\mathbb{H}} - A_t^{\tau,P,\mathbb{G}} \quad (40)$$

considering that by construction  $H$  is a  $(P^*, \mathbb{G})$ -martingale under  $P^*$ , as well as under  $P$ , and the process  $A^{\tau,P,\mathbb{H}} - A^{\tau,P,\mathbb{G}}$  is  $\mathbb{G}$ -predictable and of finite variation.  $\square$

**Remark 5.5.** Since  $\mathbb{I}_{\{\tau_h^{dp} \leq t\}}$  is  $\mathcal{G}_{\tau_h^{dp}-}$ -measurable and  $\{\tau^{dp} < +\infty\} \subset \{\tau = \tau^{dp}\}$ , (39) can be rewritten as

$$A_t^{dp,P,\mathbb{G}} = \sum_{h \in \mathbb{N}} P(\tau = \tau_h^{dp} \mid \mathcal{G}_{\tau_h^{dp}-}) \mathbb{I}_{\{\tau_h^{dp} \leq t\}}.$$

According to Definition 2.11 we introduce a new condition.

(A2)  $P$  is the minimal martingale measure for  $H'$  under  $P^*$ .

Let us analyze condition (A2) and in particular its relationship with the *immersion property*  $\mathbb{F} \xhookrightarrow{P} \mathbb{G}$ , that is the inclusion  $\mathcal{M}_{loc}(P, \mathbb{F}) \subset \mathcal{M}_{loc}(P, \mathbb{G})$  (see Section 5.9.1, page 315, in [32]).

It is to stress that under (A2) the process  $L$  defined by  $L_t := \frac{dP}{dP^*} \Big|_{\mathcal{G}_t}$  satisfies

$$L_t = 1 - \int_0^t \gamma_s L_{s-} dH_s \quad (41)$$

where  $\gamma$  is a suitable  $\mathbb{G}$ -predictable process (see Remark 2 in [39] or Theorem 9 in [5]).

### Proposition 5.6.

- (i) (A2) holds if and only if  $M$  and  $[M, H]$  are  $(P, \mathbb{G})$ -local martingales.
- (ii) (A2) implies  $\mathbb{F} \xhookrightarrow{P} \mathbb{G}$ .
- (iii) When  $\tau$  is totally inaccessible, (A2) is equivalent to  $\mathbb{F} \xhookrightarrow{P} \mathbb{G}$ .

*Proof.* (i) Assuming (A2) by point (i) of Proposition 5.3 it follows immediately that  $M$  and  $[M, H]$  are  $(P, \mathbb{G})$ -local martingales. More precisely, it is easy to check that  $M$  and  $[M, H]$  belong to  $\mathcal{M}^2(P, \mathbb{G})$ . Viceversa, let  $V$  be any element of  $\mathcal{M}_{loc}^2(P^*, \mathbb{G})$  which is orthogonal to the martingale part  $H$  of the special  $(P^*, \mathbb{G})$ -martingale  $H'$ . We have to

show that  $V$  belongs to  $\mathcal{M}(P, \mathbb{G})$ . This is enough since  $P|_{\mathcal{G}_0} = P^*|_{\mathcal{G}_0}$ . By part i) of Proposition 5.3 there exist two predictable processes  $\alpha$  and  $\beta$  such that

$$V_t = \int_0^t \alpha_s dM_s + \int_0^t \beta_s d[M, H]_s.$$

Since equation (41) holds, the processes  $LM$  e  $L[M, H]$  are  $(P^*, \mathbb{G})$ -local martingales and therefore also  $[L, M]$  e  $[L, [M, H]]$  are  $(P^*, \mathbb{G})$ -local martingales (see Lemma 15.2.1, page 373, in [18]). Moreover

$$\begin{aligned} L_t V_t &= \int_0^t L_{s-} dV_s + \int_0^t V_{s-} dL_s + [L, V]_t \\ &= \int_0^t L_{s-} dV_s + \int_0^t V_{s-} dL_s + \int_0^t \alpha_s d[L, M]_s + \int_0^t \beta_s d[L, [M, H]]_s, \end{aligned}$$

so that  $LV$  is a  $(P^*, \mathbb{G})$ -local martingale and therefore  $V$  is a  $(P, \mathbb{G})$ -local martingale.

(ii) This point follows immediately considering that, since  $M$  enjoys the  $(P, \mathbb{F})$ -PRP, any  $(P, \mathbb{F})$ -local martingale can be represented as integral w.r.t.  $M$ , which by assumption is a  $(P, \mathbb{G})$ -martingale.

(iii) By Proposition 3.3  $\tau$  totally inaccessible implies that  $[M, H] \equiv 0$  and  $\mathbb{F} \xrightarrow{P} \mathbb{G}$  implies that  $M$  is a  $(P, \mathbb{G})$ -local martingale, so that by point (i) (A2) holds. The proof ends using point (ii).  $\square$

We state now our main result in the framework of progressive enlargement completed with all the assumptions.

**Theorem 5.7.** *If conditions (D), (A1) and (A2) hold, then the triplet  $(M, H', [M, H])$  enjoys the  $(P, \mathbb{G})$ -PRP.*

*Proof.* By point (i) of Proposition 5.3 and Proposition 2.5 we derive the  $(P, \mathbb{G})$ -PRP for the triplet  $(\tilde{M}, \tilde{H}, \tilde{K})$  defined by

$$\begin{aligned} \tilde{M}_t &:= M_t - \int_0^t \frac{1}{L_{s-}} d\langle L, M \rangle_s^{P^*, \mathbb{G}}, \\ \tilde{H}_t &:= H_t - \int_0^t \frac{1}{L_{s-}} d\langle L, H \rangle_s^{P^*, \mathbb{G}}, \\ \tilde{K}_t &:= [M, H]_t - \int_0^t \frac{1}{L_{s-}} d\langle L, [M, H] \rangle_s^{P^*, \mathbb{G}}, \end{aligned} \tag{42}$$

provided the sharp brackets  $\langle L, M \rangle^{P^*, \mathbb{G}}$ ,  $\langle L, H \rangle^{P^*, \mathbb{G}}$  and  $\langle L, [M, H] \rangle^{P^*, \mathbb{G}}$  exist. Since equation (41) implies that  $LM$  and  $L[M, H]$  are  $(P^*, \mathbb{G})$ -local martingales then

$$\langle L, M \rangle^{P^*, \mathbb{G}} = \langle L, [M, H] \rangle^{P^*, \mathbb{G}} \equiv 0$$

(see Lemma 15.2.1, page 373, in [18]). Moreover, equation (40) implies that  $H$  is a special  $(P, \mathbb{G})$ -semi-martingale and therefore the existence of  $\langle L, H \rangle^{P^*, \mathbb{G}}$  follows by Lemma 2.1 and subsequent Remark 2.1 in [13].

Then by the equations (42) we derive immediately that  $\tilde{M} = M$  and  $\tilde{K} = [M, H]$  and also that  $\tilde{H} = H'$  considering that

$$\tilde{H}_t - H'_t = A_t^{\tau, P, \mathbb{G}} - A_t^{\tau, P, \mathbb{H}} - \int_0^t \frac{1}{L_{s-}} d\langle L, H \rangle_s^{P^*, \mathbb{G}},$$

and therefore  $\tilde{H} - H'$  turns out to be null since it is a predictable  $(P, \mathbb{G})$ -martingale of finite variation (see Lemma 22.11, page 416, in [34]).  $\square$

**Corollary 5.8.** *Assume (D), (A1) and (A2). If  $\tau$  is totally inaccessible, then the pair  $(M, H')$  is a  $(P, \mathbb{G})$ -basis.*

*Proof.* Proposition 3.3 applies to prove that  $[M, H]$  is identically zero. Moreover, under the assumptions of Theorem 5.7, the square-integrable  $(P, \mathbb{G})$ -martingale  $H'$  is strongly orthogonal to  $M$ . In fact it is enough to observe that (40) implies

$$[M, H'] = [M, H] + [M, A^{\tau, P, \mathbb{H}} - A^{\tau, P, \mathbb{G}}] \quad (43)$$

and that by Lemma 2.3 in [40] the process  $[M, A^{\tau, P, \mathbb{H}} - A^{\tau, P, \mathbb{G}}]$  is a  $(P, \mathbb{G})$ -local martingale.  $\square$

When  $[M, H]$  is not identically zero, the triplet  $(M, H', [M, H])$  is not a  $(P, \mathbb{G})$ -basis. In fact the  $(P, \mathbb{G})$ -martingales  $M$  and  $[M, H]$  are not strongly orthogonal and the same happens for  $H'$  and  $[M, H]$ . Let us briefly discuss this point.

$M$  and  $[M, H]$  are strongly orthogonal  $(P, \mathbb{G})$ -martingales if and only if  $M[M, H]$  is a  $(P, \mathbb{G})$ -martingale and the latter holds if and only if  $LM[M, H]$  is a  $(P^*, \mathbb{G})$ -local martingale. Moreover

$$L_t M_t = \int_0^t L_{s-} dM_s + \int_0^t M_{s-} \gamma_s L_{s-} dH_s + \int_0^t \gamma_s L_{s-} d[M, H]_s,$$

so that

$$\begin{aligned} [LM, [M, H]]_t &= \int_0^t L_{s-} d[M, [M, H]]_s + \int_0^t M_{s-} \gamma_s L_{s-} d[H, [M, H]]_s + \\ &\quad + \int_0^t \gamma_s L_{s-} d[[M, H], [M, H]]_s. \end{aligned}$$

The right hand side of previous equality is a  $(P^*, \mathbb{G})$ -local martingale if and only if  $[M, H]$  is identically zero. Similarly we derive that  $H'$  and  $[M, H]$  are not strongly orthogonal.

**Remark 5.9.** *Theorem 5.7 generalizes Proposition 5.3 (ii) in [10]. In that paper, in addition to (A1), the authors assume  $\mathbb{F} \xrightarrow{P} \mathbb{G}$  and Jacod's equivalence hypothesis for  $\tau$  w.r.t. a non-atomic measure  $\nu$  (see Condition (A) and Proposition 1.5 in [28]). Last condition implies the density hypothesis for  $\tau$  (see e.g. [23]) and as a consequence its total inaccessibility (see Remark 2.16). Corollary 2.8 in [4] assures condition (D). Then by point (iii) of Proposition 5.6 it follows that (A2) holds. So, when the processes involved live on  $[0, T]$ , our assumptions are satisfied and, by Corollary 5.8,  $(M, H')$  is a  $(P, \mathbb{G})$ -basis.*

For the sake of completeness we note that in general, under  $\mathbb{F} \xrightarrow{P} \mathbb{G}$ , if

$$\int_0^\cdot \frac{dP[\tau \leq u | \mathcal{F}_u]}{1 - P[\tau < u | \mathcal{F}_u]},$$

is a  $\mathbb{F}$ -predictable process, then its value at  $t \wedge \tau$  coincides  $P$ -a.s. with  $A_t^{\tau, P, \mathbb{G}}$  (see Proposition 6.3 and previous comment in [30]). This happens for example under the density hypothesis for  $\tau$ , so that

$$A_t^{\tau, P, \mathbb{G}} = \int_0^{\tau \wedge t} \lambda_u \nu(du),$$

where the intensity  $\lambda$  is defined by

$$\lambda_t := \frac{p_t(t)}{P(\tau > t \mid \mathcal{F}_t)},$$

with  $(p_t(u))_{u \in [0, T]}$  the  $\mathbb{F}$ -conditional density of  $\tau$ , that is

$$P(\tau > t \mid \mathcal{F}_t) = \int_t^{+\infty} p_t(u) \nu(du)$$

(see formulas (3) and (5) in [10]). So equality (10) reduces to

$$H'_t = \mathbb{I}_{\{\tau \leq t\}} - \int_0^{\tau \wedge t} \lambda_u \nu(du).$$

**Remark 5.10.** It may be worthwhile to observe that, dropping condition (D), which is a standing hypothesis in this paper, and denoting by  $\tau^{dp}$  and  $\tau^{dq}$  the  $\mathbb{G}$ -accessible and the  $\mathbb{G}$ -totally inaccessible component of  $\tau$ , respectively (possibly different from the  $\mathbb{F}$ -accessible and the  $\mathbb{F}$ -totally inaccessible component of  $\tau$ ), the expression of the  $\mathbb{G}$ -compensator of  $\tau$  given by (38) and (39) still holds.

In this more general framework if, for any  $h$ ,  $\tau_h^{dp}$ , is an  $\mathbb{F}$ -stopping time then  $\mathcal{F}_{\tau_h^{dp}-}$  is well-defined and, as it can be easily proved,

$$\mathcal{G}_{\tau_h^{dp}-} \cap \{\tau = \tau_h^{dp}\} = \mathcal{F}_{\tau_h^{dp}-} \cap \{\tau = \tau_h^{dp}\}$$

and

$$P(\tau = \tau_h^{dp} \mid \mathcal{G}_{\tau_h^{dp}-}) \mathbb{I}_{\{\tau_h^{dp} \leq t\}} = P(\tau = \tau_h^{dp} \mid \mathcal{F}_{\tau_h^{dp}-}) \mathbb{I}_{\{\tau_h^{dp} \leq t\}}.$$

As a consequence, if moreover  $\tau^{dq}$  satisfies the Jacod's equivalence hypothesis, then the  $\mathbb{F}$ -conditional law of  $\tau$ , besides an absolutely continuous part which has a density, contains a discontinuous part jumping on  $\mathbb{F}$ -predictable stopping times. In particular, when the enveloping sequence of  $\tau^{dp}$  is finite,  $\tau$  is under the generalized density hypothesis introduced in [33].

## 5.2 Enlargement by a general filtration

We now extend Theorem 5.7 to the more general setting of Section 3 and 4, that is to the case where  $\mathbb{H}$  is the reference filtration of a square integrable martingale  $N$ , which enjoys the  $\mathbb{H}$ -PRP. We recall that conditions (D) and (A1) are in force and the decomposition (26) holds. We indicate with  $\{\tau_l^{dp}\}_{l \in \mathbb{N}}$  the jump times of  $N^{dp}$  and, for each  $l \in \mathbb{N}$ , with  $\{\tau_{l,h}\}_{h \in \mathbb{N}}$  an enveloping sequence of  $\tau_l^{dp}$ . As before  $L = (L_t)_{t \in [0, T]}$  denotes the derivative process  $dP/dP^*|_{\mathcal{G}_t}$ ,  $t \in [0, T]$ .

**Lemma 5.11.** *Let  $L$  be an element of  $\mathcal{M}_{loc}^2(P^*, \mathbb{G})$ . Then*

- (i)  $N^c, N^{dq}$  and  $N^{dp}$  are special  $(P, \mathbb{G})$ -semi-martingales;
- (ii) if  $N^{dp}$  is of integrable variation under  $P$ , then the local martingale part  $N'$  of the special  $(P, \mathbb{G})$ -semi-martingale  $N$  satisfies

$$N' = N - A^{N^c, P, \mathbb{G}} - A^{N^{dq}, P, \mathbb{G}} - \sum_{l, h \in \mathbb{N}} E^P \left[ \Delta N_{\tau_{l, h}} \mathbb{I}_{\tau_l^{dp} = \tau_{l, h}} \mid \mathcal{G}_{\tau_{l, h}^-} \right] \mathbb{I}_{\tau_{l, h} \leq t}, \quad (44)$$

where  $A^{N^c, P, \mathbb{G}}$  and  $A^{N^{dq}, P, \mathbb{G}}$  are the predictable finite variation terms of the special  $(P, \mathbb{G})$ -semi-martingales  $N^c$  and  $N^{dq}$ , respectively.

*Proof.* (i)  $N^c, N^{dq}$  and  $N^{dp}$  are  $(P^*, \mathbb{G})$ -square-integrable martingales so that the hypothesis on  $L$  assures that  $\langle N^c, L \rangle^{P^*, \mathbb{G}}$ ,  $\langle N^{dq}, L \rangle^{P^*, \mathbb{G}}$  and  $\langle N^{dp}, L \rangle^{P^*, \mathbb{G}}$  exist (see VII.39 and subsequent discussion, page 227, in [21]). The statement follows by Remark 2.1 in [13].

(ii)  $N$  is a special  $(P, \mathbb{G})$ -semi-martingale as sum of special  $(P, \mathbb{G})$ -semi-martingales. Moreover, by the uniqueness of the canonical decomposition the  $(P, \mathbb{G})$ -predictable finite variation part of  $N$  coincides with the sum of the predictable finite variation parts of  $N^c, N^{dq}$  and  $N^{dp}$ . In particular, since  $N^{dp}$  is of integrable variation under  $P$ , then last term can be expressed applying Lemma 2.13, so that equality (44) is proved.  $\square$

Hypothesis (A2) in this more general framework reads as follows.

(A2<sub>N</sub>)  $P$  is the minimal martingale measure for  $N'$  under  $P^*$ .

One immediately gets the generalization of Proposition 5.6.

**Proposition 5.12.**

- (i) (A2<sub>N</sub>) holds if and only if  $M$  and  $[M, N]$  are  $(P, \mathbb{G})$ -local martingales.
- (ii) (A2<sub>N</sub>) implies  $\mathbb{F} \xhookrightarrow{P} \mathbb{G}$ .
- (iii) When  $[M, N] \equiv 0$ , (A2<sub>N</sub>) is equivalent to  $\mathbb{F} \xhookrightarrow{P} \mathbb{G}$ .

Let us now present the main theorem.

**Theorem 5.13.** Assume that (D), (A1) and (A2<sub>N</sub>) hold,  $L$  belongs to  $\mathcal{M}_{loc}^2(P^*, \mathbb{G})$  and the accessible martingale part of  $N$ ,  $N^{dp}$ , is of integrable variation. Then the triplet  $(M, N', [M, N])$  enjoys the  $(P, \mathbb{G})$ -PRP, where  $N'$  can be represented as in (44).

*Proof.*  $(M, N, [M, N])$  enjoys the  $(P^*, \mathbb{G})$ -PRP, that is the analogous of point (i) of Proposition 5.3 holds. The assumption (A2<sub>N</sub>) implies that  $\langle L, M \rangle^{P^*, \mathbb{G}} = \langle L, N \rangle^{P^*, \mathbb{G}} \equiv 0$  (see point i) of Proposition 5.12) and  $\langle L, [M, N] \rangle^{P^*, \mathbb{G}}$  exist by the regularity hypothesis on  $L$ . Then Proposition 2.5 can be applied, so that, following the same steps of the proof of Theorem 5.7, we derive the thesis.  $\square$

Under the same assumptions of previous theorem following corollary holds.

**Corollary 5.14.** If  $[M, N] \equiv 0$ , then the pair  $(M, N')$  is a  $(P, \mathbb{G})$ -basis.

*Proof.* By previous theorem  $(M, N')$  enjoys the  $(P, \mathbb{G})$ -PRP. By Proposition 2.12 the local martingale part  $N'$  of the special  $(P, \mathbb{G})$ -semi-martingale  $N$  is square-integrable. Moreover following the same argument of previous section (see (43)) we derive  $M$  and  $N'$  to be orthogonal.  $\square$



An example of martingale representation on progressive enlargement by an accessible random time is studied in [12]. Here we propose a slight generalization in order to get an example of martingales representation on a filtration enlarged through a full process.

On  $(\Omega, \mathcal{F}, P)$  let  $W$  be a standard Brownian motion and  $H^\eta$  the compensated occurrence process of a random time  $\eta$  with values in the set  $\{1, 2, 3\}$ . Assume  $W$  and  $\eta$  independent. Applying point i) of Proposition 4.2 to  $W$  and  $H^\eta$  one gets condition (A1) for  $M$  and  $\mathbb{F}$  defined by

$$M := W + H^\eta, \quad \mathbb{F} := \mathbb{F}^W \vee \sigma(\eta \wedge \cdot),$$

where  $\mathbb{F}^W$  denotes the natural filtration of  $W$ .

On the same probability space let  $H$  be the compensated occurrence process of a random time  $\tau$  taking values in the set  $\{2, 4\}$  and let  $\Pi$  be a compensated Poisson martingale independent of  $(W, \eta, \tau)$  with natural filtration  $\mathbb{F}^\Pi$ . Set

$$N := H + \Pi, \quad \mathbb{H} := \sigma(\tau \wedge \cdot) \vee \mathbb{F}^\Pi.$$

$N$  is a  $(P, \mathbb{H})$ -martingale and by point (i) of Proposition 4.2  $N$  enjoys the  $(P, \mathbb{H})$ -PRP. If  $W$  is independent of  $(\eta, \tau)$  and the joint law  $p_{\eta, \tau}$  of  $(\eta, \tau)$  is strictly positive on the set  $\{1, 2, 3\} \times \{2, 4\}$ , then condition (D) holds and the martingale preserving measure  $P^*$  is defined on  $(\Omega, \mathcal{G}_T)$ , up to a standard extension procedure, by the rule

$$P^*(A \cap C \cap D \cap E) := P(A)P(C)P(D),$$

for any  $A \in \mathcal{F}_T^W, C \in \sigma(\eta \wedge T), D \in \sigma(\tau \wedge T), E \in \mathcal{F}_T^\Pi$ . In this case  $L$  is a bounded process.

$[M, \Pi] \equiv 0$  so that  $[M, N] \equiv [M, H]$ . Therefore condition (A2<sub>N</sub>) is in force as soon as  $[M, H]$  is a  $(P, \mathbb{G})$ -martingale. It can be easily derived that this happens if

$$P(\tau = 2 \mid \eta = 2) = P(\tau = 2 \mid \eta \neq 1) = P(\tau = 2 \mid \eta = 3). \quad (45)$$

So, since  $\Pi$  is a  $(P, \mathbb{G})$ -martingale then  $N'$  coincides with  $H' + \Pi$  and the above theorem states the  $(P, \mathbb{G})$ -driving martingale is the triplet  $(M, H' + \Pi, [M, H])$ . We refer to [12] for the computation of the conditions (45) and for the explicit expression of  $H'$  and  $[M, H]$ .

## 6 Conclusions and perspectives

Our paper has been mainly inspired by the literature of credit risk theory, where the problem of martingale representation on a market with full information has been largely studied. Actually in real markets the default time may be predictable, may overlap the jump of the asset price process or even the insider may be aware of more than a single default. That is why, working in a purely theoretical framework, we have assumed that the filtration enlargement is induced by a not necessarily quasi-left continuous random process, possibly with jump times common to the processes adapted to the basic filtration and more general than the occurrence process of a random time. As a byproduct we have investigated the interplay between two notions specific of mathematical finance: the

*minimal martingale measure and the immersion property.*

A natural development of the current result, which is object of ongoing research, is to get rid of the decoupling assumption (D) by moving on the study of martingale representation on enlarged filtrations in the light of the recent paper by Aksamit, Chouilly and Jeanblanc on *thin-thick decomposition of a random time* (see [1]).

## Funding

Partially supported by the MIUR Excellence Department Project MatMod@TOV awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C23000330006.

## References

- [1] AKSAMIT, A., CHOULLI, T., AND JEANBLANC, M. Thin times and random times' decomposition. *Electron. J. Probab.* 26 (2021), Paper No. 31, 22.
- [2] AKSAMIT, A., AND FONTANA, C. Martingale spaces and representations under absolutely continuous changes of probability. *Electron. Commun. Probab.* 24 (2019), Paper No. 62, 13.
- [3] AKSAMIT, A., JEANBLANC, M., AND RUTKOWSKI, M. Integral representations of martingales for progressive enlargements of filtrations. *Stochastic Process. Appl.* 129, 4 (2019), 1229–1258.
- [4] AMENDINGER, J., BECHERER, D., AND SCHWEIZER, M. A monetary value for initial information in portfolio optimization. *Finance Stoch.* 7, 1 (2003), 29–46.
- [5] ANSEL, J.-P., AND STRICKER, C. Lois de martingale, densités et décomposition de Föllmer-Schweizer. *Ann. Inst. H. Poincaré Probab. Statist.* 28, 3 (1992), 375–392.
- [6] BANDINI, E., CALVIA, A., AND COLANERI, K. Stochastic filtering of a pure jump process with predictable jumps and path-dependent local characteristics. *Stochastic Process. Appl.* 151 (2022), 396–435.
- [7] BANDINI, E., CONFORTOLA, F., AND DI TELLA, P. Progressively enlargement of filtrations and control problems for step processes. Tech. rep., arXiv:2112.12884v1, 2021.
- [8] BIAGINI, F., MAZZON, A., AND OBERPRILLER, K. Reduced-form framework for multiple ordered default times under model uncertainty. *Stochastic Process. Appl.* 156 (2023).
- [9] BILLINGSLEY, P. *Probability and measure*, third ed. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1995. A Wiley-Interscience Publication.
- [10] CALLEGARO, G., JEANBLANC, M., AND ZARGARI, B. Carthaginian enlargement of filtrations. *ESAIM Probab. Stat.* 17 (2013), 550–566.

- 
- [11] CALZOLARI, A., AND TORTI, B. Enlargement of filtration and predictable representation property for semi-martingales. *Stochastics* 88, 5 (2016), 680–698.
  - [12] CALZOLARI, A., AND TORTI, B. An example of martingale representation in progressive enlargement by an accessible random time. In *Frontiers in Stochastic Analysis-BSDEs, SPDEs and their Applications, Springer Proceedings in Mathematics and Statistics*, vol. 289. Springer, 2019, pp. 109—121.
  - [13] CALZOLARI, A., AND TORTI, B. Martingale representations in progressive enlargement by the reference filtration of a semi-martingale: a note on the multidimensional case. *Stochastics* 91, 2 (2019), 265–287.
  - [14] CALZOLARI, A., AND TORTI, B. Martingale representations in progressive enlargement by multivariate point processes. *Intern. Journal of Theor. and Appl. Finance* 25, 03 (2022), 1–21.
  - [15] CHATELAIN, M., AND STRICKER, C. On componentwise and vector stochastic integration. *Math. Finance* 4, 1 (1994), 57–65.
  - [16] CHOU, C. S., AND MEYER, P. A. Sur la représentation des martingales comme intégrales stochastiques dans les processus ponctuels. In *Séminaire de Probabilités, IX (Seconde Partie, Univ. Strasbourg, Strasbourg, années universitaires 1973/1974 et 1974/1975)*. Springer, Berlin, 1975, pp. 226–236. Lecture Notes in Math., Vol. 465.
  - [17] COCULESCU, D., JEANBLANC, M., AND NIKEGHBALI, A. Default times, no-arbitrage conditions and changes of probability measures. *Finance Stoch.* 16, 3 (2012), 513–535.
  - [18] COHEN, S. N., AND ELLIOTT, R. J. *Stochastic calculus and applications*, second ed. Probability and its Applications. Springer, Cham, 2015.
  - [19] DAVIS, M. H. A., AND VARAIYA, P. The multiplicity of an increasing family of  $\sigma$ -fields. *Ann. Probability* 2 (1974), 958–963.
  - [20] DELLACHERIE, C., AND MEYER, P.-A. Probabilities and potential. A. vol. 29 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1978, pp. xvii+463.
  - [21] DELLACHERIE, C., AND MEYER, P.-A. Probabilities and potential. B. vol. 72 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1982, pp. xvii+463. Theory of martingales, Translated from the French by J. P. Wilson.
  - [22] DI TELLA, P., AND JEANBLANC, M. Martingale representation in the enlargement of the filtration generated by a point process. *Stochastic Process. Appl.* 131 (2021), 103–121.
  - [23] EL KAROUI, N., JEANBLANC, M., AND JIAO, Y. What happens after a default: the conditional density approach. *Stochastic Process. Appl.* 120, 7 (2010), 1011–1032.
  - [24] EL KAROUI, N., JEANBLANC, M., AND JIAO, Y. Density approach in modeling successive defaults. *SIAM J. Financial Math.* 6, 1 (2015), 1–21.

- 
- [25] FONTANA, C., AND SCHMIDT, T. General dynamic term structures under default risk. *Stochastic Process. Appl.* 128, 10 (2018), 3353–3386.
  - [26] HE, S. W., WANG, J. G., AND YAN, J. A. Semimartingale theory and stochastic calculus. Kexue Chubanshe (Science Press), Beijing; CRC Press, Boca Raton, FL, 1992, pp. xiv+546.
  - [27] JACOD, J. Calcul stochastique et problèmes de martingales. vol. 714 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979, pp. x+539.
  - [28] JACOD, J. Grossissement initial, hypothèse (H'), et théorème de Girsanov. In *Thierry Jeulin and Marc Yor editors, Grossissements de Filtrations: Exemples et Applications*, vol. 1118 of *Lecture Notes in Math.* Springer, Berlin, 1985, pp. 15–35.
  - [29] JACOD, J., AND SHIRYAEV, A. N. Limit theorems for stochastic processes. Second ed., vol. 288 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2003, pp. xx+661.
  - [30] JEANBLANC, M., AND RUTKOWSKI, M. Models for default risk: an overview. In *Mathematical finance: theory and practise.*, J. Yong and R. Cont, Eds. Higher education press, 1999.
  - [31] JEANBLANC, M., AND SONG, S. Martingale representation property in progressively enlarged filtrations. *Stochastic Process. Appl.* 125, 11 (2015), 4242–4271.
  - [32] JEANBLANC, M., YOR, M., AND CHESNEY, M. Mathematical methods for financial markets. Springer Finance. Springer-Verlag London Ltd., London, 2009, pp. xxvi+732.
  - [33] JIAO, Y., AND LI, S. The generalized density approach in progressive enlargement of filtrations. *Electron. J. Probab.* 20 (2015), no. 85, 21.
  - [34] KALLENBERG, O. Foundations of modern probability. Probability and its Applications (New York). Springer-Verlag, New York, 1997, pp. xii+523.
  - [35] KALLENBERG, O. *Random measures, theory and applications*, vol. 77 of *Probability Theory and Stochastic Modelling*. Springer, Cham, 2017.
  - [36] KCHIA, Y., AND PROTTER, P. Progressive filtration expansions via a process, with applications to insider trading. *Int. J. Theor. Appl. Finance* 18, 4 (2015), 1550027, 48.
  - [37] MANSUY, R., AND YOR, M. Random times and enlargements of filtrations in a Brownian setting. vol. 1873 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2006, pp. xiv+158.
  - [38] PROTTER, P. E. Stochastic integration and differential equations. Second ed., vol. 21 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 2004, pp. xiv+415. Stochastic Modelling and Applied Probability.
  - [39] SCHWEIZER, M. Martingale densities for general asset prices. *J. Math. Econom.* 21, 4 (1992), 363–378.

- [40] YOEURP, C. Décompositions des martingales locales et formules exponentielles. In *Séminaire de Probabilités, X (Seconde partie: Théorie des intégrales stochastiques, Univ. Strasbourg, Strasbourg, année universitaire 1974/1975)*. Springer, Berlin, 1976, pp. 432–480. Lecture Notes in Math., Vol. 511.