DILATION, FUNCTIONAL MODEL AND A COMPLETE UNITARY INVARIANT FOR $C_{.0}$ Γ_n -CONTRACTIONS

SOURAV PAL

ABSTRACT. A commuting tuple of operators $(S_1, \ldots, S_{n-1}, P)$, defined on a Hilbert space \mathcal{H} , for which the closed symmetrized polydisc

$$\Gamma_n = \left\{ \left(\sum_{1 \le i \le n} z_i, \sum_{1 \le i < j \le n} z_i z_j, \dots, \prod_{i=1}^n z_i \right) \colon |z_i| \le 1, i = 1, \dots, n \right\}$$

is a spectral set, is called a Γ_n -contraction. A Γ_n -contraction is said to be *pure* or C_0 if P is C_0 , that is, if $P^{*n} \to 0$ strongly as $n \to \infty$. We show that for any Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$, there is a unique operator tuple (A_1, \ldots, A_{n-1}) that satisfies the operator identities

 $S_i - S_{n-i}^* P = D_P A_i D_P, \qquad i = 1, \dots, n-1.$

This unique tuple is called the *fundamental operator tuple* or \mathscr{F}_O -tuple of $(S_1, \ldots, S_{n-1}, P)$. With the help of the \mathscr{F}_O -tuple, we construct an operator model for a $C_{\cdot 0}$ Γ_n -contraction and show that there exist n-1 operators C_1, \ldots, C_{n-1} such that each S_i can be represented as $S_i = C_i + PC_{n-i}^*$. We find an explicit minimal dilation for a class of $C_{\cdot 0}$ Γ_n -contractions whose \mathscr{F}_O -tuples satisfy a certain condition. Also we establish that the \mathscr{F}_O -tuple of $(S_1^*, \ldots, S_{n-1}^*, P^*)$ together with the characteristic function of P constitute a complete unitary invariant for the $C_{\cdot 0}$ Γ_n -contractions. The entire program is an analogue of the Nagy-Foias theory for $C_{\cdot 0}$ contractions.

1. INTRODUCTION

Throughout the paper all operators are bounded linear operators defined on complex Hilbert spaces. A contraction is an operator whose norm is not greater than 1. We denote by \mathbb{C}, \mathbb{N} the set of complex numbers and positive integers respectively. The open unit disc and the unit circle with center at the origin in \mathbb{C} are denoted by \mathbb{D} and \mathbb{T} respectively.

For studying an operator, it suffices to consider only contractions because an operator is just a scalar multiple of a contraction. A few decades ago Sz.-Nagy and Foias initiated a program of determining the structure of a contraction and modeling it as a compression of a unitary operator. As a consequence numerous novel results were achieved and their beautiful constructive proofs were witnessed. A keen reader is referred to the classic [7] and references there in. In 1951, von Neumann, [26], introduced the notion of spectral set for an operator and described the contractions as operators having $\overline{\mathbb{D}}$ as a spectral set.

Theorem 1.1 (von Neumann, 1951). An operator T is a contraction if and only if \mathbb{D} is a spectral set for T.

The notion of spectral set, which was later defined by Arveson (e.g., [6]) for any finite tuple of commuting operators, became more popular and effective in deciphering the interplay between

²⁰¹⁰ Mathematics Subject Classification. 47A13, 47A20, 47A25, 47A45.

Key words and phrases. Γ_n -contraction, Fundamental operator tuple, Operator model, Complete unitary invariant.

The author was supported by the Seed Grant of IIT Bombay, the CPDA of Govt. of India, the INSPIRE Faculty Award (Award No. DST/INSPIRE/04/2014/001462) of DST, India and the MATRICS Award of SERB, (Award No. MTR/2019/001010) of DST, India.

the intrinsic properties of an operator tuple and the complex geometry of an underlying compact subset of \mathbb{C}^n associated with the tuple (e.g., see [1, 22]). In 1953, Sz.-Nagy published a very influential article, [24], where he established the following dilation theorem whose impact is extraordinary till date.

Theorem 1.2 (Sz.-Nagy, 1953). If T is a contraction acting on a Hilbert space \mathcal{H} , then there exists a Hilbert space $\mathcal{K} \supseteq \mathcal{H}$ and a unitary U on \mathcal{K} such that

$$p(T) = P_{\mathscr{H}}p(U)|_{\mathscr{H}}$$
 for every polynomial $p \in \mathbb{C}[z]$.

A major and important class of operators that has been extensively studied by Sz.-Nagy, Foias and many other mathematicians (e.g., [7]), is the $C_{.0}$ class of contractions. A contraction T, defined on a Hilbert space \mathscr{H} , is said to be $C_{.0}$ or *pure*, if $T^{*n}h \to 0$ as $n \to \infty$ for all $h \in \mathscr{H}$. The aim of this article is to study the $C_{.0}$ operator tuples associated with the symmetrized polydisc \mathbb{G}_n , where

$$\mathbb{G}_n = \left\{ \left(\sum_{1 \le i \le n} z_i, \sum_{1 \le i < j \le n} z_i z_j, \dots, \prod_{i=1}^n z_i \right) : |z_i| < 1, i = 1, \dots, n \right\}.$$

The symmetrized polydisc \mathbb{G}_n and its closure Γ_n , given by

$$\Gamma_n = \left\{ \left(\sum_{1 \le i \le n} z_i, \sum_{1 \le i < j \le n} z_i z_j, \dots, \prod_{i=1}^n z_i \right) : |z_i| \le 1, i = 1, \dots, n \right\},\$$

are the images of polydisc \mathbb{D}^n and its closure $\overline{\mathbb{D}^n}$ respectively under the symmetrization map $\pi_n : \mathbb{C}^n \to \mathbb{C}^n$ defined by

$$\pi_n(z_1,\ldots,z_n) = \left(\sum_{1\leq i\leq n} z_i, \sum_{1\leq i< j\leq n} z_i z_j,\ldots,\prod_{i=1}^n z_i\right).$$

The family of domains { $\mathbb{G}_n : n \in \mathbb{N}$ } was introduced in [11] to study the spectral Nevanlinna-Pick problem. Indeed, the family of domains { $\mathbb{G}_n : n \in \mathbb{N}$ } is naturally associated with spectral interpolation in the following way: if $\mathcal{M}_n(\mathbb{C})$ is the space of all $n \times n$ complex matrices and if \mathcal{B}_1 is its spectral unit ball, then $A \in \mathcal{B}_1$ (that is, the spectral radius r(A) < 1) if and only if $\Pi_n(A) \in \mathbb{G}_n$, where $\Pi_n(A) = \pi_n(\sigma(A)), \sigma(A)$ being the spectrum of A. The domain \mathbb{G}_n is of importance because, apart from the derogatory matrices, the $n \times n$ spectral Nevanlinna-Pick problem is equivalent to a similar interpolation problem of \mathbb{G}_n (see [3], Theorem 2.1). Note that a bounded domain like \mathbb{G}_n , which has complex-dimension n, is much easier to deal with than an unbounded n^2 -dimensional object like \mathcal{B}_1 . The symmetrized polydisc has attracted considerable attention in past two decades because of its rich function theory, beautiful complex geometry and appealing operator theory (see [11, 14, 2, 4, 8, 9, 10] and references there in).

In this paper, we analyze and develop a Nagy-Foias type operator theory for the commuting operator tuples having Γ_n as a spectral set.

Definition 1.3. A commuting tuple of operators $(S_1, \ldots, S_{n-1}, P)$ that has Γ_n as a spectral set is called a Γ_n -contraction, that is, $(S_1, \ldots, S_{n-1}, P)$ is a Γ_n -contraction if the Taylor joint spectrum $\sigma_T(S_1, \ldots, S_{n-1}, P) \subseteq \Gamma_n$ and von Neumann's inequality

$$||f(S_1,...,S_{n-1},P)|| \le \sup_{\mathbf{z}\in\Gamma_n} |f(\mathbf{z})| = ||f||_{\infty,\Gamma_n} \qquad [\mathbf{z}=(s_1,...,s_{n-1},p)]$$

holds for all rational functions f = p/q, $(p, q \in \mathbb{C}[z_1, ..., z_n])$ such that f does not have any pole in Γ_n . Also, a Γ_n -contraction $(S_1, ..., S_{n-1}, P)$ is called *pure* or $C_{\cdot 0}$ if P is a $C_{\cdot 0}$ contraction.

$C_{.0}\Gamma_n$ -CONTRACTIONS

Here $f(S_1, \ldots, S_{n-1}, P) = p(S_1, \ldots, S_{n-1}, P) q(S_1, \ldots, S_{n-1}, P)^{-1}$ in order to maintain the standard convention. In Section 2, we shall explain the motivation behind defining $C_{\cdot 0} \Gamma_n$ -contraction in terms of the last component P. A Nagy-Foias type operator theoretic program for the pure Γ_2 contractions was initiated in [8] and was further carried out in [9]. In this paper, we generalize those results for any arbitrary $n \ge 2$ and our methods are also a generalization of the techniques that were used in [8, 9]. Operator theory on Γ_2 was simpler because rational dilation succeeded on Γ_2 (see [8]) and a concrete operator model was obtained as a consequence of dilation, [9]. Since rational dilation fails on \mathbb{G}_n for $n \ge 3$ (see [19]), only conditional dilation and functional model can be achieved when $n \ge 3$.

In [7], Nagy and Foias showed that a $C_{.0}$ contraction T, defined on a Hilbert space \mathscr{H} , can be realized as the compression of the shift operator M_z defined on the vectorial Hardy space $H^2(\mathscr{D}_{T^*})$, where $\mathscr{D}_{T^*} = \overline{\operatorname{Ran}} D_P = \overline{\operatorname{Ran}} (I - TT^*)^{\frac{1}{2}}$. To obtain this representation, they first showed that M_z on $H^2(\mathscr{D}_{P^*})$ is the minimal isometric dilation of T. In Theorem 4.3, we construct an analogous minimal Γ_n -isometric dilation of a $C_{.0} \Gamma_n$ -contraction $(S_1, \ldots, S_{n-1}, P)$ under certain conditions. The minimal dilation space for $(S_1, \ldots, S_{n-1}, P)$ is no bigger than the Nagy-Foias minimal dilation space for P, which is $H^2(\mathscr{D}_{P^*})$. As a consequence of this Γ_n -isometric dilation, we achieve in Theorem 4.9 a concrete functional model for $(S_1, \ldots, S_{n-1}, P)$. Also in Theorem 4.6, we independently produce an operator model for a $C_{.0} \Gamma_n$ -contraction without assuming any condition on $(S_1, \ldots, S_{n-1}, P)$. This model may or may not be a commutative one. A unique operator tuple (B_1, \ldots, B_{n-1}) associated with $(S_1^*, \ldots, S_{n-1}^*, P^*)$, which satisfies

$$S_i^* - S_{n-i}P^* = D_{P^*}B_iD_{P^*}$$
 $i = 1, \dots, n-1,$

plays the central role in the constructions of the dilation and the models. The existence and uniqueness of such an (n-1)-tuple are proved in Theorem 3.3. Indeed, in Theorem 3.3 we consider the general case, that is, we prove that for every Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$ there exists a unique tuple (A_1, \ldots, A_{n-1}) such that

$$S_i - S_{n-i}^* P = D_P A_i D_P$$
 $i = 1, ..., n - 1$

For its pivotal role in operator theory on Γ_n , (A_1, \ldots, A_{n-1}) is called the *fundamental operator tuple* or shortly the \mathscr{F}_O -tuple of $(S_1, \ldots, S_{n-1}, P)$. In [11], Costara showed that for any $(s_1, \ldots, s_{n-1}, p) \in \Gamma_n$ there is a unique $(\beta_1, \ldots, \beta_{n-1}) \in \Gamma_{n-1}$ such that $s_i = \beta_i + p\overline{\beta}_{n-i}$ for $i = 1, \ldots, n-1$. With the help of the \mathscr{F}_O -tuple, we find an operator theoretic analogue of this result for a C_0 Γ_n -contraction in Corollaries 4.7, 4.8. In view of the existence and uniqueness of the \mathscr{F}_O -tuple of a Γ_n -contraction (as in Theorem 3.3), a natural question arises: given n-1 operators A_1, \ldots, A_{n-1} , can we find a Γ_n -contraction for which (A_1, \ldots, A_{n-1}) is the \mathscr{F}_O -tuple? We provide a partial answer to this question in Theorem 3.7.

One of the most wonderful discoveries in operator theory is the characteristic function of a contraction due to Nagy and Foias (defined in Subsection 4.2) which is a complete unitary invariant for the completely non-unitary (c.n.u) contractions in the sense that two c.n.u contractions T_1, T_2 are unitarily equivalent if and only if their characteristic functions coincide. In Theorem 5.2, we find a complete unitary invariant for the $C_{.0}$ Γ_n -contractions. We show that for a $C_{.0}$ Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$, the \mathscr{F}_O -tuple (B_1, \ldots, B_{n-1}) of $(S_1^*, \ldots, S_{n-1}^*, P^*)$ and the characteristic function Θ_P of P constitute a complete unitary invariant.

In Section 2, we accumulate a few definitions and results from the literature which will be used in sequel.

Note. The present article is an updated version of a part of the author's unpublished paper [20]. We learned that Theorem 3.3 and the preparatory result Proposition 4.2 of this article were independently proved by A. Pal in [16].

Acknowledgement. The author is thankful to Orr Shalit for several fruitful comments on this article.

2. A BRIEF LITERATURE AND PRELIMINARIES

Unitary, isometry and co-isometry are special classes of contractions. There are natural analogues of these classes for Γ_n -contractions in the literature (see [4, 10, 19]). It was established in [14] that the *distinguished boundary* of Γ_n , denoted by $b\Gamma_n$, is the symmetrization of the distinguished boundary of the polydisc, which is the *n*-torus \mathbb{T}^n , and thus $b\Gamma_n$ is the set

$$b\Gamma_n = \left\{ \left(\sum_{i=1}^n z_i, \sum_{1 \le i < j \le n} z_i z_j, \dots, \prod_{i=1}^n z_i \right) : |z_i| = 1, i = 1, \dots, n \right\}$$

Definition 2.1. Let S_1, \ldots, S_{n-1}, P be commuting operators on \mathcal{H} . Then $(S_1, \ldots, S_{n-1}, P)$ is called

- (i) a Γ_n -unitary if S_1, \ldots, S_{n-1}, P are normal operators and $\sigma_T(S_1, \ldots, S_{n-1}, P) \subseteq b\Gamma_n$;
- (ii) a Γ_n -isometry if there exist a Hilbert space $\mathscr{K} \supseteq \mathscr{H}$ and a Γ_n -unitary $(T_1, \ldots, T_{n-1}, U)$ on \mathscr{K} such that \mathscr{H} is a joint invariant subspace of T_1, \ldots, T_{n-1}, U and

$$(T_1|_{\mathscr{H}},\ldots,T_{n-1}|_{\mathscr{H}},U|_{\mathscr{H}})=(S_1,\ldots,S_{n-1},P);$$

(iii) a Γ_n -co-isometry if the adjoint $(S_1^*, \ldots, S_{n-1}^*, P^*)$ is a Γ_n -isometry.

The following theorems from [19] provide clear descriptions of a Γ_n -unitary and a Γ_n -isometry.

Theorem 2.2 ([19], Theorems 4.2 & 4.4). A commuting tuple of operators $(S_1, \ldots, S_{n-1}, P)$ is a Γ_n -unitary (or, a Γ_n -isometry) if and only if $(S_1, \ldots, S_{n-1}, P)$ is a Γ_n -contraction and P is a unitary (isometry).

Needless to mention that $(S_1, \ldots, S_{n-1}, P)$ is a Γ_n -co-isometry if and only if $(S_1, \ldots, S_{n-1}, P)$ is a Γ_n -contraction and P is a co-isometry. So, it is evident that the nature of a Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$ is highly influenced by the nature of its last component P. In [18], the author of this paper showed that for a given Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$ on \mathcal{H} , if $P = P|_{\mathcal{H}_1} \oplus P|_{\mathcal{H}_2}$ is the canonical decomposition of the contraction P with respect to $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$ such that $P|_{\mathcal{H}_1}$ is a unitary and $P|_{\mathcal{H}_2}$ is a c.n.u contraction, then both $\mathcal{H}_1, \mathcal{H}_2$ reduce S_1, \ldots, S_{n-1} and $(S_1|_{\mathcal{H}_1}, \ldots, S_{n-1}|_{\mathcal{H}_1}, P|_{\mathcal{H}_1})$ is a Γ_n -unitary whereas $(S_1|_{\mathcal{H}_2}, \ldots, S_{n-1}|_{\mathcal{H}_2}, P|_{\mathcal{H}_2})$ is a Γ_n -contraction for which $P|_{\mathcal{H}_2}$ is a c.n.u contraction. This unique decomposition was named the "canonical decomposition" of a Γ_n -contraction (S_1, \ldots, S_{n-1}, P) for which P is a c.n.u contraction and indeed such a definition is justified. Taking cue from such dominant roles of P in determining the special classes of a Γ_n -contraction (S_1, \ldots, S_{n-1}, P) we are led to the following definition.

Definition 2.3. A Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$ acting on \mathscr{H} is said to be $C_{\cdot 0}$ or *pure* if P is a $C_{\cdot 0}$ contraction, that is, if $P^{*n}h \to 0$ as $n \to \infty$ for all $h \in \mathscr{H}$.

The following theorem, which provides a characterization for the Γ_n -unitaries, will be used in sequel.

Theorem 2.4 ([10], Theorem 4.2). Let $(S_1, \ldots, S_{n-1}, P)$ be a commuting tuple of bounded operators. Then the following are equivalent.

- (1) $(S_1, \ldots, S_{n-1}, P)$ is a Γ_n -unitary,
- (2) *P* is a unitary, $(\frac{n-1}{n}S_1, \frac{n-2}{n}S_2, \dots, \frac{1}{n}S_{n-1})$ is a Γ_{n-1} -contraction and $S_i = S_{n-i}^* P$ for $i = 1, \dots, n-1$.
 - 3. The fundamental operator tuple (\mathscr{F}_O -tuple) of a Γ_n -contraction

In [18], we introduced the following n-1 operator pencils $\Phi_1, \ldots, \Phi_{n-1}$ in order to determine the structure of a Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$:

$$\Phi_{i}(S_{1},...,S_{n-1},P) = (\tilde{n}_{i}-S_{i})^{*}(\tilde{n}_{i}-S_{i}) - (\tilde{n}_{i}P-S_{n-i})^{*}(\tilde{n}_{i}P-S_{n-i})$$

$$= \tilde{n}_{i}^{2}(I-P^{*}P) + (S_{i}^{*}S_{i}-S_{n-i}^{*}S_{n-i}) - \tilde{n}_{i}(S_{i}-S_{n-i}^{*}P)$$

$$- \tilde{n}_{i}(S_{i}^{*}-P^{*}S_{n-i}), \quad \text{where } \tilde{n}_{i} = \binom{n}{i}.$$
(3.1)

We mention here to the readers that while defining Φ_i in [18], \tilde{n}_i was mistakenly displayed as *n* and that was a typographical error. From the definition it is clear that in particular when S_1, \ldots, S_{n-1}, P are scalars, i.e, points in Γ_n , the above operator pencils take the following form for each *i*:

$$\Phi_i(s_1, \dots, s_{n-1}, p) = \tilde{n}_i^2 (1 - |p|^2) + (|s_i|^2 - |s_{n-i}|^2) - \tilde{n}_i(s_i - \bar{s}_{n-i}p) - \tilde{n}_i(\bar{s}_i - \bar{p}s_{n-i}).$$
(3.2)

The following result appeared in [18] and is extremely important in the context of this paper.

Proposition 3.1 (Proposition 2.6, [18]). Let $(S_1, \ldots, S_{n-1}, P)$ be a Γ_n -contraction. Then for $i = 1, \ldots, n-1, \ \Phi_i(\alpha S_1, \ldots, \alpha^{n-1} S_{n-1}, \alpha^n P) \ge 0$ for all $\alpha \in \overline{\mathbb{D}}$.

The positivity of the operator pencils Φ_i will determine a certain and unique operator tuple (A_1, \ldots, A_{n-1}) associated with each Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$. We shall call (A_1, \ldots, A_{n-1}) the fundamental operator tuple or the \mathscr{F}_O -tuple of $(S_1, \ldots, S_{n-1}, P)$ and the underlying reason is that it plays the central role in constructing the explicit dilation and operator models and in determining the complete unitary invariant for a C_0 Γ_n -contraction.

Recall that the *numerical radius* of an operator A on a Hilbert space \mathcal{H} is defined by

$$\boldsymbol{\omega}(A) = \sup\{|\langle Ax, x\rangle| : \|x\|_{\mathscr{H}} = 1\}.$$

It is well known that

$$r(A) \le \omega(A) \le ||A|| \text{ and } \frac{1}{2} ||A|| \le \omega(A) \le ||A||,$$
 (3.3)

where r(A) is the spectral radius of A. We state a basic lemma on the numerical radius whose proof is a routine exercise. We shall use this lemma in sequel.

Lemma 3.2. The numerical radius of an operator A is not greater than 1 if and only if Re $(\alpha A) \leq I$ for all complex numbers α of modulus 1.

Theorem 3.3. (Existence and Uniqueness). Let $(S_1, \ldots, S_{n-1}, P)$ be a Γ_n -contraction on a Hilbert space \mathcal{H} . Then there are unique operators $A_1, \ldots, A_{n-1} \in \mathcal{B}(\mathcal{D}_P)$ such that

$$S_i - S_{n-i}^* P = D_P A_i D_P$$
 for $i = 1, ..., n-1$.

Moreover, for each i and for all $z \in \mathbb{T}$, $\omega(A_i + A_{n-i}z) \leq \tilde{n}_i$. $[\tilde{n}_i = \binom{n}{i}]$.

Proof. We apply Proposition 3.1 to $(S_1, \ldots, S_{n-1}, P)$ and obtain for each $i = 1, \ldots, n-1$,

$$\Phi_i(\alpha S_1,\ldots,\alpha^{n-1}S_{n-1},\alpha^n P) \ge 0, \qquad (3.4)$$

for all $\alpha \in \overline{\mathbb{D}}$. Therefore, in particular for $\beta, \gamma \in \mathbb{T}$ we have from (3.4) for Φ_i and Φ_{n-i} respectively

$$\tilde{n}_{i}^{2}(I-P^{*}P) + (S_{i}^{*}S_{i} - S_{n-i}^{*}S_{n-i}) \geq \tilde{n}_{i}\beta^{i}(S_{i} - S_{n-i}^{*}P) + \tilde{n}_{i}\bar{\beta}^{i}(S_{i}^{*} - P^{*}S_{n-i}), \qquad (3.5)$$

$$\tilde{n}_{i}^{2}(I-P^{*}P) + (S_{n-i}^{*}S_{n-i} - S_{i}^{*}S_{i}) \geq \tilde{n}_{i}\gamma^{n-i}(S_{n-i} - S_{i}^{*}P) + \tilde{n}_{i}\bar{\gamma}^{n-i}(S_{n-i}^{*} - P^{*}S_{i}).$$
(3.6)

We choose $\beta, \gamma \in \mathbb{T}$ such that $\gamma^{n-i} = \beta^i = \eta$ and then by adding we get

$$2\tilde{n}_{i}(I - P^{*}P) \geq \eta \{ (S_{i} - S_{n-i}^{*}P) + (S_{n-i} - S_{i}^{*}P) \} + \bar{\eta} \{ (S_{i}^{*} - P^{*}S_{n-i}) + (S_{n-i}^{*} - P^{*}S_{i}) \}.$$
(3.7)

This shows that the Laurent polynomial

$$\xi(z) = 2\tilde{n}_i(I - P^*P) - z\{(S_i - S_{n-i}^*P) + (S_{n-i} - S_i^*P)\} - \bar{z}\{(S_i^* - P^*S_{n-i}) + (S_{n-i}^* - P^*S_i)\}$$
(3.8)

is non-negative for all $z \in \mathbb{T}$. Therefore, by the Operator Fejer-Riesz Theorem (see Theorem 1.2 in [13]) there is a polynomial of degree 1, say $P(z) = X_0 + X_1 z$, such that for all $z \in \mathbb{T}$,

$$\xi(z) = P(z)^* P(z) = (X_0^* + X_1^* \bar{z})(X_0 + X_1 z)$$

= $(X_0^* X_0 + X_1^* X_1) + (X_0^* X_1) z + X_1^* X_0 \bar{z}.$ (3.9)

Comparing (3.8) and (3.9) we obtain

$$2\tilde{n}_i D_P^2 = X_0^* X_0 + X_1^* X_1 \tag{3.10}$$

$$(S_i - S_{n-i}^* P) + (S_{n-i} - S_i^* P) = -X_0^* X_1.$$
(3.11)

Again putting $\gamma = -\beta$ in (3.6) and adding (3.5) and (3.6) we have that

$$2\tilde{n}_{i}(I - P^{*}P) \ge \beta \{ (S_{i} - S_{n-i}^{*}P) - (S_{n-i} - S_{i}^{*}P) \} + \bar{\beta} \{ (S_{i}^{*} - P^{*}S_{n-i}) - (S_{n-i}^{*} - P^{*}S_{i}) \}.$$
(3.12)

This shows that the Laurent polynomial

$$\eta(z) = 2\tilde{n}_i(I - P^*P) - z\{(S_i - S_{n-i}^*P) - (S_{n-i} - S_i^*P)\} - \bar{z}\{(S_i^* - P^*S_{n-i}) - (S_{n-i}^* - P^*S_i)\}$$
(3.13)

is non-negative for all $z \in \mathbb{T}$. Therefore, applying the Operator Fejer-Riesz Theorem again we have a polynomial of degree 1, say $Q(z) = Y_0 + Y_1 z$, such that for all $z \in \mathbb{T}$,

$$\eta(z) = Q(z)^* Q(z) = (Y_0^* + Y_1^* \bar{z})(Y_0 + Y_1 z)$$

= $(Y_0^* Y_0 + Y_1^* Y_1) + (Y_0^* Y_1) z + Y_1^* Y_0 \bar{z}.$ (3.14)

Comparing (3.13) and (3.14) we have

$$2\tilde{n}_i D_P^2 = Y_0^* Y_0 + Y_1^* Y_1 \tag{3.15}$$

$$(S_i - S_{n-i}^* P) - (S_{n-i} - S_i^* P) = -Y_0^* Y_1.$$
(3.16)

Adding (3.10) and (3.15) we have

$$4\tilde{n}_i D_P^2 = (X_0^* X_0 + X_1^* X_1) + (Y_0^* Y_0 + Y_1^* Y_1).$$
(3.17)

Similarly adding (3.11) and (3.16) we have

$$2(S_i - S_{n-i}^* P) = -(X_0^* X_1 + Y_0^* Y_1).$$
(3.18)

We obtain from (3.17) that

$$4\tilde{n}_i D_P^2 \ge X_0^* X_0, \ 4\tilde{n}_i D_P^2 \ge X_1^* X_1, \ 4\tilde{n}_i D_P^2 \ge Y_0^* Y_0 \text{ and } 4\tilde{n}_i D_P^2 \ge Y_1^* Y_1$$

So, by Douglas's lemma (see Lemma 2.1 in [12]) there are contractions Z_0, Z_1, Z_2, Z_3 such that

$$\begin{split} X_0^* &= 2\sqrt{\tilde{n}_i} D_P Z_0, \quad X_1^* &= 2\sqrt{\tilde{n}_i} D_P Z_1, \\ Y_0^* &= 2\sqrt{\tilde{n}_i} D_P Z_2, \quad Y_1^* &= 2\sqrt{\tilde{n}_i} D_P Z_3. \end{split}$$

Substituting these values in (3.18) we have that

$$S_i - S_{n-i}^* P = D_P[-2\tilde{n}_i(Z_0 Z_1^* + Z_2 Z_3^*)]D_P.$$

Setting $A_i = P_{\mathscr{D}_P}[-2\tilde{n}_i(Z_0Z_1^* + Z_2Z_3^*)]|_{\mathscr{D}_P}$, we have that $S_i - S_{n-i}^*P = D_PA_iD_P$,

which is true for all i = 1, ..., n - 1.

Adding (3.7) and (3.12) we have that

$$4\tilde{n}_i D_P^2 \ge 4 \operatorname{Re} \beta(S_i - S_{n-i}^* P) = 4 \operatorname{Re} \beta D_P A_i D_P.$$

Therefore,

$$D_P^2 \geq \operatorname{Re} \beta D_P(\frac{1}{\tilde{n}_i}A_i)D_P,$$

and thus

$$D_P[I_{\mathscr{D}_P} - \operatorname{Re} \beta(\frac{1}{\tilde{n}_i}A_i)]D_P \ge 0$$

This implies that

$$I_{\mathscr{D}_P} - \operatorname{Re} \beta(\frac{1}{\tilde{n}_i}A_i) \ge 0$$

because A_i is defined on \mathcal{D}_P . Therefore, by Lemma 3.2, we have

$$\boldsymbol{\omega}(A_i) \leq \tilde{n}_i = \binom{n}{i}.$$

Uniqueness. Let there be two solutions A_i, C_i of the equation $S_i - S_{n-i}^* P = D_P X_i D_P$. Then $D_P(A_i - C_i)D_P = 0$, which shows that $A_i - C_i = 0$ as $A_i - C_i$ is defined on \mathcal{D}_P . Thus A_i is unique for each i = 1, ..., n-1.

We shall see in the next subsection a partial converse to the existence-uniqueness of the \mathscr{F}_{O} -tuple. Under a certain condition, an operator tuple (A_1, \ldots, A_{n-1}) becomes the \mathscr{F}_{O} -tuple of a Γ_n -contraction (see Theorem 3.7).

Note 3.4. The \mathscr{F}_O -tuple of a Γ_n -isometry or a Γ_n -unitary $(S_1, \ldots, S_{n-1}, P)$ is defined to be $(0, \ldots, 0)$ because the \mathscr{F}_O -tuple is defined on the space \mathscr{D}_P and in such cases $\mathscr{D}_P = \{0\}$.

Proposition 3.5. If two Γ_n -contractions are unitarily equivalent then so are their \mathscr{F}_O -tuples.

Proof. Suppose $(S_{11}, \ldots, S_{1(n-1)}, P_1)$ and $(S_{21}, \ldots, S_{2(n-1)}, P_2)$ are two unitarily equivalent Γ_n contractions acting on Hilbert spaces \mathscr{H}_1 and \mathscr{H}_2 respectively with \mathscr{F}_O -tuples (F_1, \ldots, F_{n-1}) and (G_1, \ldots, G_{n-1}) . Then there is a unitary U from \mathscr{H}_1 to \mathscr{H}_2 such that

$$US_{11} = S_{21}U, \dots, US_{1(n-1)} = S_{2(n-1)}U$$
 and $UP_1 = P_2U$.

Obviously $UP_1^* = P_2^*U$ and consequently

$$UD_{P_1}^2 = U(I - P_1^*P_1) = U - P_2^*P_2U = D_{P_2}^2U.$$

Therefore, $UD_{P_1} = D_{P_2}U$. Let $V = U|_{\mathscr{D}_{P_1}}$. Then $V \in \mathscr{L}(\mathscr{D}_{P_1}, \mathscr{D}_{P_2})$ and $VD_{P_1} = D_{P_2}V$. Thus, using the fact that $S_{1i} - S^*_{1(n-i)}P_1$ and $S_{2i} - S^*_{2(n-i)}P_2$ are equal to 0 on the orthogonal complement of \mathscr{D}_{P_1} and \mathscr{D}_{P_2} respectively we have

$$D_{P_2}VF_iV^*D_{P_2} = VD_{P_1}F_iD_{P_1}V^* = V(S_{1i} - S_{1(n-i)}^*P_1)V^* = S_{2i} - S_{2n-i}^*P_2 = D_{P_2}G_iD_{P_2}.$$

So, F_i and G_i are unitarily equivalent for i = 1, ..., n-1 and the proof is complete.

Remark 3.6. The converse to the above result does not hold, i.e, two non-unitarily equivalent Γ_n -contractions can have unitarily equivalent \mathscr{F}_O -tuples. For example, if we consider a Γ_n -isometry on a Hilbert space which is not a Γ_n -unitary, then its \mathscr{F}_O -tuple is $(0, \ldots, 0)$ which is same as the \mathscr{F}_O -tuple of any Γ_n -unitary on the same Hilbert space.

A partial converse to the Existence-Uniqueness Theorem for the \mathscr{F}_O -tuple. The existence and uniqueness of the \mathscr{F}_O -tuple (Theorem 3.3) is in the center of all results of this article. Here we provide a partial converse to that result.

Theorem 3.7. Let A_1, \ldots, A_{n-1} be operators defined on a Hilbert space E such that

$$\left(\frac{n-1}{n}(A_1^*+A_{n-1}z),\frac{n-2}{n}(A_2^*+A_{n-2}z),\dots,\frac{1}{n}(A_{n-1}^*+A_{1}z)\right)$$

is a Γ_{n-1} -contraction for all $z \in \mathbb{T}$. Then there is a Γ_n -contraction for which (A_1, \ldots, A_{n-1}) is the \mathscr{F}_O -tuple.

Proof. Let us consider the vectorial Hardy-Hilbert space $H^2(E)$ and the Toeplitz operator tuple $(T_{A_1^*+A_{n-1}z}, \ldots, T_{A_{n-1}^*+A_1z}, T_z)$ acting on it. Here T_z on $H^2(E)$ is the shift operator. We shall show that $(T_{A_1^*+A_{n-1}z}^*, \ldots, T_{A_{n-1}^*+A_1z}^*, T_z^*)$ is a Γ_n -co-isometry and (A_1, \ldots, A_{n-1}) is the \mathscr{F}_O -tuple of it. Since for all $z \in \mathbb{T}$,

$$\left(\frac{n-1}{n}(A_1^*+A_{n-1}z),\frac{n-2}{n}(A_2^*+A_{n-2}z),\dots,\frac{1}{n}(A_{n-1}^*+A_{1}z)\right)$$

is a Γ_{n-1} -contraction, so is

$$\left(\frac{n-1}{n}M_{A_1^*+A_{n-1}z},\frac{n-2}{n}M_{A_2^*+A_{n-2}z},\ldots,\frac{1}{n}M_{A_{n-1}^*+A_1z}\right),$$

where each of the multiplication operators is defined on $L^2(E)$. It is evident that for each $z \in \mathbb{T}$, $M_{A_i^*+A_{n-iz}} = M_{A_{n-i}^*+A_{iz}}^*M_z$ and M_z on $L^2(E)$ is unitary. So by Theorem 2.4, the multiplication operator tuple $(M_{A_1^*+A_{n-1}z}, \ldots, M_{A_{n-1}^*+A_{1z}}, M_z)$ on $L^2(E)$ is a Γ_n -unitary and the Toeplitz operator tuple $(T_{A_1^*+A_{n-1}z}, \ldots, T_{A_{n-1}^*+A_{1z}}, T_z)$, being the restriction of $(M_{A_1^*+A_{n-1}z}, \ldots, M_{A_{n-1}^*+A_{1z}}, M_z)$ to the common invariant subspace $H^2(E)$, is a Γ_n -isometry. Therefore, $(T_{A_1^*+A_{n-1}z}^*, \ldots, T_{A_{n-1}^*+A_{1z}}, T_z^*)$ is a Γ_n -co-isometry. We now compute the \mathscr{P}_O -tuple of $(T_{A_1^*+A_{n-1}z}^*, \ldots, T_{A_{n-1}^*+A_{1z}}, T_z^*)$. Now for each $i = 1, \ldots, n-1$,

$$T_{A_{i}^{*}+A_{n-i}z}^{*}-T_{A_{n-i}^{*}+A_{i}z}T_{z}^{*}=T_{A_{i}+A_{n-i}^{*}\bar{z}}-T_{A_{n-i}^{*}+A_{i}z}T_{\bar{z}}=T_{A_{i}}=A_{i}.$$

Again since $I - T_z T_z^*$ is the projection onto the space $\mathscr{D}_{T_z^*}(=E)$,

$$(I - T_z T_z^*)^{\frac{1}{2}} A_i (I - T_z T_z^*)^{\frac{1}{2}} = A$$

Therefore, by the uniqueness of the \mathscr{F}_O -tuple, (A_1, \ldots, A_{n-1}) is the \mathscr{F}_O -tuple of the Γ_n -coisometry $(T^*_{A_1^*+A_{n-1}z}, \ldots, T^*_{A_{n-1}^*+A_1z}, T^*_z)$.

$C_{\cdot 0}\Gamma_n$ -CONTRACTIONS

4. DILATION AND MODEL THEORY FOR $C_{.0} \Gamma_n$ -contractions

4.1. **Dilation.** We have witnessed in [19] that in general rational dilation fails on the symmetrized polydisc in any dimension greater than 2. In this subsection, we shall determine a class of $C_{.0} \Gamma_n$ -contractions that dilate to the distinguished boundary $b\Gamma_n$. Indeed, we impose certain conditions on the \mathscr{F}_0 -tuple of a $C_{.0} \Gamma_n$ -contraction to obtain a Γ_n -isometric dilation and then extend that Γ_n -isometry to a Γ_n -unitary which eventually becomes a Γ_n -unitary dilation.

Definition 4.1. Let $(S_1, \ldots, S_{n-1}, P)$ be a Γ_n -contraction on \mathscr{H} . A commuting operator tuple $(T_1, \ldots, T_{n-1}, V)$ defined on \mathscr{K} is said to be a Γ_n -isometric dilation of $(S_1, \ldots, S_{n-1}, P)$ if $\mathscr{H} \subseteq \mathscr{K}, (T_1, \ldots, T_{n-1}, V)$ is a Γ_n -isometry and

$$P_{\mathscr{H}}(T_1^{m_1}\dots T_{n-1}^{m_{n-1}}V^n)|_{\mathscr{H}}=S_1^{m_1}\dots S_{n-1}^{m_{n-1}}P^n,$$

for all non-negative integers m_1, \ldots, m_{n-1}, n . Moreover, the dilation is called *minimal* if the following holds:

$$\mathscr{H} = \overline{\operatorname{span}} \{ T_1^{m_1} \dots T_{n-1}^{m_{n-1}} V^n h : h \in \mathscr{H} \text{ and } m_1, \dots, m_{n-1}, n \in \mathbb{N} \cup \{0\} \}.$$

In a similar fashion one can define a Γ_n -unitary dilation of a Γ_n -contraction.

Needless to mention that a Γ_n -unitary dilation of a Γ_n -contraction is a normal dilation to the distinguished boundary $b\Gamma_n$. Before going to the explicit dilation, we state and prove the following result which we use in the proof of the dilation theorem.

Proposition 4.2. Let $(T_1, \ldots, T_{n-1}, V)$ on \mathscr{K} be a Γ_n -isometric dilation of a Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$ on \mathscr{H} . If $(T_1, \ldots, T_{n-1}, V)$ is minimal, then $(T_1^*, \ldots, T_{n-1}^*, V^*)$ is a Γ_n -co-isometric extension of $(S_1^*, \ldots, S_{n-1}^*, P^*)$. Conversely, the adjoint of a Γ_n -co-isometric extension of $(S_1, \ldots, S_{n-1}, P)$ is a Γ_n -isometric dilation of $(S_1, \ldots, S_{n-1}, P)$.

Proof. We first prove that $S_i P_{\mathcal{H}} = P_{\mathcal{H}} T_i$ for each *i* and $PP_{\mathcal{H}} = P_{\mathcal{H}} V$. Clearly

 $\mathscr{K} = \overline{\operatorname{span}} \{ T_1^{m_1} \dots T_{n-1}^{m_{n-1}} V^n h : h \in \mathscr{H} \text{ and } m_1, \dots, m_{n-1}, n \in \mathbb{N} \cup \{0\} \}.$

Now for $h \in \mathscr{H}$ we have that

$$S_{i}P_{\mathscr{H}}(T_{1}^{m_{1}}\dots T_{n-1}^{m_{n-1}}V^{n}h) = S_{i}(S_{1}^{m_{1}}\dots S_{n-1}^{m_{n-1}}P^{n}h)$$

= $S_{1}^{m_{1}}\dots S_{i}^{m_{i}+1}\dots S_{n-1}^{m_{n-1}}P^{n}h$
= $P_{\mathscr{H}}(T_{1}^{m_{1}}\dots T_{i}^{m_{i}+1}\dots T_{n-1}^{m_{n-1}}V^{n}h)$
= $P_{\mathscr{H}}T_{i}(T_{1}^{m_{1}}\dots T_{n-1}^{m_{n-1}}V^{n}h).$

Thus, $S_i P_{\mathcal{H}} = P_{\mathcal{H}} T_i$. Similarly we can prove that $PP_{\mathcal{H}} = P_{\mathcal{H}} V$. Also for $h \in \mathcal{H}$ and $k \in \mathcal{K}$ we have that

$$\langle S_i^*h,k\rangle = \langle P_{\mathscr{H}}S_i^*h,k\rangle = \langle S_i^*h,P_{\mathscr{H}}k\rangle = \langle h,S_iP_{\mathscr{H}}k\rangle = \langle h,P_{\mathscr{H}}T_ik\rangle = \langle T_i^*h,k\rangle.$$

Hence $S_i^* = T_i^*|_{\mathscr{H}}$ and similarly $P^* = V^*|_{\mathscr{H}}$. Therefore, $(T_1^*, \ldots, T_{n-1}^*, V^*)$ is a Γ_n -co-isometric extension of $(S_1^*, \ldots, S_{n-1}^*, P^*)$. The converse part is obvious.

We are now in a position to present the desired dilation theorem which is one of the main results of this article.

Theorem 4.3. Let $(S_1, \ldots, S_{n-1}, P)$ be a C_0 Γ_n -contraction on a Hilbert space \mathscr{H} and let the \mathscr{F}_O -tuple (B_1, \ldots, B_{n-1}) of $(S_1^*, \ldots, S_{n-1}^*, P^*)$ be such that

$$\left(\frac{n-1}{n}(B_1^*+B_{n-1}z),\frac{n-2}{n}(B_2^*+B_{n-2}z),\ldots,\frac{1}{n}(B_{n-1}^*+B_1z)\right)$$

is a Γ_{n-1} -contraction for all $z \in \mathbb{T}$. Consider the operators T_1, \ldots, T_{n-1}, V on $\mathscr{K} = H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*}$ defined by $T_i = I \otimes B_i^* + M_z \otimes B_{n-i}$ for $i = 1, \ldots, n-1$ and $V = M_z \otimes I$. Then the n-tuple $(T_1, \ldots, T_{n-1}, V)$ is a minimal $C_{\cdot 0} \Gamma_n$ -isometric dilation of $(S_1, \ldots, S_{n-1}, P)$.

Proof. We have from the Sz.-Nagy-Foias theory for $C_{.0}$ contractions [7] that \mathscr{K} is the minimal isometric dilation space and the operator V is the minimal isometric dilation of P. So, the minimality of the dilation follows trivially. It suffices to prove that $(T_1, \ldots, T_{n-1}, V)$ is a Γ_n -isometric dilation of $(S_1, \ldots, S_{n-1}, P)$. By virtue of Lemma 4.2, it suffices to show that $(T_1^*, \ldots, T_{n-1}^*, V^*)$ is a Γ_n -co-isometric extension of $(S_1^*, \ldots, S_{n-1}^*, P^*)$. Since

$$\left(\frac{n-1}{n}(B_1^*+B_{n-1}z),\frac{n-2}{n}(B_2^*+B_{n-2}z),\dots,\frac{1}{n}(B_{n-1}^*+B_1z)\right)$$

is a Γ_{n-1} -contraction for all $z \in \mathbb{T}$, it follows from the proof of Theorem 3.7 that the Toeplitz operator tuple $(T_{B_1^*+B_{n-1}z},\ldots,T_{B_{n-1}^*+B_1z},T_z)$ on $H^2(\mathscr{D}_{P^*})$ is a $C_{\cdot 0}$ Γ_n -isometry with (B_1,\ldots,B_{n-1}) being the \mathscr{F}_O -tuple of its adjoint, $(T_{B_1^*+B_{n-1}z}^*,\ldots,T_{B_{n-1}^*+B_1z}^*,T_z^*)$. Also it is a $C_{\cdot 0}$ Γ_n -isometry as T_z is a $C_{\cdot 0}$ isometry. Again since the Toeplitz operator tuple $(T_{B_1^*+B_{n-1}z},\ldots,T_{B_{n-1}^*+B_1z},T_z)$ on $H^2(\mathscr{D}_{P^*})$ is unitarily equivalent to (T_1,\ldots,T_2,V) on $\mathscr{K} = H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*}, (T_1,\ldots,T_{n-1},V)$ is a $C_{\cdot 0}$ Γ_n -isometry. All we have to prove now is that $(T_1^*,\ldots,T_{n-1}^*,V^*)$ is a Γ_n -co-isometric extension of $(S_1^*,\ldots,S_{n-1}^*,P^*)$.

Let us define $W : \mathscr{H} \to \mathscr{K}$ by $Wh = \sum_{n=0}^{\infty} z^n \otimes D_{P^*} P^{*n} h$. Now

$$\begin{split} Wh\|^{2} &= \|\sum_{n=0}^{\infty} z^{n} \otimes D_{P^{*}} P^{*n} h\|^{2} \\ &= \langle \sum_{n=0}^{\infty} z^{n} \otimes D_{P^{*}} P^{*n} h, \sum_{m=0}^{\infty} z^{m} \otimes D_{P^{*}} P^{*m} h \rangle \\ &= \sum_{m,n=0}^{\infty} \langle z^{n}, z^{m} \rangle \langle D_{P^{*}} P^{*n} h, D_{P^{*}} P^{*m} h \rangle \\ &= \sum_{n=1}^{\infty} \langle P^{n} D_{P^{*}}^{2} P^{*n} h, h \rangle \\ &= \sum_{n=0}^{\infty} \langle P^{n} (I - PP^{*}) P^{*n} h, h \rangle \\ &= \sum_{n=0}^{\infty} \{ \langle P^{n} P^{*n} h, h \rangle - \langle P^{n+1} P^{*n+1} h, h \rangle \} \\ &= \|h\|^{2} - \lim_{n \to \infty} \|P^{*n} h\|^{2}. \end{split}$$

Since *P* is a pure contraction, $\lim_{n\to\infty} ||P^{*n}h||^2 = 0$ and thus ||Wh|| = ||h||. Therefore *W* is an isometry. For a basis vector $z^n \otimes \eta$ of \mathscr{K} we have that

$$\langle W^*(z^n\otimes\eta),h\rangle = \langle z^n\otimes\eta,\sum_{k=0}^{\infty}z^k\otimes D_{P^*}P^{*k}h\rangle = \langle \eta,D_{P^*}P^{*n}h\rangle = \langle P^nD_{P^*}\xi,h\rangle.$$

Therefore,

$$W^*(z^n \otimes \eta) = P^n D_{P^*} \eta, \quad \text{for } n = 0, 1, 2, 3, \dots$$
 (4.1)

and hence

$$PW^*(z^n \otimes \eta) = P^{n+1}D_{P^*}\eta$$
, for $n = 0, 1, 2, 3, ...$

Again by (4.1),

$$W^*V(z^n \otimes \eta) = W^*(M_z \otimes I)(z^n \otimes \eta) = W^*(z^{n+1} \otimes \eta) = P^{n+1}D_{P^*}\eta = PW^*(z^n \otimes \eta).$$

Consequently, $W^*V = PW^*$, i.e, $V^*W = WP^*$ and hence $V^*|_{W(\mathscr{H})} = WP^*W^*|_{W(\mathscr{H})}.$

We now show that $W^*T_i = S_iW^*$ for each i = 1, ..., n-1; $W^*T_i(z^n \otimes \eta) = W^*(I \otimes B_i^* + M_z \otimes B_{n-i})(z^n \otimes \eta)$ $= W^*(z^n \otimes B_i^*\eta) + W^*(z^{n+1} \otimes B_{n-i}\eta)$ $= P^n D_{P^*}B_i^*\eta + P^{n+1}D_{P^*}B_{n-1}\eta.$

Also for each *i*,

$$S_i W^*(z^n \otimes \eta) = S_i P^n D_{P^*} \eta = P^n S_i D_{P^*} \eta.$$

$$(4.2)$$

Claim 1. $S_i D_{P^*} = D_{P^*} B_i^* + P D_{P^*} B_{n-i}$. Proof of Claim 1. Since (B_1, \ldots, B_{n-1}) is the \mathscr{F}_O -tuple of $(S_1^*, \ldots, S_{n-1}^*, P^*)$, we have

$$(D_{P^*}B_i^* + PD_{P^*}B_{n-i})D_{P^*} = (S_i - PS_{n-i}^*) + P(S_{n-i}^* - S_iP^*) = S_iD_{P^*}^2$$

Now if $G = S_i D_{P^*} - D_{P^*} B_i^* - P D_{P^*} B_{n-i}$, then *G* is defined from \mathscr{D}_{P^*} to \mathscr{H} and $G D_{P^*} h = 0$ for every $h \in \mathscr{D}_{P^*}$. Thus the claim is proved.

So from (4.2) we have $S_iW^*(z^n \otimes \eta) = P^n(D_{P^*}B_i^* + PD_{P^*}B_{n-i})$. Therefore, $W^*T_i = S_iW^*$ and hence $T_i^*|_{W(\mathscr{H})} = WS_i^*W^*|_{W(\mathscr{H})}$ for each *i*. Hence the proof is complete.

Corollary 4.4. Let $(S_1, \ldots, S_{n-1}, P)$ be a Γ_n -contraction which satisfies the hypotheses of Theorem 4.3. Then the multiplication operator tuple $(M_{B_1^*+B_{n-1}z}, \ldots, M_{B_{n-1}^*+B_1z}, M_z)$ on $L^2(\mathcal{D}_{P^*})$ is a Γ_n -unitary dilation of $(S_1, \ldots, S_{n-1}, P)$.

Proof. Since the Γ_n -isometric dilation that is obtained in Theorem 4.3 is unitarily equivalent to Toeplitz operator tuple $\left(T_{B_1^*+B_{n-1}z},\ldots,T_{B_{n-1}^*+B_1z},T_z\right)$ on $H^2(\mathscr{D}_{P^*})$ and since the multiplication operator tuple $\left(M_{B_1^*+B_{n-1}z},\ldots,M_{B_{n-1}^*+B_1z},M_z\right)$ on $L^2(\mathscr{D}_{P^*})$ is a natural Γ_n -unitary extension of it, the assertion follows obviously.

4.2. **Operator models.** We recall from [7] the notion of the characteristic function of a contraction *T*. For a contraction *T* defined on a Hilbert space \mathscr{H} , let Λ_T be the set of all complex numbers for which the operator $I - zT^*$ is invertible. For $z \in \Lambda_T$, the characteristic function of *T* is defined as

$$\Theta_T(z) = [-T + zD_{T^*}(I - zT^*)^{-1}D_T]|_{\mathscr{D}_T}.$$
(4.3)

By virtue of the relation $TD_T = D_{T^*}P$ (section I.3 of [7]), $\Theta_T(z)$ maps $\mathscr{D}_T = \overline{\operatorname{Ran}}D_T$ into $\mathscr{D}_{T^*} = \overline{\operatorname{Ran}}D_{T^*}$ for every z in Λ_T . Let us define

$$\mathscr{H}_P = (H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*}) \ominus M_{\Theta_P}(H^2(\mathbb{D}) \otimes \mathscr{D}_P).$$

In [7], Sz.-Nagy and Foias showed that every $C_{.0}$ contraction P defined on a Hilbert space \mathscr{H} is unitarily equivalent to the operator $P_{\mathscr{H}_P}(M_z \otimes I_{\mathscr{D}_{P^*}})|_{\mathscr{H}_P}$ on the Hilbert space \mathscr{H}_P , where $P_{\mathscr{H}_P}$ is the projection of $H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*}$ onto \mathscr{H}_P .

In this subsection, we independently find a model, which is not necessarily commutative, for a $C_{.0} \Gamma_n$ -contraction $(S_1, \ldots, S_{n-1}, P)$ without assuming any condition on it. On the minimal

 \square

dilation space \mathscr{H}_P of P let R_1, \ldots, R_{n-1} be the compressions of T_1, \ldots, T_{n-1} (as in Theorem 4.3) respectively to \mathscr{H}_P . Evidently R_1, \ldots, R_{n-1} do not commute in general, yet we shall see that they are individually unitarily equivalent to S_1, \ldots, S_{n-1} respectively. For P, we choose the Nagy-Foias model $R = P_{\mathscr{H}_P}(M_z \otimes I_{\mathscr{D}_{P^*}})|_{\mathscr{H}_P}$. We also show that under the hypotheses of Theorem 4.3, $(R_1, \ldots, R_{n-1}, R)$ on \mathscr{H} is unitarily equivalent to $(S_1, \ldots, S_{n-1}, P)$ and thus in this case it provides a commutative model. Before going to the main results we state a lemma whose proof could be found in [9] (see Lemma 3.3 in [9]). For the sake of completeness we include the proof here too.

Lemma 4.5. For every contraction P, the identity

$$WW^* + M_{\Theta_P} M_{\Theta_P}^* = I_{H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*}}$$

$$\tag{4.4}$$

holds, where W is the isometry mentioned in the proof of Theorem 4.3.

Proof. The operator W^* satisfies the identity

$$W^*(k_z \otimes \xi) = (I - \overline{z}P)^{-1}D_{P^*}\xi \text{ for } z \in \mathbb{D} \text{ and } \xi \in \mathscr{D}_{P^*},$$

where $k_z(w) := (1 - \langle w, z \rangle)^{-1}$ for all $w \in \mathbb{D}$. For a proof one can see Theorem 1.2 in [6]. Therefore we have

$$\langle (WW^* + M_{\Theta_P}M_{\Theta_P}^*)(k_z \otimes \xi), (k_w \otimes \eta) \rangle$$

$$= \langle W^*(k_z \otimes \xi), W^*(k_w \otimes \eta) \rangle + \langle M_{\Theta_P}^*(k_z \otimes \xi), M_{\Theta_P}^*(k_w \otimes \eta) \rangle$$

$$= \langle (I - \bar{z}P)^{-1}D_{P^*}\xi, (I - \bar{w}P)^{-1}D_{P^*}\eta \rangle + \langle k_z \otimes \Theta_P(z)^*\xi, k_w \otimes \Theta_P(w)^*\eta \rangle$$

$$= \langle D_{P^*}(I - wP^*)^{-1}(I - \bar{z}P)^{-1}D_{P^*}\xi, \eta \rangle + \langle k_z, k_w \rangle \langle \Theta_P(w)\Theta_P(z)^*\xi, \eta \rangle$$

$$= \langle k_z \otimes \xi, k_w \otimes \eta \rangle \text{ for all } z, w \in \mathbb{D} \text{ and } \xi, \eta \in \mathscr{D}_{P^*}.$$

The last equality follows from the following well-known identity

$$I - \Theta_P(w)\Theta_P(z)^* = (1 - w\overline{z})D_{P^*}(I - wP^*)^{-1}(I - \overline{z}P)^{-1}D_{P^*}.$$

Now using the fact that $\{k_z : z \in \mathbb{D}\}$ forms a total set of $H^2(\mathbb{D})$, the assertion follows.

Theorem 4.6. Let $(S_1, \ldots, S_{n-1}, P)$ be a C_0 Γ_n -contraction on a Hilbert space \mathcal{H} . Then the operators S_i and P are unitarily equivalent to $P_{\mathcal{H}_P}(I \otimes B_i^* + M_z \otimes B_{n-i})|_{\mathcal{H}_P}$ and $P_{\mathcal{H}_P}(M_z \otimes I_{\mathcal{D}_{P^*}})|_{\mathcal{H}_P}$ respectively, where (B_1, \ldots, B_{n-1}) is the \mathcal{F}_O -tuple of $(S_1^*, \ldots, S_{n-1}^*, P^*)$.

Proof. Since W is an isometry, WW^* is the projection onto the range of W and since P is pure, M_{Θ_P} is also an isometry. So by Lemma 4.5, we have that

$$W(\mathscr{H}_P) = (H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*}) \ominus M_{\Theta_P}(H^2(\mathbb{D}) \otimes \mathscr{D}_P).$$

So, we have

$$W^{*}(I \otimes B_{i}^{*} + M_{z} \otimes B_{n-i})(z^{n} \otimes \xi) = W^{*}(z^{n} \otimes B_{i}^{*}\xi) + W^{*}(z^{n+1} \otimes B_{n-i}\xi)$$

= $P^{n}D_{P^{*}}B_{i}^{*}\xi + P^{n+1}D_{P^{*}}B_{n-i}\xi$
= $P^{n}(D_{P^{*}}B_{i}^{*} + PD_{P^{*}}B_{n-i})\xi$
= $P^{n}S_{i}D_{P^{*}}\xi$ [by Claim 1 in Theorem 4.3]
= $S_{i}P^{n}D_{P^{*}}\xi = S_{i}W^{*}(z^{n} \otimes \xi).$

So for each *i*, we have $W^*(I \otimes B_i^* + M_z \otimes B_{n-i}) = S_i W^*$ for the vectors of the form $z^n \otimes \xi$, for all $n \ge 0$ and $\xi \in \mathscr{D}_{P^*}$ which span $H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*}$. Hence we have $W^*(I \otimes B_i^* + M_z \otimes B_{n-i}) = S_i W^*$, which implies that $W^*(I \otimes B_i^* + M_z \otimes B_{n-i})W = S_i$. Therefore, S_i is unitarily equivalent to $P_{\mathscr{H}_P}(I \otimes B_i^* + M_z \otimes B_{n-i})|_{\mathscr{H}_P}$. Again

$$W^*(M_z \otimes I)(z^n \otimes \xi) = W^*(z^{n+1} \otimes \xi) = P^{n+1}D_{P^*}\xi = PW^*(z^n \otimes \xi).$$

Therefore by the same argument as above, P is unitarily equivalent to $P_{\mathcal{H}_P}(M_z \otimes I_{\mathcal{D}_{P^*}})|_{\mathcal{H}_P}$. \Box

Corollary 4.7. For any $C_{\cdot 0}$ Γ_n -contraction $(S_1, \ldots, S_{n-1}, P)$, there are operators C_1, \ldots, C_{n-1} satisfying $\omega(C_i + C_{n-i}z) \leq \binom{n}{i}$ for all $z \in \mathbb{T}$ and such that $S_i = C_i + PC_{n-i}^*$ for $i = 1, \ldots, n-1$.

Proof. From the previous theorem we have that

$$W^*(I \otimes B_i^* + M_z \otimes B_{n-i}) = S_i W^*$$

or $W^*(I \otimes B_i^* + M_z \otimes B_{n-i})W = S_i$
or $W^*(I \otimes B_i^*)W + W^*(M_z \otimes I)(I \otimes B_{n-i})W = S_i$
or $W^*(I \otimes B_i^*)W + PW^*(I \otimes B_{n-i})W = S_i$, since $W^*(M_z \otimes I) = PW^*$.

Taking $C_i = W^*(I \otimes B_i^*)W$ for each i = 1, ..., n-1, we get $S_i = C_i + PC_{n-i}^*$. The fact that $\omega(C_i + C_{n-i}z) \leq {n \choose i}$ for all $z \in \mathbb{T}$ is obvious.

Corollary 4.8. If $(S_1, \ldots, S_{n-1}, P)$ is a Γ_n -contraction with ||P|| < 1, then there are n-1 unique operators C_1, \ldots, C_{n-1} such that $S_i = C_i + PC_{n-i}^*$.

Proof. Let there be C_1, \ldots, C_{n-1} and $\tilde{C}_1, \ldots, \tilde{C}_{n-1}$ such that $S_i = C_i + PC_{n-i}^*$ and $S_i = \tilde{C}_i + P\tilde{C}_{n-i}^*$. Then we have $D_i + PD_{n-i}^* = 0$, where $D_i = C_i - \tilde{C}_{n-i}$. Now

$$||D_i|| = ||-PD_{n-i}^*|| \le ||P|| ||D_{n-i}|| < ||D_{n-i}||$$
 as $||P|| < 1$.

Since this holds for each *i*, replacing *i* by n - i we get $||D_{n-i}|| < ||D_i||$. This shows that $D_i = 0$ for each *i* and consequently $C_i = \tilde{C}_i$.

Theorem 4.9. Let $(S_1, \ldots, S_{n-1}, P)$ be a $C_{\cdot 0}$ Γ_n -contraction on a Hilbert space \mathscr{H} that satisfies the hypotheses of Theorem 4.3. Then $(S_1, \ldots, S_{n-1}, P)$ is unitarily equivalent to the tuple $(R_1, \ldots, R_{n-1}, R)$ on the Hilbert space $\mathscr{H}_P = (H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*}) \ominus M_{\Theta_P}(H^2(\mathbb{D}) \otimes \mathscr{D}_P)$ defined by


$$R_{i} = P_{\mathcal{H}_{P}}(I \otimes B_{i}^{*} + M_{z} \otimes B_{n-i})|_{\mathcal{H}_{P}}, \text{ for } i = 1, \dots, n-1$$

and $R = P_{\mathcal{H}_{P}}(M_{z} \otimes I)|_{\mathcal{H}_{P}}.$

Proof. Following the proof of Theorem 4.3, it suffices to show that $W(\mathscr{H}) = \mathscr{H}_P$, which follows from the fact that $WW^* + M_{\Theta_P}M^*_{\Theta_P} = I_{H^2(\mathbb{D})\otimes\mathscr{D}_{P^*}}$. Hence the proof is complete.

5. A COMPLETE UNITARY INVARIANT FOR $C_{\cdot 0} \Gamma_n$ -contractions

A complete unitary invariant for a class of operator tuples, defined on a Hilbert space \mathscr{H} , is a necessary and sufficient condition under which any two such operator tuples are unitarily equivalent, that is, there is a unitary on \mathscr{H} that intertwines the corresponding components of the two tuples. For a pair of contractions P, P' acting on Hilbert spaces \mathscr{H} and \mathscr{H}' respectively, we say that the characteristic functions of P and P' coincide if there are unitary operators u: $\mathscr{D}_P \to \mathscr{D}_{P'}$ and $u_* : \mathscr{D}_{P^*} \to \mathscr{D}_{P'^*}$ such that the following diagram commutes for all $z \in \mathbb{D}$,



A few decades ago, Sz.-Nagy and Foias proved the following theorem, which asserts that the characteristic function is a complete unitary invariant for the c.n.u contractions.

Theorem 5.1 (Nagy-Foias, [7]). *Two completely non-unitary contractions are unitarily equivalent if and only if their characteristic functions coincide.*

Here we present a complete unitary invariant for the $C_{\cdot 0}$ Γ_n -contractions.

Theorem 5.2. Let $(S_1, \ldots, S_{n-1}, P)$ and $(S'_1, \ldots, S'_{n-1}, P')$ be two C_0 Γ_n -contractions defined on \mathscr{H} and \mathscr{H}' respectively. Suppose (B_1, \ldots, B_{n-1}) and (B'_1, \ldots, B'_{n-1}) are the \mathscr{F}_0 -tuples of $(S^*_1, \ldots, S^*_{n-1}, P^*)$ and $(S'^*_1, \ldots, S'^*_{n-1}, P'^*)$ respectively. Then $(S_1, \ldots, S_{n-1}, P)$ is unitarily equivalent to $(S'_1, \ldots, S'_{n-1}, P')$ if and only if $(B_1, \ldots, B_{n-1}, \Theta_P)$ and $(B'_1, \ldots, B'_{n-1}, \Theta_{P'})$ are unitarily equivalent.

Remark 5.3. The sense carried out by the unitary equivalence of

$$(B_1,\ldots,B_{n-1},\Theta_P)$$
 and $(B'_1,\ldots,B'_{n-1},\Theta_{P'})$

is that the characteristic functions of P and P' coincide and (B_1, \ldots, B_{n-1}) is unitarily equivalent to (B'_1, \ldots, B'_{n-1}) by the unitary $u_* : \mathscr{D}_{P^*} \to \mathscr{D}_{P'^*}$ that is involved in the coincidence of the characteristic functions of P and P'.

Proof. First let us assume that $(S_1, \ldots, S_{n-1}, P)$ and $(S'_1, \ldots, S'_{n-1}, P')$ are unitarily equivalent and let $U : \mathscr{H} \to \mathscr{H}'$ be a unitary such that $US_i = S'_i U$ for each *i* and UP = P'U. Since *P* is a *C*₀ contraction, it is a completely non-unitary contraction and hence by Theorem 5.1, the characteristic functions of *P* and *P'* coincides. The unitary $u_* : \mathscr{D}_{P^*} \to \mathscr{D}_{P'^*}$ that is involved in the coincidence of the characteristic functions Θ_P and $\Theta_{P'}$ is nothing but the restriction of *U* to \mathscr{D}_{P^*} that takes \mathscr{D}_{P^*} onto $\mathscr{D}_{P'^*}$. An interested reader can see Chapter VI of [7] for a proof of this fact. We now prove that the same unitary intertwines the \mathscr{F}_O -tuples of $(S^*_1, \ldots, S^*_{n-1}, P^*)$ and $(S'_1^*, \ldots, S'_{n-1}^*, P'^*)$. We have

$$UD_{P^*}^2 = U(I - PP^*) = U - P'P'^*U = D_{P'^*}^2U,$$

which gives $UD_{P^*} = D_{P'^*}U$. Let $\tilde{U} = U|_{\mathscr{D}_{P^*}}$. Then note that $\tilde{U} \in \mathscr{B}(\mathscr{D}_{P^*}, \mathscr{D}_{P'^*})$ and $\tilde{U}D_{P^*} = D_{P'^*}\tilde{U}$. Now for each *i*,

$$D_{P'^*}\tilde{U}B_i\tilde{U}^*D_{P'^*} = \tilde{U}D_{P^*}B_iD_{P^*}\tilde{U}^* = \tilde{U}(S_i^* - S_{n-i}P^*)\tilde{U}^*$$

= $S_i'^* - S_{n-i}'P'^* = D_{P'^*}B_i'D_{P'^*}$

Therefore we have $\tilde{U}B_i\tilde{U}^* = B'_i$ for each i = 1, ..., n-1.

We prove the converse part. Let $u : \mathscr{D}_P \to \mathscr{D}_{P'}$ and $u_* : \mathscr{D}_{P^*} \to \mathscr{D}_{P'^*}$ be unitary operators such that for each *i*

$$u_*B_i = B'_i u_*$$
 and $u_*\Theta_P(z) = \Theta_{P'}(z)u$ for all $z \in \mathbb{D}$.

The unitary operator $u_*: \mathscr{D}_{P^*} \to \mathscr{D}_{P'^*}$ induces the following unitary operator

$$U_*: H^2(\mathbb{D}) \otimes \mathscr{D}_{P^*} \to H^2(\mathbb{D}) \otimes \mathscr{D}_{P'^*}$$
$$(z^n \otimes \xi) \mapsto (z^n \otimes u_*\xi)\xi \in \mathscr{D}_{P^*}, \ n \ge 0.$$

We note here that

$$U_*(M_{\Theta_P}f(z)) = u_*\Theta_P(z)f(z) = \Theta_{P'}(z)uf(z) = M_{\Theta_{P'}}(uf(z)),$$

for all $f \in H^2(\mathbb{D}) \otimes \mathcal{D}_P$ and $z \in \mathbb{D}$. Hence U_* takes $\operatorname{Ran} M_{\Theta_P}$ onto $\operatorname{Ran} M_{\Theta_{P'}}$. Since U_* is unitary, we have

$$U_*(\mathscr{H}_P) = U_*((\operatorname{Ran} M_{\Theta_P})^{\perp}) = (U_* \operatorname{Ran} M_{\Theta_P})^{\perp} = (\operatorname{Ran} M_{\Theta_{P'}})^{\perp} = \mathscr{H}_{P'}.$$

Again from the definition of U_* we have that

$$U_*(I \otimes B_i^* + M_z \otimes B_{n-i})^* = (I \otimes u_*)(I \otimes B_i + M_z^* \otimes B_{n-i}^*)$$

= $I \otimes u_* B_i + M_z^* \otimes u_* B_{n-i}^*$
= $I \otimes B_i' u_* + M_z^* \otimes B_{n-i}' u_*$
= $(I \otimes B_i' + M_z^* \otimes B_{n-i}')(I \otimes u_*)$
= $(I \otimes B_i'^* + M_z \otimes B_{n-i}')^* U_*.$

Therefore, $\mathscr{H}_{P'} = U_*(\mathscr{H}_P)$ is a joint co-invariant subspace of $(I \otimes B'_i + M_z \otimes B'_{n-i})$ for i = 1, ..., n-1. Hence $P_{\mathscr{H}_P}(I \otimes B^*_i + M_z \otimes B_{n-i})|_{\mathscr{H}_P}$ and $P_{\mathscr{H}_{P'}}(I \otimes B'^*_i + M_z \otimes B'_{n-i})|_{\mathscr{H}_P}$ are unitarily equivalent for each *i*. It is evident that the unitary operator that intertwines them is $U_*|_{\mathscr{H}_P}$: $\mathscr{H}_P \to \mathscr{H}_{P'}$. Also we have

$$U_*(M_z \otimes I_{\mathscr{D}_{P^*}}) = (I \otimes u_*)(M_z \otimes I_{\mathscr{D}_{P^*}}) = (M_z \otimes I_{\mathscr{D}_{P'^*}})(I \otimes u_*) = (M_z \otimes I_{\mathscr{D}_{P'^*}})U_*.$$

So $P_{\mathcal{H}_P}(M_z \otimes I_{\mathcal{D}_{P^*}})|_{\mathcal{H}_P}$ and $P_{\mathcal{H}_{P'}}(M_z \otimes I_{\mathcal{D}_{P'^*}})|_{\mathcal{H}_{P'}}$ are unitarily equivalent by the same unitary $U_*|_{\mathcal{H}_P}$. Therefore $(S_1, \ldots, S_{n-1}, P)$ and $(S'_1, \ldots, S'_{n-1}, P')$ are unitarily equivalent and the proof is complete.

REFERENCES

- [1] J. Agler and J. McCarthy, *Pick Interpolation and Hilbert Function Spaces*, Graduate Studies in Mathematics 44, Amer. Math. Soc., Providence, R.I. 2002.
- [2] J. Agler and N. J. Young, A commutant lifting theorem for a domain in \mathbb{C}^2 and spectral interpolation, J. *Funct. Anal.* 161 (1999), 452 477.
- [3] J. Agler and N. J. Young, The two-point spectral Nevanlinna-Pick problem, *Integral Equations Operator Theory* 37 (2000), 375 385.
- [4] J. Agler and N. J. Young, A model theory for Γ -contractions, J. Operator Theory 49 (2003), 45 60.
- [5] H. Alexander and J. Wermer, Several complex variables and Banach algebras, Graduate Texts in Mathematics, 35; 3rd Edition, Springer, 1997.
- [6] W. B. Arveson, Subalgebras of C*-algebra. II., Acta Math. 128 (1972), 271 308.
- [7] H. Bercovici, C. Foias, L. Kerchy and B. Sz.-Nagy, Harmonic analysis of operators on Hilbert space, Universitext, Springer, New York, 2010.
- [8] T. Bhattacharyya, S. Pal and S. Shyam Roy, Dilations of Γ-contractions by solving operator equations, Adv. Math. 230 (2012), 577 – 606.
- [9] T. Bhattacharyya and S. Pal, A functional model for pure Γ-contractions, J. Operator Theory 71 (2014), 327 339.
- [10] S. Biswas and S. Shyam Roy, Functional models for Γ_n -contractions and characterization of Γ_n -isometries, *J. Func. Anal.* 266 (2014), 6224 – 6255.
- [11] C. Costara, On the spectral Nevanlinna-Pick problem, *Studia Math.* 170 (2005), 23 55.

- [12] R. G. Douglas, P. S. Muhly and C. Pearcy, Lifting Commuting Operators, *Michigan Math. J.* 15 (1968), 385-395.
- [13] M. A. Dritschel and J. Rovnyak, The operator Fejér-Riesz theorem, Oper. Theory Adv. Appl. 207 (2010), 223 – 254.
- [14] A. Edigarian and W. Zwonek, Geometry of the symmetrized polydisc, *Archiv der Mathematik* 84 (2005), 364 374.
- [15] C. Foias and A. E. Frazho, The commutant lifting approach to interpolation problems, *Operator Theory: Advances and Applications* 44, Birkhuser Verlag, Basel, 1990.
- [16] A. Pal, On Γ_n -contractions and their conditional dilations, *https://arxiv.org/abs/1704.04508*.
- [17] S. Pal, From Stinespring dilation to Sz.-Nagy dilation on the symmetrized bidisc and operator models, *New York J. Math.* 20 (2014), 545 564.
- [18] S. Pal, Canonical decomposition of operators associated with the symmetrized polydisc, *Complex Anal. Oper. Theory*, 12 (2018), 931 943.
- [19] S. Pal, The failure of rational dilation on the symmetrized *n*-disk for any $n \ge 3$, *https://arxiv.org/abs/1712.05707*.
- [20] S. Pal, Operator theory and distinguished varieties in the symmetrized *n*-disk, *https://arxiv.org/abs/1708.06015*.
- [21] S. Pal and O. M. Shalit, Spectral sets and distinguished varieties in the symmetrized bidisc, *J. Funct. Anal.* 266 (2014), 5779 5800.
- [22] V. Paulsen, Completely Bounded Maps and Operator Algebras, Cambridge University Press 2002.
- [23] J. Sarkar, Operator theory on symmetrized bidisc, Indiana Univ. Math. J. 64 (2015), 847 873.
- [24] Bela Sz.-Nagy, Sur les contractions de l'espace de Hilbert, Acta Sci. Math. 15 (1953), 87 92.
- [25] F. H. Vasilescu, Analytic Functional Calculus and Spectral Decompositions, *Editura Academiei: Bucuresti, Romania and D. Reidel Publishing Company*, 1982.
- [26] J. von Neumann, Eine Spektraltheorie f
 ür allgemeine Operatoren eines unit
 ären Raumes, Math. Nachr. 4 (1951), 258 – 281.

(Sourav Pal) MATHEMATICS DEPARTMENT, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUM-BAI - 400076, INDIA.

Email address: souravmaths@gmail.com , sourav@math.iitb.ac.in