

**Globally positive solutions of linear parabolic PDEs
of second order with Robin boundary conditions**

JANUSZ MIERCZYŃSKI*

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with boundary $\partial\Omega$ of class C^2 .

We consider a linear parabolic differential equation of second order

$$(1a) \quad u_t = \sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left(a_{ij}(x) \frac{\partial u}{\partial x_j} \right) + a_0(t, x)u, \quad t \in \mathbb{R}, x \in \Omega,$$

where $a_{ij} = a_{ji} \in C^1(\overline{\Omega})$, $\sum_{i,j=1}^n a_{ij}(x)\xi_i\xi_j > 0$ for each $x \in \overline{\Omega}$ and each nonzero $(\xi_1, \dots, \xi_n) \in \mathbb{R}^n$, and $a_0 \in L^\infty(\mathbb{R} \times \Omega)$.

Equation (1a) is complemented with regular oblique (Robin) homogeneous boundary conditions

$$(1b) \quad \frac{\partial u}{\partial \nu}(t, x) + c(x)u(t, x) = 0, \quad t \in \mathbb{R}, x \in \partial\Omega,$$

where $\nu \in C^1(\partial\Omega, \mathbb{R}^n)$ is a vector field pointing out of Ω and $c \in C^1(\partial\Omega)$ is a nonnegative function.

Motivated by deep results on one-dimensional parabolic PDE's contained in a series of papers [4], [5] by S.-N. Chow, K. Lu and J. Mallet-Paret, I investigate the set of solutions to (1a)+(1b) that are globally positive.

Our main result is the following (see Corollary 2.4).

Theorem. *Let u_1, u_2 be solutions of (1a)+(1b) defined and positive for all $t \in \mathbb{R}$ and all $x \in \overline{\Omega}$. Then there is a positive constant κ such that $u_1(t, x) = \kappa u_2(t, x)$ for all $t \in \mathbb{R}$ and all $x \in \overline{\Omega}$.*

A key idea in the proofs is to consider a family of linear parabolic PDEs obtained by letting the coefficient $a_0(t, x)$ vary in some metrizable compact subset of the vector space $L^\infty(\mathbb{R} \times \Omega)$ endowed with the weak* topology. That family generates in a natural way a compact linear skew-product semiflow on a (product) vector bundle with generic fiber $L^1(\Omega)$.

For n arbitrary, the results obtained seem to be new even in the case of autonomous equations (compare the remarks on p. 288 in [5]).

© 1997. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Published in *Journal of Mathematical Analysis and Applications* **209**(1) (1997), pp. 47–59.

<https://doi.org/10.1006/jmaa.1997.5323>

*Research supported by KBN grant 2 P03A 076 08.

I would like to mention that it was John Mallet-Paret who asked me about an “easy” way to establish uniqueness (up to multiplication) of globally positive solutions. I am grateful for his hospitality during my short stay at the Lefschetz Center for Dynamical Systems, Brown University.

0. PRELIMINARIES

Throughout the paper, for a Banach space X the symbol $\mathcal{L}(X)$ stands for the Banach space of bounded linear maps from X into itself, endowed with the uniform operator topology.

Let \mathcal{A} be the differential operator

$$(0.1) \quad \mathcal{A}u(x) := - \sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left(a_{ij}(x) \frac{\partial u}{\partial x_j} \right)$$

Denote by \tilde{A}_0 the linear operator $\tilde{A}_0 u := \mathcal{A}u$ acting on the Banach space of functions from $C^2(\overline{\Omega})$ satisfying the boundary condition (1b). For each $1 \leq p \leq \infty$ the operator \tilde{A}_0 is closable and densely defined in $L^p(\Omega)$. Let \tilde{A}_p denote the closure of \tilde{A}_0 in $L^p(\Omega)$. We write $\|\cdot\|_p$ for the norm on the Banach space $L^p(\Omega)$, $1 \leq p \leq \infty$.

It is well known (see e. g. Pazy [10]) that for each $1 \leq p < \infty$ the (unbounded) linear operator \tilde{A}_p is the infinitesimal generator of an analytic semigroup $\{e^{-\tilde{A}_p t}\}_{t \geq 0}$ of compact linear operators from $\mathcal{L}(L^p(\Omega))$. Put $A_p := \tilde{A}_p + \text{Id}$.

Lemma 0.1. *For each $1 \leq p < \infty$ the spectrum $\sigma(A_p)$ of A_p lies in the halfspace $\{\zeta \in \mathbb{C} : \text{Re } \zeta \geq 1\}$.*

Proof. As $-A_p$ is the infinitesimal generator of an analytic semigroup $e^{-A_p t}$ of compact linear operators, for each $t > 0$ the spectrum $\sigma(e^{-A_p t})$ consists of eigenvalues and $\{0\}$. By the standard regularity theory the solutions of the abstract parabolic equation $u_t + A_p u = 0$ are classical ones, hence we can consider $e^{-A_p \cdot 1} \in \mathcal{L}(C(\overline{\Omega}))$. The Kreĭn–Rutman theorem (see e. g. Hess [8]) implies that the spectral radius of $e^{-A_p \cdot 1}$ is a positive eigenvalue corresponding to a positive eigenfunction v_0 . Multiplying v_0 by a positive constant, if necessary, we can assume that $v_0(x) \leq 1$ for each $x \in \overline{\Omega}$.

For $x \in \overline{\Omega}$, $t \geq 0$, put $\tilde{v}(t, x) := e^{-t}$. The function \tilde{v} is a supersolution (for the definition see e. g. Hess [8] or Smith [14]) for the initial boundary value problem

$$\begin{aligned} u_t + \mathcal{A}u &= -u, \\ \frac{\partial u}{\partial \nu} + c(x)u &= 0, \\ u(0, x) &= 1 \text{ for } x \in \overline{\Omega}. \end{aligned}$$

An application of the parabolic maximum principle to \tilde{v} and to the solution $v(t, x)$ to the initial boundary value problem

$$\begin{aligned} v_t + \mathcal{A}v &= -v, \\ \frac{\partial v}{\partial \nu} + c(x)v &= 0, \\ v(0, x) &= v_0(x) \text{ for } x \in \overline{\Omega}. \end{aligned}$$

yields that $v(t, x)$ converges to zero as $t \rightarrow \infty$, uniformly in $x \in \overline{\Omega}$, with the exponential rate of decay not larger than -1 . So the spectral radius of $e^{-A_p \cdot 1} \in \mathcal{L}(C(\overline{\Omega}))$ (hence of $e^{-A_p \cdot 1} \in \mathcal{L}(L^p(\Omega))$) is not larger than e^{-1} . As the spectral mapping theorem (see e. g. [10]) asserts that $\sigma(e^{-A_p \cdot 1}) = \exp(\sigma(-A_p)) \cup \{0\}$, the desired result follows. \square

For $\alpha \geq 0$ denote by A_p^α the fractional power of A_p , and by $L^p(\Omega)^\alpha$ its domain endowed with the graph norm $\|\cdot\|_{p,\alpha} := \|A_p^\alpha \cdot\|_p$. Obviously, $A_p^1 = A_p$ and $L^p(\Omega)^1$ is the domain of A_p .

Theorem 0.2. *For each $1 \leq p < \infty$ the following holds:*

(a) *there is a positive constant $C = C(p)$ such that*

$$\|e^{-A_p t} u\|_p \leq C \|u\|_p \quad \text{for all } u \in L^p(\Omega) \text{ and } t \geq 0,$$

(b) *for each $u \in L^p(\Omega)$, $t > 0$ and $\alpha \geq 0$, $e^{-A_p t} u \in L^p(\Omega)^\alpha$,*

(c) *for each $0 \leq \alpha \leq 1$ there is a positive constant $C_1 = C_1(p, \alpha)$ such that*

$$\|e^{-A_p t} u\|_{p,\alpha} \leq \frac{C_1}{t^\alpha} \|u\|_p \quad \text{for all } u \in L^p(\Omega) \text{ and } t > 0,$$

(d) *for each $0 < \alpha \leq 1$ there is a positive constant $C_2 = C_2(p, \alpha)$ such that*

$$\|(\text{Id} - e^{-A_p t})u\|_p \leq C_2 t^\alpha \|u\|_{p,\alpha} \quad \text{for all } u \in L^p(\Omega)^\alpha \text{ and } t > 0.$$

Proof. All the facts stated above are standard (see e. g. Henry [7] or Pazy [10]) \square

For $t \in \mathbb{R}$ and $b \in L^\infty(\mathbb{R} \times \Omega)$, denote by $b \cdot t$ the translate of b , $(b \cdot t)(s, x) := b(t + s, x)$. Put

$$\mathbb{B} := w^* \text{-cl}\{a_0 \cdot t : t \in \mathbb{R}\},$$

where $w^* \text{-cl}$ stands for the closure in the weak* topology of $L^\infty(\mathbb{R} \times \Omega)$. Since \mathbb{B} is bounded, it is a compact metrizable space in the weak* topology of $L^\infty(\mathbb{R} \times \Omega)$.

For $b \in \mathbb{B}$ denote by $\mathbf{M}_p(b \cdot t)$ the multiplication operator by $b(t, \cdot)$:

$$(\mathbf{M}_p(b \cdot t)\phi)(x) := b(t, x)\phi(x), \quad \phi \in L^p(\Omega), x \in \Omega.$$

The following result is straightforward.

Lemma 0.3. *For each $b \in \mathbb{B}$, each $1 \leq p < \infty$ and each $\phi \in L^p(\Omega)$ we have $\|\mathbf{M}_p(b \cdot t)\phi\|_p \leq R \|\phi\|_p$, where R is the essential supremum of a_0 .*

1. LINEAR SKEW-PRODUCT (SEMI)DYNAMICAL SYSTEMS

We consider the following abstract parabolic equation

$$(1.1) \quad u_t + A_p u = \mathbf{M}_p(b \cdot t)u,$$

together with an initial condition

$$(1.2) \quad u(0) = u_0.$$

In the present section we investigate, for $1 \leq p < \infty$, the dependence of solutions to (1.1) satisfying (1.2) on the coefficient $b \in \mathbb{B}$ and the initial condition $u_0 \in L^p(\Omega)$.

The abstract initial value problem (1.1)+(1.2) can be written in the form of the following integral equation

$$(1.3) \quad u(t) = e^{-A_p t} u_0 + \int_0^t e^{-A_p(t-s)} \mathbf{M}_p(b \cdot s) u(s) ds.$$

We refer to solutions of (1.3), that is, functions $u \in C([0, \infty), L^p(\Omega))$ satisfying (1.3) for each $t \geq 0$ as *mild solutions* of (1.1)+(1.2). If a (mild) solution is defined for all $t \in \mathbb{R}$, we say it is a *global solution*.

Proposition 1.1. *For each $b \in L^\infty(\mathbb{R} \times \Omega)$, each $1 \leq p < \infty$ and each $u_0 \in L^p(\Omega)$ there exists precisely one solution $u(\cdot; u_0, b) : [0, \infty) \rightarrow L^p(\Omega)$ to (1.3).*

Proof. Applying the contraction mapping principle yields the desired solution. For a similar result, see Problem 7.1.5 in Henry's book [7]. \square

The proof of the following proposition rests on the estimates occurring in the proofs of Lemma 3.2 and Thm. 3.3 in [5], and we did not repeat it here.

Proposition 1.2. a) *For $T > 0$ and $1 \leq p < \infty$ the mapping $L^p(\Omega) \times \mathbb{B} \ni (u_0, b) \mapsto u(\cdot; u_0, b) \in C([0, T], L^p(\Omega))$ is continuous.*

b) *For $0 < t_1 \leq t_2$, $1 \leq p < \infty$ and $0 \leq \alpha < 1$ the mapping $L^p(\Omega) \times \mathbb{B} \ni (u_0, b) \mapsto u(\cdot; u_0, b) \in C([t_1, t_2], (L^p(\Omega))^\alpha)$ is continuous.*

Theorem 1.3. a) *For $t > 0$, $u_0 \in L^1(\Omega)$ and $b \in \mathbb{B}$ we have $u(t; u_0, b) \in C^1(\overline{\Omega})$.*

b) *For $0 < t_1 \leq t_2$ the mapping $L^1(\Omega) \times \mathbb{B} \ni (u_0, b) \mapsto u(\cdot; u_0, b) \in C([t_1, t_2], C^1(\overline{\Omega}))$ is continuous.*

Proof. By Amann [2] or Pazy [10], the domain of A_1 embeds continuously in the Sobolev space $W^{1,1}(\Omega)$. A corollary of the Nirenberg–Gagliardo inequality (see e. g. [7]) states that

$$L^1(\Omega)^{1/2} \subset L^q(\Omega) \quad \text{if } -n/q \leq 1/2 - n,$$

that is, $L^1(\Omega)^{1/2} \subset L^q(\Omega)$ for each $1 < q < 2n/(2n-1)$. From Proposition 1.2 we deduce that $u(t; u_0, b) \in L^q(\Omega)$ for each $t > 0$, each $u_0 \in L^1(\Omega)$ and each $b \in \mathbb{B}$.

By [10], for $1 < p < \infty$ the domain of A_p embeds continuously in the Sobolev space $W^{2,p}(\Omega)$. A corollary of the Nirenberg–Gagliardo inequality states that

$$L^p(\Omega)^{1/2} \subset L^q(\Omega) \quad \text{if } -n/q \leq 1 - n/p.$$

Therefore, $L^p(\Omega)^{1/2} \subset L^q(\Omega)$ for each $p < q < np/(n-p)$. Consequently, $L^p(\Omega)^{1/2} \subset L^r(\Omega)$, where $r := n/(n-1)$. Repeating the reasoning sufficiently many times, we prove that $u(t; u_0, b) \in L^p(\Omega)$ for each $t > 0$, each $u_0 \in L^1(\Omega)$ and each $b \in \mathbb{B}$ and $1 \leq p < \infty$.

For $p > 3n$ a corollary of the Nirenberg–Gagliardo inequality states that

$$L^p(\Omega)^{3/4} \subset C^\mu(\overline{\Omega}) \quad \text{if } 0 \leq \mu < 3/2 - 1/3 = 7/6.$$

It follows that $u(t; u_0, b) \in C^1(\overline{\Omega})$ for each $t > 0$, each $u_0 \in L^1(\Omega)$ and each $b \in \mathbb{B}$.

Part b) follows similarly by Proposition 1.2. \square

We recall now the definition of a skew-product dynamical system (with discrete time).

Let $B \times X$ be a product Banach bundle, where the base space B is a compact metrizable space and the fiber X is a Banach space. For a homeomorphism $\phi : B \rightarrow B$ the iterates ϕ^k , $k \in \mathbb{Z}$, form a (discrete time) dynamical system on B . A *compact linear skew-product dynamical system* $\{\Psi^k\}_{k=1}^\infty$ on $B \times X$, *covering* ϕ , is given by a family of compact linear operators $\{\psi(b) : b \in B\}$ depending continuously on $b \in B$ in the uniform operator topology, in the following way

$$\Psi(b, u) = (\phi(b), \psi(b)u), \quad b \in B, u \in X.$$

In other words, Ψ is a vector bundle endomorphism. The iterates Ψ^k , $k \in \mathbb{N}$, are given by

$$\Psi^k(b, u) = (\phi^k(b), \psi^{(k)}(b)u),$$

where we denote

$$\psi^{(k)}(b) := \psi(\phi^{k-1}(b)) \circ \psi(\phi^{k-2}(b)) \circ \dots \circ \psi(\phi(b)) \circ \psi(b).$$

(the *cocycle identity*).

Let X^* stand for the Banach space dual to X . For a compact linear skew-product dynamical system $\{\Psi^k\}_{k=1}^\infty$ on $B \times X$ we define its *dual system* $\{(\Psi^*)^k\}$ on $B \times X^*$ by

$$\Psi^*(b, u^*) := (\phi^{-1}(b), \psi^*(b)u^*), \quad b \in B, u^* \in X^*,$$

where $\psi^*(b) \in \mathcal{L}(X^*)$ is the dual operator to $\psi(b)$. It is straightforward that $\{(\Psi^*)^k\}_{k=1}^\infty$ is a compact linear skew-product dynamical system covering ϕ^{-1} .

Theorem 1.4. *Equation (1a)+(1b) generates a compact linear skew-product dynamical system $\{\Psi^k\}_{k=1}^\infty$ (resp. $\{\hat{\Psi}^k\}_{k=1}^\infty$) on the Banach bundle $\mathcal{B} := \mathbb{B} \times L^1(\Omega)$ (resp. $\hat{\mathcal{B}} := \mathbb{B} \times C^1(\overline{\Omega})$), where $\phi(b) := b \cdot 1$ for $b \in \mathbb{B}$, and*

$$\psi(b)u_0 := u(1; b, u_0) \quad \text{for } (b, u_0) \in \mathcal{B}.$$

$$\text{(resp. } \hat{\psi}(b)u_0 := u(1; b, u_0) \quad \text{for } (b, u_0) \in \hat{\mathcal{B}}.)$$

The system $\{\Psi^k\}_{k=1}^\infty$ factorizes through $\{\hat{\Psi}^k\}_{k=1}^\infty$ in the sense that $\psi(b)u \in C^1(\overline{\Omega})$ for each $(b, u) \in \mathcal{B}$ and

$$\hat{\psi}(b) = \psi(b) \circ i \quad \text{for each } b \in \mathbb{B},$$

where i stands for the natural embedding $C^1(\overline{\Omega}) \subset L^1(\Omega)$.

Proof. The continuity of the assignment $b \mapsto \psi(b)$ or $b \mapsto \hat{\psi}(b)$ as a mapping from \mathbb{B} into the space of bounded linear operators on $L^1(\Omega)$ or $C^1(\overline{\Omega})$ with the strong operator topology follows by Theorem 1.3. The compactness follows by Proposition 1.2 and by the fact that for p sufficiently large the space $L^p(\Omega)^{3/4}$ embeds continuously in $C^\mu(\overline{\Omega})$. We derive the continuity in the uniform operator topology along the lines of Thm. 2.3.2. in Pazy [10]. \square

An immediate consequence is that for the dual skew-product dynamical system, $\psi^*(b) : C^1(\overline{\Omega})^* \rightarrow L^\infty(\Omega)$ for each $b \in \mathbb{B}$. However, by the Green's formula it follows that $\psi^*(b)u^*$, $u^* \in C^1(\overline{\Omega})^*$, equals the value at time 1 of a solution of the adjoint equation. Since the adjoint equation satisfies all the assumptions, we have the following

Theorem 1.5. *The linear skew-product dynamical system $\{(\Psi^*)^k\}_{k=1}^\infty$ dual to the system $\{\Psi^k\}$ generated by equation (1a)+(1b) has the property that $\psi^*(b) : C^1(\overline{\Omega})^* \rightarrow C^1(\overline{\Omega})$ for each $b \in \mathbb{B}$.*

2. ORDER AND MONOTONICITY

For a Banach space X consisting of functions defined on Ω , by X_+ we denote the cone of nonnegative functions. It is straightforward that X_+ is a closed convex set such that

- a) For each $u \in X_+$ and $\alpha \geq 0$ one has $\alpha u \in X_+$, and
- b) X_+ contains no one-dimensional subspace.

The cones in $L^p(\Omega)$, $1 \leq p \leq \infty$, as well as in $C^1(\overline{\Omega})$ are *generating*, that is, $X_+ + X_+ = X$.

We write $v \leq u$ if $u - v \in X_+$, and $v < u$ if $v \leq u$ and $v \neq u$.

A cone X_+ is called *solid* if its interior X_{++} is nonempty. We write $v \ll u$ if $u - v \in X_{++}$. The standard cone in $L^p(\Omega)$ is solid if and only if $p = \infty$. The standard cone in $C^1(\overline{\Omega})$ is solid.

We say that a bounded linear functional $u^* \in X^*$ is *nonnegative* if $\langle u^*, u \rangle \geq 0$ for each $u \in X_+$, where $\langle \cdot, \cdot \rangle$ denotes the duality pairing. When the cone X_+ is generating, the set of all nonnegative functionals forms a cone $(X^*)_+$, called the *dual cone*.

A functional $u^* \in (X^*)_+$ is called *uniformly positive* if there is a constant $K > 0$ such that $\langle u^*, u \rangle \geq K\|u\|$ for each nonzero $u \in X_+$, where $\|\cdot\|$ stands for the norm in X . Among the spaces $L^p(\Omega)$, only $L^1(\Omega)$ admits uniformly positive functionals: They are represented by functions in $L^\infty(\Omega)$ whose values are a.e. larger than some $K > 0$ (in other words, elements of $L^\infty(\Omega)_{++}$). For more on cones the reader is referred to Amann's paper [1].

Theorem 2.1. *Let $\{\Psi^k\}_{k=1}^\infty$ be the linear skew-product dynamical system generated on \mathcal{B} by (1a)+(1b). There exists an invariant decomposition $\mathcal{B} = \mathcal{B}_1 \oplus \mathcal{B}_2$, such that*

- (i) $\text{codim } \mathcal{B}_1 = 1$, and for each $b \in \mathbb{B}$ the fiber $\mathcal{B}_1(b)$ is represented as the nullspace of a uniformly positive functional $v^*(b) \in L^1(\Omega)^*$.
- (ii) $\mathcal{B}_2 \setminus \mathcal{Z} \subset \mathbb{B} \times (L^1(\Omega)_+ \cup -L^1(\Omega)_+)$, where \mathcal{Z} denotes the null section of \mathcal{B} .
- (iii) The mapping $\Psi|_{\mathcal{B}_2}$ is a bundle automorphism.

(iv) There are constants $D \geq 1$ and $0 < \lambda < 1$ such that

$$\frac{\|\psi^{(k)}(b)v_1\|_{L^1(\Omega)}}{\|\psi^{(k)}(b)v_2\|_{L^1(\Omega)}} \leq D\lambda^k \frac{\|v_1\|_{L^1(\Omega)}}{\|v_2\|_{L^1(\Omega)}}$$

for each $(b, v_1) \in \mathcal{B}_1$, $(b, v_2) \in \mathcal{B}_2 \setminus \mathcal{Z}$, and each $k \in \mathbb{N}$.

Proof. By Theorem 1.3, for each $b \in \mathbb{B}$, each $u_0 \in L^1(\Omega)$ and each $t > 0$ we have $u(t; u_0, b) \in C^1(\overline{\Omega})$. Assume that a nonzero $u_0 \in L^1(\Omega)_+$. As $u(t; u_0, b)$ may fail to be a classical solution, we cannot apply the standard parabolic strong maximum principle as put forward e. g. in Protter and Weinberger [12]. However, since u as well as its spatial derivatives are continuous, we can establish the parabolic strong maximum principle by reasoning along the lines of Chapters 8 and 9 in Gilbarg and Trudinger [6], which enables us to show after the pattern of Thm. 4.1 in Hirsch [9] or Cor. 7.2.3 in Smith [14] that $u(t; u_0, b) \gg 0$ for $t > 0$.

Applying Thm. 1 in Poláčik and Tereščák [11] to the linear skew-product dynamical system $\{\hat{\Psi}^k\}_{k=1}^\infty$, we obtain the existence of two $\hat{\Psi}$ -invariant subbundles \mathcal{S} and \mathcal{T} of $\hat{\mathcal{B}}$ such that:

(A1) $\hat{\mathcal{B}} = \mathbb{B} \times C^1(\overline{\Omega})$ is the direct sum of \mathcal{S} and \mathcal{T} .

(A2) $\dim \mathcal{S} = 1$, and $\mathcal{S} \setminus \hat{\mathcal{Z}} \subset \mathbb{B} \times (C^1(\overline{\Omega})_{++} \cup -C^1(\overline{\Omega})_{++})$, where $\hat{\mathcal{Z}}$ stands for the null section of $\hat{\mathcal{B}}$.

(A3) For each $b \in \mathbb{B}$ the fiber $\mathcal{T}(b)$ can be (uniquely) represented by a normalized functional $\hat{v}^*(b) \in C^1(\overline{\Omega})_+^*$. In particular, $\mathcal{T} \cap (\mathbb{B} \times C^1(\overline{\Omega})_+) = \hat{\mathcal{Z}}$.

(A4) $\hat{\Psi}|_{\mathcal{S}}$ is a bundle automorphism.

(A5) There are constants $d \geq 1$ and $0 < \lambda < 1$ such that

$$\frac{\|\hat{\psi}^{(k)}(b)v_1\|_{C^1(\overline{\Omega})}}{\|\hat{\psi}^{(k)}(b)v_2\|_{C^1(\overline{\Omega})}} \leq d\lambda^k \frac{\|v_1\|_{C^1(\overline{\Omega})}}{\|v_2\|_{C^1(\overline{\Omega})}}$$

for each $(b, v_1) \in \mathcal{T}$, $(b, v_2) \in \mathcal{S}$, $v_2 \neq 0$ and each $k \in \mathbb{N}$.

Now, the desired bundle \mathcal{B}_2 equals simply \mathcal{S} considered a subbundle of the bundle $\mathcal{B} = \mathbb{B} \times L^1(\Omega)$.

In order to construct \mathcal{B}_1 , notice first that the mapping $\mathbb{B} \ni b \mapsto \hat{v}^*(b) \in C^1(\overline{\Omega})^*$ is continuous. Moreover, the $\hat{\Psi}$ -invariance of \mathcal{T} means that $\hat{\psi}^*(\phi^{-1}(b))\hat{v}^*(\phi^{-1}(b)) = \gamma\hat{v}^*(b)$ for some $\gamma = \gamma(b) > 0$. From Theorem 1.5 we derive that for each $b \in \mathbb{B}$ the functional $\hat{v}^*(b)$ is represented as a function from $C^1(\overline{\Omega})$ depending continuously on b . Denote by $v^*(b)$ the functional $\hat{v}^*(b)$ viewed as an element of $C^1(\overline{\Omega})$. Applying the parabolic strong maximum principle to the adjoint equation we get that $v^*(b) \in C^1(\overline{\Omega})_{++}$, consequently it is a uniformly positive functional from $L^1(\Omega)^*$.

To prove (iii) and (iv), use (A4), (A5) and the fact that since \mathcal{B}_1 has finite dimension, both the $C^1(\overline{\Omega})$ -norm and the $L^1(\Omega)$ -norm on it are equivalent. \square

The property described in (iv) is usually referred to as *exponential separation* (continuous separation in [11]). The direct sum decomposition $\mathcal{B} = \mathcal{B}_1 \oplus \mathcal{B}_2$ uniquely defines the bundle projection P with image \mathcal{B}_2 and kernel \mathcal{B}_1 . The exponential separation can be formulated in the following way:

There are constants $D \geq 1$ and $0 < \lambda < 1$ such that

$$\frac{\|(\text{Id} - P(\phi^k(b)))\psi^{(k)}(b)v\|_{L^1(\Omega)}}{\|P(\phi^k(b))\psi^{(k)}(b)v\|_{L^1(\Omega)}} \leq D\lambda^k \frac{\|(\text{Id} - P(b))v\|_{L^1(\Omega)}}{\|P(b)v\|_{L^1(\Omega)}}$$

for each $(b, v) \in \mathcal{B} \setminus \mathcal{B}_1$ and each $k \in \mathbb{N}$.

For $t \in \mathbb{R}$ and $b \in \mathbb{B}$ put $\phi_t b$ to be $b \cdot t$. The family $\{\phi_t\}_{t \in \mathbb{R}}$ forms a flow (= continuous-time dynamical system) on the compact metrizable space \mathbb{B} . Define for $t \geq 0$ and $b \in \mathbb{B}$ a linear operator $\psi(t, b) \in \mathcal{L}(L^1(\Omega))$ by the formula

$$\psi(t, b)u_0 := u(t; b, u_0) \quad \text{for } u_0 \in L^1(\Omega).$$

For each $t \geq 0$ the mapping Ψ_t defined as

$$\Psi_t(b, u) := (\phi_t b, \psi(t, b)u)$$

is a bundle endomorphism of \mathcal{B} . The family $\{\Psi_t\}_{t \geq 0}$ forms a *linear skew-product semiflow* on the product bundle \mathcal{B} . A consequence of the semiflow axioms is the following *cocycle identity*

$$(2.1) \quad \psi(t_1 + t_2, b) = \psi(t_1, \phi_{t_2} b) = \psi(t_2, \phi_{t_1} b) \quad \text{for } t_1 \geq 0, t_2 \geq 0.$$

For basic properties of linear skew-products semiflows the reader is referred to Sacker and Sell [13] or to Chow and Leiva [3].

Theorem 2.2. *Let $\{\Psi_t\}_{t \geq 0}$ be the linear skew-product semiflow generated on the product bundle $\mathcal{B} = \mathbb{B} \times L^1(\Omega)$ by Equation (1a)+(1b). Then*

- (i) *The subbundles \mathcal{B}_1 and \mathcal{B}_2 are Ψ_t -invariant in the sense that if $(b, u) \in \mathcal{B}_i$ then $(\phi_t b, \psi(t, b)u) \in \mathcal{B}_i$ for $t \geq 0$, where $i = 1, 2$.*
- (ii) *$\{\Psi_t|_{\mathcal{B}_2}\}$ extends uniquely to a linear skew-product flow on \mathcal{B}_2 .*
- (iii) *There are constants $D' \geq 1$ and $\mu > 0$ ($\mu = -\log \lambda$) such that*

$$\frac{\|\psi(t, b)v_1\|_{L^1(\Omega)}}{\|\psi(t, b)v_2\|_{L^1(\Omega)}} \leq D' e^{-\mu t} \frac{\|v_1\|_{L^1(\Omega)}}{\|v_2\|_{L^1(\Omega)}}$$

for each $(b, v_1) \in \mathcal{B}_1$, $(b, v_2) \in \mathcal{B}_2 \setminus \mathcal{Z}$, and each $t \geq 0$.

Proof. We start by proving (iii). Indeed, by the cocycle identity we have $\psi(t, b) = \psi(t - [t], \phi(t, b)) \circ \psi([t], b)$, where $[t]$ stands for the integer part of t . By the standard argument (compare e. g. the proof of Lemma 3.3 in Sacker and Sell [13]) there is a positive constant C' such that $\|\psi(s, b)u\|_{L^1(\Omega)} \leq C'\|u\|_{L^1(\Omega)}$ for $s \in [0, 1]$, $b \in \mathbb{B}$ and $u \in L^1(\Omega)$. Further, as \mathcal{B}_2 is one-dimensional, there is a positive constant C'' such that $\|\psi(s, b)u\|_{L^1(\Omega)} \geq C''\|u\|_{L^1(\Omega)}$ for $s \in [0, 1]$ and $(b, u) \in \mathcal{B}_2$. From this it follows that for $(b, v_1) \in \mathcal{B}_1$, $(b, v_2) \in \mathcal{B}_2 \setminus \mathcal{Z}$, $t \geq 0$, we have

$$\frac{\|\psi(t, b)v_1\|_{L^1(\Omega)}}{\|\psi(t, b)v_2\|_{L^1(\Omega)}} \leq \frac{C'\|\psi([t], b)v_1\|_{L^1(\Omega)}}{C''\|\psi([t], b)v_2\|_{L^1(\Omega)}} \leq \frac{C'D}{C''}\lambda^{[t]} \frac{\|v_1\|_{L^1(\Omega)}}{\|v_2\|_{L^1(\Omega)}} \leq \frac{C'D}{C''}\lambda^{t-1} \frac{\|v_1\|_{L^1(\Omega)}}{\|v_2\|_{L^1(\Omega)}}.$$

Putting $D' := C'C/\lambda C''$ and $\mu := -\log \lambda$ gives the desired result.

Part (ii) is a consequence of the fact that $\Psi|_{\mathcal{B}_2}$ is a bundle automorphism and the cocycle property.

The proof of the Ψ_t -invariance of \mathcal{B}_2 is straightforward: $(b, u) \in \mathcal{B}_2$ is equivalent to $(\text{Id} - P(b))u = 0$, which yields, by part (iii), $(\text{Id} - P(\phi_t b))\psi(t, b)u$ for all $t \geq 0$. In order to establish the Ψ_t -invariance of \mathcal{B}_1 , suppose by way of contradiction that for some $(b', v) \in \mathcal{B}_1$ and some $t' > 0$, $t' \notin \mathbb{N}$, we have $\psi(t', b')v \notin \mathcal{B}_1$. This means that $P(\phi_{t'} b')v \neq 0$. As $\psi([t'] + 1, b')v = \psi([t'] + 1 - t', \phi_{t'} b')v$, from (iii) we deduce that the ratio

$$\frac{\|(\text{Id} - P(\phi_{[t'] + 1} b'))\psi([t'] + 1, b')v\|_{L^1(\Omega)}}{\|P(\phi_{[t'] + 1} b')\psi([t'] + 1, b')v\|_{L^1(\Omega)}}$$

is finite, which contradicts the fact that $(\phi_{[t'] + 1} b', \psi([t'] + 1, b')v) \in \mathcal{B}_1$. \square

Theorem 2.3. *Let $u : \mathbb{R} \times \overline{\Omega} \rightarrow \mathbb{R}$ be a nonzero global solution to (1a)+(1b) such that $u(t, x) \geq 0$ for each $t \in \mathbb{R}$ and each $x \in \overline{\Omega}$. Then for each $t \in \mathbb{R}$ the pair $(a_0 \cdot t, u(t, \cdot))$ belongs to the one-dimensional subbundle \mathcal{B}_2 .*

Proof. An application of the parabolic strong maximum principle yields $u(t, \cdot) \in C^1(\overline{\Omega})_{++}$ for each $t \in \mathbb{R}$.

We claim that there is a positive constant L such that

$$(2.2) \quad \|P(b)w\|_{L^1(\Omega)} \geq L\|w\|_{L^1(\Omega)} \quad \text{for each } b \in \mathbb{B} \text{ and each } w \in L^1(\Omega)_+.$$

Indeed, as the fiber $\mathcal{B}_1(b)$ is the nullspace of the functional $\hat{v}^*(b)$, one has $\langle \hat{v}^*(b), w \rangle = \langle \hat{v}^*(b), P(b)w \rangle$ for all $w \in L^1(\Omega)$. Further, since \mathcal{B}_2 has dimension one and the functional $\hat{v}^*(b)$ is uniformly positive, there is a positive number $h(b)$ such that $\|w\|_{L^1(\Omega)} = h(b)\langle \hat{v}^*(b), w \rangle$ for all $w \in L^1(\Omega)_+$ with $(b, w) \in \mathcal{B}_2$. Consequently $\|P(b)w\|_{L^1(\Omega)} = h(b)\langle \hat{v}^*(b), P(b)w \rangle$ for all $w \in L^1(\Omega)_+$. The positive function $h(\cdot)$ is easily seen to be continuous, so there is $h > 0$, $h = \min\{h(b) : b \in \mathbb{B}\}$, such that $\|P(b)w\|_{L^1(\Omega)} \geq h\langle \hat{v}^*(b), w \rangle$ for all $b \in \mathbb{B}$ and $w \in \mathbb{B}$. Now it remains to notice that by the continuity of the mapping $\mathbb{B} \ni b \mapsto \hat{v}^*(b) \in L^1(\Omega)^*$ and the fact that uniformly positive functionals form an open set in $L^1(\Omega)^*$, the positive constant in the definition of uniform positivity can be chosen independent of b . Formula (2.2) follows immediately.

As a consequence, for each nonzero $w \in L^1(\Omega)_+$ and each $b \in \mathbb{B}$ one has

$$\frac{\|(\text{Id} - P(b))w\|_{L^1(\Omega)}}{\|P(b)w\|_{L^1(\Omega)}} \leq \frac{1 + N}{L},$$

where $N := \sup\{\|P(b)w\|_{L^1(\Omega)} : b \in \mathbb{B}, \|w\|_{L^1(\Omega)} = 1\}$.

For each $t \in \mathbb{R}$ denote $\tilde{u}(t) = u(t, \cdot)$ regarded as an element of $L^1(\Omega)$. Suppose to the contrary that there is t' such that $(a_0 \cdot t', \tilde{u}(t'))$ does not belong to \mathcal{B}_2 . Put

$$M := \frac{\|(\text{Id} - P(a_0 \cdot t'))\tilde{u}(t')\|_{L^1(\Omega)}}{\|P(a_0 \cdot t')\tilde{u}(t')\|_{L^1(\Omega)}} > 0,$$

and

$$t'' := t' + \frac{1}{\mu} \log \frac{ML}{2D'(1 + N)},$$

where μ and D' are constants from Theorem 2.2(iii). As $M \leq (1 + N)/L$, we have $t'' < t'$.

An application of Theorem 2.2(iii) yields

$$\begin{aligned}
& \frac{\|(\text{Id} - P(a_0 \cdot t'))\tilde{u}(t')\|_{L^1(\Omega)}}{\|P(a_0 \cdot t')\tilde{u}(t')\|_{L^1(\Omega)}} \\
&= \frac{\|\psi(t' - t'', a_0 \cdot t'')(\text{Id} - P(a_0 \cdot t''))\tilde{u}(t'')\|_{L^1(\Omega)}}{\|\psi(t' - t'', a_0 \cdot t'')P(a_0 \cdot t'')\tilde{u}(t'')\|_{L^1(\Omega)}} \\
&\leq D'e^{-\mu(t' - t'')} \frac{\|\psi(t' - t'', a_0 \cdot t'')(\text{Id} - P(a_0 \cdot t''))\tilde{u}(t'')\|_{L^1(\Omega)}}{\|\psi(t' - t'', a_0 \cdot t'')P(a_0 \cdot t'')\tilde{u}(t'')\|_{L^1(\Omega)}} \\
&\leq D'e^{-\mu(t' - t'')} \frac{1 + N}{L} \\
&\leq \frac{M}{2},
\end{aligned}$$

a contradiction. \square

Finally, we formulate now our main result.

Corollary 2.4. *Assume that u_1, u_2 are nonzero global solutions of (1a)+(1b) such that $u_1(t, x) \geq 0$ and $u_2(t, x) \geq 0$ for all $t \in \mathbb{R}$ and all $x \in \overline{\Omega}$. Then there is a positive constant κ such that $u_1(t, x) = \kappa u_2(t, x)$ for all $t \in \mathbb{R}$ and all $x \in \overline{\Omega}$.*

REFERENCES

1. H. Amann, *Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces*, SIAM Rev. **18** (1976), 620–709.
2. H. Amann, *Dual semigroups and second order linear elliptic boundary value problems*, Israel J. Math. **45** (1983), 225–254.
3. S.-N. Chow and H. Leiva, *Existence and roughness of the exponential dichotomy for skew-product semiflow in Banach spaces*, J. Differential Equations **120** (1995), 429–477.
4. S.-N. Chow, K. Lu and J. Mallet-Paret, *Floquet theory for parabolic differential equations*, J. Differential Equations **109** (1994), 147–200.
5. S.-N. Chow, K. Lu and J. Mallet-Paret, *Floquet bundles for scalar parabolic equations*, Arch. Rational Mech. Anal. **129** (1995), 245–304.
6. D. Gilbarg and N. W. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Second Edition, Springer, Berlin–Heidelberg–New York, 1983.
7. D. Henry, *Geometric Theory of Semilinear Parabolic Equations*, Lecture Notes in Math., **840**, Springer, Berlin–New York, 1981.
8. P. Hess, *Periodic-Parabolic Boundary Value Problems and Positivity*, Pitman Res. Notes Math. Ser., **247**, Longman Sci. Tech., Harlow, 1991.
9. M. W. Hirsch, *Stability and convergence in strongly monotone dynamical systems*, J. Reine Angew. Math. **383** (1988), 1–53.
10. A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Appl. Math. Sci., **44**, Springer, New York–Berlin–Heidelberg–Tokyo, 1983.
11. P. Poláčik and I. Tereščák, *Exponential separation and invariant bundles for maps in ordered Banach spaces with applications to parabolic equations*, J. Dynamics Differential Equations **5** (1993), 279–303.
12. M. H. Protter and H. F. Weinberger, *Maximum Principles in Differential Equations*, Prentice–Hall, Englewood Cliffs, NJ, 1967.
13. R. J. Sacker and G. R. Sell, *Dichotomies for linear evolutionary equations in Banach spaces*, J. Differential Equations **113** (1994), 17–67.
14. H. L. Smith, *Monotone Dynamical Systems. An Introduction to the Theory of Competitive and Cooperative Systems*, Math. Surveys Monogr., Vol. **41**, Amer. Math. Soc., Providence, R.I., 1995.

WYBRZEŻE WYSPIAŃSKIEGO 27

PL-50-370 WROCLAW

POLAND

E-mail address: `mierzyn@banach.im.pwr.wroc.pl`