

Parametric Summability and Its Applications

Jinlu Li, Robert Mendris

April 11, 2015

Summary: In this paper we study summability based on double sequences of complex constants as it is defined in “Linear Operators, General Theory” by N. Dunford and J. T. Schwartz. We define “power double sequences” or infinite “power matrices” as certain generalizations of double sequences and power series.

We relate the summability and boundedness of the power double sequences to the summability and boundedness of the double sequence A .

While others do investigate “power matrices” their definitions, as far as we were able to find, differ from our definitions. Using these definitions we extend some summability results for double sequences of constants to our power double sequences.

We investigate also other possible generalizations of double sequences and assess their usefulness in summability study.

1 Introduction

The only (most recent) related work we found is ...

There are different reasons for transforming one sequence into another and most of them are related to convergence. The practical need to improve convergence gave the impuls to study sequence transformations already in 17th century and resulted in the creation of summability theory at the end of 19th century. Before the invention of computers, mainly linear sequence transformations were studied. Approaches based on classical analysis culminated when [1] was published. After that modern approaches based on functional analysis appeared. For a comprehensive review of classical and modern methods in summability see [2]. From practical point of view, regular linear transformations are in general at most moderately powerful in improving convergence, and the popularity of most linear transformations has declined considerably in recent years. It seems, however, that the limiting factor is regularity not linearity. Recently also new powerful non-linear sequence transformations attracted research and applications. This is discussed in a nice historical review [3].

In this paper, we contribute to the classical summability methods of double sequences. As a review of these methods see [4].

Let $A = \{a_{ij}\}$, $i = 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex constants, that is,

$$A = \begin{pmatrix} a_{10} & a_{11} & a_{12} & \dots \\ a_{20} & a_{21} & a_{22} & \dots \\ a_{30} & a_{31} & a_{32} & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

Let Δ be the set of all double sequences of complex constants and c be the space of all convergent sequences of scalars.

Definition 1.1 Suppose that a double sequence $\{a_{ij}\}$ defines a linear transformation T of c onto itself by means of formula

$$T[s_0, s_1, \dots] = [t_0, t_1, \dots] = \left[\sum_{i=0}^{\infty} a_{0i} s_i, \sum_{i=0}^{\infty} a_{1i} s_i, \dots \right].$$

If T preserves limits of sequences (i.e. if $\lim_{i \rightarrow \infty} t_i = \lim_{i \rightarrow \infty} s_i$ for every $[s_i] \in c$), then the double sequence (or matrix) $A = \{a_{ij}\}$ is said to define a regular method of summability.

Lemma 1.2 $A(a)$ defines a bounded linear map of c into c , if and only if the following three conditions hold:

1. $\sup_{1 \leq i < \infty} \sum_{j=0}^{\infty} |f_{ij}(a)| = M < \infty$;
2. $\lim_{i \rightarrow \infty} f_{ij}(a)$ exists for $j = 1, 2, \dots$;
3. $\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} f_{ij}(a)$ exists.

Once $A(a)$ satisfies the above three conditions, for any column vector $s = [s_1, s_2, s_3, \dots] \in c$, the bounded linear map, $A(a)$, of c into c is defined by

$$A(a)(s)_i = f_{i0}(a) \lim_{i \rightarrow \infty} s_i + \sum_{j=1}^{\infty} f_{ij}(a) s_j, \text{ for } i = 1, 2, \dots,$$

where $A(a)(s) = [A(a)(s)_1, A(a)(s)_2, A(a)(s)_3, \dots]$.

Let B_c denote the space of all linear bounded maps of c into c . From Lemma 1, if $A(a) \in B_c$, then it has norm

$$|A(a)| = M$$

Theorem 1.3 Silverman-Toeplitz. $A(a)$ defines a regular method of summability, if and only if the following three conditions hold:

1. $\sup_{1 \leq i < \infty} \sum_{j=0}^{\infty} |f_{ij}(a)| = M < \infty$;
2. $\lim_{i \rightarrow \infty} f_{ij}(a) = 0$ for $j = 1, 2, \dots$;
3. $\lim_{i \rightarrow \infty} \sum_{j=1}^{\infty} f_{ij}(a) = 1$.

It is clear that if $A(a)$ defines a regular method of summability, then $A(a) \in B_c$.

One can find all the Definition, the Lemma 1. and Silverman-Toeplitz Theorem in [Dunford-Schwartz]. ("Linear Operators, General Theory" by N. Dunford and J. T. Schwartz.)

Let $A(z) = \{f_{ij}(z)\}$, $i = 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of functions with same domain D , which is a subset of complex numbers, that is,

$$A(z) = \begin{pmatrix} f_{10}(z) & f_{11}(z) & f_{12}(z) & \dots \\ f_{20}(z) & f_{21}(z) & f_{22}(z) & \dots \\ f_{30}(z) & f_{31}(z) & f_{32}(z) & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}.$$

It is clear that for any input $a \in D$, the output of $A(a) \in \Delta$.

2 Preliminary and Review

Power Matrices

Let $A = \{ a_{ij} \}$, $i = 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex constants.

The column power matrix induced by A is defined as

$$P_A^C(z) = \{ a_{ij} z^i \}, i = 1, 2, \dots, j = 0, 1, 2, \dots, \text{ that is,}$$

$$P_A^C(z) = \begin{pmatrix} a_{10}z & a_{11}z & a_{12}z & \dots \\ a_{20}z^2 & a_{21}z^2 & a_{22}z^2 & \dots \\ a_{30}z^3 & a_{31}z^3 & a_{32}z^3 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}.$$

The row power matrix induced by A is defined as

$$P_A^R(z) = \{ a_{ij} z^j \}, i = 1, 2, \dots, j = 0, 1, 2, \dots, \text{ that is,}$$

$$P_A^R(z) = \begin{pmatrix} a_{10} & a_{11}z & a_{12}z^2 & \dots \\ a_{20} & a_{21}z & a_{22}z^2 & \dots \\ a_{30} & a_{31}z & a_{32}z^2 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}.$$

The double power function matrix induced from A is defined as

$$P_A(z) = \{ a_{ij} z^{i+j} \}, i = 1, 2, \dots, j = 0, 1, 2, \dots, \text{ that is,}$$

$$P_A(z) = \begin{pmatrix} a_{10}z & a_{11}z^2 & a_{12}z^3 & \dots \\ a_{20}z^2 & a_{21}z^3 & a_{22}z^4 & \dots \\ a_{30}z^3 & a_{31}z^4 & a_{32}z^5 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}.$$

We will immediately generalize these definitions in the following section.

General Power Matrices

We will define power matrices of the first type now.

Definition. Let $A = \{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex constants. Let $g(z) = \sum_{i=0}^{\infty} g_i z^i$ be a complex power series. Denote its radius of convergence as R_g .

The *column power matrix* induced by A and associated with $g(z)$ is defined as

$$P_{A;g}^C(z) = \{a_{ij}g_i z^i\}, i = 0, 1, 2, \dots, j = 0, 1, 2, \dots, \text{ that is,}$$

$$P_{A;g}^C(z) = \begin{pmatrix} a_{00}g_0 & a_{01}g_0 & a_{02}g_0 & \dots \\ a_{10}g_1 z & a_{11}g_1 z & a_{12}g_1 z & \dots \\ a_{20}g_2 z^2 & a_{21}g_2 z^2 & a_{22}g_2 z^2 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}.$$

Definition. Let $A = \{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex constants. Let $h(z) = \sum_{j=0}^{\infty} h_j z^j$ be a complex power series. Denote its radius of convergence as R_h .

The *row power matrix* induced by A and associated with $h(z)$ is defined as

$$P_{A;h}^R(z) = \{a_{ij}h_j z^j\}, i = 0, 1, 2, \dots, j = 0, 1, 2, \dots, \text{ that is,}$$

$$P_{A;h}^R(z) = \begin{pmatrix} a_{00}h_0 & a_{01}h_1 z & a_{02}h_2 z^2 & \dots \\ a_{10}h_0 & a_{11}h_1 z & a_{12}h_2 z^2 & \dots \\ a_{20}h_0 & a_{21}h_1 z & a_{22}h_2 z^2 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

We will generalize the double power function matrix $P_A(z)$ now.

Definition. Let $A = \{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex constants. Let $g(z) = \sum_{i=0}^{\infty} g_i z^i$ be a complex power series. Denote its radius of convergence respectively as r_g . The *power double sequence of second type* induced by A and associated with $g(z)$ is defined as

$$P_{A;g}(z) = \{a_{ij}g_i z^{i+j}\}, i = 0, 1, 2, \dots, j = 0, 1, 2, \dots :$$

$$P_{A;g}(z) = \begin{pmatrix} a_{00}g_0 & a_{01}g_1 z & a_{02}g_2 z^2 & \dots \\ a_{10}g_1 z & a_{11}g_2 z^2 & a_{12}g_3 z^3 & \dots \\ a_{20}g_2 z^2 & a_{21}g_3 z^3 & a_{22}g_4 z^4 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

Definition. Let $A = \{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex constants. Let $g(z) = \sum_{i=0}^{\infty} g_i z^i$ and $h(z) = \sum_{j=0}^{\infty} h_j z^j$ be two complex power series. Denote their radius of convergence respectively as r_g and r_h . The *power double sequence of third type* induced by A and associated with $g(z)$ and $h(z)$ is defined as

$$P_{A;g,h}(z) = \{a_{ij}g_i h_j z^{i+j}\}, i = 0, 1, 2, \dots, j = 0, 1, 2, \dots :$$

$$P_{A;g,h}(z) = \begin{pmatrix} a_{00}g_0 h_0 & a_{01}g_0 h_1 z & a_{02}g_0 h_2 z^2 & \dots \\ a_{10}g_1 h_0 z & a_{11}g_1 h_1 z^2 & a_{12}g_1 h_2 z^3 & \dots \\ a_{20}g_2 h_0 z^2 & a_{21}g_2 h_1 z^3 & a_{22}g_2 h_2 z^4 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

Remark. More general definition would consider $\{g_i\}$ and $\{h_j\}$ to be two arbitrary number sequences.

3 Summability Results

For power double sequences of first type we have:

Proposition 1(C). Let the double sequence of complex constants $\{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a regular method of summability and $g(z) = \sum_{i=0}^{\infty} g_i z^i$ be a complex function with its power series. Then the following two conditions are equivalent for any complex number z :

(i) $\lim_{i \rightarrow \infty} g_i z^i = 1$.

(ii) The power double sequence of first type $\left\{ \left(P_{A;g}^C(z) \right)_{ij} \right\}$ is a regular method of summability.

Proof. (i) implies (ii) is a straightforward verification of the three conditions of Silverman-Toeplitz Theorem.

(ii) implies (i). The third condition of Silverman-Toeplitz Theorem for $P_{A;g}^C$ and for A gives (i).

Proposition 1(R). Let the double sequence of complex constants $\{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a regular method of summability and $h(z) = \sum_{j=0}^{\infty} h_j z^j$ be a complex function with its power series. Then the following two conditions are equivalent for any complex number z , which the sequence $\{h_j z^j\}$ is convergent for:

(i) $\lim_{j \rightarrow \infty} h_j z^j = 1$.

(ii) The power double sequence of first type $\left\{ \left(P_{A;h}^R(z) \right)_{ij} \right\}$ is a regular method of summability.

Proof. (i) implies (ii) is a straightforward verification of the three conditions of Silverman-Toeplitz Theorem.

(ii) implies (i). The third condition of Silverman-Toeplitz Theorem for $P_{A;h}^R$ and for A along with the convergence of the sequence $\{h_j z^j\}$ gives (i) (see the proof of (ii) implies (i) in the Proposition 1(II)).

For power double sequences of second type we have:

Theorem 1(II). Let double sequence of complex constants $\{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a regular method of summability and $g(z) = \sum_{i=0}^{\infty} g_i z^i$ be a complex function with its power series.

Then the following two conditions are equivalent for any complex number z , which the sequence $\{g_i z^i\}$ is convergent for:

(i) $\lim_{i \rightarrow \infty} g_i z^i = 1$.

(ii) The power double sequence of first type $\left\{ (P_{A;g}(z))_{ij} \right\}$ is a regular method of summability.

Proof. Let's show first that (ii) implies (i). From (ii) and the condition 3. of Silverman-Toeplitz theorem, we have

$\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_{i+j} z^{i+j} = 1$. Set $k = i + j$. This changes into $\lim_{i \rightarrow \infty} \sum_{k=i}^{\infty} a_{i,k-i} g_k z^k = 1$. Set $b_{ik} = a_{i,k-i}$ for $i \leq k$ and zero otherwise. Observe $\{b_{ik}\}$ is also a regular method of summability. Then by Silverman-Toeplitz theorem the limits of convergent sequences are preserved: $\lim_{k \rightarrow \infty} g_k z^k = \lim_{i \rightarrow \infty} \sum_{k=0}^{\infty} b_{i,k} g_k z^k$. But the right hand side is equal to $\lim_{i \rightarrow \infty} \sum_{k=i}^{\infty} a_{i,k-i} g_k z^k = 1$ and (i) immediately follows.

Now we show that (i) implies (ii). We will use Silverman-Toeplitz theorem again and we need to prove its three conditions:

$$1. \sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |(P_{A;g}(z))_{ij}| = \sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} g_{i+j} z^{i+j}| \text{ by the definition of } P_{A;g}.$$

Set $k = i + j$. Then the above supremum is

$$\sup_{0 \leq i < \infty} \sum_{k=i}^{\infty} |a_{i,k-i} g_k z^k| \leq \sup_{0 \leq i < \infty} \sum_{k=i}^{\infty} |a_{i,k-i}| \cdot \sup_{0 \leq k < \infty} |g_k z^k| < \infty$$

since the first supremum is finite by the first condition of $\{a_{ij}\}$ being a regular method of summability and the second one by the existence of the limit in (i).

$$2. \lim_{i \rightarrow \infty} (P_{A;g}(z))_{ij} = \lim_{i \rightarrow \infty} a_{ij} g_{i+j} z^{i+j} \text{ for } j = 0, 1, 2, \dots \text{ by the definition.}$$

Set $k = i + j$. Then the absolute value of the above limit is $\left| \lim_{k \rightarrow \infty} a_{k-j,j} g_k z^k \right| \leq \lim_{k \rightarrow \infty} |a_{k-j,j}| \cdot$

$\lim_{k \rightarrow \infty} |g_k z^k| = 0$ since the first limit is zero by the second condition of $\{a_{ij}\}$ being a regular method of summability and the second limit is 1 by (i).

$$3. \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} (P_{A;g}(z))_{ij} = \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_{i+j} z^{i+j} \text{ by the definition.}$$

Set $k = i + j$. Starting with (i) following the first part of this proof ((ii) implies (i)) backwards we have

$$1 = \lim_{k \rightarrow \infty} g_k z^k = \lim_{i \rightarrow \infty} \sum_{k=0}^{\infty} b_{i,k} g_k z^k = \lim_{i \rightarrow \infty} \sum_{k=i}^{\infty} a_{i,k-i} g_k z^k = \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_{i+j} z^{i+j}.$$

And this finishes the proof that $\{(P_{A;g}(z))_{ij}\}$ is a regular method of summability by Silverman-Toeplitz theorem.

Remarks. The condition (i) in Theorem A. implies $|z| = r_g$, where r_g is the radius of convergence. The requirement in the Theorem A. that the sequence $\{g_i z^i\}$ must be convergent seems to be too restrictive but the condition (ii) does not guarantee its convergence. There are examples of non-convergent sequences $\{g_i z^i\}$ (for both bounded and unbounded case) and regular methods of summability that map these sequences to convergent ones (= that sum them). Then by choosing $z = 1$ one gets a counterexample for each case.

Corollary. From the proof of the Theorem A. it is clear that for $|z| < r_g$, conditions 1. and 2. hold but the limit in 3. is zero and we don't get a regular method of summability in that case.

For power double sequences of third type we have:

Theorem 1(III). Let double sequence of complex constants $\{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a regular method of summability, and $g(z) = \sum_{i=0}^{\infty} g_i z^i$ and $h(z) = \sum_{j=0}^{\infty} h_j z^j$ be two complex power

series. If $\lim_{i \rightarrow \infty} g_i z^i$ and $\lim_{j \rightarrow \infty} h_j z^j$ exist then the following two conditions are equivalent for any such complex number z :

(i) $\lim_{i \rightarrow \infty} g_i z^i \cdot \lim_{j \rightarrow \infty} h_j z^j = 1.$

(ii) The power double sequence of second type $\left\{ (P_{A;g,h}(z))_{ij} \right\}$ is a regular method of summability.

Proof. The main structure of this proof is similar to the one of the Theorem A. Let's show first that (ii) implies (i). From (ii) and the condition 3. of Silverman-Toeplitz theorem, we have

$1 = \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_i h_j z^{i+j}.$ This equals to $\lim_{i \rightarrow \infty} g_i z^i \cdot \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} h_j z^j = \lim_{i \rightarrow \infty} g_i z^i \cdot \lim_{j \rightarrow \infty} h_j z^j,$ where we also used that $\{a_{ij}\}$ as a linear operator preserves limits by Silverman-Toeplitz theorem.

Now we show that (i) implies (ii). We will use again Silverman-Toeplitz theorem and need to prove its three conditions:

1. $\sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |(P_{A;g,h}(z))_{ij}| = \sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} g_i h_j z^{i+j}|$ by the definition of $P_{A;g,h}.$

The above supremum equals to

$$\sup_{0 \leq i < \infty} |g_i z^i| \cdot \sum_{j=0}^{\infty} |a_{ij} h_j z^j| \leq \sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |a_{i,j}| \cdot \sup_{0 \leq i < \infty} |g_i z^i| \cdot \sup_{0 \leq j < \infty} |h_j z^j| < \infty$$

since the first supremum is finite by the first condition of $\{a_{ij}\}$ being a regular method of summability and the other two are finite by the existence of the limit in (i).

2. $\lim_{i \rightarrow \infty} (P_{A;g,h}(z))_{ij} = \lim_{i \rightarrow \infty} a_{ij} g_i h_j z^{i+j}$ for $j = 0, 1, 2, \dots$ by the definition.

The absolute value of the above limit is $\left| h_j z^j \cdot \lim_{i \rightarrow \infty} a_{ij} g_i z^i \right| \leq \sup_{0 \leq i < \infty} |a_{i,j}| \cdot \left| h_j z^j \cdot \lim_{i \rightarrow \infty} g_i z^i \right| = 0$

for $j = 0, 1, 2, \dots$

since the first limit is zero by the second condition of $\{a_{ij}\}$ being a regular method of summability and the second limit is finite by (i).

3. $\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} (P_{A;g,h}(z))_{ij} = \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_i h_j z^{i+j}$ by the definition.

Starting with (i) following the proof of necessary condition backwards we have:

$$1 = \lim_{i \rightarrow \infty} g_i z^i \cdot \lim_{j \rightarrow \infty} h_j z^j = \lim_{i \rightarrow \infty} g_i z^i \cdot \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} h_j z^j = \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_i h_j z^{i+j}.$$

And this finishes the proof that $\left\{ (P_{A;g,h}(z))_{ij} \right\}$ is a regular method of summability by Silverman-Toeplitz theorem.

Remarks. (i) implies $|z| = r_g = r_h,$ where r_g and r_h are the radii of convergence.

Corollary. From the proof it is clear that for $|z| < r_g$ and $|z| < r_h,$ conditions 1. and 2. hold but the limit in 3. is zero and we don't get a regular method of summability in that case.

4 Boundedness Results

Assume $A \in B_c$ now, but not necessarily a regular method of summability. It is clear that $P_{A;h}^R(0) \in B_c$ and $P_{A;h}^R(1) \in B_c.$ Also $A = P_{A;g}^C(1) \in B_c.$ On the other hand, for a given z we can ask: Does

$P_{A;g}^C(z) \in B_c$ or $P_{A;h}^R(z) \in B_c$ hold? And because the conditions 1 and 2 clearly hold this is equivalent to: Does

$$\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_i z^i \text{ or } \lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} h_j z^j \text{ exist?}$$

The next propositions provide an answer for these two kinds of power matrices. For power double sequences of first type we have:

Proposition 2(C). Let $A = \{ a_{ij} \}$, $i = 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex scalars satisfying $A \in B_c$. If $P_{A;g}^C(a) \in B_c$, for some $a \neq 0$, then $P_{A;g}^C(z) \in B_c$, for all z satisfying $\|z\| < \|a\|$.

Proof. It is a straightforward verification of Silverman-Toeplitz Theorem conditions.

Proposition 2(R). Let $A = \{ a_{ij} \}$, $i = 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex scalars satisfying $A \in B_c$. If $P_{A;h}^R(a) \in B_c$, for some $a \neq 0$, then $P_{A;h}^R(z) \in B_c$, for all z satisfying $|z| < |a|$.

Proof. From the above argument, we have $P_A^R(0) \in B_c$. We only need to prove $P_A^R(z) \in B_c$, for all $|z| < |a|$ and $z \neq 0$.

From the hypothesis $P_A^R(a) \in B_c$, we have

1. $\sup_{1 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} a^j| = M < \infty$;
2. $\lim_{i \rightarrow \infty} a_{ij} a^j$ exists for $j = 1, 2, \dots$;
3. $\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} a^j$ exists.

We have to show

1. $\sup_{1 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} z^j| < \infty$;
2. $\lim_{i \rightarrow \infty} a_{ij} z^j$ exists for $j = 1, 2, \dots$;
3. $\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} z^j$ exists.

In fact, from the first condition for $P_A^R(a)$, we obtain

$$\sup_{1 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} z^j| = \sup_{1 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} a^j| \left| \frac{z}{a} \right|^j \leq \sup_{1 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} a^j| = M < \infty.$$

So $P_A^R(z)$ satisfies its first condition. Similarly, from the Condition 2 of $P_A^R(a)$, we have

$$\lim_{i \rightarrow \infty} a_{ij} z^j = \lim_{i \rightarrow \infty} a_{ij} a^j \left(\frac{z}{a} \right)^j = 0, \text{ for } j = 1, 2, \dots$$

So $P_A^R(z)$ satisfies its second condition. Next we show that $P_A^R(z)$ satisfies its condition 3 from Lemma 1.

For any given $\varepsilon > 0$, there exists N , such that $|\frac{z}{a}|^{N-1} < \frac{\varepsilon}{4M}$. From Conditions 1 and 2 above, there exists $K > 0$ such that for all $m, n > K$, the following inequality holds

$$|a_{mj}a^j - a_{nj}a^j| < \frac{\varepsilon}{2N}.$$

Now for all $m, n > K$, we have

$$\begin{aligned} & \left| \sum_{j=0}^{\infty} a_{mj}z^j - \sum_{j=0}^{\infty} a_{nj}z^j \right| \leq \\ & \leq \left| \sum_{j=0}^{N-1} a_{mj}z^j - \sum_{j=0}^{N-1} a_{nj}z^j \right| + \left| \sum_{j=N}^{\infty} a_{mj}z^j - \sum_{j=N}^{\infty} a_{nj}z^j \right| \\ & \leq \sum_{j=0}^{N-1} |a_{mj}a^j - a_{nj}a^j| \left| \frac{z}{a} \right|^j + \\ & \quad \left| \frac{z}{a} \right|^{N-1} \left(\sum_{j=N}^{\infty} |a_{mj}a^j| \left| \frac{z}{a} \right|^{j-N+1} + \sum_{j=N}^{\infty} |a_{nj}a^j| \left| \frac{z}{a} \right|^{j-N+1} \right) \\ & \leq \sum_{j=0}^{N-1} |a_{mj}a^j - a_{nj}a^j| + \left| \frac{z}{a} \right|^{N-1} \left(\sum_{j=N}^{\infty} |a_{mj}a^j| + \sum_{j=N}^{\infty} |a_{nj}a^j| \right) \\ & < \frac{N\varepsilon}{2N} + \frac{\varepsilon}{4M} (M + M) \\ & = \varepsilon. \end{aligned}$$

This proposition is proved.

Proposition 2(R) indicates that, the row power functions matrix $P_A^R(z)$ has a similar property to power series: If there exists a number $a \neq 0$, such that $P_A^R(a) \in B_c$, then there exists a positive number r_A such that, $P_A^R(z) \in B_c$, for all $|z| < r_A$, and $P_A^R(z) \notin B_c$, for all $|z| > r_A$. r_A is called the radius of summability of the matrix A . The radius of summability of the matrix A is 0, if there does not exist a number $a \neq 0$, such that $P_A^R(a) \in B_c$; The radius of summability of the matrix A is ∞ , if $P_A^R(a) \in B_c$ for all numbers a .

The following corollary follows immediately from Proposition 2(R) and the above notations.

Corollary 3. Let $A = \{ a_{ij} \}$, $i = 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex scalars. If $A \in B_c$ then $r_A \geq 1$.

For any given row power matrix $P_A^R(z)$, the entries of any fixed row, i , can be treated as the terms of a power series

$$\sum_{j=0}^{\infty} a_{ij}z^j$$

Its radius of convergence is denoted by r_A^i , for $i = 1, 2, \dots$.

Proposition 4. $r_A \leq \inf_{1 \leq i < \infty} r_A^i$.

Proof. For any given $|z| < r_A$, $P_A^R(z) \in B_c$, we have $\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} z^j$ exists. The series $\sum_{j=0}^{\infty} a_{ij} z^j$ is convergent, for $i = 1, 2, \dots$. It implies that $|z| \leq r_A^i$, for $i = 1, 2, \dots$. It completes the proof of this proposition.

For power double sequences of second type we have:

Proposition 2(II). Let $\{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex scalars and g a complex power series.

If $P_{A;g}(a) \in B_c$, for some $a \neq 0$, then $P_{A;g}(z) \in B_c$, for all z satisfying $|z| < |a|$.

Proof. It is clear that $P_{A;g}(0) \in B_c$ and we only need to prove $P_{A;g}(z) \in B_c$, for all $0 < |z| < |a|$.

From the hypothesis $P_{A;g}(a) \in B_c$, we have

- (a) $\sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} g_{i+j} a^{i+j}| = M < \infty$;
- (b) $\lim_{i \rightarrow \infty} a_{ij} g_{i+j} a^{i+j}$ exists for $j = 1, 2, \dots$;
- (c) $\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_{i+j} a^{i+j}$ exists.

We have to show that the above three conditions are also true for $P_{A;g}(z)$:

- (a) $\sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} g_{i+j} z^{i+j}| < \infty$;
- (b) $\lim_{i \rightarrow \infty} a_{ij} g_{i+j} z^{i+j}$ exists for $j = 0, 1, 2, \dots$;
- (c) $\lim_{i \rightarrow \infty} \sum_{j=0}^{\infty} a_{ij} g_{i+j} z^{i+j}$ exists.

In fact, from the condition (a) for $P_{A;g}(a)$, we obtain

$$\begin{aligned} \sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} g_{i+j} z^{i+j}| &= \sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} g_{i+j} a^{i+j}| \left| \frac{z}{a} \right|^{i+j} \\ &\leq \sup_{0 \leq i < \infty} \sum_{j=0}^{\infty} |a_{ij} g_{i+j} a^{i+j}| = M < \infty \end{aligned}$$

So $P_{A;g}(z)$ satisfies its first condition. Similarly, from the condition (b) of $P_{A;g}(a)$, we have

$$\lim_{i \rightarrow \infty} a_{ij} g_{i+j} z^{i+j} = \lim_{i \rightarrow \infty} a_{ij} g_{i+j} a^{i+j} \left(\frac{z}{a} \right)^{i+j} = 0, \text{ for } j = 0, 1, 2, \dots$$

So $P_{A;g}(z)$ satisfies its second condition. Next we show that $P_{A;g}(z)$ satisfies its condition (c).

For any given $\varepsilon > 0$, there exists N , such that $\left| \frac{z}{a} \right|^{N-1} < \frac{\varepsilon}{4M}$. From conditions (a) and (b) above, where we set $k = i + j$, there exists $K > 0$ such that $\left| \frac{z}{a} \right|^K < \frac{\varepsilon}{4MN}$ and for all $m, n > K$, the following inequalities hold (without loss of generality $m < n$)

$$\begin{aligned} \sum_{j=0}^{\infty} |a_{m-j,j} g_m a^m| &< M, \\ \sum_{j=0}^{\infty} |a_{n-j,j} g_n a^n| &< M, \\ |a_{m-j,j} g_m a^m - a_{n-j,j} g_n a^n| &< \frac{\varepsilon}{4N}. \end{aligned}$$

Now for all $m, n > K$, we have

$$\begin{aligned} & \left| \sum_{j=0}^{\infty} a_{m-j,j} g_m z^m - \sum_{j=0}^{\infty} a_{n-j,j} g_n z^n \right| \leq \\ & \leq \left| \sum_{j=0}^{N-1} a_{m-j,j} g_m z^m - \sum_{j=0}^{N-1} a_{n-j,j} g_n z^n \right| + \left| \sum_{j=N}^{\infty} a_{m-j,j} g_m z^m - \sum_{j=N}^{\infty} a_{n-j,j} g_n z^n \right| \\ & \leq \sum_{j=0}^{N-1} \left| a_{m-j,j} g_m a^m \left| \frac{z}{a} \right|^m - a_{n-j,j} g_n a^n \left| \frac{z}{a} \right|^n \right| + \\ & \quad \left| \frac{z}{a} \right|^{N-1} \left(\sum_{j=N}^{\infty} |a_{m-j,j} g_m a^m| \left| \frac{z}{a} \right|^{m-N+1} + \sum_{j=N}^{\infty} |a_{n-j,j} g_n a^n| \left| \frac{z}{a} \right|^{n-N+1} \right) \\ & \leq \sum_{j=0}^{N-1} \left(|a_{m-j,j} g_m a^m - a_{n-j,j} g_n a^n| \cdot \left| \frac{z}{a} \right|^m + |a_{n-j,j} g_n a^n| \cdot (1 - \left| \frac{z}{a} \right|^{n-m}) \cdot \left| \frac{z}{a} \right|^m \right) + \\ & \quad \left| \frac{z}{a} \right|^{N-1} \left(\sum_{j=N}^{\infty} |a_{m-j,j} g_m a^m| + \sum_{j=N}^{\infty} |a_{n-j,j} g_n a^n| \right) \\ & < \frac{N\varepsilon}{4N} + NM \frac{\varepsilon}{4MN} + \frac{\varepsilon}{4M} (M + M) \\ & = \varepsilon. \end{aligned}$$

This proposition is proved.

For power double sequences of third type we have:

Proposition 2(III). Let $\{a_{ij}\}$, $i = 0, 1, 2, \dots$, $j = 0, 1, 2, \dots$, be a double sequence of complex scalars and g, h complex power series. If $P_{A;g,h}(a) \in B_c$, for some $a \neq 0$, then $P_{A;g,h}(z) \in B_c$, for all z satisfying $|z| < |a|$.

Proof. This proof is very similar to the one of the Proposition 2(II) above and will be omitted.

Remarks. Propositions 2(II) and 2(III) might seem to generalize the Proposition 2(C) or Proposition 2(R) but they don't. We cannot write $P_A^R(z)$ nor $P_A^C(z)$ in the form of $P_{A;g}$ or $P_{A;g,h}$. But we can do so for $P_A(z)$ by choosing $g_i = 1$ or $g_i = h_i = 1$ respectively.

Obviously $P_{A;g}(0) \in B_c$, $A = P_{A;g}(1) \in B_c$, $P_{A;g,h}(0) \in B_c$, and $A = P_{A;g,h}(1) \in B_c$. Then for these power matrices $r_A \geq 1$.

5 Examples

For the following examples, one can check the conditions listed.

Example 1. Let $A = \{i + j\}$, $i = 1, 2, \dots, j = 0, 1, 2, \dots$:

- (a) $A \notin B_c$;
- (b) $r_A^i = 1$, for $i = 1, 2, \dots$;
- (c) $r_A = 1$.

Example 2. Let $A = \{ \frac{1}{i+j} \}$, $i = 1, 2, \dots, j = 0, 1, 2, \dots$:

- (a) $A \notin B_c$;
- (b) $r_A^i = 1$, for $i = 1, 2, \dots$;
- (c) $r_A = 1$.

Example 3. Let $A = \{ (i + j)! \}$, $i = 1, 2, \dots, j = 0, 1, 2, \dots$:

- (a) $A \notin B_c$;
- (b) $r_A^i = 0$, for $i = 1, 2, \dots$;
- (c) $r_A = 0$.

Example 4. Let $A = \{ \frac{1}{(i+j)!} \}$, $i = 1, 2, \dots, j = 0, 1, 2, \dots$:

- (a) $A \in B_c$, with $M = \sum_{j=0}^{\infty} \frac{1}{j!}$;
- (b) $r_A^i = \infty$, for $i = 1, 2, \dots$;
- (c) $r_A = \infty$.

Example 5. For a given $b > 0$, let $A = \{ b^{i+j} \}$, $i = 1, 2, \dots, j = 0, 1, 2, \dots$:

- (a) $A \in B_c$, if $b < 1$, with $M = \frac{1}{1-b}$, and $A \notin B_c$, if $b \geq 1$;
- (b) $r_A^i = b^{-1}$, for $i = 1, 2, \dots$;
- (c) $r_A = b^{-1}$.

Example 6. Let $[p_k]$ be a sequence of positive numbers satisfying $\sum_{k=1}^{\infty} p_k < \infty$, and let $P_i = \sum_{k=1}^i p_k$.

Define $A = a_{ij}$, $i = 1, 2, \dots, j = 0, 1, 2, \dots$, as follows

$$a_{ij} = \begin{cases} \frac{1}{P_i}, & j < i \\ 0, & j \geq i \end{cases}$$

- (a) $A \in B_c$, with $M = \frac{1}{p_1}$;
- (b) $r_A^i = \infty$, for $i = 1, 2, \dots$;
- (c) $r_A = 1$;

(d) A is not a regular method of summability.

Example 7. Let $[p_k]$ be a sequence of positive numbers satisfying $\sum_{k=1}^{\infty} p_k = \infty$, and let $P_i = \sum_{k=1}^i p_k$. Define $A = a_{ij}$, as in Example 6. Then:

- (a) $A \in B_c$, with $M = \frac{1}{p_1}$;
- (b) $r_A^i = \infty$, for $i = 1, 2, \dots$;
- (c) $r_A = \sup\{b > 0 : \lim_{i \rightarrow \infty} \frac{b^i}{P_i}$ exists $\} < \infty$;
- (d) A is a regular method of summability.

Proposition 6. Let $f(z)$ be an analytic function with power series expansion $\sum_{k=0}^{\infty} p_k z^k$ about point 0 with the radius of convergence r . Then, for any $s = [s_1, s_2, s_3, \dots] \in c$ and for any $|z| < r$, the sequence $[\sum_{k=0}^n p_k s_{k+1} z^k], n = 1, 2, \dots$, is also convergent.

Proof. Define $A = \{a_{ij}\}$, $i = 1, 2, 3, \dots, j = 0, 1, 2, \dots$, as follows:

$$a_{ij} = \begin{cases} p_j, & j < i \\ 0, & j \geq i \end{cases}$$

One can check that $A, A(z)$ have the following properties:

- (a) $A \in B_c$, with $M = \sum_{k=0}^{\infty} p_k$;
- (b) $r_A^i = \infty$, for $i = 1, 2, \dots$;
- (c) $r_A = r$;
- (d) $A(z) \in B_c$.

Then this proposition follows immediately from the property (d).

References

- [1] G.H. Hardy. *Divergent series*. Clarendon Press Oxford, 1949.
- [2] J. Boos. *Classical and Modern Methods in Summability*. Oxford Mathematical Monographs, Oxford University Press Oxford, 2000.
- [3] Ernst Joachim Weniger. *Nonlinear Sequence Transformations: Computational Tools for the Acceleration of Convergence and the Summation of Divergent Series*. eprint arXiv:math/0107080, 2001
- [4] V.G. Chelidze. *Summability Methods for Double Series and Double Integrals*. Tbilisi University Press Tbilisi, 1977.

Additional References:

- [5] A. Pringsheim. *Divergente Reihen. Encycl. math. Wiss. I A. 3*, 105111 (1898).
- [6] A. Peyerimhoff. *Lectures on Summability. Lecture Notes in Mathematics*, vol. 107, Springer-Verlag, Berlin 1969.
- [7] K. Zeller und W. Beekmann. *Theorie der Limitierungsverfahren (2. Aufl.)*. Springer, Berlin Heidelberg New York 1970.
- [8] K. Knopp. *Weierstrass Double Series Theorem ...*
- [9] *Theory of Functions*. Dover, New York 1996.
- [10] Review of [2] by Billy Rhoades:
<http://www.zentralblatt-math.org/zmath/en/search/?q=an:0954.40001&format=complete>
- [11] S.A. Baron. *Introduction to the theory of summability of series*. Tartu (1966) (In Russian)
- [12] C. Belen, M. Mursaleen and M. Yildirim. *Statistical A-summability of double sequences and a Korovkin type approximation theorem*. *Bull. Korean Math. Soc.* 49(4), 851861 (2012)
- [13] J. Boos. *Classical and Modern Methods in Summability*. Oxford University Press, New York, 2000
- [14] A.L. Brudno. *Summation of bounded sequences by matrices*. *Mat. Sb.* 16 (58) : 2 (1945) pp. 191247 (In Russian, English abstract.)
- [15] J.S. Connor. *The statistical and strong p -Cesro convergence of sequences*. *Anal. (Munich)* 8, 4763 (1988)
- [16] R.G. Cooke. *Infinite matrices and sequence spaces*. Macmillan (1950)
- [17] O.H.H. Edely, M. Mursaleen. *On statistical A-summability*. *Math. Comput. Model.* 49, 672680 (2009)
- [18] N. Etemadi. *Convergence of weighted averages of random variables revisited*. *Proceedings of the American Mathematical Society*, Volume 134, Number 9, 27392744 (2006)
- [19] H. Fast. *Sur la convergence statistique*. *Colloq. Math.* 2, 241244 (1951)
- [20] W. Feller. *An Introduction to Probability Theory and Its Applications, Volume II*. John Wiley & Sons Inc., 2nd Ed., New York, 1971
- [21] J.A. Fridy. *On statistical convergence*. *Anal. (Munich)* 5, 301313 (1985)
- [22] J.A. Fridy, H.I. Miller. *A matrix characterization of statistical convergence*. *Analysis* 11 (1991), pp. 5966
- [23] H. J. Hamilton. *Transformations of multiple sequences*. *Duke Math. J.* 2 (1936), no. 1, 2960.
- [24] G. H. Hardy. *Divergent Series*. Oxford Univ. Press, London, 1949.
- [25] G.F. Kangro. *Theory of summability of sequences and series*. *J. Soviet Math.* 5 : 1 (1970) pp. 145
- [26] K. Knopp. *Theorie und Anwendung der unendlichen Reihen*. Springer (1964) (English translation: Blackie, 1951 & Dover, reprint, 1990)

- [27] K. Knopp. *Theory and Application of Infinite Series*. Dover Publishing Company, 1990
- [28] E. Kolk. *Matrix summability of statistically convergent sequences*. *Anal. (Munich)* 13, 7783 (1993)
- [29] F. Móricz. *Tauberian conditions under which statistical convergence follows from statistical summability (C,I)*. *J. Math. Anal. Appl.* 275, 277287 (2002)
- [30] F. Móricz. *Statistical convergence of multiple sequences*. *Arch. Math. (Basel)* 81, 8289 (2003)
- [31] M. Mursaleen. *-Statistical convergence*. *Math. Slovaca* 50, 111115 (2000)
- [32] M. Mursaleen, O.H.H. Edely. *Statistical convergence of double sequences*. *J. Math. Anal. Appl.* 288, 223231 (2003)
- [33] T. Neubrunn, J. Smital, T. Salat. *On the structure of the space $M(0,1)$* . *Rev. Roumaine Math. Pures Appl.* 13 (1968), pp. 337386
- [34] A. Peyerimhoff. *Lectures on summability*. Springer (1969)
- [35] G. M. Robison. *Divergent double sequences and series*. *Trans. Amer. Math. Soc.* 28 (1926), no. 1, 5073.
- [36] T. Šalát. *On statistically convergent sequences of real numbers*. *Math. Slovaca* 30, 139150 (1980)
- [37] A. Zygmund. *On the convergence and summability of power series on the circle of convergence*. *Proc. London Math. Soc.* 47, 326350 (1941)

Jinlu Li
Department of Mathematics
Shawnee State University
940 Second Street
Portsmouth, OH 45662, USA
jli@shawnee.edu

Robert Mendris
Department of Mathematics
Shawnee State University
940 Second Street
Portsmouth, OH 45662, USA
rmendris@shawnee.edu