Komlós-Major-Tusnády approximations to increments of uniform empirical processes

Abdelhakim Necir*

Laboratory of Applied Mathematics, Mohamed Khider University, Biskra, Algeria

arXiv:1709.00747v4 [math.ST] 8 Dec 2017

Abstract

The well-known Komlós-Major-Tusnády inequalities [Z. Wahrsch. Verw. Gebiete 32 (1975) 111-131; Z. Wahrsch. Verw. Gebiete 34 (1976) 33-58] provide sharp inequalities to partial sums of iid standard exponential random variables by a sequence of standard Brownian motions. In this paper, we employ these results to establish Gaussian approximations to weighted increments of uniform empirical and quantile processes. This approach provides rates to the approximations which, among others, have direct applications to statistics of extreme values for randomly censored data.

Keywords: KMT approximation; empirical process; quantile process, random censoring.

AMS 2010 Subject Classification: 60F17, 62G15, 62G50, 62P05.

*Corresponding author: necirabdelhakim@yahoo.fr

1. Introduction

Csörgő *et al.* (1986) have constructed a probability space, denoted by $(\Omega, \mathcal{A}, \mathbb{P})$, carrying a sequence of independent random variables (rv's) $U_1, U_2, ...$ uniformly distributed on $(0,1)$ and a sequence of Brownian bridges $\{B_n(s); 0 \leq s \leq 1\}_{n\geq 1}$ such that for the empirical process

$$\alpha_n(s) := \sqrt{n}\,(G_n(s) - s),\ 0 \leq s \leq 1$$

and the quantile process

$$\beta_n(s) := \sqrt{n}\left(s - G_n^{-1}(s)\right),\ 0 \leq s \leq 1$$

where $G_n(s) := n^{-1}\sum_{i=1}^{n} \mathbf{1}\{U_i \leq s\}$ and

$$G_n^{-1}(s) := \inf\{t,\ G_n(s) \geq s\}, 0 \leq s \leq 1,$$

with $G_n^{-1}(0) := G_n^{-1}(0+)$, for universal positive constants a, b and c

$$\mathbb{P}\left\{\sup_{0\leq s\leq d/n} |\beta_n(s) - B_n(s)| \geq n^{-1/2}(a\log d + x)\right\} \leq be^{-cx}, \tag{1.1}$$

for all $0 \leq x \leq d^{1/2}$ and $1 \leq d \leq n$, with the same inequality holding for the supremum taken over $1 - d/n \leq s \leq 1$. Thereby, they showed that

$$\sup_{\lambda/n\leq s\leq 1-\lambda/n} \frac{n^{\eta}\,|\beta_n(s) - B_n(s)|}{[s(1-s)]^{1/2-\eta}} = O_{\mathbb{P}}(1), \tag{1.2}$$

as $n \to \infty$, for every fixed $0 < \lambda < \infty$ and $0 \leq \eta < 1/2$, leading to

$$\sup_{\lambda/n\leq s\leq 1-\lambda/n} \frac{n^{\nu}\,|\alpha_n(s) - B_n(s)|}{[s(1-s)]^{1/2-\nu}} = O_{\mathbb{P}}(1), \tag{1.3}$$

for every fixed $0 \leq \nu < 1/4$. The inequality (1.1) is a result of Theorem 1.1 will approximations (1.2) and (1.3) contain, respectively, in Theorem 2.1 and Corollary 2.1 of the above paper. Similar results may be found in Mason and van Zwet (1987). These two Gaussian approximations remain powerful tools to establish the asymptotic normality, among others, in statistics of extreme values, see, e.g., Csörgő *et al* (1985) and Peng (2001). In this paper, we are concerned with Gaussian approximations of the increments

$$\alpha_n(s;t) := \alpha_n(t) - \alpha_n(t-s),\ 0 \leq s < t < 1$$

and

$$\beta_n(s;t) := \beta_n(t) - \beta_n(t-s),\ 0 \leq s < t < 1.$$

Such processes are used, for example, in goodness of fit test statistics (see, e.g., Section 2 in Shorack and Wellner, 1982) and in nonparametric statistics for censored data (see, e.g., Deheuvels and Einmahl, 1996). For convenience, we next use the notation $f(s;t) := f(t) - f(t-s)$, $0 \leq s < t < 1$, for any measurable function f. Shorack and Wellner (1982) (Theorem 1.2) showed that there exist another Brownian bridge $\widetilde{B}(s)$, $0 \leq s \leq 1$ such that

$$\sup_{cn^{-1}\log n \leq s < t} \frac{\left|\alpha_n(s;t) - \widetilde{B}(s;t)\right|}{s^{\nu}} = o_{\mathbb{P}}(1), \tag{1.4}$$

for every $0 < t < 1$ and $c > 0$. By using Komlós-Major-Tusnády inequalities, Csörgő *et al.* (1986) (Theorem 4.6.1) also obtained a similar result and proved that, in the probability space $(\Omega, \mathcal{A}, \mathbb{P})$, we have

$$\sup_{cn^{-1}\log n \leq s < t} \frac{|\alpha_n(s;t) - B_n(s;t)|}{s^{\nu}} = o_{\mathbb{P}}(1), \tag{1.5}$$

where B_n is the same Brownian bridge as used in both approximations (1.2) and (1.3). The authors are noticed, in their Remark 4.6.1, that (1.5) is equivalent to

$$\sup_{cn^{-1}\log n \leq s < t} \frac{|\beta_n(s;t) - B_n(s;t)|}{s^{\nu}} = o_{\mathbb{P}}(1). \tag{1.6}$$

Otherwise, Alexander (1987) (Remark 2.7, Assertion 2.7) gave a refinement of (1.4) and (1.5) to prove that there exists another Brownian bridge $\widehat{B}(s)$, $0 \leq s \leq 1$ such that, for every $0 < \lambda < \infty$

$$\sup_{\lambda/n \leq s < t} \frac{\left|\alpha_n(s;t) - \widehat{B}(s;t)\right|}{s^{\nu}} = o_{\mathbb{P}}(1). \tag{1.7}$$

It is worth mentioning, that the three Brownian bridges B_n, $\widetilde{B}$ and $\widehat{B}$ are not necessarily the same. Note also that, for all large n, $cn^{-1}\log n > n^{-1}$, then approximation (1.7) is less restrictive and more useful than (1.5). But when we deal, for instance, to statistics of extreme values for randomly censored data (see, e.g., Brahimi *et al.*, 2015) the rate of this approximation is needed. This, to our knowledge, does not discussed yet in literature. In the following theorem we answer to this issue by providing a new Gaussian approximation it term of a sequence of Brownian bridges instead of their increments.

Theorem 1.1. *On the probability space* $(\Omega, \mathcal{A}, \mathbb{P})$, *carrying the sequence of iid rv's* $U_1, U_2, \ldots$ *uniformly distributed on* $(0,1)$ *and the sequence of Brownian bridges*

$B_1, B_2, ...$, *for every* $0 < \lambda < \infty$, $0 \leq \eta < 1/2$, *and* $0 \leq \nu < 1/4$, *we have approximations* (1.2) *and* (1.3), *together with*

$$\sup_{\lambda/n \leq s < t} \frac{n^{\eta} \left| \beta_n (s;t) - B_n (s) \right|}{s^{1/2-\eta}} = O_{\mathbb{P}} (1) \tag{1.8}$$

and

$$\sup_{\lambda/n \leq s < t} \frac{n^{\nu} \left| \alpha_n (s;t) - B_n (s) \right|}{s^{1/2-\nu}} = O_{\mathbb{P}} (1) . \tag{1.9}$$

2. **Application to statistics for censored data**

Let $X_1, ..., X_n$ be $n \geq 1$ independent copies of a non-negative continuous random variable (rv) X, defined over the probability space with cumulative distribution function (cdf) F. These rv's are censored to the right by a sequence of independent copies $Y_1, ..., Y_n$ of a non-negative continuous rv Y, independent of X and having a cdf G. At each stage $1 \leq j \leq n$, we only can observe the rv's $Z_j := \min (X_j, Y_j)$ and $\delta_j := \mathbf{1} \{X_j \leq Y_j\}$. If we denote by H the cdf of the observed $Z's$, then, in virtue of the independence of X and Y, we have $1 - H = (1 - F)(1 - G)$. We introduce two very crucial sub-distribution functions $H^{(i)} (z) := \mathbf{P} \{Z_1 \leq z, \delta_1 = i\}$, $i = 0, 1$, for $z > 0$, so that one have $H (z) = H^{(0)} (z) + H^{(1)} (z)$. The empirical counterparts are, respectively, defined by

$$H_n^{(0)} (z) := \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \{Z_i \leq z\} (1 - \delta_i), \ H_n^{(1)} (z) := \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \{Z_i \leq z\} \delta_i,$$

and therefore

$$H_n (z) := n^{-1} \sum_{i=1}^{n} \mathbf{1} \{Z_i \leq z\} = H_n^{(0)} (z) + H_n^{(1)} (z) .$$

Let

$$\xi_i := \delta_i H^{(1)} (Z_i) + (1 - \delta_i) \left(\theta + H^{(0)} (Z_i) \right), \ i = 1, ..., n,$$

be a sequence iid rv's uniformly distributed on $(0, 1)$ (Einmahl and Koning, 1992), and define the corresponding empirical cdf and empirical process by

$$\mathbb{U}_n (s) := \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \{\xi_i \leq s\} \text{ and } \alpha_n^* (s) := \sqrt{n} (\mathbb{U}_n (s) - s), \ 0 \leq s \leq 1,$$

respectively. Thereby we may represent, almost surely (a.s.), both $H_n^{(0)}$ and $H_n^{(1)}$ in term of $\mathbb{U}_n$, as follows $H_n^{(0)} (v) = \mathbb{U}_n \left(H^{(0)} (v) + \theta \right) - \mathbb{U}_n (\theta)$, for $0 < H^{(0)} (v) < 1 - \theta$, and $H_n^{(1)} (v) = \mathbb{U}_n \left(H^{(1)} (v) \right)$, for $0 < H^{(1)} (v) < \theta$. For further details, see for

instance Deheuvels and Einmahl (1996). From the previous representations, a.s., we may write

$$\sqrt{n}\left(\overline{H}_n^{(0)}(v)-\overline{H}^{(0)}(v)\right)=-\alpha_n^*\left(1-\overline{H}^{(0)}(v)\right),\ \text{for } 0<\overline{H}^{(0)}(v)<1-\theta$$

and

$$\sqrt{n}\left(\overline{H}_n^{(1)}(v)-\overline{H}^{(1)}(v)\right)=\alpha_n^*\left(\overline{H}^{(1)}(v);\theta\right),\ \text{for } 0<\overline{H}^{(1)}(v)<\theta.$$

By applying two approximations (1.3) and (1.9) , there exists a sequence of Brownian bridges $\{\mathcal{B}_n(s);\ 0\leq s\leq 1\}$ such that for every $0<\lambda<\infty$ and $0\leq\xi<1/4$,

$$\sup_{\lambda/n\leq\overline{H}^{(0)}(v)\leq 1}\frac{n^{\xi}\left|\alpha_n^*\left(1-\overline{H}^{(0)}(v)\right)-\mathcal{B}_n\left(1-\overline{H}^{(0)}(v)\right)\right|}{\left[\overline{H}^{(0)}(v)\right]^{1/2-\xi}}=O_{\mathbb{P}}(1)$$

and

$$\sup_{\lambda/n\leq\overline{H}^{(1)}(v)<\theta}\frac{n^{\xi}\left|\alpha_n^*\left(\overline{H}^{(1)}(v);\theta\right)-\mathcal{B}_n\left(\overline{H}^{(1)}(v)\right)\right|}{\left[\overline{H}^{(1)}(v)\right]^{1/2-\xi}}=O_{\mathbb{P}}(1).$$

These approximations will be useful tools for asymptotic results to statistics of extreme values for censored data, see for instance Brahimi *et al.* (2015).

3. **Proof of Theorem** 1.1

Let $Y_1^{(i)},Y_2^{(i)},...,\ (i=1,2)$, be two independent sequences of iid exponential rv's with mean 1. From Komlós *et al.* (1975) inequalities, there exist two independent copies $W^{(i)}(z)$, $0\leq z<\infty$, $(i=1,2)$, of standard Brownian motion defined on a probability space, such that for all real x, we have

$$\mathbb{P}\left\{\max_{1\leq k\leq m}\left|S_k^{(i)}-k-W^{(i)}(k)\right|\geq C\log m+x\right\}\leq Ke^{-\mu x},\tag{3.10}$$

for $m=1,2,...,$ where $S_k^{(i)}:=\sum_{j=1}^{k}Y_j^{(i)}$, with C, K and μ are positive universal constants independent of i and m. For each integer $n\geq 2$, we set

$$Y_j(n):=\begin{cases}Y_{[n/2]-j+1}^{(1)} & \text{for } j=1,...,[n/2],\\ Y_{n-j+2}^{(2)} & \text{for } j=[n/2]+1,...,n+1.\end{cases}$$

Then $Y_1(n),Y_2(n),...,Y_{n+1}(n)$ are iid sequence of exponential rv's with mean 1. For further use, we set $S_m(n):=\sum_{j=1}^{m}Y_j(n)$, $m=1,...,n+1$, and for the sake of notational simplicity, we will write from now on, S_m and Y_j instead of $S_m(n)$ and $Y_j(n)$, respectively, and will also use the usual convention $S_0=0$. It is easy to

verify that, for each integer $n \geq 2$, the following process is a sequence of standard Brownian motions on $[0, n+1]$:

$$W_n(s) := \begin{cases} W^{(1)}(s) & , \text{ for } 0 \leq s \leq \left[\frac{n}{2}\right], \\ W^{(1)}\left(\left[\frac{n}{2}\right]\right) + W^{(2)}\left(n+1-\left[\frac{n}{2}\right]\right) & \\ \qquad -W^{(2)}(n+1-s) & , \text{ for } \left[\frac{n}{2}\right] < s \leq n+1. \end{cases}$$

Let us define the following two processes

$$\widetilde{B}_n(s) := n^{-1/2}\left(sW_n(n) - W_n(sn)\right), \ 0 \leq s \leq 1$$

and

$$\widetilde{\beta}_n(s) := \sqrt{n}\left(s - \widetilde{U}_{[sn]:n}\right), \ 0 \leq s \leq 1,$$

where $\widetilde{U}_{k:n} := S_k/S_{n+1}$, for $k = 1, ..., n$, be a sequence of the uniform order statistics, with the convention $\widetilde{U}_{0:n} = S_0 \equiv 0$. We also define the uniform empirical process, corresponding to $\widetilde{U}_{1:n}, ..., \widetilde{U}_{n:n}$, by

$$\widetilde{\alpha}_n(s) := n^{1/2}\left(\widetilde{G}_n(s) - s\right), \ 0 \leq s \leq 1,$$

where $\widetilde{G}_n(s) := n^{-1}\sum_{i=1}^{n} \mathbf{1}\left\{\widetilde{U}_{i:n} \leq s\right\}$. In their inequalities (1.23) and (1.24), Csörgő *et al.* (1986), showed that

$$\mathbb{P}\left\{\sup_{0\leq s\leq d/n}\left|\widetilde{\beta}_n(s) - \widetilde{B}_n(s)\right| \geq 2n^{-1/2}\left(a\log d + x\right)\right\}$$

and

$$\mathbb{P}\left\{\sup_{1-d/n\leq s\leq 1}\left|\widetilde{\beta}_n(s) - \widetilde{B}_n(s)\right| \geq 2n^{-1/2}\left(a\log d + x\right)\right\},$$

whenever $n_0 < d < n$ and $0 \leq x \leq d^{1/2}$ for suitably chosen positive constants n_0, a, b and c. Thereby they stated that

$$\sup_{\lambda/n\leq s\leq 1-\lambda/n}\frac{n^{\eta}\left|\widetilde{\beta}_n(s) - \widetilde{B}_n(s)\right|}{\left[s(1-s)\right]^{1/2-\eta}} = O_{\mathbb{P}}(1) = \sup_{\lambda/n\leq s\leq 1-\lambda/n}\frac{n^{\nu}\left|\widetilde{\alpha}_n(s) - \widetilde{B}_n(s)\right|}{\left[s(1-s)\right]^{1/2-\nu}},$$

for every $0 < \lambda < \infty$, $0 \leq \eta < 1/2$, and $0 \leq \nu < 1/4$. Next we establish similar results to the increments

$$\widetilde{\beta}_n(s;t) = \widetilde{\beta}_n(t) - \widetilde{\beta}_n(t-s) = \sqrt{n}\left(s - \widetilde{U}_{[nt]:n} + \widetilde{U}_{[n(t-s)]:n}\right), \ 0 \leq s < t < 1$$

and

$$\widetilde{\alpha}_n(s;t) = \widetilde{\alpha}_n(t) - \widetilde{\alpha}_n(t-s) = \sqrt{n}\left(\widetilde{G}_n(s) - \widetilde{G}_n(t-s) - s\right), \ 0 \leq s < t < 1.$$

To this end, we will follow similar steps as used for the proof of Theorem 1.1 (inequality 1.1) in Csörgő *et al.* (1986). Let both d and n be sufficiently large and $n_0 > 1$, so that $n_0 < d < n$. For $a > 0$ and $0 \leq x \leq d^{1/2}$, we set

$$\mathbf{P}_n(x;d) := \mathbb{P}\left\{\sup_{0\leq s\leq d/n}\left|\widetilde{\beta}_n(s;t) - \widetilde{B}_n(s)\right| \geq 2n^{-1/2}(a\log d + x)\right\},$$

which is less than or equal to the sum of

$$\mathbf{P}_{1,n}(x;d) := \mathbb{P}\left\{\sup_{0\leq s\leq d/n}\left|\widehat{\beta}_n(s;t) - \widetilde{\beta}_n(s)\right| \geq n^{-1/2}(a\log d + x)\right\},$$

and

$$\mathbf{P}_{2,n}(x;d) := \mathbb{P}\left\{\sup_{0\leq s\leq d/n}\left|\widetilde{\beta}_n(s) - \widetilde{B}_n(s)\right| \geq n^{-1/2}(a\log d + x)\right\}.$$

Next we show that $\mathbf{P}_{1,n}(x;d) \xrightarrow{\mathbb{P}} 0$, as $n \to \infty$. Indeed, let us write

$$\left|\widehat{\beta}_n(s;t) - \widehat{\beta}_n(s)\right| = \sqrt{n}\left|\widetilde{U}_{[nt]:n} - \widetilde{U}_{[n(t-s)]:n} - \widetilde{U}_{[ns]:n}\right|,$$

which equals

$$\sqrt{n}\frac{\left|S_{[nt]} - S_{[n(t-s)]} - S_{[ns]}\right|}{S_{n+1}} = \sqrt{n}\frac{S_{|[nt]-[n(t-s)]-[ns]|}}{S_{n+1}},$$

thus

$$\mathbf{P}_{1,n}(x;d) = \mathbb{P}\left\{\frac{n}{S_{n+1}}\sup_{0\leq s\leq d/n} S_{|[nt]-[n(t-s)]-[ns]|} \geq a\log d + x\right\}.$$

Since $u \leq [u] \leq u+1$, then it is easy to check that

$$-2 \leq [nt] - [n(t-s)] - [ns] \leq 1,$$

this implies that $|[nt] - [n(t-s)] - [ns]| \leq 2$, it follows that for $0 \leq s < t < 1$, we have $S_{|[nt]-[n(t-s)]-[ns]|} \leq S_2$, therefore

$$\mathbf{P}_{1,n}(x;d) \leq \mathbb{P}\left\{\frac{n}{S_{n+1}}S_2 \geq a\log d + x\right\}.$$

By the law of large numbers, $\mathbb{P}\{|n/S_{n+1} - 1| \geq \epsilon\} \to 0$, for any fixed $0 < \epsilon < 1$, this implies that

$$\mathbf{P}_{1,n}(x;d) \leq \mathbb{P}\{(1-\epsilon)S_2 \geq a\log d + x\} + \mathbb{P}\{|n/S_{n+1} - 1| \geq \epsilon\}.$$

Note that S_2 is a sum of two iid standard exponential rv's, this means that it follows the Gamma cdf with two parameters $(2,1)$, that is $\mathbb{P}(S_2 > u) = (u+1)e^{-u}$, therefore

$$\mathbb{P}\{(1-\epsilon)S_2 \geq a\log d + x\} = \left(\frac{a\log d + x}{1-\epsilon} + 1\right)\exp\left(-\frac{a\log d + x}{1-\epsilon}\right),$$

which tends to zero as $d \to \infty$, hence $\mathbf{P}_{1,n}(x;d) \to 0$. On the other hand, from inequality (1.23) in Csörgő *et al.* (1986), we have $\mathbf{P}_{2,n}(x;d) \leq b\exp(-cx)$, thus $\mathbf{P}_{n}(x;d) \leq b\exp(-cx)$, too. Thereby, by using the latter inequality with similar arguments as used for the proof of Theorem 2.1 (statement 2.2) of the same paper, we end up with

$$\sup_{\lambda/n \leq s < t} \frac{n^{\eta}\left|\widetilde{\beta}_{n}(s;t) - \widetilde{B}_{n}(s)\right|}{s^{1/2-\eta}} = O_{\mathbb{P}}(1), \tag{3.11}$$

for every $0 \leq \eta < 1/2$ and $0 < \lambda < \infty$. Next we show that for every $0 < t < 1$ and $0 \leq \nu < 1/4$, we also have

$$A_{n,\nu}(t) := \sup_{\widetilde{U}_{1:n} \leq s < \widetilde{U}_{t_n:n}} \frac{n^{\nu}\left|\widetilde{\alpha}_{n}(s;t) - \widetilde{B}_{n}(s)\right|}{s^{1/2-\nu}} = O_{\mathbb{P}}(1), \tag{3.12}$$

where $t_n := [nt]$. Indeed, let us write

$$A_{n,\nu}(t) = \max_{1 \leq k \leq t_n - 1} \left\{ \sup_{\widetilde{U}_{k:n} \leq s < \widetilde{U}_{k+1:n}} \frac{n^{\nu}\left|\widetilde{\alpha}_{n}(s;t) - \widetilde{B}_{n}(s)\right|}{s^{1/2-\nu}} \right\},$$

and, for $0 < \tau \leq 1$, set

$$A_{n,\nu}(t;\tau) := \max_{1 \leq k \leq t_n - 1} \left\{ \sup_{\widetilde{U}_{k:n} \leq s < \widetilde{U}_{k+1:n}} \frac{n^{\nu}\left|\widetilde{\alpha}_{n}(s;t) - \widetilde{B}_{n}(s)\right|}{(\tau k/n)^{1/2-\nu}} \right\}.$$

Observe that

$$\left\{ \min_{1 \leq k \leq n} \widetilde{U}_{k:n}/k \geq \tau \right\} \subset \left\{ A_{n,\nu}(t) \leq \tau^{\nu-1/2} A_{n,\nu}(t;1) \right\},$$

and from assertion (2.9) in Csörgő *et al.* (1986), we have

$$\mathbb{P}\left\{ \min_{1 \leq k \leq n} \widetilde{U}_{k:n}/k \leq \tau \right\} = \tau,$$

it follows that

$$\mathbb{P}\left\{ A_{n,\nu}(t) \leq \tau^{\nu-1/2} A_{n,\nu}(t;1) \right\} \geq 1 - \tau.$$

Hence, to show that $A_{n,\nu}(t) = O_{\mathbb{P}}(1)$, it suffices to verify that $A_{n,\nu}(t;1) = O_{\mathbb{P}}(1)$ for sufficiently small τ. To this end, we will first state that for $1 \leq k \leq t_n - 1$ and $\widetilde{U}_{k:n} \leq s < \widetilde{U}_{k+1:n}$, we have

$$\widetilde{\alpha}_{n}(s;t) - \widetilde{\beta}_{n}\left(\frac{t_n}{n}, \frac{k}{n}\right) = O_{\mathbb{P}}\left(n^{-1/2}\right). \tag{3.13}$$

Indeed, let us fix $\epsilon > 0$ be small such that

$$\widetilde{U}_{t_n+1:n} < \widetilde{U}_{t_n+2:n} - \epsilon < \widetilde{U}_{t_n+2:n} + \epsilon < \widetilde{U}_{t_n+3:n},$$

and set

$$\mathcal{A}_{\epsilon,n}(t) := \left\{ \left| \widetilde{U}_{t_n+2:n} - t \right| < \epsilon \right\}.$$

Since $\widetilde{U}_{t_n+2:n} \xrightarrow{\mathbb{P}} t$ as $n \to \infty$ then $\mathbb{P}(\mathcal{A}_{\epsilon,n}(t)) \downarrow 1$, as $n \to \infty$. Hence, in the set $\mathcal{A}_{\epsilon,n}(t)$, we have $\widetilde{U}_{t_n+2:n} - \epsilon < t < \widetilde{U}_{t_n+2:n} + \epsilon$, which implies that

$$\widetilde{U}_{t_n:n} < t < \widetilde{U}_{t_n+3:n}. \tag{3.14}$$

Then, for $\widetilde{U}_{k:n} \leq s < \widetilde{U}_{k+1:n}$, we have

$$\widetilde{G}_n(t) - \widetilde{G}_n(t-s) - s \geq \widetilde{G}_n\left(\widetilde{U}_{t_n:n}\right) - \widetilde{G}_n\left(\widetilde{U}_{t_n+3:n} - \widetilde{U}_{k:n}\right) - \widetilde{U}_{k+1:n}.$$

Note that

$$\widetilde{U}_{t_n+3:n} - \widetilde{U}_{k:n} = \frac{S_{t_n+3} - S_k}{S_{n+1}} = \frac{S_{t_n-k+3}}{S_{n+1}} = \widetilde{U}_{t_n-k+3:n},$$

$$\widetilde{G}_n\left(\widetilde{U}_{t_n:n}\right) = t_n/n \text{ and } \widetilde{G}_n\left(\widetilde{U}_{t_n-k+3:n}\right) = \frac{t_n - k + 3}{n}.$$

Then the right-side of the previous inequality is equal to

$$\frac{k-3}{n} - \widetilde{U}_{k+1:n} = \frac{k}{n} - \widetilde{U}_{k:n} - \left(\widetilde{U}_{k+1:n} - \widetilde{U}_{k:n}\right) - \frac{3}{n}.$$

Note also $\widetilde{U}_{k+1:n} - \widetilde{U}_{k:n} = S_1/S_{n+1}$ and $\widetilde{U}_{k:n} = \widetilde{U}_{t_n:n} - \widetilde{U}_{t_n-k:n}$, it follows that

$$\widetilde{\alpha}_n(s;t) \geq \widetilde{\beta}_n\left(\frac{k}{n}, \frac{t_n}{n}\right) - \sqrt{n} S_1/S_{n+1} - 3/\sqrt{n}.$$

By using the law of large numbers, we have with large probability $n/S_{n+1} < 2$, then without loss of generality, we get

$$\widetilde{\alpha}_n(s;t) \geq \widetilde{\beta}_n\left(\frac{k}{n}, \frac{t_n}{n}\right) - 3(S_1+1)/\sqrt{n}. \tag{3.15}$$

Likewise, by using similar arguments as above we get

$$\widetilde{G}_n(t) - \widetilde{G}_n(t-s) - s \leq \widetilde{G}_n\left(\widetilde{U}_{t_n+3:n}\right) - \widetilde{G}_n\left(\widetilde{U}_{t_n:n} - \widetilde{U}_{k+1:n}\right) - \widetilde{U}_{k:n},$$

which implies that

$$\widetilde{\alpha}_n(s;t) \leq \widetilde{\beta}_n\left(\frac{k}{n}, \frac{t_n}{n}\right) + 4/\sqrt{n}. \tag{3.16}$$

By letting $\zeta := \max(3(S_1+1), 4)$, the inequalities (3.15) and (3.16) together give

$$\left| \widetilde{\alpha}_n(s;t) - \widetilde{\beta}_n\left(\frac{k}{n}, \frac{t_n}{n}\right) \right| < \zeta/\sqrt{n}.$$

Since $\zeta = O_{\mathbb{P}}(1)$, hence $\widetilde{\alpha}_n(s;t) - \widetilde{\beta}_n\left(\frac{k}{n}, \frac{t_n}{n}\right) = O_{\mathbb{P}}\left(n^{-1/2}\right)$ which meets (3.13). It is clear that $A_{n,\nu}(t;1)$ is less than or equal to the sum of

$$L_n := \max_{1\leq k\leq t_n-1}\left\{\sup_{\widetilde{U}_{k:n}\leq s<\widetilde{U}_{k+1:n}} \frac{n^{\nu}\left|\widetilde{\alpha}_n(s;t) - \widetilde{\beta}_n\left(\frac{k}{n}, \frac{t_n}{n}\right)\right|}{(k/n)^{1/2-\nu}}\right\}$$

and

$$T_n := \max_{1\leq k\leq t_n-1}\left\{\sup_{\widetilde{U}_{k:n}\leq s<\widetilde{U}_{k+1:n}} \frac{n^{\nu}\left|\widetilde{\beta}_n\left(\frac{k}{n}, \frac{t_n}{n}\right) - \widetilde{B}_n(s)\right|}{(k/n)^{1/2-\nu}}\right\}.$$

Making use of (3.13), we infer that $L_n = O_{\mathbb{P}}(1)$. Observe now that T_n is less than or equal to the sum of

$$T_{n1} := \max_{1\leq k\leq t_n-1}\left\{\frac{n^{\nu}\left|\widetilde{\beta}_n\left(\frac{k}{n}, \frac{t_n}{n}\right) - \widetilde{B}_n\left(\frac{k}{n}\right)\right|}{(k/n)^{1/2-\nu}}\right\}$$

and

$$T_{n2} := \max_{1\leq k\leq t_n-1}\left\{\sup_{\widetilde{U}_{k:n}\leq s<\widetilde{U}_{k+1:n}} \frac{n^{\nu}\left|\widetilde{B}_n(s) - \widetilde{B}_n\left(\frac{k}{n}\right)\right|}{(k/n)^{1/2-\nu}}\right\}.$$

By letting $k/n = s$ and $t_n/n = t^*$, we may write

$$T_{n1} \leq \sup_{1/n\leq s<t^*} \frac{n^{\nu}\left|\widetilde{\beta}_n(s,t^*) - \widetilde{B}_n(s)\right|}{s^{1/2-\nu}},$$

which, by (3.11), is equal to $O_{\mathbb{P}}(1)$. Let us now show that $T_{n2} = O_{\mathbb{P}}(1)$ too. To this end, we will follow similar procedures are those used for the proof of assertion (2.21) in Csörgő *et al.* (1986). Let us choose $0 < \nu < 1/4$ and set $\delta := (1/4 - \nu)/2$. For any $1 \leq k \leq n-1$ and $b \geq 1$, let $c_{k,n}^{(\delta)} := k^{2\delta+1/2}/n$ and

$$I_{k,n}(b) := \left[k/n - 3bc_{k,n}^{(\delta)}, k/n - 3bc_{k,n}^{(\delta)}\right]$$

and

$$D_{n,v}(b) := \max_{1\leq k\leq t_n-1}\left\{\sup_{s\in I_{k,n}(b)} \frac{n^{\nu}\left|\widetilde{B}_n(k/n) - \widetilde{B}_n(s)\right|}{(k/n)^{1/2-\nu}}\right\}.$$

Assertion (2.25) in Csörgő *et al.* (1986) states that

$$\lim_{b\to\infty} \mathbb{P}\left\{T_{n2} \geq D_{n,v}(b)\right\} = 0. \tag{3.17}$$

Then we have to show that $D_{n,v}(b) = O_{\mathbb{P}}(1)$. Let us write

$$D_{n,v}(b) = \max_{1\leq k\leq t_n-1}\left\{\sup_{\mathbf{s}\in[\mathbf{a}_k-h,\mathbf{a}_k+h]\cap[0,1]}\frac{n^{\nu}\left|\widetilde{B}_n(\mathbf{a}_k)-\widetilde{B}_n(\mathbf{s})\right|}{(k/n)^{1/2-\nu}}\right\},$$

where $\mathbf{a}_k = t-s$, $\mathbf{s} = t^*-k/n$ and $h = |\mathbf{a}_k-\mathbf{s}|$. Since $|t-t^*|\leq n^{-1}\leq bc_{k,n}^{(\delta)}$ and $|s-k/n|\leq 3bc_{k,n}^{(\delta)}$, then $h\leq |t-t^*|+|s-k/n| = 4bc_{k,n}^{(\delta)} =: h_*$. Let us write

$$D_{n,v}^*(b) := \max_{1\leq k\leq t_n-1}\left\{\sup_{\mathbf{s}\in[\mathbf{a}_k-h_*,\mathbf{a}_k+h_*]\cap[0,1]}\frac{n^{\nu}\left|\widetilde{B}_n(\mathbf{a}_k)-\widetilde{B}_n(\mathbf{s})\right|}{(k/n)^{1/2-\nu}}\right\}.$$

It is clear that $D_{n,v}^*(b)\leq D_{n,v}(b)$. Hence, from (3.17), it remain to show that $D_{n,v}^*(b) = O_{\mathbb{P}}(1)$. Indeed, for $d>0$ arbitrarily chosen, we have

$$\begin{aligned}&\mathbb{P}\left\{D_{n,v}^*(b)\geq d(4b)^{1/2}\right\}\\ &\leq\sum_{k=1}^{t_n-1}\mathbb{P}\left\{\sup_{\mathbf{s}\in[\mathbf{a}_k-h^*,\mathbf{a}_k+h^*]\cap[0,1]}\left|\widetilde{B}_n(\mathbf{a}_k)-\widetilde{B}_n(\mathbf{s}_k)\right|\geq d(4b)^{1/2}k^{1/2-\nu}n^{-1/2}\right\}\end{aligned}$$

which may be rewritten into

$$\sum_{k=1}^{t_n-1}\mathbb{P}\left\{\sup_{\mathbf{s}\in[\mathbf{a}_k-h^*,\mathbf{a}_k+h^*]\cap[0,1]}\left|\widetilde{B}_n(\mathbf{a}_k)-\widetilde{B}_n(\mathbf{s}_k)\right|\geq dk^{1/4-\nu-\delta}h_*^{1/2}\right\}. \tag{3.18}$$

From inequality (1.11) in Csörgő *et al.* (1986), for a given Brownian bridge $\mathbf{B}(s)$, $0\leq s\leq 1$ defined on $(\Omega,\mathcal{A},\mathbb{P})$, we have

$$\mathbb{P}\left\{\sup_{\mathbf{s}\in[\mathbf{a}-h,\mathbf{a}+h]\cap[0,1]}|\mathbf{B}(\mathbf{a})-\mathbf{B}(\mathbf{s})|\geq uh^{1/2}\right\}\leq Au^{-1}\exp\left(-u^2/8\right),$$

for any $0<\mathbf{a}<1$, $h>0$ and $0<u<\infty$, with a suitably chosen universal constant A. By applying this inequality we infer that (3.18) is less than or equal to

$$\sum_{k=1}^{\infty}\frac{d^{-1}\exp\left(-d^2k^{1/2-2\nu-2\delta}/8\right)}{k^{1/4-\nu-\delta}} =: \mathbf{P}(d).$$

Note that the series $\mathbf{P}(d)$ is uniformly convergent on $d\geq 1$, and $\lim_{d\to\infty}\mathbf{P}(d)=0$, it follows that $D_{n,v}^*(b) = O_{\mathbb{P}}(1)$, as sought. Let us now show that

$$\sup_{\lambda/n\leq s<t}\frac{n^{\nu}\left|\widetilde{\alpha}_n(s;t)-\widetilde{B}_n(s)\right|}{s^{1/2-\nu}} = O_{\mathbb{P}}(1).$$

Indeed, let us write

$$\sup_{\lambda/n\leq s<t}\frac{n^{\nu}\left|\widetilde{\alpha}_n(s;t)-\widetilde{B}_n(s)\right|}{s^{1/2-\nu}}\leq A_{n,\nu}(t)+B_{n,\nu}(t)+C_{n,\nu}(t)+D_{n,\nu}(t),$$

where $A_{n,\nu}(t)$ is that of (3.12),

$$B_{n,\nu}(t) := \sup_{(\lambda/n)\wedge\widetilde{U}_{1:n}\leq s<\widetilde{U}_{1:n}} \frac{n^{\nu}\left|\widetilde{\alpha}_n(s;t)\right|}{s^{1/2-\nu}},$$

$$C_{n,\nu}(t) := \sup_{\widetilde{U}_{t_n:n}\leq s<\widetilde{U}_{t_n:n}\vee t} \frac{n^{\nu}\left|\widetilde{\alpha}_n(s;t)-\widetilde{B}_n(s)\right|}{s^{1/2-\nu}}$$

and

$$D_{n,\nu}(t) := \sup_{(\lambda/n)\wedge\widetilde{U}_{1:n}\leq s<\widetilde{U}_{1:n}} \frac{n^{\nu}\left|\widetilde{B}_n(s)\right|}{s^{1/2-\nu}}.$$

It is clear that $B_{n,\nu}(t)$ is less than or equal to

$$\sup_{(\lambda/n)\wedge\widetilde{U}_{1:n}\leq s<\widetilde{U}_{1:n}} \frac{n^{\nu+1/2}\left|G_n(t)-G_n(t-s)\right|}{s^{1/2-\nu}} + n^{\nu+1/2}\sup_{(\lambda/n)\wedge\widetilde{U}_{1:n}\leq s<\widetilde{U}_{1:n}} s^{1/2+\nu}.$$

Recall (3.14) and observe that for $(\lambda/n)\wedge\widetilde{U}_{1:n}\leq s<\widetilde{U}_{1:n}$ (sufficiently small), we have

$$G_n(t)-G_n(t-s)\leq G_n\left(\widetilde{U}_{t_n+3:n}\right)-G_n\left(\widetilde{U}_{t_n:n}-\widetilde{U}_{1:n}\right)=\frac{t_n+3}{n}-\frac{t_n+1}{n}=\frac{2}{n}$$

and

$$G_n(t)-G_n(t-s)\geq G_n\left(\widetilde{U}_{t_n:n}\right)-G_n\left(\widetilde{U}_{t_n+3:n}\right)=\frac{t_n}{n}-\frac{t_n+3}{n}=-\frac{3}{n}.$$

It follows that $\left|G_n(t)-G_n(t-s)\right|\leq 3/n$, then it is easy to verify that

$$\sup_{(\lambda/n)\wedge\widetilde{U}_{1:n}\leq s<\widetilde{U}_{1:n}} \frac{n^{\nu+1/2}\left|G_n(t)-G_n(t-s)\right|}{s^{1/2-\nu}} = O_{\mathbb{P}}(1),$$

and since $n\widetilde{U}_{1:n}\xrightarrow{\mathbb{P}}1$, we get

$$n^{\nu+1/2}\sup_{(\lambda/n)\wedge\widetilde{U}_{1:n}\leq s<\widetilde{U}_{1:n}} s^{1/2+\nu}=\left(n\widetilde{U}_{1:n}\right)^{1/2+\nu}=O_{\mathbb{P}}(1),$$

therefore $B_{n,\nu}(t)=o_{\mathbb{P}}(1)$. For the second term, we write

$$C_{n,\nu}(t)\leq\left(\widetilde{U}_{t_n:n}\right)^{1/2-\nu}\sup_{\widetilde{U}_{1:n}\leq s<\widetilde{U}_{n:n}} n^{\nu}\left|\widetilde{\alpha}_n(s;t)-\widetilde{B}_n(s)\right|.$$

In view of assertion (2.6) of Theorem 2.2 in Csörgő *et al.* (1986), we have

$$\sup_{0\leq s\leq 1}\left|\widetilde{\alpha}_n(s;t)-\widetilde{B}_n(s)\right|=O\left(\frac{(\log n)^{1/2}(\log\log n)^{1/4}}{n^{1/4}}\right),\ \text{almost surely,}$$

it follows, since $0\leq\nu<1/4$, that

$$\sup_{\widetilde{U}_{1:n}\leq s<\widetilde{U}_{n:n}} n^{\nu}\left|\widetilde{\alpha}_n(s;t)-\widetilde{B}_n(s)\right|=o_{\mathbb{P}}(1).$$

On the other hand $\widetilde{U}_{t_n:n} \xrightarrow{\mathbb{P}} t$, it follows that $C_{n,\nu}(t) = o_{\mathbb{P}}(1)$. We have $n\widetilde{U}_{1:n} \xrightarrow{\mathbb{P}} 1$, then it is easy to show that $D_{n,\nu}(t) = O_{\mathbb{P}}(1)$, that we omit further details. To summarize, we briefly stated that

$$\sup_{\lambda/n \le s \le 1-\lambda/n} \frac{n^{\eta} \left| \widetilde{\beta}_n(s) - \widetilde{B}_n(s) \right|}{[s(1-s)]^{1/2-\eta}} = O_{\mathbb{P}}(1) = \sup_{\lambda/n \le s \le 1-\lambda/n} \frac{n^{\nu} \left| \widetilde{\alpha}_n(s) - \widetilde{B}_n(s) \right|}{[s(1-s)]^{1/2-\nu}}. \tag{3.19}$$

and showed that

$$\sup_{\lambda/n \le s < t} \frac{n^{\eta} \left| \widetilde{\beta}_n(s;t) - \widetilde{B}_n(s) \right|}{s^{1/2-\eta}} = O_{\mathbb{P}}(1) = \sup_{\lambda/n \le s < t} \frac{n^{\nu} \left| \widetilde{\alpha}_n(s;t) - \widetilde{B}_n(s) \right|}{s^{1/2-\nu}}. \tag{3.20}$$

On the other hand, we have for every $0 < t < 1$,

$$\{\widetilde{\alpha}_n(s); 0 \le s \le 1\} \overset{\mathcal{D}}{=} \{\alpha_n(s); 0 \le s \le 1\}$$

and

$$\left\{\widetilde{\beta}_n(s); 0 \le s \le 1\right\} \overset{\mathcal{D}}{=} \{\beta_n(s); 0 \le s \le 1\}.$$

It follows that

$$\{\widetilde{\alpha}_n(s;t); 0 \le s < t\} \overset{\mathcal{D}}{=} \{\alpha_n(s;t); 0 \le s < t\}$$

and

$$\left\{\widetilde{\beta}_n(s;t); 0 \le s < t\right\} \overset{\mathcal{D}}{=} \{\beta_n(s;t); 0 \le s < t\},$$

with $\widetilde{B}_n$ is Brownian bridge for each n. Then, having established Gaussian approximations above one may construct a sequence $U_1, U_2, ...$ of iid rv's uniformly distributed on $[0,1]$ and a sequence of Brownian bridges $B_1, B_2, ...$ defining on the probability space $(\Omega, \mathcal{A}, \mathbb{P})$ such that both (3.19) and (3.20) hold with $\widetilde{\alpha}_n$, $\widetilde{\beta}_n$ and $\widetilde{B}_n$ replaced respectively by α_n, β_n and B_n. This technique for constructing a such probability space, described in Lemma 3.1.1 in M. Csörgő (1983), is used for instance in both Csörgő *et al.* (1986) and Mason and van Zwet (1987).

References

Alexander, Kenneth S., 1987. The central limit theorem for weighted empirical processes indexed by sets. *J. Multivariate. Anal.* **22**, 313-339.

Csörgő, M., 1983. *Quantile processes with statistical applications.* Regional Conference Series in App. Math. SIAM, Philadelphia.

Brahimi, B., Meraghni, D., Necir, A., 2015. Approximations to the tail index estimator of a heavy-tailed distribution under random censoring and application. *Math. Methods Statist.* **24,** 266-279.

Csörgő, M., Révész, P., 1981. Strong approximations in probability and statistics. Probability and Mathematical Statistics. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London.

Csörgő, M., Csörgő, S., Horváth, L., Mason, D.M., 1986. Weighted empirical and quantile processes. *Ann. Probab.* **14**, 31-85.

Csörgő, S., Deheuvels, P., Mason, D., 1985. Kernel estimates of the tail index of a distribution. *Ann. Statist.* **13**, 1050-1077.

Deheuvels, P., Einmahl, J.H.J., 1996. On the strong limiting behavior of local functionals of empirical processes based upon censored data. *Ann. Probab.* **24**, 504-525.

Einmahl, J.H.J. and Koning, A.J., 1992. Limit theorems for a general weighted process under random censoring. *Canad. J. Statist.* **20**, 77-89.

Komlós, J., Major, P., Tusnády, G., 1975. An approximation of partial sums of independent 's and the sample . *I. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **32**, 111-131.

Komlós, J., Major, P., Tusnády, G., 1976. An approximation of partial sums of independent RV's, and the sample DF. II. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **34**, 33-58.

Mason, D. M., van Zwet, W. R., 1987. A refinement of the KMT inequality for the uniform empirical process. *Ann. Probab.* **15**, 871-884.

Peng, L., 2001. Estimating the mean of a heavy tailed distribution. *Statist. Probab. Lett.* **52**, 255-264.

Shorack, Galen R., Wellner, Jon A., 1982. Limit theorems and inequalities for the uniform empirical process indexed by intervals. *Ann. Probab.* **10**, 639-652.