

Quantum coherence of cosmological perturbations

Massimo Giovannini ¹

Department of Physics, Theory Division, CERN, 1211 Geneva 23, Switzerland

INFN, Section of Milan-Bicocca, 20126 Milan, Italy

Abstract

The degrees of quantum coherence of cosmological perturbations of different spins are computed in the large-scale limit and compared with the standard results holding for a single mode of the electromagnetic field in an optical cavity. The degree second-order coherence of curvature inhomogeneities (and, more generally, of the scalar modes of the geometry) reproduces faithfully the optical limit. For the vector and tensor fluctuations the numerical values of the normalized degrees of second-order coherence in the zero-time delay limit are always larger than unity (which is the Poisson benchmark value) but differ from the corresponding expressions obtainable in the framework of the single-mode approximation. General lessons are drawn on the quantum coherence of large-scale cosmological fluctuations.

¹Electronic address: massimo.giovannini@cern.ch

In quantum optics [1] the degrees of second-order coherence are determined by correlating the intensities of the radiation emitted by a source. This observation is at the heart of the Hanbury Brown-Twiss interferometry which is customarily used, with complementary purposes, both in quantum optics [2] and in high-energy physics [3]. The logic of interfering intensities (as opposed to amplitudes) is also applicable to large-scale cosmological perturbations [4] where the second-order interference effects can be used to establish, in a model independent manner, the classical or quantum origin of curvature fluctuations which have been directly probed by Cosmic Microwave Background experiments [5, 6].

The adjective *coherent* is often employed with different meanings: a signal is sometimes said to be coherent as opposed to noisy. The word coherent may also indicate the correlation between two or more functions although the functions themselves may have some random properties. The theory of quantum coherence [1], originally conceived and developed by various authors including Glauber and Sudarshan [7], gives an unambiguous answer by relating the coherence properties of a state (or of a source) to the minimization of the indetermination relations². The purpose of this analysis is to investigate, in a unified perspective, the quantum coherence of the cosmological perturbations of different spins and relate the obtained results to the conventional quantum optical approach based on the single-mode approximation.

The single-mode approximation is widely used to interpret a variety of observations both in Mach-Zender and Hanbury Brown-Twiss interferometry [1]. Since many experiments use plane parallel light beams whose transverse intensity profiles are not important for the measured quantities, it is often sufficient in interpreting the data to consider the light beams as exciting a single mode of the field. In this sense the viewpoint of quantum optics is *exclusive* insofar as the degrees of first- and second-order coherence are exclusively defined for a single mode of the field and they are given by[1]:

$$\bar{g}^{(1)}(\tau_1, \tau_2) = \frac{\langle \hat{a}^\dagger(\tau_1) \hat{a}(\tau_2) \rangle}{\sqrt{\langle \hat{a}^\dagger(\tau_1) \hat{a}(\tau_1) \rangle} \sqrt{\langle \hat{a}^\dagger(\tau_2) \hat{a}(\tau_2) \rangle}}, \quad (1)$$

$$\bar{g}^{(2)}(\tau_1, \tau_2) = \frac{\langle \hat{a}^\dagger(\tau_1) \hat{a}^\dagger(\tau_2) \hat{a}(\tau_2) \hat{a}(\tau_1) \rangle}{\langle \hat{a}^\dagger(\tau_1) \hat{a}(\tau_1) \rangle \langle \hat{a}^\dagger(\tau_2) \hat{a}(\tau_2) \rangle}. \quad (2)$$

Equations (1) and (2) define, respectively, the degrees of first and second-order temporal coherence. The expectation values of Eqs. (1)–(2) are evaluated on a specific quantum state. A single-mode coherent state is, by definition, an eigenstate of the annihilation operator (i.e. $\hat{a}|\alpha\rangle = \alpha|\alpha\rangle$). Therefore, in the zero time-delay limit (i.e. $\tau_1 - \tau_2 \rightarrow 0$) Eqs. (1) and (2) imply $\bar{g}^{(1)} = \bar{g}^{(2)} = 1$. According to Glauber theory, the property expressed by Eqs. (1) and

²A coherent state (characterized by a Poissonian statistics) is, by definition, coherent to all orders and the corresponding normalized degrees of quantum coherence always coincide with unity.

(2) holds, for a coherent state, to any order (i.e. $\bar{g}^{(1)} = \bar{g}^{(2)} = \dots = \bar{g}^{(n-1)} = \bar{g}^{(n)} = 1$). For a chaotic state with statistical weights provided by the Bose-Einstein distribution³, Eqs. (1) and (2) demand $\bar{g}^{(1)} = 1$ but $\bar{g}^{(2)} = 2$. This result is often dubbed by saying that chaotic (i.e. white) light is bunched and it exhibits super-Poissonian statistics [1, 7]. In the case of a single Fock state we have instead $\bar{g}^{(2)} = (1 - 1/n) < 1$ showing that Fock states always lead to sub-Poissonian behaviour [1]. While chaotic light is an example of bunched quantum state (i.e. $\bar{g}^{(2)} > 1$ implying more degree of second-order coherence than in the case of a coherent state), Fock states are instead antibunched (i.e. $\bar{g}^{(2)} < 1$) indicating a degree of second-order coherence smaller than in the case of a coherent state.

In the zero time-delay limit the degree of second-order coherence of Eq. (2) has a simple relation with the variance of the probability distribution associated with a given quantum state. Defining $\hat{N} = \hat{a}^\dagger \hat{a}$, Eq. (2) implies $\bar{g}^{(2)} = (D^2 - \bar{n})/\bar{n}^2$ where $\bar{n} = \langle \hat{N} \rangle$ and $D^2 = \langle \hat{N}^2 \rangle - \langle \hat{N} \rangle^2$. It is sometimes useful to define the so-called Mandel parameter [1] whose expression is $\mathcal{Q} = \bar{n}[\bar{g}^{(2)} - 1] = D^2/\bar{n} - 1$. The parameter \mathcal{Q} is directly related to the way second-order correlations are parametrized in subatomic physics [3]. In the case of a coherent state we have that $D^2 = \langle \hat{N} \rangle$ while $\mathcal{Q} = 0$; this means that the distribution underlying this state is just the Poisson distribution (i.e. the variance coincides with the mean value).

For the present ends it is important to appreciate that, in the Heisenberg description, the evolution of cosmological perturbations of different spins can be parametrized, in a unified perspective, as:

$$\begin{aligned}\hat{a}_{\vec{p}\alpha} &= e^{-i\varphi_p} \left[\cosh r_p \hat{b}_{\vec{p}\alpha} - e^{i\gamma_p} \sinh r_p \hat{b}_{-\vec{p}\alpha}^\dagger \right], \\ \hat{a}_{-\vec{p}\alpha}^\dagger &= e^{i\varphi_p} \left[\cosh r_p \hat{b}_{-\vec{p}\alpha}^\dagger - e^{-i\gamma_p} \sinh r_p \hat{b}_{\vec{p}\alpha} \right].\end{aligned}\quad (3)$$

Since $[\hat{a}_{\vec{k}\alpha}, \hat{a}_{\vec{p}\beta}^\dagger] = \delta_{\alpha\beta} \delta^{(3)}(\vec{k} - \vec{p})$ and $[\hat{b}_{\vec{k}\alpha}, \hat{b}_{\vec{p}\beta}^\dagger] = \delta_{\alpha\beta} \delta^{(3)}(\vec{k} - \vec{p})$ the transformation connecting the two sets of creation and annihilation operators is unitary. Equation (3) describes, in one shot, the scalar, vector and tensor case with the proviso that, for scalar fluctuations, the polarization index must be dropped; moreover, in the vector and tensor cases, α corresponds to the two (massless) vector or tensor polarizations, as it will be clear in a moment. The notations of Eq. (3) are conventional and the only essential point is that the unitary transformation connecting the two sets of creation and annihilation operators must be parametrized by two complex numbers

$$u_k(\tau) = e^{-i\varphi_k(\tau)} \cosh r_k(\tau), \quad v_k(\tau) = e^{-i(\varphi_k(\tau) - \gamma_k(\tau))} \sinh r_k(\tau), \quad (4)$$

³To evaluate Eqs. (1)–(2) for a chaotic mixture, the single mode density matrix in the number basis can be written as $\hat{\rho} = \sum_n p_n |n\rangle \langle n|$ where the statistical weights are given by $p_n = \bar{n}^n / (\bar{n} + 1)^{n+1}$ and \bar{n} is the mean number of quanta (coinciding with the Bose-Einstein occupation number only in the thermal case).

subjected to the condition $|u_k(\tau)|^2 - |v_k(\tau)|^2 = 1$. In the single-mode approximation described by Eqs. (1)–(2), the expression of Eq. (3) can be written as $\hat{a} = \cosh r \hat{b} - \sinh r \hat{b}^\dagger$ where the phases, for the sake of simplicity, have been fixed to zero. Taking the limit of zero time-delay and inserting these expressions in Eq. (2) we have that: $\bar{g}^{(2)} = 3 + 1/\bar{n}$ with $\bar{n} = \sinh^2 r$. Whenever $\bar{n} \gg 1$ (which corresponds to the physical situation in a cosmological setting) we have that $\bar{g}^{(2)} \rightarrow 3$, as implied for a squeezed state in the single-mode approximation [1].

In the single-mode approximation the degrees of second-order coherence range⁴ between $\bar{g}^{(2)} \rightarrow 1$ (in the case of a coherent state) to $\bar{g}^{(2)} \rightarrow 3$ (in the case of a squeezed vacuum state). The single-mode thermal state is super-Poissonian (i.e. $\bar{g}^{(2)} \rightarrow 2$) but less correlated than a squeezed vacuum state. In the case of large-scale cosmological perturbations, thanks to the results of the single-mode approximation, the normalized degrees of second-order coherence should all be close to 3 (and anyway greatly exceed 1). Before the direct scrutiny of this plausible expectation, we have to remark that the degrees of coherence and the corresponding Glauber correlation functions change in the scalar, vector and tensor cases since they are sensitive to the polarizations. The Glauber correlation in the scalar case can be written as:

$$\begin{aligned} \mathcal{S}^{(n,m)}(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m}) \\ = \text{Tr} \left[\hat{\rho} \hat{q}^{(-)}(x_1) \dots \hat{q}^{(-)}(x_n) \hat{q}^{(+)}(x_{n+1}) \dots \hat{q}^{(+)}(x_{n+m}) \right], \end{aligned} \quad (5)$$

where $x_i \equiv (\vec{x}_i, \tau_i)$ and $\hat{\rho}$ is the density operator representing the (generally mixed) state of the field \hat{q} . The field operator can always be expressed as $\hat{q}(x) = \hat{q}^{(+)}(x) + \hat{q}^{(-)}(x)$, with $\hat{q}^{(+)}(x) = \hat{q}^{(-)\dagger}(x)$. By definition we will have that $\hat{q}^{(+)}(x)|\text{vac}\rangle = 0$ and also that $\langle \text{vac} | \hat{q}^{(-)}(x) = 0$. It is understood that the state $|\text{vac}\rangle$ minimizes the appropriate Hamiltonian which is, in general, different, in the scalar, vector and tensor cases. These issues have been addressed in the current literature and more specific discussions can be found, for instance, in [4, 8, 9]. The mode expansion in the scalar case can be written as:

$$\hat{q}(\vec{x}, \tau) = \frac{1}{\sqrt{V}} \sum_{\vec{p}} \hat{q}_{\vec{p}}(\tau) e^{-i\vec{p}\cdot\vec{x}}, \quad \hat{q}_{\vec{p}} = \frac{1}{\sqrt{2p}} (\hat{a}_{\vec{p}} + \hat{a}_{-\vec{p}}^\dagger), \quad (6)$$

where V represents a fiducial (normalization) volume. As it is clear from Eq. (6) we also have that $\hat{q}_{\vec{p}}^\dagger = \hat{q}_{-\vec{p}}$. By switching from discrete to continuous modes the creation and annihilation operators obey $[\hat{a}_{\vec{k}}, \hat{a}_{\vec{p}}^\dagger] = \delta^{(3)}(\vec{k} - \vec{p})$ and the sums are replaced by integrals according to $\sum_{\vec{k}} \rightarrow V \int d^3k / (2\pi)^3$.

We shall always be concerned with the case of a conformally flat background geometry $\bar{g}_{\mu\nu} = a^2(\tau)\eta_{\mu\nu}$ where $\eta_{\mu\nu}$ is the Minkowski metric (with signature $(+, -, -, -)$) and $a(\tau)$

⁴Antibunched sources (like for instance Fock states) will be disregarded since they have no analog in cosmological applications where the occupation numbers of the field are always large.

is the scale factor in the conformal time parametrization. The field \hat{q} may represent various physical quantities: it could simply be a spectator field (minimally or non-minimally coupled to the conformally flat background geometry) but Eq. (5) applies to the case of the curvature perturbations where $\hat{q} = -\hat{\mathcal{R}}z$ and $\hat{\mathcal{R}}$ is the field operator corresponding to the perturbations of the spatial curvature on comoving orthogonal hypersurfaces [8]. If the source of inhomogeneity is represented by a scalar degree of freedom ϕ (not necessarily identified with the inflaton) we will have, as usual, that $z = a\phi'/\mathcal{H}$ where $\mathcal{H} = a'/a$ and the prime denotes a derivation with respect to the conformal time coordinate τ . In a complementary perspective, when the source of scalar fluctuations is given by a perfect and irrotational fluid, z will have a different analytic expression given by $z = a^2\sqrt{p_t + \rho_t}/(c_{st}\mathcal{H})$ (as originally suggested by Lukash [8]) where (p_t, ρ_t) are the total pressure and energy density of the fluid and $c_{st}^2 = p'_t/\rho'_t$ is the total sound speed. In both cases the evolution of $u_k(\tau)$ and $v_k(\tau)$ appearing in Eq. (4) can be written as:

$$u'_k = -ik u_k - \frac{z'}{z} v_k^*, \quad v'_k = -ik v_k - \frac{z'}{z} u_k^*, \quad (7)$$

where, as already mentioned, the prime denotes a derivation with respect to τ .

The Glauber correlation function has been originally defined in the vector case [7] and in the framework of quantum electrodynamics; the analog of Eq. (5) in the vector case is therefore given by:

$$\begin{aligned} & \mathcal{V}_{i_1, \dots, i_n, i_{n+1}, \dots, i_{n+m}}^{(n,m)}(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m}) \\ &= \text{Tr} \left[\hat{\rho} \hat{\mathcal{A}}_{i_1}^{(-)}(x_1) \dots \hat{\mathcal{A}}_{i_n}^{(-)}(x_n) \hat{\mathcal{A}}_{i_{n+1}}^{(+)}(x_{n+1}) \dots \hat{\mathcal{A}}_{i_{n+m}}^{(+)}(x_{n+m}) \right], \end{aligned} \quad (8)$$

where $x_i \equiv (\vec{x}_i, \tau_i)$ and $\hat{\rho}$ is the density operator representing the (generally mixed) state of the field⁵ $\hat{\mathcal{A}}_i(\vec{x}, \tau)$. The difference between Eqs. (5) and (8) is represented by the vector polarizations denoted, in Eq. (8), by $(i_1, \dots, i_n, i_{n+1}, \dots, i_{n+m})$. With the same notations of Eq. (6) the mode expansion in the vector case can be written as:

$$\hat{\mathcal{A}}_i(\vec{x}, \tau) = \frac{1}{\sqrt{V}} \sum_{\vec{p}, \alpha} e_i^{(\alpha)} \hat{\mathcal{A}}_{\vec{p}, \alpha}(\tau) e^{-i\vec{p} \cdot \vec{x}}, \quad \hat{\mathcal{A}}_{\vec{p}, \alpha} = \frac{1}{\sqrt{2p}} (\hat{a}_{\vec{p}\alpha} + \hat{a}_{-\vec{p}\alpha}^\dagger). \quad (9)$$

where $e_i^{(\alpha)}(\hat{k})$ (with $\alpha = 1, 2$) are the two polarizations which are mutually orthogonal and orthogonal to \hat{k} ; as in the case of Eq. (6) we have that $\hat{\mathcal{A}}_{\vec{p}, \alpha}^\dagger = \hat{\mathcal{A}}_{-\vec{p}, \alpha}$. Since we shall be mainly concerned with massless and divergence-less vectors the sum over the polarizations will be given by $\sum_\alpha e_i^{(\alpha)}(\hat{k}) e_j^{(\alpha)}(\hat{k}) = P_{ij}(\hat{k})$ where $P_{ij}(\hat{k}) = \delta_{ij} - \hat{k}_i \hat{k}_j$. In a cosmological

⁵The field $\hat{\mathcal{A}}_i(\vec{x}, \tau)$ can always be expressed as $\hat{\mathcal{A}}_i(x) = \hat{\mathcal{A}}_i^{(+)}(x) + \hat{\mathcal{A}}_i^{(-)}(x)$, with $\hat{\mathcal{A}}_i^{(+)}(x) = \hat{\mathcal{A}}_i^{(-)\dagger}(x)$. By definition we will have that $\hat{\mathcal{A}}_i^{(+)}(x)|\text{vac}\rangle = 0$ and also that $\langle \text{vac} | \hat{\mathcal{A}}_i^{(-)}(x) = 0$; the state $|\text{vac}\rangle$ denotes the vacuum.

context the vector fluctuations may come either from the metric or from gauge fields [4]. In the four-dimensional case there are actually two divergence-less vectors related to the vector fluctuations fluctuations of the metric; they would correspond to $\delta_v g_{0i} = -a^2 Q_i$ and to $\delta_v g_{ij} = a^2(\partial_i W_j + \partial_j W_i)$ with $\partial_i Q^i = 0, \partial_i W^i = 0$. These two vectors, however, are not amplified in the concordance paradigm [5, 6] and shall not be specifically considered here. Conversely the vector fluctuations possibly relevant for our purposes are the ones of gauge fields; in this case $u_k(\tau)$ and $v_k(\tau)$ obey an equation analog to Eq. (7) but with $z'/z \rightarrow \chi'/\chi$ where χ denotes the susceptibility of the Abelian gauge field (see e.g. [4]).

As in the scalar and vector cases the (divergenceless and traceless) tensor field $\hat{\mu}_{ij}(\vec{x}, \tau)$ can always be expressed as $\hat{\mu}_{ij}(x) = \hat{\mu}_{ij}^{(+)}(x) + \hat{\mu}_{ij}^{(-)}(x)$ and their mode expansion is⁶

$$\hat{\mu}_{ij}(\vec{x}, \tau) = \frac{\sqrt{2\ell_P}}{\sqrt{V}} \sum_{\vec{p}, \alpha} e_{ij}^{(\alpha)} \hat{\mu}_{\vec{p}, \alpha}(\tau) e^{-i\vec{p} \cdot \vec{x}}, \quad \hat{\mu}_{\vec{p}, \alpha} = \frac{1}{\sqrt{2p}} (\hat{a}_{\vec{p}\alpha} + \hat{a}_{-\vec{p}\alpha}^\dagger), \quad (10)$$

where $\ell_P = 8\pi G$; following the standard practice, in Eq. (10) and elsewhere we shall adopt units $16\pi G = 1$. Furthermore, as in the scalar and vector cases we shall have that $\hat{\mu}_{\vec{p}, \alpha}^\dagger = \hat{\mu}_{-\vec{p}, \alpha}$. The two polarizations of the gravitons in a conformally flat background geometry are:

$$e_{ij}^{(\oplus)}(\hat{k}) = (\hat{m}_i \hat{m}_j - \hat{n}_i \hat{n}_j), \quad e_{ij}^{(\otimes)}(\hat{k}) = (\hat{m}_i \hat{n}_j + \hat{n}_i \hat{m}_j), \quad (11)$$

where $\hat{k}_i = k_i/|\vec{k}|$, $\hat{m}_i = m_i/|\vec{m}|$ and $\hat{n}_i = n_i/|\vec{n}|$ denote three mutually orthogonal directions. It follows from Eq. (11) that $e_{ij}^{(\lambda)} e_{ij}^{(\lambda')} = 2\delta_{\lambda\lambda'}$ while the sum over the polarizations gives:

$$A_{ijmn}(\hat{k}) = \sum_{\lambda} e_{ij}^{(\lambda)}(\hat{k}) e_{mn}^{(\lambda)}(\hat{k}) = \left[P_{mi}(\hat{k}) P_{nj}(\hat{k}) + P_{mj}(\hat{k}) P_{ni}(\hat{k}) - P_{ij}(\hat{k}) P_{mn}(\hat{k}) \right]; \quad (12)$$

where, as already mentioned, $P_{ij}(\hat{k}) = (\delta_{ij} - \hat{k}_i \hat{k}_j)$. In the tensor case the Glauber correlation function is finally given by:

$$\begin{aligned} & \mathcal{T}_{(i_1 j_1), \dots, (i_n j_n), (i_{n+1} j_{n+1}), \dots, (i_{n+m} j_{n+m})}^{(n,m)}(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m}) \\ &= \text{Tr} \left[\hat{\rho} \hat{\mu}_{i_1 j_1}^{(-)}(x_1) \dots \hat{\mu}_{i_n j_n}^{(-)}(x_n) \hat{\mu}_{i_{n+1} j_{n+1}}^{(+)}(x_{n+1}) \dots \hat{\mu}_{i_{n+m} j_{n+m}}^{(+)}(x_{n+m}) \right], \end{aligned} \quad (13)$$

where $x_i \equiv (\vec{x}_i, \tau_i)$ and $\hat{\rho}$ is the density operator representing the (generally mixed) state of the field $\hat{\mu}_{ij}$. In Eq. (13) the polarization structure is different from the vector case of Eq. (8): instead of $n + m$ vector indices we have $n + m$ tensor indices.

⁶The tensor modes arise naturally in the framework of the concordance scenario since their evolution is not Weyl invariant as established long ago by Grishchuk [9]. In the case of the tensor modes the evolution of $u_k(\tau)$ and $v_k(\tau)$ is given by Eq. (7) but with z'/z replaced by $a'/a = \mathcal{H}$.

The general expressions of the Glauber correlators defined in Eqs. (5), (8) and (13) contain a wealth of informations. For the present purposes the intensity correlators (also relevant in the context of the Hanbury Brown-Twiss interferometry) can be deduced from Eqs. (5), (8) and (13) by identifying the corresponding space-time points as follows:

$$x_1 \equiv x_{n+1}, \quad x_2 \equiv x_{n+2}, \quad \dots \quad x_n \equiv x_{2n}. \quad (14)$$

In this case the original Glauber correlator will effectively be a function of n points and it will describe the correlation of n intensities. Using Eq. (14) into Eq. (5) in the case $n = 2$ we have that the correlation of the scalar intensities can be written as:

$$\begin{aligned} \mathcal{S}^{(2)}(x_1, x_2) &= \langle \hat{\mathcal{I}}_{\mathcal{S}}(\vec{x}_1, \tau_1) \hat{\mathcal{I}}_{\mathcal{S}}(\vec{x}_2, \tau_2) \rangle = \int \frac{d^3 k_1}{2k_1(2\pi)^3} \int \frac{d^3 k_2}{2k_2(2\pi)^3} \\ &\times \left[|v_{k_1}(\tau_1)|^2 |v_{k_2}(\tau_2)|^2 + v_{k_1}^*(\tau_1) v_{k_1}(\tau_2) v_{k_2}^*(\tau_2) v_{k_2}(\tau_1) e^{-i(\vec{k}_1 - \vec{k}_2) \cdot \vec{r}} \right. \\ &\left. + v_{k_1}^*(\tau_1) u_{k_1}^*(\tau_2) u_{k_2}(\tau_1) v_{k_2}(\tau_2) e^{-i(\vec{k}_1 + \vec{k}_2) \cdot \vec{r}} \right], \end{aligned} \quad (15)$$

where $\vec{r} = \vec{x}_1 - \vec{x}_2$. Similarly, using Eq. (14) into Eq. (8), the correlation of the intensities in the vector case is given by

$$\begin{aligned} \mathcal{V}^{(2)}(x_1, x_2) &= \langle \hat{\mathcal{I}}_{\mathcal{V}}(\vec{x}_1, \tau_1) \hat{\mathcal{I}}_{\mathcal{V}}(\vec{x}_2, \tau_2) \rangle = \int \frac{d^3 k_1}{2k_1(2\pi)^3} \int \frac{d^3 k_2}{2k_2(2\pi)^3} \\ &\times \left\{ |v_{k_1}(\tau_1)|^2 |v_{k_2}(\tau_2)|^2 P_{ii}(\hat{k}_1) P_{jj}(\hat{k}_2) + \right. \\ &+ P_{ij}(\hat{k}_1) P_{ij}(\hat{k}_2) \left[v_{k_1}^*(\tau_1) v_{k_1}(\tau_2) v_{k_2}^*(\tau_2) v_{k_2}(\tau_1) e^{-i(\vec{k}_1 - \vec{k}_2) \cdot \vec{r}} \right. \\ &\left. \left. + v_{k_1}^*(\tau_1) u_{k_1}^*(\tau_2) u_{k_2}(\tau_1) v_{k_2}(\tau_2) e^{-i(\vec{k}_1 + \vec{k}_2) \cdot \vec{r}} \right] \right\}. \end{aligned} \quad (16)$$

Finally, in the tensor case, the analog of Eqs. (15) and (16) is given by:

$$\begin{aligned} \mathcal{T}^{(2)}(x_1, x_2) &= \langle \hat{\mathcal{I}}_{\mathcal{T}}(\vec{x}_1, \tau_1) \hat{\mathcal{I}}_{\mathcal{T}}(\vec{x}_2, \tau_2) \rangle = \int \frac{d^3 k_1}{2k_1(2\pi)^3} \int \frac{d^3 k_2}{2k_2(2\pi)^3} \\ &\times \left\{ |v_{k_1}(\tau_1)|^2 |v_{k_2}(\tau_2)|^2 A_{i j i j}(\hat{k}_1) A_{\ell m \ell m}(\hat{k}_2) \right. \\ &+ A_{i j \ell m}(\hat{k}_1) A_{i j \ell m}(\hat{k}_2) \left[v_{k_1}^*(\tau_1) v_{k_1}(\tau_2) v_{k_2}^*(\tau_2) v_{k_2}(\tau_1) e^{-i(\vec{k}_1 - \vec{k}_2) \cdot \vec{r}} \right. \\ &\left. \left. + v_{k_1}^*(\tau_1) u_{k_1}^*(\tau_2) u_{k_2}(\tau_1) v_{k_2}(\tau_2) e^{-i(\vec{k}_1 + \vec{k}_2) \cdot \vec{r}} \right] \right\}. \end{aligned} \quad (17)$$

We are now in condition of comparing the degree of second-order coherence obtained in the framework of the single-mode approximation (see Eq. (2)) with the results of Eqs. (15), (16) and (17) which determine the corresponding degrees of second-order coherence:

$$g_{\mathcal{S}}^{(2)}(x_1, x_2) = \frac{\mathcal{S}^{(2)}(x_1, x_2)}{\mathcal{S}^{(1)}(x_1, x_1) \mathcal{S}^{(1)}(x_1, x_1)} = \frac{\langle \hat{\mathcal{I}}_{\mathcal{S}}(\vec{x}_1, \tau_1) \hat{\mathcal{I}}_{\mathcal{S}}(\vec{x}_2, \tau_2) \rangle}{\langle \hat{\mathcal{I}}_{\mathcal{S}}(\vec{x}_1, \tau_1) \rangle \langle \hat{\mathcal{I}}_{\mathcal{S}}(\vec{x}_2, \tau_2) \rangle}, \quad (18)$$

$$g_{\mathcal{V}}^{(2)}(x_1, x_2) = \frac{\mathcal{V}^{(2)}(x_1, x_2)}{\mathcal{V}^{(1)}(x_1, x_1)\mathcal{V}^{(1)}(x_1, x_1)} = \frac{\langle \hat{\mathcal{I}}_{\mathcal{V}}(\vec{x}_1, \tau_1) \hat{\mathcal{I}}_{\mathcal{V}}(\vec{x}_2, \tau_2) \rangle}{\langle \hat{\mathcal{I}}_{\mathcal{V}}(\vec{x}_1, \tau_1) \rangle \langle \hat{\mathcal{I}}_{\mathcal{V}}(\vec{x}_2, \tau_2) \rangle}, \quad (19)$$

$$g_{\mathcal{T}}^{(2)}(x_1, x_2) = \frac{\mathcal{T}^{(2)}(x_1, x_2)}{\mathcal{T}^{(1)}(x_1, x_1)\mathcal{T}^{(1)}(x_1, x_1)} = \frac{\langle \hat{\mathcal{I}}_{\mathcal{T}}(\vec{x}_1, \tau_1) \hat{\mathcal{I}}_{\mathcal{T}}(\vec{x}_2, \tau_2) \rangle}{\langle \hat{\mathcal{I}}_{\mathcal{T}}(\vec{x}_1, \tau_1) \rangle \langle \hat{\mathcal{I}}_{\mathcal{T}}(\vec{x}_2, \tau_2) \rangle}. \quad (20)$$

In the zero time-delay limit $\tau_1 - \tau_2 \rightarrow 0$ the scalar degree of quantum coherence becomes

$$g_S^{(2)}(\vec{r}, \tau) = 1 + \frac{\int k_1 dk_1 |v_{k_1}(\tau)|^2 j_0(k_1 r) \int k_2 dk_2 |v_{k_2}(\tau)|^2 j_0(k_2 r)}{\int k_1 dk_1 |v_{k_1}(\tau)|^2 \int k_2 dk_2 |v_{k_2}(\tau)|^2} + \frac{\int k_1 dk_1 u_{k_1}^*(\tau) v_{k_1}^*(\tau) j_0(k_1 r) \int k_2 dk_2 u_{k_2}(\tau) v_{k_2}(\tau) j_0(k_2 r)}{\int k_1 dk_1 |v_{k_1}(\tau)|^2 \int k_2 dk_2 |v_{k_2}(\tau)|^2}, \quad (21)$$

where $j_0(k_1 r)$ and $j_0(k_2 r)$ denote the spherical Bessel function of zeroth order. In the large-scale limit $k_1 r \ll 1$ and $k_2 r \ll 1$, Eq. (21) implies that $g_S^{(2)}(\vec{r}, \tau) \rightarrow 3$. The reason for this result stems from the observation that $u_{k_1}^*(\tau) v_{k_1}^*(\tau) u_{k_2}(\tau) v_{k_2}(\tau) = |v_{k_1}(\tau)|^2 |v_{k_2}(\tau)|^2 [1 + \mathcal{O}(k_1 \tau) + \mathcal{O}(k_2 \tau)]$. This result can be easily obtained in the case of a quasi de Sitter evolution where the solution of Eq. (7) with the correct boundary conditions implies:

$$v_k(\tau) = -\frac{e^{-ik\tau}}{2k^2\tau^2}, \quad u_k(\tau) = e^{-ik\tau} \left[1 - \frac{i}{k\tau} - \frac{1}{2k^2\tau^2} \right]. \quad (22)$$

In the vector and tensor cases the analysis proceeds along the same lines, with the difference that the angular integrals appearing in Eqs. (16) and (17) are more complicated. This is due to the presence of the terms

$$P_{ij}(\hat{k}_1)P_{ij}(\hat{k}_2) = 1 + (\hat{k}_1 \cdot \hat{k}_2)^2, \quad A_{ij\ell m}(\hat{k}_1)A_{ij\ell m}(\hat{k}_2) = [1 + (\hat{k}_1 \cdot \hat{k}_2)^2][1 + 3(\hat{k}_1 \cdot \hat{k}_2)^2]; \quad (23)$$

needless to say that, by definition, $P_{ii}(\hat{k}) = 2$ and that $A_{ijij} = 4$. After performing some lengthy (but straightforward) angular integrals, the results of Eq. (23) imply that, in the zero-time delay limit $g_{\mathcal{V}}^{(2)}(x_1, x_2)$ and $g_{\mathcal{T}}^{(2)}(x_1, x_2)$ go, respectively, to $5/3$ and to $71/60$.

All in all Eqs. (18), (19) and (20) in the zero time-delay limit (i.e. $\tau_1 \rightarrow \tau_2$) and for large-scales imply the following results for the degrees of quantum coherence:

$$\lim_{\tau_1 \rightarrow \tau_2, kr \ll 1} g_S^{(2)}(r, \tau_1, \tau_2) = g_S^{(2)}(r, \tau) \rightarrow 3, \quad (24)$$

$$\lim_{\tau_1 \rightarrow \tau_2, kr \ll 1} g_{\mathcal{V}}^{(2)}(r, \tau_1, \tau_2) = g_{\mathcal{V}}^{(2)}(r, \tau) \rightarrow \frac{5}{3}, \quad (25)$$

$$\lim_{\tau_1 \rightarrow \tau_2, kr \ll 1} g_{\mathcal{T}}^{(2)}(r, \tau_1, \tau_2) = g_{\mathcal{T}}^{(2)}(r, \tau) \rightarrow \frac{71}{60}. \quad (26)$$

The quantum optical results (obtained in the single-mode approximation) is only recovered in the scalar case where the degree of second-order coherence goes to 3 exactly as in the case of a (single-mode) squeezed state. When the vector polarizations are taken into account

the degree of quantum coherence is always super-Poissonian but is given by $5/3$. Finally, in the tensor case, the degree of quantum coherence is $71/60$ (i.e. slightly above 1). We have therefore that $g_S^{(2)}(r, \tau) > g_V^{(2)}(r, \tau) > g_T^{(2)}(r, \tau)$. In the vector and the tensor cases the degree of second-order approaches 1 from above, as implied by the super-Poissonian character of the original quantum state of relic photons [4] and of relic gravitons [9].

It is well known that first order interference effects between the amplitudes cannot be used to distinguish the nature of different quantum states of the radiation field. Young interferometry is not able, by itself, to provide information on the statistical properties of the quantum correlations since various states with diverse physical properties (such as laser light and chaotic light) may lead to comparable degrees of first-order coherence [1]. The statistical properties of the quantum states can be disambiguated by examining the higher degrees of coherence. This program has been specifically suggested also in the case of the large-scale curvature perturbations determining the temperature and polarization anisotropies of the Cosmic Microwave Background [4]. For these analyses the new generations of CMB detectors and the Hanbury Brown-Twiss interferometry in the THz region could be a plausible framework. In this respect the present findings suggest that the degree of second-order coherence for the curvature perturbations is roughly thrice the one of the relic gravitational waves. A full account of this discussion is beyond the scopes of this paper.

The normalized degrees of second-order coherence for the cosmological perturbations have been computed with the purpose of comparing the obtained results with the standard quantum optical derivations obtained in the single-mode approximation. Since the large-scale cosmological perturbations are typically described by squeezed quantum states, it could be naively expected that their degree of second-order coherence should always equal 3 in spite of the spin of the fluctuation. While this is true in the scalar case, for the vector and tensor fluctuations the presence of the polarizations reduces the degree of second-order coherence in comparison with the scalar case and brings the result much closer to the Poissonian limit. In quantum optics the single-mode approximation is fully justified when the experimental set-up effectively involves a single mode of the electromagnetic field in a cavity. In the cosmological context this exclusive perspective is not typical and the sum over the polarizations cannot be neglected since the quantum state of large-scale fluctuations is generally unpolarized. Taken at face value the results presented here suggest that the single-mode approximation cannot be used in a cosmological situation. All in all we can instead observe that the effect of the polarizations is a progressive reduction of the degree of second-order coherence. This reduction preserves the super-Poissonian character of the quantum state so that the Poissonian limit (typical of the coherent state) is never reached.

The author wishes to thank J. Vigen and T. Basaglia of the CERN scientific information service for their kind assistance.

References

- [1] J. Klauder and E. Sudarshan, *Fundamentals of quantum optics* (Benjamin, New York, 1968); R. Loudon, *The quantum theory of light* (Clarendon Press, Oxford, third edition 2000).
- [2] R. Hanbury Brown and R. Q. Twiss, *Nature* **178**, 1046 (1956); R. Hanbury Brown and R. Q. Twiss, *Proc. Roy. Soc. (London)* **A242**, 300 (1957); *Proc. Roy. Soc. (London)* **A243**, 291 (1958).
- [3] G. I. Kopylov, M. I. Podgoretsky, *Sov. J. Nucl. Phys.* **15**, 219-223 (1972) [*Yad. Fiz.* **15**, 392 (1972)]; *Sov. J. Nucl. Phys.* **18**, 336 (1973) [*Yad. Fiz.* **18**, 656 (1973)]; G. Cocconi, *Phys. Lett.* **B49**, 459 (1974); D. H. Boal, C. K. Gelbke, B. K. Jennings, *Rev. Mod. Phys.* **62**, 553 (1990); G. Baym, *Acta Phys. Polon. B* **29**, 1839 (1998).
- [4] M. Giovannini, *Phys. Rev. D* **61**, 087306 (2000); *Phys. Rev. D* **83**, 023515 (2011); *Class. Quant. Grav.* **34**, no. 3, 035019 (2017).
- [5] D. N. Spergel *et al.*, *Astrophys. J. Suppl.* **148**, 175 (2003); D. N. Spergel *et al.*, *ibid.* **170**, 377 (2007); L. Page *et al.* *ibid.* **170**, 335 (2007).
- [6] B. Gold *et al.*, *ibid.* **192**, 15 (2011); D. Larson, *et al.*, *ibid.* **192**, 16 (2011); C. L. Bennett *et al.*, *ibid.* **192**, 17 (2011); G. Hinshaw *et al.*, *ibid.* **208** 19 (2013); C. L. Bennett *et al.*, *ibid.* **208** 20 (2013); P. A. R. Ade *et al.* [BICEP2 Collaboration], *Phys. Rev. Lett.* **112**, 241101 (2014); P. A. R. Ade *et al.* [Planck Collaboration], *Astron. Astrophys.* **571**, A22 (2014); *Astron. Astrophys.* **571**, A16 (2014); *Astron. Astrophys.* **594**, A20 (2016).
- [7] R. J. Glauber, *Phys. Rev. Lett.* **10**, 84 (1963); E. C. Sudarshan, *Phys. Rev. Lett.* **10**, 277 (1963); R. J. Glauber, *Phys. Rev.* **130**, 2529 (1963); *Phys. Rev.* **131**, 2766 (1963).
- [8] V. N. Lukash, *Sov. Phys. JETP* **52**, 807 (1980) [*Zh. Eksp. Teor. Fiz.* **79**, 1601 (1980)]; G. V. Chibisov, V. F. Mukhanov, *Mon. Not. Roy. Astron. Soc.* **200**, 535-550 (1982); H. Kodama, M. Sasaki, *Prog. Theor. Phys. Suppl.* **78**, 1 (1984); M. Sasaki, *Prog. Theor. Phys.* **76**, 1036 (1986). V. F. Mukhanov, *Sov. Phys. JETP* **67**, 1297 (1988) [*Zh. Eksp. Teor. Fiz.* **94**, 1 (1988)].
- [9] L. P. Grishchuk, *Sov. Phys. JETP* **40**, 409 (1975) [*Zh. Eksp. Teor. Fiz.* **67**, 825 (1974)]; *Annals N. Y. Acad. Sci.* **302**, 439 (1977); L. P. Grishchuk and Yu. V. Sidorov *Phys. Rev. D* **42**, 3413 (1990); L. P. Grishchuk, *Class. Quantum Grav.* **10**, 2449 (1993).