

A Truncated Cramér-von Mises test of normality

Juan Kalemkerian

Universidad de la República, Facultad de Ciencias.

December 25, 2021

Abstract

A new test of normality based on a standardised empirical process is introduced in this article. The first step is to introduce a Cramér–von Mises type statistic with weights equal to the inverse of the standard normal density function supported on a symmetric interval $[-a_n, a_n]$ depending on the sample size n . The sequence of end points a_n tends to infinity, and is chosen so that the statistic goes to infinity at the speed of $\ln \ln n$. After subtracting the mean, a suitable test statistic is obtained, with the same asymptotic law as the well-known Shapiro–Wilk statistic. The performance of the new test is described and compared with three other well-known tests of normality, namely, Shapiro–Wilk, Anderson–Darling and that of del Barrio-Matrán, Cuesta Albertos, and Rodríguez Rodríguez, by means of power calculations under many alternative hypotheses.

Keywords: empirical process, normality tests, Cramér-von-Mises statistic. 60G10

1 Definitions and notation

In this section we will introduce some definitions and notation to be used throughout this paper. The notation $:=$ means a definition. Given $X_1, X_2, \dots, X_n$ iid, we define $\bar{X}_n := \frac{X_1 + X_2 + \dots + X_n}{n}$ the sample mean, $S_n^2 := \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2$ the sample variance, and $Y_i := \frac{X_i - \bar{X}_n}{S_n}$ for $i = 1, 2, \dots, n$, and we define the standardised empirical process $\hat{b}_n(x) := \frac{1}{\sqrt{n}} \sum_{i=1}^n (1_{\{Y_i \leq x\}} - \phi(x))$ for all $x \in \mathbb{R}$. Also we write $F_n(x) := \frac{1}{n} \sum_{i=1}^n 1_{\{X_i \leq x\}}$ for the empirical distribution, and $b_n(x) := \frac{1}{\sqrt{n}} \sum_{i=1}^n (1_{\{X_i \leq x\}} - \phi(x))$.

We write $\varphi(x) := \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2\sigma^2}}$ and $\phi(x) := \int_{-\infty}^x \varphi(t) dt$ for the density and distribution functions of $N(0, 1)$. The notation $\xrightarrow{P}$, $\xrightarrow{a.s.}$ and $\xrightarrow{w}$ means convergence in probability, almost surely and in distribution, respectively. Throughout this paper, we write a_n for the sequence $a_n := \phi^{-1}(1 - 1/n)$.

2 Introduction

In the present paper, we study the Cramér–von Mises statistic to test normality when the weight function is $\psi(t) = \frac{1}{\varphi^2(\phi^{-1}(t))}$ and we integrate over the interval $(-a_n, a_n)$ instead of over all the real line. The main contribution of this paper is that we will obtain the asymptotic distribution of this ‘truncated’ kind of Cramér–von–Mises statistic and we will prove that its limit distribution is equivalent to that of the Shapiro–Wilk test. Also, for fixed sample sizes, we will simulate the behaviour of its power, for a wide class of alternative hypotheses, and we will see that its performance is comparable to that of the Shapiro–Wilk and Anderson–Darling tests, improving them in many cases. Given $X_1, X_2, \dots, X_n$ iid with distribution F when the null hypothesis is $F = F_0$ for a certain F_0 (fixed), Cramér (1928) postulated the statistic $n \int_{-\infty}^{+\infty} (F_n(x) - F_0(x))^2 dx$, and more generally von Mises (1931) took $n \int_{-\infty}^{+\infty} (F_n(x) - F_0(x))^2 \rho(x) dx$ when ρ is some non-negative weight function. Later Smirnov (1936, 1937) proposed working with the statistic $n \int_{-\infty}^{+\infty} (F_n(x) - F_0(x))^2 \psi(F_0(x)) dF_0(x)$, for certain ψ . Nowadays all these statistics (varying ψ) are called Cramér–von Mises statistics. The two most popular cases are when $\psi(t) = 1$ (because of its simplicity and similarity with the original Cramér statistic), i.e. $w_n^2 := n \int_{-\infty}^{+\infty} (F_n(x) - F_0(x))^2 dF_0(x)$, which is called the Cramér–von Mises statistic; and when $\psi(t) = \frac{1}{t(1-t)}$, proposed by Anderson and Darling (1954), i.e. $A_n^2 := n \int_{-\infty}^{+\infty} \frac{(F_n(x) - F_0(x))^2}{F_0(x)(1-F_0(x))} dF_0(x)$. In this case, the quadratic difference $(F_n(x) - F_0(x))^2$ is normalized by dividing it by its expected value, and this is called the Anderson–Darling statistic.

A_n^2 has the advantage of being generally more powerful than w_n^2 , for a wide class of alternative hypotheses, see, for instance, Stephens (1986).

When the null hypothesis is that the distribution lies in a certain parametric family, i.e., $F(x) = F(x, \theta)$ for certain θ , then it is natural to extend the Cramér–von Mises statistics used for a simple hypothesis in the form

$$n \int_{-\infty}^{+\infty} \left(F_n(x) - F(x, \hat{\theta}) \right)^2 \psi \left(F(x, \hat{\theta}) \right) dF(x, \hat{\theta})$$

where $\hat{\theta}$ is a suitable estimator of θ . Then we obtain $\hat{A}_n^2 := n \int_{-\infty}^{+\infty} \frac{(F_n(x) - F(x, \hat{\theta}))^2}{F(x, \hat{\theta})(1 - F(x, \hat{\theta}))} dF(x, \hat{\theta})$ and $\hat{w}_n^2 := n \int_{-\infty}^{+\infty} \left(F_n(x) - F(x, \hat{\theta}) \right)^2 dF(x, \hat{\theta})$. Again, generally, $\hat{A}_n^2$ performs better than $\hat{w}_n^2$.

The general study of empirical processes when parameters are estimated was conducted by Durbin (1973). He used the theory of weak convergence in $D[0, 1]$. Durbin's results give the asymptotic distribution of a wide class of Cramér–von Mises statistics and many others.

Normality tests are a particular case of goodness of fit tests. Many tests of normality have been designed. Stephens (1986), for instance, provides a good summary. The theory of tests for normality was initiated by Pearson (1895, 1930), Fisher (1930) and Williams (1935), with the $\sqrt{\beta_1}$ and β_2 tests (third and fourth standardised moments). Of course, the performance and the asymptotic distribution of the classical Kolmogorov–Smirnov, χ^2 , Cramér–von Mises, Anderson–Darling and many other tests of normality, have been extensively studied. Further, with the Shapiro–Wilk (1965) test for normality, there began the development of correlation and regression tests; see, for instance, Lockhart and Stephens (1998) for a good review of this subject. In the normal case, the Cramér–von Mises statistics are $n \int_{-\infty}^{+\infty} \left(F_n(t) - \phi \left(\frac{t - \bar{X}_n}{S_n} \right) \right)^2 \psi \left(\phi \left(\frac{t - \bar{X}_n}{S_n} \right) \right) \varphi \left(\frac{t - \bar{X}_n}{S_n} \right) dt$.

In the normal case, there have also been extensively studied, both theoretically and empirically, the statistics $\hat{w}_n^2$ and $\hat{A}_n^2$.

In Csörgő and Faraway (1996) the authors find an exact and asymptotic distribution for w_n^2 . In Pettitt and Stephens (1976) the authors modified the Cramér–von Mises statistic to allow them to test the normality of censored samples. The asymptotic theory for this is found in Pettitt (1976). Stephens (1974), found that in the case of the normality tests $\hat{w}_n^2$ and $\hat{A}_n^2$, especially the latter, are comparable in terms of power with the Shapiro–Wilk test.

The paper is organized as follows. In Section 3, we will examine the reasons for taking this particular weight function ψ . In Section 4 we will find the asymptotic distribution of the proposed test and give a proof that it is equivalent to the asymptotic distribution of the Shapiro–Wilk test. Our concluding remarks are in section 5. In Appendix 1 we will calculate the power of this test under certain alternative hypotheses, and we will compare it with other tests, like the Shapiro–Wilk, Anderson–Darling, and others. In Appendix 2 we give indications for calculating the statistic and in Appendix 3 we give tables with their critical values for different possible levels of significance and for different values of sample size n .

3 Test Aims and Approach

Given $X_1, X_2, \dots, X_n$ iid random variables with distribution F , we want to test $H_0 : F(x) = \phi\left(\frac{x-\mu}{\sigma}\right)$ for certain μ and σ . In del Barrio et al. (1999), a test of goodness of fit for normal distributions based on the Wasserstein distance is presented, and they prove that that test statistic has an asymptotic distribution equivalent to that of the Shapiro–Wilk test (1965). This limiting distribution is $\int_{-\infty}^{+\infty} \frac{b^2(x) - \mathbb{E}b^2(x)}{\varphi(x)} dx - \left(\int_{-\infty}^{+\infty} xb(x)dx\right)^2 - \left(\int_{-\infty}^{+\infty} b(x)dx\right)^2$ where $\{b(x)\}_{x \in \mathbb{R}}$ is a ϕ -Brownian bridge. The first summand of this decomposition allows us to propose $T_n^* := \int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x)}{\varphi(x)} dx$ to test normality. Although $\int_{-\infty}^{+\infty} \frac{\widehat{b}_n^2(x)}{\varphi(x)} dx < +\infty$ for each n , Kalemkerian (2016) proves that $\int_{-a_n}^{a_n} \frac{\mathbb{E}(\widehat{b}_n^2(x))}{\varphi(x)} dx = \ln(\ln n) + c_n$, where $\{c_n\}$ is a convergent sequence. This result is similar to the equality $\int_{-a_n}^{a_n} \frac{\mathbb{E}b^2(x)}{\varphi(x)} dx = \ln(\ln n) + \ln 2 + \gamma + o(1)$ where γ is Euler's constant, proved in de Wet and Venter (1972). Then $\int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x)}{\varphi(x)} dx$ goes to infinity as $n \rightarrow \infty$. Hence we cannot expect T_n^* to converge. So if we follow the ideas of del Barrio et al. (1999), we can propose studying

$$\int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x) - \mathbb{E}\widehat{b}_n^2(x)}{\varphi(x)} dx \text{ or } \int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x) - \mathbb{E}b^2(x)}{\varphi(x)} dx$$

as a test statistic with the intention of obtaining an asymptotic distribution, which means that the rejection zone is

$$\left\{ \int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x) - \mathbb{E}\widehat{b}_n^2(x)}{\varphi(x)} dx \geq c \right\} \text{ or } \left\{ \int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x) - \mathbb{E}b^2(x)}{\varphi(x)} dx \geq c \right\}.$$

Then for fixed n , we propose $\int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x)}{\varphi(x)} dx$ instead $\int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x) - \mathbb{E}\widehat{b}_n^2(x)}{\varphi(x)} dx$ or $\int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x) - \mathbb{E}b^2(x)}{\varphi(x)} dx$ to test normality. Finally, if $t = S_n x + \overline{X}_n$, then $\widehat{b}_n(x) = \sqrt{n} \left(F_n(t) - \phi\left(\frac{t - \overline{X}_n}{S_n}\right) \right)$, so $T_n^* = \int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x)}{\varphi(x)} dx = \frac{n}{S_n} \int_{\overline{X}_n - a_n S_n}^{\overline{X}_n + a_n S_n} \frac{\left(F_n(t) - \phi\left(\frac{t - \overline{X}_n}{S_n}\right) \right)^2}{\varphi\left(\frac{t - \overline{X}_n}{S_n}\right)} dt$. Then, estimating μ by $\overline{X}_n$ and σ by S_n , it follows that T_n^* can be seen as a kind of Cramér–von Mises statistic to test normality when the weight function is $\psi(t) = \frac{1}{\varphi^2(\frac{1}{\phi^{-1}(t)})}$ and integrated over the interval $(-a_n, a_n)$ instead of over all the real line.

4 Asymptotic distribution of the TCVM test statistic

In this section, given $X_1, X_2, \dots, X_n$ iid, we propose $\{T_n \geq c\}$ to test normality, where $T_n := \int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x) - \mathbb{E}b^2(x)}{\varphi(x)} dx$, and we will obtain the asymptotic distribution of T_n under H_0 . Since $\widehat{b}_n(x)$ is invariant under changes of location and scale, to test normality, we

may assume (if the null hypothesis is true) that $X_1, X_2, \dots, X_n$ are independent $N(0, 1)$. We can express $\widehat{b}_n(x)$ and $b_n(x)$ (defined in Section 1) as

$$\widehat{b}_n(x) = b_n(x_n) + \sqrt{n}(\phi(x_n) - \phi(x))$$

where $x_n = S_n x + \overline{X}_n$. Then, given x , $\widehat{b}_n(x)$ converges to $b(x)$ with $\{b(x)\}_{x \in \mathbb{R}}$ being the Brownian bridge associated with ϕ , the asymptotic limit of the processes $\{b_n(x)\}_{x \in \mathbb{R}}$. Moreover, Kalemkerian (2016) proved that

$$\int_{-a_n}^{a_n} \frac{\mathbb{E}(\widehat{b}_n^2(x))}{\varphi(x)} dx = \ln(\ln n) + c_n \text{ where } \{c_n\} \text{ converges.}$$

Hence, our test statistic goes to infinity. In Lemma 4 we give meaning to the expression $\int_{-\infty}^{+\infty} \frac{b^2(x) - \phi(x)(1 - \phi(x))}{\varphi(x)} dx$, therefore recalling that $\mathbb{E}(b^2(x)) = \phi(x)(1 - \phi(x))$, it is reasonable to assume that if we subtract from our statistic its expected value or an equivalent quantity, we should find a non-trivial asymptotic distribution of the sequence

$$T_n = \int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x) - \mathbb{E}(b^2(x))}{\varphi(x)} dx.$$

Indeed, in the following theorem, we summarize the main result of this section.

Theorem 1. T_n converges in law to

$$\int_{-\infty}^{+\infty} \frac{b^2(x) - \mathbb{E}(b^2(x))}{\varphi(x)} dx - \left(\int_{-\infty}^{+\infty} x db(x) \right)^2 - \left(\int_{-\infty}^{+\infty} x^2 db(x) \right)^2 + c,$$

where c is a constant and $\{b(x)\}_{x \in \mathbb{R}}$ is a ϕ -Brownian bridge.

To obtain this result, we use the Skorokhod construction (Skorokhod, 1956). Thus, there exists a triangular array of row independent random variables $X_{n1}, \dots, X_{nn}$ and $\{b(x)\}_{x \in \mathbb{R}}$ a ϕ -Brownian bridge, all defined on the same sample space, such that $\|b_n - b\| \xrightarrow{a.s.} 0$ where $\|f\| = \sup_{x \in \mathbb{R}} |f(x)|$ and $\{b_n(x)\}_{x \in \mathbb{R}}$ are the empirical processes associated to $X_{n1}, \dots, X_{nn}$. To simplify the notation, we will continue to write $X_1, \dots, X_n$ instead $X_{n1}, \dots, X_{nn}$ throughout the rest of this paper.

Theorem 1 follows from the following succession of lemmas and propositions.

We begin by noting that

$$\widehat{b}_n^2(x) = b_n^2(x_n) + n(\phi(x_n) - \phi(x))^2 + 2\sqrt{n}b_n(x_n)(\phi(x_n) - \phi(x)), \quad (1)$$

adding and subtracting the expression $\mathbb{E}(b_n^2(x_n))$ to the numerator of the integrand in T_n , we obtain the following decomposition

$$T_n = H_n + I_n + J_n \quad (2)$$

where

$$H_n := \int_{-a_n}^{a_n} \frac{b_n^2(x_n) - \mathbb{E}(b_n^2(x_n))}{\varphi(x)} dx,$$

$$I_n := \int_{-a_n}^{a_n} \frac{n(\phi(x_n) - \phi(x))^2 + 2\sqrt{n}b_n(x_n)(\phi(x_n) - \phi(x))}{\varphi(x)} dx,$$

$$J_n := \int_{-a_n}^{a_n} \frac{\mathbb{E}(b_n^2(x_n)) - \mathbb{E}(b^2(x))}{\varphi(x)} dx.$$

Lemma 1.

$$\frac{x\varphi(x)}{1+x^2} \leq \phi(-x) \leq \frac{\varphi(x)}{x} \text{ for all } x > 0. \quad (3)$$

Proof. $\varphi(x) = \int_x^{+\infty} t\varphi(t) dt \geq x \int_x^{+\infty} \varphi(t) dt = x(1 - \phi(x))$. Then, the second inequality in (3) is true. For the other inequality, observe that $1 - \phi(x) = \int_x^{+\infty} \varphi(t) dt = \int_x^{+\infty} t\varphi(t) \frac{1}{t} dt$. If we integrate by parts, we obtain that $1 - \phi(x) = \frac{\varphi(x)}{x} - \int_x^{+\infty} \frac{\varphi(t)}{t^2} dt \geq \frac{\varphi(x)}{x} - \frac{1-\phi(x)}{x^2}$, then $1 - \phi(x) \geq \frac{x\varphi(x)}{1+x^2}$. $\square$

Lemma 2. $\frac{a_n^p}{n^q} \rightarrow 0$ for all $p, q > 0$.

Proof. From the previous lemma, we obtain that if $x \rightarrow +\infty$, then $\phi(-x) \sim \frac{\varphi(x)}{x}$, where $f(x) \sim g(x)$ means that $\frac{f(x)}{g(x)} \rightarrow 1$ when $x \rightarrow +\infty$. Then

$$\frac{a_n^p}{n^q} = \phi^q(-a_n) a_n^p \sim \varphi^q(a_n) a_n^{p-1} \rightarrow 0.$$

$\square$

Lemma 3. For any $\{b(x)\}_{x \in \mathbb{R}}$ ϕ -Brownian bridge, it holds that

$$\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{\mathbb{COV}(b^2(x); b^2(y))}{\varphi(x)\varphi(y)} dx dy < +\infty. \quad (4)$$

Proof. We use that if (X, Y) are centered Gaussian random variables in $\mathbb{R}^2$, then $\mathbb{COV}(X^2, Y^2) = 2(\mathbb{COV}(X, Y))^2$. Then

$$\begin{aligned} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{\mathbb{COV}(b^2(x); b^2(y))}{\varphi(x)\varphi(y)} dx dy &= 2 \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{\phi^2(x \wedge y) (1 - \phi(x \vee y))^2}{\varphi(x)\varphi(y)} dx dy = \\ &= 4 \int_{-\infty}^{+\infty} dx \int_{-\infty}^x \frac{\phi^2(y) (1 - \phi(x))^2}{\varphi(x)\varphi(y)} dy = 4 \int_{-\infty}^{+\infty} \frac{(1 - \phi(x))^2}{\varphi(x)} dx \int_{-\infty}^x \frac{\phi^2(y)}{\varphi(y)} dy. \end{aligned}$$

If $x \rightarrow -\infty$, then

$$\begin{aligned} \frac{(1 - \phi(x))^2}{\varphi(x)} \int_{-\infty}^x \frac{\phi^2(y)}{\varphi(y)} dy &\leq \frac{(1 - \phi(x))^2}{\varphi(x)} \int_{-\infty}^x \frac{\varphi(y)}{y^2} dy \leq \\ &\leq \frac{1}{\varphi(x)} \frac{\phi(x)}{x^2} \leq \frac{1}{|x^3|}. \end{aligned}$$

If $x \rightarrow +\infty$, then

$$\frac{(1 - \phi(x))^2}{\varphi(x)} \int_{-\infty}^x \frac{\phi^2(y)}{\varphi(y)} dy \leq \frac{(1 - \phi(x))^2}{\varphi(x)} \left[\int_{-\infty}^1 \frac{\phi^2(y)}{\varphi(y)} dy + \int_1^x \frac{\phi^2(y)}{\varphi(y)} dy \right].$$

To complete the proof, it is enough to observe that

$$\begin{aligned} \frac{(1 - \phi(x))^2}{\varphi(x)} \int_1^x \frac{\phi^2(y)}{\varphi(y)} dy &\leq \frac{\varphi(x)}{x^2} \int_1^x \frac{dy}{\varphi(y)} \leq \\ \frac{\varphi(x)}{x^2} \int_1^x \frac{y dy}{\varphi(y)} &= \frac{1 - e^{-(x^2-1)/2}}{x^2} \leq \frac{1}{x^2}. \end{aligned}$$

□

Lemma 4. Define the sequence

$$W_n := \int_{-a_n}^{a_n} \frac{b^2(x) - \mathbb{E}(b^2(x))}{\varphi(x)} dx = \int_{-a_n}^{a_n} \frac{b^2(x) - \phi(x)(1 - \phi(x))}{\varphi(x)} dx. \quad (5)$$

Then exists a random variable Y in L^2 such that $\mathbb{E}(W_n - Y)^2 \rightarrow 0$.

Proof. We will demonstrate that $\{W_n\}$ is an L^2 -Cauchy sequence.

$$\begin{aligned} \mathbb{E}(W_n W_m) &= \int_{-a_n}^{a_n} \int_{-a_m}^{a_m} \mathbb{E} \left(\frac{b^2(x) - \phi(x)(1 - \phi(x))}{\varphi(x)} \frac{b^2(y) - \phi(y)(1 - \phi(y))}{\varphi(y)} \right) dx dy = \\ &\int_{-a_n}^{a_n} \int_{-a_m}^{a_m} \frac{\mathbb{E}(b^2(y)b^2(x)) - \phi(x)\phi(y)(1 - \phi(x))(1 - \phi(y))}{\varphi(x)\varphi(y)} dx dy \end{aligned}$$

which goes to

$$\int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{\mathbb{C}\text{OV}(b^2(y)b^2(x))}{\varphi(x)\varphi(y)} dx dy < +\infty$$

for $n, m \rightarrow +\infty$.

Then $\mathbb{E}(W_n - W_m)^2 = \mathbb{E}(W_n^2 + W_m^2 - 2W_n W_m) \rightarrow 0$, and $\{W_n\}$ is an L^2 -Cauchy sequence. □

From now, we call $Y := \int_{-\infty}^{+\infty} \frac{b^2(x) - \phi(x)(1 - \phi(x))}{\varphi(x)} dx$.

Lemma 5. There exist Z_1 and Z_2 centered, independent Gaussian random variables such that

$$(\sqrt{n}\bar{X}_n, \sqrt{n}(S_n - 1)) \xrightarrow{P} (Z_1, Z_2).$$

Proof. We define $Y_n := \sup_{x \in \mathbb{R}} \frac{|b_n(x) - b(x)|}{\sqrt{\varphi(x)}}$, and we will apply theorem 1.1 of Shorack & Wellner (1982) to $q(t) = \sqrt{\varphi(\phi^{-1}(t))}$. Set $\|f\| = \sup_{x \in [0,1]} |f(x)|$, and $B_n(t) := b_n(\phi^{-1}(t))$, $B(t) := b(\phi^{-1}(t))$. Then $Y_n = \left\| \frac{B_n - B}{q} \right\|$. Using Lemma 2, that $\frac{q^2(t)}{t \ln(-\ln t)} \rightarrow +\infty$ for $t \rightarrow 0$, then $Y_n \xrightarrow{P} 0$.

$\int_{-\infty}^{+\infty} |x|^\alpha |b_n(x) - b(x)| dx \leq Y_n \int_{-\infty}^{+\infty} |x|^\alpha \sqrt{\varphi(x)} dx \xrightarrow{P} 0$ for all $\alpha > 0$. Then,

$$\sqrt{n} \bar{X}_n = \sqrt{n} \int_{-\infty}^{+\infty} x d(F_n(x) - \phi(x)) = \int_{-\infty}^{+\infty} x db_n(x) \xrightarrow{P} \int_{-\infty}^{+\infty} x db(x) := Z_1.$$

$$\sqrt{n}(S_n - 1) = \frac{\sqrt{n}(S_n^2 - 1)}{S_n + 1} = \frac{1}{S_n + 1} \left(\int_{-\infty}^{+\infty} x^2 db_n(x) - \sqrt{n} \bar{X}_n^2 \right) \xrightarrow{P} \frac{1}{2} \int_{-\infty}^{+\infty} x^2 db(x) := Z_2.$$

Observe that Z_1, Z_2 are centered Gaussian random variables, and $\mathbb{V}(Z_1) = 1$, $\mathbb{V}(Z_2) = 1/2$.

As $\sqrt{n} \bar{X}_n$ and S_n are independent for all n , it follows that Z_1 and Z_2 are independent too. $\square$

Lemma 6.

$$\mathbb{E}(b_n^2(x)b_n^2(y)) = \mathbb{E}(b^2(x)b^2(y)) + \frac{p(x,y)}{n}$$

where

$$p(x,y) = \phi(x \wedge y)(1 - \phi(x \vee y)) \frac{1 + 6\phi(x \vee y)\phi(x \wedge y) - 2\phi(x \vee y) - 4\phi(x \wedge y)}{n}$$

and $\{b(x)\}_{x \in \mathbb{R}}$ is any ϕ -Brownian motion.

Proof. For simplicity in notation, we will call $z = \phi(x \wedge y)$ and $t = \phi(x \vee y)$.

$$\mathbb{E}(b_n^2(x)b_n^2(y)) = n^2 \mathbb{E}(F_n(x) - \phi(x))^2 (F_n(y) - \phi(y))^2 =$$

$$n^2 \mathbb{E}(F_n^2(x) - 2F_n(x)\phi(x) + \phi^2(x)) (F_n^2(y) - 2F_n(y)\phi(y) + \phi^2(y)) \quad (6)$$

To calculate (6) we will use that $\mathbb{E}(F_n(x)) = \phi(x)$ and $\mathbb{E}(F_n^2(x)) = \frac{\phi(x)(1-\phi(x))}{n} + \phi^2(x)$ for all x , and the formulas for $\mathbb{E}(F_n(x)F_n(y))$, $\mathbb{E}(F_n^2(x)F_n(y))$ and $\mathbb{E}(F_n^2(x)F_n^2(y))$ that we will obtain. First, we observe that

$$n^2 \mathbb{E}(F_n(x)F_n(y)) = \sum_{i,j=1}^n P(X_i \leq x, X_j \leq y) = nz + n(n-1)zt. \quad (7)$$

If we define $I := -2\phi(y) \mathbb{E}(F_n^2(x)\phi(y)) - 2\phi(x) \mathbb{E}(F_n^2(y)\phi(x))$ then (6) (after some calculation) is equal to

$$n^2 \mathbb{E}F_n^2(x)F_n^2(y) + n^2 I + 5nz^2t + nzt^2 + (3n^2 - 6n)z^2t^2.$$

If we proceed similarly to (7), we obtain that

$$n^2 \mathbb{E}(F_n^2(x)F_n(y)) = \frac{1}{n} \sum_{i,j,k=1}^n P(X_i \leq x, X_j \leq x, X_k \leq y) =$$

$$(n-1)(n-2)\phi^2(x)\phi(y) + (n-1)\phi(x)\phi(y) + 2(n-1)\phi(x\wedge y)\phi(x) + \phi(x\wedge y),$$

and

$$n^2\mathbb{E}(F_n^2(x)F_n^2(y)) = \frac{1}{n^2} \sum_{i,j,k,l=1}^n P(X_i \leq x, X_j \leq x, X_k \leq y, X_l \leq y)$$

$$\frac{n-1}{n} (4z^2 + 3zt + 5(n-2)z^2t + (n-2)zt^2 + (n^2 - 5n + 6)z^2t^2) + z/n$$

Then

$$n^2I = -4(n^2 - 3n + 2)z^2t^2 + (10 - 10n)z^2t + (2 - 2n)zt^2 - 2z^2 - 2zt.$$

Finally, $\mathbb{E}(b_n^2(x)b_n^2(y))$ is equal to

$$\begin{aligned} & -5tz^2 - t^2z + 3t^2z^2 + tz + 2z^2 + z(t-1) \frac{2t + 4z - 6tz - 1}{n} = \\ & \mathbb{E}(b^2(x)b^2(y)) + z(t-1) \frac{2t + 4z - 6tz - 1}{n}. \end{aligned} \quad (8)$$

where $\{b(x)\}_{x \in \mathbb{R}}$ is any ϕ -Brownian motion. $\square$

Lemma 7.

$$\int_{-a_n}^{a_n} \frac{b_n^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx \xrightarrow{P} \int_{-\infty}^{+\infty} \frac{b^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx. \quad (9)$$

Proof. We decompose

$$\int_{-a_n}^{a_n} \frac{b_n^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx := Y_{n,\delta} + Z_{n,\delta} + W_{n,\delta}$$

where $Y_{n,\delta} := \int_{-\delta}^{\delta} \frac{b_n^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx$, $Z_{n,\delta} := \int_{-a_n}^{-\delta} \frac{b_n^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx$ and $W_{n,\delta} := \int_{\delta}^{a_n} \frac{b_n^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx$. We will prove that $\lim_{\delta \rightarrow +\infty} \lim_{n \rightarrow +\infty} Y_{n,\delta} = Y$, $\lim_{\delta \rightarrow +\infty} \lim_{n \rightarrow +\infty} Z_{n,\delta} = 0$ and $\lim_{\delta \rightarrow +\infty} \lim_{n \rightarrow +\infty} W_{n,\delta} = 0$ in probability.

Indeed

$$\begin{aligned} \lim_{\delta \rightarrow +\infty} \lim_{n \rightarrow +\infty} Y_{n,\delta} &= \lim_{\delta \rightarrow +\infty} \int_{-\delta}^{\delta} \frac{b^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx = \\ &= \int_{-\infty}^{+\infty} \frac{b^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx. \end{aligned}$$

To prove that $\lim_{\delta \rightarrow +\infty} \lim_{n \rightarrow +\infty} W_{n,\delta} = 0$ in probability, it is enough to show that $\lim_{\delta \rightarrow +\infty} \lim_{n \rightarrow +\infty} \mathbb{E}(W_{n,\delta}^2) = 0$.

$$\mathbb{E}(W_{n,\delta}^2) = \mathbb{E}\left(\int_{\delta}^{a_n} \frac{b_n^2(x) - \phi(x)(1 - \phi(x))}{\phi(x)} dx \int_{\delta}^{a_n} \frac{b_n^2(y) - \phi(y)(1 - \phi(y))}{\phi(y)} dy\right) =$$

$$\int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{\mathbb{E}(b_n^2(x)b_n^2(y)) - \phi(x)(1-\phi(x))\phi(y)(1-\phi(y))}{\varphi(x)\varphi(y)} dx dy \quad (10)$$

Then, using Lemma 6 we obtain that (10) is equal to

$$\int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{\mathbb{COV}(b^2(x), b^2(y))}{\varphi(x)\varphi(y)} dx dy + \frac{1}{n} \int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{p(x, y)}{\varphi(x)\varphi(y)} dx dy.$$

From Lemma 3 we obtain that

$$\lim_{\delta \rightarrow +\infty} \lim_{n \rightarrow +\infty} \int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{\mathbb{COV}(b^2(x), b^2(y))}{\varphi(x)\varphi(y)} dx dy = 0.$$

Finally, we will show that $\lim_{n \rightarrow +\infty} \frac{1}{n} \int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{p(x, y)}{\varphi(x)\varphi(y)} dx dy = 0$ for all $\delta > 0$.
Indeed, for certain $c > 0$,

$$\frac{1}{n} \int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{|p(x, y)|}{\varphi(x)\varphi(y)} dx dy \leq \frac{c}{n} \int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{\phi(x \wedge y)(1 - \phi(x \vee y))}{\varphi(x)\varphi(y)} dx dy.$$

Then, applying L'Hôpital rule, we obtain that

$$\begin{aligned} & \lim_{n \rightarrow +\infty} \frac{1}{n} \int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{\phi(x \wedge y)(1 - \phi(x \vee y))}{\varphi(x)\varphi(y)} dx dy = \\ & \lim_{n \rightarrow +\infty} \frac{\partial}{\partial n} \left(\int_{\delta}^{a_n} \int_{\delta}^{a_n} \frac{\phi(x \wedge y)(1 - \phi(x \vee y))}{\varphi(x)\varphi(y)} dx dy \right) = \\ & \lim_{n \rightarrow +\infty} \frac{2}{n^2 \varphi^2(a_n)} \int_{\delta}^{a_n} \frac{\phi(x)(1 - \phi(a_n))}{\varphi(x)} dx = \lim_{n \rightarrow +\infty} \frac{2}{n^3 \varphi^2(a_n)} \int_{\delta}^{a_n} \frac{\phi(x)}{\varphi(x)} dx = 0. \end{aligned}$$

The case $\lim_{\delta \rightarrow +\infty} \lim_{n \rightarrow +\infty} Z_{n, \delta} = 0$ is solved similarly. □

In the next lemmas and propositions, we will use the following equalities:

$$\phi(x_n) - \phi(x) = \varphi(c_n)(x_n - x) \text{ for } c_n \in (x_n \wedge x, x_n \vee x). \quad (11)$$

$$\frac{\varphi^2(c_n)}{\varphi^2(x)} = e^{x^2 - c_n^2} = 1 + e^{d_n}(x^2 - c_n^2) \text{ for } |d_n| \leq |x^2 - c_n^2|. \quad (12)$$

$$\frac{\varphi(c_n)}{\varphi(x)} = e^{\frac{x^2 - c_n^2}{2}} = 1 + e^{d'_n} \frac{(x^2 - c_n^2)}{2} \text{ for } |d'_n| \leq \frac{|x^2 - c_n^2|}{2}. \quad (13)$$

Observe that if $x \in (-a_n, a_n)$, then $|d_n| \leq |x^2 - x_n^2| \leq \frac{A a_n^2}{\sqrt{n}}$ for certain random variable A , and $|d'_n| \leq \frac{|x^2 - x_n^2|}{2} \leq \frac{A' a_n^2}{\sqrt{n}}$ for certain random variable A' .

In the next proposition, we prove that the term H_n in decomposition (2) converges in law.

Proposition 1.

$$H_n = \int_{-a_n}^{a_n} \frac{b_n^2(x_n) - \mathbb{E}(b_n^2(x_n))}{\varphi(x)} dx \xrightarrow{P} \int_{-\infty}^{+\infty} \frac{b^2(x) - \mathbb{E}(b^2(x))}{\varphi(x)} dx.$$

Proof. Since

$$\int_{-a_n}^{a_n} \frac{b_n^2(x_n) - \mathbb{E}(b_n^2(x_n))}{\varphi(x_n)} dx = \frac{1}{S_n} \int_{-S_n a_n + \bar{X}_n}^{S_n a_n + \bar{X}_n} \frac{b_n^2(t) - \mathbb{E}(b_n^2(t))}{\varphi(t)} dt$$

converges in probability (by Lemma 7) to $\int_{-\infty}^{+\infty} \frac{b^2(x) - \mathbb{E}(b^2(x))}{\varphi(x)} dx$,

$$H_n = \int_{-a_n}^{a_n} \frac{b_n^2(x_n) - \mathbb{E}(b_n^2(x_n))}{\varphi(x_n)} \frac{\varphi(x_n)}{\varphi(x)} dx$$

and due to (13) (with $c_n = x_n$) and Lemma 3 we obtain that H_n converges in probability to $\int_{-\infty}^{+\infty} \frac{b^2(x) - \mathbb{E}(b^2(x))}{\varphi(x)} dx$. $\square$

Lemma 8.

$$\int_{-a_n}^{a_n} \frac{n(\phi(x_n) - \phi(x))^2}{\varphi(x)} dx = n(S_n - 1)^2 \alpha_n + n\bar{X}_n^2 \beta_n + Y_n$$

where $\alpha_n = \int_{-a_n}^{a_n} x^2 \varphi(x) dx$, $\beta_n = \int_{-a_n}^{a_n} \varphi(x) dx$ and $\mathbb{E}(|Y_n|) \rightarrow 0$.

Proof. Using (11) and (12), we obtain that

$$\begin{aligned} \int_{-a_n}^{a_n} \frac{n(\phi(x_n) - \phi(x))^2}{\varphi(x)} dx &= \int_{-a_n}^{a_n} \frac{n\varphi^2(c_n)(x_n - x)^2}{\varphi^2(x)} \varphi(x) dx = \\ &= \int_{-a_n}^{a_n} n(x_n - x)^2 \varphi(x) dx + \int_{-a_n}^{a_n} n e^{d_n} (x^2 - c_n^2) (x_n - x)^2 \varphi(x) dx. \\ \int_{-a_n}^{a_n} n(x_n - x)^2 \varphi(x) dx &= \int_{-a_n}^{a_n} n((S_n - 1)x + \bar{X}_n)^2 \varphi(x) dx = \\ &= n(S_n - 1)^2 \alpha_n + n\bar{X}_n^2 \beta_n. \end{aligned}$$

Define $Y_n := \int_{-a_n}^{a_n} n e^{d_n} (x^2 - c_n^2) (x_n - x)^2 \varphi(x) dx$. Then, using that $|d_n| \leq \frac{A a_n^2}{\sqrt{n}}$ and Lemma 3,

$$\mathbb{E}(|Y_n|) \leq \mathbb{E} \left(\int_{-a_n}^{a_n} n e^{d_n} |x^2 - c_n^2| (x_n - x)^2 \varphi(x) dx \right) \leq \frac{a_n^2}{\sqrt{n}} \mathbb{E} \left(A e^{\frac{A a_n^2}{\sqrt{n}}} \right) \rightarrow 0.$$

$\square$

Lemma 9.

$$\int_{-a_n}^{a_n} \frac{n(\phi(x_n) - \phi(x))^2}{\varphi(x)} dx \xrightarrow{P} Z_1^2 + Z_2^2.$$

Proof. This follows directly from Lemma 8 and Lemma 5. $\square$

Lemma 10.

$$\int_{-a_n}^{a_n} \frac{\sqrt{n}b_n(x_n)(\phi(x_n) - \phi(x))}{\varphi(x)} dx = \sqrt{n} \frac{(S_n - 1)}{S_n^2} Z'_n + \sqrt{n} \frac{\bar{X}_n}{S_n^2} Z''_n + Y'_n$$

where $Z'_n = \int_{\bar{X}_n - a_n S_n}^{\bar{X}_n + a_n S_n} tb_n(t) dt$, $Z''_n = \int_{\bar{X}_n - a_n S_n}^{\bar{X}_n + a_n S_n} b_n(t) dt$ and $\mathbb{E}(|Y'_n|) \rightarrow 0$.

Proof. Using (11) and (13), observe that

$$\begin{aligned} \int_{-a_n}^{a_n} \frac{\sqrt{n}b_n(x_n)(\phi(x_n) - \phi(x))}{\varphi(x)} dx &= \int_{-a_n}^{a_n} \frac{\sqrt{n}b_n(x_n)\varphi(c_n)(x_n - x)}{\varphi(x)} dx = \\ &= \int_{-a_n}^{a_n} \sqrt{n}b_n(x_n)(x_n - x) dx + \frac{1}{2} \int_{-a_n}^{a_n} \sqrt{n}b_n(x_n)e^{d'_n}(x^2 - c_n^2)(x_n - x) dx. \\ \sqrt{n} \int_{-a_n}^{a_n} b_n(x_n)(x_n - x) dx &= \frac{\sqrt{n}}{S_n^2} \int_{-S_n a_n + \bar{X}_n}^{S_n a_n + \bar{X}_n} b_n(t) [(S_n - 1)t + \bar{X}_n] dt = \\ &= \frac{\sqrt{n}(S_n - 1)}{S_n^2} \int_{-S_n a_n + \bar{X}_n}^{S_n a_n + \bar{X}_n} tb_n(t) dt + \frac{\sqrt{n}\bar{X}_n}{S_n^2} \int_{-S_n a_n + \bar{X}_n}^{S_n a_n + \bar{X}_n} b_n(t) dt. \end{aligned}$$

Define $Y'_n := \frac{1}{2} \int_{-a_n}^{a_n} \sqrt{n}b_n(x_n)e^{d'_n}(x^2 - c_n^2)(x_n - x) dx$. Then working similarly to Lemma 8, we obtain that $\mathbb{E}(|Y'_n|) \rightarrow 0$. $\square$

Lemma 11.

$$\int_{-a_n}^{a_n} \frac{\sqrt{n}b_n(x_n)(\phi(x_n) - \phi(x))}{\varphi(x)} dx \xrightarrow{P} -Z_1^2 - Z_2^2.$$

Proof.

$$Z'_n = \int_{\bar{X}_n - a_n S_n}^{\bar{X}_n + a_n S_n} tb_n(t) dt = \int_{-\infty}^{+\infty} tb_n(t) dt + \varepsilon_n$$

where $\varepsilon_n \xrightarrow{P} 0$.

Integrating by parts and using Lemma 5, we obtain that

$$Z'_n = -\frac{1}{2} \int_{-\infty}^{+\infty} t^2 db_n(t) + \varepsilon_n \xrightarrow{P} -Z_2.$$

Analogously $Z''_n \xrightarrow{P} -Z_1$, and the result of the lemma it follows directly from Lemma 10. $\square$

Proposition 2.

$$I_n = \int_{-a_n}^{a_n} \frac{n(\phi(x_n) - \phi(x))^2 + 2\sqrt{n}b_n(x_n)(\phi(x_n) - \phi(x))}{\varphi(x)} dx \xrightarrow{P} -Z_1^2 - Z_2^2.$$

Proof. This follows directly from lemmas 9 and 11. $\square$

Finally, to prove that the term J_n in decomposition (2) converges, we need the following lemma.

Lemma 12. *The sequence*

$$\int_{-a_n}^{a_n} \frac{\mathbb{E}(\widehat{b}_n^2(x)) - \mathbb{E}(b^2(x))}{\varphi(x)} dx$$

converges.

Proof. By symmetry it is enough to take the integral between 1 and a_n . By using the results in section 3 of Kalemkerian (2016), one gets

$$\begin{aligned} \int_1^{a_n} \frac{\mathbb{E}(\widehat{b}_n^2(x)) - \mathbb{E}(b^2(x))}{\varphi(x)} dx &= \frac{1}{2} \ln \ln n + c_n - \int_1^{a_n} \frac{\phi(x)(1 - \phi(x))}{\varphi(x)} dx = \\ &= \int_1^{a_n} \left(\frac{1}{x} - \frac{\phi(x)(1 - \phi(x))}{\varphi(x)} \right) dx + c'_n. \end{aligned}$$

where $\{c_n\}$ and $\{c'_n\}$ are convergent sequences.

Define $f(x) := \frac{1 - \phi(x)}{\varphi(x)} - \frac{1}{x}$, and apply L'Hôpital rule twice to show

$$\frac{f(x)}{\frac{1}{x^2}} \xrightarrow{x \rightarrow +\infty} 0.$$

$\square$

Proposition 3. *The sequence*

$$J_n = \int_{-a_n}^{a_n} \frac{\mathbb{E}(b_n^2(x_n)) - \mathbb{E}(b^2(x))}{\varphi(x)} dx$$

converges.

Proof. We will see that

$$\int_{-a_n}^{a_n} \frac{\mathbb{E}(b_n^2(x_n)) - \mathbb{E}(\widehat{b}_n^2(x))}{\varphi(x)} dx \text{ and } \int_{-a_n}^{a_n} \frac{\mathbb{E}(\widehat{b}_n^2(x)) - \mathbb{E}(b^2(x))}{\varphi(x)} dx \text{ are convergent.}$$

The second term converges due to Lemma 12. Observe that from (2) one gets

$$\int_{-a_n}^{a_n} \frac{\mathbb{E}(b_n^2(x_n)) - \mathbb{E}(\widehat{b}_n^2(x))}{\varphi(x)} dx = -\mathbb{E}(I_n) \rightarrow 3/2 \quad (14)$$

due to lemmas 8 and 10. $\square$

Note that the constant appearing in Theorem 1 is unknown. As seen in Lemma 12 and Proposition 3, one gets

$$c = \lim J_n = \lim \int_{-a_n}^{a_n} \frac{\mathbb{E}(\widehat{b}_n^2(x)) - \mathbb{E}(b^2(x))}{\varphi(x)} dx + 3/2.$$

In Table 2, we present the results, from $m = 1000$ simulations, the estimated values of the constant c for different values of n .

Table 2. Estimated values of constant c for different values of n .

n	500	1000	10000	20000	50000	100000
c	0.0258	0.009137	-0.022793	-0.01752	0.002268	0.00093

5 Conclusions

This paper has presented a test of normality with μ and σ unknown, based on the statistic $\int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x)}{\varphi(x)} dx$, which is a type of Cramér–von Mises statistic, except that it is integrated over a particular interval, which is a function of the sample size. This statistic was inspired by del Barrio et al. (1999). We analysed its asymptotic behavior, which has a limit distribution equivalent to that of S–W statistic. A comparative study of its behaviour over a wide range of alternative hypotheses found that the test proposed here is often better than the tests of Shapiro–Wilk and Anderson–Darling, which are two tests that have very good performance as tests of normality.

6 Appendix 1. Performance of the proposed test

We present a table, for sample size of $n = 50$, where we compare the performance of our test, namely the Truncated Cramér–von Mises test (TCVM), with the following four tests: 1) the non-truncated Cramér–von Mises test, which integrates over all the real line (CVM); 2) the test proposed by del Barrio, Matrán-Cuesta, and Rodríguez (BCMR); 3) the Anderson–Darling test (AD) and 4) the Shapiro–Wilk test (SW).

The power of these tests was compared based on 10,000 replications. We chose the level of significance $\alpha = 0.05$ and the empirical critical values were obtained after 50,000 replications.

The list of 35 alternative hypotheses considered in the present section are borrowed from Gan and Koelher (1990) and are a summary of a total of 69.

We use the notation of Gan and Koelher (1990), and the alternative hypotheses are considered in the same order as in the cited reference.

LoConN(p, a) indicates a mixture of $N(0; 1)$ with probability $1 - p$ and $N(a; 1)$ with probability p . ScConN(p, a) indicates the mixture of $N(0; 1)$ with probability $1 - p$ and $N(0; a)$ with probability p . TruncN($a; b$) is $N(0; 1)$ truncated to the interval $(a; b)$. SB($a; b$) is the Johnson bounded distribution with parameters a, b , and SU($a; b$) is the Johnson unbounded distribution with parameters a, b . Triangle I(a) is the distribution with density $f(x) = 1/a - |x|/a^2$ for $|x| \leq a$, and Triangle II(a) is the distribution with density

Table 1. Powers under 35 alternative hypotheses at the 5% level, $n = 50$.

Set	N ^o	Alternative	TCVM	CVM	BCMR	AD	SW
1	1	LoConN (0.5; 4)	0.935	0.432	0.883	0.956	0.783
	2	LoConN (0.5; 3)	0.439	0.044	0.341	0.480	0.212
	3	LoConN (0.5; 2)	0.084	0.009	0.053	0.093	0.033
2	4	SB (0; 0.5)	0.958	0.496	0.957	0.926	0.880
	5	Unif (0; 1)	0.708	0.124	0.689	0.616	0.466
	6	SB (0, 0.707)	0.553	0.063	0.508	0.495	0.309
	7	TruncN (−1; 1)	0.876	0.375	0.735	0.350	0.197
	8	Beta (2; 2)	0.163	0.005	0.117	0.155	0.051
	9	Triangle I (1)	0.061	0.002	0.034	0.055	0.015
3	10	t (10)	0.098	0.199	0.169	0.113	0.186
	11	Logist (0; 1)	0.123	0.248	0.204	0.156	0.243
4	12	ScConN (0.05; 3)	0.063	0.114	0.099	0.071	0.116
	13	ScConN (0.05; 5)	0.114	0.243	0.207	0.142	0.234
5	14	ScConN (0.1; 5)	0.168	0.351	0.298	0.202	0.340
	15	ScConN (0.1; 7)	0.297	0.518	0.467	0.337	0.494
6	16	ScConN (0.2; 3)	0.099	0.228	0.189	0.126	0.216
	17	ScConN (0.2; 7)	0.426	0.649	0.596	0.306	0.464
7	18	Laplace (0; 1)	0.455	0.562	0.539	0.526	0.581
	19	SU (0; 1)	0.688	0.788	0.770	0.752	0.808
	20	t (2)	0.818	0.881	0.871	0.854	0.892
8	21	Beta (2; 1)	0.815	0.310	0.811	0.750	0.702
	22	TruncN (−2; 1)	0.642	0.242	0.549	0.620	0.470
	23	Beta (3; 2)	0.227	0.022	0.177	0.007	0.095
9	24	SB (1; 2)	0.105	0.052	0.094	0.089	0.072
	25	Weibull (2)	0.344	0.237	0.394	0.393	0.355
	26	HalfN (0; 1)	0.891	0.665	0.922	0.815	0.883
10	27	LoConN (0.2; 3)	0.668	0.369	0.606	0.589	0.554
	28	LoConN (0.2; 5)	0.999	0.994	0.999	0.865	0.989
11	29	LoConN (0.1; 3)	0.489	0.487	0.559	0.610	0.569
	30	LoConN (0.1; 5)	0.960	0.988	0.986	0.988	0.989
12	31	LoConN (0.05; 3)	0.263	0.427	0.402	0.429	0.418
	32	LoConN (0.05; 5)	0.758	0.894	0.882	0.878	0.888
13	33	Triangle II (1)	0.818	0.314	0.809	0.499	0.689
	34	χ_4^2	0.916	0.818	0.945	0.921	0.927
	35	Lognormal (0; 1)	0.999	0.999	0.999	1.000	1.000

$f(x) = 2/a - 2x/a^2$ for $0 \leq x \leq a$.

For others values of n , the comparison was similar to the one presented here. The 35 alternative hypotheses are ordered in 13 types of distributions. Type 1 includes symmetric bimodal distributions with low skewness. Type 2 includes symmetric multimodal distributions with low tails. Type 3 includes distributions with slightly heavier tails than

the normal distribution. Types 4, 5 and 6 are contaminations of the normal distribution. Type 7 includes symmetrical distributions with high skewness. The remaining types are asymmetrical distributions. Types 8 and 9 are distributions with low skewness with opposite signs. Types 10, 11 and 12 are bimodal distributions with positive coefficients of symmetry. Type 13 contains distributions with extreme values of coefficients of skewness or symmetry. In the following graphs, we compare the powers (at the 5% level) of our test (TCVM), Anderson–Darling (AD) and Shapiro–Wilk (SW), for sample sizes $n = 20$ and $n = 50$, when the alternative hypotheses are in the lambda Tukey family. The critical values were obtained by simulations, with 50,000 replications. The various powers, presented in the graphs, were found by simulations, with 10,000 replications.




Figure 1: Tukey(d) alternative, $n = 20$ and $n = 50$.

In Figure 1, the alternative hypotheses belong to Tukey's family, for sample sizes $n = 20$ and $n = 50$. In Figure 2, the alternative hypotheses belong to the LoConN(0.3, p) family, for $n = 20$ and $n = 50$. We see that for distribution Type 1, the TCVM powers are close to the AD that are the best. For distribution Type 2, TCVM is the test with the best performance, and CVM performs badly. In Types 3, 4, 5 and 6, the performance is the opposite (CVM is the best test while TCVM performs badly). We can see that in many cases, TCVM is better than the others. Besides, in several of the cases in which TCVM performs badly, CVM is competitive with SW, AD and BCMR. Taking into account all 35 alternative hypotheses, we can see that in many cases, TCVM is better than the others, but under some alternative hypotheses, like symmetrical distributions with high skewness (Type 7), and contaminations in the variance of normal distributions (Types 4, 5 and 6), TCVM performs badly. However, in Figure 2 (contaminations in the mean of normal distributions), TCVM performs very well. It should be noted that in most cases, either TCVM or CVM results in the best test.




Figure 2: LoConN(0.3, p) alternative, $n = 20$ and $n = 50$.

7 Appendix 2. Computing the statistic T_n^* .

$$T_n^* = \int_{-a_n}^{a_n} \frac{\widehat{b}_n^2(x)}{\varphi(x)} dx = \int_{-a_n}^{a_n} \frac{\frac{1}{n} \left(\sum_{i=1}^n \mathbf{1}_{\{Y_i \leq x\}} \right)^2 - 2\phi(x) \sum_{i=1}^n \mathbf{1}_{\{Y_i \leq x\}} + n\phi^2(x)}{\varphi(x)} dx =$$

$$\frac{1}{n} \int_{-a_n}^{a_n} \frac{\left(\sum_{i=1}^n \mathbf{1}_{\{Y_i \leq x\}} \right)^2}{\varphi(x)} dx - 2 \int_{-a_n}^{a_n} \frac{\phi(x) \sum_{i=1}^n \mathbf{1}_{\{Y_i \leq x\}}}{\varphi(x)} dx + n \int_{-a_n}^{a_n} \frac{\phi^2(x)}{\varphi(x)} dx.$$

Then, given $X_1, X_2, \dots, X_n$, the calculation of the statistic T_n^* can be summarized in the following steps.

Step 1. Compute $\bar{X}_n$, S_n and delete all data such that $X_i \leq \bar{X}_n - a_n S_n$ or $X_i \geq \bar{X}_n + a_n S_n$ where the values of a_n are shown in ninth column of Appendix 2.

Step 2. Define $k := \# \{j : X_j \leq \bar{X}_n - a_n S_n\}$ and call $\tilde{X}_1, \tilde{X}_2, \dots, \tilde{X}_m$ to the values that resulted from step 1, and sort them: $\tilde{X}_{1:m} \leq \tilde{X}_{2:m} \leq \dots \leq \tilde{X}_{m:m}$.

Step 3. Compute $\tilde{Y}_0 = -a_n$, $\tilde{Y}_{m+1} = a_n$, $\tilde{Y}_j = \frac{\tilde{X}_{j:m} - \bar{X}_n}{S_n}$ for $j = 1, 2, \dots, m$.

Step 4. Compute $A_j = \int_{\tilde{Y}_j}^{\tilde{Y}_{j+1}} \frac{dx}{\varphi(x)}$ and $B_j = \int_{\tilde{Y}_j}^{\tilde{Y}_{j+1}} \frac{\phi(x) dx}{\varphi(x)}$ for $j = 0, 1, 2, \dots, m$ (for example in *R* you can use the “integrate” function to compute A_j and B_j values).

Step 5. Compute $T_n^* = \frac{1}{n} \sum_{j=0}^m (j+k)^2 A_j - 2 \sum_{j=0}^m (j+k) B_j + C_n$, where $C_n := n \int_{-a_n}^{a_n} \frac{\phi^2(x)}{\varphi(x)} dx$ and their values for different sample sizes are shown in the last column of Appendix 2.

Step 6. The critical values for rejection the null hypothesis for different levels of significance are in Appendix 3. When T_n^* is greater than the critical value, we must reject the hypothesis of normality.

8 Appendix 3. Tables

In this section we present a table with critical values of T_n^* for different values of n . The values were obtained by simulation from 50000 replications. From columns 2 to 8, we find the levels of significance of the test. In columns 9 and 10, we find the values of a_n and C_n respectively.

n	0.15	0.1	0.075	0.05	0.025	0.01	0.001	a_n	C_n
10	0.7547	0.8525	0.92590	1.0203	1.1917	1.4128	1.9025	1.2816	28.5798
11	0.7909	0.8961	0.9714	1.0754	1.2470	1.4733	2.0513	1.3352	34.3184
12	0.8179	0.9254	0.9996	1.1050	1.2860	1.5043	2.0086	1.3830	40.4855
13	0.8489	0.9597	1.0410	1.1456	1.3263	1.5679	2.1420	1.4261	47.0700
14	0.8728	0.9840	1.0632	1.1723	1.3571	1.5931	2.1457	1.4652	54.0622
15	0.8879	1.00573	1.08548	1.2007	1.3975	1.6664	2.2531	1.5341	61.4534
16	0.9132	1.0305	1.1154	1.2308	1.4238	1.6776	2.2766	1.5011	69.2355
17	0.9300	1.0491	1.1289	1.2381	1.4288	1.6714	2.2844	1.5647	77.4015
18	0.9522	1.0769	1.1624	1.2814	1.4854	1.7259	2.4215	1.5932	85.9446
19	0.9691	1.0945	1.1823	1.3062	1.5030	1.7649	2.4001	1.6199	94.8590
20	0.9857	1.1091	1.1963	1.312	1.5213	1.7918	2.4935	1.6448	104.1389
21	1.0076	1.1320	1.2174	1.3381	1.5482	1.8287	2.4845	1.6684	113.7793
22	1.0142	1.1414	1.2330	1.3607	1.5687	1.8671	2.5638	1.6906	123.7752
23	1.0287	1.1549	1.2454	1.3729	1.5899	1.8607	2.5877	1.7117	134.1223
24	1.0434	1.1731	1.2664	1.3940	1.6186	1.9087	2.5657	1.7317	144.8161
25	1.0538	1.1876	1.2837	1.4213	1.6322	1.9017	2.5965	1.7507	155.8529
26	1.0587	1.1905	1.2856	1.4124	1.6442	1.9479	2.6597	1.7688	167.2289
27	1.0745	1.2076	1.3000	1.4348	1.6762	1.9522	2.7323	1.7862	178.9405
28	1.0847	1.2190	1.3199	1.4525	1.6863	1.9890	2.6675	1.8027	190.9843
29	1.0942	1.2286	1.3213	1.4607	1.6994	2.0166	2.7327	1.8186	203.3574
30	1.1102	1.2441	1.3391	1.4708	1.6967	1.9752	2.6952	1.8339	216.0565
31	1.1192	1.2600	1.3598	1.5001	1.7279	2.0439	2.7540	1.8486	229.0789
32	1.1226	1.2671	1.3668	1.5062	1.7311	2.0420	2.7798	1.8627	242.4218
33	1.1363	1.2760	1.3730	1.5112	1.7519	2.0624	2.9159	1.8764	256.0826
34	1.1408	1.2788	1.3781	1.5166	1.7584	2.0704	2.8223	1.8895	270.0589
35	1.1518	1.2886	1.3852	1.5342	1.7691	2.1021	2.8338	1.9022	284.3482
36	1.1603	1.3033	1.4066	1.5445	1.7852	2.1074	2.9445	1.9145	298.9482
37	1.1666	1.3092	1.4067	1.5474	1.7937	2.1235	2.8274	1.9264	313.8568
38	1.1730	1.3191	1.4193	1.5632	1.8161	2.1436	2.9951	1.9379	329.0718
39	1.1908	1.3345	1.4365	1.5804	1.8254	2.1241	2.9245	1.9491	344.5912
40	1.1970	1.3367	1.4392	1.5841	1.8324	2.1714	2.9444	1.9600	360.4130
41	1.2026	1.3437	1.4469	1.5920	1.8416	2.1833	3.0389	1.9705	376.5354
42	1.2111	1.3603	1.4613	1.6042	1.8393	2.1652	2.9808	1.9808	392.9565
43	1.2015	1.3467	1.4528	1.6031	1.8512	2.1814	2.9263	1.9907	409.6745
44	1.2213	1.3708	1.4767	1.6147	1.8585	2.1950	3.0911	2.0004	426.6878
45	1.2285	1.3771	1.4843	1.6341	1.8750	2.1900	3.0056	2.0099	443.9947
46	1.2350	1.3875	1.4955	1.6477	1.9053	2.2467	3.0391	2.0191	461.5935
47	1.2342	1.3844	1.4903	1.6384	1.8839	2.2215	3.0267	2.0281	479.4828
48	1.2469	1.4040	1.5141	1.6680	1.9346	2.2634	3.0625	2.0368	497.6611
49	1.2535	1.4063	1.5159	1.6730	1.9293	2.2552	3.0640	2.0454	516.1269
50	1.2629	1.4271	1.5304	1.6897	1.9490	2.2770	3.1309	2.0537	534.8787
51	1.2645	1.4220	1.5301	1.6947	1.9627	2.3063	3.2487	2.0619	553.9153
52	1.2708	1.4243	1.5399	1.6938	1.9521	2.3066	3.1432	2.0699	573.2352
53	1.2801	1.4340	1.5445	1.7008	1.9603	2.3068	3.1140	2.0777	592.8372

n	0.15	0.1	0.075	0.05	0.025	0.01	0.001	a_n	C_n
54	1.2772	1.4313	1.5422	1.6962	1.9627	2.3047	3.1660	2.0854	612.7200
55	1.2854	1.4378	1.5531	1.7133	1.9860	2.3389	3.2503	2.0928	632.8823
56	1.2902	1.4408	1.5486	1.7107	1.9917	2.3399	3.2057	2.1002	653.3230
57	1.2936	1.4500	1.5653	1.7184	1.9894	2.3707	3.2309	2.1073	674.0410
58	1.2986	1.4590	1.5700	1.7210	1.9798	2.3287	3.2403	2.1144	695.0349
59	1.3175	1.4762	1.5872	1.7379	2.0025	2.3765	3.3164	2.1213	716.3038
60	1.3056	1.4609	1.5736	1.7346	2.0055	2.3570	3.2960	2.1280	737.8466
61	1.3125	1.4747	1.5869	1.7424	2.0206	2.3778	3.3615	2.1347	759.6622
62	1.3120	1.4715	1.5852	1.7434	2.0084	2.3791	3.3989	2.1412	781.7495
63	1.3282	1.4904	1.5979	1.7568	2.0256	2.3538	3.2608	2.1476	804.1076
64	1.3275	1.4875	1.6038	1.7623	2.0344	2.4059	3.3059	2.1539	826.7354
65	1.3259	1.4876	1.6038	1.7701	2.0566	2.4492	3.4750	2.1600	849.6320
66	1.3280	1.4886	1.6081	1.7691	2.0498	2.4100	3.2162	2.1661	872.7964
67	1.3444	1.5083	1.6218	1.7782	2.0520	2.4257	3.4441	2.1721	896.2277
68	1.3521	1.5184	1.6297	1.7866	2.0490	2.4221	3.4079	2.1779	919.9251
69	1.3492	1.5077	1.6262	1.7902	2.0887	2.4654	3.4971	2.1837	943.8875
70	1.3507	1.5173	1.6381	1.8052	2.0929	2.4834	3.3568	2.1893	968.1142
71	1.3485	1.5160	1.6333	1.7975	2.0825	2.4569	3.4083	2.1949	992.6042
72	1.3643	1.5318	1.6550	1.8260	2.1092	2.4882	3.3969	2.2004	1017.3568
73	1.3668	1.5324	1.6465	1.8064	2.0829	2.4538	3.3884	2.2058	1042.3711
74	1.3643	1.5258	1.6434	1.8053	2.0907	2.4632	3.4550	2.2111	1067.6463
75	1.3698	1.5353	1.6487	1.8207	2.1057	2.5039	3.3958	2.2164	1093.1817
76	1.3619	1.5231	1.6324	1.8007	2.0775	2.4609	3.4476	2.2215	1118.9764
77	1.3931	1.5646	1.6840	1.8451	2.1306	2.5129	3.5805	2.2266	1145.0297
78	1.3896	1.5560	1.6807	1.8527	2.1487	2.5420	3.4796	2.2316	1171.3408
79	1.3930	1.5669	1.6900	1.8486	2.1391	2.5279	3.4635	2.2365	1197.9091
80	1.3873	1.5528	1.6711	1.8431	2.1262	2.5284	3.4714	2.2414	1224.7337
81	1.3923	1.5573	1.6734	1.8414	2.1218	2.4825	3.4870	2.2462	1251.8140
82	1.3857	1.5559	1.6767	1.8532	2.1422	2.4974	3.4783	2.2509	1279.1492
83	1.3860	1.5579	1.6799	1.8568	2.1377	2.5472	3.4902	2.2556	1306.7388
84	1.4178	1.5832	1.7032	1.8737	2.1635	2.5825	3.5271	2.2602	1334.5820
85	1.4066	1.5793	1.7007	1.8776	2.1598	2.5435	3.4956	2.2647	1362.6781
86	1.4097	1.5776	1.6929	1.8562	2.1554	2.5406	3.5896	2.2692	1391.0266
87	1.4173	1.5900	1.7129	1.8834	2.1823	2.6082	3.6324	2.2736	1419.6267
88	1.4149	1.5912	1.7170	1.8921	2.1844	2.6253	3.6058	2.2780	1448.4778
89	1.4159	1.5853	1.7026	1.8711	2.1765	2.5753	3.5992	2.2823	1477.5794
90	1.4191	1.5880	1.7085	1.8794	2.1779	2.5629	3.5503	2.2865	1506.9307
91	1.4419	1.6148	1.7353	1.9100	2.2180	2.6173	3.6562	2.2907	1536.5313
92	1.4381	1.6061	1.7302	1.9133	2.2286	2.6358	3.7350	2.2949	1566.3805
93	1.4378	1.6117	1.7277	1.8948	2.2087	2.5941	3.6177	2.2990	1596.4777
94	1.4356	1.6133	1.7393	1.9103	2.2107	2.5887	3.6782	2.3030	1626.8223
95	1.4363	1.6112	1.7321	1.9039	2.2049	2.6048	3.6679	2.3070	1657.4138
96	1.4375	1.6138	1.7321	1.9096	2.2107	2.6169	3.6031	2.3110	1688.2517
97	1.4355	1.6057	1.7274	1.9046	2.2146	2.6168	3.6890	2.3149	1719.3353
98	1.4445	1.6137	1.7390	1.9171	2.2214	2.6184	3.6037	2.3188	1750.6642
99	1.4338	1.6128	1.7343	1.9024	2.2065	2.6307	3.6025	2.3226	1782.2377
100	1.4568	1.6358	1.7637	1.9396	2.2606	2.6495	3.6210	2.3263	1814.0555
101	1.4648	1.6426	1.7661	1.9444	2.2449	2.6978	3.7069	2.3301	1846.1168
102	1.4579	1.6328	1.7594	1.9336	2.2381	2.6473	3.7758	2.3338	1878.4214
103	1.4656	1.6398	1.7669	1.9521	2.2556	2.6575	3.6712	2.3374	1910.9685
104	1.4652	1.6465	1.7742	1.9433	2.2514	2.6530	3.7589	2.3410	1943.7578
105	1.4668	1.6380	1.7589	1.9397	2.2514	2.6561	3.6646	2.3446	1976.7887

n	0.15	0.1	0.075	0.05	0.025	0.01	0.001	a_n	C_n
106	1.4653	1.6402	1.7629	1.9431	2.2407	2.6590	3.7376	2.3481	2010.0608
107	1.4624	1.6360	1.7586	1.9389	2.2620	2.6619	3.7710	2.3516	2043.5736
108	1.4611	1.6334	1.7577	1.9301	2.2403	2.6498	3.8194	2.3551	2077.3266
109	1.4634	1.6421	1.7680	1.9488	2.2502	2.6373	3.7118	2.3585	2111.3193
110	1.4654	1.6381	1.7622	1.9379	2.2344	2.6400	3.6154	2.3619	2145.5513
111	1.4671	1.6525	1.7806	1.9570	2.2594	2.6291	3.7433	2.3652	2180.0221
112	1.4914	1.6740	1.7934	1.9840	2.2849	2.7172	3.7062	2.3686	2214.7313
113	1.4973	1.6782	1.8027	1.9823	2.3010	2.7150	3.7486	2.3719	2249.6784
114	1.4936	1.6668	1.7955	1.9757	2.2761	2.7000	3.7684	2.3751	2284.8629
115	1.4859	1.6651	1.7934	1.9717	2.2655	2.6743	3.7925	2.3783	2320.2846
116	1.4931	1.6768	1.8051	1.9844	2.2938	2.7092	3.7641	2.3815	2355.9428
117	1.4982	1.6737	1.8047	1.9934	2.2950	2.7069	3.8541	2.3847	2391.8373
118	1.4916	1.6766	1.8010	1.9803	2.2962	2.7081	3.7584	2.3878	2427.9675
119	1.4936	1.6765	1.8043	1.9907	2.2986	2.7223	3.8190	2.3909	2464.3331
120	1.4914	1.6709	1.7969	1.9796	2.2913	2.6762	3.6763	2.3940	2500.9337
121	1.4992	1.6804	1.8054	1.9775	2.2979	2.7028	3.8052	2.3970	2537.7688
122	1.5008	1.6747	1.7973	1.9802	2.2697	2.6783	3.7898	2.4000	2574.8381
123	1.4952	1.6757	1.8048	1.9860	2.2866	2.7052	3.7794	2.4030	2612.1411
124	1.4959	1.6759	1.8052	1.9868	2.2933	2.7154	3.6734	2.4060	2649.6775
125	1.5318	1.7153	1.8474	2.0254	2.3491	2.7653	3.9034	2.4089	2687.4469
126	1.5294	1.7087	1.8409	2.0284	2.3443	2.7871	3.8894	2.4118	2725.4489
127	1.5381	1.7214	1.8481	2.0238	2.3321	2.7497	3.8923	2.4147	2763.6831
128	1.5309	1.7123	1.8400	2.0182	2.3288	2.7619	3.8255	2.4176	2802.1492
129	1.5409	1.7260	1.8572	2.0429	2.3626	2.7835	3.7827	2.4204	2840.8467
130	1.5261	1.7104	1.8356	2.0192	2.3348	2.7624	3.9036	2.4232	2879.7754
131	1.5235	1.7010	1.8275	2.0085	2.3269	2.7192	3.8243	2.4260	2918.9348
132	1.5336	1.7112	1.8436	2.0235	2.3412	2.7569	3.7583	2.4287	2958.3246
133	1.5288	1.7149	1.8497	2.0347	2.3422	2.7812	3.7633	2.4315	2997.9444
134	1.5335	1.7190	1.8476	2.0265	2.3437	2.8052	3.8702	2.4342	3037.7939
135	1.5384	1.7142	1.8415	2.0223	2.3308	2.7512	3.8024	2.4369	3077.8728
136	1.5266	1.7098	1.8395	2.0177	2.3359	2.7451	3.8569	2.4395	3118.1807
137	1.5298	1.7110	1.8388	2.0191	2.3386	2.7836	3.8683	2.4422	3158.7172
138	1.5304	1.7133	1.8412	2.0182	2.3396	2.7247	3.7756	2.4448	3199.4820
139	1.5280	1.7143	1.8458	2.0307	2.3437	2.8075	3.8780	2.4474	3240.4748
140	1.5367	1.7242	1.8548	2.0318	2.3468	2.7696	3.8377	2.4500	3281.6953
141	1.5285	1.7122	1.8489	2.0288	2.3198	2.7296	3.9228	2.4526	3323.1431
142	1.5277	1.7122	1.8403	2.0260	2.3461	2.7479	3.8630	2.4551	3364.8179
143	1.5703	1.7560	1.8874	2.0776	2.4052	2.8404	3.9334	2.4576	3406.7194
144	1.5686	1.7593	1.8872	2.0716	2.3878	2.8007	4.1252	2.4601	3448.8472
145	1.5750	1.7631	1.8959	2.0823	2.3833	2.8230	4.0015	2.4626	3491.2011
146	1.5679	1.7588	1.8976	2.0884	2.4081	2.8252	3.8971	2.4651	3533.7808
147	1.5641	1.7590	1.8962	2.0794	2.4078	2.8128	3.9899	2.4675	3576.5858
148	1.5628	1.7477	1.8827	2.0588	2.3786	2.8044	4.0139	2.4699	3619.6160
149	1.5781	1.7605	1.8931	2.0821	2.4324	2.8661	4.0380	2.4723	3662.8710
150	1.5693	1.7627	1.8942	2.0844	2.4212	2.8560	3.9064	2.4747	3706.3505
151	1.5664	1.7507	1.8861	2.0802	2.4117	2.8842	4.0000	2.4771	3750.0543
152	1.5685	1.7542	1.8842	2.0725	2.4020	2.8504	3.9861	2.4795	3793.9820
153	1.5670	1.7486	1.8777	2.0670	2.3894	2.8398	3.9350	2.4818	3838.1333
154	1.5736	1.7672	1.8997	2.0929	2.4096	2.8565	3.9442	2.4841	3882.5079
155	1.5720	1.7637	1.8945	2.0866	2.4006	2.8125	3.8631	2.4864	3927.1056
156	1.5713	1.7670	1.8988	2.0980	2.4320	2.8729	3.9977	2.4887	3971.9261
157	1.5689	1.7527	1.8830	2.0711	2.3973	2.8370	3.8880	2.4910	4016.9691

n	0.15	0.1	0.075	0.05	0.025	0.01	0.001	a_n	C_n
158	1.5723	1.7604	1.8990	2.0775	2.4074	2.8411	3.9726	2.4932	4062.2343
159	1.5728	1.7598	1.8960	2.0764	2.3939	2.8315	4.0027	2.4955	4107.7214
160	1.5662	1.7526	1.8812	2.0655	2.4023	2.8322	4.0344	2.4977	4153.4301
161	1.5673	1.7568	1.8910	2.0793	2.4079	2.8703	3.9389	2.4999	4199.3603
162	1.5670	1.7538	1.8878	2.0758	2.4150	2.8757	4.0885	2.5021	4245.5115
163	1.5731	1.7620	1.8952	2.0779	2.3952	2.8549	3.9610	2.5043	4291.8836
164	1.5638	1.7507	1.8851	2.0779	2.4150	2.8065	3.8816	2.5064	4338.4763
165	1.5700	1.7558	1.8872	2.0719	2.3851	2.8230	3.9072	2.5086	4385.2894
166	1.5566	1.7460	1.8801	2.0689	2.3815	2.8411	3.9225	2.5107	4432.3224
167	1.6161	1.8091	1.9469	2.1541	2.4844	2.9439	4.0744	2.5128	4479.5753
168	1.6217	1.8117	1.9473	2.1437	2.4647	2.9140	4.0022	2.5150	4527.0477
169	1.6170	1.8016	1.9385	2.1341	2.4764	2.9199	4.0265	2.5170	4574.7394
170	1.6114	1.8041	1.9389	2.1331	2.4608	2.9014	4.1730	2.5191	4622.6501
171	1.6161	1.8114	1.9454	2.1368	2.4676	2.9217	4.0926	2.5212	4670.7796
172	1.6106	1.7986	1.9399	2.1326	2.4624	2.8999	4.1438	2.5232	4719.1276
173	1.6192	1.8214	1.9553	2.1474	2.4929	2.9706	4.0761	2.5253	4767.6939
174	1.6157	1.8069	1.9431	2.1289	2.4482	2.8809	4.0472	2.5273	4816.4782
175	1.6082	1.8052	1.9447	2.1369	2.4672	2.9281	3.9707	2.5293	4865.4804
176	1.6173	1.8097	1.9474	2.1422	2.4750	2.9213	4.0997	2.5313	4914.7001
177	1.6078	1.7979	1.9427	2.1378	2.4589	2.9142	4.0648	2.5333	4964.1371
178	1.6102	1.8069	1.9417	2.1439	2.4849	2.9138	3.9984	2.5353	5013.7912
179	1.6150	1.8043	1.9347	2.1237	2.4533	2.9027	4.0803	2.5372	5063.6621
180	1.6133	1.8025	1.9414	2.1374	2.4735	2.9096	3.9865	2.5392	5113.7496
181	1.6080	1.7933	1.9204	2.1121	2.4501	2.8787	4.0807	2.5411	5164.0535
182	1.6149	1.8081	1.9445	2.1443	2.4809	2.9197	4.0438	2.5430	5214.5735
183	1.6136	1.8006	1.9423	2.1373	2.4674	2.8889	4.2658	2.5450	5265.3095
184	1.6041	1.7966	1.9325	2.1395	2.4730	2.9267	4.1624	2.5469	5316.2611
185	1.6135	1.8104	1.9452	2.1380	2.4619	2.8934	4.3550	2.5488	5367.4282
186	1.6122	1.8053	1.9433	2.1357	2.4744	2.9524	4.0716	2.5506	5418.8106
187	1.6162	1.8079	1.9412	2.1267	2.4492	2.9129	4.0421	2.5525	5470.4079
188	1.6070	1.7985	1.9401	2.1441	2.4866	2.9596	4.0490	2.5544	5522.2200
189	1.6244	1.8170	1.9510	2.1456	2.4839	2.9188	4.0255	2.5562	5574.2468
190	1.6234	1.8161	1.9584	2.1561	2.4949	2.9272	4.0296	2.5580	5626.4878
191	1.6099	1.8018	1.9372	2.1328	2.4302	2.8630	4.0091	2.5599	5678.9431
192	1.6190	1.8098	1.9478	2.1408	2.4843	2.9319	4.1542	2.5617	5731.6122
193	1.6219	1.8135	1.9520	2.1423	2.4679	2.9339	4.1218	2.5635	5784.4951
194	1.6181	1.8069	1.9465	2.1347	2.4615	2.8779	4.0501	2.5653	5837.5914
195	1.6056	1.8000	1.9323	2.1229	2.4512	2.8756	4.1073	2.5671	5890.9011
196	1.6090	1.8070	1.9426	2.1445	2.4596	2.9005	4.1082	2.5688	5944.4238
197	1.6227	1.8158	1.9551	2.1444	2.4778	2.9440	4.1214	2.5706	5998.1594
198	1.6224	1.8160	1.9520	2.1429	2.4777	2.9228	4.1046	2.5724	6052.1077
199	1.6171	1.8062	1.9428	2.1411	2.4806	2.9012	4.0380	2.5741	6106.2685
200	1.6534	1.8537	1.9956	2.1848	2.5180	2.9667	4.2543	2.5758	6160.6415
n	0.15	0.1	0.075	0.05	0.025	0.01	0.001	a_n	C_n
250	1.6854	1.8946	2.0403	2.2348	2.5882	3.0200	4.4252	2.6521	9145.8
500	1.8803	2.1020	2.2570	2.4775	2.8403	3.3546	4.6692	2.8782	31419.2
750	1.9601	2.1819	2.3448	2.5749	2.9473	3.4903	4.8916	3.0038	65019.6
1000	2.0218	2.2488	2.4066	2.6323	3.0151	3.5407	4.9433	3.0902	109204
10000	2.7769	3.0512	3.2332	3.4776	3.8861	4.4679	5.6259	3.7190	7439183

9 Acknowledgements

The author wishes to thank Enrique Cabaña for your help in the preparation of this work.

10 References

1. Anderson, T. W. and Darling, D. A. (1954). A test of goodness of fit. *J. Amer. Statist. Assoc.* 49, 765–769.
2. del Barrio, E., Cuesta Albertos, J. A., Matrán, C. and Rodríguez Rodríguez, J. (1999). Tests of fit based on the L2-Wasserstein distance. *Ann. Statist.* 27, 1230–1239.
3. Cramér, H. (1928). On the composition of elementary errors. Second paper: Statistical applications. *Skand. Aktuarietidskr.* 11, 141–180.
4. Csörgö³, S. and Faraway, J. (1996). The Exact and Asymptotic Distributions of Cramér–von Mises Statistics *J. R. Statist. Soc. B* 58(1), 221–234.
5. Durbin, J. (1973). Weak convergence of the sample distribution when parameters are estimated. *Ann. Statist.* 1, 219.
6. Fisher, R. A., (1930). The moments of the distribution for normal samples of measures of departure from normality. *Proceedings of Royal Society, A*, 130, 16.
7. Gan, F. F., Koelher, K. J. (1990). Goodness of fit tests based on P–P probability plots. *Technometrics* 32, 289–303.
8. Kalemkerian, J., (2017). An integral formula for the distribution of self-normalized Gaussian random samples. *Communications in Statistics. Theory and Methods* 46(10), 4671–4685.
9. Lockhart, R. A. and Stephens, M. A., (1998). *The probability plot: Test of fit based on the correlation coefficient*. In *Order statistics: Applications. Handbook of Statistics* 17, 453–473. North Holland, Amsterdam.
10. Pearson, K. (1895). Contributions to the mathematical theory of evolution. *Philosophical Transactions of the Royal Society* 91, 343.
11. Pearson, E. S., (1930). A further development of tests for normality. *Biometrika* 22, 239–249.
12. Pettitt, A. N. (1976). Cramér–von Mises statistics for testing normality with censored samples. *Biometrika*, 63(3), 475–481.
13. Pettitt, A. N. and Stephens, M. A. (1976). Modified Cramér–von Mises statistics with censored samples. *Biometrika* 63(2), 291–298.

14. Shapiro, S. S. and Wilk, M. B. (1965). An analysis of variance test for normality (complete samples). *Biometrika* 64, 415–418.
15. Shorack, G. and Wellner, J. (1982). Limit theorems and inequalities for the uniform empirical processes indexed by intervals. *The Annals of Probability* 10(3), 639–652.
16. Skorokhod, A. (1956). Limit theorem for stochastic processes. *Theor. Probability Appl.* 1, 261–290.
17. Smirnov, N. V. (1936). Sur la distribution de w^2 (Critérium de M. R. von Mises). *Comptes Rendus de l'Académie des Sciences* 202, 449–452.
18. Smirnov, N. V. (1937). Sur la distribution de w^2 (Critérium de M. R. von Mises). *Matematicheskij Sbornik* (in Russian with French summary) 2, 973–993.
19. Stephens, M. A. (1974). EDF Statistics for Goodness of Fit and Some Comparisons. *JASA* 79, 730–737.
20. Stephens, M. A. (1986). Tests based on EDF statistics. In R. B. D'Agostino and M. A. Stephens, eds.. *Goodness of Fit Techniques*, North-Holland, Amsterdam.
21. de Wet and Venter, J. (1973). Asymptotic distributions for quadratic forms with applications to test of fit. *Ann. Statist.* 2, 380–387.
22. von Mises, R. (1931). *Wahrscheinlichkeitsrechnung*, Vienna.
23. Williams, P. (1935). Note on the sampling distribution of $\sqrt{\beta_2}$, where the population is normal. *Biometrika* 27, 269–271.