

On the Optimality of Kernel-Embedding Based Goodness-of-Fit Tests

Krishnakumar Balasubramanian*, Tong Li[†] and Ming Yuan[‡]

*Princeton University and ^{†,‡}Columbia University

(November 16, 2021)

Abstract

The reproducing kernel Hilbert space (RKHS) embedding of distributions offers a general and flexible framework for testing problems in arbitrary domains and has attracted considerable amount of attention in recent years. To gain insights into their operating characteristics, we study here the statistical performance of such approaches within a minimax framework. Focusing on the case of goodness-of-fit tests, our analyses show that a vanilla version of the kernel-embedding based test could be suboptimal, and suggest a simple remedy by moderating the embedding. We prove that the moderated approach provides optimal tests for a wide range of deviations from the null and can also be made adaptive over a large collection of interpolation spaces. Numerical experiments are presented to further demonstrate the merits of our approach.

[‡]Address for Correspondence: Department of Statistics, Columbia University, 1255 Amsterdam Avenue, New York, NY 10027.

1 Introduction

In recent years, statistical tests based on the reproducing kernel Hilbert space (RKHS) embedding of distributions have attracted much attention because of their flexibility and broad applicability. Like other kernel methods, RKHS embedding based tests present a general and unifying framework for testing problems in arbitrary domains by using appropriate kernels defined on those domains. See, *e.g.*, [Muandet et al. \(2017\)](#), for a recent review and detailed discussion about the applications of kernel embeddings. The idea of using kernel embedding for comparing probability distributions was initially introduced by [Smola et al. \(2007\)](#); [Gretton et al. \(2007, 2012\)](#). Related extensions were also proposed by [Harchaoui et al. \(2007\)](#); [Zaremba et al. \(2013\)](#). Furthermore, [Sejdinovic et al. \(2013\)](#) established a close relationship between kernel-based hypothesis tests and energy distanced based test introduced by [Székely et al. \(2007\)](#). See also [Lyons \(2013\)](#). More recently, motivated by several applications based on quantifying the convergence of Monte Carlo simulations, [Liu et al. \(2016\)](#), [Chwialkowski et al. \(2016\)](#) and [Gorham and Mackey \(2017\)](#) proposed goodness-of-fit tests which were based on combining the kernel based approach with Stein’s identity. A linear-time method for goodness-of-fit was also proposed by [Jitkrittum et al. \(2017\)](#) recently. Finally, the idea of kernel-embedding has also been used for constructing implicit generative models (*e.g.*, [Dziugaite et al., 2015](#); [Li et al., 2015](#)).

Despite their popularity, fairly little is known about the statistical performance of these kernel-embedding based tests. Our goal is to fill in this void. In particular, we focus on kernel-embedding based goodness-of-fit tests and investigate their power under a general composite alternative. Our results not only provide new insights on the operating characteristics of these kernel-embedding based tests but also suggest improved testing procedures that are minimax optimal and adaptive over a large collection of alternatives.

The problem of testing for goodness-of-fit has a long and illustrious history in statistics and is often associated with household names such as *Kolmogrov-Smirnov tests*, *Pearson’s Chi-square test* or *Neyman’s smooth test*. A plethora of other techniques have also been proposed over the years in both parametric and non-parametric settings (*e.g.*, [Ingster and Suslina, 2003](#); [Lehmann and Romano, 2008](#)). Most of the existing techniques are developed with the domain $\mathcal{X} = \mathbb{R}$ or $[0, 1]$ in mind and work the best in these cases. Modern applications, however, oftentimes involve domains different from these traditional ones. For example,

when dealing with directional data, which arise naturally in applications such as diffusion tensor imaging, it is natural to consider \mathcal{X} as the unit sphere in \mathbb{R}^3 (e.g., Jupp, 2005). Another example occurs in the context of ranking or preference data (e.g., Ailon et al., 2008). In these cases, \mathcal{X} can be taken as the group of permutations. Furthermore, motivated by several applications, combinatorial testing problems have been investigated recently (e.g., Addario-Berry et al., 2010), where the spaces under consideration are specific combinatorially structured spaces.

A particularly attractive approach to goodness-of-fit testing problems in general domains is through RKHS embedding of distributions. Specifically, let $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ be a Mercer kernel that is symmetric, positive (semi-)definite and square integrable. The RKHS embedding of a probability measure P on $(\mathcal{X}, \mathcal{B})$, with respect to K , is given by

$$\mu_P(\cdot) := \int_{\mathcal{X}} K(x, \cdot) P(dx).$$

The Moore-Aronszajn Theorem indicates that there is an RKHS, denoted by $(\mathcal{H}(K), \langle \cdot, \cdot \rangle_K)$, uniquely identified with the kernel K (e.g., Aronszajn, 1950). It is clear that $\mu_P \in \mathcal{H}(K)$, and hence the notion of RKHS embedding. The RKHS embedding of probability measures is closely related to a certain integral probability metric. The so-called *maximum mean discrepancy* (MMD) between two probability measures P and Q is defined as

$$\gamma_K(P, Q) := \sup_{f \in \mathcal{H}(K) : \|f\|_K \leq 1} \int_{\mathcal{X}} f d(P - Q),$$

where $\|\cdot\|_K$ is the norm associated with $(\mathcal{H}(K), \langle \cdot, \cdot \rangle_K)$. It is not hard to see (e.g., Gretton et al., 2012) that

$$\gamma_K(P, Q) = \|\mu_P - \mu_Q\|_K.$$

The goodness-of-fit test can be carried out conveniently through RKHS embeddings of P and P_0 by first constructing an estimate of $\gamma_K(P, P_0)$:

$$\gamma_K(\hat{P}_n, P_0) := \sup_{f \in \mathcal{H}(K) : \|f\|_K \leq 1} \int_{\mathcal{X}} f d(\hat{P}_n - P_0),$$

where \hat{P}_n is the empirical distribution of X_1, \dots, X_n , and then rejecting H_0 if the estimate exceeds a threshold calibrated to ensure a certain significance level, say α ($0 < \alpha < 1$).

In this paper, we investigate the power of the above discussed testing strategy under a general composite alternative. Following the spirit of Ingster and Suslina (2003), we consider

in particular a set of alternatives that are increasingly close to the null hypothesis. To fix ideas, we assume hereafter that P is dominated by P_0 under the alternative so that the Radon-Nikodym derivative dP/dP_0 is well defined. Recall that the χ^2 divergence between P and P_0 is defined as

$$\chi^2(P, P_0) := \int_{\mathcal{X}} \left(\frac{dP}{dP_0} \right)^2 dP_0 - 1.$$

We are particularly interested in the detection boundary, namely how close P and P_0 can be in terms of χ^2 distance, under the alternative, so that a test based on a sample of n observations can still consistently distinguish between the null hypothesis and the alternative. For example, in the parametric setting where P is known up to a finite dimensional parameters under the alternative, the detection boundary of the likelihood ratio test is n^{-1} under mild regularity conditions (*e.g.*, [Lehmann and Romano, 2008](#)). We are concerned here with alternatives that are nonparametric in nature. Our first result suggests that the detection boundary for aforementioned $\gamma_K(\hat{P}_n, P_0)$ based test is of the order $n^{-1/2}$. However, our main results indicate, perhaps surprising at first, that this rate is far from optimal and the gap between it and the usual parametric rate can be largely bridged.

In particular, we argue that the distinguishability between P and P_0 depends on how close $u := dP/dP_0 - 1$ is to the RKHS $\mathcal{H}(K)$. The closeness of u to $\mathcal{H}(K)$ can be measured by the distance from u to an arbitrary ball in $\mathcal{H}(K)$. In particular, we shall consider the case where $\mathcal{H}(K)$ is dense in $L_2(P_0)$, and focus on functions that are polynomially approximable by $\mathcal{H}(K)$ for concreteness. More precisely, for some constants $M, \theta > 0$, denote by $\mathcal{F}(\theta; M)$ the collection of functions $f \in L_2(P_0)$ such that for any $R > 0$, there exists an $f_R \in \mathcal{H}(K)$ such that

$$\|f_R\|_K \leq R, \quad \text{and} \quad \|f - f_R\|_{L_2(P_0)} \leq MR^{-1/\theta}.$$

See, *e.g.*, [Cucker and Zhou \(2007\)](#) for further discussion on these so-called interpolation spaces and their use in statistical learning. We shall also adopt the convention that

$$\mathcal{F}(0; M) = \{f \in \mathcal{H}(K) : \|f\|_K \leq M\}.$$

We investigate the optimal rate of detection for testing H_0 against

$$H_1(\Delta_n, \theta, M) : P \in \mathcal{P}(\Delta_n, \theta, M), \tag{1}$$

where $\mathcal{P}(\Delta_n, \theta, M)$ is the collection of distributions P on $(\mathcal{X}, \mathcal{B})$ satisfying:

$$dP/dP_0 - 1 \in \mathcal{F}(\theta; M), \quad \text{and} \quad \chi^2(P, P_0) \geq \Delta_n.$$

We call r_n the optimal rate of detection if for any $c > 0$, there exists no consistent test whenever $\Delta_n \leq cr_n$; and on the other hand, a consistent test exists as long as $\Delta_n \gg r_n$.

Although one could consider a more general setup, for concreteness, we assume that the eigenvalues of K with respect to $L_2(P_0)$ decays polynomially in that $\lambda_k \asymp k^{-2s}$. We show that the optimal rate of detection for testing H_0 against $H_1(\Delta_n, \theta, M)$ for any $\theta \geq 0$ is $n^{-\frac{4s}{4s+\theta+1}}$. The rate of detection, although not achievable with a $\gamma_K(\hat{P}_n, P_0)$ based test, can be attained via a moderated version of the MMD based approach. A practical challenge to the approach, however, is its reliance on the knowledge of θ . Unlike s which is determined by K and P_0 and therefore known apriori, θ depends on u and is not known in advance. This naturally brings about the issue of adaptation – is there an agnostic approach that can adaptively attain the optimal detection boundary without the knowledge of θ . We show that the answer is affirmative although a small price in the form of $\log \log n$ is required to achieve such adaptation.

The rest of the paper is organized as follows. We first analyze the power of MMD based tests in Section 2. This analysis reveals a significant gap between the detection boundary achieved by the MMD based test and the usual parametric $1/n$ rate. In turn, this prompts us to introduce, in Section 3, a new class of tests based on a modified MMD. We show that the new tests are rate optimal. To address the practical challenge of choosing an appropriate tuning parameter for these tests, we investigate the issue of optimal adaptation in Section 4, where we establish the optimal rates of detection for adaptively testing H_0 against a broader set of alternatives and propose a test based on the modified MMD that can attain these rates. Numerical experiments are presented in Section 5. All proofs are relegated to Section 6.

2 Operating characteristics of MMD based test

2.1 Background and notation

In this section, we investigate the performance of the MMD based test. As shown in Gretton et al. (2012), the squared MMD between two probability distributions P and P_0

can be expressed as

$$\gamma_K^2(P, P_0) = \int K(x, x') d(P - P_0)(x) d(P - P_0)(x'). \quad (2)$$

Write

$$\bar{K}(x, x') = K(x, x') - \mathbb{E}_{P_0} K(x, X) - \mathbb{E}_{P_0} K(X, x') + \mathbb{E}_{P_0} K(X, X'),$$

where the subscript P_0 signifies the fact that the expectation is taken over $X, X' \sim P_0$ independently. By (2), $\gamma_K^2(P, P_0) = \gamma_{\bar{K}}^2(P, P_0)$. Therefore, without loss of generality, we shall assume in what follows that K is degenerate under P_0 , *i.e.*,

$$\mathbb{E}_{P_0} K(X, \cdot) = 0. \quad (3)$$

For brevity, we shall omit the subscript K in γ in the rest of the paper, unless it is necessary to emphasize the dependence of MMD on the reproducing kernel.

Assuming that K is square integrable, by Mercer's theorem, it can be decomposed as

$$K(x, x') = \sum_{k \geq 1} \lambda_k \varphi_k(x) \varphi_k(x'), \quad (4)$$

where the limit is in the sense of $L_2(P_0)$, $\lambda_1 > \lambda_2 > \dots > 0$ are the positive eigenvalues of the integral operator induced by K , and $\{\varphi_k : k \geq 1\}$ are the corresponding orthonormal eigenfunctions, *i.e.*, $\langle \varphi_k, \varphi_{k'} \rangle_{L_2(P_0)} = \delta_{k,k'}$ and $\langle \varphi_k, \varphi_{k'} \rangle_K = \lambda_k^{-1} \delta_{k,k'}$, with δ representing the Kronecker delta. For the sake of concreteness, we shall assume K is universal in that $\{\varphi_k : k \geq 1\}$ forms an orthonormal basis of $L_2(P_0)$, and has infinitely many positive eigenvalues decaying polynomially, that is,

$$0 < \underline{\lim}_{k \rightarrow \infty} k^{2s} \lambda_k \leq \overline{\lim}_{k \rightarrow \infty} k^{2s} \lambda_k < \infty \quad (5)$$

for some $s > 1/2$. Moreover, we assume the eigenfunctions are uniformly bounded, *i.e.*,

$$\sup_{k \geq 1} \|\varphi_k\|_\infty < \infty. \quad (6)$$

Assumptions (5) and (6) ensure that the spectral decomposition (4) holds both pointwisely and uniformly.

Note that (3) implies $\mathbb{E}_{P_0} \varphi_k(X) = 0$, $\forall k \geq 1$, and (4) gives

$$\gamma^2(P, P_0) = \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2$$

for any P . Accordingly, when P is replaced by the empirical distribution \hat{P}_n , the empirical squared MMD can be expressed as

$$\gamma^2(\hat{P}_n, P_0) = \sum_{k \geq 1} \lambda_k \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) \right]^2.$$

Classic results on the asymptotics of V-statistic ([Serfling, 2009](#)) imply that

$$n\gamma^2(\hat{P}_n, P_0) \xrightarrow{d} \sum_{k \geq 1} \lambda_k Z_k^2 := W$$

under H_0 , where $Z_k \stackrel{i.i.d.}{\sim} N(0, 1)$. Let T_{MMD} be an MMD based test, which rejects H_0 if and only if $n\gamma^2(\hat{P}_n, P_0)$ exceeds the upper α quantile $q_{w, 1-\alpha}$ of W , *i.e.*,

$$T_{\text{MMD}} = \mathbb{1}_{\{n\gamma^2(\hat{P}_n, P_0) > q_{w, 1-\alpha}\}}.$$

The above limiting distribution of $n\gamma^2(\hat{P}_n, P_0)$ immediately suggests that T_{MMD} is an asymptotic α -level test.

2.2 Power analysis for MMD based tests

We now investigate the power of T_{MMD} in testing H_0 against $H_1(\Delta_n, \theta, M)$ given by (1). Recall that the type II error of a test $T : \mathcal{X}^n \rightarrow [0, 1]$ for testing H_0 against a composite alternative $H_1 : P \in \mathcal{P}$ is given by

$$\beta(T; \mathcal{P}) = \sup_{P \in \mathcal{P}} \mathbb{E}_P[1 - T(X_1, \dots, X_n)],$$

where \mathbb{E}_P means taking expectation over $X_1, \dots, X_n \stackrel{i.i.d.}{\sim} P$. For brevity, we shall write $\beta(T; \Delta_n, \theta, M)$ instead of $\beta(T; \mathcal{P}(\Delta_n, \theta, M))$ in what follows. The performance of a test T can then be evaluated by its detection boundary, that is, the smallest Δ_n under which the type II error converges to 0 as $n \rightarrow \infty$. Our first result establishes the convergence rate of the detection boundary for T_{MMD} in the case when $\theta = 0$. Hereafter, we abbreviate M in $\mathcal{P}(\Delta_n, \theta, M)$, $H_1(\Delta_n, \theta, M)$ and $\beta(T; \Delta_n, \theta, M)$, unless it is necessary to emphasize the dependence.

Theorem 1. *Consider testing $H_0 : P = P_0$ against $H_1(\Delta_n, 0)$ by T_{MMD} .*

(i) If $\sqrt{n}\Delta_n \rightarrow \infty$, then

$$\beta(T_{\text{MMD}}; \Delta_n, 0) \rightarrow 0 \quad \text{as } n \rightarrow \infty;$$

(ii) conversely, there exists a constant $c_0 > 0$ such that

$$\lim_{n \rightarrow \infty} \beta(T_{\text{MMD}}; c_0 n^{-1/2}, 0) > 0.$$

Theorem 1 shows that when the alternative $H_1(\Delta_n, 0)$ is considered, the detection boundary of T_{MMD} is of the order $n^{-1/2}$. It is of interest to compare the detection rate achieved by T_{MMD} with that in a parametric setting where consistent tests are available if $n\Delta_n \rightarrow \infty$ (Lehmann and Romano, 2008). It is natural to raise the question to what extent such a gap can be entirely attributed to the fundamental difference between parametric and non-parametric testing problems. We shall now argue that this gap actually is largely due to the sup-optimality of T_{MMD} , and the detection boundary of T_{MMD} could be significantly improved through a slight modification of the MMD.

3 Optimal tests based on moderated MMD

3.1 Moderated MMD test statistic

The basic idea behind MMD is to project two probability measures onto a unit ball in $\mathcal{H}(K)$ and use the distance between the two projections to measure the distance between the original probability measures. If the two probability measures are far away from $\mathcal{H}(K)$, the distance between the two projections may not honestly reflect the distance between them. More specifically, $\gamma^2(P, P_0) = \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2$, while the χ^2 distance between P and P_0 is $\chi^2(P, P_0) = \sum_{k \geq 1} [\mathbb{E}_P \varphi_k(X)]^2$. Considering that λ_k decreases with k , $\gamma^2(P, P_0)$ can be much smaller than $\chi^2(P, P_0)$. To overcome this problem, we consider a moderated version of the MMD which allows us to project the probability measures onto a larger ball in $\mathcal{H}(K)$. The new class of integral probability metric between two distributions P and Q is given as

$$\eta_{K, \varrho}(P, Q; P_0) = \sup_{f \in \mathcal{H}(K): \|f\|_{L_2(P_0)}^2 + \varrho^2 \|f\|_K^2 \leq 1} \int_{\mathcal{X}} f d(P - Q) \quad (7)$$

for a given distribution P_0 and a constant $\varrho > 0$. It should be noted that a related test statistics was proposed previously by [Harchaoui et al. \(2007\)](#) from a completely different viewpoint.

It is worth noting that $\eta_{K,\varrho}(P, Q; P_0)$ can also be identified with a particular type of RKHS embedding. Specifically, $\eta_{K,\varrho}(P, Q; P_0) = \gamma_{\tilde{K}_\varrho}(P, Q)$, where

$$\tilde{K}_\varrho(x, x') := \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho^2} \varphi_k(x) \varphi_k(x').$$

We shall abbreviate the dependence of η on K and P_0 unless necessary. The unit ball in (7) is defined in terms of both RKHS norm and L^2 norm. Recall that $u = dP/dP_0 - 1$ so that

$$\sup_{\|f\|_{L_2(P_0)} \leq 1} \int_{\mathcal{X}} f d(P - P_0) = \sup_{\|f\|_{L_2(P_0)} \leq 1} \int_{\mathcal{X}} f u dP_0 = \|u\|_{L_2(P_0)} = \chi(P, P_0).$$

We can therefore expect that a smaller ϱ will make $\eta_\varrho^2(P, P_0)$ closer to $\chi^2(P, P_0)$, since the unit ball to be considered will become more similar to the unit ball in $L_2(P_0)$. This can also be verified by noticing that $\eta_\varrho^2(P, P_0)$ could be expressed as

$$\eta_\varrho^2(P, P_0) = \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho^2} [\mathbb{E}_P \varphi_k(X)]^2.$$

Therefore, we choose ϱ converging to 0 when constructing our test statistic.

Hereafter we shall attach the subscript n to ϱ to signify its dependence on n . We now argue that letting ϱ_n converge to 0 at an appropriate rate indeed results in a test more powerful than T_{MMD} . The test statistic we propose is the empirical version of $\eta_{\varrho_n}^2(P, P_0)$,

$$\eta_{\varrho_n}^2(\hat{P}_n, P_0) = \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) \right]^2. \quad (8)$$

3.2 Operating characteristics of $\eta_{\varrho_n}^2(\hat{P}_n, P_0)$ based tests

Although the expression for $\eta_{\varrho_n}^2(\hat{P}_n, P_0)$ given by (8) looks similar to that of $\gamma^2(\hat{P}_n, P_0)$, their asymptotic behaviors are quite different. At a technical level, this is due to the fact that the eigenvalues of the underlying kernel

$$\tilde{\lambda}_{nk} := \frac{\lambda_k}{\lambda_k + \varrho_n^2}$$

depend on n and may not be uniformly summable over n . As presented in the following theorem, a certain type of asymptotic normality, instead of a sum of chi-squares as in the case of $\gamma^2(\widehat{P}_n, P_0)$, holds for $\eta_{\varrho_n}^2(\widehat{P}_n, P_0)$ under P_0 , which helps determine the rejection region of the $\eta_{\varrho_n}^2$ based test.

Theorem 2. *Assume that $\varrho_n \rightarrow 0$ as $n \rightarrow \infty$ in such a fashion that $n\varrho_n^{1/(2s)} \rightarrow \infty$. Then under H_0 where $X_1, \dots, X_n \stackrel{i.i.d.}{\sim} P_0$,*

$$v_n^{-1/2}[n\eta_{\varrho_n}^2(\widehat{P}_n, P_0) - A_n] \xrightarrow{d} N(0, 2),$$

where

$$v_n = \sum_{k \geq 1} \left(\frac{\lambda_k}{\lambda_k + \varrho_n^2} \right)^2, \quad \text{and} \quad A_n = \frac{1}{n} \sum_{i=1}^n \tilde{K}_{\varrho_n}(X_i, X_i).$$

In the light of Theorem 2, a test that rejects H_0 if and only if

$$2^{-1/2}v_n^{-1/2}[n\eta_{\varrho_n}^2(\widehat{P}_n, P_0) - A_n]$$

exceeds $z_{1-\alpha}$ is an asymptotic α -level test, where $z_{1-\alpha}$ stands for the $1 - \alpha$ quantile of a standard normal distribution. We refer to this test as T_{M^3d} . The performance of T_{M^3d} under the alternative hypothesis is characterized by the following theorem, showing that its detection boundary is much improved when compared with that of T_{MMD} .

Theorem 3. *Consider testing H_0 against $H_1(\Delta_n, \theta)$ by T_{M^3d} with $\varrho_n = cn^{-\frac{2s(\theta+1)}{4s+\theta+1}}$ for an arbitrary constant $c > 0$. If $n^{\frac{4s}{4s+\theta+1}}\Delta_n \rightarrow \infty$, then*

$$\beta(T_{M^3d}; \Delta_n, \theta) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Theorem 3 indicates that the detection boundary for T_{M^3d} is $n^{-4s/(4s+\theta+1)}$. In particular, when testing H_0 against $H_1(\Delta_n, 0)$, i.e., $\theta = 0$, it becomes $n^{-4s/(4s+1)}$. This is to be contrasted with the detection boundary for T_{MMD} , which, as suggested by Theorem 1, is of the order $n^{-1/2}$. It is also worth noting that the detection boundary for T_{M^3d} deteriorates as θ increases, implying that it is harder to test against a larger interpolation space.

3.3 Minimax optimality

It is of interest to investigate if the detection boundary of T_{M^3d} can be further improved. We now show that the answer is negative in a certain sense. More specifically, we shall

follow the minimax framework for nonparametric hypothesis testing pioneered by Ingster (see, *e.g.*, [Ingster, 1993, 1995](#)) and show that T_{M^3d} attains the optimal rate of detection for testing H_0 against $H_1(\Delta_n, \theta)$ in that no consistent test exists if there exists $c > 0$ such that $\Delta_n \leq cn^{-\frac{4s}{4s+\theta+1}}$.

Theorem 4. *Consider testing $H_0 : P = P_0$ against $H_1(\Delta_n, \theta)$, for some $\theta < 2s - 1$. If $\overline{\lim}_{n \rightarrow \infty} \Delta_n n^{\frac{4s}{4s+\theta+1}} < \infty$, then*

$$\lim_{n \rightarrow \infty} \inf_{T \in \mathcal{T}_n} [\mathbb{E}_{P_0} T + \beta(T; \Delta_n, \theta)] > 0,$$

where \mathcal{T}_n denotes the collection of all test functions based on X_1, \dots, X_n .

Recall that for a test T , $\mathbb{E}_{P_0} T$ is its Type I error. Theorem 4 shows that, if $\Delta_n = O(n^{-4s/(4s+\theta+1)})$, then the sum of Type I and Type II errors of any test does not vanish as n increases. In other words, there is no consistent test if $\Delta_n = O(n^{-4s/(4s+\theta+1)})$. Together with Theorem 3, this suggests that T_{M^3d} is rate optimal in the minimax sense.

4 Adaptation

Despite the minimax optimality of T_{M^3d} , a practical challenge in using it is the choice of an appropriate tuning parameter ϱ_n . In particular, Theorem 3 suggests that ϱ_n needs to be taken at the order of $n^{-2s(\theta+1)/(4s+\theta+1)}$ which depends on the value of s and θ . On the one hand, since P_0 and K are known apriori, so is s . On the other hand, θ reflects the property of dP/dP_0 which is typically not known in advance. This naturally brings about the issue of adaptation (see, *e.g.*, [Spokoiny, 1996; Ingster, 2000](#)). In other words, we are interested in a single testing procedure that can achieve the detection boundary for testing H_0 against $H_1(\Delta_n(\theta), \theta)$ simultaneously over all $\theta \geq 0$. We emphasize the dependence of Δ_n on θ since the detection boundary may depend on θ , as suggested by the results from the previous section. In fact, we should build upon the test statistic introduced before.

More specifically, write

$$\rho_* = \left(\frac{\sqrt{\log \log n}}{n} \right)^{2s},$$

and

$$m_* = \left\lceil \log_2 \left[\rho_*^{-1} \left(\frac{\sqrt{\log \log n}}{n} \right)^{\frac{2s}{4s+1}} \right] \right\rceil.$$

Then our test statistics is taken to be the maximum of T_{n,ϱ_n} for $\rho_n = \rho_*, 2\rho_*, 2^2\rho_*, \dots, 2^{m_*}\rho_*$:

$$\tilde{T}_n := \sup_{0 \leq k \leq m_*} T_{n,2^k \varrho_*},$$

where, with slight abuse of notation,

$$T_{n,\varrho_n} = (2v_n)^{-1/2} [n\eta_{\varrho_n}^2(\hat{P}_n, P_0) - A_n].$$

It turns out if an appropriate rejection threshold is chosen, \tilde{T}_n can achieve a detection boundary very similar to the one we have before, but now simultaneously over all $\theta > 0$.

Theorem 5. (i) Under H_0 ,

$$\lim_{n \rightarrow \infty} P\left(\tilde{T}_n \geq \sqrt{3 \log \log n}\right) = 0;$$

(ii) on the other hand, there exists a constant $c_1 > 0$ such that,

$$\lim_{n \rightarrow \infty} \inf_{P \in \cup_{\theta \geq 0} \mathcal{P}(\Delta_n(\theta))} P\left(\tilde{T}_n \geq \sqrt{3 \log \log n}\right) = 1,$$

provided that $\Delta_n(\theta) \geq c_1(n^{-1} \sqrt{\log \log n})^{\frac{4s}{4s+\theta+1}}$.

Theorem 5 immediately suggests that a test rejects H_0 if and only if $\tilde{T}_n \geq \sqrt{3 \log \log n}$ is consistent for testing it against $H_1(\Delta_n(\theta), \theta)$ for all $\theta \geq 0$ provided that $\Delta_n(\theta) \geq c_1(n^{-1} \sqrt{\log \log n})^{\frac{4s}{4s+\theta+1}}$. We can further calibrate the rejection region to yield a test at a given significance level. More precisely, let \tilde{q}_α be the upper α quantile of \tilde{T}_n , we can proceed to reject H_0 whenever the observed test statistic exceeds \tilde{q}_α . Denote such a test by \tilde{T}_{M^3d} . By definition, \tilde{T}_{M^3d} is an α -level test. Theorem 5 implies that the type II error of \tilde{T}_{M^3d} vanishes as $n \rightarrow \infty$ uniformly over all $\theta \geq 0$. In practice, the quantile \tilde{q}_α can be evaluated by Monte Carlo methods as we shall discuss in further details in the next section. We note that the detection boundary given in Theorem 5 is similar, but inferior by a factor of $(\log \log n)^{\frac{2s}{4s+\theta+1}}$, to that from Theorem 4. This turns out to be the price one needs to pay for adaptation.

Theorem 6. Let $0 < \theta_1 < \theta_2 < 2s - 1$. Then there exists a positive constant c_2 such that

$$\overline{\lim}_{n \rightarrow \infty} \sup_{\theta \in [\theta_1, \theta_2]} \left\{ \Delta_n(\theta) \left(\frac{n}{\sqrt{\log \log n}} \right)^{\frac{4s}{4s+\theta+1}} \right\} \leq c_2$$

implies that

$$\lim_{n \rightarrow \infty} \inf_{T \in \mathcal{T}_n} \left[\mathbb{E}_{P_0} T + \sup_{\theta \in [\theta_1, \theta_2]} \beta(T; \Delta_n(\theta), \theta) \right] = 1.$$

Similar to Theorem 4, Theorem 6 shows that there is no consistent test for H_0 against $H_1(\Delta_n, \theta)$ simultaneously over all $\theta \in [\theta_1, \theta_2]$, if $\Delta_n(\theta) \leq c_2 (n^{-1} \sqrt{\log \log n})^{\frac{4s}{4s+\theta+1}} \forall \theta \in [\theta_1, \theta_2]$ for a sufficiently small c_2 . Together with Theorem 5, this suggests that the test $\tilde{T}_{\text{M}^3\text{d}}$ is indeed rate optimal.

5 Numerical Experiments

To complement the earlier theoretical development, we also performed several sets of simulation experiments to demonstrate the merits of the proposed adaptive test based on \tilde{T}_n . To do so, we need to first address a practical issue of computing the test statistic \tilde{T}_n : how to compute $\eta_{\varrho_n}^2(\hat{P}_n, P_0)$ for a given ϱ_n .

5.1 Computing \tilde{T}_n

Though the form of $\eta_{\varrho_n}^2(\hat{P}_n, P_0)$ looks similar to that of $\gamma^2(\hat{P}_n, P_0)$, from the point of view of computing it numerically, there is a subtle issue. The kernel $\tilde{K}_{\varrho_n}(x, x')$ is defined only in its Mercer decomposed form, which is based on the Mercer decomposition of $K(x, x')$. Hence, in order to compute the kernel $\tilde{K}_{\varrho_n}(x, x')$, we need to first choose a kernel $K(x, x')$ and compute its Mercer decomposition numerically. Specifically, we use *chebfun* framework in Matlab (with slight modifications) to compute Mercer decompositions associated with kernels based on their integral operator representations Driscoll et al. (2014); Trefethen and Battles (2004). Once we compute λ_k and the associated $\varphi_k(\cdot)$, we approximately compute $\tilde{K}_{\varrho_n}(x, x')$ based on the top K eigenvalues and eigenfunctions. This provides a numerical framework for computing $\tilde{K}_{\varrho_n}(x, x')$ once we fix a kernel $K(x, x')$. In the cases when the eigenvalues and eigenfunction are known, for example when using polynomial kernels, from our experiments we found that using the top few numerical eigenvalues gives a good approximation to the actual value of the kernel. Given a way to compute kernel evaluations, computing $\eta_{\varrho_n}^2(\hat{P}_n, P_0)$ follows similarly.

5.2 Power comparison

Once we are able to compute \tilde{T}_n , we can assess its null distribution by simulating it under the null hypothesis H_0 . In particular, we repeated for each case 200 runs and estimated the 95%

quantile of \tilde{T}_n under H_0 by the corresponding sample quantile. We then proceeded to reject H_0 when an observed test statistic exceeds the estimate 95% quantile. By construction, the procedure gives a 5%-level test, up to Monte Carlo error.

Euclidean data: We first consider using both the test T_{MMD} and $\tilde{T}_{\text{M}^3\text{d}}$ to test the hypothesis P_0 is uniform on $[0, 1]^d$ given a sample of observations $\{X_1, \dots, X_n\}$. The dimensionality used are 100 and 200. We followed the examples for densities put forward in [Marron and Wand \(1992\)](#) in the context of nonparametric density estimation, for the alternatives. Specifically we set the alternative hypothesis to be (1) mixture of five Gaussians, (2) skewed unimodal, (3) asymmetric claw density and (4) smooth comb density. The value of α is set to 0.05. The sample size n is varied from 200 to 1000 (in steps of 200) and for each value of sample size 100 simulations are conducted to estimate the probability of rejecting a false null hypothesis.

We use a Gaussian kernel K to compute the $\gamma^2(\hat{P}_n, P_0)$ and use the procedure outlined in [section 5.1](#) to compute $\eta_{\varrho_n}^2(\hat{P}_n, P_0)$. The issue of choosing the kernel is subtle when using T_{MMD} . For simplicity, we fixed the value of bandwidth of Gaussian kernel (which corresponds to choosing the kernel in this case) to a fixed value, that corresponds to the best performance of T_{MMD} . With the fixed value of the bandwidth, to fix ϱ , we tried values over a grid and set it to the value that performed best. [Figure 1](#) illustrates a plot of the estimated probability of accepting the null hypothesis when it is false for different values of sample size n for the proposed test $\tilde{T}_{\text{M}^3\text{d}}$ along with T_{MMD} and the more classical Kolmogorov-Smirnov (K-S, for short) goodness-of-fit test. We note from [Figure 1](#) that the estimated error probability converges to zero at a faster rate for the adaptive M^3D test compared to the MMD test and the Kolmogorov-Smirnov test on all the different simulation settings that are considered. Note that it has been previously observed that MMD test performs better than K-S test in various setting in [Gretton et al. \(2012\)](#), which we observe in our setting as well.

Directional data: One of the advantages of the proposed RKHS embedding based approach is that it could be used on domains other than the d -dimensional Euclidean space. For example when $\mathcal{X} = \mathbb{S}^{d-1}$ where \mathbb{S}^{d-1} corresponds to the d -dimensional unit sphere, one can perform hypothesis testing using the above framework, as long as we can compute the Mercer decomposition of a kernel K defined on the domain. In several applications, like




Figure 1: Error versus Sample Size: mixture of Gaussian (row 1), skewed unimodal (row 2), asymmetric claw (row 3) and smooth comb (row 4) with dimensionality 100 (left) and 200 (right).

protein folding, often times data are modeled as coming from the unit-sphere and testing goodness-of-fit for such data needs specialized methods different from the standard nonparametric testing methods [Mardia and Jupp \(2009\)](#); [Jupp \(2005\)](#).

In order to highlight the advantage of the proposed approach, we assume P_0 is uniform distribution on the unit sphere of dimension 100 and test it against the alternative that data are from:

- (1) multivariate von Mises-Fisher distribution (which is the Gaussian analogue on the unit-sphere) given by $f_{vM-F}(x, \mu, \kappa) = C_{vM-F}(\kappa) \exp(\kappa \mu^\top x)$ for data $x \in \mathbb{S}^{d-1}$, where $\kappa \geq 0$ is concentration parameter and μ is the mean parameter. The term C_{vM-F} is the normalization constant given by $\frac{\kappa^{d/2-1}}{2\pi^{d/2}I_{d/2-1}(\kappa)}$ where I is modified Bessel function;
- (2) multivariate Watson distribution (used to model axially symmetric data on sphere) given by $f_W(x, \mu, \kappa) = C_W(\kappa) \exp(\kappa(\mu^\top x)^2)$ for data $x \in \mathbb{S}^{d-1}$, where $\kappa \geq 0$ is concentration parameter and μ is the mean parameter as before. The term $C_W(\kappa)$ is the normalization constant given by $\frac{\Gamma(d/2)}{2\pi^{d/2}M(1/2, d/2, \kappa)}$ where M is Kummer's confluent hypergeometric function;
- (3) mixture of five von Mises-Fisher distribution which are used in modeling and clustering spherical data [Banerjee et al. \(2005\)](#);
- (4) mixture of five Watson distribution which are used in modeling and clustering spherical data [Sra and Karp \(2013\)](#).

Note that in this setup one can analytically compute the Mercer decomposition of the Gaussian kernel on the unit sphere with respect to the uniform distribution. Specifically, the eigenvalues are given by Theorem 2 in [Minh et al. \(2006\)](#) and the eigenfunctions are the standard spherical harmonics of order k (see section 2.1 in [Minh et al. \(2006\)](#) for details). Rest of the simulation setup is similar to the previous setting (of Euclidean data) and we compared \tilde{T}_{M^3d} against T_{MMD} and the Sobolev test approach (denoted as ST hereafter) proposed in [Jupp \(2005\)](#). Figure 2 illustrates a plot of estimated probability of accepting null hypothesis when it is false for different values of sample size, from which we see the adaptive M³D test performs better.




Figure 2: Error versus Sample Size: von Mises-Fisher distribution (row 1), Watson distribution (row 2), Mixture of von Mises-Fisher distribution (row 3) and mixture of Watson distribution (row 4) on sphere for 100 dimensions (left) and 150 dimensions (right).

5.3 Real data experiments

In addition to the simulation examples, we also performed experiments on several real-world data examples. Similar to before, in the case of Euclidean data, we compared \tilde{T}_{M^3d} against the standard MMD and Kolmogorov-Smirnov test, and in the case of spherical data, we used the Sobolev test [Jupp \(2005\)](#), instead of the Kolmogorov-Smirnov test.

For the case of Euclidean data, we used the MNIST digits data set from the following webpage: <http://yann.lecun.com/exdb/mnist/>. Model-based clustering [Fraley and Raftery \(2002\)](#) is a widely-used and practical successful clustering technique in the literature. Furthermore, the MNIST data set is a standard data set for testing clustering algorithms and consists of image of digits. Several works have implicitly assumed that the data come from a mixture of Gaussian distributions, because of the observed superior empirical performance under such an assumption. But the validity of such a mixture model assumption is invariably not tested statistically. In this experiment we selected three digits (which correspond to a cluster) randomly and conditioned on the selected digit (cluster), we test the hypothesis that the data come from a Gaussian distribution (that is, P_0 is Gaussian). For our experiments, we down sampled the images and use pixels as feature vectors with dimensionality 64 as is commonly done in the literature. Table 1 reports the probability with which the null hypothesis is accepted. The observed result reiterates in a statistically significant way that it is reasonable to make a mixture of Gaussian assumption in this case.

For the spherical case, we use the Human Fibroblasts dataset from [Iyer et al. \(1999\)](#); [Dhillon et al. \(2003\)](#), Yeast Cell Cycle dataset from [Spellman et al. \(1998\)](#) and the Rosetta yeast gene expression dataset [Hughes et al. \(2000\)](#). The Fibroblast data set contains 12 expression data corresponding to 517 samples (genes) report in the response of human fibroblasts following addition of serum to the growth media. We refer to [Iyer et al. \(1999\)](#) for more details about the scientific procedure with which these data were obtained. The Yeast Cell Cycle dataset consists of 82-dimensional data corresponding to 696 subjects. The Rosetta yeast dataset contains 300-dimensional element vector for around 6000 yeast genes. Previous data analysis studies [Sra and Karp \(2013\)](#); [Dhillon et al. \(2003\)](#) have used mixtures of spherical distributions for clustering the above data set. Specifically, it has been observed in [Sra and Karp \(2013\)](#) that clustering using a mixture of Watson distribution has superior performance. While that has proved to be useful scientifically, it was not statistically tested

Sample size =	300	400	500	300	400	500	300	400	500
K-S	0.86	0.91	0.94	0.83	0.88	0.93	0.84	0.88	0.92
MMD	0.90	0.93	0.95	0.89	0.92	0.95	0.88	0.92	0.94
M^3D	0.94	0.96	0.98	0.93	0.95	0.98	0.93	0.95	0.98

Table 1: The values reported are the estimated probability with which the corresponding hypothesis test accepts the null hypothesis when it is true. The level of the test $\alpha = 0.05$. Digit 4 on left, Digit 6 on the middle and Digit 7 on right, for various values of sample size.

Sample size =	75	150	200	150	200	250	400	500	600
ST	0.87	0.93	0.98	0.82	0.87	0.91	0.76	0.84	0.91
MMD	0.90	0.94	0.98	0.85	0.92	0.94	0.79	0.87	0.92
M^3D	0.92	0.96	0.99	0.88	0.93	0.96	0.81	0.90	0.95

Table 2: The values reported are the estimated probability with which the corresponding hypothesis test accepts the null hypothesis when it is true. The level of the test $\alpha = 0.05$. Human Fibroblasts dataset on left, Yeast Cell Cycle dataset on the middle and Rosetta Yeast dataset on the right, for various values of sample size.

if such an assumption is valid. Here, we test for goodness of fit of Watson distribution (that is, P_0 is a Watson distribution) for the largest cluster from the above data sets. Table 2 shows the estimated probability of acceptance of the null hypothesis when it is assumed to be true. The values reported are averages from 50 random trails of the same dataset. The observed results provide a statistical justification for the use of Watson distribution in modeling the above data sets.

We note that for both situations, the tests considered tend to agree that the true hypothesis is true when there are more samples as indicated by Table 1 and 2. But, the probability of acceptance is higher for low sample sizes for the adaptive M^3D test \tilde{T}_{M^3D} in all cases, showing that the method works better with finite sample sizes. This highlights the advantage of the \tilde{T}_{M^3D} in a finite sample setting confirming the better rates of convergence obtained in theory.

6 Proofs

Proof of Theorem 1. Part (i). The proof of the first part consists of two key steps. First, we show that the population counterpart $n\gamma^2(P, P_0)$ of the test statistic converges to ∞ uniformly, *i.e.*,

$$n \inf_{P \in \mathcal{P}(\Delta_n, 0)} \gamma^2(P, P_0) \rightarrow \infty.$$

Then, we argue that the deviation from $\gamma^2(P, P_0)$ to $\gamma^2(\hat{P}_n, P_0)$ is uniformly negligible compared with $\gamma^2(P, P_0)$ itself.

It is not hard to see that

$$\begin{aligned} \gamma(\hat{P}_n, P_0) &= \sqrt{\sum_{k \geq 1} \lambda_k \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) \right]^2} \\ &\geq \sqrt{\sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2} - \sqrt{\sum_{k \geq 1} \lambda_k \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) - \mathbb{E}_P \varphi_k(X) \right]^2}. \end{aligned}$$

Thus,

$$\begin{aligned} &P \left\{ n\gamma^2(\hat{P}_n, P_0) < q_{w, 1-\alpha} \right\} \\ &\leq P \left\{ \sqrt{n \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2} - \sqrt{n \sum_{k \geq 1} \lambda_k \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) - \mathbb{E}_P \varphi_k(X) \right]^2} < \sqrt{q_{w, 1-\alpha}} \right\} \\ &= P \left\{ \sqrt{n \sum_{k \geq 1} \lambda_k \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) - \mathbb{E}_P \varphi_k(X) \right]^2} > \sqrt{n \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2} - \sqrt{q_{w, 1-\alpha}} \right\}. \end{aligned}$$

Suppose that

$$n \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2 > q_{w, 1-\alpha}.$$

Then

$$P \left\{ n\gamma^2(\hat{P}_n, P_0) < q_{w, 1-\alpha} \right\} \leq \frac{\mathbb{E}_P \left\{ n \sum_{k \geq 1} \lambda_k \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) - \mathbb{E}_P \varphi_k(X) \right]^2 \right\}}{\left\{ \sqrt{n \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2} - \sqrt{q_{w, 1-\alpha}} \right\}^2}.$$

Observe that for any $P \in \mathcal{P}(\Delta_n, 0)$,

$$\begin{aligned}\mathbb{E}_P \left\{ n \sum_{k \geq 1} \lambda_k \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) - \mathbb{E}_P \varphi_k(X) \right]^2 \right\} &= \sum_{k \geq 1} \lambda_k \text{Var}[\varphi_k(X)] \\ &\leq \sum_{k \geq 1} \lambda_k \mathbb{E}_P \varphi_k^2(X) \\ &\leq \left(\sup_{k \geq 1} \|\varphi_k\|_\infty \right)^2 \sum_{k \geq 1} \lambda_k < \infty.\end{aligned}$$

This implies that

$$\begin{aligned}\lim_{n \rightarrow \infty} \beta(T_{\text{MMD}}; \Delta_n, 0) &= \lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}(\Delta_n, 0)} P \left\{ n\gamma^2(\hat{P}_n, P_0) < q_{w, 1-\alpha} \right\} \\ &\leq \lim_{n \rightarrow \infty} \frac{\sup_{P \in \mathcal{P}(\Delta_n, 0)} \mathbb{E}_P \left\{ n \sum_{k \geq 1} \lambda_k \left[\frac{1}{n} \sum_{i=1}^n \varphi_k(X_i) - \mathbb{E}_P \varphi_k(X) \right]^2 \right\}}{\inf_{P \in \mathcal{P}(\Delta_n, 0)} \left\{ \sqrt{n \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2} - \sqrt{q_{w, 1-\alpha}} \right\}^2} \\ &= 0,\end{aligned}$$

provided that

$$\inf_{P \in \mathcal{P}(\Delta_n, 0)} n \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2 \rightarrow \infty, \quad \text{as } n \rightarrow \infty. \quad (9)$$

It now suffices to show that (9) holds if $n\Delta_n^2 \rightarrow \infty$ as $n \rightarrow \infty$.

To this end, let $u = dP/dP_0 - 1$ and

$$a_k = \langle u, \varphi_k \rangle_{L_2(P_0)} = \mathbb{E}_P \varphi_k(X) - \mathbb{E}_{P_0} \varphi_k(X) = \mathbb{E}_P(\varphi_k(X)).$$

It is clear the that

$$\sum_{k \geq 1} \lambda_k^{-1} a_k^2 = \|u\|_K^2, \quad \text{and} \quad \sum_{k \geq 1} a_k^2 = \|u\|_{L_2(P_0)}^2 = \chi^2(P, P_0).$$

By the definition of $\mathcal{P}(\Delta_n, 0)$,

$$\sup_{P \in \mathcal{P}(\Delta_n, 0)} \sum_{k \geq 1} \lambda_k^{-1} a_k^2 \leq M^2, \quad \text{and} \quad \inf_{P \in \mathcal{P}(\Delta_n, 0)} \sum_{k \geq 1} a_k^2 \geq \Delta_n.$$

Since $n\Delta_n^2 \rightarrow \infty$ as $n \rightarrow \infty$, we get

$$\begin{aligned} \inf_{P \in \mathcal{P}(\Delta_n, 0)} n \sum_{k \geq 1} \lambda_k [\mathbb{E}_P \varphi_k(X)]^2 &= \inf_{P \in \mathcal{P}(\Delta_n, 0)} n \sum_{k \geq 1} \lambda_k a_k^2 \\ &\geq \inf_{P \in \mathcal{P}(\Delta_n, 0)} n \frac{\left(\sum_{k \geq 1} a_k^2 \right)^2}{\sum_{k \geq 1} \lambda_k^{-1} a_k^2} \\ &\geq \frac{n\Delta_n^2}{M^2} \rightarrow \infty \end{aligned}$$

as $n \rightarrow \infty$.

Part (ii). In proving the second part, we will make use of the following lemma that can be obtained by adapting the argument in [Gregory \(1977\)](#). It gives the limit distribution of V-statistic under P_n such that P_n converges to P_0 in the order $n^{-1/2}$.

Lemma 1. *Consider a sequence of probability measures $\{P_n : n \geq 1\}$ contiguous to P_0 satisfying $u_n = dP_n/dP_0 - 1 \rightarrow 0$ in $L^2(P_0)$. Suppose that for any fixed k ,*

$$\lim_{n \rightarrow \infty} \sqrt{n} \langle u_n, \varphi_k \rangle_{L^2(P_0)} = \tilde{a}_k, \quad \text{and} \quad \lim_{n \rightarrow \infty} \sum_{k \geq 1} \lambda_k (\sqrt{n} \langle u_n, \varphi_k \rangle_{L^2(P_0)})^2 = \sum_{k \geq 1} \lambda_k \tilde{a}_k^2 + \tilde{a}_0 < \infty,$$

for some sequence $\{\tilde{a}_k : k \geq 0\}$, then

$$\frac{1}{n} \sum_{k \geq 1} \lambda_k \left[\sum_{i=1}^n \varphi_k(X_i) \right]^2 \xrightarrow{d} \sum_{k \geq 1} \lambda_k (Z_k + \tilde{a}_k)^2 + \tilde{a}_0,$$

where $X_1, \dots, X_n \stackrel{i.i.d}{\sim} P_n$, and Z_k s are independent standard normal random variables.

Write $L(k) = \lambda_k k^{2s}$. By assumption (5),

$$0 < \underline{L} := \inf_{k \geq 1} L(k) \leq \sup_{k \geq 1} L(k) := \bar{L} < \infty.$$

Consider a sequence of $\{P_n : n \geq 1\}$ such that

$$dP_n/dP_0 - 1 = C_1 \sqrt{\lambda_{k_n}} [L(k_n)]^{-1} \varphi_{k_n},$$

where C_1 is a positive constant and $k_n = \lfloor C_2 n^{\frac{1}{4s}} \rfloor$ for some positive constant C_2 . Both C_1 and C_2 will be determined later. Since $\sup_{k \geq 1} \|\varphi_k\|_\infty < \infty$ and $\lim_{k \rightarrow \infty} \lambda_k = 0$, there exists $N_0 > 0$ such that P_n 's are well-defined probability measures for any $n \geq N_0$.

Note that

$$\|u_n\|_K^2 = \frac{C_1^2}{L^2(k_n)} \leq \underline{L}^{-2} C_1^2$$

and

$$\|u_n\|_{L^2(P_0)}^2 = \frac{C_1^2 \lambda_{k_n}}{L^2(k_n)} = \frac{C_1^2}{L(k_n)} k_n^{-2s} \geq \bar{L}^{-1} C_1^2 k_n^{-2s} \sim \bar{L}^{-1} C_1^2 C_2^{-2s} n^{-1/2},$$

where $A_n \sim B_n$ means that $\lim_{n \rightarrow \infty} A_n/B_n = 1$. Thus, by choosing C_1 sufficiently small and $c_0 = \frac{1}{2} \bar{L}^{-1} C_1^2 C_2^{-2s}$, we ensure that $P_n \in \mathcal{P}(c_0 n^{-1/2}, 0)$ for sufficiently large n .

To apply Lemma 1, we note that

$$\lim_{n \rightarrow \infty} \|u_n\|_{L^2(P_0)}^2 = \lim_{n \rightarrow \infty} \frac{C_1^2 \lambda_{k_n}}{L^2(k_n)} = 0.$$

In addition, for any fixed k ,

$$\tilde{a}_{n,k} = \sqrt{n} \langle u_n, \varphi_k \rangle_{L^2(P_0)} = 0$$

for sufficiently large n , and

$$\sum_{k \geq 1} \lambda_k \tilde{a}_{n,k}^2 = \frac{n C_1^2 \lambda_{k_n}^2}{L^2(k_n)} = n C_1^2 k_n^{-4s} \rightarrow C_1^2 C_2^{-4s}$$

as $n \rightarrow \infty$. Thus, Lemma 1 implies that

$$n\gamma(\hat{P}_n, P_0) \xrightarrow{d} \sum_{k \geq 1} \lambda_k Z_k^2 + C_1^2 C_2^{-4s}.$$

Now take $C_2 = (2C_1^2/q_{w,1-\alpha})^{1/4s}$ so that $C_1^2 C_2^{-4s} = \frac{1}{2} q_{w,1-\alpha}$. Then

$$\begin{aligned} \lim_{n \rightarrow \infty} \beta(T_{\text{MMD}}; c_0 n^{-1/2}, 0) &\geq \lim_{n \rightarrow \infty} P_n(n\gamma(\hat{P}_n, P_0) < q_{w,1-\alpha}) \\ &= P\left(\sum_{k \geq 1} \lambda_k Z_k^2 < \frac{1}{2} q_{w,1-\alpha}\right) > 0, \end{aligned}$$

which concludes the proof. □

Proof of Theorem 2. Let $\tilde{K}_n(\cdot, \cdot) := \tilde{K}_{\varrho_n}(\cdot, \cdot)$. Note that

$$\begin{aligned} n\hat{\eta}_{\varrho_n}^2(P, P_0) &= \frac{1}{n} \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} \left[\sum_{i=1}^n \varphi_k(X_i) \right]^2 \\ &= \frac{1}{n} \sum_{i,j=1}^n \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} \varphi_k(X_i) \varphi_k(X_j) \\ &= \frac{1}{n} \sum_{i,j=1}^n \tilde{K}_n(X_i, X_j). \end{aligned}$$

Thus

$$v_n^{-1/2} [n\hat{\eta}_{\varrho_n}^2(P, P_0) - A_n] = 2(n^2 v_n)^{-1/2} \sum_{j=2}^n \sum_{i=1}^{j-1} \tilde{K}_n(X_i, X_j).$$

Let $\zeta_{nj} = \sum_{i=1}^{j-1} \tilde{K}_n(X_i, X_j)$. Consider a filtration $\{\mathcal{F}_j : j \geq 1\}$ where $\mathcal{F}_j = \sigma\{X_i : 1 \leq i \leq j\}$. Due to the assumption that K is degenerate, we have $\mathbb{E}\varphi_k(X) = 0$ for any $k \geq 1$, which implies that

$$\mathbb{E}(\zeta_{nj} | \mathcal{F}_{j-1}) = \sum_{i=1}^{j-1} \mathbb{E}[\tilde{K}_n(X_i, X_j) | \mathcal{F}_{j-1}] = \sum_{i=1}^{j-1} \mathbb{E}[\tilde{K}_n(X_i, X_j) | X_i] = 0,$$

for any $j \geq 2$.

Write

$$U_{nm} = \begin{cases} 0 & m = 1 \\ \sum_{j=2}^m \zeta_{nj} & m \geq 2 \end{cases}.$$

Then for any fixed n , $\{U_{nm}\}_{m \geq 1}$ is a martingale with respect to $\{\mathcal{F}_m : m \geq 1\}$ and

$$v_n^{-1/2} [n\hat{\eta}_{\varrho_n}^2(P, P_0) - A_n] = 2(n^2 v_n)^{-1/2} U_{nn}.$$

We now apply martingale central limit theorem to U_{nn} . Following the argument from [Hall \(1984\)](#), it can be shown that

$$\left[\frac{1}{2} n^2 \mathbb{E} \tilde{K}_n^2(X, X') \right]^{-1/2} U_{nn} \xrightarrow{d} N(0, 1), \quad (10)$$

provided that

$$[\mathbb{E} G_n^2(X, X') + n^{-1} \mathbb{E} \tilde{K}_n^2(X, X') \tilde{K}_n^2(X, X'') + n^{-2} \mathbb{E} \tilde{K}_n^4(X, X')] / [\mathbb{E} \tilde{K}_n^2(X, X')]^2 \rightarrow 0, \quad (11)$$

as $n \rightarrow \infty$, where $G_n(x, x') = \mathbb{E}\tilde{K}_n(X, x)\tilde{K}_n(X, x')$. Since

$$\mathbb{E}\tilde{K}_n^2(X, X') = \sum_{k \geq 1} \left(\frac{\lambda_k}{\lambda_k + \varrho_n^2} \right)^2 = v_n,$$

(10) implies that

$$v_n^{-1/2} [n\hat{\eta}_{\varrho_n}^2(P, P_0) - A_n] = \sqrt{2} \cdot \left(\frac{1}{2} n^2 \mathbb{E}\tilde{K}_n^2(X, X') \right)^{-1/2} U_{nn} \xrightarrow{d} N(0, 2).$$

It therefore suffices to verify (11).

Note that

$$\begin{aligned} \mathbb{E}\tilde{K}_n^2(X, X') &= \sum_{k \geq 1} \left(\frac{\lambda_k}{\lambda_k + \varrho_n^2} \right)^2 \geq \sum_{\lambda_k \geq \varrho_n^2} \frac{1}{4} + \frac{1}{4\varrho_n^4} \sum_{\lambda_k < \varrho_n^2} \lambda_k^2 \\ &= \frac{1}{4} |\{k : \lambda_k \geq \varrho_n^2\}| + \frac{1}{4\varrho_n^4} \sum_{\lambda_k < \varrho_n^2} \lambda_k^2 \asymp \varrho_n^{-1/s}, \end{aligned}$$

where the last step holds by considering that $\lambda_k \asymp k^{-2s}$. Hereafter, we shall write $a_n \asymp b_n$ if $0 < \varliminf_{n \rightarrow \infty} a_n/b_n \leq \overline{\lim}_{n \rightarrow \infty} a_n/b_n < \infty$, for two positive sequences $\{a_n\}$ and $\{b_n\}$. Similarly,

$$\mathbb{E}G_n^2(X, X') = \sum_{k \geq 1} \left(\frac{\lambda_k}{\lambda_k + \varrho_n^2} \right)^4 \leq |\{k : \lambda_k \geq \varrho_n^2\}| + \varrho_n^{-8} \sum_{\lambda_k < \varrho_n^2} \lambda_k^4 \asymp \varrho_n^{-1/s},$$

and

$$\begin{aligned} \mathbb{E}\tilde{K}_n^2(X, X')\tilde{K}_n^2(X, X'') &= \mathbb{E} \left\{ \sum_{k \geq 1} \left(\frac{\lambda_k}{\lambda_k + \varrho_n^2} \right)^2 \varphi_k^2(X) \right\}^2 \\ &\leq \left(\sup_{k \geq 1} \|\varphi_k\|_\infty \right)^4 \left\{ \sum_{k \geq 1} \left(\frac{\lambda_k}{\lambda_k + \varrho_n^2} \right)^2 \right\}^2 \asymp \varrho_n^{-2/s}. \end{aligned}$$

Thus there exists a positive constant C_3 such that

$$\mathbb{E}G_n^2(X, X') / [\mathbb{E}\tilde{K}_n^2(X, X')]^2 \leq C_3 \varrho_n^{1/s} \rightarrow 0, \quad (12)$$

and

$$n^{-1} \mathbb{E}\tilde{K}_n^2(X, X')\tilde{K}_n^2(X, X'') / [\mathbb{E}\tilde{K}_n^2(X, X')]^2 \leq C_3 n^{-1} \rightarrow \infty, \quad (13)$$

as $n \rightarrow \infty$. On the other hand,

$$\mathbb{E}\tilde{K}_n^4(X, X') \leq \|\tilde{K}_n\|_\infty^2 \mathbb{E}\tilde{K}_n^2(X, X'),$$

where

$$\|\tilde{K}_n\|_\infty = \sup_x \left\{ \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} \varphi_k^2(x) \right\} \leq \left(\sup_{k \geq 1} \|\varphi_k\|_\infty \right)^2 \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} \asymp \varrho_n^{-1/s}.$$

This implies that for some positive constant C_4 ,

$$n^{-2} \mathbb{E} \tilde{K}_n^4(X, X') / [\mathbb{E} \tilde{K}_n^2(X, X')]^2 \leq n^{-2} \|\tilde{K}_n\|_\infty^2 / \mathbb{E} \tilde{K}_n^2(X, X') \leq C_4 (n^2 \varrho_n^{1/s})^{-1} \rightarrow 0. \quad (14)$$

as $n \rightarrow \infty$. Together, (12), (13) and (14) ensure that condition (11) holds. \square

Proof of Theorem 3. Note that

$$\begin{aligned} & n\eta_{\varrho_n}^2 (\hat{P}_n, P_0) - \frac{1}{n} \sum_{i=1}^n \tilde{K}_n(X_i, X_j) \\ &= \frac{1}{n} \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} \sum_{\substack{1 \leq i, j \leq n \\ i \neq j}} \varphi_k(X_i) \varphi_k(X_j) \\ &= \frac{1}{n} \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} \sum_{\substack{1 \leq i, j \leq n \\ i \neq j}} [\varphi_k(X_i) - \mathbb{E}_P \varphi_k(X)] [\varphi_k(X_j) - \mathbb{E}_P \varphi_k(X)] \\ &\quad + \frac{2(n-1)}{n} \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] \sum_{1 \leq i \leq n} [\varphi_k(X_i) - \mathbb{E}_P \varphi_k(X)] \\ &\quad + \frac{n(n-1)}{n} \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)]^2 \\ &:= V_1 + V_2 + V_3. \end{aligned}$$

Obviously, $\mathbb{E}_P V_1 V_2 = 0$. We first argue that the following three statements together implies the desired result:

$$\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} v_n^{-1/2} V_3 = \infty, \quad (15)$$

$$\sup_{P \in \mathcal{P}(\Delta_n, \theta)} (\mathbb{E}_P V_1^2 / V_3^2) = o(1), \quad (16)$$

$$\sup_{P \in \mathcal{P}(\Delta_n, \theta)} (\mathbb{E}_P V_2^2 / V_3^2) = o(1). \quad (17)$$

To see this, note that (15) implies that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} P(v_n^{-1/2} [n\hat{\eta}_{\varrho_n}^2(P, P_0) - A_n] \geq \sqrt{2}z_{1-\alpha}) \\
& \geq \lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} P\left(v_n^{-1/2} V_3 \geq 2\sqrt{2}z_{1-\alpha}, V_1 + V_2 + V_3 \geq \frac{1}{2}V_3\right) \\
& = \lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} P\left(V_1 + V_2 + V_3 \geq \frac{1}{2}V_3\right).
\end{aligned}$$

On the other hand, (16) and (17) imply that

$$\begin{aligned}
\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} P\left(V_1 + V_2 + V_3 \geq \frac{1}{2}V_3\right) &= 1 - \lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}(\Delta_n, \theta)} P\left(V_1 + V_2 + V_3 < \frac{1}{2}V_3\right) \\
&\geq 1 - \lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}(\Delta_n, \theta)} \frac{\mathbb{E}_P(V_1 + V_2)^2}{(V_3/2)^2} = 1.
\end{aligned}$$

This immediately suggests that $T_{M^3_d}$ is consistent. We now show that (15)-(17) indeed hold.

Verifying (15). We begin with (15). Since $v_n \asymp \varrho_n^{-1/s}$ and $V_3 = (n-1)\eta_{\varrho_n}^2(P, P_0)$, (15) is equivalent to

$$\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} n\varrho_n^{\frac{1}{2s}} \eta_{\varrho_n}^2(P, P_0) = \infty.$$

For any $P \in \mathcal{P}(\Delta_n, \theta)$, let $u = dP/dP_0 - 1$ and $a_k = \langle u, \varphi_k \rangle_{L_2(P_0)} = \mathbb{E}_P \varphi_k(X)$. Based on the assumption that K is universal, $u = \sum_{k \geq 1} a_k \varphi_k$. We consider the case $\theta = 0$ and $\theta > 0$ separately.

(1) First consider $\theta = 0$. It is clear that

$$\begin{aligned}
\eta_{\varrho_n}^2(P, P_0) &= \sum_{k \geq 1} a_k^2 - \sum_{k \geq 1} \frac{\varrho_n^2}{\lambda_k + \varrho_n^2} a_k^2 \\
&\geq \|u\|_{L_2(P_0)}^2 - \varrho_n^2 \sum_{k \geq 1} \frac{1}{\lambda_k} a_k^2 \\
&\geq \|u\|_{L_2(P_0)}^2 - \varrho_n^2 M^2.
\end{aligned}$$

Take $\varrho_n \leq \sqrt{\Delta_n/(2M^2)}$ so that $\varrho_n^2 M^2 \leq \frac{1}{2}\Delta_n$. Then we have

$$\inf_{P \in \mathcal{P}(\Delta_n, 0)} \eta_{\varrho_n}^2(P, P_0) \geq \frac{1}{2} \inf_{P \in \mathcal{P}(\Delta_n, 0)} \|u\|_{L_2(P_0)}^2 = \frac{1}{2} \Delta_n.$$

(2) Now consider the case when $\theta > 0$. For $P \in \mathcal{P}(\Delta_n, \theta)$, $\forall R > 0$, $\exists f_R \in \mathcal{H}(K)$ such that $\|u - f_R\|_{L_2(P_0)} \leq MR^{-1/\theta}$ and $\|f_R\|_K \leq R$. Let $b_k = \langle f_R, \varphi_k \rangle_{L_2(P_0)}$.

$$\begin{aligned}
\eta_{\varrho_n}^2(P, P_0) &= \sum_{k \geq 1} a_k^2 - \sum_{k \geq 1} \frac{\varrho_n^2}{\lambda_k + \varrho_n^2} a_k^2 \\
&\geq \|u\|_{L_2(P_0)}^2 - 2 \sum_{k \geq 1} \frac{\varrho_n^2}{\lambda_k + \varrho_n^2} (a_k - b_k)^2 - 2 \sum_{k \geq 1} \frac{\varrho_n^2}{\lambda_k + \varrho_n^2} b_k^2 \\
&\geq \|u\|_{L_2(P_0)}^2 - 2 \sum_{k \geq 1} (a_k - b_k)^2 - 2\varrho_n^2 \sum_{k \geq 1} \frac{1}{\lambda_k} b_k^2 \\
&= \|u\|_{L_2(P_0)}^2 - 2\|u - f_R\|_{L_2(P_0)}^2 - 2\varrho_n^2 \|f_R\|_K^2.
\end{aligned}$$

Taking $R = (2M/\|u\|_{L_2(P_0)})^\theta$ yields that

$$\eta_{\varrho_n}^2(P, P_0) \geq \|u\|_{L_2(P_0)}^2 - 2M^2 R^{-2/\theta} - 2\varrho_n^2 R^2 = \frac{1}{2} \|u\|_{L_2(P_0)}^2 - 2\varrho_n^2 R^2.$$

Now by choosing

$$\varrho_n \leq \frac{1}{2\sqrt{2}} (2M)^{-\theta} \Delta_n^{\frac{1+\theta}{2}},$$

we can ensure that

$$2\varrho_n^2 R^2 \leq \frac{1}{4} \|u\|_{L_2(P_0)}^2.$$

So that

$$\inf_{P \in \mathcal{P}(\Delta_n, \theta)} \eta_{\varrho_n}^2(P, P_0) \geq \inf_{P \in \mathcal{P}(\Delta_n, \theta)} \frac{1}{4} \|u\|_{L_2(P_0)}^2 \geq \frac{1}{4} \Delta_n.$$

In both cases, with $\varrho_n \leq C\Delta_n^{\frac{\theta+1}{2}}$ for a sufficiently small $C = C(M) > 0$, $\lim_{n \rightarrow \infty} \varrho_n^{\frac{1}{2s}} n \Delta_n = \infty$ suffices to ensure (15) holds. Under the condition that $\lim_{n \rightarrow \infty} \Delta_n n^{\frac{4s}{4s+\theta+1}} = \infty$,

$$\varrho_n = cn^{-\frac{2s(\theta+1)}{4s+\theta+1}} \leq C\Delta_n^{\frac{\theta+1}{2}}$$

for sufficiently large n and $\lim_{n \rightarrow \infty} \varrho_n^{\frac{1}{2s}} n \Delta_n = \infty$ holds as well.

Verifying (16). Rewrite V_1 as

$$\begin{aligned}
V_1 &= \frac{1}{n} \sum_{\substack{1 \leq i, j \leq n \\ i \neq j}} \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\varphi_k(X_i) - \mathbb{E}_P \varphi_k(X)] [\varphi_k(X_j) - \mathbb{E}_P \varphi_k(X)] \\
&:= \frac{1}{n} \sum_{\substack{1 \leq i, j \leq n \\ i \neq j}} F_n(X_i, X_j).
\end{aligned}$$

Then

$$\begin{aligned}
\mathbb{E}_P V_1^2 &= \frac{1}{n^2} \sum_{\substack{i \neq j \\ i' \neq j'}} \mathbb{E}_P F_n(X_i, X_j) F_n(X_{i'}, X_{j'}) \\
&= \frac{2n(n-1)}{n^2} \mathbb{E}_P F_n^2(X, X') \\
&\leq 2 \mathbb{E}_P F_n^2(X, X').
\end{aligned}$$

Recall that, for any two random variables Y_1, Y_2 such that $\mathbb{E}Y_1^2 < \infty$,

$$\mathbb{E}[Y_1 - \mathbb{E}(Y_1|Y_2)]^2 = \mathbb{E}Y_1^2 - \mathbb{E}[\mathbb{E}(Y_1|Y_2)^2] \leq \mathbb{E}Y_1^2.$$

Therefore,

$$\mathbb{E}_P F_n^2(X, X') \leq \mathbb{E}_P \{\tilde{K}_n(X, X') - \mathbb{E}_P[\tilde{K}_n(X, X')|X]\}^2 \leq \mathbb{E}_P \tilde{K}_n^2(X, X').$$

Thus, to prove (16), it suffices to show that

$$\lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}(\Delta_n, \theta)} \mathbb{E}_P \tilde{K}_n^2(X, X') / V_3^2 = 0.$$

For any $g \in L_2(P_0)$ and positive definite kernel $G(\cdot, \cdot)$ such that $\mathbb{E}_{P_0} G^2(X, X') < \infty$, let

$$\|g\|_G := \sqrt{\mathbb{E}_{P_0}[g(X)g(X')G(X, X')]}.$$

By the positive definiteness of $G(\cdot, \cdot)$, triangular inequality holds for $\|\cdot\|_G$, i.e., for any $g_1, g_2 \in L_2(P_0)$,

$$|\|g_1\|_G - \|g_2\|_G| \leq \|g_1 - g_2\|_G,$$

which implies that

$$\left| \sqrt{\mathbb{E}_P \tilde{K}_n^2(X, X')} - \sqrt{\mathbb{E}_{P_0} \tilde{K}_n^2(X, X')} \right| \leq \sqrt{\mathbb{E}_{P_0}[u(X)u(X')\tilde{K}_n^2(X, X')]}. \quad (18)$$

We now appeal to the following lemma to bound the right hand side of (18):

Lemma 2. *Let G be a Mercer kernel defined over $\mathcal{X} \times \mathcal{X}$ with eigenvalue-eigenfunction pairs $\{(\mu_k, \varphi_k) : k \geq 1\}$ with respect to $L_2(P)$ such that $\mu_1 \geq \mu_2 \geq \dots$. If G is a trace kernel in that $\mathbb{E}G(X, X) < \infty$, then for any $g \in L_2(P)$*

$$\mathbb{E}_P[g(X)g(X')G^2(X, X')] \leq \mu_1 \left(\sum_{k \geq 1} \mu_k \right) \left(\sup_{k \geq 1} \|\varphi_k\|_\infty \right)^2 \|g\|_{L_2(P)}^2.$$

By Lemma 2, we get

$$\mathbb{E}_{P_0}[u(X)u(X')\tilde{K}_n^2(X, X')] \leq C_5 \left(\sum_k \frac{\lambda_k}{\lambda_k + \varrho_n^2} \right) \|u\|_{L_2(P_0)}^2 \asymp \varrho_n^{-1/s} \|u\|_{L_2(P_0)}^2.$$

Recall that

$$\mathbb{E}_{P_0}\tilde{K}_n^2(X, X') = \sum_k \left(\frac{\lambda_k}{\lambda_k + \varrho_n^2} \right)^2 \asymp \varrho_n^{-1/s}.$$

In the light of (18), they imply that

$$\mathbb{E}_P \tilde{K}_n^2(X, X') \leq 2\{\mathbb{E}_{P_0}\tilde{K}_n^2(X, X') + \mathbb{E}_{P_0}[u(X)u(X')\tilde{K}_n^2(X, X')]\} \leq C_6 \varrho_n^{-1/s} [1 + \|u\|_{L_2(P_0)}^2].$$

On the other hand, it is not hard to verify that with our choice of ϱ_n ,

$$\frac{1}{4} \|u\|_{L_2(P_0)}^2 \leq \eta_{\varrho_n}^2(P, P_0) \leq \|u\|_{L_2(P_0)}^2,$$

for any $P \in \mathcal{P}(\Delta_n, \theta)$. Thus

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}(\Delta_n, \theta)} \mathbb{E}_P \tilde{K}_n^2(X, X') / V_3^2 \\ & \leq 16C_6 \left\{ \left(\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} \varrho_n^{1/s} n^2 \|u\|_{L_2(P_0)}^4 \right)^{-1} + \left(\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} \varrho_n^{1/s} n^2 \|u\|_{L_2(P_0)}^2 \right)^{-1} \right\} = 0 \end{aligned}$$

provided that $\lim_{n \rightarrow \infty} n^{\frac{4s}{4s+\theta+1}} \Delta_n = \infty$. This immediately implies (16).

Verifying (17). Observe that

$$\begin{aligned} \mathbb{E}_P V_2^2 & \leq 4n \mathbb{E}_P \left\{ \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] [\varphi_k(X) - \mathbb{E}_P \varphi_k(X)] \right\}^2 \\ & \leq 4n \mathbb{E}_P \left\{ \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] [\varphi_k(X)] \right\}^2 \\ & = 4n \mathbb{E}_{P_0} \left([1 + u(X)] \left\{ \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] [\varphi_k(X)] \right\}^2 \right). \end{aligned}$$

It is clear that

$$\begin{aligned} & \mathbb{E}_{P_0} \left\{ \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] [\varphi_k(X)] \right\}^2 \\ & = \sum_{k, k' \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} \frac{\lambda_{k'}}{\lambda_{k'} + \varrho_n^2} \mathbb{E}_P \varphi_k(X) \mathbb{E}_P \varphi_{k'}(X) \mathbb{E}_{P_0} [\varphi_k(X) \varphi_{k'}(X)] \\ & = \sum_{k \geq 1} \left(\frac{\lambda_k}{\lambda_k + \varrho_n^2} \right)^2 [\mathbb{E}_P \varphi_k(X)]^2 \leq \eta_{\varrho_n}^2(P, P_0). \end{aligned}$$

On the other hand,

$$\begin{aligned}
& \mathbb{E}_{P_0} \left(u(X) \left\{ \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] [\varphi_k(X)] \right\}^2 \right) \\
& \leq \sqrt{\mathbb{E}_{P_0} \left(u^2(X) \left\{ \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] [\varphi_k(X)] \right\}^2 \right)} \times \\
& \quad \times \sqrt{\mathbb{E}_{P_0} \left\{ \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] [\varphi_k(X)] \right\}^2} \\
& \leq \|u\|_{L_2(P_0)} \sup_x \left| \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)] [\varphi_k(x)] \right| \cdot \eta_{\varrho_n}(P, P_0) \\
& \leq \left(\sup_k \|\varphi_k\|_\infty \right) \|u\|_{L_2(P_0)} \sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} |\mathbb{E}_P \varphi_k(X)| \cdot \eta_{\varrho_n}(P, P_0) \\
& \leq \left(\sup_k \|\varphi_k\|_\infty \right) \|u\|_{L_2(P_0)} \sqrt{\sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2}} \sqrt{\sum_{k \geq 1} \frac{\lambda_k}{\lambda_k + \varrho_n^2} [\mathbb{E}_P \varphi_k(X)]^2} \cdot \eta_{\varrho_n}(P, P_0) \\
& \leq C_7 \|u\|_{L_2(P_0)} \varrho_n^{-\frac{1}{2s}} \eta_{\varrho_n}^2(P, P_0).
\end{aligned}$$

Together, they imply that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}(\Delta_n, \theta)} \mathbb{E}_P V_1^2 / V_3^2 \\
& \leq 4 \max\{1, C_7\} \left\{ \left(\lim_{n \rightarrow \infty} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} n \eta_{\varrho_n}^2(P, P_0) \right)^{-1} + \lim_{n \rightarrow \infty} \sup_{P \in \mathcal{P}(\Delta_n, \theta)} \left(\frac{\|u\|_{L_2(P_0)}}{\varrho_n^{\frac{1}{2s}} n \eta_{\varrho_n}^2(P, P_0)} \right) \right\} = 0,
\end{aligned}$$

under the assumption that $\lim_{n \rightarrow \infty} n^{\frac{4s}{4s+\theta+1}} \Delta_n = \infty$. \square

Proof of Theorem 4. Without loss of generality, assume $M = 1$ and $\Delta_n = cn^{-\frac{4s}{4s+\theta+1}}$ for some $c > 0$. The main idea behind our proof is to carefully construct a finite subset of $\mathcal{P}(\Delta_n, \theta) \setminus \{P_0\}$, and show that one can not reliably distinguish P_0 from an unknown instance from this subset based on a sample of n observations. We shall consider the cases of $\theta = 0$ and $\theta > 0$ separately.

The case of $\theta = 0$. We first treat the case when $\theta = 0$. Let $K_n = \lfloor C_8 \Delta_n^{-\frac{1}{2s}} \rfloor$ for a sufficiently small constant $C_8 > 0$ and $a_n = \sqrt{\Delta_n / K_n}$. For any $\xi_n := (\xi_{n1}, \xi_{n2}, \dots, \xi_{nK_n})^\top \in$

$\{\pm 1\}^{K_n}$, write

$$u_{\xi_n} = a_n \sum_{k=1}^{K_n} \xi_{nk} \varphi_k.$$

It is clear that

$$\|u_{n,\xi_n}\|_{L_2(P_0)}^2 = K_n a_n^2 = \Delta_n$$

and

$$\|u_{n,\xi_n}\|_{\infty} \leq a_n K_n \left(\sup_k \|\varphi_k\|_{\infty} \right) \asymp \Delta_n^{\frac{2s-1}{4s}} \rightarrow 0.$$

By taking C_8 small enough, we can also ensure

$$\|u_{\xi_n}\|_K^2 = a_n^2 \sum_{k=1}^{K_n} \lambda_k^{-1} \leq 1,$$

Therefore, there exists a probability measure $P_{\xi_n} \in \mathcal{P}(\Delta_n, 0)$ such that $dP/dP_0 = 1 + u_{n,\xi_n}$. Following a standard argument for minimax lower bound, it suffices to show that

$$\overline{\lim}_{n \rightarrow \infty} \mathbb{E}_{P_0} \left(\frac{1}{2^{K_n}} \sum_{\xi_n \in \{\pm 1\}^{K_n}} \left\{ \prod_{i=1}^n [1 + u_{\xi_n}(X_i)] \right\} \right)^2 < \infty. \quad (19)$$

See, *e.g.*, [Ingster and Suslina \(2003\)](#); [Tsybakov \(2008\)](#).

Note that

$$\begin{aligned} & \mathbb{E}_{P_0} \left(\frac{1}{2^{K_n}} \sum_{\xi_n \in \{\pm 1\}^{K_n}} \left\{ \prod_{i=1}^n [1 + u_{\xi_n}(X_i)] \right\} \right)^2 \\ &= \mathbb{E}_{P_0} \left(\frac{1}{2^{2K_n}} \sum_{\xi_n, \xi'_n \in \{\pm 1\}^{K_n}} \left\{ \prod_{i=1}^n [1 + u_{\xi_n}(X_i)] \right\} \left\{ \prod_{i=1}^n [1 + u_{\xi'_n}(X_i)] \right\} \right) \\ &= \frac{1}{2^{2K_n}} \sum_{\xi_n, \xi'_n \in \{\pm 1\}^{K_n}} \prod_{i=1}^n \mathbb{E}_{P_0} \left\{ [1 + u_{\xi_n}(X_i)] [1 + u_{\xi'_n}(X_i)] \right\} \\ &= \frac{1}{2^{2K_n}} \sum_{\xi_n, \xi'_n \in \{\pm 1\}^{K_n}} \left(1 + a_n^2 \sum_{k=1}^{K_n} \xi_{nk} \xi'_{nk} \right)^n \\ &\leq \frac{1}{2^{2K_n}} \sum_{\xi_n, \xi'_n \in \{\pm 1\}^{K_n}} \exp \left(n a_n^2 \sum_{k=1}^{K_n} \xi_{n,k} \xi'_{n,k} \right) \\ &= \left\{ \frac{\exp(n a_n^2) + \exp(-n a_n^2)}{2} \right\}^{K_n} \\ &\leq \exp \left\{ K_n \left(\frac{\exp(n a_n^2) + \exp(-n a_n^2)}{2} - 1 \right) \right\}. \end{aligned}$$

An application of Taylor expansion shows that there exist $t_0 > 0$ and $C_9 > 0$ such that

$$t^2 - C_9|t|^3 \leq \exp(t) + \exp(-t) \leq t^2 + C_9|t|^3$$

for any $|t| \leq t_0$. With the particular choice of K_n , a_n , and the conditions on Δ_n , this immediately implies (19).

The case of $\theta > 0$. The main idea is similar to before. To find a set of probability measures in $\mathcal{P}(\Delta_n, \theta)$, we appeal to the following lemma.

Lemma 3. *Let $u = \sum_k a_k \varphi_k$. If*

$$\left(\sum_{k=1}^K \frac{a_k^2}{\lambda_k} \right)^{2/\theta} \left(\sum_{k \geq K} a_k^2 \right) \leq M^2,$$

then $u \in \mathcal{F}(\theta, M)$.

Similar to before, we shall now take $K_n = \lfloor C_{10} \Delta_n^{-\frac{\theta+1}{2s}} \rfloor$ and $a_n = \sqrt{\Delta_n / K_n}$. By Lemma 3, we can find $P_{\xi_n} \in \mathcal{P}(\Delta_n, \theta)$ such that $dP/dP_0 = 1 + u_{n,\xi_n}$, for appropriately chosen C_{10} . Following the same argument as in the previous case, we can again verify (19). \square

Proof of Theorem 5. Without loss of generality, assume that $\Delta_n(\theta) = c_1(n^{-1}\sqrt{\log \log n})^{\frac{4s}{4s+\theta+1}}$ for some constant $c_1 > 0$ to be determined later.

Type I Error. We first prove the first statement which shows that the Type I error converges to 0. Following the same notations as defined in Theorem 2, let

$$N_{n,2} = \mathbb{E} \left\{ \sum_{j=2}^n \mathbb{E} \left(\tilde{\zeta}_{nj}^2 | \mathcal{F}_{j-1} \right) - 1 \right\}^2, \quad L_{n,2} = \sum_{j=2}^n \mathbb{E} \tilde{\zeta}_{nj}^4$$

where $\tilde{\zeta}_{nj} = \sqrt{2} \zeta_{nj} / (n \sqrt{v_n})$. As shown by Haeusler (1988),

$$\sup_t |P(T_{n,\varrho_n} > t) - \bar{\Phi}(t)| \leq C_{11}(L_{n,2} + N_{n,2})^{1/5},$$

where $\bar{\Phi}(t)$ is the survival function of the standard normal, *i.e.*, $\bar{\Phi}(t) = P(Z > t)$ where $Z \sim N(0, 1)$. Again by the argument from Hall (1984),

$$\mathbb{E} \left\{ \sum_{j=2}^n \mathbb{E}(\zeta_{nj}^2 | \mathcal{F}_{j-1}) - \frac{1}{2} n(n-1)v_n \right\}^2 \leq C_{12} [n^4 \mathbb{E} G_n^2(X, X') + n^3 \mathbb{E} \tilde{K}_n^2(X, X') \tilde{K}_n^2(X, X'')],$$

where $G_n(\cdot, \cdot)$ is defined in the proof of Theorem 2, and

$$\sum_{j=2}^n \mathbb{E} \zeta_{nj}^4 \leq C_{13} [n^2 \mathbb{E} \tilde{K}_n^4(X, X') + n^3 \mathbb{E} \tilde{K}_n^2(X, X') \tilde{K}_n^2(X, X'')],$$

which ensure

$$\begin{aligned} N_{n,2} &= \frac{4 \mathbb{E} \left\{ \sum_{j=2}^n \mathbb{E}(\zeta_{nj}^2 | \mathcal{F}_{j-1}) - \frac{1}{2} n(n-1) v_n - \frac{1}{2} n v_n \right\}^2}{n^4 v_n^2} \\ &\leq 8 \max \left\{ C_{12}, \frac{1}{4} \right\} \left\{ \frac{\mathbb{E} G_n^2(X, X')}{v_n^2} + \frac{\mathbb{E} \tilde{K}_n^2(X, X') \tilde{K}_n^2(X, X'')}{n v_n^2} + \frac{1}{n^2} \right\}, \end{aligned}$$

and

$$L_{n,2} = \frac{4 \sum_{j=2}^n \mathbb{E} \tilde{\zeta}_{nj}^4}{n^4 v_n^2} \leq 4 C_{13} \left\{ \frac{\mathbb{E} \tilde{K}_n^4(X, X')}{n^2 v_n^2} + \frac{\mathbb{E} \tilde{K}_n^2(X, X') \tilde{K}_n^2(X, X'')}{n v_n^2} \right\}.$$

As shown in the proof of Theorem 2,

$$\frac{\mathbb{E} G_n^2(X, X')}{v_n^2} \leq C_3 \varrho_n^{1/s}, \quad \frac{\mathbb{E} \tilde{K}_n^4(X, X')}{n^2 v_n^2} \leq C_4 n^{-2} \varrho_n^{-1/s}, \quad \text{and} \quad \frac{\mathbb{E} \tilde{K}_n^2(X, X') \tilde{K}_n^2(X, X'')}{n v_n^2} \leq C_3 n^{-1}.$$

Therefore,

$$\sup_t |P(T_{n, \varrho_n} > t) - \bar{\Phi}(t)| \leq C_{14} (\varrho_n^{\frac{1}{5s}} + n^{-\frac{1}{5}} + n^{-\frac{2}{5}} \varrho_n^{-\frac{1}{5s}}),$$

which implies that

$$P \left(\sup_{\varrho_n \in \Lambda_n} T_{n, \varrho_n} > t \right) \leq m_* \bar{\Phi}(t) + C_{15} (2^{\frac{m_*}{5s}} \varrho_*^{\frac{1}{5s}} + m_* n^{-\frac{1}{5}} + n^{-\frac{2}{5}} \varrho_*^{-\frac{1}{5s}}), \quad \forall t.$$

It is not hard to see, by the definitions of m_* ,

$$2^{m_*} \varrho_* \leq 2 \left(\frac{\sqrt{\log \log n}}{n} \right)^{\frac{2s}{4s+1}}$$

and

$$\begin{aligned} m_* &= (\log 2)^{-1} \{ 2s \log n - \frac{2s}{4s+1} \log n + o(\log n) \} \\ &= (\log 2)^{-1} \frac{8s^2}{4s+1} \log n + o(\log n) \asymp \log n. \end{aligned}$$

Together with the fact that $\bar{\Phi}(t) \leq \frac{1}{2} e^{-t^2/2}$ for $t \geq 0$, we get

$$\begin{aligned} &P \left(\sup_{0 \leq k \leq m_*} T_{n, 2^k \varrho_*} > \sqrt{3 \log \log n} \right) \\ &\leq C_{16} \left[e^{-\frac{3}{2} \log \log n} \log n + \left(\frac{\sqrt{\log n}}{n} \right)^{\frac{2}{5(4s+1)}} + n^{-\frac{1}{5}} \log \log n + n^{-\frac{2}{5}} \left(\frac{\sqrt{\log \log n}}{n} \right)^{-\frac{2}{5}} \right] \rightarrow 0, \end{aligned}$$

as $n \rightarrow \infty$.

Type II Error. Next consider Type II error. To this end, write $\varrho_n(\theta) = \left(\frac{\sqrt{\log \log n}}{n}\right)^{\frac{2s(\theta+1)}{4s+\theta+1}}$. Let

$$\tilde{\varrho}_n(\theta) = \sup_{0 \leq k \leq m_*} \{2^k \varrho_* : \varrho_n \leq \varrho_n(\theta)\}.$$

It is clear that $\tilde{T}_n \geq T_{n, \tilde{\varrho}_n(\theta)}$ for any $\theta \geq 0$. It therefore suffices to show that for any $\theta \geq 0$,

$$\lim_{n \rightarrow \infty} \inf_{\theta \geq 0} \inf_{P \in \mathcal{P}(\Delta_n, \theta)} P \left\{ T_{n, \tilde{\varrho}_n(\theta)} \geq \sqrt{3 \log \log n} \right\} = 1.$$

By Markov inequality, this can be accomplished by verifying

$$\inf_{\theta \in [0, \infty)} \inf_{P \in \mathcal{P}(\Delta_n(\theta), \theta)} \mathbb{E}_P T_{n, \tilde{\varrho}_n(\theta)} \geq \tilde{M} \sqrt{\log \log n} \quad (20)$$

for some $\tilde{M} > \sqrt{3}$; and

$$\lim_{n \rightarrow \infty} \sup_{\theta \geq 0} \sup_{P \in \mathcal{P}(\Delta_n(\theta), \theta)} \frac{\text{Var}(T_{n, \tilde{\varrho}_n(\theta)})}{(\mathbb{E}_P T_{n, \tilde{\varrho}_n(\theta)})^2} = 0. \quad (21)$$

We now show that both (20) and (21) hold with

$$\Delta_n(\theta) = c_1 \left(\frac{\sqrt{\log \log n}}{n} \right)^{\frac{4s}{4s+\theta+1}}$$

for a sufficiently large $c_1 = c_1(M, \tilde{M})$.

Note that $\forall \theta \in [0, \infty)$,

$$\frac{1}{2} \varrho_n(\theta) \leq \tilde{\varrho}_n(\theta) \leq \varrho_n(\theta), \quad (22)$$

which immediately suggests

$$\eta_{\tilde{\varrho}_n(\theta)}^2(P, P_0) \geq \eta_{\varrho_n(\theta)}^2(P, P_0). \quad (23)$$

Following the arguments in the proof of Theorem 3,

$$\mathbb{E}_P T_{n, \tilde{\varrho}_n(\theta)} \geq C_{17} n [\tilde{\varrho}_n(\theta)]^{1/(2s)} \eta_{\tilde{\varrho}_n(\theta)}^2(P, P_0) \geq 2^{-1/(2s)} C_{17} n [\varrho_n(\theta)]^{1/2s} \eta_{\varrho_n(\theta)}^2(P, P_0),$$

and $\forall P \in \mathcal{P}(\Delta_n(\theta), \theta)$,

$$\eta_{\varrho_n(\theta)}^2(P, P_0) \geq \frac{1}{4} \|u\|_{L_2(P_0)}^2 \quad (24)$$

provided that $\Delta_n(\theta) \geq C'(M) \left(\frac{\sqrt{\log \log n}}{n} \right)^{\frac{4s}{4s+\theta+1}}$.

Therefore,

$$\inf_{P \in \mathcal{P}(\Delta_n(\theta), \theta)} \mathbb{E}_P T_{n, \tilde{\varrho}_n(\theta)} \geq C_{18} n [\varrho_n(\theta)]^{1/(2s)} \Delta_n(\theta) \geq C_{18} c_1 \sqrt{\log \log n} \geq \tilde{M} \sqrt{\log \log n}$$

if $c_1 \geq C_{18}^{-1} \tilde{M}$. Hence to ensure (20) holds, it suffices to take

$$c_1 = \max\{C'(M), C_{18}^{-1} \tilde{M}\}.$$

With (22), (23) and (24), the results in Theorem 3 imply that for sufficiently large n

$$\begin{aligned} \sup_{P \in \mathcal{P}(\Delta_n^*(\theta), \theta, M)} \frac{\text{Var}(T_{n, \tilde{\varrho}_n(\theta)})}{(\mathbb{E}_P T_{n, \tilde{\varrho}_n(\theta)})^2} &\leq C_{19} \left\{ \left([\varrho_n(\theta)]^{\frac{1}{2s}} n \Delta_n^*(\theta) \right)^{-2} + \left([\varrho_n(\theta)]^{\frac{1}{s}} n^2 \Delta_n^*(\theta) \right)^{-1} \right. \\ &\quad \left. + (n \Delta_n^*(\theta))^{-1} + \left([\varrho_n(\theta)]^{\frac{1}{2s}} n \sqrt{\Delta_n^*(\theta)} \right)^{-1} \right\} \\ &\leq 2C_{19} \left([\varrho_n(\theta)]^{\frac{1}{2s}} n \Delta_n^*(\theta) \right)^{-1} = 2C_{19} (c_1 \log \log n)^{-\frac{1}{2}} \rightarrow \infty, \end{aligned}$$

which shows (21). \square

Proof of Theorem 6. The main idea of the proof is similar to that for Theorem 4. To this end, assume, without loss of generality, that

$$\Delta_n(\theta) = c_2 \left(\frac{n}{\sqrt{\log \log n}} \right)^{-\frac{4s}{4s+\theta+1}}, \quad \forall \theta \in [\theta_1, \theta_2],$$

where $c_2 > 0$ is a sufficiently small constant to be determined later.

Let $r_n = \lfloor C_{20} \log n \rfloor$ and $K_{n,1} = \lfloor C_{21} \Delta_n^{-\frac{\theta_1+1}{2s}}(\theta_1) \rfloor$ for sufficiently small $C_{20}, C_{21} > 0$. Set $\theta_{n,1} = \theta_1$. For $2 \leq r \leq r_n$, let

$$K_{n,r} = 2^{r-2} K_{n,1}$$

and $\theta_{n,r}$ is selected such that the following equation holds.

$$K_{n,r} = \left\lfloor C_{21} [\Delta_n(\theta_{n,r})]^{-\frac{\theta_{n,r}+1}{2s}} \right\rfloor.$$

Note that by choosing C_{20} sufficiently small,

$$\begin{aligned} K_{n,r_n} = 2^{r_n-2} K_{n,1} &\leq \left\lfloor c_2^{\frac{2(\theta_1+1)}{4s+\theta+1}} \left(\frac{n}{\sqrt{\log \log n}} \right)^{\frac{2(\theta_1+1)}{4s+\theta+1}} \cdot 2^{r_n-2} \right\rfloor \\ &= \left\lfloor c_2^{\frac{2(\theta_1+1)}{4s+\theta+1}} \exp \left(\log \left(\frac{n}{\sqrt{\log \log n}} \right) \cdot \frac{2(\theta_1+1)}{4s+\theta_1+1} + (r_n-2) \log 2 \right) \right\rfloor \\ &\leq \left\lfloor C_{21} \exp \left(\log \left(\frac{n}{\sqrt{\log \log n}} \right) \cdot \frac{2(\theta_2+1)}{4s+\theta_2+1} \right) \right\rfloor = \lfloor C_{21} [\Delta_n(\theta_2)]^{-\frac{\theta_2+1}{2s}} \rfloor \end{aligned}$$

for sufficiently large n . Thus, we can guarantee that $\forall 1 \leq r \leq r_n, \theta_{n,r_n} \in [\theta_1, \theta_2]$.

We now construct a finite subset of $\cup_{\theta \in [\theta_1, \theta_2]} \mathcal{P}(\Delta_n(\theta), \theta)$ as follows. For each $\xi_{n,r} = (\xi_{n,r,1}, \dots, \xi_{n,r,K_{n,r}}) \in \{\pm 1\}^{K_{n,r}}$, let

$$f_{n,r,\xi_{n,r}} = 1 + \sum_{k=K_{n,r-1}^*+1}^{K_{n,r}^*} a_{n,r} \xi_{n,r,k} \varphi_k,$$

where $K_{n,r}^* = K_{n,1} + \dots + K_{n,r}$, and $a_{n,r} = \sqrt{\Delta_n(\theta_{n,r})/K_{n,r}}$. Following the same argument as that in the proof of Theorem 4, we can verify that with a sufficiently small C_{21} , each $P_{n,r,\xi_{n,r}} \in \mathcal{P}(\Delta_n(\theta_{n,r}), \theta_{n,r})$, where $f_{n,r,\xi_{n,r}}$ is the Radon-Nikodym derivative $dP_{n,r,\xi_{n,r}}/dP_0$. With slight abuse of notation, write

$$f_n(X_1, X_2, \dots, X_n) = \frac{1}{r_n} \sum_{r=1}^{r_n} f_{n,r}(X_1, X_2, \dots, X_n),$$

where

$$f_{n,r}(X_1, X_2, \dots, X_n) = \frac{1}{2^{K_{n,r}}} \sum_{\xi_{n,r} \in \{\pm 1\}^{K_{n,r}}} \prod_{i=1}^n f_{n,r,\xi_{n,r}}(X_i).$$

It now suffices to show that

$$\|f_n\|_{L_2(P_0)} := \mathbb{E}_{P_0} f_n^2(X_1, X_2, \dots, X_n) \rightarrow 1, \quad \text{as } n \rightarrow \infty.$$

Note that

$$\begin{aligned} \|f_n\|_{L_2(P_0)}^2 &= \frac{1}{r_n^2} \sum_{1 \leq r, r' \leq r_n} \langle f_{n,r}, f_{n,r'} \rangle_{L_2(P_0)} \\ &= \frac{1}{r_n^2} \sum_{1 \leq r \leq r_n} \|f_{n,r}\|_{L_2(P_0)}^2 + \frac{1}{r_n^2} \sum_{\substack{1 \leq r, r' \leq r_n \\ r \neq r'}} \langle f_{n,r}, f_{n,r'} \rangle_{L_2(P_0)}. \end{aligned}$$

It is easy to verify that, for any $r \neq r'$,

$$\langle f_{n,r}, f_{n,r'} \rangle_{L_2(P_0)} = 1.$$

It therefore suffices to show that

$$\sum_{1 \leq r \leq r_n} \|f_{n,r}\|_{L_2(P_0)}^2 = o(r_n).$$

Following the same derivation as that in the proof of Theorem 4, we can show that

$$\|f_{n,r}\|_{L_2(P_0)}^2 \leq \left(\frac{\exp(na_{n,r}^2) + \exp(-na_{n,r}^2)}{2} \right)^{K_{n,r}} \leq \exp(K_{n,r}n^2a_{n,r}^4)$$

for sufficiently large n . By setting c_2 in the expression of $\Delta_n(\theta)$ sufficiently small, we have

$$K_{n,r}n^2a_{n,r}^4 \leq \frac{1}{2} \log r_n,$$

which ensures that

$$\|f_{n,r}\|_{L_2(P_0)}^2 \leq r_n^{1/2} = o(r_n).$$

□

References

- L. Addario-Berry, N. Broutin, L. Devroye, and G. Lugosi. On combinatorial testing problems. *The Annals of Statistics*, 38(5):3063–3092, 2010.
- N. Ailon, M. Charikar, and A. Newman. Aggregating inconsistent information: ranking and clustering. *Journal of ACM*, 55(5):23:1–23:27, 2008.
- N. Aronszajn. Theory of reproducing kernels. *Transactions of the American Mathematical Society*, 68(3):337–404, 1950.
- A. Banerjee, I. S. Dhillon, J. Ghosh, and S. Sra. Clustering on the unit hypersphere using von mises-fisher distributions. In *Journal of Machine Learning Research*, 1345–1382, 2005.
- K. Chwialkowski, H. Strathmann, and A. Gretton. A kernel test of goodness of fit. In *International Conference on Machine Learning*, 2606–2615, 2016.
- F. Cucker and D. Zhou. *Learning Theory: an Approximation Theory Viewpoint*. Cambridge University Press, New York, NY, 2007.
- I. S. Dhillon, E. M. Marcotte, and U. Roshan. Diametrical clustering for identifying anti-correlated gene clusters. *Bioinformatics*, 19(13):1612–1619, 2003.

- T. A. Driscoll, N. Hale, and L. N. Trefethen. *Chebfun Guide*. Pafnuty Publications, Oxford, 2014. URL <http://www.chebfun.org/docs/guide/>.
- G. K. Dziugaite, D. M. Roy, and Z. Ghahramani. Training generative neural networks via maximum mean discrepancy optimization. In *Conference on Uncertainty in Artificial Intelligence*, 258–267, 2015.
- C. Fraley and A. E. Raftery. Model-based clustering, discriminant analysis, and density estimation. *Journal of the American Statistical Association*, 97(458):611–631, 2002.
- J. Gorham and L. Mackey. Measuring sample quality with kernels. In *International Conference on Machine Learning*, 2017.
- G. G. Gregory. Large sample theory for U-statistics and tests of fit. *The Annals of Statistics*, 5(1):110–123, 1977.
- A. Gretton, K. M. Borgwardt, M. Rasch, B. Schölkopf, and A. J. Smola. A kernel method for the two-sample-problem. In *Advances in Neural Information Processing Systems*, 513–520, 2007.
- A. Gretton, K. M. Borgwardt, M. J. Rasch, B. Schölkopf, and A. Smola. A kernel two-sample test. *Journal of Machine Learning Research*, 13(Mar):723–773, 2012.
- E. Haeusler. On the rate of convergence in the central limit theorem for martingales with discrete and continuous time. *The Annals of Probability*, 16(1):275–299, 1988.
- P. Hall. Central limit theorem for integrated square error of multivariate nonparametric density estimators. *Journal of Multivariate Analysis*, 14(1):1–16, 1984.
- Z. Harchaoui, F. Bach, and E. Moulines. Testing for homogeneity with kernel fisher discriminant analysis. In *Advances in Neural Information Processing Systems*, 609–616, 2007.
- T. R. Hughes, M. J. Marton, A. R. Jones, C. J. Roberts, R. Stoughton, C. D. Armour, H. A. Bennett, E. Coffey, H. Dai, and Y. D. He. Functional discovery via a compendium of expression profiles. *Cell*, 102(1):109–126, 2000.
- Yu. I. Ingster. Asymptotically minimax hypothesis testing for nonparametric alternatives. i, ii, iii. *Mathematical Methods of Statistics*, 2(2):85–114, 1993.

- Yu. I. Ingster. Minimax testing of hypotheses on the distribution density for ellipsoids in l_p . *Theory of Probability & Its Applications*, 39(3):417–436, 1995.
- Yu. I. Ingster. Adaptive chi-square tests. *Journal of Mathematical Sciences*, 99(2):1110–1119, 2000.
- Yu. I. Ingster and I. A. Suslina. *Nonparametric Goodness-of-Fit Testing under Gaussian Models*. Springer, New York, NY, 2003.
- V. R. Iyer, M. B. Eisen, D. T. Ross, G. Schuler, T. Moore, J. Lee, J. M. Trent, L. M. Staudt, J. Hudson, and M. S. Boguski. The transcriptional program in the response of human fibroblasts to serum. *Science*, 283(5398):83–87, 1999.
- W. Jitkrittum, W. Xu, Z. Szabo, K. Fukumizu, and A. Gretton. A linear-time kernel goodness-of-fit test. In *Advances in Neural Information Processing Systems (to appear)*, 2017.
- P. E. Jupp. Sobolev tests of goodness of fit of distributions on compact riemannian manifolds. *The Annals of Statistics*, 33(6):2957–2966, 2005.
- E. L. Lehmann and J. P. Romano. *Testing Statistical Hypotheses*. Springer Science & Business Media, New York, NY, 2008.
- Y. Li, K. Swersky, and R. Zemel. Generative moment matching networks. In *International Conference on Machine Learning*, 1718–1727, 2015.
- Q. Liu, J. Lee, and M. Jordan. A kernelized stein discrepancy for goodness-of-fit tests. In *International Conference on Machine Learning*, 276–284, 2016.
- R. Lyons. Distance covariance in metric spaces. *The Annals of Probability*, 41(5):3284–3305, 2013.
- K. V. Mardia and P. E. Jupp. *Directional Statistics*. John Wiley & Sons, New York, NY, 2009.
- J. S. Marron and M. P. Wand. Exact mean integrated squared error. *The Annals of Statistics*, 20(2):712–736, 1992.

- H. Q. Minh, P. Niyogi, and Y. Yao. Mercers theorem, feature maps, and smoothing. In *Conference on Learning Theory*, 154–168, 2006.
- K. Muandet, K. Fukumizu, B. Sriperumbudur, and B. Schölkopf. Kernel mean embedding of distributions: a review and beyond. *Foundations and Trends® in Machine Learning*, 10(1-2):1–141, 2017.
- D. Sejdinovic, B. Sriperumbudur, A. Gretton, and K. Fukumizu. Equivalence of distance-based and RKHS-based statistics in hypothesis testing. *The Annals of Statistics*, 41(5):2263–2291, 2013.
- R. J. Serfling. *Approximation Theorems of Mathematical Statistics*. John Wiley & Sons, New York, NY, 2009.
- A. J. Smola, A. Gretton, L. Song, and B. Schölkopf. A Hilbert space embedding for distributions”. In *Proc. 18th International Conference on Algorithmic Learning Theory*, 13–31. Springer-Verlag, Berlin, Germany, 2007.
- P. T. Spellman, G. Sherlock, M. Q. Zhang, V. R. Iyer, K. Anders, M. B. Eisen, P. O. Brown, D. Botstein, and B. Futcher. Comprehensive identification of cell cycle-regulated genes of the yeast *saccharomyces cerevisiae* by microarray hybridization. *Molecular Biology of the Cell*, 9(12):3273–3297, 1998.
- V. G. Spokoiny. Adaptive hypothesis testing using wavelets. *The Annals of Statistics*, 24(6):2477–2498, 1996.
- S. Sra and D. Karp. The multivariate Watson distribution: maximum-likelihood estimation and other aspects. *Journal of Multivariate Analysis*, 114(C):256–269, 2013.
- G. J. Székely, M. L. Rizzo, and N. K. Bakirov. Measuring and testing dependence by correlation of distances. *The Annals of Statistics*, 35(6):2769–2794, 2007.
- L. N. Trefethen and Z. Battles. An extension of MATLAB to continuous functions and operators. *SIAM Journal on Scientific Computing*, 25:1743–1770, 2004. doi: 10.1137/S1064827503430126. URL <http://link.aip.org/link/?SCE/25/1743/1>.

- A. B. Tsybakov. *Introduction to Nonparametric Estimation*. Springer Science & Business Media, New York, NY, 2008.
- W. Zaremba, A. Gretton, and M. Blaschko. B-test: a non-parametric, low variance kernel two-sample test. In *Advances in Neural Information Processing Systems*, 755–763, 2013.

Appendices

Proof of Lemma 2. We have

$$G^2(x, x') = \sum_{k,l} \mu_k \mu_l \varphi_k(x) \varphi_l(x) \varphi_k(x') \varphi_l(x').$$

Thus

$$\begin{aligned} & \int g(x) g(x') G^2(x, x') dP(x) dP(x') \\ &= \sum_{k,l} \mu_k \mu_l \left(\int g(x) \varphi_k(x) \varphi_l(x) dP(x) \right)^2 \\ &\leq \mu_1 \sum_k \mu_k \sum_l \left(\int g(x) \varphi_k(x) \varphi_l(x) dP(x) \right)^2 \\ &\leq \mu_1 \left(\sum_k \mu_k \int g^2(x) \varphi_k^2(x) dP(x) \right) \\ &\leq \mu_1 \left(\sum_k \mu_k \right) \left(\sup_k \|\varphi_k\|_\infty \right)^2 \|g\|_{L_2(P)}^2. \end{aligned}$$

□

Proof of Lemma 3. For brevity, write

$$l_K = \sum_{k=1}^K \frac{a_k^2}{\lambda_k}.$$

By definition, it suffices to show that $\forall R > 0, \exists f_R \in \mathcal{H}(K)$ such that $\|f_R\|_K^2 \leq R^2$ and $\|u - f_R\|_{L_2(P_0)}^2 \leq M^2 R^{-2/\theta}$.

To this end, let K be such that $l_K^2 \leq R^2 \leq l_{K+1}^2$, and denote by

$$f_R = \sum_{k=1}^K a_k \varphi_k + a_{K+1}^*(R) \varphi_{K+1},$$

where

$$a_{K+1}^*(R) = \text{sgn}(a_{K+1}) \sqrt{\lambda_{K+1}(R^2 - l_K^2)}.$$

Clearly,

$$\|f_R\|_K^2 = \sum_{k=1}^K \frac{a_k^2}{\lambda_k} + \frac{(a_{K+1}^*(R))^2}{\lambda_{K+1}} = R^2,$$

and

$$\|u - f_R\|_{L_2(P_0)}^2 = \sum_{k>K+1} a_k^2 + \left(|a_{K+1}| - \sqrt{\lambda_{K+1}(R^2 - l_K^2)} \right)^2 \leq \sum_{k \geq K+1} a_k^2.$$

To ensure $u \in \mathcal{F}(\theta, M)$, it suffices to have

$$\sup_{l_K^2 \leq R^2 \leq l_{K+1}^2} \|u - f_R\|_{L_2(P_0)}^2 R^{2/\theta} \leq M^2, \quad \forall K \geq 0,$$

which concludes the proof. □