

CONIC BUNDLES AND ITERATED ROOT STACKS

JAKOB OESINGHAUS

ABSTRACT. We generalize a classical result by V. G. Sarkisov about conic bundles to the case of a not necessarily algebraically closed perfect field, using iterated root stacks, destackification, and resolution of singularities. More precisely, we prove that whenever resolution of singularities is available, over a general perfect base field, any conic bundle is birational to a standard conic bundle.

CONTENTS

1. Introduction	1
2. Preliminaries	2
2.1. Conic bundles	2
2.2. Gerbes	3
2.3. Root stacks and the Brauer group	4
3. Proof the of the main result	5
References	9

1. INTRODUCTION

In this paper, we study the geometry of conic bundles, that is, fibrations whose generic fiber is a smooth conic. They have been widely studied in the context of rationality problems, notably the classic result of Artin and Mumford, who computed their Brauer groups in [5] to produce unirational non-rational conic bundles over rational surfaces. In order to better understand these bundles, it is desirable to bring a conic bundle into a standard form where the locus of degeneration can be controlled.

Over an algebraically closed field k and assuming resolution of singularities, a classical result by Sarkisov ([17]) states every conic bundle $\pi : V \rightarrow S$ of irreducible varieties such that S is smooth and π is projective can be brought into a standard form. Concretely, this means that there exist smooth varieties \tilde{V} and \tilde{S} and a projective morphism $\tilde{\pi} : \tilde{V} \rightarrow \tilde{S}$ fitting

2010 *Mathematics Subject Classification.* 14J10 (Primary) 14A20, 14E05, 14F22 (Secondary).

Key words and phrases. Conic bundles, root stacks, destackification, resolution of singularities.

into a commutative square

$$\begin{array}{ccc} \tilde{V} & \dashrightarrow & V \\ \downarrow \tilde{\pi} & & \downarrow \pi \\ \tilde{S} & \longrightarrow & S \end{array}$$

such that the rational map $\tilde{V} \dashrightarrow V$ and the projective morphism $\tilde{S} \rightarrow S$ are birational, and such that

- the generic fiber of $\tilde{\pi}$ is a smooth conic,
- the discriminant divisor of $\tilde{\pi}$ is a simple normal crossing divisor,
- the general fiber of $\tilde{\pi}$ along every irreducible component of the discriminant divisor is a singular irreducible reduced conic, and
- the fibers of $\tilde{\pi}$ over the singular locus of the discriminant divisor are non-reduced conics, i.e. double lines.

We use root stacks, resolution of singularities, and the destackification procedure ([7]) to generalize Sarkisov's result to a general perfect base field in Theorem 4. An analogous result for Brauer-Severi surface bundles, i.e. fibrations whose generic fiber is a form of \mathbb{P}^2 , has been proven by Kresch and Tschinkel in [12], also using root stack techniques.

Acknowledgements. I would like to thank my advisor Andrew Kresch for his guidance. I am supported by Swiss National Science Foundation grant 156010.

2. PRELIMINARIES

Throughout this section, when X is a Noetherian Deligne-Mumford (DM) stack, we will let n be a positive integer, invertible in the local rings of an étale atlas of X . A *sheaf* on X is a sheaf of abelian groups on the étale site of X ; all cohomology will be étale cohomology.

2.1. Conic bundles.

Definition 1. Let S be a regular scheme such that 2 is invertible in its local rings. A *regular conic bundle over S* is a flat, projective morphism $\pi : V \rightarrow S$ from a regular scheme V such that the generic fiber over every irreducible component is smooth and such that all fibers are isomorphic to a conic in \mathbb{P}^2 . A regular conic bundle is called *standard* if π is relatively minimal, i.e. if the preimage of an irreducible divisor under π is an irreducible divisor; equivalently, if there exists a reduced divisor $D \subset S$ whose singular locus is regular, such that

- The morphism π is smooth over $S \setminus D$ and the generic fiber over every irreducible component is a smooth conic.
- The generic fiber of π over every irreducible component of D is a singular reduced irreducible conic, i.e. the union of two lines with conjugate slopes.

- The fiber of π over every point of D^{sing} is non-reduced, i.e. a double line.

We remark that we put no further requirements on D , although our construction actually produces standard conic bundles with simple normal crossing discriminant divisor.

2.2. Gerbes. Let X be a Noetherian Deligne-Mumford stack.

Definition 2. A gerbe over X banded by μ_n , or simply a μ_n -gerbe over X , is the data of a Deligne-Mumford stack H and a morphism $H \rightarrow X$ that is étale locally isomorphic to a product with $B\mu_n$, together with compatible identifications of the automorphism groups of local sections with μ_n .

We can classify μ_n -gerbes by their class in $H^2(X, \mu_n)$. We use the notion of the *residual gerbe* \mathcal{G}_x of X at a point $x \in |X|$, an étale gerbe over the residue field $\kappa(x)$ satisfying certain universal properties ([16, App. B]).

Lemma 1. *Let $x \in |X|$. Then $H^1(\mathcal{G}_x, \mathbb{Z}) = 0$.*

Proof. The Leray spectral sequence for $f : \mathcal{G}_x \rightarrow \text{Spec } \kappa(x)$ gives a monomorphism

$$H^1(\mathcal{G}_x, \mathbb{Z}) \rightarrow H^0(\text{Spec } \kappa(x), R^1 f_* \mathbb{Z}).$$

Let $K/\kappa(x)$ be a finite separable extension such that the gerbe

$$\mathcal{Y} := \text{Spec } K \times_{\text{Spec } \kappa(x)} \mathcal{G}_x$$

has a section. This implies that $\mathcal{Y} \cong BG$ for a finite étale group scheme G over K . But then

$$H^1(\mathcal{Y}, \mathbb{Z}) = H^1(BG, \mathbb{Z}) = \text{Hom}(G, \mathbb{Z}) = 0.$$

Hence, we have $R^1 f_* \mathbb{Z} = 0$. □

We will also frequently make use of the *Kummer sequence*

$$0 \rightarrow \mu_n \rightarrow \mathbb{G}_m \xrightarrow{(\cdot)^n} \mathbb{G}_m \rightarrow 0,$$

which is an exact sequence of sheaves on X .

Suppose now that X is regular and integral, with trivial stabilizer at the generic point $\iota_\eta : \text{Spec}(\eta) \rightarrow X$. By [10, (2)-(3)], the following is an exact sequence of sheaves on X :

$$(1) \quad 0 \rightarrow \mathbb{G}_m \rightarrow (\iota_\eta)_* \mathbb{G}_m \rightarrow \bigoplus_{x \in X^{(1)}} (\iota_x)_* \mathbb{Z} \rightarrow 0.$$

By Lemma 1, this implies that $H^2(X, \mathbb{G}_m) \rightarrow H^2(X, (\iota_\eta)_* \mathbb{G}_m)$ is injective. The Leray spectral sequence for ι_η and Hilbert's Theorem 90 imply that

$$H^2(X, (\iota_\eta)_* \mathbb{G}_m) \rightarrow H^2(\text{Spec}(\eta), \mathbb{G}_m)$$

is injective. Hence, we can infer that $H^2(X, \mathbb{G}_m)$ is a torsion group, which we call the *Brauer group* of X . This motivates the following definition.

Definition 3. The *Brauer group* $\mathrm{Br}(X)$ of a Noetherian DM stack X is defined to be the torsion subgroup of $H^2(X, \mathbb{G}_m)$.

This reduces to the classical definition of the Brauer group if X is the spectrum of a field.

2.3. Root stacks and the Brauer group. Given effective Cartier divisors

$$D_1, \dots, D_\ell$$

on a Noetherian scheme or Deligne-Mumford stack X , we can define the *iterated n -th root stack* of X along those divisors ([8]), denoted by

$$\sqrt[n]{(X, \{D_1, \dots, D_\ell\})} \rightarrow X.$$

This construction adds stacky structure along the divisors, and is an isomorphism outside of the union of divisors. It should be noted that when any intersection $D_i \cap D_j$ with $i \neq j$ is nonempty, this is not isomorphic to the root stack along the union of the divisors. For any $i \in \{1, \dots, \ell\}$, the iterated root stack

$$\sqrt[n]{(X, \{D_1, \dots, \widehat{D_i}, \dots, D_\ell\})}$$

is a relative coarse moduli space for $\sqrt[n]{(X, \{D_1, \dots, D_\ell\})}$ in the sense of [1, §3]. In particular, a locally free sheaf on $\sqrt[n]{(X, \{D_1, \dots, D_\ell\})}$ such that the associated linear μ_n -representation is trivial at a general point of any irreducible component of D_i descends to

$$\sqrt[n]{(X, \{D_1, \dots, \widehat{D_i}, \dots, D_\ell\})}.$$

To prove this, we can apply a relative version of [2, Thm. 10.3] for good moduli space morphisms, which are relative versions of good moduli spaces. Note that

- (1) the condition of trivial stabilizer actions at closed points is replaced by trivial action of the *relative inertia stack* at closed points;
- (2) due to the construction of root stacks, this action will be trivial if it is trivial at a general point of every irreducible component of D_i ;
- (3) since the stacks involved are tame, the relative coarse moduli space mentioned above is a good moduli space morphism.

We will use this several times throughout.

Lemma 2. *Let S be a regular integral Noetherian scheme of dimension 1, let n be invertible in the local rings of S , and let D_1, \dots, D_ℓ be distinct closed points of S . Then we have*

$$\mathrm{Br}(\sqrt[n]{(S, \{D_1, \dots, D_\ell\})}[n]) \cong \mathrm{Br}(S \setminus \{D_1, \dots, D_\ell\})[n].$$

Proof. Let X be any regular integral Noetherian DM stack with trivial generic stabilizer such that $\dim(X) = 1$, and such that n is invertible in the local rings of an étale atlas of X . Then the results from [11, 2.] and

the Leray spectral sequence for ι_η imply that $H^2(X, (\iota_\eta)_* \mathbb{G}_m)[n] \cong \text{Br}(\eta)[n]$. Moreover, if $x \in X^{(1)}$, the vanishing of $H^1(\mathcal{G}_x, \mathbb{Z})$ implies that

$$H^1(\mathcal{G}_x, \mathbb{Z}/n\mathbb{Z}) \cong H^2(\mathcal{G}_x, \mathbb{Z})[n].$$

The long exact sequence of cohomology of (1) then gives rise to an exact sequence

$$(2) \quad 0 \rightarrow \text{Br}(X)[n] \rightarrow \text{Br}(\eta)[n] \rightarrow \bigoplus_{x \in X^{(1)}} H^1(\mathcal{G}_x, \mathbb{Z}/n\mathbb{Z}).$$

Taking $X = \sqrt[n]{(S, \{D_1, \dots, D_\ell\})}$, with codimension 1 point x_i over D_i for all i , we compare the exact sequence (2) with the analogous exact sequence for S (loc. cit.) to obtain the vanishing of the right-hand map in (2) after projection to the factor x_i for any i [13, §3.2]. Comparison with the exact sequence for $S \setminus \{D_1, \dots, D_\ell\}$ gives the result. \square

Lemma 3. *Let k be a field and let X be integral, smooth and of finite type over k with trivial generic stabilizer. For any positive integer n with $\text{char}(k) \nmid n$ and open substack $U \subset X$ whose complement has codimension at least 2, we have $H^2(X, \mu_n) \cong H^2(U, \mu_n)$, and therefore $\text{Br}(X)[n] \cong \text{Br}(U)[n]$.*

Proof. By [14, Rem II.3.17] there is no loss of generality in assuming that k is perfect. By shrinking X if necessary and iterating the process for large open substacks of X , we can assume that the complement $Y = X \setminus U$ is smooth and of constant codimension $c \geq 2$ everywhere. In this situation, we know by [4, §XVI.3] that $H_Y^i(X, \mu_n) = 0$ for $i \neq 2c$. Combining this with the exact sequence for cohomology with support ([14, Prop III.1.25]) and the local-to-global spectral sequence ([14, §VI.5]) gives the result; cf. [11, Cor 6.2]. \square

3. PROOF THE OF THE MAIN RESULT

Here we state and prove the main result.

Theorem 4. *Let k be a perfect field of characteristic different from 2 and S a smooth projective algebraic variety over k . Assume that embedded resolution of singularities for reduced subschemes of S of pure codimension 1 and desingularization of reduced finite-type Deligne-Mumford stacks of pure dimension equal to $\dim(S)$ are known. Let*

$$\pi : V \rightarrow S$$

be a morphism of projective varieties over k whose generic fiber is a smooth conic. Then there exists a commutative diagram

$$\begin{array}{ccc} \tilde{V} & \xrightarrow{\rho_V} & V \\ \downarrow \tilde{\pi} & & \downarrow \pi \\ \tilde{S} & \xrightarrow{\rho_S} & S \end{array}$$

where ρ_S is a projective birational morphism, ρ_V is a birational map, and $\tilde{\pi}$ is a standard conic bundle with simple normal crossing discriminant divisor.

Remark 1. Embedded resolution of singularities is known in characteristic 0 for all dimensions by Hironaka's celebrated result. In positive characteristic, embedded resolution of singularities for both curves and surfaces is known (cf. [9]). Since the resolutions commute with smooth base change, the assumptions about desingularization of reduced finite-type Deligne-Mumford stacks are also true in all of these cases (apply resolution of singularities to a presentation).

Before embarking on the proof we make several observations of a general nature. Let k be a perfect field, let S be a smooth projective algebraic variety over k , let n be a positive integer such that $\text{char}(k) \nmid n$, and let $\alpha \in \text{Br}(k(S))[n]$. Then there exists a dense open $U \subset S$ and $\beta \in \text{Br}(U)$ such that α is the restriction of β . Taking U to be maximal, by Lemma 3, the complement $S \setminus U$ is a finite union of divisors.

Assume embedded resolution of singularities for reduced subschemes of S of pure codimension 1.

Then, upon replacing S by a smooth projective variety with birational morphism to S , we may suppose that the complement of U is a simple normal crossing divisor $D_1 \cup \dots \cup D_\ell$. Let

$$X := \sqrt[n]{(S, \{D_1, \dots, D_\ell\})},$$

the iterated root stack of S along the divisors D_i . We apply Lemma 2 to the scheme obtained by gluing the local rings at the generic points of the D_i along the generic point of S . Then [3, §VII.5, Thm 5.7] implies that α extends to an open neighborhood of the root stack over this scheme. Hence, by Lemma 3, there is a unique $\beta \in \text{Br}(X)[n]$ that restricts to α .

Now suppose that α is the class of a central simple algebra A of dimension d^2 as a $k(S)$ -vector space, with $n \mid d$. Let $\beta_0 \in H^2(X, \mu_n)$ be a lift of $\beta \in \text{Br}(X)[n]$ with corresponding gerbe G_0 banded by μ_n . Now A is the fiber at the generic point of some sheaf of Azumaya algebras \mathcal{A} on an open $W \subset X$ with complement of codimension at least 3 ([10, Thm 2.1]). The Brauer class of the pullback of \mathcal{A} to $W \times_X G_0$ is trivial, hence this pullback is the endomorphism algebra of a locally free coherent sheaf of rank d , which is the restriction of a coherent sheaf \mathcal{E}_0 on G_0 .

Assume resolution of singularities for reduced Noetherian DM stacks of finite type over k of pure dimension $\dim(S)$.

The identity of $\mathcal{E}_0|_{W \times_X G_0}$ induces a morphism to the Grassmannian of rank d quotients of \mathcal{E}_0 ([15]). Apply resolution of singularities to the closure of the image to obtain a smooth DM stack Y with a projective morphism to X that restricts to an isomorphism over W , a gerbe $H_0 := Y \times_X G_0$, and a locally free coherent sheaf \mathcal{F}_0 on H_0 whose restriction over W is isomorphic to the restriction of \mathcal{E}_0 . In this situation, let $\gamma_0 := \beta_0|_Y$.

In the proof of Theorem 4 we specialize the above to $n = d = 2$.

Proof. The proof begins with a series of reductions steps, starting with Y as above, equipped with a sheaf of Azumaya algebras, restricting to the quaternion algebra A over $k(S)$, associated with the generic fiber of π .

Step 1. We may suppose that $Y \cong \sqrt{(T, \{E_1, \dots, E_m\})}$ for some smooth projective variety T with birational morphism to S and irreducible divisors E_i such that $E_1 \cup \dots \cup E_m$ is a simple normal crossing divisor and such that at the generic point of each E_i , the projective representation $\mu_2 \rightarrow \mathrm{PGL}_2$ given by the sheaf of Azumaya algebras is nontrivial. Indeed, the destackification program ([7, Thm 1.2]) yields a morphism $\tilde{Y} \rightarrow Y$ that is a composition of blow-ups with smooth centers, such that

$$\tilde{Y} \cong \sqrt{(T, \{E_1, \dots, E_m\})}$$

for a smooth projective variety T and irreducible divisors E_i such that $E_1 \cup \dots \cup E_m$ is a simple normal crossing divisor. We pull back the sheaf of Azumaya algebras to \tilde{Y} . If there is an i such that the projective representation $\mu_2 \rightarrow \mathrm{PGL}_2$ over a general point of E_i is trivial, the sheaf of Azumaya algebras descends to

$$\sqrt{(T, \{E_1, \dots, \widehat{E_i}, \dots, E_m\})}.$$

Step 2. We may suppose, additionally, that generically along every component of $E_i \cap E_{i'}$ for $i \neq i'$ the projective representation of $\mu_2 \times \mu_2$ is faithful. Let $F \subset E_i \cap E_{i'}$ be an irreducible component with non-faithful representation. Let \tilde{T} be the blow-up of T along F . For every $j \in \{1, \dots, m\}$, we denote the proper transform of E_j by \tilde{E}_j , and we denote the exceptional divisor of the blow-up by E' . We let \tilde{Y} be the normalization of $\tilde{T} \times_T Y$. Then \tilde{Y} is isomorphic to the blow-up of Y at the corresponding component of the fiber product of the gerbes of the root stacks, which is itself isomorphic to the root stack

$$\sqrt{(\tilde{T}, \{\tilde{E}_1, \dots, \tilde{E}_m, E'\})}.$$

The projective representation over a general point of E' is trivial, so the sheaf of Azumaya algebras descends to

$$\sqrt{(\tilde{T}, \{\tilde{E}_1, \dots, \tilde{E}_m\})}.$$

Step 3. We may suppose, furthermore, that all triple intersections $E_i \cap E_{i'} \cap E_{i''}$ are empty, where i, i' , and i'' are distinct. Since there can never be more than 2 independent commuting subgroups of order 2 in PGL_2 ([6]), the projective representation $(\mu_2)^3 \rightarrow \mathrm{PGL}_2$ over a general point of $E_i \cap E_{i'} \cap E_{i''}$ has kernel equal to the diagonal μ_2 . We blow up T along $E_i \cap E_{i'} \cap E_{i''}$ and proceed as in Step 2.

Step 4. We may suppose, furthermore, that the Brauer class $[A] \in \mathrm{Br}(k(S))$ does not extend across the generic point of E_i for any i . Assume that it does, for some i . Let

$$\tilde{Y} := \sqrt{(T, \{E_1, \dots, \widehat{E_i}, \dots, E_m\})}.$$

Then by Lemma 3, the Brauer class is restriction of an element $\delta \in \text{Br}(\tilde{Y})$. Let $\varepsilon \in H^2(\tilde{Y}, \mu_2)$ denote an arbitrary lift of δ , with corresponding gerbe H_1 , such that if we let H_0 denote the base-change

$$Y \times_{\tilde{Y}} H_1,$$

then on H_0 the sheaf of Azumaya algebras is identified with endomorphism algebra of some locally free coherent sheaf \mathcal{F}_0 . Notice that H_0 is a root stack over H_1 . The relative stabilizer acts with eigenvalues 1 and -1 on fibers of \mathcal{F}_0 . The (-1) -eigensheaf is a quotient sheaf \mathcal{L}_{-1} supported on the gerbe of the root stack, such that the kernel \mathcal{F}_1 in

$$0 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_0 \rightarrow \mathcal{L}_{-1} \rightarrow 0$$

is again locally free and descends to H_1 , yielding a sheaf of Azumaya algebras on \tilde{Y} .

We now have $Y = \sqrt{(T, \{E_1, \dots, E_m\})}$, equipped with a sheaf of Azumaya algebras \mathcal{A} , such that the projective representations at a general point of every E_i are nontrivial, the projective representations at a general point of every intersection is faithful, there are no triple intersections, and such that the Brauer class does not extend over any of the generic points of the E_i . Let $P \rightarrow Y$ be the smooth \mathbb{P}^1 -fibration associated with \mathcal{A} .

Let T_0 denote the complement of the intersections of pairs of divisors,

$$T_0 := T \setminus \bigcup_{1 \leq i < i' \leq m} E_i \cap E_{i'}.$$

We apply [12, Proposition 3.1] to $T_0 \times_T P$ to obtain a regular conic bundle

$$\pi_0: V_0 \rightarrow T_0.$$

This factors canonically through $\mathbb{P}(\pi_{0*}(\omega_{V_0/T_0}^\vee))$. Let $i: T_0 \rightarrow T$ denote the inclusion. We claim that $i_*(\pi_{0*}(\omega_{V_0/T_0}^\vee))$ is a locally free coherent sheaf and, denoting this by \mathcal{E} , the closure $V = \overline{V_0}$ of V_0 in $\mathbb{P}(\mathcal{E})$ is a regular conic bundle over T . It suffices to verify these assertions after passing to an algebraic closure of k . Then there is a unique faithful projective representation $\mu_2 \times \mu_2 \rightarrow \text{PGL}_2$ (up to conjugacy), cf. [6]. So, by [12, Lemma 2.8] after base change to a suitable affine étale neighborhood $T' = \text{Spec}(B') \rightarrow T$ of a given point of an intersection $E_i \cap E_{i'}$, we have

$$Y' \cong \sqrt{(T', \{E_i, E_{i'}\})} = [\text{Spec}(B'[t, t']/(t^2 - x, t'^2 - x'))/\mu_2 \times \mu_2],$$

where x and x' are the respective defining equations for the preimage in T' of E_i and $E_{i'}$, with P' obtained by pulling back $[\mathbb{P}^1/\mu_2 \times \mu_2]$. Here, on $B'[t, t']/(t^2 - x, t'^2 - x')$, the action of the factors of $\mu_2 \times \mu_2$ is by respective scalar multiplication of t and t' , while the action on \mathbb{P}^1 corresponds to the faithful projective representation $\mu_2 \times \mu_2 \rightarrow \text{PGL}_2$. Over T' we compute $\overline{V'_0} \cong \text{Proj}(B'[u, v, w]/(xu^2 + x'v^2 - w^2))$, which is regular.

The fact that the Brauer class does not extend across the generic point of any E_i ensures that the conic bundle $V \rightarrow T$ is standard. \square

Remark 2. While the general destackification process outlined in [7] requires stacky blow-ups, it is never necessary to take root stacks when all stabilizers are powers of μ_2 .

REFERENCES

- [1] Dan Abramovich, Martin Olsson, and Angelo Vistoli. Twisted stable maps to tame Artin stacks. *J. Algebraic Geom.*, 20(3):399–477, 2011.
- [2] Jarod Alper. Good moduli spaces for Artin stacks. *Ann. Inst. Fourier (Grenoble)*, 63(6):2349–2402, 2013.
- [3] M. Artin, A. Grothendieck, and J. L. Verdier. *Théorie des topos et cohomologie étale des schémas (SGA 4), tome 2*. Lecture Notes in Mathematics, Vol. 270. Springer-Verlag, Berlin-New York, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat.
- [4] M. Artin, A. Grothendieck, and J. L. Verdier. *Théorie des topos et cohomologie étale des schémas (SGA 4), tome 3*. Lecture Notes in Mathematics, Vol. 305. Springer-Verlag, Berlin-New York, 1973.
- [5] M. Artin and D. Mumford. Some elementary examples of unirational varieties which are not rational. *Proc. London Math. Soc. (3)*, 25:75–95, 1972.
- [6] Arnaud Beauville. Finite subgroups of $\mathrm{PGL}_2(K)$. In *Vector bundles and complex geometry*, volume 522 of *Contemp. Math.*, pages 23–29. Amer. Math. Soc., Providence, RI, 2010.
- [7] Daniel Bergh. Functorial destackification of tame stacks with abelian stabilisers. *Compositio Mathematica*, 153(6):1257–1315, 2017.
- [8] Charles Cadman. Using stacks to impose tangency conditions on curves. *Amer. J. Math.*, 129(2):405–427, 2007.
- [9] Vincent Cossart, Uwe Jannsen, and Shuji Saito. Canonical embedded and non-embedded resolution of singularities for excellent two-dimensional schemes, 2009, [arXiv:0905.2191](#).
- [10] Alexander Grothendieck. Le groupe de Brauer. II. Théorie cohomologique. In *Dix exposés sur la cohomologie des schémas*, volume 3 of *Adv. Stud. Pure Math.*, pages 67–87. North-Holland, Amsterdam, 1968.
- [11] Alexander Grothendieck. Le groupe de Brauer. III. Exemples et compléments. In *Dix exposés sur la cohomologie des schémas*, volume 3 of *Adv. Stud. Pure Math.*, pages 88–188. North-Holland, Amsterdam, 1968.
- [12] A. Kresch and Y. Tschinkel. Models of Brauer-Severi surface bundles, August 2017, [arXiv:1708.06277](#).
- [13] Max Lieblich. Period and index in the Brauer group of an arithmetic surface. *J. Reine Angew. Math.*, 659:1–41, 2011. With an appendix by Daniel Krashen.
- [14] James S. Milne. *Étale cohomology*, volume 33 of *Princeton Mathematical Series*. Princeton University Press, Princeton, N.J., 1980.
- [15] Nitin Nitsure. Construction of Hilbert and Quot schemes. In *Fundamental algebraic geometry*, volume 123 of *Math. Surveys Monogr.*, pages 105–137. Amer. Math. Soc., Providence, RI, 2005.
- [16] David Rydh. Étale dévissage, descent and pushouts of stacks. *J. Algebra*, 331:194–223, 2011.
- [17] V. G. Sarkisov. On conic bundle structures. *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(2):371–408, 432, 1982.