

Approximation of the Masses and other properties of Unknown Hyperons in Standard Model

Imran Khan

*Department of Physics, University of Science and Technology,
Bannu 28100, Pakistan*

Email address: immarwat@yahoo.com

Abstract—In particle physics, study of the symmetry and its breaking play very important role in order to get useful information about the nature. The classification and arrangements of subatomic particles is also necessary to study particle physics. Particles which are building blocks of nature are quarks, gluons and leptons. Baryons and Mesons composed of quarks were arranged by Gell-Mann and Okubo in their well-known Eight-Fold way up to SU(3) symmetry. Standard model of particles is composed of these particles. Particles in SU(4) also make some beautiful pattern. These make some multiplets. but all the baryons with spin $J^P = 3/2^+$ in these multiplets have not been observed till date. In this paper the SU(4) multiplets have been organized and studied in an easy and new way. As a result we obtained some clues about the masses and other characteristics of the unknown hyperons. These approximations about the characteristics of the unidentified baryons have been recorded in this article. Mass formula for the baryon SU(4) multiplets have been obtained.

Keywords—Baryons, SU(3), SU(4), Hyperons, Standard Model, Mass Formula.

I. INTRODUCTION

Baryons are made of three quarks (qqq). The three flavors up u , down d , and strange s , imply an approximate flavor SU(3), which requires that baryons made of these quarks belong to the multiplets on the right side of the ‘equation’ $3 \otimes 3 \otimes 3 = 10_S \oplus 8_M \oplus 8_M \oplus 1_A$. Here the subscripts indicate symmetric, mixed symmetry, or anti-symmetric states under interchange of any two quarks [1]. These were classified and arranged by Gell-Mann and Okubo in De-Couplets ($J = 3/2, l = 0$) and Octets ($J = 1/2, l = 0$) with +1 parities [2–4]. The Gell-Mann / Okubo mass formula which relates the masses of members of the baryon octet is given by; [2–4]

$$2(m_N + m_{\Xi}) = 3m_{\Lambda} + m_{\Sigma} \quad (1)$$

While mass formula for De-couplets consists of equal spacing between the rows. The spaces are equal to an average value $-151 \text{ MeV}/c^2$.

$$M_{\Lambda} - M_{\Sigma^*} = M_{\Sigma^*} - M_{\Xi^*} = M_{\Xi^*} - M_{\Omega} \quad (2)$$

Gell-Mann used this formula and predicted the mass of the Ω^- baryon in 1962, equal to $M_{\Omega} = 1685 \text{ MeV}/c^2$ [2]. Whereas the actual mass of the Ω^- is $1672 \text{ MeV}/c^2$ which was observed in 1964 [4]. Their mass difference is only 0.72 %, or in other words it was 99% true guess.

Now let's move towards baryon types made from the combination of four quarks, i.e. up u , down d , strange s and charm c . These belong to $SU(4)$ multiplets. The $SU(4)$ multiplets numerology is given by

$$4 \otimes 4 \otimes 4 = 20_S \oplus 20_M \oplus 20_M \oplus 4_A \text{ [5].}$$

We are interested in twenty particles having spin $3/2$ and even parity $+1$ forming one of the $SU(4)$ multiplets. These particles are in their ground states, that is with $l = 0$. Their masses and other properties are listed in the Table 1.

Since the mass splitting due to strangeness is typically of order $151 \text{ MeV}/c^2$ in the de-couplet, which is an effect of about 13% of the masses it contains. The splitting due to charm in the $SU(4)$ particles multiplets is expected to be larger [6]. The ratio $\frac{m_2}{m_1} = \frac{m_4}{m_3}$ indicates the direction of the increasing mass in the Strangeness, or Charm-ness planes in $SU(3)$ and $SU(4)$ multiplets respectively. It gives the mass symmetry breaking ratio for the multiplet [6]. In the $SU(3)$ framework the Gell-Mann / Okubo relation for the $J^P = 1/2^+$ Octets, Eq. (1) and the equal spacing rule for the $J^P = 3/2^+$ de-couplets, Eq. (2) work so nicely that we cannot abandon linear mass formulae for baryons [6]. Using the above relation for baryons of $J^P = 3/2^+$ de-couplet of $SU(3)$ gives a mass symmetry breaking ratio of order 1.0.

Table 1. List of the spin $J^P = 3/2^+$ particles formed from three quarks combinations (qqq) of four quarks u, d, s and c [5].

S.#	Particle Name	Symbol	Quark contents	Rest Mass (MeV/c^2)
1	Delta $^{++}$	$\Delta^{++} (1232)$	uuu	1232 ± 2
2	Delta Plus	$\Delta^+ (1232)$	uud	1232 ± 2
3	Delta Zero	$\Delta^0 (1232)$	udd	1232 ± 2
4	Delta Minus	$\Delta^- (1232)$	ddd	1232 ± 2
5	Sigma Plus	$\Sigma^{*+} (1385)$	uus	1382.8 ± 0.4
6	Sigma Zero	$\Sigma^{*0} (1385)$	uds	1383.7 ± 1.0
7	Sigma Minus	$\Sigma^{*-} (1385)$	dds	1387.2 ± 0.5
8	charmed Sigma $^{++}$	$\Sigma_c^{*++} (2520)$	uuc	2517.9 ± 0.6
9	charmed Sigma $^{+}$	$\Sigma_c^{*+} (2520)$	udc	2517.5 ± 2.3
10	charmed	$\Sigma_c^{*0} (2520)$	ddc	2518.8 ± 0.6

II. METHOD

There are four particles in the Table 1, whose masses are unknown. Let's try to make some predictions about the unknown masses of the Ξ_{cc}^{*++} , Ξ_{cc}^{*+} , Ω_{cc}^{*+} , and Ω_{ccc}^{*++} hyperons. Let the particles listed in Table 1 may be grouped in pyramid shape of increasing charm number from $C = 0$ to $C = 3$ as shown in Fig. 1 [5]. Masses of the particles are expressed in round numbers. The bottom of the 'Pyramid' is the $SU(3)$ de-couplet of $J^P = 3/2^+$ particles at $C = 0$. Average masses written on the right side of the Fig. 1 are discussed later in the text.

We can expand this 20-plet given in Fig. 1 in the shape as shown in Fig. 2, with increasing Charge Number from bottom to top. Also there becomes a beautiful pattern of Isospin numbers, I of the particles. We can treat it as follows: Adding masses of the Δ^{++} and Δ^+ in the first row of Charm Number $C = 0$, gives $2464 \text{ MeV}/c^2$, which is 54 less than the mass of the Σ_c^{*+} (2518) particle in 2nd row of Charm Number $C = 1$. Hence approximately equal to its mass. Similarly Adding masses of Δ^+ and Δ^0 particles gives $2464 \text{ MeV}/c^2$, which is 55 less than the mass of the Σ_c^{*0} (2519) particle in 2nd row of Charm Number $C = 1$. Same can be done for other particles. It gives approximate mass of the particle in the adjacent row with higher charm-ness.




Fig. 2. $SU(4)$ 20-plet of Baryons ($J^P=3/2^+$) made of u , d , s and c quarks, arrows represent lines of the addition of particles masses. Indexed with a - f (See Table 2), and numbers beneath the arrows are mass difference between theoretical and experimentally observed masses of the particles.

II MASSES OF THE Ξ_{cc}^{*+} AND Ξ_{cc}^{*++}

Assuming the same behavior of the masses, we can sum up masses of two particles in 2nd row of Charm Number $C = 1$, and get mass of the particle in 3rd row with charm Number $C = 2$.

Sum of the masses calculated in such a manner are indexed from (a) to (i) in Fig. 2; and are listed in Table 2. Masses of the Particles in 3rd row of charm Number $C = 2$ are unknown. Using this method the masses of these unknown particles Ξ_{cc}^{+} and Ω_{cc}^{+} have been obtained and are equal to $5036 \text{ MeV}/c^2$ and $5164 \text{ MeV}/c^2$ respectively, as can be seen in Table 2. Since Ξ_{cc}^{*+} and Ξ_{cc}^{*++} are two different states of same particle, i.e. their Iso-spin is equal to $1/2$, therefore both will approximately have equal masses, equal to $5036 \text{ MeV}/c^2$. But how could we guess that these two particles should have same iso-spin? Let us check.

ISO-SPIN OF THE PARTICLES

If we see Fig. 2, values of Iso-spin numbers are decreasing from $I = +3/2$ to $I = 0$ by $-1/2$, from left to right, and then increasing by $+1/2$ till $+3/2$. Similarly, values of the third component of Iso-spin, I_z or I_3 , decreases for particles of same Iso-spin quantum number, at upper horizontal row to particles at lower horizontal row by 1. $SU(4)$ 20-plet of Baryons ($J^P=3/2^+$) shown in figure 2 may be presented with details of I and I_3 , as shown in figure (3). Here iso-spin I and its third component I_3 are presented in following figures as (I, I_3).




Figure 3. $SU(4)$ 20-plet of Baryons ($J^P=3/2^+$) presented with details of I and I_3 . Here iso-spin I and its third component I_3 are presented as (I, I_3).

Here in this figure it may be observed that charge value in layers is increasing from bottom to top. Iso-spin has also a definite value in any diagonal layer (from top left side) from top to

bottom. It is found from figure (2) and (3) that iso-spins of the undiscovered exotic hyperons Ξ_{cc}^{*++} , Ξ_{cc}^{*+} , Ω_{cc}^{*+} and Ω_{ccc}^{++} should be 1/2, 1/2, 0, and 0 respectively. Similarly in order to find third component of iso-spin (I_3) of these particles, in figure (3) it may be observed that I_3 of particles in diagonal layers (from top right side) is decreasing by 1/2 for each layer from top to bottom. By setting the values of I and I_3 for undiscovered hyperons according to the above procedure, it is found that Ξ_{cc}^{*++} , Ξ_{cc}^{*+} , Ω_{cc}^{*+} and Ω_{ccc}^{++} should have values of I_3 equal to 1/2, -1/2, 0 and 0 respectively.

Table 2. Calculations of masses of the particles from sum of other particles, indexed in Fig. 2.

S#	Index	Particle 1 (m_1)	Particle 2 (m_2)	$m_1 + m_2$ (MeV/c ²)	Resultant Particle (m)	$(m_1 + m_2) - m$
1	<i>a</i>	Δ^{++} (1232)	Δ^+ (1232)	2464	Σ_c^{*+} (2518)	-54
2	<i>b</i>	Δ^+ (1232)	Δ^0 (1232)	2464	Σ_c^{*0} (2519)	-55
3	<i>c</i>	Δ^- (1232)	Σ^{*-} (1387)	2619	Ξ_c^{*0} (2646)	-27
4	<i>d</i>	Σ^{*-} (1387)	Ξ^{*-} (1535)	2922	Ω_c^{*0} (2766)	156
5	<i>e</i>	Σ^{*+} (1383)	Ξ^{*0} (1532)	2915	Ω_c^{*0} (2766)	149
6	<i>f</i>	Δ^{++} (1232)	Σ^{*+} (1383)	2615	Ξ_c^{*+} (2646)	-31
7	<i>g</i>	Σ_c^{*++} (2518)	Σ_c^{*+} (2518)	5036	Ξ_{cc}^{*+} (??)	??
8	<i>h</i>	Σ_c^{*0} (2519)	Ξ_c^{*0} (2646)	5165	Ω_{cc}^{*+} (??)	??
9	<i>i</i>	Σ_c^{*++} (2518)	Ξ_c^{*+} (2646)	5164	Ω_{cc}^{*+} (??)	??

Mass of the double charmed Ξ_{cc}^{*++} (probably with $J^P = 1/2^+$) is approximately equal to 3621 MeV/c² [6]. We can compare this value with the theoretical mass of the Ξ_{cc}^{*++} particle with $J^P = 3/2^+$, we extracted i.e. 5036 MeV/c². Percentage difference between these two masses is $\frac{5036-3621}{3621}$, which is equal to 39%. It is in agreement with the percentage mass difference between masses of proton/neutron and delta resonances (having same quark contents, but different spins) $\frac{1232-938}{938}$ equal to 31%. It is acceptable, and hence we can take 5036 MeV/c² as approximate mass of the Ξ_{cc}^{*+} and Ξ_{cc}^{*++} particles with some uncertainty equal to its square root, given by ± 71 MeV/c².




Figure 4. A portion of the SU(4) Multiplets, with increasing Mass, Charge number and Charmness from bottom to top.

ANOTHER TEST

Mass of the Ω_{cc}^+ hyperon having $(J)^P=(1/2)^+$ has been predicted [7] to be $3916 \pm 62 \text{ MeV}/c^2$. Percentage difference between particles with same quark contents but different total spin in average, calculated above (0.39 and 0.31, is equal to 0.35. The mass of the Ω_{cc}^{+*} having $(J)^P=(3/2)^+$ should be:

$$3916 \times 0.35(3916) = 5286 \text{ MeV}/c^2 \approx 5165 \pm 72 \text{ MeV}/c^2$$

Or if we consider it 0.33, it gives exact mass of the Ω_{cc}^{+*} particle:

$$3916 \times 0.33(3916) = 5208 \text{ MeV}/c^2 \approx 5165 \pm 72 \text{ MeV}/c^2$$

III MASS OF THE Ω_{ccc}^{++} PARTICLE

Mass of the Ω_{ccc}^{++} particle can be estimated using Fig. 4. Taking average mass value of first three rows i.e. rows with $C = 0, 1, 2$ and finding their difference may lead us to the mass of the particle at the top, with $C = 3$.

$$M_1 \text{ (Average mass of row with } C = 0) = 1456 \text{ MeV}/c^2$$

$$M_2 \text{ (Average mass of row with } C = 1) = 2644 \text{ MeV}/c^2$$

$$M_3 \text{ (Average mass of row with } C = 2) = 5100 \text{ MeV}/c^2$$

$$M_4 = \text{mass of } \Omega_{ccc}^{++} \text{ particle.}$$

Table 3. Approximated masses and other properties of the unknown particles in Standard Model of $SU(4)$.

S.#	Particle	Mass (MeV/c^2)	Iso spin		Parity (P)	$I(J)^P$
			I	I_z		
1	Ξ_{cc}^{*++}	5036 ± 71	$\frac{1}{2}$	$+\frac{1}{2}$	+1	$\frac{1}{2}(\frac{3}{2})^+$
2	Ξ_{cc}^{*+}	5036 ± 71	$\frac{1}{2}$	$-\frac{1}{2}$	+1	$\frac{1}{2}(\frac{3}{2})^+$
3	Ω_{cc}^{*+}	5165 ± 72	0	0	+1	$0(\frac{3}{2})^+$
4	Ω_{ccc}^{++}	8710 ± 93	0	0	+1	$0(\frac{3}{2})^+$

Difference between M_2 and M_1 :

$$M_2 - M_1 = 2644 - 1456 = 1188 MeV / c^2$$

Similarly;

$$M_3 - M_2 = 5100 - 2644 = 2456 MeV / c^2$$

$$\text{Also } 1188 \times 2 = 2376 MeV / c^2,$$

which is very near to $2456 MeV/c^2$ (with only difference of $80 MeV/c^2$). It means;

$$M_2 - M_1 = 1188 MeV / c^2$$

$$M_3 - M_2 = 2(M_2 - M_1) + 80 = (2 \times 1188) + 80 MeV / c^2$$

and

$$M_4 - M_3 = 3(M_2 - M_1) + 46 = (3 \times 1188) + 46 MeV / c^2$$

Now we can generalize these in one equation:

$$M_n = M_{n-1} + Charmness \times (1188) + Charge \times (\sqrt{1188}) \quad (3)$$

Where $n=2, 3, 4$

That is;

$$\begin{aligned} M_2 &= M_1 + 1 \times (1188) + 0 \times (\sqrt{1188}) \\ &= 1456 + 1188 = 2644 MeV / c^2 \end{aligned}$$

$$\begin{aligned} M_3 &= M_2 + 2 \times (1188) + 1 \times (\sqrt{1188}) \\ &= 2644 + 2376 + 34 = 5054 MeV / c^2 \end{aligned}$$

Here value of M_3 will be average of both the theoretical values, as there are no experimentally observed values for particles in this row. i.e.

$$M_3 = \frac{5100 + 5054}{2} = 5077 MeV / c^2$$

hence

$$M_4 = M_3 + 3 \times (1188) + 2 \times (\sqrt{1188})$$

$$= 5077 + 3564 + 69 = 8710 \text{ MeV} / c^2$$

Thus the mass of Ω_{ccc}^{++} particle, estimated using the above method is equal to $8710 \pm 93 \text{ MeV}/c^2$. Statistical Errors are the square root values of the particle masses. Experimentally observed Particle masses may be slightly differ from the above estimated values of masses of all the four particles, due to rough estimates. However in that case the values obtained experimentally should agree with theoretical mass values with in two standard errors. All the predicted masses of the four unknown particles and their other properties are listed in Table 3.

Now let us check what is the mass difference, $M_2 - M_1 = 1188 \text{ MeV} / c^2$. It is actually mass of the Charm quark, which is the constituent of the particles. Charm number increases by +1 as we go up in the particles pyramid of Fig. 1, and Fig. 4. So this increase in the masses of the particles is due to the mass of the charm quark. Comparing this with the evaluated mass of the charm quark $M_c = 1270 \pm 30$ and $M_c = 1196 \pm 59 \pm 50 \text{ MeV} / c^2$ listed in Ref. [5] gives satisfactory result. Hence it is also proved that splitting due to charm quarks in the $SU(4)$ particles multiplets is about an order of magnitude, equal to an average value of 80%, as was expected [8].

The calculations may also be expanded to the whole 20- multiplet, as shown in Fig. 1, so that a combined formula for all the baryons masses of this multiplet may be extracted.

In the Fig. 1, m_1 shows average mass of the particles forming de-couplet, with $C = 0$, and is equal to $1382 \text{ MeV}/c^2$, $m_2 = 2602 \text{ MeV}/c^2$ is average mass of the particles with $C = 1$, $m_3 = 5079 \text{ MeV}/c^2$ is the average mass of particles with $C = 2$ and $m_4 = 8710 \text{ MeV}/c^2$ is the average mass of particle having $C = 3$.

General expression for these masses may be derived as under;

$$m'_n = m_{n-1} + \text{Charmness} \times (1188) + \text{Charmness} \times (\sqrt{1188})$$

Where $n=2, 3, 4$

Or more generally,

$$\boxed{m'_n = m_{n-1} + \text{Charmness} \times (1218)} \quad (4)$$

with $n=2, 3, 4$

Let's try to find average mass of the particles multiplets using this relation;

$$m'_2 = m_1 + C \times (1218) = 1382 + 1 \times (1218)$$

$$= 2600 \text{ MeV} / c^2 \approx m_2$$

$$\begin{aligned} m'_3 &= m_2 + C \times (1218) = 2602 + 2(1218) \\ &= 5038 \text{ MeV} / c^2 \approx m_3 \end{aligned}$$

and

$$\begin{aligned} m'_4 &= m_3 + C \times (1218) = 5079 + 3(1218) \\ &= 8733 \text{ MeV} / c^2 \approx m_4 \end{aligned}$$

Or equation (4) can also be written as under:

$$\boxed{m'_n - m_{n-1} = 1218(n-1) \text{ MeV} / c^2} \quad (5)$$

with $n = 2, 3, 4$

Mass difference term $1218 \text{ MeV}/c^2$ is also equal to the mass of the charm quark [5]. The mass symmetry braking ratio $\frac{m_2}{m_1} = \frac{m_4}{m_3}$ in the direction of the increasing mass in the Charm plane for $SU(4)$ multiplets, is of the order of an average value of 1.8 for baryons of $J^P=3/2^+$.

III. CONCLUSIONS

$SU(4)$ multiplets of Standard Model have been analyzed in this article and some predictions have been made about the unknown masses and other properties of the Ξ_{cc}^{*++} , Ξ_{cc}^{*+} , Ω_{cc}^{*+} , and Ω_{ccc}^{++} hyperons. Their masses are estimated to be $\Xi_{cc}^{*++} (5036 \pm 71)$, $\Xi_{cc}^{*+} (5036 \pm 71)$, $\Omega_{cc}^{*+} (5165 \pm 72)$, and $\Omega_{ccc}^{++} (8710 \pm 93) \text{ MeV}/c^2$. Statistical errors are the square root values of the estimated masses of particles. Experimentally observed Particles' masses may slightly differ from the estimated values of masses of all the four particles, due to rough estimates. However in that case the values obtained experimentally should agree with theoretical mass values with in two standard errors. The mass symmetry braking ratio for $SU(4)$ multiplets, is estimated to be of the order of an average value of 1.8 for baryons of $J^P=3/2^+$. It is found that Ξ_{cc}^{*++} and Ξ_{cc}^{*+} have same isospin, equal to $1/2$, so both would have approximately equal masses. Ξ_{cc}^{*++} has third component of isospin I_z equal to $+1/2$, while that of Ξ_{cc}^{*+} is equal to $-1/2$. While other two particles have I and I_3 equal to zero. A general formula for masses has been derived for the $SU(4)$ multiplets, which gives satisfactory results for observed and estimated masses of the particles of $SU(4)$. The mass difference in the multiplets of the $SU(4)$ is due to the mass of Charm quark, and its value is also equal to mass of the charm quark. The results in this research, if proved correct, will be helpful in understanding the particle physics, and will strengthen the Standard Model of particles.

ACKNOWLEDGMENTS

Thanks are due to Dr. Khusniddin Olimov who is my Ph.D Supervisor and to Dr. Farida Tahir, who introduced me with Particle Physics and inspired me during a lecture to the need for some mass formula and predictions about masses of missing hyperons of the $SU(4)$ multiplets.

REFERENCES

- [1] Introduction to Elementary Particles, David J. Griffiths, John Wiley & sons, INC, New York, 2008.
- [2] M. Gell-Mann, Proceedings of the International Conference on High-Energy Nuclear Physics, Geneva, Switzerland, 1962.
- [3] S. Okubo, “Note on Unitary Symmetry in Strong Interactions”, *Progr. Theoret. Phys.* (Kyoto) 27, 949 (1962)
- [4] V. E. Barnes *et.al.*, “Observation of a hyperon with strangeness minus three”, *Physical Review Letters*, 12 (8), 204, 1964.
- [5] C. Patrignani *et al.* (Particle Data Group), *Chin. Phys. C*, 40, 100001 (2016)
- [6] R. Aaij *et al.*, “Observation of the doubly charmed baryon Ξ_{cc}^{++} ” *PRL* 119, 112001 (2017)
- [7] Imran Khan, “Expression for $SU(4)$ multiplet and Masses of undiscovered Baryons in Standard Model”, arxiv.org/abs/1904.05155
- [8] M. Gourdin, “Mass Formulae and Mixing in $SU(4)$ Symmetry”, (France) *Pages_from_C75-03-02-1_229.pdf*.