

# A Bayesian inference approach for determining player abilities in soccer

Gavin A. Whitaker<sup>1,2\*</sup>

Ricardo Silva<sup>1,3</sup>

Daniel Edwards<sup>2</sup>

<sup>1</sup>Department of Statistical Science, University College London, London, WC1E 6BT

<sup>2</sup>Stratagem Technologies, 19 Eastbourne Terrace, London, W2 6LG

<sup>3</sup>The Alan Turing Institute, 96 Euston Road, London, NW1 2DB

## Abstract

We consider the task of determining a soccer player’s ability for a given event type, for example, scoring a goal. We propose an interpretable Bayesian inference approach that centres on variational inference methods. We implement a Poisson model to capture occurrences of event types, from which we infer player abilities. Our approach also allows the visualisation of differences between players, for a specific ability, through the marginal posterior variational densities. We then use these inferred player abilities to extend the Bayesian hierarchical model of [Baio and Blangiardo \(2010\)](#), which captures a team’s scoring rate (the rate at which they score goals). We apply the resulting scheme to the English Premier League, capturing player abilities over the 2013/2014 season, before using output from the hierarchical model to predict whether over or under 2.5 goals will be scored in a given fixture or not in the 2014/2015 season.

**Keywords:** Variational inference; Bayesian hierarchical modelling; Soccer; Bayesian inference.

## 1 Introduction

Within this paper we look to determine the ability of those players who play in the English Premier League. The Premier League is an annual soccer league established in 1992 and is the most watched soccer league in the world ([Yueh, 2014](#); [Curley and Roeder, 2016](#)). It is made up of 20 teams, who, over the course of a season, play every other team twice (both home and away), giving a total of 380 fixtures each year. It is the top division of English soccer, and every year the bottom 3 teams are relegated to be replaced by 3 teams from the next division down (the Championship). In recent times the Premier League has also become known as the richest league in the world ([Deloitte, 2016](#)), through both foreign investment and a lucrative deal for television rights ([Cave and Miller, 2016](#); [Rumsby, 2016](#); [BBC Business, 2016](#)). Whilst there is growing financial competition from China, the Premier league arguably still attracts some of the best players in the world. Staying in the Premier league (by avoiding relegation) is worth a large amount of money, therefore teams are looking for any advantage when accessing a player’s ability to ensure they sign the best players. With (enormously) large sums of money spent to buy/transfer these players, it is natural to ask “How good are they at a specific skill, for example, passing a ball, scoring a goal or making a tackle?” Here, we present a method to access this ability, whilst quantifying the uncertainty around any given player.

The statistical modelling of sports has become a topic of increasing interest in recent times, as more data is collected on the sports we love, coupled with a heightened interest in the outcome of these sports, that is, the continuous rise of online betting. Soccer is providing an area of rich

---

\*email: [gavin.whitaker@ucl.ac.uk](mailto:gavin.whitaker@ucl.ac.uk)

research, with the ability to capture the goals scored in a match being of particular interest. [Reep et al. \(1971\)](#) used a negative binomial distribution to model the aggregate goal counts, before [Maher \(1982\)](#) used independent Poisson distributions to capture the goals scored by competing teams on a game by game basis. [Dixon and Coles \(1997\)](#) also used the Poisson distribution to model scores, however they departed from the assumption of independence; the model is extended in [Dixon and Robinson \(1998\)](#). The model of [Dixon and Coles \(1997\)](#) is also built upon in [Karlis and Ntzoufras \(2000, 2003\)](#), who inflate the probability of a draw. [Baio and Blangiardo \(2010\)](#) consider this model in the Bayesian paradigm, implementing a Bayesian hierarchical model for goals scored by each team in a match. Other works to investigate the modelling of soccer scores include ([Lee, 1997](#); [Joseph et al., 2006](#); [Karlis and Ntzoufras, 2009](#)).

A player performance rating system (the EA Sports Player Performance Index) is developed by [McHale et al. \(2012\)](#). The rating system is developed in conjunction with the English Premier League, the English Football League, Football DataCo and the Press Association, and aims to represent a player’s worth in a single number. There is some debate within the soccer community on the weightings derived in the paper, and as [McHale et al. \(2012\)](#) point out, the players who play for the best teams lead the index. There is also some questions raised as to whether reducing the rating to a single number (whilst easy to understand), masks a player’s ability in a certain skill, whether good or bad. Finally, as mentioned by the authors, the rating system does not handle those players who sustain injuries (and therefore have little playing time) well. [McHale and Szczepański \(2014\)](#) attempt to identify the goal scoring ability of players. Spatial methods to capture a team’s style/behaviour are explored in [Lucey et al. \(2013\)](#), [Bialkowski et al. \(2014\)](#) and [Bojinov and Bornn \(2016\)](#). Here, our interest lies in defining that player ability, addressing some of the issues raised by [McHale et al. \(2012\)](#), before attempting to capture the goals scored in a game, taking into account these abilities.

To infer player abilities we appeal to variational inference (VI) methods, an alternative strategy to Markov chain Monte Carlo (MCMC) sampling, which can be advantageous to use when datasets are large and/or models have high complexity. Popularised in the machine learning literature ([Jordan et al., 1999](#); [Wainwright and Jordan, 2008](#)), VI transforms the problem of approximate posterior inference into an optimisation problem, meaning it is easier to scale to large data and tends to be faster than MCMC. Some application areas and indicative references where VI has been used include sports ([Kitani et al., 2011](#); [Ruiz and Perez-Cruz, 2015](#); [Franks et al., 2015](#)), computational biology ([Jojic et al., 2004](#); [Stegle et al., 2010](#); [Carbonetto and Stephens, 2012](#); [Raj et al., 2014](#)), computer vision ([Bishop and Winn, 2000](#); [Likas and Galatsanos, 2004](#); [Blei and Jordan, 2006](#); [Cummins and Newman, 2008](#); [Sudderth and Jordan, 2009](#); [Du et al., 2009](#)) and language processing ([Reyes-Gomez et al., 2004](#); [Wang and Blunsom, 2013](#); [Yogatama et al., 2014](#)). For a discussion on VI techniques as a whole, see [Blei et al. \(2017\)](#) and the references therein.

The remainder of this article is organised as follows. The data is presented in [Section 2](#). In [Section 3](#) we outline our model to define player abilities before discussing a variational inference approach; we finish the section by offering our extension to the Bayesian hierarchical model of [Baio and Blangiardo \(2010\)](#). Applications are considered in [Section 4](#) and a discussion is provided in [Section 5](#).

## 2 The data

The data available to us is a collection of touch-by-touch data, which records every touch in a given fixture, noting the time, team, player, type of event and outcome. A section of the data is shown in [table 1](#). The data covers the 2013/2014 and 2014/2015 English Premier League seasons, and consists of roughly 1.2 million events in total, which equates to approximately 1600 for each fixture in the dataset. There are 39 event types in the dataset, which we list in [table 2](#). The nature of most of these event types is self-explanatory, that is, Goal indicates that a player scored a goal at

minute	second	period	team id	player id	type	outcome
0	1	FirstHalf	663	91242	Pass	Successful
0	2	FirstHalf	663	23736	Pass	Successful
0	3	FirstHalf	663	17	Pass	Successful
0	4	FirstHalf	663	14230	Pass	Successful
0	5	FirstHalf	663	7398	Pass	Successful
0	6	FirstHalf	663	31451	Pass	Successful
0	9	FirstHalf	663	7398	Pass	Successful
0	10	FirstHalf	690	38772	Tackle	Successful
0	10	FirstHalf	663	80767	Dispossessed	Successful
0	12	FirstHalf	690	8505	Pass	Successful

Table 1: A section of the touch-by-touch data.

Stop	Control	Disruption	Miscellanea
Card	Aerial	BlockedPass	CornerAwarded
End	BallRecovery	Challenge	CrossNotClaimed
FormationChange	BallTouch	Claim	KeeperSweeper
FormationSet	ChanceMissed	Clearance	ShieldBallOpp
OffsideGiven	Dispossessed	Interception	
PenaltyFaced	Error	KeeperPickup	
Start	Foul	OffsideProvoked	
SubstitutionOff	Goal	Punch	
SubstitutionOn	GoodSkill	Save	
	MissedShots	Smother	
	OffsidePass	Tackle	
	Pass		
	SavedShot		
	ShotOnPost		
	TakeOn		

Table 2: Event types contained within the data.

that event time. Throughout this paper we will mainly concern ourselves with event types which are self-evident, and will define the more subtle event types as and when needed.

We can split the event types into 4 categories.

1. **Stop:** An event corresponding to a stoppage in play such as a substitution or offside decision.
2. **Control:** An event where a team is perceived to be in control of the ball, these are mainly seen as attacking events.
3. **Disruption:** An event where a team is perceived to be disrupting the current play within a game, these can generally be seen as defensive events.
4. **Unclear:** These events could be classified in any of the other three categories.

In this paper we are interested in those events which correspond to a player during active game-play, hence we remove Stop events from the data. Instead we focus on those event types categorised as either Control or Disruption, or, when a team is attempting to score a goal and when a team is attempting to stop the opposition from scoring a goal respectively.

It should be noted that OffsideGiven is the inverse of OffsideProvoked and as such we remove one of these events from the data. Henceforth, it is assumed that the event type OffsideGiven is removed




Figure 1: Frequency of each event type observed in the Liverpool vs Stoke 2013/2014 English Premier League match, 17th August 2013. The event type Pass is removed for clarity, it occurs with a ratio of approximately 10:1 over BallRecovery.

fixture id	player id	team id	Count for each event type			
			Goal	Pass	Tackle	...
1483412	17	663	0	97	3	
1483412	3817	663	0	37	3	
1483412	4574	663	0	73	3	
⋮	⋮	⋮	⋮	⋮	⋮	
1483831	10136	676	1	36	4	
1483831	12267	676	0	45	0	
1483831	12378	676	0	52	2	
⋮	⋮	⋮	⋮	⋮	⋮	

Table 3: A section of the count data derived from the data of table 1.

from the data, rewarding the defensive side for provoking an offside through OffsideProvoked. The frequency of each event type (after removing Pass) during the Liverpool vs Stoke match, which occurred on the 17th August 2013, is shown in figure 1. The match is typical of any fixture within in the dataset. Pass dominates the data over all other event types recorded, with a ratio of approximately 10:1 to BallRecovery, and hence is removed for clarity. This is not surprising given the make up of a soccer match (where teams mainly pass the ball).

In determining a player’s ability for a given event type we make the assumption that the more times a player is involved, the better they are at that event type; for example, a player who makes more passes than another player is assumed to be the better passer. On this basis, we can transform the data displayed in table 1 to represent the number of each event type each player is involved in, at a fixture by fixture level. This count data is illustrated in table 3. It is to this data, which the methods of Section 3 will be applied.

### 3 Bayesian inference

Consider the case where we have  $K$  matches, numbered  $k = 1, \dots, K$ . We denote the set of teams in fixture  $k$  as  $T_k$ , with  $T_k^H$  and  $T_k^A$  representing the home and away teams respectively. Explicitly,  $T_k = \{T_k^H, T_k^A\}$ . We take  $P$  to be the set of all players who feature in the dataset, and  $P_k^j \in P$  to be the subset of players who play for team  $j$  in fixture  $k$ . We may want to consider how players' abilities over different event types interact, for this we group event types to create meaningful interactions. For simplicity, we describe the model for a single pair of event types which are deemed to interact, for example, Pass and Interception, we denote these event types  $e_1$  and  $e_2$ , such that  $E = \{e_1, e_2\}$ .

Taking  $X_{i,k}^e$  as the number of occurrences (counts) of event type  $e$ , by player  $i$  (who plays for team  $j$ ), in match  $k$ , we have

$$X_{i,k}^e \sim \text{Pois}(\eta_{i,k}^e \tau_{i,k}), \quad (1)$$

where

$$\eta_{i,k}^e = \exp \left\{ \Delta_i^e + \tau_{i,k} \left( \lambda_1^e \sum_{i' \in P_k^j} \Delta_{i'}^e - \lambda_2^e \sum_{i' \in P_k^{T_k \setminus j}} \Delta_{i'}^{E \setminus e} \right) + \left( \delta_{T_k^H, j} \right) \gamma^e \right\}, \quad (2)$$

$\delta_{r,s}$  is the Kronecker delta and  $\tau_{i,k}$  is the fraction of time player  $i$  (playing for team  $j$ ), spent on the pitch in match  $k$ , with  $\tau_{i,k} \in [0, 1]$ . Explicitly, if a player plays 60 minutes of a 90 minute match then  $\tau_{i,k} = 2/3$ . The home effect is represented by  $\gamma^e$ . The home effect reflects the (supposed) advantage the home team has over the away team in event type  $e$ . The  $\Delta_i^e$  represent the (latent) ability of each player for a specific event type, where we let  $\Delta$  be the vector of all players abilities. The impact of a player's own team on the number of occurrences is captured through  $\lambda_1^e$ , with  $\lambda_2^e$  describing the opposition's ability to stop the player in that event type. For identifiability purposes, we impose the constraint that the  $\lambda$ s must be positive. Figure 2 illustrates the model for one fixture, allowing for some abuse in notation, and where we assume each team consists of 11 players only (that is, we ignore substitutions) and suppress the time dependence ( $\tau$ ). From (1) and (2), the log-likelihood is given by

$$\ell = \sum_{e \in E} \sum_{k=1}^K \sum_{j \in T_k} \sum_{i \in P_k^j} X_{i,k}^e \log(\eta_{i,k}^e \tau_{i,k}) - \eta_{i,k}^e \tau_{i,k} - \log(X_{i,k}^e!). \quad (3)$$

Interest lies in estimating this model using a Bayesian approach. We put independent Gaussian priors over all abilities, whilst treating the remaining unknown parameters as hyperparameters — to be fitted by the marginal likelihood function. Given the size of the data and the number of parameters needing to be estimated to fit equation 3, we appeal to variational inference techniques, which are the subject of the next section.

#### 3.1 Variational inference

For a general introduction to variational inference (VI) methods we direct the reader to [Blei et al. \(2017\)](#) (and the references therein). Variational inference is paired with automatic differentiation in [Kucukelbir et al. \(2016\)](#), leading to a technique known as automatic differentiation variational inference (ADVI). ADVI provides an automated solution to VI and is built upon recent approaches in *black-box* VI, see ([Ranganath et al., 2014](#); [Mnih and Gregor, 2014](#); [Kingma and Welling, 2014](#)). [Duvenaud and Adams \(2015\)](#) present some `Python` code (5 lines) for implementing black-box VI. A good overview to VI can be found in Chapter 19 of [Goodfellow et al. \(2016\)](#). Given the set-up of the model above though, it is sufficient within this paper, to require only *standard* variational inference methods. We briefly outline these below, and refer the reader to [Blei et al. \(2017\)](#) for a more complete derivation.




Figure 2: Pictorial representation of the model for one fixture. For ease we assume that only 11 players play for each team in the fixture (that is, we ignore substitutions) and suppress the time dependence ( $\tau$ ).

In contrast to some other techniques for Bayesian inference, such as MCMC, in VI we specify a variational family of densities over the latent variables ( $\nu$ ). We then aim to find the best candidate approximation,  $q(\nu)$ , to minimise the Kullback-Leibler (KL) divergence to the posterior

$$\nu^* = \operatorname{argmin}_{\nu} \operatorname{KL} \{q(\nu) \parallel \pi(\nu|x)\},$$

where  $x$  denotes the data. Unfortunately, due to the analytic intractability of the posterior distribution, the KL divergence is not available in closed (analytic) form. However, it is possible to maximise the evidence lower bound (ELBO). The ELBO is the expectation of the joint density under the approximation minus the entropy of the variational density and is given by

$$\operatorname{ELBO}(\nu) = \mathbb{E}_{\nu} [\log \{\pi(\nu, x)\}] - \mathbb{E}_{\nu} [\log \{q(\nu)\}]. \quad (4)$$

The ELBO is the equivalent of the negative KL divergence up to the constant  $\log\{\pi(x)\}$ , and from [Jordan et al. \(1999\)](#) and [Bishop \(2006\)](#) we know that, by maximising the ELBO we minimise the KL divergence.

In performing VI, assumptions must be made about the variational family to which  $q(\nu)$  belongs. Here we consider the *mean-field variational family*, in which the latent variables are assumed to be mutually independent. Moreover, each latent variable  $\nu_r$  is governed by its own variational parameters ( $\phi_r$ ), which determine  $\nu_r$ 's variational factor, the density  $q(\nu_r|\phi_r)$ . Specifically, for  $R$  latent variables

$$q(\nu|\phi) = \prod_{r=1}^R q_r(\nu_r|\phi_r). \quad (5)$$

We note that the complexity of the variational family determines the complexity of the optimisation, and hence impacts the computational cost of any VI approach. In general, it is possible to impose any graphical structure on  $q(\nu_r|\phi_r)$ ; a fully general graphical approach leads to structured variational inference, see [Saul and Jordan \(1996\)](#). Furthermore, the data ( $x$ ) does not feature in equation 5, meaning the variational family is not a model of the observed data; it is in fact the ELBO which connects the variational density,  $q(\nu|\phi)$ , to the data and the model.

For the model outlined at the beginning of this section, let  $\nu = \Delta$ , and set

$$q(\Delta_i^e|\phi_i^e) \sim N(\mu_{\Delta_i^e}, \sigma_{\Delta_i^e}^2). \quad (6)$$

Our aim is to find suitable candidate values for the variational parameters

$$\phi_i^e = (\mu_{\Delta_i^e}, \sigma_{\Delta_i^e})^T, \quad \forall i, \forall e.$$

Explicitly  $q(\Delta|\phi)$  follows (5). Whence

$$q(\Delta|\phi) = \prod_{e \in E} \prod_{j \in T} \prod_{i \in P^j} q(\Delta_i^e | \phi_i^e), \quad (7)$$

where  $T$  is the set of all teams and  $P^j$  are the players who play for team  $j$ . Finally we take  $\psi = (\lambda_1^{e1}, \lambda_2^{e1}, \gamma^{e1}, \lambda_1^{e2}, \lambda_2^{e2}, \gamma^{e2})^T$  to be fixed parameters, and assume each  $\Delta_i^e$  follows a  $N(m, s^2)$  prior distribution, fully specifying the model given by equations (1)–(3). Thus, the ELBO (4) is given by

$$\begin{aligned} \text{ELBO}(\Delta) &= \sum_{e \in E} \sum_{k=1}^K \sum_{j \in T_k} \sum_{i \in P_k^j} \mathbb{E}_{\Delta_i^e} [\log \{\pi(\Delta_i^e, \phi_i^e, \psi, x)\}] - \mathbb{E}_{\Delta_i^e} [\log \{q(\Delta_i^e | \phi_i^e)\}] \\ &= \sum_{e \in E} \sum_{k=1}^K \sum_{j \in T_k} \sum_{i \in P_k^j} \mathbb{E}_{\Delta_i^e} [\log \{\pi(\Delta_i^e)\}] + \mathbb{E}_{\Delta_i^e} [\log \{\pi(x | \Delta_i^e, \phi_i^e, \psi)\}] \\ &\quad - \mathbb{E}_{\Delta_i^e} [\log \{q(\Delta_i^e | \phi_i^e)\}]. \end{aligned} \quad (8)$$

The above is available in closed-form (see Appendix A), avoiding the need for black-box VI. We do however incorporate the techniques of automatic differentiation for computational ease, and use the Python package `autograd` (Maclaurin et al., 2015) to fit the model.

### 3.2 Hierarchical model

Building on the methods of Section 3.1, we wish to discover whether the inferred  $\Delta$ s have any impact on our ability to predict the goals scored in a soccer match. As a baseline model we consider the work of Baio and Blangiardo (2010), who present the model of Karlis and Ntzoufras (2003) in a Bayesian framework. The model has close ties with (Dixon and Coles, 1997; Lee, 1997; Karlis and Ntzoufras, 2000) which have all previously been used to predict soccer scores. We first briefly outline the model of Baio and Blangiardo (2010), before offering our extension to include the imputed  $\Delta$ s.

The model is a Poisson-log normal model, see for example Aitchison and Ho (1989), Chib and Winkelmann (2001) or Tunaru (2002) (amongst others). For a particular fixture  $k$ , we let  $y^k = (y_h^k, y_a^k)^T$  be the total number of goals scored, where  $y_h^k$  is the number of goals scored by the home team, and  $y_a^k$ , the number by the away team. Inherently, we let  $h$  denote the home team and  $a$  the away team for the given fixture  $k$ . The goals of each team are modelled by independent Poisson distributions, such that

$$y_t^k | \theta_t \stackrel{\text{indep}}{\sim} \text{Pois}(\theta_t), \quad t \in \{h, a\}, \quad (9)$$

where

$$\log(\theta_h) = \text{home} + \text{att}_h + \text{def}_a, \quad (10)$$

$$\log(\theta_a) = \text{att}_a + \text{def}_h. \quad (11)$$

Each team has their own team-specific attack and defence ability, `att` and `def` respectively, which form the scoring intensities ( $\theta_t$ ) of the home and away teams. A constant home effect (`home`), which is assumed to be constant across all teams and across the time-span of the data, is also included in the rate of the home team's goals.




Figure 3: Pictorial representation of the Bayesian hierarchical model. Removing both  $f(\Delta)_h$  and  $f(\Delta)_a$  gives the baseline model of [Baio and Blangiardo \(2010\)](#).

For identifiability, we follow [Baio and Blangiardo \(2010\)](#) and [Karlis and Ntzoufras \(2003\)](#), and impose sum-to-zero constraints on the attack and defence parameters

$$\sum_{t \in T} \text{att}_t = 0 \quad \text{and} \quad \sum_{t \in T} \text{def}_t = 0,$$

where  $T$  is the set of all teams to feature in the dataset. Furthermore, the attack and defence parameters for each team are seen to be draws from a common distribution

$$\text{att}_t \sim N(\mu_{\text{att}}, \sigma_{\text{att}}^2) \quad \text{and} \quad \text{def}_t \sim N(\mu_{\text{def}}, \sigma_{\text{def}}^2).$$

We follow the prior set-up of [Baio and Blangiardo \(2010\)](#) and assume that home follows a  $N(0, 100^2)$  distribution *a priori*, with the hyper parameters having the priors

$$\begin{aligned} \mu_{\text{att}} &\sim N(0, 100^2), & \mu_{\text{def}} &\sim N(0, 100^2), \\ \sigma_{\text{att}} &\sim \text{Inv-Gamma}(0.1, 0.1), & \sigma_{\text{def}} &\sim \text{Inv-Gamma}(0.1, 0.1). \end{aligned}$$

A graphical representation of the model is given in figure 3.

As an extension to the model of [Baio and Blangiardo \(2010\)](#) we propose to include the latent  $\Delta$ s of Section 3.1 in the scoring intensities of both the home and away teams. Explicitly (10) and (11) become

$$\log(\theta_h) = \text{home} + \text{att}_h + \text{def}_a + f(\Delta)_h, \quad (12)$$

$$\log(\theta_a) = \text{att}_a + \text{def}_h + f(\Delta)_a, \quad (13)$$

where  $f(\Delta)$  is to be determined. For a single pair of event types (as outlined at the start of this section), a sensible choice for  $f(\Delta)$  could be

$$f(\Delta)_h = \sum_{i \in I_k^{T^H}} \mu_{\Delta_i^e} - \sum_{i \in I_k^{T^A}} \mu_{\Delta_i^{E \setminus e}} \quad (14)$$

and

$$f(\Delta)_a = \sum_{i \in I_k^{T^A}} \mu_{\Delta_i^e} - \sum_{i \in I_k^{T^H}} \mu_{\Delta_i^{E \setminus e}}, \quad (15)$$



2013/2014 English Premier League teams			
Arsenal	Everton	Manchester United	Sunderland
Aston Villa	Fulham	Newcastle United	Swansea City
Cardiff City	Hull City	Norwich City	Tottenham Hotspur
Chelsea	Liverpool	Southampton	West Bromwich Albion
Crystal Palace	Manchester City	Stoke City	West Ham United

Table 4: The teams which constituted the 2013/2014 English Premier League.

with  $I_k^j$  being the initial eleven players who start fixture  $k$  for team  $j$  and  $\mu_\Delta$  being the mean of the marginal posterior variational densities. This extension is also illustrated in figure 3. We fit both the baseline model of Baio and Blangiardo (2010) and our extension using PyStan (Stan Development Team, 2016). We note that it may be desirable to fit both the model of Section 3.1 and the Bayesian hierarchical model concurrently. However, we find that in reality this is infeasible as the latter can be fit using MCMC, whilst it is difficult to fit the model of Section 3.1 using MCMC due to the large number of parameters.

## 4 Applications

Having outlined our approach to determine a player’s ability in a given event type, and offered an extension to the model of Baio and Blangiardo (2010) to capture the goals scored in a specific fixture, we wish to test the proposed methods in real world scenarios. We therefore consider two applications. In the first we use data from the 2013/2014 English Premier League to learn players abilities across the season as a whole for a number of event types, including the ability to score a goal. The second example concerns the number of goals observed in a given fixture, specifically, we predict whether a certain number of goals will be scored (or not) in each fixture.

### 4.1 Determining a player’s ability

In this section we consider the touch-by-touch data described in Section 2 and consider data for the 2013/2014 English Premier League season only. We look to create an ordering of players abilities, from which we hope to extract meaning based on what we know of the season. We also have data on the amount of time each player spent on the pitch in each match and this information is factored in accordingly through  $\tau_{i,k}$ . The season consisted of 380 matches for the 20 team league, with 544 different players used during matches. The teams are listed in table 4, with the final league table shown in figure 4. From figure 4 we note that Manchester City and Liverpool were the teams who scored the most goals, with Chelsea conceding the least. These teams did well over the season and we expect players from these teams to have high abilities. The teams to do worst (and got relegated), were Norwich City, Fulham and Cardiff; we do not expect players from these teams to feature highly in any ordering created. A final note is that, in this season, Manchester United underperformed (given past seasons) under new manager David Moyes. Whence,  $k = 1, \dots, 380$ ,  $j \in T_k$  where  $T_k$  consists of a subset of  $\{1, \dots, 20\}$  and  $i \in P_k^j$  where  $P_k^j$  is a subset of  $P = \{1, \dots, 544\}$ .

For a pair of interacting event types we fit the model defined by (1)–(3), by maximising (8) where  $q(\cdot)$  follows (6). This model set-up has 2182 parameters governing any two interacting event types. We take the (reasonably uninformative) prior

$$\pi(\Delta_i^e) \sim N(-2, 2^2), \quad (16)$$

where -2 represents the ability of an average player. We found little difference in results for alternative priors. We begin by considering occurrences of Goal and GoalStop. GoalStop is an event type of our own creation (in conjunction with expert soccer analysts), made up of many other event

English Premier League 2013/2014									
#	Team	Pl	W	D	L	F	A	GD	Pts
1	Manchester City	38	27	5	6	102	37	65	86
2	Liverpool	38	26	6	6	101	50	51	84
3	Chelsea	38	25	7	6	71	27	44	82
4	Arsenal	38	24	7	7	68	41	27	79
5	Everton	38	21	9	8	61	39	22	72
6	Tottenham Hotspur	38	21	6	11	55	51	4	69
7	Manchester United	38	19	7	12	64	43	21	64
8	Southampton	38	15	11	12	54	46	8	56
9	Stoke City	38	13	11	14	45	52	-7	50
10	Newcastle United	38	15	4	19	43	59	-16	49
11	Crystal Palace	38	13	6	19	33	48	-15	45
12	Swansea City	38	11	9	18	54	54	0	42
13	West Ham United	38	11	7	20	40	51	-11	40
14	Sunderland	38	10	8	20	41	60	-19	38
15	Aston Villa	38	10	8	20	39	61	-22	38
16	Hull City	38	10	7	21	38	53	-15	37
17	West Bromwich Albion	38	7	15	16	43	59	-16	36
18	Norwich City	38	8	9	21	28	62	-34	33
19	Fulham	38	9	5	24	40	85	-45	32
20	Cardiff City	38	7	9	22	32	74	-42	30

Figure 4: Final league table for the 2013/2014 English Premier League.  $Pl$  matches played,  $W$  matches won,  $D$  matches drawn,  $L$  matches lost,  $F$  goals scored,  $A$  goals conceded,  $GD$  goal difference (scored – conceded),  $Pts$  final points total.

types (BallRecovery, Challenge, Claim, Error, Interception, KeeperPickup, Punch, Save, Smother, Tackle), with BallRecovery being the event type where a player collects the ball after it has gone loose. GoalStop aims to represent all the things a team can do to stop the other team from scoring a goal. A Monte Carlo simulation of the prior for  $\eta_{i,k}^{\text{Goal}}$  using 100K draws of  $\Delta_i^{\text{Goal}}$  from (16) is shown in figure 5, where most players are viewed to score 0 or 1 goal (as expected).

We ran the model for 7000 iterations to achieve convergence. Trace plots of the ELBO and a selection of model parameters are shown in figure 6. It is clear that convergence has been achieved (measured via the ELBO). For completeness the values of the fixed parameters ( $\psi$ ) under the model are given in table 5, where all respective parameters appear to be on the same scale. We observe small differences in the parameters dictating the amount of impact both a player’s own team, and the opposing team has on occurrences of an event type. There are more noticeable differences in the home effects of each event type, with the home effect for Goal being much larger than that of GoalStop. This is in line with other research around the goals scored in a match, where a clear home effect is acknowledged, see Dixon and Coles (1997), Karlis and Ntzoufras (2003) and Baio and Blangiardo (2010) (amongst others) for further discussion of this home effect. The home effect for GoalStop is closer to zero, suggesting the number of attempts a team makes to stop a goal is similar whether they are playing at home or away.

Figure 7 shows the  $\eta_{i,k}^e$  (2) we obtain when the model parameters are combined for 2 randomly selected matches, where we set  $\Delta_i^e$  to be  $\mu_{\Delta_i^e}$ . We plot these against the observed counts and




Figure 5: Monte Carlo simulation of the prior for  $\eta_{i,k}^{\text{Goal}}$  using 100K draws of  $\Delta_i^{\text{Goal}}$  from (16).




Figure 6: Trace plots for the ELBO and a selection of the model parameters outputted every 100 iterations.

Event type ( $e$ )	Fixed parameter		
	$\lambda_1^e$	$\lambda_2^e$	$\gamma^e$
Goal	$2.907 \times 10^{-8}$	0.041	0.165
GoalStop	$1.621 \times 10^{-7}$	0.009	0.003

Table 5: Values of the fixed parameters ( $\psi$ ) for interacting event types Goal and GoalStop.




Figure 7: Within sample predictive distributions for the number of goals/goal-stops in the 2013/2014 English Premier League for 2 randomly selected matches. *Cross* model combinations of  $\eta_{i,k}^e$ , *circle* observed counts, *dashed bars* 95% prediction interval for each  $\eta_{i,k}^e$ . The solid line separates the players from the two teams.

include the 95% prediction intervals for each  $\eta_{i,k}^e$  to add further clarity. The solid line on each plot separates the players from the two opposing teams. A large number of the model  $\eta$ s are close to the observed counts (especially for GoalStop), and nearly all observed values fall within the 95% prediction intervals, showing a reasonable model fit. The number of goal-stops across teams is not particularly variable, however there is a suggestion of player variability (although this is somewhat clouded by the fact that substitutes are not specifically marked, as we would expect them to register lower counts, by virtue of less playing time).

We sample the marginal posterior variational densities,  $q(\Delta_i^e)$ , 10K times (constructing the corresponding  $\eta_{i,k}^e$ ), and simulate from the relevant Poisson distributions (with mean  $\eta_{i,k}^e \tau_{i,k}$ ). This gives a Monte Carlo simulation of each player's number of goals and goal-stops for each fixture in the 2013/2014 English Premier League. Summing over the players who played in a given match, gives an in-sample prediction of the total number of goals/goal-stops for each team, in every fixture. We present box-plots of these totals in figure 8, where for reference, we also include box-plots for each team's total number of goals/goal-stops in each fixture constructed from the touch-by-touch data. The model is clearly capturing the patterns between differing teams (and the patterns observable within the data), especially for goal-stop. The teams who scored the most goals over the season, Manchester City and Liverpool, have higher Goal box-plots than other teams, encompassing a larger range of goals scored. On the other hand, teams who scored few goals over the season, such as Norwich City, have the lowest box-plots, which cover a small range of goals scored in a match. Slightly surprisingly, there appears to be no connection between the occurrences of GoalStop and




Figure 8: Box-plots of the total number of goals/goal-stops in a game for each team in the 2013/2014 English Premier League observed under the model and from the data. *Top row* Goal, *bottom row* GoalStop.

the goals a team concedes, with both Chelsea and Norwich City having similar box-plots, despite conceding a vastly different number of goals, 27 and 62 respectively. There is some suggestion that such observations may be used to determine a team’s style of play, for example, whether they are a *passing* team or follow the *long ball* philosophy; however, we leave such questions for future investigation given the set-up we derive here. We can conclude, nevertheless, that the model is capturing the trends observed in the touch-by-touch data well.

Marginal posterior variational densities of Goal,  $q(\Delta_i^{\text{Goal}})$ , for two players are presented in figure 9, where  $q(\Delta_i^{\text{Goal}})$  takes the form of (6) and the prior is (16). The two players shown are Daniel Sturridge and Harrison Reed. Sturridge played 29 times over the season, totalling 2414 minutes of match time, scoring 21 goals, whereas Reed played 4 times, totalling 23 minutes, scoring zero goals. These attributes are clearly captured by the posteriors; the greater number of observations for Sturridge leading to a posterior with a much smaller variance. The high number of goals scored by Sturridge leads to him having a higher value of  $\mu_{\Delta_i^{\text{Goal}}}$  (with reasonable certainty), whilst the lack of both goals and playing time leads to a posterior for Reed which resembles the prior.

The model is clearly capturing differences between players abilities, as evidenced by the posteriors of figure 9. Thus, the natural question to ask is whether these differences are sensible, and, if we were to order the players by their inferred ability, would this ordering agree with (a debatable) reality. Hence, we construct the marginal posterior variational densities for all players and rank them according to the 2.5% quantile of these densities. A top 10 for Goal is presented in table 6, with a ranking for GoalStop given in table 7. We present top 10 lists for other event types in




Figure 9: Marginal posterior variational densities of Goal for 2 players in the 2013/2014 English Premier League. *Dashed* prior, *solid* posterior.

Goal - top 10								
Rank	Player	2.5% quantile	Mean	Standard deviation	Observed	Observed rank	Rank difference	Time played
1	Suarez	0.508	0.869	0.184	31	1	0	3185
2	Sturridge	0.176	0.617	0.225	21	2	0	2414
3	Aguero	0.147	0.636	0.250	17	4	+1	1616
4	Y. Toure	-0.043	0.395	0.224	20	3	-1	3113
5	Rooney	-0.056	0.421	0.243	17	5	0	2625
6	Dzeko	-0.065	0.424	0.249	16	8	+2	2128
7	van Persie	-0.136	0.430	0.289	12	15	+8	1690
8	Remy	-0.230	0.302	0.271	14	11	+3	2274
9	Bony	-0.257	0.238	0.252	16	7	-2	2644
10	Rodriguez	-0.354	0.161	0.263	15	10	0	2758

Table 6: Top 10 goal scorers in the 2013/2014 English Premier League based on the 2.5% quantile of the marginal posterior variational density for each player,  $q(\Delta_i^{\text{Goal}})$ .

Appendix B. The ranking shown in table 6 appears sensible, and comprises of those players who were the main goal scorers (strikers) for the best teams, and the players who scored nearly all the goals a lesser team scored over the season. The ranking is very close to that obtained by ranking players on the total number of goals scored over the season (although there is some debate in the soccer community as to whether this is a sensible way of ranking, with some suggesting a ranking based on a per 90 minute statistic, however, this can be distorted by those with very little playing time, see Chapter 3 of [Anderson and Sally \(2013\)](#) or [AGR Analytics \(2016\)](#) for further discussion). The questionable deviations from this ranking are Aguero (ranked third) and van Persie (ranked seventh). Both these players have less playing time over the season compared to their competitors, and thus, the model highlights them as better goal scorers, given the time available to them, than other players based on total goals scored. Expert soccer analysts agreed with this view when we showed them these rankings. Suarez has an inferred ability much greater than any other player, which is evidenced by the 31 goals he scored (10 more than any other player). At points the difference between successive ranks is small, suggesting some players are harder to distinguish between. Finally, we note that the standard deviations for all players in the top 10 are roughly the same, meaning we have similar confidence in the ability of any of these players.

GoalStop - top 10								
Rank	Player	2.5% quantile	Mean	Standard deviation	Observed	Observed rank	Rank difference	Time played
1	Mulumbu	2.575	2.653	0.040	631	1	0	3319
2	Kallstrom	2.553	2.900	0.177	33	405	+403	144
3	Mannone	2.528	2.615	0.044	508	12	+9	2767
4	Yacob	2.510	2.614	0.053	359	43	+39	1979
5	Tiote	2.474	2.560	0.044	517	8	+3	2988
6	Lewis	2.446	2.863	0.213	23	436	+430	98
7	Palacios	2.441	2.638	0.101	100	286	+279	585
8	Jedinak	2.420	2.500	0.041	603	2	-6	3651
9	Ruddy	2.411	2.491	0.041	600	3	-6	3679
10	Arteta	2.409	2.503	0.048	431	21	+11	2615

Table 7: Top 10 goal-stoppers in the 2013/2014 English Premier League based on the 2.5% quantile of the marginal posterior variational density for each player,  $q(\Delta_i^{\text{GoalStop}})$ .

The ranking of GoalStop (table 7) appears, at first glance, to be less sensible than that of table 6. It features 3 players with comparatively larger standard deviations, Kallstrom (rank 2), Lewis (rank 6) and Palacios (rank 7). Whilst these players did well with the little playing time afforded to them, it is somewhat presumptuous to postulate that they would maintain a similar level of ability given more game time, leading to their ranking slipping. Unfortunately, this is a by product of inducing a ranking from the 2.5% quantile — ideally we would provide several tables for each event type, filtering players by the amount of uncertainty surrounding them, although such an approach would be unwieldy given the large number of players in the dataset. Moreover, fully factorised mean-field approximations are known to underestimate the uncertainty of the posterior (Bishop, 2006). Although comparative uncertainty between players is easier to gauge, it is less clear how to quantify how much bias is being added to the variances of each latent variable individually. In a future work, this could be mitigated by adopting a variational approximation that accounts for some correlations of the latent variables. However, the rest of the list appears sensible and is made up mainly of defensive midfielders (whose main role it is to disrupt the oppositions play); only Mannone and Ruddy are goalkeepers (discounting the 3 players with large standard deviations). This suggests, that to stop a goal, it is more prudent to invest in a better defensive midfielder than it is a goalkeeper, presuming you can not just buy the best player in each position. Here, the differences between successive ranks are much smaller than in table 6, implying it is harder to distinguish between player ability to perform goal-stops than it is the ability to score goals.

Overall though, the model provides a good fit to the data and suggests a reasonable prowess to determine a player’s ability in a specific event type, with marginal posterior variational densities providing a good visual comparison between different players abilities (and the confidence surrounding that ability). In the next section we look to utilise these player abilities in the prediction of goals in a soccer match.

## 4.2 Prediction

A key betting market, stemming from the rise of online betting, is the over/under market (Betfair, 2017; betHQ, 2017; SPORTINGINDEX, 2017), where people bet whether a certain number of goals will (over) or won’t (under) be scored in a match. Here we attempt to predict whether 2.5 goals (a number of goals common with online betting) will be scored or not in a given fixture. To predict the goals scored in a fixture which takes place in the future we first fit the model on all the past available to us. We use a whole season of data to train the model, before predicting the following




Figure 10: Illustration of the approach to prediction. The model is fit on all past data, before predictions are made for a future block of fixtures. *Solid* fit, *dashed* predict.

season in incremental blocks. Here, we use the entirety of the 2013/2014 English Premier League season (380 fixtures) to train the model, before attempting to predict the goals scored in each match of the 2014/2015 English Premier League season. We introduce the fixtures (on which we predict) in blocks of size 80, with a final block of 60 fixtures to total 380 (the number of fixtures over a season). In each case we use all of the available past to fit the model, that is, in predicting the second block of 80 fixtures in the 2014/2015 season, we use all of the 2013/2014 season and the first block of the 2014/2015 season to fit the model. Figure 10 shows a graphical representation of this approach.

For the extension to the model of Baio and Blangiardo (2010) (which we consider to be the baseline model), we include the latent player abilities for the event types Goal, Shots and ChainEvents, with their counterparts being GoalStop, ShotStop and AntiPass respectively. Goal and GoalStop are as defined in Section 4.1, whilst Shots and ShotStop have homogeneous roots to Goal and GoalStop, that being the ability to shoot or to stop a shot. ChainEvents represents how prevalent a player is in the lead up to a good attacking chance, with AntiPass being a player’s ability to stop the other team from passing the ball. We refer the reader to Appendix B for the more technical definitions of these event types. Explicitly (14) and (15) are given by

$$f(\Delta)_h = \sum_{i \in I_k^{T^H}} \left( \Delta_i^{\text{Goal}} + \Delta_i^{\text{Shots}} + \Delta_i^{\text{ChainEvents}} \right) - \sum_{i \in I_k^{T^A}} \left( \Delta_i^{\text{GoalStop}} + \Delta_i^{\text{ShotStop}} + \Delta_i^{\text{AntiPass}} \right) \quad (17)$$

and

$$f(\Delta)_a = \sum_{i \in I_k^{T^A}} \left( \Delta_i^{\text{Goal}} + \Delta_i^{\text{Shots}} + \Delta_i^{\text{ChainEvents}} \right) - \sum_{i \in I_k^{T^H}} \left( \Delta_i^{\text{GoalStop}} + \Delta_i^{\text{ShotStop}} + \Delta_i^{\text{AntiPass}} \right), \quad (18)$$

where  $I_k^j$  is the initial eleven players who start fixture  $k$  for team  $j$ . We also considered including a player’s ability to pass, but found this led to no increase in predictive power (and in some instances diminished it). We found little difference when setting  $\Delta_i^e$  to be either  $\mu_{\Delta_i^e}$  or the 2.5% quantile of  $q(\Delta_i^e)$  in (17) and (18), and so here report results for the mean ( $\mu_{\Delta_i^e}$ ).

Both the models were fit using PyStan, and were run long enough to yield a sample of approximately 10K independent posterior draws, after an initial burn in period. The teams which feature in the data are those in table 4, with the addition of Burnley, Leicester City and Queens



Park Rangers, who replaced the relegated teams of Cardiff City, Fulham and Norwich City for the 2014/2015 season. The set-up outlined at the beginning of this section allows us to view the evolution of a team’s attack/defence parameter or a player’s latent ability through time after different fitting blocks. We denote block 0 to be all the fixtures in the 2013/2014 English Premier League season, block 1 to be block 0 plus the first 80 fixtures of the 2014/2015 season, block 2 to be block 1 plus the next 80 fixtures, block 3 to include the next 80 fixtures, with block 4 including the next 80 fixtures again. The attack and defence parameters through time for both the baseline model and the model including the latent player abilities for selected teams are shown in figure 11, where we plot negative defence so that positive values indicate increased ability. Recall that these parameters for all teams must sum-to-zero. We see similar, but not identical, patterns under both models. The model including latent player abilities reduces the variance of the attack and defence parameters compared to the baseline model, suggesting the inclusion of the  $\Delta$ s accounts for some of a team’s attack and defensive ability. Manchester City and Chelsea follow similar patterns under both models, with Chelsea clearly having the best defence parameter. Including the  $\Delta$ s impacts Liverpool’s attacking ability, where the removal of Suarez (Liverpool’s best attacking player, who transferred to Barcelona between the 2013/2014 and 2014/2015 seasons), clearly reduces Liverpool’s attacking threat at a more drastic rate than the baseline model. This is inline with reality, where Liverpool only scored 52 goals over the 2014/2015 season compared to 101 goals the previous year. Cardiff City, who got relegated after the 2013/2014 season, but feature in all blocks despite not being used for prediction (as fixtures involving them can inform the attack and defence abilities of other teams), have relatively constant parameters under both models, accounting for the reduction in variance. Notable for Burnley is the peak/trough observed after block 3; this is due to the fact that Burnley were starting to look at the prospect of relegation and needed to start winning games, hence, they tried (and succeeded) to score more goals in order to win games, but found themselves more likely to concede goals in the process of doing so.

The mean of  $q(\Delta_i^{\text{Goal}})$  through time for a selection of players are illustrated in figure 12, we let this value represent a player’s ability to score a goal. If a player does not feature in a block we represent their ability by the mean of the prior distribution (-2). We see that the model is quick to identify a given player’s ability. To elucidate, Costa is immediately (after block 1) identified as one of the top goal scorers, despite not featuring in the 2013/2014 season. The same can be said for Sanchez, who takes longer to establish his ability after a less impressive start to the season. Aguero was one of the best goal scorers across all the data and has a constant ability near the top. Kane had little playing time until block 2 where the model starts to increase his ability to score a goal. Defoe spent most of the 2013/2014 season on the bench before transferring to Toronto FC; he returned to the English Premier League with Sunderland in January 2015, where he scored a number of goals, saving Sunderland from relegation. The model rightly acknowledges this and raises his ability as a goal scorer (a trait he is well known for). Given a player scores a small number of goals, relative to other event types, we include G. Johnson to show the effect of scoring a goal. Johnson scored 1 goal in the 2013/2014 and 2014/2015 seasons (during block 2), and his ability rises by a large jump because of this; such jumps are not evident for players who score a reasonable number of goals (5+).

The mean of  $q(\Delta_i^{\text{Control}})$  and the mean of  $q(\Delta_i^{\text{Disruption}})$  are plotted against each other through time for a selection of players in figure 13. Control and Disruption comprise of the event types listed in table 2. It is evident that for the majority of players, their Control and Disruption abilities do not vary much through time (from block to block). This is perhaps unsurprising, given we do not expect a player’s ability to change dramatically from fixture to fixture. Those that vary the most are the players with fewer minutes played in the earlier blocks, but have much more playing time as time progresses, for example, Kane (see figure 13). The figure does however show clear distinction between players, with defenders tending to occupy the top half of the graph, and strikers the bottom half. An interesting extension to this work would be to see whether a clustering analysis




Figure 11: Attack and defence parameters through time under the baseline model and the model including the latent player abilities for selected teams. *Top row* attack, *bottom row* negative defence. *Black-solid* Liverpool, *black-dashed* Chelsea, *black-dotted* Manchester City, *grey-solid* Cardiff City, *grey-dashed* Burnley.




Figure 12: The mean of  $q(\Delta_i^{\text{Goal}})$  through time for a selection of players. *Black-solid* Costa, *black-dashed* A. Sanchez, *black-dotted* Aguero, *grey-solid* Defoe, *grey-dashed* G. Johnson, *grey-dotted* Kane.




Figure 13: The mean of  $q(\Delta_i^{\text{Control}})$  versus the mean of  $q(\Delta_i^{\text{Disruption}})$  through time for a selection of players. *Triangle* block 0, *square* block 4. *Black-solid* Agüero, *black-dashed* A. Carroll, *black-dotted* Koscielny, *grey-solid* G. Johnson, *grey-dashed* Silva, *grey-dotted* Kane.

of these latent player abilities would reveal player positions, that is, central defender or wing-back for example.

To form our predictions of whether over or under 2.5 goals are scored in a given fixture, we take each of our posterior draws (fitted using the previous block) and construct the  $\theta_t$  of (9) via (10) and (11) (baseline model), or (12) and (13) (including latent player abilities) for the fixtures in the following block (our prediction block). Our prediction blocks are formed of fixtures between teams we have already seen in the previous (fitting) blocks, hence prediction block 1 consists of 57 fixtures, prediction blocks 2-4 are made up of 80 fixtures, with 60 fixtures in prediction block 5. We use a predicted starting line-up from expert soccer analysts to determine  $I_k^j$ , the players who enter (17) and (18); these are usually quite accurate (86% accuracy over the season) and vary little from the players who start a particular game. We then combine the  $\theta_t$  for the home and away teams to give an overall scoring rate for each fixture,  $\theta = \theta_h + \theta_a$ , from which we calculate the probability of there being over 2.5 goals in the match. We average these probabilities across the posterior sample. ROC curves based on these averaged probabilities, for each prediction block, are presented in figure 14. For clarity we also present the area under the curve (AUC) values in table 8.

It is evident from both the figure and the table, that including the latent player abilities in the model leads to a better predictive performance. We observe this increase across all blocks, although the difference between the models in block 5 is severely reduced compared to other blocks. The reasons for this reduction are twofold, the first being that given a near full season of data (2014/2015) on which we are predicting, the baseline model can reasonably accurately capture a team's attack and defence parameters better than it can towards the start of the season. Secondly the last block of a season tends to be more volatile, as some teams try out younger players (who are not observed in the data previously), and others have increased motivation to score more goals to try and win games, for example, to avoid relegation. Whence, we observe similar behaviour under both models, as we observe less players in a starting line-up, moving the model including player abilities towards the baseline model. However, overall, we can conclude that the inclusion of the latent player abilities in the model results in a better predictive performance throughout the 2014/2015 season.




Figure 14: ROC curves based on averaged probabilities for each prediction block. *Black* model including the latent player abilities, *grey* baseline model, the *dashed line* is the line  $y = x$ .

Area under the curve values					
Model	Block				
	1	2	3	4	5
Baseline	0.47	0.60	0.53	0.55	0.61
Including latent player abilities	0.54	0.65	0.58	0.68	0.62

Table 8: Area under the ROC curves based on averaged probabilities for each prediction block under both models.

## 5 Discussion

We have provided a framework to establish player abilities in a Bayesian inference setting. Our approach is computationally efficient and centres on variational inference methods. By adopting a Poisson model for occurrences of event types we are able to infer a player’s ability for a multitude of event types. These inferences are reasonably accurate and have close ties to reality, as seen in Section 4.1. Furthermore, our approach allows the visualisation of differences between players, for a specific ability, through the marginal posterior variational densities.

We also extended the Bayesian hierarchical model of Baio and Blangiardo (2010) to include these latent player abilities. Through this model we captured a team’s propensity to score goals, including a team’s attacking ability, defensive ability and accounting for a home effect. We used output from this model to predict whether 2.5 goals would be scored in a fixture or not, observing an improvement in performance over the baseline model. A benefit of the prediction approach (and the block structure we implemented), is that, it allowed us to see how our inference about a players ability evolved through time, explicitly highlighting what impact fringe players can have when they start getting regular playing time, for example, Kane in Section 4.2.

We plan three major ways of extending the current work. First, we intend to extend the variational approximation to allow for dependency among the latent abilities in the posterior. Allowing for correlations in  $q(\cdot)$  will let the model infer higher posterior variances, resulting in a more robust ranking of players, and possibly improved predictive power for tasks such as providing probabilities on the number of goals in a future match. From a modelling perspective, an extension is to let abilities change over time using a random walk across seasons and within seasons, which will be particularly useful when a substantial number of years of historical touch-by-touch data eventually becomes available. Finally, as the model gets applied to more competitions simultaneously, it will be important to propose ways of scaling up the procedure. A topic worth investigating is how to best iteratively subsample the data for stochastic optimisation of the variational objective function.

## A Closed-form expression for the ELBO

Recall the ELBO of Section 3.1 (4) is available in closed-form. Below we consider the terms in (8) on an individual basis to derive this closed-form. Let us begin by considering  $E_{\Delta_i^e}[\log\{q(\Delta_i^e|\phi_i^e)\}]$ . From (6) we have

$$\log\{q(\Delta_i^e)\} = -\frac{1}{2}\log\left(2\pi\sigma_{\Delta_i^e}^2\right) - \frac{(\Delta_i^e - \mu_{\Delta_i^e})^2}{2\sigma_{\Delta_i^e}^2}.$$

Taking expectations gives

$$\begin{aligned} E_{\Delta_i^e} \left[ \log \left\{ q \left( \Delta_i^e \mid \phi_i^e \right) \right\} \right] &= -\frac{1}{2} \log \left( 2\pi\sigma_{\Delta_i^e}^2 \right) \\ &\quad - \frac{1}{2\sigma_{\Delta_i^e}^2} \left[ E \left\{ (\Delta_i^e)^2 \right\} - 2E(\Delta_i^e) \mu_{\Delta_i^e} + (\mu_{\Delta_i^e})^2 \right] \\ &= -\frac{1}{2} \log \left( 2\pi\sigma_{\Delta_i^e}^2 \right) \\ &\quad - \frac{1}{2\sigma_{\Delta_i^e}^2} \left\{ \sigma_{\Delta_i^e}^2 + \mu_{\Delta_i^e}^2 - 2\mu_{\Delta_i^e}^2 + \mu_{\Delta_i^e}^2 \right\} \\ &= -\frac{1}{2} \log \left( 2\pi\sigma_{\Delta_i^e}^2 \right) - \frac{1}{2}, \end{aligned} \tag{19}$$

which is the negative entropy of the Gaussian distribution.

Let us now turn our attention to evaluating  $E_{\Delta_i^e}[\log\{\pi(\Delta_i^e)\}]$ , where  $\Delta_i^e$  follows a  $N(m, s^2)$  prior. Therefore

$$\log\{\pi(\Delta_i^e)\} = -\frac{1}{2}\log(2\pi s^2) - \frac{(\Delta_i^e - m)^2}{2s^2}.$$

Hence

$$\begin{aligned} E_{\Delta_i^e}[\log\{\pi(\Delta_i^e)\}] &= -\frac{1}{2}\log(2\pi s^2) \\ &\quad - \frac{1}{2s^2} \left[ E\{(\Delta_i^e)^2\} - 2E(\Delta_i^e)m + m^2 \right] \\ &= -\frac{1}{2}\log(2\pi s^2) - \frac{\sigma_{\Delta_i^e}^2 + \mu_{\Delta_i^e}^2 - 2m\mu_{\Delta_i^e} + m^2}{2s^2}. \end{aligned} \quad (20)$$

Finally, let us consider  $E_{\Delta_i^e}[\log\{\pi(x|\Delta_i^e, \phi_i^e, \psi)\}]$ . From (3) we have

$$E_{\Delta_i^e} \left[ \log \left\{ \pi \left( x | \Delta_i^e, \phi_i^e, \psi \right) \right\} \right] = \sum_{k=1}^K \underbrace{E \left\{ X_{i,k}^e \log(\eta_{i,k}^e \tau_{i,k}) \right\}}_{\star} - \underbrace{E(\eta_{i,k}^e \tau_{i,k})}_{\dagger} - \log(X_{i,k}^e!).$$

Evaluating  $\star$  first gives

$$\begin{aligned} \star &= E \left[ X_{i,k}^e \left\{ \Delta_i^e + \tau_{i,k} \left( \lambda_1^e \sum_{i' \in P_k^j} \Delta_{i'}^e - \lambda_2^e \sum_{i' \in P_k^{T_k \setminus j}} \Delta_{i'}^{E \setminus e} \right) + (\delta_{T_k^H, j}) \gamma^e + \log(\tau_{i,k}) \right\} \right] \\ &= X_{i,k}^e \left\{ \mu_{\Delta_i^e} + \tau_{i,k} \left( \lambda_1^e \sum_{i' \in P_k^j} \mu_{\Delta_{i'}^e} - \lambda_2^e \sum_{i' \in P_k^{T_k \setminus j}} \mu_{\Delta_{i'}^{E \setminus e}} \right) + (\delta_{T_k^H, j}) \gamma^e + \log(\tau_{i,k}) \right\}. \end{aligned}$$

Turning to  $\dagger$ , we have

$$\begin{aligned} \dagger &= E \left[ \exp \left\{ \Delta_i^e + \tau_{i,k} \left( \lambda_1^e \sum_{i' \in P_k^j} \Delta_{i'}^e - \lambda_2^e \sum_{i' \in P_k^{T_k \setminus j}} \Delta_{i'}^{E \setminus e} \right) + (\delta_{T_k^H, j}) \gamma^e \right\} \tau_{i,k} \right] \\ &= E \left[ \exp \left\{ (1 + \tau_{i,k} \lambda_1^e) \Delta_i^e \right\} \exp \left( \tau_{i,k} \lambda_1^e \sum_{i' \in P_k^j \setminus i} \Delta_{i'}^e \right) \exp \left( -\tau_{i,k} \lambda_2^e \sum_{i' \in P_k^{T_k \setminus j}} \Delta_{i'}^{E \setminus e} \right) \exp \left\{ (\delta_{T_k^H, j}) \gamma^e \right\} \tau_{i,k} \right] \\ &= \exp \left\{ (1 + \tau_{i,k} \lambda_1^e) \mu_{\Delta_i^e} + \frac{(1 + \tau_{i,k} \lambda_1^e)^2 \sigma_{\Delta_i^e}^2}{2} \right\} \exp \left\{ \tau_{i,k} \lambda_1^e \sum_{i' \in P_k^j \setminus i} \mu_{\Delta_{i'}^e} + \frac{(\tau_{i,k} \lambda_1^e)^2 \sum_{i' \in P_k^j \setminus i} \sigma_{\Delta_{i'}^e}^2}{2} \right\} \\ &\quad \times \exp \left\{ -\tau_{i,k} \lambda_2^e \sum_{i' \in P_k^{T_k \setminus j}} \mu_{\Delta_{i'}^{E \setminus e}} + \frac{(-\tau_{i,k} \lambda_2^e)^2 \sum_{i' \in P_k^{T_k \setminus j}} \sigma_{\Delta_{i'}^{E \setminus e}}^2}{2} \right\} \exp \left\{ (\delta_{T_k^H, j}) \gamma^e \right\} \tau_{i,k}. \end{aligned}$$

Therefore

$$\begin{aligned}
E_{\Delta_i^e} \left[ \log \left\{ \pi \left( x \mid \Delta_i^e, \phi_i^e, \psi \right) \right\} \right] &= \sum_{k=1}^K \left( X_{i,k}^e \left\{ \mu_{\Delta_i^e} + \tau_{i,k} \left( \lambda_1^e \sum_{i' \in P_k^j} \mu_{\Delta_{i'}^e} - \lambda_2^e \sum_{i' \in P_k^{T_k \setminus j}} \mu_{\Delta_{i'}^{E \setminus e}} \right) \right. \right. \\
&\quad \left. \left. + \left( \delta_{T_k^H, j} \right) \gamma^e + \log \left( \tau_{i,k} \right) \right\} \right. \\
&\quad \left. - \left[ \exp \left\{ \left( 1 + \tau_{i,k} \lambda_1^e \right) \mu_{\Delta_i^e} + \frac{\left( 1 + \tau_{i,k} \lambda_1^e \right)^2 \sigma_{\Delta_i^e}^2}{2} \right\} \right. \right. \\
&\quad \left. \left. \times \exp \left\{ \tau_{i,k} \lambda_1^e \sum_{i' \in P_k^j \setminus i} \mu_{\Delta_{i'}^e} + \frac{\left( \tau_{i,k} \lambda_1^e \right)^2 \sum_{i' \in P_k^j \setminus i} \sigma_{\Delta_{i'}^e}^2}{2} \right\} \right. \right. \\
&\quad \left. \left. \times \exp \left\{ -\tau_{i,k} \lambda_2^e \sum_{i' \in P_k^{T_k \setminus j}} \mu_{\Delta_{i'}^{E \setminus e}} + \frac{\left( -\tau_{i,k} \lambda_2^e \right)^2 \sum_{i' \in P_k^{T_k \setminus j}} \sigma_{\Delta_{i'}^{E \setminus e}}^2}{2} \right\} \right. \right. \\
&\quad \left. \left. \times \exp \left\{ \left( \delta_{T_k^H, j} \right) \gamma^e \right\} \tau_{i,k} \right] - \log \left\{ X_{i,k}^e ! \right\} \right). \quad (21)
\end{aligned}$$

Thus the closed form of the ELBO (8) is obtained through a combination of (19)–(21) whilst summing over  $i, j, k$  and  $e$ , or explicitly

$$\text{ELBO}(\Delta) = \sum_{e \in E} \sum_{j \in T_k} \sum_{i \in P_k^j} (\text{20}) + (\text{21}) - (\text{19}). \quad (22)$$

## B Top 10 results

In this section we detail top 10 rankings for a number of event types not considered in Section 4.1, namely Shots, ShotStop, ChainEvents and AntiPass, which are presented in tables 9, 10, 11 and 12 respectively. All 4 event types are of our own creation, made up of many other event types.

- **Shots:** Goal, MissedShots, SavedShot, ShotOnPost.
- **ShotStop:** Challenge, Claim, Interception, KeeperPickup, Punch, Save, Smother, Tackle.
- **AntiPass:** BallRecovery, BlockedPass, Claim, Clearance, CornerAwarded, CrossNotClaimed, Interception, KeeperPickup, OffsideProvoked, Punch, Smother, Tackle.

Goal features in Shots, as a successful shot on target leads to a goal, unless it becomes a SavedShot. ChainEvents is created by counting the number of instances a player is involved in the last 5 successful events leading to an event type contained within Shots, that is, the number of times a player is involved in a chain leading to a good attacking chance. The last 5 events were chosen as the length of the chain after discussion with expert soccer analysts, who thought that any further events back from the chance would have had little impact in creating it.

The top 10 for Shots consists entirely of strikers, the person seen as the main scorer of goals in a team, and thus, the person likely to have the most shots. The ranking appears sensible, with the players heightened in our ranking (Aguero, Kane, Jovetic and A. Carroll), playing less time over the season due to injury, or mainly featuring as a substitute. The model suggests they took a large

Shots - top 10								
Rank	Player	2.5% quantile	Mean	Standard deviation	Observed	Observed rank	Rank difference	Time played
1	Suarez	1.426	1.571	0.074	181	1	0	3185
2	Aguero	1.269	1.481	0.108	86	12	+10	1616
3	Dzeko	1.220	1.414	0.099	103	5	+2	2128
4	Kane	1.038	1.413	0.191	28	126	+122	549
5	Bony	1.036	1.225	0.096	108	3	-2	2644
6	Sturridge	1.035	1.233	0.101	99	9	+3	2414
7	Jovetic	1.027	1.441	0.211	23	150	+143	440
8	Remy	0.998	1.206	0.106	90	11	+3	2274
9	A. Carroll	0.982	1.258	0.141	51	53	+44	1200
10	Jelavic	0.978	1.210	0.118	72	18	+8	1804

Table 9: Top 10 shooters in the 2013/2014 English Premier League based on the 2.5% quantile of the marginal posterior variational density for each player,  $q(\Delta_i^{\text{Shots}})$ .

number of shots with the limited time they played. Suarez has an ability greater than any other player by a reasonable amount, which is expected given he had nearly 70 more shots than anyone else over the season. Over the 2013/2014 English Premier League season Suarez was regarded as the best player, winning many awards, it is therefore unsurprising that he features highly in many of the top 10 rankings.

The ranking for ShotStop is made up completely from goalkeepers, a natural conclusion given the event type. Lewis tops the ranking, although he only played 1 game and has a much larger standard deviation than anyone else. The goalkeepers for Fulham (Stockdale and Stekelenburg) played roughly half the season each, both stopping shots well (and at a similar level) when playing, for this reason they feature higher in our rankings than the observed order (determined by the total shots stopped over the season) suggests.

The top 10 for ChainEvents is similar in ways to that of GoalStop (table 6). It features a number of players with less playing time and therefore larger standard deviations. Whilst most of these players play a reasonable amount of time, from which to draw conclusions about their ability, the obvious outlier is Teixeira who played only 14 minutes (and has a very large standard deviation, comparatively). Again Suarez features highly in the rankings. The top 10 contains the *creative* players for each team, with that player for the top teams all featuring, Silva - Manchester City, Coutinho - Liverpool, Hazard - Chelsea. When we showed this ranking to expert soccer analysts there was a consensus that the ordering made sense (with the obvious exception of Teixeira).

The ranking for AntiPass consists of both defenders and defensive midfielders, both types of player whose job it is to disrupt play. Alcaraz and Kallstrom have comparatively larger standard deviations, but the remainder of the ranking appears sensible. The difference between successive rankings is small, and there is some suggestion that it is easier to distinguish between player attacking ability than player defensive ability (GoalStop, ShotStop, AntiPass). This would agree with some in the soccer community who view attacking as an individual ability, whereas defending is more of a team ability. Overall, all 4 of the rankings presented in this appendix appear largely sensible, and agree with expert soccer analysts views.



ShotStop - top 10								
Rank	Player	2.5% quantile	Mean	Standard deviation	Observed	Observed rank	Rank difference	Time played
1	Lewis	2.413	2.823	0.209	23	369	+368	98
2	Mannone	2.394	2.485	0.046	453	8	+6	2767
3	Ruddy	2.312	2.394	0.042	555	1	-2	3679
4	Guzan	2.258	2.343	0.043	512	2	-2	3684
5	Stockdale	2.225	2.343	0.060	267	19	+14	1866
6	Marshall	2.223	2.310	0.044	497	3	-3	3594
7	Stekelenburg	2.218	2.340	0.062	252	24	+17	1790
8	Howard	2.218	2.306	0.045	483	5	-3	3575
9	Szczesny	2.199	2.286	0.045	484	4	-5	3594
10	Adrian	2.187	2.306	0.061	262	20	+10	1943

Table 10: Top 10 shot-stoppers in the 2013/2014 English Premier League based on the 2.5% quantile of the marginal posterior variational density for each player,  $q(\Delta_i^{\text{ShotStop}})$ .

ChainEvents - top 10								
Rank	Player	2.5% quantile	Mean	Standard deviation	Observed	Observed rank	Rank difference	Time played
1	Teixeira	2.419	3.291	0.444	5	474	+473	14
2	Suarez	2.408	2.486	0.040	546	1	-1	3185
3	Eikrem	2.349	2.622	0.139	43	323	+320	229
4	Jovetic	2.303	2.525	0.113	72	250	+246	440
5	Silva	2.299	2.394	0.049	372	5	0	2308
6	Coutinho	2.241	2.338	0.049	362	6	0	2473
7	Taarabt	2.230	2.426	0.100	89	214	+207	639
8	Ramirez	2.218	2.415	0.101	92	208	+200	601
9	Aguero	2.213	2.330	0.060	251	32	+23	1616
10	Hazard	2.197	2.282	0.043	471	2	-8	3100

Table 11: Top 10 players involved in the last 5 interactions leading to a chance in the 2013/2014 English Premier League based on the 2.5% quantile of the marginal posterior variational density for each player,  $q(\Delta_i^{\text{ChainEvents}})$ .

AntiPass - top 10								
Rank	Player	2.5% quantile	Mean	Standard deviation	Observed	Observed rank	Rank difference	Time played
1	Alcaraz	2.864	3.031	0.085	135	292	+291	532
2	Vidic	2.855	2.940	0.043	520	35	+33	2256
3	Skrtel	2.814	2.885	0.036	759	2	-1	3468
4	Kallstrom	2.793	3.107	0.160	39	419	+415	144
5	Lovren	2.789	2.865	0.039	640	10	+5	2993
6	Koscielny	2.783	2.860	0.039	632	11	+5	2980
7	Mulumbu	2.778	2.852	0.038	700	5	-2	3319
8	Azpilicueta	2.769	2.853	0.043	528	34	+26	2522
9	Jedinak	2.763	2.833	0.036	771	1	-8	3651
10	Fonte	2.762	2.835	0.037	709	4	-6	3430

Table 12: Top 10 anti-passers in the 2013/2014 English Premier League based on the 2.5% quantile of the marginal posterior variational density for each player,  $q(\Delta_i^{\text{AntiPass}})$ .

## References

- AGR Analytics (2016). Explaining and examining per 90. <http://alexrathke.net/2016/07/explaining-and-examining-per-90>. 14
- Aitchison, J. and Ho, C. H. (1989). The multivariate poisson-log normal distribution. *Biometrika*, 76(4):643–653. 7
- Anderson, C. and Sally, D. (2013). *The Numbers Game: Why Everything You Know About Football is Wrong*. Penguin Books Limited. 14
- Baio, G. and Blangiardo, M. (2010). Bayesian hierarchical model for the prediction of football results. *Journal of Applied Statistics*, 37(2):253–264. 1, 2, 7, 8, 9, 10, 16, 21
- BBC Business (2016). Premier League in record £5.14bn TV rights deal. <http://www.bbc.co.uk/news/business-31379128>. 1
- Betfair (2017). Over Under 2.5 Goals Betting Advice on Betfair. <https://betting.betfair.com/over-under-25-goals-betting-advice-on-betfair.html>. 15
- betHQ (2017). Over/under goals betting. <https://www.bethq.com/how-to-bet/articles/overunder-goals-betting>. 15
- Bialkowski, A., Lucey, P., Carr, P., Yue, Y., Sridharan, S., and Matthews, I. (2014). Identifying team style in soccer using formations learned from spatiotemporal tracking data. In *Data Mining Workshop (ICDMW), 2014 IEEE International Conference on*, pages 9–14. IEEE. 2
- Bishop, C. and Winn, J. (2000). Non-linear Bayesian image modelling. *Computer Vision-ECCV 2000*, pages 3–17. 2
- Bishop, C. M. (2006). *Pattern Recognition and Machine Learning*. Information Science and Statistics. Springer. 6, 15
- Blei, D. M. and Jordan, M. I. (2006). Variational inference for Dirichlet process mixtures. *Bayesian analysis*, 1(1):121–143. 2
- Blei, D. M., Kucukelbir, A., and McAuliffe, J. D. (2017). Variational inference: A review for statisticians. *Journal of the American Statistical Association*, (just-accepted). 2, 5
- Bojinov, I. and Bornn, L. (2016). The pressing game: Optimal defensive disruption in soccer. In *MIT Sloan Sports Analytics Conference*. MIT SSAC. 2
- Carbonetto, P. and Stephens, M. (2012). Scalable variational inference for Bayesian variable selection in regression, and its accuracy in genetic association studies. *Bayesian analysis*, 7(1):73–108. 2
- Cave, A. and Miller, A. (2016). Why football’s TV deal is a game changer. <http://www.telegraph.co.uk/investing/business-of-sport/premier-league-investors/>. 1
- Chib, S. and Winkelmann, R. (2001). Markov chain Monte Carlo analysis of correlated count data. *Journal of Business & Economic Statistics*, 19(4):428–435. 7
- Cummins, M. and Newman, P. (2008). FAB-MAP: Probabilistic localization and mapping in the space of appearance. *The International Journal of Robotics Research*, 27(6):647–665. 2
- Curley, J. P. and Roeder, O. (2016). English soccer’s mysterious worldwide popularity. <https://contexts.org/articles/english-soccers-mysterious-worldwide-popularity/>. 1

- Deloitte (2016). Deloitte’s annual review of football finance. <https://www2.deloitte.com/uk/en/pages/sports-business-group/articles/annual-review-of-football-finance.html>. 1
- Dixon, M. J. and Coles, S. G. (1997). Modelling association football scores and inefficiencies in the football betting market. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 46(2):265–280. 2, 7, 10
- Dixon, M. J. and Robinson, M. (1998). A birth process model for association football matches. *Journal of the Royal Statistical Society: Series D (The Statistician)*, 47(3):523–538. 2
- Du, L., Ren, L., Carin, L., and Dunson, D. B. (2009). A Bayesian model for simultaneous image clustering, annotation and object segmentation. In *Advances in neural information processing systems*, pages 486–494. 2
- Duvenaud, D. and Adams, R. P. (2015). Black-box stochastic variational inference in five lines of python. In *NIPS Workshop on Black-box Learning and Inference*. 5
- Franks, A., Miller, A., Bornn, L., and Goldsberry, K. (2015). Characterizing the spatial structure of defensive skill in professional basketball. *The Annals of Applied Statistics*, 9(1):94–121. 2
- Goodfellow, I., Bengio, Y., and Courville, A. (2016). *Deep Learning*. MIT Press. <http://www.deeplearningbook.org>. 5
- Jojic, V., Jojic, N., Meek, C., Geiger, D., Siepel, A., Haussler, D., and Heckerman, D. (2004). Efficient approximations for learning phylogenetic HMM models from data. *Bioinformatics*, 20:161–168. 2
- Jordan, M. I., Ghahramani, Z., Jaakkola, T. S., and Saul, L. K. (1999). An introduction to variational methods for graphical models. *Machine Learning*, 37(2):183–233. 2, 6
- Joseph, A., Fenton, N. E., and Neil, M. (2006). Predicting football results using Bayesian nets and other machine learning techniques. *Knowledge-Based Systems*, 19(7):544–553. 2
- Karlis, D. and Ntzoufras, I. (2000). On modelling soccer data. *Student*, 3:229–245. 2, 7
- Karlis, D. and Ntzoufras, I. (2003). Analysis of sports data by using bivariate Poisson models. *Journal of the Royal Statistical Society: Series D (The Statistician)*, 52(3):381–393. 2, 7, 8, 10
- Karlis, D. and Ntzoufras, I. (2009). Bayesian modelling of football outcomes: using the skellam’s distribution for the goal difference. *IMA Journal of Management Mathematics*, 20(2):133–145. 2
- Kingma, D. P. and Welling, M. (2014). Auto-encoding variational Bayes. In *The International Conference on Learning Representations (ICLR), Banff*. 5
- Kitani, K. M., Okabe, T., Sato, Y., and Sugimoto, A. (2011). Fast unsupervised ego-action learning for first-person sports videos. In *Computer Vision and Pattern Recognition (CVPR), IEEE Conference on*, pages 3241–3248. 2
- Kucukelbir, A., Tran, D., Ranganath, R., Gelman, A., and Blei, D. M. (2016). Automatic differentiation variational inference. *arXiv preprint arXiv:1603.00788*. 5
- Lee, A. J. (1997). Modeling scores in the Premier League: Is Manchester United really the best? *Chance*, 10(1):15–19. 2, 7
- Likas, A. C. and Galatsanos, N. P. (2004). A variational approach for Bayesian blind image deconvolution. *IEEE transactions on signal processing*, 52(8):2222–2233. 2

- Lucey, P., Oliver, D., Carr, P., Roth, J., and Matthews, I. (2013). Assessing team strategy using spatiotemporal data. In *Proceedings of the 19th ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 1366–1374. ACM. 2
- Maclaurin, D., Duvenaud, D., and Adams, R. P. (2015). Autograd: Effortless gradients in numpy. In *ICML 2015 AutoML Workshop*. 7
- Maher, M. J. (1982). Modelling association football scores. *Statistica Neerlandica*, 36(3):109–118. 2
- McHale, I. G., Scarf, P. A., and Folker, D. E. (2012). On the development of a soccer player performance rating system for the English Premier League. *Interfaces*, 42(4):339–351. 2
- McHale, I. G. and Szczepański, Ł. (2014). A mixed effects model for identifying goal scoring ability of footballers. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 177(2):397–417. 2
- Mnih, A. and Gregor, K. (2014). Neural variational inference and learning in belief networks. In *Proceedings of the 31st International Conference on Machine Learning*, pages 1791–1799. 5
- Raj, A., Stephens, M., and Pritchard, J. K. (2014). fastSTRUCTURE: Variational inference of population structure in large SNP data sets. *Genetics*, 197(2):573–589. 2
- Ranganath, R., Gerrish, S., and Blei, D. M. (2014). Black box variational inference. In *AISTATS*, pages 814–822. 5
- Reep, C., Pollard, R., and Benjamin, B. (1971). Skill and chance in ball games. *Journal of the Royal Statistical Society. Series A (General)*, pages 623–629. 2
- Reyes-Gomez, M. J., Ellis, D. P. W., and Jojic, N. (2004). Multiband audio modeling for single-channel acoustic source separation. In *Acoustics, Speech, and Signal Processing. IEEE International Conference on*. 2
- Ruiz, F. J. R. and Perez-Cruz, F. (2015). A generative model for predicting outcomes in college basketball. *Journal of Quantitative Analysis in Sports*, 11(1):39–52. 2
- Rumsby, B. (2016). Premier League clubs to share £8.3 billion TV windfall. <http://www.telegraph.co.uk/sport/football/12141415/Premier-League-clubs-to-share-8.3-billion-TV-windfall.html>. 1
- Saul, L. and Jordan, M. I. (1996). Exploiting tractable substructures in intractable networks. In *Advances in Neural Information Processing Systems 8*, pages 486–492. MIT Press. 6
- SPORTINGINDEX (2017). Most popular spread betting markets. <https://www.sportingindex.com/#/o/learn/training-centre/most-popular-markets>. 15
- Stan Development Team (2016). PyStan: the Python interface to Stan, version 2.15.0.0. 9
- Stegle, O., Parts, L., Durbin, R., and Winn, J. (2010). A Bayesian framework to account for complex non-genetic factors in gene expression levels greatly increases power in eQTL studies. *PLoS Comput Biol*, 6(5):e1000770. 2
- Sudderth, E. B. and Jordan, M. I. (2009). Shared segmentation of natural scenes using dependent Pitman-Yor processes. In *Advances in Neural Information Processing Systems*, pages 1585–1592. 2

- Tunaru, R. (2002). Hierarchical Bayesian models for multiple count data. *Austrian Journal of statistics*, 31(3):221–229. 7
- Wainwright, M. J. and Jordan, M. I. (2008). Graphical models, exponential families, and variational inference. *Foundations and Trends in Machine Learning*, 1(1-2):1–305. 2
- Wang, P. and Blunsom, P. (2013). Collapsed variational Bayesian inference for hidden Markov models. In *Artificial Intelligence in Statistics*, pages 599–607. 2
- Yogatama, D., Wang, C., Routledge, B. R., Smith, N. A., and Xing, E. P. (2014). Dynamic language models for streaming text. *Transactions of the Association for Computational Linguistics*, 2:181–192. 2
- Yueh, L. (2014). Exporting football. Why does the world love the English Premier League? <http://www.bbc.co.uk/news/business-27369580>. 1