

Fully Bayesian Estimation Under Informative Sampling

Luis G. León-Novelo[‡] and Terrance D. Savitsky[§]

*1200 Pressler St. Suite E805, Houston, TX, 77030, USA and
PSB Suite 1950, 2 Massachusetts Avenue, NE Washington, DC 20212-0001, USA.
e-mail: Luis.G.LeonNovelo@uth.tmc.edu; Savitsky.Terrance@bls.gov*

Abstract: Survey data are often collected under informative sampling designs where subject inclusion probabilities are designed to be correlated with the response variable of interest. The data modeler seeks to estimate the parameters of a population model they specify from these data. Sampling weights constructed from marginal inclusion probabilities are typically used to form an exponentiated pseudo likelihood that adjusts the population likelihood for estimation on the sample. We propose an alternative adjustment based on a Bayes rule construction that simultaneously performs weight smoothing and estimates the population model parameters in a fully Bayesian construction. We formulate conditions on known marginal and pairwise inclusion probabilities that define a class of sampling designs where L_1 consistency of the joint posterior is guaranteed. We compare performances between the two approaches on synthetic data. We demonstrate that the credibility intervals under our fully Bayesian method achieve nominal coverage. We apply our method to data from the National Health and Nutrition Examination Survey to explore the relationship between caffeine consumption and systolic blood pressure.

Keywords and phrases: Bayesian penalized B-splines, Informative sampling, Inclusion probabilities, NHANES, Sampling weights, Survey sampling.

Contents

1	Introduction	2
1.1	Review of Methods to Account for Informative Sampling	3
2	Fully Bayesian Estimator under Bayes Rule	5
2.1	Linear Regression Population Model	8
2.2	Other Population Models	9
2.3	Splines Basis Population Model	9
2.4	Estimation of Population Distribution of Inclusion Probabilities	10
2.5	Pseudolikelihood Alternative to the Fully Bayes Estimator	12
3	Posterior Consistency	12
3.1	Set-up	13
3.2	Main Result	14
4	Simulation Study	17

[‡]Department of Biostatistics and Data Science, University of Texas Health Science Center at Houston- School of Public Health

[§]Office of Survey Methods Research, Bureau of Labor Statistics

4.1	Simple Linear Regression (SLR): PPS Design	17
4.1.1	Scenarios SLR: π -skewed.	17
4.1.2	Scenario SLR: π -symmetric.	20
4.1.3	Scenario non-linear: π -skewed.	21
5	Application	25
6	Discussion	29
	Supplementary Material	29
7	Proof of Theorem 1	30
8	Closed-Form for log likelihood of Simple Linear Regression Model . . .	31
8.1	Enabling Lemmas for Theorem 2	31
9	Data preprocessing in Section 5	35
	References	36

1. Introduction

Our focus is on the data analyst who seeks to conduct inference about a finite population generating model, P_0 , where the observed data are acquired through a survey sample taken from a finite population of interest. The data analyst has a chosen population model, P , about which they desire to perform inference or prediction and seeks to estimate its parameters taking into account an informative sampling design that governs the observed sample (because an informative sampling design is not ignorable as it is under simple random sampling). Our focus on population model estimation contrasts with the usual design-based estimation of simple domain statistics (*e.g.* employment rate by county) that are often of interest to statistical agencies. Of course, a well-constructed population model, P , that accounts for dependencies within the observed data may also be used to construct simple domain-level statistics in an efficient manner.

Bayesian hierarchical model specifications are increasingly used to perform inference about P_0 due to their flexibility for modeling dependencies among observations and the advent of “black box” solvers, especially “Stan” [Carpenter et al., 2016], which performs an efficiently-mixing Hamiltonian Monte Carlo sampling algorithm with a feature that non-conjugate model specifications are readily accommodated. Bayesian population models can be flexibly parameterized and, since they do not rely on asymptotic theory, are suitable for small sample inference.

A survey sampling procedure randomly draws the observed sample from a target finite population on which inference is performed about P_0 . A known sampling design distribution, P_ν , defines a joint distribution over the random inclusions of units from the finite population. The sampling design distribution often intentionally constructs a correlation between specified unit inclusion probabilities and the response variable of interest; for example, the U.S. Bureau of Labor Statistics (BLS) employs proportion-to-size (PPS) sampling design in the Current Employment Statistics (CES) survey. The CES is administered with the purpose to construct total employment statistics for area and industry-indexed domains. Establishments with larger employment values are assigned

higher unit inclusion probabilities since these larger establishments drive more of the variance in total employment point estimates for domains. Survey designs that instantiate a correlation between unit inclusions and the response variable of interest are termed, “informative”, because the inclusion probabilities are informed by the response variable. The balance of information in the observed sample is different from that in the population under informative sampling, such that naively estimating a proposed population model, P , on the observed sample will result in biased inference about P_0 , that is supposed to govern the finite population [Savitsky and Toth, 2016]. We construct a fully Bayesian estimator in the sequel that accounts for joint distribution, (P_0, P_ν) , over population generation and the taking of an informative sample

1.1. Review of Methods to Account for Informative Sampling

There are three broad classes of approaches to adjust model estimation on the observed sample acquired under an informative sampling (IS) design for inference about P_0 . One approach parameterizes the sampling design into the model estimated on the sampled data [Little, 2004]. The sampling design, itself, is often a nuisance to the data analyst, however, and their focus on the parameters of P_0 requires them to marginalize over parameters indexing the sampling design distribution. It is also often the case that the analyst does not know the sampling design to parameterize it.

The next two classes of modeling adjustments employ a sampling weight that is constructed to be inversely proportional to the marginal inclusion probability, $\pi_i = P(\delta_i = 1)$, for each unit, $i \in (1, \dots, n)$, where n denotes the number of units in the observed sample, S . The inclusion of unit, i , from the population, U , in the sample, $S \subset U$ is indexed by the random variables, $\delta_i \in \{0, 1\}$, governed by P_ν . The likelihood contribution for each unit in the observed sample is adjusted by its associated sampling weight, such that the joint adjusted likelihood over the sample provides an approximation to the balance of information in the finite population.

The second class of modeling approaches employs a particular form for the likelihood [Dong et al., 2014, Kunihama et al., 2016, Rao and Wu, 2010, Si et al., 2015], rather than allowing the analyst to specify a population model, P , because the inferential focus is not on parameters of the generating model, P_0 , but on domain-level estimation of simple mean and total statistics, such as total employment for a geographic area and industry combination.

In the third class of modeling approaches, Savitsky and Toth [2016] construct a sampling-weighted pseudo posterior distribution by exponentiating each unit likelihood contribution, under the analyst-specified model, by its sampling weight, to produce, $p(y_i | \delta_i = 1, \boldsymbol{\lambda})^{w_i}$. Exponentiating by the sampling weight, $w_i \propto 1/\pi_i$, constructs the pseudo likelihood used to estimate the pseudo posterior when convolved with the prior distributions for model parameters, $\boldsymbol{\lambda}$. Savitsky and Toth [2016] demonstrate that estimation of (the parameters of) P_0 from the pseudo posterior distribution is asymptotically unbiased. This approach provides a “plug-in” approximation to the population likelihood (for n

observations), in that the sampling inclusion probabilities, (π_i) , are assumed fixed. On the one hand, one may suppose that a sequential generation for the finite population under P_0 and the subsequent taking of the sample under P_ν (where the two distributions are convolved under informative sampling). In this case, one may view the information contained in the generated finite population as fixed. Inclusion probabilities are assigned by P_ν to the units comprising the population, and so, are also viewed as fixed (given the underlying finite population). From a Bayesian perspective, on the other hand, there is a dynamic process of population generation and the taking of an informative sample. If the population changes, then sample inclusion probabilities, which depend on the population response values, will also change.

The absence of an established procedure to incorporate the inclusion probabilities into the likelihood of a particular population model chosen by the data analyst hinders the use of Bayesian statistics in the analysis of survey data released with sampling weights. To address this lack of a fully Bayesian treatment for modeling data under informative sampling, this paper formulates a fully Bayesian set-up to jointly model (y_i, π_i) from P on the observed sample. Our approach extends the set-up of Pfeffermann et al. [1998] to a fully Bayesian formulation by specifying a conditional population model, $p(\pi_i | y_i, \boldsymbol{\kappa})$, for the inclusion probabilities, $(\pi_i)_{i \in U}$. We then apply a Bayes rule approach to construct, $p(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}, \delta_i = 1)$, that conditions on the observed sample, $(\delta_i = 1)_{i=1, \dots, N}$, where N denotes the population size, $|U|$. We employ this full likelihood defined on the observed sample to estimate the joint posterior distribution for model parameters, $(\boldsymbol{\theta}, \boldsymbol{\kappa})$. We demonstrate in the sequel that the informativeness of the sampling design improves estimation efficiency for our Fully Bayes approach that jointly models (y_i, π_i) as compared to the plug-in, pseudo posterior framework. We formulate conditions that guarantee the L_1 frequentist consistency of our fully Bayes estimator. We further conduct a simulation study that demonstrates the credibility sets generated under our fully Bayes construction achieve nominal coverage (and we also demonstrate, by contrast, that credibility sets of the sampling-weighted, pseudo posterior distribution do not achieve nominal coverage).

Pfeffermann et al. [1998] focus on maximum likelihood point estimation from $p(y_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}, \delta_i = 1)$, rather than the joint distribution over the response and inclusion probabilities, as do we. We also relax their theoretical condition of independence among the sampled units that they employ to guarantee consistency of the point estimate to asymptotic independence in our result for the L_1 consistency of our fully Bayes posterior distribution.

Pfeffermann et al. [2006] also extend Pfeffermann et al. [1998] to Bayesian estimation, but they employ $p(y_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}, \delta_i = 1)$, (without specifying a model for the (π_i)) rather than our formulation for the joint likelihood, which perhaps conforms to the view that the inclusion probabilities, (π_i) , are *not* generated along with the population, $(y_i, \mathbf{x}_i)_{i=1, \dots, N}$, $N = |U|$, but are fixed under specification of the sampling design distribution, P_ν . So their approach may be viewed to be *not* fully Bayesian as is ours because it does not specify a model for $p(\pi_i | y_i, \boldsymbol{\kappa})$.

We derive the joint likelihood for the observed sample, $p(y_i, \pi_i | \delta_i = 1, \dots)$,

using a Bayes rule approach that adjusts the analyst-specified population likelihood, $p(y_i, \pi_i | \dots)$, from which we formulate our joint posterior distribution for the observed sample in Section 2. Conditions are constructed that guarantee a frequentist L_1 contraction of our fully Bayesian posterior distribution, P , onto the true generating distribution in Section 3. This section can be skipped without loss of understanding the practical application of the proposed method. We conduct a simulation study under a synthetically generated population from a known, P_0 , from which we draw samples under an informative, proportion-to-size (PPS) sampling design, to compare the performances of our fully Bayesian estimator and the pseudo posterior in Section 4. We reveal that: (1) fully Bayes credible intervals achieve nominal frequentist coverage while the pseudo posterior under covers; and (2) the fully Bayes point estimates are robust, in terms of bias and mean square error (MSE), against high variability of inclusion probabilities in contrast to the pseudo posterior. We illustrate our fully Bayesian estimator by assessing the relationship between caffeine consumption and systolic blood pressure using NHANES data in Section 5. The paper concludes with a discussion in Section 6. Supplementary Material is provided in Sections 7-9. This includes the proofs for the main result, along with two enabling results in Section 8.

2. Fully Bayesian Estimator under Bayes Rule

We proceed to formulate a posterior distribution for the *observed* sample taken under an informative design as a function of inclusion probabilities to produce unbiased inference and correct uncertainty quantification for the population model. Our approach first constructs a joint distribution for the response and inclusion probabilities for the *population* in a fully Bayesian specification. We then use Bayes rule to adjust the joint population likelihood to an expression for the observed sample by incorporating conditional distributions for inclusion indicators, ($\delta_i \in \{0, 1\}$). The adjusted likelihood is unbiased for the population model parameters estimated on the sample. We later show that credibility sets of the posterior distribution estimated from the fully Bayes adjusted likelihood achieve nominal coverage.

From Bayes rule,

$$p(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}, \delta_i = 1) = \frac{\Pr(\delta_i = 1 | y_i, \pi_i, \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) \times p(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa})}{\Pr(\delta_i = 1 | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa})}, \quad (1)$$

where the expression on the left-hand side of Equation (1) adjusts the joint population likelihood, $p(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa})$, specified on the right-hand side for the realized sample.

We may simplify the numerator by plugging in [See also equation (7.1) in [Pfeffermann et al., 1998](#)],

$$\Pr(\delta_i = 1 | y_i, \pi_i, \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) = \pi_i. \quad (2)$$

We next compute the double expectation with respect to $(\boldsymbol{\kappa}, \boldsymbol{\theta})$ in the denominator,

$$\begin{aligned} \Pr(\delta_i = 1 | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) &= \int \int \Pr(\delta_i = 1 | y_i, \pi_i, \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) p(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) d\pi_i dy_i \quad (3) \\ &= \int \int \pi_i p(\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa}) d\pi_i p(y_i | \mathbf{x}_i, \boldsymbol{\theta}) dy_i \\ &= \int E(\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa}) p(y_i | \mathbf{x}_i, \boldsymbol{\theta}) dy_i \\ &= E_{y_i | \mathbf{x}_i, \boldsymbol{\theta}} [E(\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa})], \end{aligned}$$

where we assume separability of the parameter sets, $\boldsymbol{\kappa}, \boldsymbol{\theta}$, in the conditional distributions for $\pi_i | y_i, \mathbf{x}_i$ and $y_i | \mathbf{x}_i$,

$$\begin{aligned} p(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) &= p(\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa}) p(y_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) \quad (4) \\ &= p(\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa}) p(y_i | \mathbf{x}_i, \boldsymbol{\theta}), \end{aligned}$$

where $p(\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa})$ and $p(y_i | \mathbf{x}_i, \boldsymbol{\theta})$ denote conditional distributions for the population. We make note that we have factorized the joint population distribution over (y_i, π_i) , such that information about the sampling design is parameterized in the conditional model for $\pi_i | y_i$, rather than in the conditional distribution for $y_i | x_i$.

We define $p_s(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) = p(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}, \delta_i = 1)$ (where subscript, s , indexes the observed sample of size, n) to be the likelihood contribution for each unit, $i \in S$, and plug in Equations (2) and (3) into Equation (1) to obtain,

$$p_s(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) = \frac{\pi_i p(\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa})}{E_{y'_i | \mathbf{x}_i, \boldsymbol{\theta}} [E(\pi'_i | y'_i, \mathbf{x}_i, \boldsymbol{\kappa})]} \times p(y_i | \mathbf{x}_i, \boldsymbol{\theta}) \quad (5)$$

Let $N = |U|$ denote the population size and $S = \{S(1), \dots, S(n)\} := \{i \in \{1, \dots, N\} : \delta_i = 1\}$, the population indices of individuals or units included in the random sample S of size $n = |S|$. Therefore, $\Pr(S = s) \propto \prod_{i=1}^n \pi_{s(i)}$. Let the generic vector $\mathbf{r}^{(s)} := \{r_1^{(s)}, \dots, r_n^{(s)}\} := \{r_{s(1)}, \dots, r_{s(n)}\}$ represent the values of variable, r , observed in the sample. With this notation the likelihood over the observed sample of size, n , is specified by,

$$\ell(\boldsymbol{\theta}, \boldsymbol{\kappa}; \mathbf{y}^{(s)}, \boldsymbol{\pi}^{(s)}, \mathbf{x}^{(s)}) = \prod_{i=1}^n [p_s(y_i^{(s)}, \pi_i^{(s)} | \mathbf{x}_i^{(s)}, \boldsymbol{\theta}, \boldsymbol{\kappa})]$$

from which we may incorporate the prior distributions to formulate the posterior distribution observed on the sample,

$$p_s(\boldsymbol{\theta}, \boldsymbol{\kappa} | \mathbf{y}^{(s)}, \boldsymbol{\pi}^{(s)}, \mathbf{x}^{(s)}) \propto \ell(\boldsymbol{\theta}, \boldsymbol{\kappa}; \mathbf{y}^{(s)}, \boldsymbol{\pi}^{(s)}, \mathbf{x}^{(s)}) \times \text{Prior}(\boldsymbol{\theta}) \times \text{Prior}(\boldsymbol{\kappa}).$$

We have now constructed the backbone of a fully Bayesian model that takes into account informative sampling designs for estimation of population model

parameters. Our result jointly models the response and the inclusion probabilities, *i.e.*, (y_i, π_i) , using only quantities observed in the sample; in particular, the joint distribution of (y_i, π_i) are different in the observed sample and in the population, and we have corrected for this difference in a way that allows us to make unbiased estimation of the parameters of the population model. In contrast to Si et al. [2015], we do not impute the values for the non sampled units. By assuming a population distribution for the inclusion probabilities, the variability of the inclusion probabilities is modeled and the noise not depending on the response (and \mathbf{x}_i) is discarded; more specifically $p_s(y_i, \pi_i | \dots)$ in Equation (5) is constructed from a conditional model that regresses π_i on y_i and uses the expected value in the denominator.

Achieving our result of Equation (5) requires specification of a joint population generating distribution, $p(y_i, \pi_i | \mathbf{x}_i, \boldsymbol{\theta}, \boldsymbol{\kappa})$. Specifying a population model for the response, $p(y_i | \mathbf{x}_i, \boldsymbol{\theta})$, is commonly done. Going a step further, as we do, to incorporate the inclusion probabilities in the population model, is not commonly done. By contrast, the pseudo posterior approach, which is formally introduced in the sequel, considers the inclusions probabilities (or equivalently, the sampling weights) as fixed. Under an informative sampling design where the (π_i) are constructed to depend on the (y_i) , each time we generate new values of the response variable we are prompted to update the inclusion probabilities for all units in the population after fixing a sampling design. So we believe the treatment of π_i as random is natural as they are conditioned on the values of population variables.

The price the modeler pays for this fully Bayesian approach is that they are required to specify a conditional distribution of the inclusion probabilities for all units in the population, $p(\pi_i | y_i, \boldsymbol{\kappa})$. This requirement raises two computational difficulties: Firstly, even if the specified population likelihood, $p(y_i | \mathbf{x}_i, \boldsymbol{\theta})$, and the associated prior, $p(\boldsymbol{\theta})$, yield a closed form conditional posterior distribution, this construction for the population model does not produce a conjugate posterior under our formulation to correct for informative sampling (IS) because the sampling design is not ignorable; Secondly, the computation of the expected value in the denominator of Equation (5) is a computational bottleneck as we are required to compute this expected value for each observation in every Gibbs sampler iteration. We cope with the first difficulty by relying on the black box solver Stan [Carpenter et al., 2016], that uses a Hamiltonian Monte Carlo approach to draw samples from the full joint posterior distribution, so that it performs well under (is insensitive to) non-conjugacy. Regarding the second computational challenge, we proceed to construct a set of conditions on $p(\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa})$ and the likelihood that guarantee the availability of a closed form expression for this expected value.

We next specify a class of conditional population distributions from Equation (4) that yield a closed form result for the expectation step in denominator of (5), which simplifies posterior computation. Let \mathbf{v}_i and \mathbf{u}_i be sub-vectors of \mathbf{x}_i , the covariates used to specify the conditional distribution of $\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa}$ and $y_i | \mathbf{x}_i, \boldsymbol{\theta}$, respectively; that is, $\pi_i | y_i, \mathbf{x}_i, \boldsymbol{\kappa} \sim \pi_i | y_i, \mathbf{v}_i, \boldsymbol{\kappa}$ and $y_i | \mathbf{x}_i, \boldsymbol{\theta} \sim y_i | \mathbf{u}_i, \boldsymbol{\theta}$. Note that we allow for \mathbf{v}_i and \mathbf{u}_i to have common

covariates. Let $\text{normal}(x \mid \mu, s^2)$ denote the normal distribution pdf with mean μ and variance s^2 evaluated at x , and $\text{lognormal}(\cdot \mid \mu, s^2)$ denote the lognormal pdf, so that $X \sim \text{lognormal}(\mu, s^2)$ is equivalent to $\log X \sim \text{normal}(\mu, s^2)$. Going forward, we assume that π_i is proportional, as opposed to exactly equal, to the inclusion probability for unit i . In other words, no restriction is imposed on $\sum_i \pi_i$ where the index i could run over the population or sample indices. We will see in the sequel that normalizing the sum of the inverse inclusion probabilities *is* required to regulate the estimation of posterior uncertainty under the pseudo likelihood approach, but such is not required for our fully Bayesian formulation.

Theorem 1. *If $p(\pi_i \mid y_i, \mathbf{v}_i, \boldsymbol{\kappa}) = \text{lognormal}(\pi_i \mid h(y_i, \mathbf{v}_i, \boldsymbol{\kappa}), \sigma_{\pi^2})$, with the function $h(y_i, \mathbf{v}_i, \boldsymbol{\kappa})$ of the form $h(y_i, \mathbf{v}_i, \boldsymbol{\kappa}) = g(y_i, \mathbf{v}_i, \boldsymbol{\kappa}) + t(\mathbf{v}_i, \boldsymbol{\kappa})$ where $\sigma_{\pi^2} = \sigma_{\pi}^2(\boldsymbol{\kappa}, \mathbf{v}_i)$, possibly a function of $(\boldsymbol{\kappa}, \mathbf{v}_i)$ then*

$$p_s(y_i, \pi_i \mid \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) = \frac{\text{normal}(\log \pi_i \mid g(y_i, \mathbf{v}_i, \boldsymbol{\kappa}) + t(\mathbf{v}_i, \boldsymbol{\kappa}), \sigma_{\pi}^2)}{\exp\{t(\mathbf{v}_i, \boldsymbol{\kappa}) + \sigma_{\pi}^2/2\}} \times p(y_i \mid \mathbf{u}_i, \boldsymbol{\theta})$$

with $M_y(\boldsymbol{\kappa}; \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}) := E_{y \mid \mathbf{u}_i, \boldsymbol{\theta}}[\exp\{g(y_i, \mathbf{v}_i, \boldsymbol{\kappa})\}]$.

The proof of this theorem is in the Supplementary Material, Section 7. If both M_y and $p(y_i \mid \dots)$ admit closed form expressions, then $p_s(y_i, \pi_i \mid \dots)$ has a closed form, as well; for example, when $g(y_i, \mathbf{v}_i, \boldsymbol{\kappa}) = \kappa_y y_i$ with $\kappa_y \in \boldsymbol{\kappa} \in \mathbb{R}$, then $M_y(\boldsymbol{\kappa}; \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta})$ is the moment generating function (MGF) of $y_i \mid \boldsymbol{\theta}$ evaluated at κ_y , which will have a closed form defined on \mathbb{R} for typically-used normal and binomial population models for the responses. The closed form for $M_y(\boldsymbol{\kappa}; \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta})$ implies a closed form for $p_s(y_i, \pi_i \mid \dots)$. Analogously, we may consider an interaction between y and \mathbf{v} , using $g(y_i, \mathbf{v}_i, \boldsymbol{\kappa}) = (\kappa_y + \mathbf{v}_i^t \boldsymbol{\kappa}_{\mathbf{v}}) y_i \equiv t y_i$ with $\boldsymbol{\kappa} = (\kappa_y, \boldsymbol{\kappa}_{\mathbf{v}})$. In this case, we achieve, $M_y(t; \dots)$, which is the MGF evaluated at t . Our assumption of a lognormal distribution for π_i is mathematically appealing since the inclusion probability, π_i , for individual, i , is composed from the product of inclusion probabilities of selection across the stages of the multi-stage survey design. If each of these stagewise probabilities are lognormal then their product, π_i , is lognormal as well. We next apply Theorem 1 to some common settings.

2.1. Linear Regression Population Model

Assume the linear regression model for the population, $p(y_i \mid \mathbf{u}_i, \boldsymbol{\theta})$, is constructed as,

$$y_i \mid \mathbf{u}_i, \boldsymbol{\theta} \sim \text{normal}(\mathbf{u}_i^t \boldsymbol{\beta}, \sigma_y^2), \quad \text{with } \boldsymbol{\theta} = (\boldsymbol{\beta}, \sigma_y^2) \quad (6)$$

and the conditional population model for inclusion probabilities is specified as,

$$\pi_i \mid y_i, \mathbf{v}_i, \boldsymbol{\kappa} \sim \text{lognormal}\left(\kappa_y y_i + \mathbf{v}_i^t \boldsymbol{\kappa}_x, \sigma_{\pi}^2\right), \quad \text{with } \boldsymbol{\kappa} = (\kappa_y, \boldsymbol{\kappa}_x, \sigma_{\pi}^2) \quad (7)$$

This construction results from setting, $g(y_i, \mathbf{v}_i, \boldsymbol{\kappa}) = \kappa_y y_i$, $t(\mathbf{v}_i, \boldsymbol{\kappa}) = \mathbf{v}_i^t \boldsymbol{\kappa}_x$, $\sigma_{\pi}^2(\boldsymbol{\kappa}, \mathbf{v}_i) = \sigma_{\pi}^2$. Here $\boldsymbol{\beta}$ and $\boldsymbol{\kappa}_x$ are vectors of regression coefficients that include

an intercept, so the first entry of both \mathbf{u}_i and \mathbf{v}_i equals 1. We select prior distributions,

$$\begin{aligned}\boldsymbol{\beta} &\sim \text{MVN}(\mathbf{0}, 100\mathbf{I}) \\ \boldsymbol{\kappa} &\sim \text{MVN}(\mathbf{0}, 100\mathbf{I}) \\ \sigma_y^2, \sigma_\pi^2 &\stackrel{\text{iid}}{\sim} \text{normal}^+(0, 1)\end{aligned}\tag{8}$$

with $\text{normal}^+(m, s^2)$ denoting a normal distribution with mean m and variance s^2 restricted to the positive real line; $\text{MVN}(\mathbf{m}, \Sigma)$ denotes the multivariate normal distribution with mean vector \mathbf{m} and variance-covariance matrix Σ ; and \mathbf{I} the identity matrix.

Since $y \sim \text{normal}(m, s^2)$ admits a closed form expression for $M_y(t) = \exp(tm + t^2s^2/2)$, we apply Theorem 1 to obtain,

$$\begin{aligned}p_s(y_i, \pi_i \mid \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) &= \frac{\text{normal}(\log \pi \mid \kappa_y y_i + \mathbf{v}_i^t \boldsymbol{\kappa}_x, \sigma_\pi^2)}{\exp\{\mathbf{v}_i^t \boldsymbol{\kappa}_x + \sigma_\pi^2/2 + \kappa_y \mathbf{u}_i^t \boldsymbol{\beta} + \kappa_y^2 \sigma_y^2/2\}} \\ &\quad \times \text{normal}(y_i \mid \mathbf{u}_i^t \boldsymbol{\beta}, \sigma_y^2)\end{aligned}\tag{9}$$

The resulting form of the expression for $\log p_s(\dots)$ is provided in the Supplementary Material, Section 8.

2.2. Other Population Models

Since a closed-form moment generating function of $y_i \mid \mathbf{u}_i, \boldsymbol{\theta}$ defined in \mathbb{R} implies a closed form for $p_s(y_i, \pi_i \mid \dots)$, extensions for the logistic, probit and Poisson regressions to perform asymptotically unbiased estimations on the observed sample are straightforward; for example, employing the conditional lognormal model, Equation (7), for the inclusion probabilities, but now selecting a probit construction for a dichotomous, $y_i \sim \text{Bernoulli}(p_i)$, with $p_i = \Phi^{-1}(\mathbf{x}_i^t \boldsymbol{\theta})$, where Φ^{-1} is the inverse of the standard normal CDF. We note the closed form MGF, $E(e^{ty}) = (1 - p + pe^t)$ for all t , and obtain,

$$p_s(y_i, \pi_i \mid \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) = \frac{\text{normal}(\log \pi_i \mid \kappa_y y_i + \mathbf{v}_i^t \boldsymbol{\kappa}_x, \sigma_\pi^2)}{\exp\{\mathbf{v}_i^t \boldsymbol{\kappa}_x + \sigma_\pi^2/2\} (1 - p_i + p_i e^{\kappa_y})} \times \text{Bernoulli}(y \mid p_i)$$

with $\text{Bernoulli}(y \mid p_i) = p_i^{y_i} (1 - p_i)^{1-y_i}$.

2.3. Splines Basis Population Model

We extend the linear regression model to a splines setting for situations where the relationship between the response and the predictor is not linear. Here, $p(y_i \mid \mathbf{u}_i, \boldsymbol{\theta})$ is constructed as,

$$y_i \sim \text{normal}(\mu(\mathbf{u}_i), \sigma_y^2) \quad \text{with } \mu(\mathbf{u}_i) := B(\mathbf{u}_i)\boldsymbol{\beta}\tag{10}$$

with $B(\mathbf{u}_i) := (B_{i1}, \dots, B_{ib})$, the vector of penalized B-spline coefficients associated with \mathbf{u}_i , where b denotes the number of spline bases. We construct a rank-deficient multivariate Gaussian prior for the coefficients, $\boldsymbol{\beta}$, which penalizes or regulates the smoothness of the estimated function, μ , by employing a $b \times b$ matrix, $Q = D^t D$, where D is the discretized k th difference operator [Speckman and Sun, 2003] in the density kernel,

$$\sigma_{\boldsymbol{\beta}}^{-(b-k)/2} \exp -\frac{1}{2\sigma_{\boldsymbol{\beta}}^2} \boldsymbol{\beta}^t Q \boldsymbol{\beta}.$$

The order, k , regulates the resulting smoothness of the fitted function or its order of differentiability. $B(\mathbf{u}_i)$ has at most k entries different from zero. The penalized pdf specified for $\boldsymbol{\beta}$ is not proper since the dimension of $\boldsymbol{\beta}$ is b , but Q is of rank $b - k$. We observe that the MLE for $(\boldsymbol{\beta}, \sigma_y^2, \sigma_{\boldsymbol{\beta}}^2)$ is a type of ridge regression estimator since it satisfies,

$$(B^t B / \hat{\sigma}_y^2 + Q / \hat{\sigma}_{\boldsymbol{\beta}}^2) \hat{\boldsymbol{\beta}} = B^t \mathbf{y} / \hat{\sigma}_y^2.$$

Our simulation scenarios in Section 4 will utilize this penalized B-spline formulation with $(b = 8, k = 4)$. The priors for model variances are set as weakly informative proper priors,

$$\sigma_y \sim \text{normal}^+(0, 10), \quad \sigma_{\boldsymbol{\beta}} \sim \text{Cauchy}^+(0, 10) \quad \text{and} \quad \sigma_{\pi} \sim \text{Cauchy}(0, 1);$$

where $\text{Cauchy}^+(m, v)$ denotes the Cauchy distribution with location and scale parameters m and v , respectively, restricted to be positive. $\tau_{\boldsymbol{\beta}} = 1/\sigma_{\boldsymbol{\beta}}^2$ is referred to as a complexity parameter.

Since the spline model is a particular case of the linear regression model in Subsection 2.1, $p_s(y_i, \pi_i \mid \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}, \boldsymbol{\kappa})$, with $\boldsymbol{\theta} := (\boldsymbol{\beta}, \sigma_y^2)$, for the spline model has the same expression as that for the linear model in Equation (9) after replacing $\mathbf{u}_i^T \boldsymbol{\beta}$ for $B_i \boldsymbol{\beta}$ (assuming our continued employment of a lognormal prior for π_i in Equation (7)).

2.4. Estimation of Population Distribution of Inclusion Probabilities

For a simple one-stage design the whole population of inclusion probabilities or equivalent is available, because their values are required to draw a sample. For a multistage cluster sample like a survey of individuals within households, we draw a sample recursively, only specifying inclusion probabilities for units selected at the previous level. This procedure provides sampling probabilities only for those lowest-level units or individuals that are selected into the sample. It may be useful for the survey administrator to estimate the population distribution of inclusion probabilities from the observed sample in order to verify a subgroup of units targeted for over-sampling do, indeed, express relatively higher inclusion probabilities.




Fig 1: Histogram of the $n = 100$ log-transformed *sampled* inclusion probabilities along with the true simulated standardized normal population density (solid line) and estimated normalized population density (dashed line) with mean and variance estimated using Equation (11).

We select a model that satisfies the conditions of Theorem 1, but set $y_i \equiv 0$. This is, $g(y_i, \mathbf{v}_i, \boldsymbol{\kappa})$ is constant and without loss of generality equal to zero. Then, $M_y(\boldsymbol{\kappa}; \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}) = 1$ and Theorem 1 implies

$$p_s(\pi_i | \mathbf{v}_i, \boldsymbol{\kappa}) = \frac{\text{normal}(\log \pi_i | t(\mathbf{v}_i, \boldsymbol{\kappa}), \sigma_\pi^2)}{\exp\{t(\mathbf{v}_i, \boldsymbol{\kappa}) + \sigma_\pi^2/2\}} \quad (11)$$

If we assume $\pi_i \sim \text{lognormal}(t(\mathbf{v}_i, \boldsymbol{\kappa}), \sigma_\pi^2)$ for $i = 1, \dots, N$ with $\Pr(\delta_i = 1) = \pi_i / \sum_{i=1}^N \pi_i$, we can make inference about the parameters $\boldsymbol{\kappa}$ and σ_π^2 using only the observed inclusion probabilities. As an example we simulate $\pi_1, \dots, \pi_{10^5} \stackrel{\text{iid}}{\sim} \text{lognormal}(\boldsymbol{\kappa} = 0, \sigma_\pi^2 = 1)$. (Our choice of the lognormal distribution implies $t(\mathbf{v}_i, \boldsymbol{\kappa}) \equiv \boldsymbol{\kappa}$.) We draw $n = 100$ inclusion probabilities (π_i) under IS, and estimate the parameter values $\boldsymbol{\kappa}$ and σ_π^2 via Equation (11) using the priors outlined in Equation (8). Figure 1 shows the histogram of the log transformed sampled inclusion probabilities observed in our sample of $n = 100$, along with the standardized density of generated inclusion probabilities for our simulated population, displayed in the solid line, and our estimated normal density with population mean $E(\boldsymbol{\kappa} | \pi_1, \dots, \pi_n)$ and variance $E(\sigma_\pi^2 | \pi_1, \dots, \pi_n)$, displayed in the dashed line. Notice that we employ the unstandardized sampling weights. If the sampling weights are multiplied by a constant $c > 0$; e.g., $c := n / \sum_{i=1}^n \pi_i$, we are then estimating $\boldsymbol{\kappa} + \log c$.

2.5. Pseudolikelihood Alternative to the Fully Bayes Estimator

The pseudo posterior approach is a plug-in method that is not fully Bayesian, which we will use to compare with the fully Bayes method we propose in this paper. The pseudo likelihood is formed by exponentiating the likelihood contribution for each observed unit in the sample by its sampling weight, $w_i^{(s)}$, in,

$$\text{pseudo}(\boldsymbol{\theta}; \mathbf{y}^{(s)}, \mathbf{w}^{(s)}) = \prod_{i=1}^n p(y_i^{(s)} | \mathbf{x}^{(s)}_i, \boldsymbol{\theta})^{w_i^{(s)}}$$

where $w_i^{(s)} \propto 1/\pi_i^{(s)}$ is the sampling weight that is standardized such that $\sum w_i^{(s)} = n$, where n denotes to the observed sample size, which is required to regulate the amount of estimated posterior uncertainty. Let $\mathbf{w}^{(s)} := (w_1^{(s)}, \dots, w_n^{(s)})$ denote the vector of these unit-indexed sampling weights that are typically published with the survey data in order to correct for IS. Inference is carried out by constructing the pseudo posterior distribution be proportional to $\text{pseudo}(\boldsymbol{\theta}; \mathbf{y}^{(s)}, \mathbf{w}^{(s)}) \times \text{Prior}(\boldsymbol{\theta})$ [Savitsky and Toth, 2016]. Although the pseudo likelihood is an improper distribution, it admits a proper joint posterior distribution under employment of proper priors. Classical implementations of the sampling weighted pseudo likelihood, by contrast to the Bayesian implementation, employ MLE estimation of population model parameters on the observed IS. The standard error of the estimate of parameter, θ , is estimated via resampling methods such as balanced repeated replication [McCarthy, 1969, Krewski and Rao, 1981] or Fray’s method [Judkins, 1990] when the sampling design and the information provided to the analyst allow it.

The advantages of the plug-in pseudo posterior method over the proposed fully Bayesian construction are: (i) Estimation does not require a custom posterior sampler. Relatively slight modifications are performed to the population model sampler to incorporate the pseudo likelihood; (ii) Specification of $\pi_i | y_i, \dots$ for the population is not required; (iii) There is no expected value in Equation (3) to compute as there is in the fully Bayes method. The disadvantages of the pseudo posterior approach are: (i) It is not fully Bayesian; (ii) It does not discard variation in weights that do not depend on the response, such that noise (unrelated to the response) is introduced into the estimation of the pseudo posterior distribution; (iii) The weights must be normalized to regulate the amount of estimated posterior uncertainty, which is not required for the fully Bayes approach. We will show in the sequel that the fully Bayes approach expresses superior performance in estimation of posterior uncertainty than does the pseudo posterior.

3. Posterior Consistency

We proceed to construct conditions on the population generating model and sample inclusion probabilities that guarantee the contraction of our fully Bayes estimator to the true joint posterior distribution. The conditions constructed for

the sample inclusion probabilities define a class of sampling designs for which we would expect asymptotically unbiased estimation of our population model parameters on the observed data.

3.1. Set-up

Let $\nu \in \mathbb{Z}^+$ index a sequence of finite populations, $\{U_\nu\}_{\nu=1, \dots, N_\nu}$, each of size, $|U_\nu| = N_\nu$, such that $N_\nu < N_{\nu'}$, for $\nu < \nu'$, so that the finite population size grows as ν increases. We employ subscript, ν , because the process of rendering a sample involves the generation of a population followed by the assignment of sample inclusion probabilities and the taking of a sample from that population using the sampling weights. The entire sequence is repeated on each increment of ν . A sampling design is defined by placing a distribution on a vector of inclusion indicators, $\boldsymbol{\delta}_\nu = (\delta_{\nu 1}, \dots, \delta_{\nu N_\nu})$, linked to the units comprising the population, U_ν , that we use to draw a sample of size $n_\nu < N_\nu$. We construct a sampling distribution by specifying marginal inclusion probabilities, $\pi_{\nu i} = \Pr\{\delta_{\nu i} = 1 | Y_{\nu, i} = y_{\nu i}\}$ for all $i \in U_\nu$ and the second-order pairwise probabilities, $\pi_{\nu ij} = \Pr\{\delta_{\nu i} = 1 \cap \delta_{\nu j} = 1 | Y_{\nu, i} = y_{\nu i}, Y_{\nu, j} = y_{\nu j}\}$ for $i, j \in U_\nu$, where $Y_{\nu, 1}, \dots, Y_{\nu, N_\nu}$ index random variables that are independently distributed according to some unknown distribution P_θ (with density, p_θ) defined on the sample space, $(\mathcal{Y}_\nu, \mathcal{A}_\nu)$.

In the usual survey sampling set-up, the inclusion probabilities are treated as *fixed*. Under the Bayesian paradigm, however, Y_ν is random variable drawn from a super-population, such that if a new population, U_ν , with associated observed values $(y_{\nu i})$, is drawn, the marginal inclusion probabilities, $(\pi_{\nu i})$, will update. So we condition the inclusion probabilities explicitly on the population response values for emphasis, unlike in Savitsky and Toth [2016]. We envision both inclusion probabilities, $(\pi_{\nu i})$, and our data, $(y_{\nu, i})$, as generated from a super-population under some true model. We construct the joint density of the model for $(y_{\nu i}, \pi_{\nu i})$ for each $i = 1, \dots, N_\nu$ as,

$$p_\lambda(y_{\nu i}, \pi_{\nu i}) = p_\theta(y_{\nu i}) \times p_\kappa(\pi_{\nu i} | Y_{\nu i} = y_{\nu i}), \quad (12)$$

where $\lambda = (\theta, \kappa)$ denotes our model parameters.

We use the Bayes rule construction we earlier illustrated to formulate a sampling design-adjusted fully Bayesian joint likelihood for the unknown population value, λ_0 , which we suppose generates the finite population values that we ob-

serve in our sample, $(y_{\nu,i}, \pi_{\nu,i})_{i \in U_\nu: \delta_{\nu,i}=1}$,

$$p_\lambda^\pi(x_{\nu i} \delta_{\nu i}) \quad (13)$$

$$:= \left[\frac{\pi_{\nu i} \times p_\lambda(y_{\nu i}, \pi_{\nu i})}{\mathbb{E}_\theta \{ \Pr_\kappa(\delta_{\nu i} = 1 | Y_{\nu i} = y_{\nu i}) \}} \right]^{\delta_{\nu i}} \quad (14)$$

$$:= \left[\frac{\pi_{\nu i}}{\mathbb{E}_\theta(\pi_{\nu i}^\kappa)} \times p_\lambda(y_{\nu i}, \pi_{\nu i}) \right]^{\delta_{\nu i}} \quad (15)$$

$$:= \left[\frac{\pi_{\nu i}}{\pi_{\nu i}^\lambda} p_\lambda(y_{\nu i}, \pi_{\nu i}) \right]^{\delta_{\nu i}}, \quad i \in U_\nu, \quad (16)$$

where we collect, $x_{\nu i} = (y_{\nu i}, \pi_{\nu i})$,

and define $\pi_{\nu i}^\kappa := \Pr_\kappa(\delta_{\nu i} = 1 | Y_{\nu i} = y_{\nu i}) = \mathbb{E}_\kappa(\pi_{\nu i} | Y_{\nu i} = y_{\nu i})$, where the second equality is derived in [Pfeffermann et al. \[1998\]](#). We further define $\pi_{\nu i}^\lambda := \mathbb{E}_\theta(\pi_{\nu i}^\kappa) = \mathbb{E}_\theta \{ \mathbb{E}_\kappa(\pi_{\nu i} | Y_{\nu i} = y_{\nu i}) \}$ for ease-of-reading.

We conduct Bayesian inference under the likelihood of Equation (16) by assigning prior, Π , on the parameter space, Λ , such that $\lambda_0 \in \Lambda$, which produces the sampling design-adjusted posterior mass,

$$\Pi^\pi(B | x_{\nu 1} \delta_{\nu 1}, \dots, x_{\nu N_\nu} \delta_{\nu N_\nu}) = \frac{\int_{\lambda \in B} \prod_{i=1}^{N_\nu} \frac{p_\lambda^\pi(x_{\nu i} \delta_{\nu i})}{p_{\lambda_0}^\pi(x_{\nu i} \delta_{\nu i})} d\Pi(\lambda)}{\int_{\lambda \in \Lambda} \prod_{i=1}^{N_\nu} \frac{p_\lambda^\pi(x_{\nu i} \delta_{\nu i})}{p_{\lambda_0}^\pi(x_{\nu i} \delta_{\nu i})} d\Pi(\lambda)}, \quad (17)$$

that we use to formulate our theoretical result. We define $p_{\lambda_0}^\pi(x_{\nu i} \delta_{\nu i}) = p_{\lambda_0}^\pi(x_{\nu i})^{\delta_{\nu i}}$, that confines evaluation to the observed sample.

We will utilize the empirical distribution construction to establish a bound used to prove our result,

$$\mathbb{P}_{N_\nu}^\pi = \frac{1}{N_\nu} \sum_{i=1}^{N_\nu} \frac{\delta_{\nu i}}{\pi_{\nu i}^\lambda} \delta(X_i), \quad (18)$$

which is nearly identical to [Savitsky and Toth \[2016\]](#), only now the denominator is a modeled quantity dependent on λ . We construct the associated expectation functional, $\mathbb{P}_{N_\nu}^\pi f = \frac{1}{N_\nu} \sum_{i=1}^{N_\nu} \frac{\delta_{\nu i}}{\pi_{\nu i}^\lambda} f(X_i)$. We make a similar adjustment to the Hellinger distance,

$$d_{N_\nu}^{\pi,2}(p_{\lambda_1}, p_{\lambda_2}) = \frac{1}{N_\nu} \sum_{i=1}^{N_\nu} \frac{\delta_{\nu i}}{\pi_{\nu i}^\lambda} d^2(p_{\lambda_1}(x_{\nu i}), p_{\lambda_2}(x_{\nu i})),$$

where $d(p_{\lambda_1}, p_{\lambda_2}) = \left[\int (\sqrt{p_{\lambda_1}} - \sqrt{p_{\lambda_2}})^2 d\mu \right]^{\frac{1}{2}}$ (for dominating measure, μ).

3.2. Main Result

We outline the 6 conditions and our result for completeness, though they are very similar to [Savitsky and Toth \[2016\]](#) save for the replacement of $\pi_{\nu i}$ with the modeled $\pi_{\nu i}^\lambda$. The first 3 conditions place restrictions on the generating distribution and prior, while the following three do the same for the sampling design distribution. Suppose we have a sequence, $\xi_{N_\nu} \downarrow 0$ and $N_\nu \xi_{N_\nu}^2 \uparrow \infty$ and $n_\nu \xi_{N_\nu}^2 \uparrow \infty$ as $\nu \in \mathbb{Z}^+ \uparrow \infty$ and any constant, $C > 0$,

(A1) (Local entropy condition - Size of model)

$$\sup_{\xi > \xi_{N_\nu}} \log N(\xi/36, \{\lambda \in \Lambda_{N_\nu} : d_{N_\nu}(P_\lambda, P_{\lambda_0}) < \xi\}, d_{N_\nu}) \leq N_\nu \xi_{N_\nu}^2,$$

(A2) (Size of space)

$$\Pi(\Lambda \setminus \Lambda_{N_\nu}) \leq \exp(-N_\nu \xi_{N_\nu}^2 (2(1 + 2C)))$$

(A3) (Prior mass covering the truth)

$$\Pi\left(P_\lambda : \left(-\mathbb{E}_{\lambda_0} \log \frac{p_\lambda}{p_{\lambda_0}} \leq \xi_{N_\nu}^2\right) \cap \left(\mathbb{E}_{\lambda_0} \left[\log \frac{p_\lambda}{p_{\lambda_0}}\right]^2 \leq \xi_{N_\nu}^2\right)\right) \geq \exp(-N_\nu \xi_{N_\nu}^2 C)$$

(A4) (Non-zero Inclusion Probabilities)

$$\sup_\nu \left[\frac{1}{\min_{i \in U_\nu} \pi_{\nu i}} \right] \leq \gamma, \text{ with } P_{\lambda_0}\text{-probability 1.}$$

(A5) (Asymptotic Independence Condition)

$$\limsup_{\nu \uparrow \infty} \max_{i \neq j \in U_\nu} \left| \frac{\pi_{\nu ij}}{\pi_{\nu i} \pi_{\nu j}} - 1 \right| = \mathcal{O}(N_\nu^{-1}), \text{ with } P_0\text{-probability 1}$$

such that $\pi_{\nu ij}$ factors to $\pi_{\nu i} \pi_{\nu j}$ for N_ν sufficiently large where there exists some $C_3 > 0$,

$$N_\nu \sup_\nu \max_{i \neq j \in U_\nu} \left[\frac{\pi_{\nu ij}}{\pi_{\nu i} \pi_{\nu j}} - 1 \right] \leq C_3.$$

(A6) (Constant Sampling fraction) For some constant, $f \in (0, 1)$, that we term the “sampling fraction”,

$$\limsup_\nu \left| \frac{n_\nu}{N_\nu} - f \right| = \mathcal{O}(1), \text{ with } P_0\text{-probability 1.}$$

(A7) (Convergence of the Point Estimate)

$$\limsup_{\nu \uparrow \infty} \max_{i \in U_\nu} |\mathbb{E}_\lambda[\pi_{\nu i}] - \mathbb{E}_{\lambda_0}[\pi_{\nu i}]| = \mathcal{O}(N_\nu^{-1}), \text{ with } P_0\text{-probability 1.}$$

Condition (A1) restricts the growth in the size of the model space [Ghosal et al., 2000] by bounding the growth in the logarithm of the covering number, $N(\xi/36, \{\lambda \in \Lambda_{N_\nu} : d_{N_\nu}(P_\lambda, P_{\lambda_0}) < \xi\})$, defined as the *minimum* number of balls of radius $\xi/36$ needed to cover $\{P \in \mathcal{P}_{N_\nu} : d_{N_\nu}(P, P_0) < \xi\}$ under distance metric, d_{N_ν} . Condition (A2) allows, but restricts, the prior mass placed on the uncountable portion of the model space, such that we may direct our inference to an approximating sieve, \mathcal{P}_{N_ν} . This sequence of spaces “trims” away a portion of the space that is not entropy bounded (in condition (A1)). Condition (A3) ensures the prior, Π , assigns mass to convex balls in the vicinity of P_{λ_0} .

The next three conditions, together, restrict the class of sampling designs under which our result is guaranteed. Condition (A4) requires the sampling design to assign a positive probability for inclusion of every unit in the population because the restriction bounds the sampling inclusion probabilities away from 0. Since the maximum inclusion probability is 1, the bound, $\gamma \geq 1$. Unlike in Savitsky and Toth [2016], however, $\pi_{\nu i}^\lambda$ is a smoothed model estimator under our fully Bayes construction, which discards the variation in $\pi_{\nu i}$ that is unrelated to $y_{\nu i}$, such that the value of γ , in practice, is expected to be lower than in Savitsky and Toth [2016]. Condition (A5) restricts the result to sampling designs where the dependence among lowest-level sampled units attenuates to 0 as $\nu \uparrow \infty$. Dependence in a multistage design is driven the higher level sampling stages; for example, PSUs. Since the number of PSUs increases in the limit of N_ν , this condition is not very restrictive and admits nearly all sampling designs used, in practice. Condition (A6) ensures that the observed sample size, n_ν , limits to ∞ along with the size of the partially-observed finite population, N_ν . The denominator of our Bayes rule posterior estimator of Equation (16) is a conditional expectation with respect to our model, P_λ . We require convergence of this point estimate in order to achieve the bound specified in condition (A5). Condition (A7) is not needed in Savitsky and Toth [2016], since the inclusion probabilities, (π_ν) , are assumed fixed.

Theorem 2. *Suppose conditions (A1)-(A7) hold. Then for sets $\Lambda_{N_\nu} \subset \Lambda$, constants, $K > 0$, and M sufficiently large,*

$$\begin{aligned} \mathbb{E}_{P_{\lambda_0}, P_\nu} \Pi^\pi (\lambda : d_{N_\nu}^\pi (P_\lambda, P_{\lambda_0}) \geq M \xi_{N_\nu} | x_{\nu 1} \delta_{\nu 1}, \dots, x_{\nu N_\nu} \delta_{\nu N_\nu}) \leq \\ \frac{16\gamma^2 [\gamma + C_3]}{(Kf + 1 - 2\gamma)^2 N_\nu \xi_{N_\nu}^2} + 5\gamma^2 \exp\left(-\frac{Kn_\nu \xi_{N_\nu}^2}{2\gamma}\right), \end{aligned} \quad (19)$$

which tends to 0 as $(n_\nu, N_\nu) \uparrow \infty$.

The rate of convergence is injured for a sampling distribution, P_ν , that assigns relatively low inclusion probabilities to some units in the finite population such that γ will be relatively larger. Our result differs from Savitsky and Toth [2016] in that 5γ that multiplies the second term is replaced by $5\gamma^2$, here, though the value γ is expected to be lower, as earlier discussed, for our modeled estimate, $\pi_{\nu i}^\lambda$, than for the raw, $\pi_{\nu i}$. Similarly, the larger the dependence among the finite population unit inclusions induced by P_ν , the higher will be C_3 and the slower will be the rate of contraction.

The proof of our main result proceeds by bounding the numerator of Equation (17) (on the set $\{\lambda : d_{N_\nu}^\pi (P_\lambda, P_{\lambda_0}) \geq M \xi_{N_\nu}\}$), from above, and the denominator, from below, and is the same as that outlined in Savitsky and Toth [2016]. So we do not repeat it here. The proof relies two enabling results, which express some differences in this work, however, from Savitsky and Toth [2016]. So those Lemma are reformulated and proved in Supplementary Material section 8.1.

4. Simulation Study

We explore the bias and coverage performance of our fully Bayes formulation and compare it to the plug-in pseudo posterior model under employment of an informative, probability proportional-to-size (PPS) sampling design; first in a simple linear regression and later in a nonlinear, spline model setting. We construct a synthetic population in each iteration of our Monte Carlo simulation, from which we draw two samples: 1. An informative sample (IS) taken under the PPS design; 2. A simple random sample (SRS). The IS is analyzed using both the fully Bayes and the pseudo posterior methods. The SRS is estimated using our population model with no needed correction. The SRS serves as a something of a gold standard comparator to assess the efficacy of bias removal and uncertainty estimations using the methods that correct for informative sampling, which are estimated on the IS. Of course, an informative sampling design can be more efficient (lower variance) than an SRS, though an IS is often constructed as much for convenience (and cost) as for efficiency. We estimate bias, mean square error (MSE), coverage of central 95% credible interval (CI) and its average length by repeating this process $M = 1,000$ times in a Monte Carlo framework. The three analysis methods (that we label, “Full” (Fully Bayesian), “Pseudo” (Pseudo Posterior) and “SRS” (simple random sampling)) will use the same population distribution for $y_i | \theta, \dots$ and prior for θ . The true value of the generic simulation model parameter, η , is denoted by η^{TRUE} .

4.1. Simple Linear Regression (SLR): PPS Design

We focus on estimating the population generating for the slope coefficient, β_1 , specified for the linear regression framework in Section 2.1. The fully Bayes estimation model assumes a lognormal conditional likelihood for $\pi_i | y_i, \dots$, which is right skewed. We explore three simulation scenarios; In the first two, the true generating distribution of the inclusion probabilities is a gamma distribution and, therefore, skewed as in the the fully Bayes analysis model. In the third simulation scenario, we explore the robustness of the fully Bayes model to this assumption by generating the inclusion probabilities from a (symmetric) normal distribution. The difference between the first and second scenarios is that we vary the rate hyperparameter of the gamma distribution to induce more variance (and skewness) into the generated (π_i) in the second scenario.

4.1.1. Scenarios SLR: π -skewed.

We call the first two simulation scenarios “SLR: π -skewed with low variance” and “SLR: π -skewed with high variance”. Both set $\pi_i \stackrel{\text{iid}}{\sim} \text{gamma}(2, \text{rate} = b_\pi^{\text{TRUE}})$ (See 1 (b) below) where $b_\pi^{\text{TRUE}} = 2$, in the former, and $b_\pi^{\text{TRUE}} = 1$, in the latter. We set $M = 10^3$ to be the total number of synthetic population datasets or Monte Carlo iterations, and $N = 10^5$ to be the number of individuals / units in the population, and we set $n = 10^3$ to be the sample size.

Let $\boldsymbol{\theta}^{\text{TRUE}} = (\beta_0^{\text{TRUE}}, \beta_1^{\text{TRUE}}, \beta_2^{\text{TRUE}}, \sigma_y^{2,\text{TRUE}}) = (0, 1, 1, 0.1^2)$ under the following simulation procedure:

- For $m = 1, \dots, M$, Monte Carlo iterations
 1. Generate the population, *i.e.*, for $i = 1, \dots, N$,
 - (a) Draw $\mathbf{u}_i \stackrel{\text{iid}}{\sim} \text{uniform}(0, 1)$
 - (b) Draw $\pi_i \stackrel{\text{iid}}{\sim} \text{gamma}(2, \text{rate} = b_\pi^{\text{TRUE}})$ (which produces unnormalized π_i).
 - (c) Generate the population response:

$$y_i \mid \mathbf{u}_i, \pi_i, \boldsymbol{\theta}^{\text{TRUE}} \sim \text{normal}(\beta_1^{\text{TRUE}} \mathbf{u}_i + \beta_2^{\text{TRUE}} \pi_i, \sigma_y^{2,\text{TRUE}})$$

2. Draw two samples of size $n = 10^3$:
 - (a) Take an IS with, $\Pr[(y_i, \pi_i) \in \text{sample}] = \pi_i / \sum_{i'=1}^N \pi_{i'}$.
 - (b) Take an SRS sample.
 3. Conduct estimation using the priors given in Equation (8).
 - (a) The informative sample is estimated using,
 - i. the fully Bayes linear regression approach outlined in Subsection 2.1 with parameters, $\boldsymbol{\theta} = (\beta_0, \beta_1, \sigma_y^2)$, $\boldsymbol{\kappa} = (\kappa_y, \sigma_\pi^2)$, where we note there is no \mathbf{v} or $\boldsymbol{\kappa}_x$.
 - ii. the pseudo posterior linear regression formulation.
 - (b) The SRS is estimated using the linear regression population model of Equation (6).
 4. (a) Store the posterior expected values of β_1 under the analyses 3(a) i, 3(a) ii and 3(b).
 - (b) Also compute the central 95% credible interval (CI) for β_1 under the three analyses and store their lengths and indicator of whether they contain β_1^{TRUE} .
- Using the values stored in step 4, (a) estimate the Bias and MSE of the estimate of β_1^{TRUE} and, (b), estimate the coverage and average length of the central 95% credible interval (CI) for β_1^{TRUE} .

Note that the synthetic data generating likelihood, $\pi_i \mid y_i, \mathbf{v}_i, \boldsymbol{\kappa}$, is *not* the lognormal distribution for the population estimation model; that is, the fully Bayes (estimation) model is misspecified. Also note that the fully Bayes population model in (6) is misspecified under the three analyses. More specifically, the expected value $y_i \mid \mathbf{u}_i, \boldsymbol{\theta}$, after integrating out π_i , matches that corresponding to a simple linear regression model with intercept $\beta_0^{\text{TRUE}} + E(\pi_i) = 0 + 2/b_\pi^{\text{TRUE}}$, slope coefficient $\beta_1^{\text{TRUE}} = 1$, but the error, $y_i - E(y_i \mid \mathbf{u}_i, \boldsymbol{\theta})$, though having mean zero and constant variance, $\text{var}(\pi_i) + \sigma_y^{2,\text{TRUE}} = 2/(b_\pi^{\text{TRUE}})^2 + 0.1^2$, is *not* normally distributed.




Fig 2: Estimation of β_1^{TRUE} under simulation scenario, SLR: π skewed with low variance. Here, $\pi_i \stackrel{\text{iid}}{\sim} \text{gamma}(2, b_\pi^{\text{TRUE}} = 2)$ in step 1(a) in Section 4.1.1. Box plots of posterior expected values of β_1 are displayed under IS for the fully Bayes and pseudo posterior approaches and under SRS. The horizontal dashed line represents simulation value $\beta_1^{\text{TRUE}} = 1$.

Figure 2 displays boxplots of the M posterior expected values of β_1 under each analysis under scenario SLR: π -skewed with low variance. Table 1 presents the bias, MSE, coverage of central 95% CI and its average length under both scenarios. The figure and the left-hand columns 2 to 4 of the table show that when the true variance of π is relatively *low* the pseudo posterior yields a good point estimator of β_1 with MSE lower even than under SRS. There is a trade-off of some bias for improved efficiency. The right-hand columns of the table, reporting the results under scenario SLR: π -skewed with relatively *high* variance, show that this property does not hold when the variance of π is large; the MSE for the pseudo posterior is now the highest, as is the bias, so there is no trade-off of one for the other. In other words, the point estimate of the fully Bayes model is robust against high variability of the inclusion probabilities while pseudo posterior estimator is not. In both scenarios, the average length of the CI is notably shorter under SRS than for the other two approaches under an IS because there is more variation in the realized samples drawn under an informative design, P_ν , than under SRS. The pseudo posterior underestimates the uncertainty (of the point estimate for β_1); its CI fails to capture the nominal 95% coverage. In contrast, the fully Bayes model estimates the uncertainty appropriately; its CI maintains a coverage similar to the model under SRS.

True variance of π_i	Low ($b_{\pi}^{\text{TRUE}} = 2$)			High ($b_{\pi}^{\text{TRUE}} = 1$)		
	Full	Pseudo	SRS	Full	Pseudo	SRS
Bias	-0.003	0.013	-0.002	-0.007	1.031	0.005
MSE	0.219	0.109	0.183	0.509	1.278	0.374
95% CI coverage	0.944	0.861	0.947	0.926	0.839	0.933
95% CI length	1.770	1.663	0.952	2.472	2.227	2.212

Table 1: Bias, MSE, coverage of central 95% CI and its average length with three analyses under simulation scenarios SLR: π -skewed with low variance and with high variance.

4.1.2. Scenario SLR: π -symmetric.

We explore the robustness of our approach in the case where the inclusion probabilities are generated as symmetric but modeled as skewed. To do so, we repeat the simulation study in subsection 4.1.1, but now with $\pi_i \stackrel{\text{iid}}{\sim} \text{normal}^+(1, \sigma_{\pi}^{2, \text{TRUE}} = 0.1^2)$ in 1 (b). We call this simulation scenario: “SLR: π -symmetric”. Notice that the fully Bayes approach, as in simulation scenario scenario SLR: π -skewed, misspecifies the distribution of $\pi_i | y_i, \boldsymbol{\kappa}$. In contrast to scenario SLR: π -skewed, the regression model for $y_i | \mathbf{u}_i, \boldsymbol{\theta}$ is correctly specified; in particular the distribution of the error is normally distributed.

Figure 3 and Table 2 are constructed in the same formats as Figure 2 and Table 1, for scenario: SLR: π -symmetric. The three methods perform similarly in terms of bias, MSE and coverage, even though, by design, the population model for the inclusion probabilities is misspecified under the fully Bayes method because we have drawn the inclusion probabilities from a symmetric distribution and modeled them with a skewed distribution. As in the previous scenario, SLR: π -skewed, the average length of the CI is notably shorter under SRS than under the other two approaches. That is, both, the fully Bayes and the pseudo posterior approaches maintain nominal coverage at the cost of wider credible intervals.




Fig 3: Estimation of β_1 under simulation scenario, SLR π -symmetric. Box plots of posterior expected values of β_1 are displayed under IS for the fully Bayes and pseudo posterior approaches and under SRS. The horizontal, dashed line represents the simulation value of $\beta_1^{\text{TRUE}} = 1$.

	Full	Pseudo	SRS
Bias	0.013	-0.011	0.002
MSE	0.123	0.042	0.115
95% CI coverage	0.957	0.948	0.950
95% CI length	1.373	1.370	0.795

Table 2: Bias, MSE, coverage of central 95% CI and its average length with three analyses under simulation scenario SLR: π -symmetric.

4.1.3. Scenario non-linear: π -skewed.

We conclude our simulation studies by exploring the situation where the relationship between the predictor and the response is *not* linear. As before, the superindex TRUE indicates the simulation model parameters. To ease the computational burden, we simulate only one synthetic population dataset with $N = 10^5$ individuals. We run this simulation twice, first with sample size $n = 100$ and last with $n = 1000$. The objective is to estimate the curve $E(y | \mathbf{u})$ on a regular grid, $\mathbf{u} \in (0, 2)$. The analysis model is the splines regression model described in Subsection 2.3.

- Generate the population (once)
 1. Draw $\mathbf{u}_i \stackrel{\text{iid}}{\sim} \text{uniform}(0, 2)$
 2. Draw $\pi_i \stackrel{\text{iid}}{\sim} \text{gamma}(2, \text{rate} = 1)$

3. Generate the population response:

$$y_i \mid \mathbf{u}_i, \pi_i, (\beta_{\mathbf{u}}^{\text{TRUE}}, \beta_{\pi}^{\text{TRUE}}, \beta_{\mathbf{u},\mathbf{u}}^{\text{TRUE}}, \sigma_y^{2,\text{TRUE}}) \sim \text{normal}(\beta_{\mathbf{u}}^{\text{TRUE}}\mathbf{u}_i + \beta_{\pi}^{\text{TRUE}}\pi_i + \beta_{\mathbf{u},\mathbf{u}}^{\text{TRUE}}\mathbf{u}_i^2, \sigma_y^{2,\text{TRUE}}),$$

$$\text{with } (\beta_{\mathbf{u}}^{\text{TRUE}}, \beta_{\pi}^{\text{TRUE}}, \beta_{\mathbf{u},\mathbf{u}}^{\text{TRUE}}, \sigma_y^{2,\text{TRUE}}) = (1, 1, -0.5, 0.1^2)$$

- For $m = 1, \dots, M$:
 1. Generate an IS and SRS of size n in the same fashion as in the SLR simulation study and, subsequently, estimate the model described in Section 2.3, using cubic B-splines, with the first and last internal knots equal to, respectively, the minimum and the maximum of the \mathbf{u}_i s in the sample.
 2. Compute the central 95% credible intervals for $E(y \mid \mathbf{u}) = B(\mathbf{u})E(\boldsymbol{\beta} \mid \mathbf{y}, \boldsymbol{\pi})$, pointwise on a grid, *i.e.*, $\mathbf{u} = 0, 1/40, 2/40, \dots, 80/40$, under the three analyses and store their credible interval lengths and an indicator for whether or not they contain the true curve given by

$$\beta_{\mathbf{u}}^{\text{TRUE}}\mathbf{u} + \beta_{\pi}^{\text{TRUE}}E(\pi_i) + \beta_{\mathbf{u},\mathbf{u}}^{\text{TRUE}}\mathbf{u}^2 = \mathbf{u} + 2 - 0.5\mathbf{u}^2$$

- Based on the values stored in step 3 (a), estimate the (pointwise) Bias and MSE curves and based on the values stored in step 3 (b), estimate the coverage and average length of the central 95% CI for the curve $E(y \mid \mathbf{u})$ with $\mathbf{u} \in (0, 2)$.

Figure 4 shows the results of the simulation under this scenario with sample size, $n = 100$. The first plot panel (from the top) displays the simulation curve and those estimated under the three methods. The next four panels (from top-to-bottom) display the MSE, Bias, coverage of central 95% CI and the average length of the CI for the three methods, respectively. The SRS method performs uniformly the best in terms of fit, bias, MSE and coverage. It is followed by the fully Bayes approach. As in scenario SLR: π -skewed the pseudo posterior approach fails to maintain a 95% coverage for its central 95% CI. The bottom panel shows that the cost of having an IS is a wider CI for the same coverage due to the larger variation in information about the population in samples drawn under an informative design, as earlier discussed. Figure 5 is the same as Figure 4 after increasing the sample size to $n = 1000$. The results are qualitatively the same as when $n = 100$ except that the pseudo posterior performs better in terms of bias; in agreement with the result that the pseudo posterior approach produces asymptotically unbiased estimators [Savitsky and Toth, 2016].




Fig 4: Simulation scenario non-linear: π -skewed with sample size $n = 100$. From top to bottom, average estimated curve, MSE, bias, coverage probability of 95% central credible intervals and their average length. The black curve in the upper panel is the true curve. For visual purposes the horizontal (discontinuous) lines at 0 and 0.95 in third and fourth panels, respectively, are depicted.




Fig 5: Simulation scenario non-linear: π -skewed with sample size $n = 1000$. Same as Figure 4 but with $n = 1000$.

5. Application

As an illustration we explore the relationship between caffeine and systolic blood pressure (SBP), applying our methodology to data from the 2013-2014 National Health and Nutrition Examination Survey (NHANES). We will perform two analyses. In the first analysis we do not adjust for any other covariate and in the second we adjust for age and gender.

Due in part to technical, budget and logistic considerations, the survey design of the NHANES is complex. NHANES is designed to assess the health and nutritional status of adults and children in the United States [CDC-A, 2016]. Data are collected from a sample of individuals from the non-institutionalized, civilian US population. Although nationally representative, the NHANES is designed to oversample specific subpopulations of interest (e.g. minorities, low income groups, pregnant women) for population-based studies using a complex, multistage, cluster sampling design [CDC-B, 2016]. To correct for informativeness (partly induced by the correlation between response values and group memberships), the NHANES survey data are released with observation-indexed sampling weights (based on marginal inclusion probabilities) corresponding to the sampled participants. These sampling weights are computed based on the sampling design (i.e. probability of the individual of being included in the sample), non-response and the masking of the individual for confidentiality [CDC-D, 2016].

NHANES participants are required to visit a mobile examination center where Health and nutrition information are collected during these visits. In particular, SBP is measured for participants of ages 8 years and older. A 24-hour dietary recall interview is taken. The results of this interview are used to estimate the participant's intakes of nutrients during the 24-hour period preceding their interview (from the period of midnight to midnight). Our dataset consists of systolic blood pressure (SBP) (mm Hg) measurements of $n = 6847$ participants 8 or older and 24-hour caffeine consumption (mg), along with their sampling weights, as estimated from the first 24-hour dietary recall interview. See Supplementary Material 9 for details regarding our data preprocessing.

We fit the spline basis model introduced in Subsection 2.3 with response, $y = \log(\text{SBP})$, inclusion probability, π , designed to be proportional to the inverse of the NHANES sampling weight, and predictor, $x = \log(\text{caffeine consumption} + 1)$. We fit both the fully Bayesian (Full) and the pseudo posterior (Pseudo) models. Additionally, against the recommendation stated in NHANES, we fit the splines model for the population that ignores IS (equivalent to the pseudo posterior model with all sampling weights equal to 1). The cubic B-spline model is constructed with the priors and hyperparameters described in Section 2.3.

After a burn-in period of 10^4 iterations, a Monte Carlo posterior sample of size 10^4 was retained for each of the model parameters and used to estimate $E(y | x)$, together with 95% credible intervals on a pointwise basis under the three models. Figure 6 displays these curves. There is a positive relationship between caffeine consumption and blood pressure that levels off in higher ranges of caffeine consumption. The Pseudo posterior estimates a gradually increasing




Fig 6: Caffeine consumption (mg) *v.s.* SBP (mm Hg) under the Full, Pseudo and Ignoring IS models.

relationship, while the Fully Bayes estimates 3 distinct regions in the support of $\log(\text{caffeine}+1)$, where each expresses a different sensitivity to $\log(\text{SBP})$; in particular, the credible intervals for these two models do not overlap in parts of the second region (from left-to-right) in a range of $2.5 \leq x \leq 4$, where the Fully Bayes model shows a notably greater sensitivity. In this case, then, the inferences performed under these two models would differ.

The resulting negative sign of κ_y , with central 95% credible interval $(-0.64, -0.35)$ and $\Pr(\kappa_y < 0 \mid \text{data}) \approx 1$, in the model for $\pi_i \mid y_i, x_i$ indicates that the sample design is informative for the logarithm of SBP (when not conditioned on predictors). The higher the value of the response the lower is the inclusion probability.

We next proceed to check if this positive association between SBP and caffeine consumption still holds after controlling for age and gender. We categorized the age into seven groups as shown in Table 3. The reference categories were the 8-17 age group and the male gender. We adjust the cubic B-spline model

$$y_i \sim \text{normal} \left(\mu(\mathbf{u}_i) + \beta_{\text{age}(i)}^{\text{age}} + \beta_{\text{gender}(i)}^{\text{gender}}, \sigma_y^2 \right)$$

with $\mu(\mathbf{u}_i)$ defined in Equation (10). $\beta_{\text{age}(i)}^{\text{age}}$ and $\beta_{\text{gender}(i)}^{\text{gender}}$ are the regression coefficients associated with the age group and gender of the participant i for groupings, $\text{age}(i) \in \{1, \dots, 7\}$ and $\text{gender}(i) \in \{1, 2\}$. The regression coefficients associated to the reference groups are set to 0 ($\beta_1^{\text{age}} \equiv \beta_1^{\text{gender}} \equiv 0$). We employ vague priors $\beta_2^{\text{age}}, \dots, \beta_7^{\text{age}}, \beta_2^{\text{gender}} \stackrel{\text{iid}}{\sim} \text{normal}(0, 10^4)$. Age group and

gender were also included as covariates in $\pi_i \mid y_i, \mathbf{v}_i$, where \mathbf{v}_i is a vector of dimension 9, by incorporating for an intercept, $x \equiv \log(\text{caffeine} + 1)$, seven age groups and gender. The estimated curves along with their 95% credible intervals for the reference group (males between 8 and 16 years old) under the three models are shown in Figure 7. The positive association between caffeine consumption and SBP vanishes when controlling for age and gender. The estimated curves under the Fully Bayes and Pseudo posterior models differ in shape but their bands overlap across all the range of x . The posterior mean curve for the fully Bayes model is, however, more smoothly centered on the horizontal line to more strongly indicate little-to-no association between systolic blood pressure and caffeine consumption for the reference group. The credible intervals for the Fully Bayes model and that ignoring IS are almost perfectly overlapping, which suggests a non-informative sampling design. The central 95% credible interval for κ_y , $(-.06, 0.28)$, contains zero, confirming that the sampling design is non-informative for SBP when controlling for age and gender. Table 3 depicts the posterior mean, standard deviation and 2.5% and 97.5% quantiles of the marginal posterior distribution of the Fully Bayes, adjusting for age and gender, model parameters. The regression coefficients in the conditional model for $y_i \mid \mathbf{u}_i$ indicate that mean SBP for females is lower and SBP increases with age.

When the sampling design is not informative for the chosen response (given the available predictors), the fully Bayes model provides similar inference than under SRS. By contrast, the pseudo posterior estimator is notably noisier than under SRS, which in some cases may lead to incorrect inference.

Steffen et al. [2012] performed a meta-analysis of randomized clinical trials and prospective studies and concluded that coffee consumption was not associated with a significant change in SBP. The Fully Bayes and ‘Ignoring IS’ curves in Figure 7 are nearly horizontal, which effectively reproduces this result. The pseudo posterior result, however, expresses a lot more oscillation around around the horizontal line.




Fig 7: Caffeine consumption (mg) *v.s.* SBP (mm Hg), or males 8-16 years old, under the Full, Pseudo and Ignoring IS models.

	mean	sd	2.5%	97.5%
Parameters of $y_i \mid \dots$				
Regression coefficients				
Gender (female)	-0.03	0.00	-0.04	-0.03
Age 17-24	0.08	0.00	0.07	0.09
Age 25-34	0.09	0.01	0.08	0.10
Age 35-39	0.12	0.01	0.11	0.13
Age 40-49	0.14	0.01	0.13	0.15
Age 50-59	0.18	0.01	0.17	0.19
Age 60 or older	0.24	0.00	0.23	0.25
σ_y	0.11	0.00	0.11	0.12
(Caffeine) Spline parameters in $y_i \mid \dots$				
β_1	4.66	0.03	4.59	4.73
\vdots				
Parameters of $\pi_i \mid y_i, \dots$				
Intercept	-0.91	0.41	-1.71	-0.11
κ_{zy}	0.11	0.09	-0.06	0.28
Caffeine	-0.05	0.01	-0.06	-0.04
Gender (Female)	-0.03	0.02	-0.07	0.01
Age 17-24	-0.55	0.04	-0.62	-0.48
Age 25-34	-0.74	0.04	-0.81	-0.67
Age 35-39	-0.67	0.05	-0.77	-0.58
Age 40-49	-0.67	0.04	-0.75	-0.59
Age 50-59	-0.72	0.04	-0.80	-0.63
Age 60 or older	-0.54	0.04	-0.62	-0.47
σ_π	0.83	0.01	0.81	0.84

Table 3: Fully Bayes model parameter estimates for caffeine consumption versus SBP. (Caffeine) spline parameters β_2, \dots, β_8 and σ_β are not depicted to save space.

6. Discussion

We have proposed a fully Bayes approach to incorporate the sampling weights into the estimation of population model parameters on data collected under an informative sampling design. The approach does not require extensive knowledge of sampling design, uses only quantities observed in sample, and discards variation in weights not dependent on response. We have shown via simulation that the method performs as well and often better in terms of bias, MSE and, the coverage of its central 95% credible interval than the non-fully Bayesian plug-in pseudo posterior. Our fully Bayes approach demonstrates a consistently superior ability to accurately measure uncertainty, in contrast with the pseudo posterior method, which is too confident such that it fails to achieve nominal coverage. The price to be paid for the achievement of unbiased inference with correct coverage of population model parameters from an informative sample is wider credible intervals to achieve the same coverage relative to simple random sampling. We applied the Fully Bayes approach to explore the relationship between caffeine consumption and SBP analyzing NHANES data. We notice a (non linear) positive association between these two variables. This association vanishes when controlling for age and gender. The Fully Bayesian approach estimates a smoother and closer to the horizontal line (indicating no association) caffeine vs SBP curve as compared to the pseudo posterior approach, which better agrees with previous research for the relationship between caffeine consumption and SBP.

The unusual assumption of our approach, from the perspective of the survey sampler, is that it considers the weights (inclusion probabilities) as random. Although the fully Bayes approach requires specification of a conditional distribution for the inclusion probabilities, we have shown that coverage properties are robust to misspecification of this distribution. A disadvantage of the proposed method is that it requires a customized posterior sampler because the posterior sampler designed for simple random sampling cannot easily be adapted to IS under the fully Bayes construction. We cope with this problem by relying on Stan [Carpenter et al., 2016]. A related issue is the required integration step for every observation at every iteration of the posterior sampler. This computation must be performed, numerically, in the general case. Theorem 1 provides some conditions in the sampling distribution and the conditional distribution of the weights given the response that allow a closed form for the fully Bayes likelihood. We applied the theorem to illustrate a collection of useful population joint distributions over the inclusion probabilities and response that all allow for a closed form for the integration step. More research is needed to apply our approach to more complicated settings. Future work will focus on performing the computation of this expected value, numerically, or circumventing it by introducing latent random variables.

Supplementary Material

Contains:

1. Proof of Theorem 1 that specifies conditions under which a closed form is available for the expectation step in the denominator of the fully Bayes estimator.
2. Derivation for the closed form expression for $\log p_s(\dots)$ in Subsection 2.1
3. Enabling Lemmas for Theorem 2.
4. Detailed data preprocessing steps for application in Section 5.

Supplementary Material: Fully Bayesian Estimation Under Informative Sampling

Luis G. León-Novelo[‡] and Terrance D. Savitsky[§]

1200 Pressler St. Suite E805, Houston, TX, 77030, USA and
 PSB Suite 1950, 2 Massachusetts Avenue, NE Washington, DC 20212-0001, USA.
 e-mail: Luis.G.LeonNovelo@uth.tmc.edu; Savitsky.Terrance@bls.gov

7, proof of Theorem 1; 8, closed form expression for $\log p_s(\dots)$ in Subsection 2.1; 8.1, enabling Lemmas for Theorem 2; and, 9, detailed data preprocessing steps for application in Section 5.

7. Proof of Theorem 1

Recall that the vector of covariates \mathbf{x}_i has been split into two sets of covariates, \mathbf{u}_i and \mathbf{v}_i , relevant for the distribution of $y_i \mid \boldsymbol{\theta}, \mathbf{u}_i$, and $\pi_i \mid y_i, \mathbf{v}_i, \boldsymbol{\kappa}$, respectively. Further recall the definition for $M_y(\boldsymbol{\kappa}; \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}) := E_{y \mid \mathbf{u}_i, \boldsymbol{\theta}}[\exp\{g(y_i, \mathbf{v}_i, \boldsymbol{\kappa})\}]$. Using the fact that $v \sim \text{lognormal}(m, v^2) \Rightarrow E(v) = \exp(m + v^2/2)$, we obtain,

$$\begin{aligned} E_{y \mid \mathbf{u}_i, \boldsymbol{\theta}}\{E[\pi_i \mid y, \mathbf{v}_i, \boldsymbol{\kappa}]\} &= E_{y \mid \mathbf{u}_i, \boldsymbol{\theta}}[\exp\{g(y_i, \mathbf{v}_i, \boldsymbol{\kappa}) + t(\mathbf{v}_i, \boldsymbol{\kappa}) + \sigma_\pi^2/2\}] \\ &= \exp[t(\mathbf{v}_i, \boldsymbol{\kappa}) + \sigma_\pi^2/2] \times E_{y \mid \mathbf{u}_i, \boldsymbol{\theta}}[\exp\{g(y_i, \mathbf{v}_i, \boldsymbol{\kappa})\}] \\ &= \exp[t(\mathbf{v}_i, \boldsymbol{\kappa}) + \sigma_\pi^2/2] \times M_y(\boldsymbol{\kappa}; \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}) \end{aligned}$$

Noting that $\text{lognormal}(\pi \mid m, v^2) = \text{normal}(\log \pi \mid m, v^2) \times 1/\pi$, plugging the expression above in equation (5) we obtain,

$$\begin{aligned} p_s(y_i, \pi_i \mid \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) &= \frac{\pi_i p(\pi_i \mid y_i, \mathbf{v}_i, \boldsymbol{\kappa})}{E_{y \mid \mathbf{u}_i, \boldsymbol{\theta}}\{E[\pi_i' \mid y, \mathbf{v}_i, \boldsymbol{\kappa}]\}} p(y_i \mid \mathbf{u}_i, \boldsymbol{\theta}) \\ &= \frac{\pi_i \times \text{lognormal}(\pi_i \mid t(\mathbf{v}_i, \boldsymbol{\kappa}) + g(y_i, \mathbf{v}_i, \boldsymbol{\kappa}), \sigma_\pi^2)}{\exp[t(\mathbf{v}_i, \boldsymbol{\kappa}) + \sigma_\pi^2/2] \times M_y(\boldsymbol{\kappa}; \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta})} \times p(y_i \mid \mathbf{u}_i, \boldsymbol{\theta}) \\ &= \frac{\cancel{\pi_i} \times (1/\cancel{\pi_i}) \times \text{normal}(\log \pi_i \mid t(\mathbf{v}_i, \boldsymbol{\kappa}) + g(y_i, \mathbf{v}_i, \boldsymbol{\kappa}), \sigma_\pi^2)}{\exp[t(\mathbf{v}_i, \boldsymbol{\kappa}) + \sigma_\pi^2/2] \times M_y(\boldsymbol{\kappa}; \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta})} \times p(y_i \mid \mathbf{u}_i, \boldsymbol{\theta}), \end{aligned}$$

[‡]Department of Biostatistics and Data Science, University of Texas Health Science Center at Houston- School of Public Health

[§]Office of Survey Methods Research, Bureau of Labor Statistics

proving the theorem.

8. Closed-Form for log likelihood of Simple Linear Regression Model

$$\begin{aligned} \log p_s(y_i, \pi_i \mid \mathbf{u}_i, \mathbf{v}_i, \boldsymbol{\theta}, \boldsymbol{\kappa}) &\propto -\frac{1}{2} \log(\sigma_\pi^2) - \frac{1}{2\sigma_\pi^2} [\log \pi_i - (\kappa_y y_i + \mathbf{v}_i^t \boldsymbol{\kappa}_x)]^2 \\ &\quad - \frac{1}{2} \log(\sigma_y^2) - \frac{1}{2\sigma_y^2} [y_i - \mathbf{u}_i^t \boldsymbol{\beta}]^2 \\ &\quad - \mathbf{v}_i^t \boldsymbol{\kappa}_x - \sigma_\pi^2/2 - \kappa_y \mathbf{u}_i^t \boldsymbol{\beta} - \kappa_y^2 \sigma_y^2/2 \end{aligned}$$

$$\begin{aligned} \log p_s(\mathbf{y}, \boldsymbol{\pi} \mid \mathbf{x}, \boldsymbol{\theta}, \boldsymbol{\kappa}) &\propto -\frac{n}{2} \log(\sigma_\pi^2) - \frac{1}{2\sigma_\pi^2} \sum_i [\log \pi_i - (\kappa_y y_i + \mathbf{v}_i^t \boldsymbol{\kappa}_x)]^2 \\ &\quad - \frac{n}{2} \log(\sigma_y^2) - \frac{1}{2\sigma_y^2} \sum_i [y_i - \mathbf{u}_i^t \boldsymbol{\beta}]^2 \\ &\quad - \sum_i \mathbf{v}_i^t \boldsymbol{\kappa}_x - n\sigma_\pi^2/2 - \kappa_y \sum_i \mathbf{u}_i^t \boldsymbol{\beta} - n\kappa_y^2 \sigma_y^2/2 \end{aligned}$$

8.1. Enabling Lemmas for Theorem 2

We next construct two enabling results needed to prove Theorem 2 to account informative sampling under (A4), (A5) and (A6). The first enabling result is used to bound from above the numerator in the expression for the expectation with respect to the joint distribution for population generation and the taking of the informative sample, (P_{λ_0}, P_ν) , of the fully Bayesian, sampling-weighted posterior distribution in Equation (17) on the restricted set of measures that includes those P_λ that are at some *minimum* distance, $\delta\xi_{N_\nu}$, from P_{λ_0} under pseudo Hellinger metric, $d_{N_\nu}^\pi$. The second result, Lemma 4, extends Lemma 8.1 of Ghosal et al. [2000] to bound the probability of the denominator of Equation (17) with respect to (P_{λ_0}, P_ν) , from below. Given these 2 Lemmas, the subsequent proof exposition for Theorem 2 is identical to Savitsky and Toth [2016].

Lemma 3. *Suppose conditions (A1) and (A4) hold. Then for every $\xi > \xi_{N_\nu}$,*

a constant, $K > 0$, and any constant, $\delta > 0$,

$$\mathbb{E}_{P_0, P_\nu} \left[\int_{P \in \mathcal{P} \setminus \mathcal{P}_{N_\nu}} \prod_{i=1}^{N_\nu} \frac{p^\pi}{p_0^\pi} (x_{\nu i} \delta_{\nu i}) d\Pi(\lambda) (1 - \phi_{n_\nu}) \right] \leq \gamma \Pi(\Lambda \setminus \Lambda_{N_\nu}) \quad (20)$$

$$\mathbb{E}_{P_{\lambda_0}, P_\nu} \left[\int_{\lambda \in \Lambda_{N_\nu} : d_{N_\nu}^\pi(P_\lambda, P_{\lambda_0}) > \delta \xi} \prod_{i=1}^{N_\nu} \frac{p^\pi}{p_0^\pi} (x_{\nu i} \delta_{\nu i}) d\Pi(\theta) (1 - \phi_{n_\nu}) \right] \leq 2\gamma^2 \exp\left(\frac{-Kn_\nu \delta^2 \xi^2}{\gamma}\right). \quad (21)$$

This result is adjusted from that in [Savitsky and Toth \[2016\]](#) by multiplying the upper bounds on the right-hand side of both equations by constant multiplier, $\gamma \geq 1$, defined in condition [\(A4\)](#).

Proof. The proof approach is the same as [Savitsky and Toth \[2016\]](#), where we first bound the left-hand sides of Equations (20) and (21) from above by an expectation of the test statistic, ϕ_{n_ν} . When then further refine this bound on the the two subsets of measures outlined in those equations. The refinement step is unchanged from [Savitsky and Toth \[2016\]](#), but the first step is revised due to our unique form for our sampling-weighted, fully Bayesian estimator, outlined in Equation (16). So we fully repeat the first step in the proof, below.

Fixing ν , we index units that comprise the population with, $U_\nu = \{1, \dots, N_\nu\}$. Next, draw a single observed sample of n_ν units from U_ν , indexed by subsequence,

$\{i_\ell \in U_\nu : \delta_{\nu i_\ell} = 1, \ell = 1, \dots, n_\nu\}$. Without loss of generality, we simplify notation to follow by indexing the observed sample, sequentially, with $\ell = 1, \dots, n_\nu$.

We next decompose the expectation under the joint distribution with respect to population generation, P_{λ_0} , and the drawing of a sample, P_ν ,

Suppose we draw λ from some set $B \subset \Lambda$. By Fubini,

$$\begin{aligned} & \mathbb{E}_{P_{\lambda_0}, P_\nu} \left[\int_{P \in B} \prod_{i=1}^{N_\nu} \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} (x_{\nu i} \delta_{\nu i}) d\Pi(\lambda) (1 - \phi_{n_\nu}) \right] \\ & \leq \int_{\lambda \in B} \left[\mathbb{E}_{P_0, P_\nu} \prod_{i=1}^{N_\nu} \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} (x_{\nu i} \delta_{\nu i}) (1 - \phi_{n_\nu}) \right] d\Pi(\lambda) \end{aligned} \quad (22)$$

$$\leq \int_{\lambda \in B} \left\{ \sum_{\delta_\nu \in \Delta_\nu} \mathbb{E}_{P_{\lambda_0}} \left[\prod_{\ell=1}^{n_\nu} \left[\frac{\pi_{\nu \ell} p_\lambda}{\pi_{\nu \ell}^\lambda p_{\lambda_0}} (x_{\nu \ell}) \right] (1 - \phi_{n_\nu}) \middle| \delta_\nu \right] P_{P_\nu}(\delta_\nu) \right\} d\Pi(\lambda) \quad (23)$$

$$\leq \int_{\lambda \in B} \left\{ \sum_{\delta_\nu \in \Delta_\nu} \mathbb{E}_{P_{\lambda_0}} \left[\prod_{\ell=1}^{n_\nu} \left[\frac{1}{\pi_{\nu \ell}^\lambda} \frac{p_\lambda}{p_{\lambda_0}} (x_{\nu \ell}) \right] (1 - \phi_{n_\nu}) \middle| \delta_\nu \right] P_{P_\nu}(\delta_\nu) \right\} d\Pi(\lambda) \quad (24)$$

$$\leq \int_{\lambda \in B} \max_{\delta_\nu \in \Delta_\nu} \mathbb{E}_{P_{\lambda_0}} \left[\prod_{\ell=1}^{n_\nu} \left[\frac{1}{\pi_{\nu \ell}^\lambda} \frac{p_\lambda}{p_{\lambda_0}} (x_{\nu \ell}) \right] (1 - \phi_{n_\nu}) \middle| \delta_\nu \right] d\Pi(\lambda) \quad (25)$$

$$\leq \int_{\lambda \in B} \mathbb{E}_{P_{\lambda_0}} \left[\prod_{\ell=1}^{n_\nu} \left[\frac{1}{\pi_{\nu \ell}^\lambda} \frac{p_\lambda}{p_{\lambda_0}} (x_{\nu \ell}) \right] (1 - \phi_{n_\nu}) \middle| \delta_\nu^* \right] d\Pi(\lambda) \quad (26)$$

$$\leq \gamma \int_{\lambda \in B} \mathbb{E}_{P_{\lambda_0}} \left[\prod_{\ell=1}^{n_\nu} \left[\frac{p_\lambda}{p_{\lambda_0}} (x_{\nu \ell}) \right] (1 - \phi_{n_\nu}) \middle| \delta_\nu^* \right] d\Pi(\lambda) \quad (27)$$

$$\leq \gamma \int_{\lambda \in B} \mathbb{E}_{\delta_\nu^*} (1 - \phi_{n_\nu}) d\Pi(\lambda),$$

where $\sum_{\delta_\nu \in \Delta_\nu} P_{P_\nu}(\delta_\nu) = 1$ [Särndal et al., 2003] and

$\delta_\nu^* \in \Delta_\nu = \left\{ \{\delta_{\nu i}^*\}_{i=1, \dots, N_\nu}, \delta_{\nu i}^* \in \{0, 1\} \right\}$ denotes that sample, drawn from the space of all possible samples, Δ_ν , which maximizes the probability under the population generating distribution for the event of interest. The inequality in Equation (27) results from the bound, $\frac{1}{\pi_{\nu \ell}^\lambda} \leq \gamma$, specified in Condition (A4).

The remainder (second step) of the proof is the same as in Savitsky and Toth [2016], except one multiplies their result by γ to compute the bounds for Equations (20) and (21). \square

Lemma 4. For every $\xi > 0$ and measure Π on the set,

$$B = \left\{ \lambda : -\mathbb{E}_{\lambda_0} \log \left(\frac{p_\lambda}{p_{\lambda_0}} \right) \leq \xi^2, \mathbb{E}_{\lambda_0} \left(\log \frac{p_\lambda}{p_{\lambda_0}} \right)^2 \leq \xi^2 \right\}$$

under the conditions (A2), (A3), (A4), and (A5), we have for every $C > 0$

and N_ν sufficiently large,

$$\Pr \left\{ \int_{\lambda \in \Lambda} \prod_{i=1}^{N_\nu} \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} (x_{\nu i} \delta_{\nu i}) d\Pi(\lambda) \leq \exp[-(1+C)N_\nu \xi^2] \right\} \leq \frac{\gamma + C_3}{C^2 N_\nu \xi^2}, \quad (28)$$

where the above probability is taken with the respect to P_{λ_0} and the sampling generating distribution, P_ν , jointly.

The sum of positive constants, $\gamma + C_3$, is greater than 1 and will be larger for sampling designs where the (modeled) inclusion probabilities, $\{\pi_{\nu i}^\lambda\}$, are more variable.

Proof. The first part of the proof bounds the integral on the left-hand side of the event over which we take the probability (in Equation (28)), from below, by a centered and scaled empirical process. This first part is altered under our construction for our sampling-weighted, fully Bayesian estimator, outlined in Equation (16). So we specify this part, here.

By Jensen's inequality,

$$\begin{aligned} \log \int_{\lambda \in \Lambda} \prod_{i=1}^{N_\nu} \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} (x_{\nu i} \delta_{\nu i}) d\Pi(\lambda) &\geq \sum_{i=1}^{N_\nu} \int_{\lambda \in \Lambda} \log \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} (x_{\nu i} \delta_{\nu i}) d\Pi(\lambda) \\ &= N_\nu \cdot \mathbb{P}_{N_\nu} \int_{\lambda \in \Lambda} \log \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} d\Pi(\lambda), \end{aligned}$$

where we recall that the last equation denotes the empirical expectation functional taken with respect to the joint distribution over population generating and informative sampling. By Fubini,

$$\mathbb{P}_{N_\nu} \int_{\lambda \in \Lambda} \log \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} d\Pi(\lambda) = \int_{\lambda \in \Lambda} \left[\mathbb{P}_{N_\nu} \log \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} \right] d\Pi(\lambda) \quad (29)$$

$$= \int_{\lambda \in \Lambda} \left[\mathbb{P}_{N_\nu} \delta_\nu \log \left\{ \frac{\pi_\nu}{\pi_\nu^\lambda} \frac{p_\lambda}{p_{\lambda_0}} \right\} \right] d\Pi(\lambda) \quad (30)$$

$$\geq \int_{\lambda \in \Lambda} \left[\mathbb{P}_{N_\nu} \frac{\delta_\nu}{\pi_\nu^\lambda} \log \frac{p_\lambda}{p_{\lambda_0}} \right] d\Pi(\lambda) \quad (31)$$

$$= \int_{\lambda \in \Lambda} \left[\mathbb{P}_{N_\nu}^\pi \log \frac{p_\lambda}{p_{\lambda_0}} \right] d\Pi(\lambda) \quad (32)$$

$$= \mathbb{P}_{N_\nu}^\pi \int_{\lambda \in \Lambda} \log \frac{p_\lambda}{p_{\lambda_0}} d\Pi(\lambda), \quad (33)$$

where we, again, apply Fubini. Equation (30) provides an upper bound for Equation (31) because $\pi_{\nu i} \leq 1$, $\frac{1}{\pi_{\nu i}^\lambda} > 0$, while $\log \frac{p_\lambda^\pi}{p_{\lambda_0}^\pi} (x_{\nu i}) < 0$.

The remainder of the proof, which uses Chebyshev to provide an upper bound on the probability of the event in Equation (28), is specified by,

$$\leq \xi^2 \sup_{\nu} \left[\frac{1}{\min_{i \in U_{\nu}} \pi_{\nu i}^{\lambda}} \right] + \xi^2 (N_{\nu} - 1) \sup_{\nu} \max_{i \neq j \in U_{\nu}} \left[\left| \frac{\mathbb{E}_{\lambda_0} [\pi_{\nu ij}]}{\pi_{\nu i}^{\lambda} \pi_{\nu j}^{\lambda}} - 1 \right| \right], \quad (34)$$

which is nearly identical to Savitsky and Toth [2016] with the raw, $\pi_{\nu i}$, replaced by the modeled, $\pi_{\nu i}^{\lambda} = \mathbb{E}_{P_{\theta}} (\pi_{\nu i}^{\kappa})$, where $\pi_{\nu i}^{\kappa} = \mathbb{E}_{\kappa} (\pi_{\nu i} | Y_{\nu i} = y_{\nu i})$ in the first term on the left of Equation (34). Under condition (A5), $\mathbb{E}_{\lambda_0} [\pi_{\nu ij}] \rightarrow \mathbb{E}_{\lambda_0} [\pi_{\nu i} \pi_{\nu j}] = \pi_{\nu i}^{\lambda_0} \pi_{\nu j}^{\lambda_0}$. Condition (A7) guarantees convergence to 1 of the ratio in the second term of Equation (34). \square

9. Data preprocessing in Section 5

Here we describe how we obtained our dataset of $n = 6847$ participants that we used in analyses in Section 5. We downloaded the ‘‘Dietary Interview - Total Nutrient Intakes, First Day’’ and ‘‘Blood Pressure’’ datasets. These datasets, named DR1TOT.H.XPT and BPX.H.XPT, are available at <https://www.cdc.gov/Nchs/Nhanes/Search/DataPage.aspx?Component=Dietary&CycleBeginYear=2013> and https://www.cdc.gov/Nchs/Nhanes/2013-2014/BPX_H.htm, respectively.

The first dataset includes the ‘‘Dietary day one sample weight’’ variable (called WTDRD1) and its documentation states that this variable should be used as a sampling weight when analyzing it, as we did in our analyzes. The dataset contains the nutrient intake information 9813 NHANES participants. This dataset is derived from questionnaire data where each participant is asked questions on salt, amounts of food and beverages consumed during the 24-hour period prior to the interview (midnight to midnight). For more details see the dataset documentation. 1152 participants have sampling weights equal to zero leaving 9813-1152=8661 (the participants with weight 0 also had missing values in all their nutrient consumption data) participants in this analysis.

The second dataset includes three consecutive blood pressure readings. According to its documentation, when a blood pressure measurement is interrupted or incomplete, a fourth attempt may be made. In our analysis SBP was the average of the not missing values of these four reads. This dataset contains the information of the 9813 participants in the first dataset but only 7818 are eight years or older (by design their SBP was not measured for younger participants), from these 807 have sampling weights equal to zero leaving 7818-807=7011. Out of these 7011, 6847 at least one (out of four) SBP measure recorded. Figure 8 in this Supplementary Material depicts the Venn diagram of the sets of NHANES participants with non-zero sampling weights, 8 years or older and SBP recorded at least once.

To obtain age and gender information of the participants we downloaded the dataset DEMO.H.XPT available at <https://www.cdc.gov/nchs/nhanes/search/datapage.aspx?Component=Demographics&CycleBeginYear=2013>.




Fig 8: Venn diagram the three sets of NHANES participants. The sets are with non-zero weights, 8 years or older and SBP recorded at least once.

References

- Bob Carpenter, Andrew Gelman, Matt Hoffman, Daniel Lee, Ben Goodrich, Michael Betancourt, Michael A Brubaker, Jiqiang Guo, Peter Li, and Allen Riddell. Stan: A probabilistic programming language. *Journal of Statistical Software*, 20:1–37, 2016.
- CDC-A. Centers for Disease Control (CDC). national health and nutrition examination survey. <https://www.cdc.gov/nchs/nhanes/index.htm>, 2016. Accessed: June 7, 2016.
- CDC-B. Centers for Disease Control (CDC). modeling usual intake using dietary recall data, task 4: Key concepts about variance estimation methods in NHANES. <https://www.cdc.gov/nchs/tutorials/dietary/Advanced/ModelUsualIntake/Info4.htm>, 2016. Accessed: June 7, 2016.
- CDC-D. Centers for Disease Control (CDC). key concepts about NHANES survey design: NHANES sampling procedure. <http://www.cdc.gov/nchs/tutorials/NHANES/SurveyDesign/SampleDesign/Info1.htm>, 2016. Accessed: June 7, 2016.
- Qi Dong, Michael R. Elliott, and Trivellore E. Raghunathan. A nonparametric method to generate synthetic populations to adjust for complex sampling design features. *Survey Methodology*, 40(1):29–46, 2014.
- Subhashis Ghosal, Jayanta K Ghosh, and Aad W Van Der Vaart. Convergence rates of posterior distributions. *Annals of Statistics*, 28(2):500–531, 2000.
- David R Judkins. Fay’s method for variance estimation. *Journal of Official Statistics*, 6(3):223, 1990.
- D Krewski and JNK Rao. Inference from stratified samples: properties of the linearization, jackknife and balanced repeated replication methods. *The Annals of Statistics*, pages 1010–1019, 1981.
- Tsuyoshi Kuniyama, AH Herring, CT Halpern, and DB Dunson. Nonparametric bayes modeling with sample survey weights. *Statistics & Probability Letters*, 113:41–48, 2016.

- Roderick J. Little. To model or not to model? Competing modes of inference for finite population sampling. *Journal of the American Statistical Association*, 99(466):546–556, 2004.
- Philip J McCarthy. Pseudo-replication: Half samples. *Revue de l'Institut International de Statistique*, pages 239–264, 1969.
- Danny Pfeiffermann, Abba M Krieger, and Yosef Rinott. Parametric distributions of complex survey data under informative probability sampling. *Statistica Sinica*, pages 1087–1114, 1998.
- Danny Pfeiffermann, Fernando Antonio Da Silva Moura, and Pedro Luis Do Nascimento Silva. Multi-level modelling under informative sampling. *Biometrika*, 93(4):943–959, 2006.
- J. N. K. Rao and Changbao Wu. Bayesian pseudo-empirical-likelihood intervals for complex surveys. *Journal of the Royal Statistical Society Series B*, 72(4):533–544, 2010.
- Carl-Erik Särndal, Bengt Swensson, and Jan Wretman. *Model Assisted Survey Sampling (Springer Series in Statistics)*. Springer Science & Business Media, 2003. ISBN 0387406204.
- Terrance D Savitsky and Daniell Toth. Bayesian estimation under informative sampling. *Electronic Journal of Statistics*, 10(1):1677–1708, 2016.
- Yajuan Si, Natesh S Pillai, Andrew Gelman, et al. Bayesian nonparametric weighted sampling inference. *Bayesian Analysis*, 10(3):605–625, 2015.
- Paul L Speckman and Dongchu Sun. Fully bayesian spline smoothing and intrinsic autoregressive priors. *Biometrika*, 90(2):289–302, 2003.
- Mark Steffen, Carol Kuhle, Donald Hensrud, Patricia J Erwin, and Mohammad H Murad. The effect of coffee consumption on blood pressure and the development of hypertension: a systematic review and meta-analysis. *Journal of hypertension*, 30(12):2245–2254, 2012.