

ON COBORDISM OF GENERALIZED (REAL) BOTT MANIFOLDS

YUXIU LU

ABSTRACT. We show that all generalized (real) Bott manifolds which are (small covers) quasitoric manifolds over a product of simplices $\Delta^{n_1} \times \cdots \times \Delta^{n_r} \times \Delta^1$ are always boundaries of some manifolds. But these manifolds with the natural $(\mathbb{Z}_2)^n$ action do not necessarily bound equivariantly. In addition, we can construct some examples of null-cobordant but not orientedly null-cobordant manifolds among quasitoric manifolds.

1. INTRODUCTION

In this paper, we will mainly investigate cobordism of certain type of manifolds called real Bott manifold and generalized real Bott manifolds (see Kamishima and Masuda [7]). A real Bott manifold is defined to be a sequence of $\mathbb{R}P^1$ -bundles

$$M_n \xrightarrow{\mathbb{R}P^1} M_{n-1} \xrightarrow{\mathbb{R}P^1} \cdots \xrightarrow{\mathbb{R}P^1} M_1 \xrightarrow{\mathbb{R}P^1} M_0 = \{a \text{ point}\}$$

such that $M_i \rightarrow M_{i-1}$ for $i = 1, \dots, n$ is the projective bundle of a Whitney sum of two real line bundles over M_i . It is well known that a real Bott manifold is a flat manifold, which means that it admits a Riemannian metric with everywhere zero sectional curvature. Then this manifold is a boundary, since any closed manifold with a flat Riemannian metric is a boundary (see G.C. Hamrick and D.C. Royster, [3]).

Any real Bott manifold bounds equivariantly (see Y. Cheng and Y. Wang [6], Z. Lü and Q. Tan [10]). But their methods are not easy to use to discuss a generalized real Bott manifold. So we need some new technique to deal with the general case.

This study of n -dimensional real Bott manifolds is equivalent to the study of small covers over an n -cube (see Kamishima [7], Suh [4], Masuda [8]). This kind of manifolds and quasitoric manifolds are defined in Davis-Januszkiewicz [5]. By a small cover one means an n -dimensional manifold M^n with \mathbb{Z}_2^n -action locally isomorphic to the standard action of \mathbb{Z}_2^n on \mathbb{R}^n for which the orbit space M^n/\mathbb{Z}_2^n is a simple convex polytope P^n . A quasitoric manifold means a smooth orientable

Date: September 17, 2017.

Key words and phrases. real Bott manifold, generalized real Bott manifold, small cover, quasitoric manifold, generalized Bott manifold, cobordism, product of simplices.

$2n$ -dimensional manifold M^{2n} with T^n -action locally isomorphic to the standard action of T^n on \mathbb{C}^n for which the orbit space M^{2n}/T^n is a simple convex polytope P^n . The facial submanifold of a quasitoric manifold or a small cover is defined to be $\pi^{-1}(F)$, where $\pi : M^{2n} \rightarrow P^n$ is a quotient map, $d = 1, 2$. The concept of characteristic map is very important in the construction of the small cover and quasitoric manifold. Take a small cover as an example. A characteristic map means a function $\lambda : \{\text{the facets of } P^n\} \rightarrow \mathbb{Z}_2^n$, which satisfies that for every vertex $v = F_1 \cap \cdots \cap F_n$ of P^n , $\lambda(F_1), \dots, \lambda(F_n)$ form a basis of \mathbb{Z}_2^n . The characteristic map can also be identified with a map mapping each facet F to an element in \mathbb{Z}_2^n , where $\lambda(F)$ is the generator of the isotropy group of the facial submanifold $\pi^{-1}(F)$ for each facet F . The characteristic map determines a small cover over P^n and that any function satisfying the above condition can be realized as the characteristic function of some small cover.

In their paper, Davis and Januszkiewicz also provide following formulas to compute the total Pontryagin class and the total Stiefel-Whitney class for small covers and quasitoric manifolds.

Theorem 1.1. (Davis and Januszkiewicz [5, corollary 6.8]) *Let M^n be a small cover over a simple polytope P^n . Let v_1, \dots, v_m denote elements in the cohomology ring of M dual to the facial submanifolds of all the facets of M . Let $j : M^n \rightarrow B_d P^n$ be inclusion of the fiber.*

(i) *If $d = 1$, then*

$$w(M^{dn}) = j^* \prod_{i=1}^m (1 + v_i), \text{ and } p(M^{dn}) = 1.$$

(ii) *If $d = 2$, then*

$$w(M^{dn}) = j^* \prod_{i=1}^m (1 + v_i) \text{ mod } 2, \text{ and } p(M^{dn}) = j^* \prod_{i=1}^m (1 - v_i^2).$$

Davis and Januszkiewicz also give a method to compute the cohomology ring of small covers and quasitoric manifolds.

Theorem 1.2. (Davis and Januszkiewicz [5, Theorem 4.14]) *If M^n is a small cover, then $H^*(M; \mathbb{Z}_2) = \mathbb{Z}_2[v_1, \dots, v_m]/(I + J)$; If M^{2n} is a quasitoric manifold, then $H^*(M; \mathbb{Z}) = \mathbb{Z}[v_1, \dots, v_m]/(I + J)$; I is the Stanley-Reisner ideal of P^n , J is generated by linear combinations of v_1, \dots, v_m which are determined by characteristic map λ .*

By these two theorems, we can prove that any real Bott manifold is cobordant to zero, and more generally, any generalized real Bott manifold over $P^n \times \Delta^1$ is cobordant to zero, where P^n is an arbitrary product of simplices (see Theorem 3.4). This gives some examples of small covers which are boundaries while not

bounding equivariantly. Furthermore, one sufficient condition for a generalized real Bott manifold over $P^n \times \Delta^k$ to be null-cobordant, is given for odd numbers $k = 2^l - 1$ (see Theorem 3.6). Finally, for any product of simplices P^n , the generalized Bott manifold over $P^n \times \Delta^1$ is proved to be orientedly cobordant to zero just as the case of generalized real Bott manifold (see Theorem 4.3), and we can confirm that a special kind of quasitoric manifolds over an even dimensional cube are all unorientedly cobordant to zero, but are not orientedly cobordant to zero (see Theorem 4.5).

2. COBORDISM OF REAL BOTT MANIFOLD

Suppose that M^n is a small cover over an n -cube. Let the first \mathbb{Z}_2 -coefficient cohomology classes v_1, \dots, v_n dual to the facial submanifolds of facets F_1, \dots, F_n , meeting at a vertex. Then we have facets F_1^*, \dots, F_n^* which are parallel to F_1, \dots, F_n respectively. Let the first \mathbb{Z}_2 -coefficient cohomology classes u_1, \dots, u_n be dual to the facial submanifolds of facets F_1^*, \dots, F_n^* . Since $\{\lambda(F_1), \dots, \lambda(F_n)\}$ is a basis of \mathbb{Z}_2^n , we can assume that $\lambda(F_1) = e_1, \dots, \lambda(F_n) = e_n$. So $\lambda(F_1^*), \dots, \lambda(F_n^*)$ can be written in the form of linear combinations of e_1, \dots, e_n , which will give a \mathbb{Z}_2 -coefficient matrix of order n . This matrix is defined to be the *reduced matrix* of the small cover M^n , denoted by $A = (a_{ij})$, where

$$\lambda(F_i^*) = \sum a_{ij} e_j. \quad (1)$$

The theorem of Davis and Januszkiewicz gives the total Stiefel-Whitney class $w = (1 + u_1)(1 + v_1) \cdots (1 + u_n)(1 + v_n)$. By the restrictions given by the Stanley-Reisner ideal I , $w = (1 + u_1 + v_1) \cdots (1 + u_n + v_n)$.

Define $y_i = u_i + v_i$, then $w = (1 + y_1) \cdots (1 + y_n)$.

The restrictions given by the ideal J determines another relation:

$$(u_1, \dots, u_n)(A + E_n) = (y_1, \dots, y_n), \quad (2)$$

where E_n is the identity matrix of order n .

Lemma 2.1. *Let A be an $n \times n$ matrix with entries in \mathbb{Z}_2 . Suppose that every principal minor of A is 1. If $\det A = 1$, then A is conjugated by a permutation matrix to a unipotent upper triangular matrix.*

Proof. see M. Masuda and T.E. Panov[8, Lemma 3.3]. \square

Lemma 2.2. *The reduced matrix of a small cover over n -cube can be conjugated by a permutation matrix to a unipotent upper triangular matrix.*

Proof. Owing to the linear independence of characteristic maps of facets in every vertex, we can easily get that every principal minor of A is 1. So, by the Lemma 2.1 above, the conclusion is obvious. \square

Lemma 2.3. *If we assume the conditions above, and if $l_1 + \cdots + l_k = k + 1$, where $1 \leq k \leq n$. Then for 1-cohomology classes u_1, \dots, u_n of M^n , $u_1^{l_1} \cdots u_k^{l_k} = 0$.*

Proof. If $k = n$, the equation is obvious because the degree of $u_1^{l_1} \cdots u_k^{l_k}$ is greater than the dimension of small cover. If $k < n$, this theorem can be proved by induction. When $k = 1$, then $l_1 = 2$, the only equation one should justify is $u_1^2 = 0$, which will be proved in Lemma 3.3 in a broader sense. By the Lemma 2.2, we can assume that the *reduced matrix* of a small cover is a unipotent upper-diagonal matrix. This means that y_k can be expressed by a linear combination of u_1, \dots, u_{k-1} , $1 \leq k \leq n$. Assume that all the equations of number k is correct, then consider equations of number $k + 1$.

$$u_1^{l_1} \cdots u_{k+1}^{l_{k+1}} = u_1^{l_1} \cdots u_k^{l_k} u_{k+1} y_{k+1}^{l_{k+1}-1}.$$

The degree of $u_1^{l_1} \cdots u_k^{l_k} y_{k+1}^{l_{k+1}-1}$ is $l_1 + \cdots + l_{k+1} - 1 = k + 1$. Since y_{k+1} can be expressed by u_1, \dots, u_k , $u_1^{l_1} \cdots u_k^{l_k} y_{k+1}^{l_{k+1}-1}$ consists of monomials with elements in u_1, \dots, u_k of degree $k + 1$. So by induction, these polynomials are all zero, and $u_1^{l_1} \cdots u_{k+1}^{l_{k+1}}$ is zero. \square

Lemma 2.4. *Given the conditions above, if $l_2 + \cdots + l_k = k$, then $y_2^{l_2} \cdots y_k^{l_k} = 0$.*

Proof. $y_2^{l_2} \cdots y_k^{l_k}$ consists of monomials with elements in u_1, \dots, u_{k-1} of degree k . So due to the above Lemma 2.3, we can obtain this equation. \square

Theorem 2.5. *All the Stiefel-Whitney numbers of a small cover M over n -cube is 0, i.e. such a small cover is a boundary.*

Proof. Since $y_1 = 0$, the total Stiefel-Whitney class of M , $w = (1 + y_2) \cdots (1 + y_n)$. It is obvious that when $i_1 + 2i_2 + \cdots + ni_n = n$, $w_1^{i_1} \cdots w_n^{i_n}$ consists of polynomials in y_2, \dots, y_n of degree n . By the Lemma 2.4 above and the definition of Stiefel-Whitney numbers, this conclusion is proved. \square

3. COBORDISM OF GENERALIZED REAL BOTT MANIFOLD

In this section, we will mainly use the terms defined in S. Choi, M. Masuda and D.Y. Suh [4]. Firstly, generalized real Bott manifold M^n is defined to be a sequence of $\mathbb{R}P^k$ -bundles of various numbers,

$$M_m \xrightarrow{\mathbb{R}P^{n_m}} M_{m-1} \xrightarrow{\mathbb{R}P^{n_{m-1}}} \cdots \xrightarrow{\mathbb{R}P^{n_2}} M_1 \xrightarrow{\mathbb{R}P^{n_1}} M_0 = \{a \text{ point}\},$$

such that $M_i \longrightarrow M_{i-1}$ for $i = 1, \dots, m$ is the projective bundle of a Whitney sum of n_i real line bundles over M_i . This definition is equivalent to another one that a generalized real Bott manifold is a small cover over a product of simplices P^n , where $P^n = \prod_{i=1}^m \Delta^{n_i}$, with $\sum_{i=1}^m n_i = n$, and Δ^{n_i} is the n_i -simplex for $i = 1 \cdots m$. Then each facet of P^n is the product of a codimension-one face of one of the Δ^{n_i} 's

and the remaining simplices. Thus there are $n+m$ facets of P , which form the set $\{F_{k_i}^i | 0 \leq k_i \leq n_i, 1 \leq i \leq m\}$, where $F_{k_i}^i = \Delta^{n_1} \times \cdots \times \Delta^{n_{i-1}} \times f_{k_i}^i \times \Delta^{n_{i+1}} \times \Delta^{n_m}$. At each vertex $v_{j_1 \dots j_m}$, all facets except $\{F_{j_i}^i | 1 \leq i \leq m\}$ intersect with this vertex, where $v_{j_1 \dots j_m} = v_{j_1}^1 \times \cdots \times v_{j_m}^m$ and $v_{j_k}^k$ is the only vertex of Δ^{n_k} not contained in $f_{j_k}^k$.

Let $\lambda : \{F_{k_i}^i | 0 \leq k_i \leq n_i, 1 \leq i \leq m\} \rightarrow \mathbb{Z}_2^n$ be the characteristic map of the small cover M^n , the linear independence condition at vertex $v_{0 \dots 0}$ means that $\lambda(F_1^1), \dots, \lambda(F_1^{n_1}), \dots, \lambda(F_m^1), \dots, \lambda(F_m^{n_m})$ consist of a basis of \mathbb{Z}_2^n , which is defined to be $\{e_1, \dots, e_n\}$. Then F_i^0 for every $1 \leq i \leq m$ can be written as a linear combination of e_1, \dots, e_n . We set $\lambda(F_i^0) = \mathbf{a}_i$ for $1 \leq i \leq m$, where \mathbf{a}_i is the coordinate of $\lambda(F_i^0)$ related to the basis $\{e_1, \dots, e_n\}$. In this way, we have the correspondence $m \times m$ reduced vector matrix A of small cover M^n .

$$A = \begin{pmatrix} \mathbf{a}_1 \\ \vdots \\ \mathbf{a}_m \end{pmatrix} = \begin{pmatrix} \mathbf{a}_1^1 & \cdots & \mathbf{a}_1^m \\ \vdots & \cdots & \vdots \\ \mathbf{a}_m^1 & \cdots & \mathbf{a}_m^m \end{pmatrix}, \text{ where } \mathbf{a}_i^j = (a_{i1}^j, \dots, a_{in_i}^j).$$

We have more general versions of Lemma 2.1 and Theorem 2.2, which can be found in S. Choi, M. Masuda and D.Y. Suh [4] and also M. Masuda and T.E. Panov [8]. To state this lemma, the principal minor of a vector matrix is defined to be a principal minor of $A_{k_1 \dots k_m}$, where $1 \leq k_1 \leq n_1, \dots, 1 \leq k_m \leq n_m$. $A_{k_1 \dots k_m}$ is defined in S. Choi, M. Masuda and D.Y. Suh [4]. For given $1 \leq k_j \leq n_j$ with $j = 1, \dots, m$, $A_{k_1 \dots k_m}$ is the $m \times m$ submatrix of A whose j -th column is the k_j -th column of the $m \times n_j$ matrix $(\mathbf{a}_1^{jT}, \dots, \mathbf{a}_m^{jT})^T$. Thus

$$A_{k_1 \dots k_m} = \begin{pmatrix} a_{1k_1}^1 & \cdots & a_{1k_m}^m \\ \vdots & & \vdots \\ a_{mk_1}^1 & \cdots & a_{mk_m}^m \end{pmatrix}.$$

Lemma 3.1. (S. Choi, M. Masuda and D.Y. Suh [4]) *Let A be an $m \times m$ reduced vector matrix with \mathbb{Z}_2 -coefficient such that all the principal minors of A are 1, then A is conjugate to a unipotent upper triangular vector matrix of the following form:*

$$\begin{pmatrix} 1 & \mathbf{b}_1^2 & \mathbf{b}_1^3 & \cdots & \mathbf{b}_1^m \\ \mathbf{0} & 1 & \mathbf{b}_2^3 & \cdots & \mathbf{b}_2^m \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ \mathbf{0} & \cdots & \cdots & 1 & \mathbf{b}_{m-1}^m \\ \mathbf{0} & \cdots & \cdots & \mathbf{0} & 1 \end{pmatrix}$$

where $\mathbf{0} = (0, \dots, 0)$, $\mathbf{1} = (1, \dots, 1)$ of appropriate sizes.

Lemma 3.2. *The reduced vector matrix of a small cover M^n over a product of simplices P^n can be conjugated by a permutation vector matrix to a unipotent upper triangular vector matrix.*

Proof. By the linear independence of characteristic maps in vertex $v_{k_1 \dots k_m}$, the determinant of $A_{k_1 \dots k_m}$ is nonzero. More generally, because of the linear independence of characteristic maps in vertex $v_{k_1 \dots k_l 0 \dots 0}$ for $l \leq m$, the principal minors of $A_{k_1 \dots k_m}$ are nonzero, which leads to the conclusion. \square

Let the 1-cohomology classes $v_i^{(0)}, \dots, v_i^{(n_i)}$ dual to the facial submanifolds of facets $F_0^i, \dots, F_{n_i}^i$ for $1 \leq i \leq m$. For convenience, we can define: $u_i = v_i^{(0)}$. From the restrictions given by J and the unipotent upper triangular vector matrix, $v_i^{(j)}$ is the sum of u_i and the linear combination of u_1, \dots, u_{i-1} . In the following theorems, we will use y_j defined in $v_i^{(j)} = u_i + y_j$ if there is no ambiguity. Then y_j is a linear combination of u_1, \dots, u_{j-1} .

Given these conditions, we can extend the Lemma 2.3 of the last section to a broader one.

Lemma 3.3. *For every positive integer $k \leq m$, if $l_1 + \dots + l_k = \sum_{i=1}^k n_i + 1$,*

$$u_1^{l_1} \dots u_k^{l_k} = 0. \quad (3)$$

Proof. It is obvious to prove the equation when $k = m$. While $k < m$, it can be proved by induction. When $k = 1$, the equation is equivalent to $u_1^{n_1+1} = 0$. Because of the elements from the first column to the n_1 -th column of the *reduced matrix*, we have: $v_1^{(j)} = v_1^{(0)} = u_1$, the restriction of the Stanley-Reisner ideal means $0 = \prod_{j=0}^{n_1} v_1^{(j)} = u_1^{n_1+1}$. Assume that the equation is true at every number $\leq k$, then consider $u_1^{l_1} \dots u_{k+1}^{l_{k+1}}$. If $l_{k+1} \leq n_{k+1}$, then $l_1 + \dots + l_k \geq n_1 + \dots + n_k + 1$, it can be proved by the induction hypothesis. If $l_{k+1} \geq n_{k+1} + 1$, we first have:

$$\begin{aligned} 0 &= v_{k+1}^{(0)} \dots v_{k+1}^{(n_{k+1})} = u_{k+1}(u_{k+1} + y_1) \dots (u_{k+1} + y_{n_{k+1}}) \\ &= u_{k+1}^{n_{k+1}+1} + u_{k+1}^{n_{k+1}} a_1 + \dots + u_{k+1} a_{n_{k+1}}, \end{aligned}$$

where $a_1, \dots, a_{n_{k+1}}$ are some homogenous polynomials in elements u_1, \dots, u_k , $\deg(a_i) = i$, since y_j is a linear combination of u_1, \dots, u_{j-1} .

$$u_{k+1}^{l_{k+1}} = -u_{k+1}^{l_{k+1}-(n_{k+1}+1)}(u_{k+1}^{n_{k+1}} a_1 + \dots + u_{k+1} a_{n_{k+1}}),$$

Iterate this process, $u_{k+1}^{l_{k+1}}$ can be written as $u_{k+1}^j f_j$, where $\deg(f_j) = l_{k+1} - j$ and $j \leq n_{k+1}$. Since f_j is a homogenous polynomial over u_1, \dots, u_k , we can compute the degree of $f_j u_1^{l_1} \dots u_k^{l_k}$:

$$\deg(f_j u_1^{l_1} \dots u_k^{l_k}) = l_1 + \dots + l_{k+1} - j \geq n_1 + \dots + n_k + 1,$$

which means $f_j u_1^{l_1} \dots u_k^{l_k}$ is zero by induction. \square

Then we can use this lemma to give a more general theorem.

Theorem 3.4. *Let P^n be any product of simplices. Then any small cover N over $P^n \times \Delta^1$ has identically zero Stiefel-Whitney numbers, i.e. is a boundary.*

Proof. The Stiefel-Whitney class of N ,

$$w(N) = (\prod_{i=1}^m (1 + v_i^{(0)}) \cdots (1 + v_i^{(n_i)}))(1 + v_{m+1}^{(0)})(1 + v_{m+1}^{(1)}).$$

$$(1 + v_{m+1}^{(0)})(1 + v_{m+1}^{(1)}) = 1 + v_{m+1}^{(0)} + v_{m+1}^{(1)},$$

where $v_{m+1}^{(0)} + v_{m+1}^{(1)}$ is a linear combination of $u_1 \cdots u_m$. So when $i_1 + 2i_2 + \cdots + (n+1)i_{n+1} = n+1$, $w_1^{i_1} \cdots w_{n+1}^{i_{n+1}}$ consists of forms like $u_1^{l_1} \cdots u_m^{l_m}$, where $l_1 + \cdots + l_m = \sum_{i=1}^m n_i + 1 = n+1$. By the Lemma 3.3 above, we can conclude that $w_1^{i_1} \cdots w_{n+1}^{i_{n+1}}$ is zero for every i_1, \dots, i_{n+1} satisfying $i_1 + 2i_2 + \cdots + (n+1)i_{n+1} = n+1$, which implies that every Stiefel-Whitney number is zero. \square

Unlike the real Bott manifolds, a small cover N over $P^n \times \Delta^1$ may not bound equivariantly. Z. Lü and L. Yu [9] give examples of small covers over $\Delta_2 \times \Delta_1$, which do not bound equivariantly.

Indeed, there is a natural $\mathbb{R}P^1$ -bundle on N . Note that $\mathbb{R}P^1 \cong S^1$, if N is a S^1 -principal bundle, we can directly deduce that N is the boundary of a disk bundle. But actually we know the fact that if this S^1 -bundle N is not an orientable bundle, then this S^1 -bundle N is not principal. Furthermore, if N is not orientable, while the related small cover over P^n is orientable, this kind of N is not a principal bundle. The following proposition tells us how to judge the orientability of a real Bott manifold in the terms of its *reduce matrix*.

Proposition 3.5. *(Y. Kamishima and M. Masuda [7, Lemma 2.2]) The real Bott manifold M^n is orientable if and only if the sum of entries is zero in \mathbb{Z}_2 for each column of $E_n + A$, where A is the reduced matrix of M^n and E_n is the identity matrix of order n .*

Proof. By the computations in Section 2, we have the first Stiefel-Whitney class $w_1(M^n) = \sum_{j=1}^n y_j$. Let k_{ij} be the i -th row and the j -th column element in the matrix $A + E_n$, then by the Equation (2) in Section 2,

$$w_1(M^n) = \sum_{j=1}^n y_j = \sum_{j=1}^n \sum_{i=1}^n u_i k_{ij} = \sum_{i=1}^n u_i \left(\sum_{j=1}^n k_{ij} \right).$$

Since a manifold is orientable if and only if its first Stiefel-Whitney class is zero, we could see that M^n is orientable if and only if $\sum_{j=1}^n k_{ij} = 0$ for every i . \square

It is easy to see that there exist many upper triangular matrices of order $n+1$ that the sum of entries is zero for the first n columns, then we can just add the $n+1$ -column with the sum not equal to 0 and get the whole matrix $A + E_n$,

where the S^1 -bundle N is not an orientable bundle. There is an example of such matrix,

$$A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

In this example, the related small cover over P^n is a 2-torus T^2 , but N is an unoriented S^1 -bundle over T^2 .

We can also give a sufficient condition to determine when the small cover over $P^n \times \Delta^l$ has identically zero Stiefel-Whitney numbers. This condition is only related to the elementary symmetric polynomials of y_1, \dots, y_l , where y_j is defined by $v_{m+1}^{(j)} = u_{m+1} + y_j$, $1 \leq j \leq l$.

Theorem 3.6. *Let $l = 2^k - 1$ be a fixed number. For any product of simplices P^n , there exists a set of equations of y_1, \dots, y_l that the small cover over $P^n \times \Delta^l$ with the reduced matrix satisfying the equations will have identically zero Stiefel-Whitney numbers.*

Proof. Assume that y_1, \dots, y_l satisfy a set of equations $\{\sigma_i = 0 | 1 \leq i \leq l, i \notin l+1-2^j, j=0 \dots k\}$, where $\sigma_1, \dots, \sigma_l$ are elementary symmetric polynomials of y_1, \dots, y_l .

The Stiefel-Whitney class of N ,

$$\begin{aligned} w(N) &= (\prod_{i=1}^m (1 + v_i^{(0)}) \cdots (1 + v_i^{(n_i)})) (1 + u_{m+1}) (1 + u_{m+1} + y_1) \cdots (1 + u_{m+1} + y_l). \\ &\quad (1 + u_{m+1}) (1 + u_{m+1} + y_1) \cdots (1 + u_{m+1} + y_l) \\ &= (1 + u_{m+1})^{l+1} + (1 + u_{m+1})^l \sigma_1 + \cdots + (1 + u_{m+1}) \sigma_l. \end{aligned}$$

Only if j is the power of 2, $(1 + u_{m+1})^j$ will exactly contain 1 and the term of the highest degree. So, since $\sigma_i = 0$, when $1 \leq i \leq l, i \notin l+1-2^j$ and $j = 0, \dots, k$, then $(1 + u_{m+1})(1 + u_{m+1} + y_1) \cdots (1 + u_{m+1} + y_l)$ will only have polynomials of elements in u_1, \dots, u_m when modulo the ideals I and J . By almost the same process as the proof of the Theorem 3.4, we can conclude that every Stiefel-Whitney number of N is zero. \square

Example 1. When $k = 2, l = 3$, and P^n is any product of simplices, the small cover over $P^n \times \Delta^3$ has identically zero Stiefel-Whitney numbers, if the corresponding *reduced matrix* satisfies $\sigma_1 = 0$, i.e. $y_1 + y_2 + y_3 = 0$.

But generally, a small cover over $P^n \times \Delta^3$ may not have identically zero Stiefel-Whitney numbers. Take $\Delta^3 \times \Delta^3$ as an example. If $y_1 = y_2 = y_3 = u_1$, then the total Stiefel-Whitney class $w = (1 + u_1)^4 (1 + u_1 + u_2)^3 (1 + u_2)$. We can compute and get: $w_3^2 = u_2^4 (u_1 + u_2)^2$. Since the cohomology ring of this manifold is $\mathbb{Z}_2[u_1, u_2] / \langle u_1^4, u_2(u_1 + u_2)^3 \rangle$, and $u_2^4 (u_1 + u_2)^2$ doesn't belong to the ideal $\langle u_1^4, u_2(u_1 + u_2)^3 \rangle$, then $w_3^2 = u_2^4 (u_1 + u_2)^2$ is not zero.

4. COBORDISM OF GENERALIZED BOTT MANIFOLD AND QUASITORIC MANIFOLD

Suppose that M^{2n} is a quasitoric manifold over the product of simplices $P^n = \Delta^{n_1} \times \cdots \times \Delta^{n_m}$ of dimension n . Then let the second integral cohomology classes v_1, \dots, v_{n+m} dual to the facial submanifolds of facets F_1, \dots, F_{n+m} . For convenience, we will directly use the terms with the coefficient \mathbb{Z}_2 defined for the generalized real Bott manifolds, if there is no ambiguity. Furthermore, we have exactly the same equations as in Lemma 3.3. In this section, the *reduced matrix* is an integer matrix, and other terms are defined with \mathbb{Z} -coefficient if necessary.

Then we have the following lemma, which is an easy corollary of Theorem 1.1.

Lemma 4.1. *If we project every element in the reduced matrix A of M^{2n} into \mathbb{Z}_2 , we can get a \mathbb{Z}_2 -coefficient matrix \tilde{A} . If this matrix determines a small cover with identically zero Stiefel-Whitney numbers, then the reduced matrix A determines a quasitoric manifold with identically zero Stiefel-Whitney numbers.*

To extend our theorems above, we need to introduce the concept of generalized Bott manifold, which means it is not only a quasitoric manifold, but also a complex manifold. There are equivalent conditions for a quasitoric manifold to be a generalized real Bott manifold.

Theorem 4.2. *(S. Choi, M. Masuda and D.Y. Suh [4, Theorem 6.4]) Let M be a quasitoric manifold over a product of simplices P^n , and let A be the reduced vector matrix associated with M , which has $\mathbf{1}$ as the diagonal entries. Then the following are equivalent:*

- (1) *M is equivalent to a generalized Bott manifold.*
- (2) *M is equivalent to a quasitoric manifold which admits an invariant almost complex structure under the action.*
- (3) *the principal minors of A are 1.*

The quasitoric version of Lemma 3.1 (see S. Choi, M. Masuda and D.Y. Suh [4, Lemma 5.1]) confirms that the reduced vector matrix of a generalized Bott manifold is conjugate to a unipotent triangular upper vector matrix. Then the Theorem 3.4 can be extended to the case of a quasitoric manifold.

Theorem 4.3. *Let P^n be a product of simplices. The generalized Bott manifold $M^{2(n+1)}$ over $P^n \times \Delta^1$ has identically zero Stiefel-Whitney numbers and Pontryagin numbers, i.e. is a boundary.*

Proof. Owing to the Theorem 3.4 and the Lemma 4.1, the generalized Bott manifold $M^{2(n+1)}$ has identically zero Stiefel-Whitney numbers. Assume that $2(n+1) = 4k$, otherwise the Pontryagin numbers will be zero. From the terms defined in the Section 3, the total Pontryagin class of $M^{2(n+1)}$ is $p(M^{2(n+1)}) =$

$(\prod_{i=1}^m (1 - (v_i^{(0)})^2) \cdots (1 - (v_i^{(n_i)})^2))(1 - (v_{m+1}^{(0)})^2)(1 - (v_{m+1}^{(1)})^2)$. By the restriction of the Stanley-Reisner ideal, $v_{m+1}^{(0)}v_{m+1}^{(1)} = 0$, then $(1 - (v_{m+1}^{(0)})^2)(1 - (v_{m+1}^{(1)})^2) = 1 - (v_{m+1}^{(0)})^2 - (v_{m+1}^{(1)})^2 = 1 - (v_{m+1}^{(0)} + v_{m+1}^{(1)})^2$. Since $v_{m+1}^{(1)}$ is the sum of $-v_{m+1}^{(0)}$ and a linear combination of u_1, \dots, u_m , $v_{m+1}^{(0)} + v_{m+1}^{(1)}$ is a linear combination of u_1, \dots, u_m . Furthermore, every square term of $p(M^{2n})$, such as $(v_1^{(0)})^2, \dots, (v_m^{(n_m)})^2, (v_{m+1}^{(0)} + v_{m+1}^{(1)})^2$, has degree 4. Then every term $p_1^{i_1} \cdots p_k^{i_k}$ of degree $2(n+1)$ consists of terms which are products of square terms. Each of these products consists of homogenous polynomials of u_1, \dots, u_m with degree $2k = n+1$. So by the Equation (3) in Lemma 3.3, every $p_1^{i_1} \cdots p_k^{i_k}$ of degree $2(n+1)$ is zero. \square

If the principal minors of the *reduced matrix* are not identically 1, the quasitoric manifold $M^{2(n+1)}$ over a cube may or may not be null-cobordant. But, for the $4k$ -dimensional quasitoric manifold M over an even dimensional n -cube, if all the proper principal minors of the *reduced matrix* are 1, while the determinant of the whole *reduced matrix* is -1 , then this manifold is never a boundary. To state this conclusion, we need a lemma.

Lemma 4.4. (*M. Masuda and T.E. Panov [8, Theorem 3.3]*) *Let A be an $n \times n$ matrix with entries in \mathbb{Z} . Suppose that every proper principal minor of A is 1. If $\det A = -1$, then A is conjugated by a permutation matrix to the following matrix*

$$\begin{pmatrix} 1 & b_1 & 0 & \cdots & 0 \\ 0 & 1 & b_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & b_{n-1} \\ b_n & 0 & \cdots & 0 & 1 \end{pmatrix}$$

where $b_i \neq 0$.

Since the dimension $n(=2k)$ of the cube is even and the determinant is -1 , we have $b_1 \cdots b_n = 2$. Just as the case of real Bott manifold, Let v_1, \dots, v_n be the 2-cohomology classes dual to the facial submanifolds of facets F_1, \dots, F_n , meeting at a vertex. Then we have facets F_1^*, \dots, F_n^* which are parallel to F_1, \dots, F_n respectively. Let the 2-cohomology classes u_1, \dots, u_n be dual to the facial submanifolds of facets F_1^*, \dots, F_n^* .

Theorem 4.5. *For the $4k$ -dimensional quasitoric manifold M over n -cube, where $n = 2k$ and k is any positive number, if all the proper principal minors of the reduced matrix of M are 1, but the determinant of the reduced matrix of M is -1 , then this manifold has nonzero pontryagin numbers, i.e. is not a boundary.*

Proof. By the Lemma 4.4, the *reduced matrix* of M is conjugated by a permutation matrix to the matrix in Lemma 4.4. By the restrictions given by J , there

exist equations that: $-v_1 = u_1 + b_n u_n$ and $-v_i = u_i + b_{i-1} u_{i-1}$, for $2 \leq i \leq n$. The total Pontryagin class of M is

$$\begin{aligned} p(M) &= (1 - u_1^2)(1 - v_1^2) \cdots (1 - u_n^2)(1 - v_n^2) = (1 - (u_1 + v_1)^2) \cdots (1 - (u_n + v_n)^2) \\ &= (1 - (b_1 u_1)^2) \cdots (1 - (b_n u_n)^2), \end{aligned}$$

given by the restrictions of Stanley-Reisner Ideal I . This ideal also gives a group of relations: $u_1(u_1 + b_n u_n) = 0$ and $u_i(u_i + b_{i-1} u_{i-1}) = 0$, which determine the cohomology ring of M . The equality $b_1 \cdots b_n = 2$ implies that there exists exactly only one b_l equal to ± 2 , and other b_j are ± 1 . Then we consider u_j^n , where $j = 1, 2, \dots, n$. By the group of relations above, every u_j^n will be equal to $u_1 \cdots u_n$ multiplied by a coefficient. For example, $u_l^n = \pm u_1 \cdots u_n$, $u_{l-1}^n = \pm 2u_1 \cdots u_n$, \dots , $u_{l+1}^n = \pm 2^{n-1} u_1 \cdots u_n$. Then $\sum_{j=1}^n (b_j u_j)^n$ can be written as the sum of $\pm u_1 \cdots u_n$, $\pm 2u_1 \cdots u_n$, \dots , $\pm 2^n u_1 \cdots u_n$. But by the equation $1 + 2 + \cdots + 2^{n-1} < 2^n$, $\sum_{j=1}^n (b_j u_j)^n$ is a term of $u_1 \cdots u_n$ multiplied by a nonzero coefficient. For elementary symmetric polynomials $\sigma_1, \dots, \sigma_k$ and $s_k = \sum_{i=1}^n t_i^k$, the Newton-Girard formula states that:

$$(-1)^k s_k / k = \sum_{i_1 + 2i_2 + \cdots + k i_k = k} (-1)^{i_1 + \cdots + i_k} \frac{(i_1 + \cdots + i_k - 1)!}{i_1! \cdots i_k!} \sigma_1^{i_1} \cdots \sigma_k^{i_k}$$

This formula can be used in the ring $H^*(M; \mathbb{Z}) \otimes \mathbb{R}$. Obviously, p_1, \dots, p_k are elementary symmetric polynomials of $(b_1 u_1)^2, \dots, (b_n u_n)^2$, maybe with a sign. If all the Pontryagin numbers of M are zero, then $\sum_{j=1}^n (b_j u_j)^{2n}$ is zero by the formula, which is a contradiction. \square

The manifolds discussed in Theorem 4.5 are interesting. If it is over an odd dimensional cube, it is cobordant to zero by Lemma 4.1. But if it is over an even dimensional cube, it is unorientably cobordant to zero, but is not orientably cobordant to zero.

However, for the $4k$ -dimensional quasitoric manifold M over $2k$ -cube, even if the proper principal minors of the *reduced matrix* of M are not all 1, M might be orientably cobordant to zero.

Example 2. Assume that the *reduced matrix* of M is the following:

$$\begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix}, \text{ where } A_1 \text{ and } A_2 \text{ are of the form in Lemma 4.4.}$$

Let the orders of A_1 and A_2 , n_1 and n_2 , be odd. The total Pontryagin class of M is $p(M) = (1 - (b_1 u_1)^2) \cdots (1 - (b_{n_1} u_{n_1})^2)(1 - (c_1 u_{n_1+1})^2) \cdots (1 - (c_{n_2} u_{n_1+n_2})^2)$. Since every polynomial of degree more than n_1 over u_1, \dots, u_{n_1} is zero, and every polynomial of degree more than n_2 over $u_{n_1+1}, \dots, u_{n_1+n_2}$ is zero, we can deduce that the cohomology class $p_1^{i_1} \cdots p_k^{i_k}$ of degree $4k$ is zero.

We can apply the method in Theorem 4.5 to give a sufficient condition for a generalized Bott manifold to have nonzero Pontryagin numbers. In the following theorem, we will use the terms defined in the discussion of the generalized real Bott manifold. To be clear, we claim that y_i is defined by $v_m^{(i)} = -(u_m + y_i)$.

Theorem 4.6. *For the $4k$ -dimensional generalized Bott manifold M over a product of simplices P^n , where $n = 2k$ and k is any positive number, if we have the equation $u_m^n + (u_m + y_1)^n + \cdots + (u_m + y_{n_m})^n \neq 0$ in the integral cohomology ring of M , then this manifold has nonzero pontryagin numbers.*

Proof. The total Pontryagin class of M ,

$$\begin{aligned} p(M) &= \left(\prod_{i=1}^{m-1} (1 + (v_i^{(0)})^2) \cdots (1 + (v_i^{(n_i)})^2) \right) (1 + (v_m^{(0)})^2) \cdots (1 + (v_m^{(n_m)})^2) \\ &= \left(\prod_{i=1}^{m-1} (1 + (v_i^{(0)})^2) \cdots (1 + (v_i^{(n_i)})^2) \right) (1 + u_m^2) \cdots (1 + (u_m + y_{n_m})^2). \end{aligned}$$

By the method of the Theorem 4.5, if $\sum_{i=1}^{m-1} ((v_i^{(0)})^n + \cdots + (v_i^{(n_i)})^n) + u_m^n + (u_m + y_1)^n + \cdots + (u_m + y_{n_m})^n \neq 0$, then there exist nonzero Pontryagin numbers. By the Lemma 3.3, every term in $\sum_{i=1}^{m-1} ((v_i^{(0)})^n + \cdots + (v_i^{(n_i)})^n)$ is zero, which ends the proof. \square

Acknowledgments This work was motivated by a talk with Prof. L. Yu, the author's master mentor in Nanjing University. The author would also like to thank him for his useful suggestions.

REFERENCES

- [1] J.W. Milnor and J.D. Stasheff, *Characteristic classes*, Princeton University Press, 1974.
- [2] V.M. Buchstabe and T.E. Panov, *Torus actions and their applications in topology and combinatorics*, American Mathematical Society, 2002.
- [3] G.C. Hamrick; D.C. Royster *Flat Riemannian manifolds are boundaries*, Invent. Math. 66 (1982), no. 3, 405-413.
- [4] S. Choi, M. Masuda, D.Y. Suh, *Quasitoric manifolds over a product of simplices*, Osaka. J. Math. 47 (2010), no.1, 109-129.
- [5] M.W. Davis, T. Januszkiewicz, *Convex polytopes, Coxeter orbifolds and torus actions*, Duke Math. J. 62 (1991) 417-451.
- [6] Y. Cheng, Y. Wang, *The number of small covers over cubes and the product of at most three simplices up to equivariant cobordism*, Proc. Japan Acad. Ser. A Math. Sci. 87(2011), no.6, 95-98.
- [7] Y. Kamishima and M. Masuda *Cohomological rigidity of real Bott manifold*, Algebr. Geom. Topol. 9(2009), no.4, 2479-2502.
- [8] M. Masuda and T.E. Panov, *Semi-free circle actions, Bott towers, and quasitoric manifolds*, Mat. Sb. 199 (2008) 95-122.

- [9] Z. Lü and L. Yu, *Topological types of 3-dimensional small covers*, Forum Math.23 (2011), no.2, 245-284.
- [10] Z. Lü and Q. Tan, *small covers and the equivariant bordism classification of 2-torus manifolds*, Int. Math. Res. Not. IMRN 2014, no.24, 6756-6797.

DEPARTMENT OF MATHEMATICS AND IMS, NANJING UNIVERSITY, NANJING, 210093,
P.R.CHINA

E-mail address: luyxnj@gmail.com